1 **Towards a genome sequence for every animal: where are we now?**
2

3 Scott Hotaling[1], Joanna L. Kelley[1], and Paul B. Frandsen[2,3,4]
4
5 **Affiliations:**
6 [1] School of Biological Sciences, Washington State University, Pullman, WA, USA
7 [2] LOEWE Centre for Translational Biodiversity Genomics (LOEWE- TBG), Frankfurt, Germany
8 [3] Department of Plant and Wildlife Sciences, Brigham Young University, Provo, UT, USA
9 [4] Data Science Lab, Smithsonian Institution, Washington, DC, USA
10
11 **Correspondence:**
12 Scott Hotaling, School of Biological Sciences, Washington State University, Pullman, WA,
13 99164, USA; Email: scott.hotaling@wsu.edu: Phone: (828) 507-9950
14
15 Paul B. Frandsen, Department of Plant and Wildlife Sciences, Brigham Young University, Provo,
16 UT, 84602, USA; Email: paul_frandsen@byu.edu; Phone: (804) 422-2283
17

18 **Keywords:** animal genomes, genome biology, Arthropoda, Mammalia, Metazoa, genomic
19 natural history
20

Abstract:
In less than 25 years, the field of animal genome science has transformed from a discipline
seeking its first glimpses into genome sequences across the Tree of Life to a global enterprise
with ambitions to sequence genomes for all of Earth's eukaryotic diversity (1). As the field
rapidly moves forward, it is important to take stock of the progress that has been made to best
inform the discipline's future. In this Perspective, we provide a contemporary, quantitative
overview of animal genome sequencing. We identified the best available genome assemblies on
GenBank, the world's most extensive genetic database, for 3,278 unique animal species across
24 phyla. We assessed taxonomic representation, assembly quality, and annotation status for
major clades. We show that while tremendous taxonomic progress has occurred, stark
disparities in genomic representation exist, highlighted by a systemic overrepresentation of
vertebrates and underrepresentation of arthropods. In terms of assembly quality, long-read
sequencing has dramatically improved contiguity, whereas gene annotations are available for
just 34.3% of taxa. Furthermore, we show that animal genome science has diversified in recent
years with an ever-expanding pool of researchers participating. However, the field still appears
to be dominated by institutions in the Global North, which have been listed as the submitting
institution for 77% of all assemblies. We conclude by offering recommendations for how we can
collectively improve genomic resource availability and value while also broadening global
representation.
40

Significance statement:
The field of animal genome science is rapidly developing, and efforts are underway to sequence
genomes for all of Earth's eukaryotic biodiversity. Here, we provide an overview of animal
genome sequencing, with emphases on taxonomic representation, assembly quality, and
geographic representation. We show that while a staggering 3,278 unique animal species have
had their genomes sequenced, massive disparities exist in terms of the taxonomic groups
receiving attention, the quality of the resources being produced, and the institutions driving the
field. We highlight areas where improvements can be made, notably by continuing to increase
the quality of genome assemblies, including by improving metadata and voucher specimen
associations, and actively developing meaningful collaborations between researchers form the
Global North and South.

52
Introduction:
The first animal genome sequence was published 23 years ago (2). The 97 million base pair (Mb) *Caenorhabditis elegans* genome assembly ushered in a new era of animal genome biology where genetic patterns and processes could be investigated at genome scales. As genome assemblies have accumulated for an increasingly diverse set of species, so too has our knowledge of how genomes vary and shape Earth's biodiversity (e.g., 3, 4). Major shifts in genome availability and quality have been driven by two key events. First, the invention of high-throughput, short-read sequencing provided an economical means to generate millions of reads for any species from which sufficient DNA could be obtained. These ~100-bp short reads could be assembled into useful, albeit fragmented, genome assemblies. Later, the rise of long-read sequencing allowed for similarly economical generation of reads that are commonly orders of magnitude longer than short reads, resulting in vastly more contiguous genome assemblies (5).

We have now entered an era of genomic natural history. Building on ~250 years of natural history efforts to describe and classify the morphological diversity of life on Earth, we are gaining a complementary genomic perspective of Earth's biodiversity. However, a baseline accounting of our progress towards a complete perspective of Earth's genomic natural history—where every species has a corresponding, reference-quality genome assembly available—has not been presented. This knowledge gap is particularly important given the momentum towards sequencing all animal genomes which is being driven by a host of sequencing consortia. For instance, the Vertebrates Genome Project seeks to generate high-quality assemblies for all vertebrates (6), the Bird10K project seeks to generate assemblies for all extant birds (7), the i5K Project plans to produce 5,000 arthropod genome assemblies (8), the Earth BioGenome Project aims to sequence all eukaryote genomes (1), and the Darwin Tree of Life project plans to sequence all eukaryotes in Britain and Ireland (https://www.darwintreeoflife.org/).

In this perspective, we curated, quantified, and summarized genomic progress for a major component of Earth's biodiversity: kingdom Animalia (Metazoa) and its roughly 1.66 million described species (9). We show that as of June 2021, 3,278 unique animals have had their nuclear genome sequenced and the assembly made publicly available on the National Center for Biotechnology Information (NCBI) GenBank database (10). This translates to 0.2% of all animal species, but when viewed through the lens of major clades, massive disparities exist. For instance, 32x more assemblies are available for chordates relative to arthropods (Fig. 1).

Methods:
To construct a database of the best available genome assembly for all animals, we downloaded metadata from GenBank for all kingdom Animalia taxa using the "summary genome" function in v.10.9.0 of the NCBI data sets command line tool on 4 February 2021. Next, we used the TaxonKit (11) "lineage" function to retrieve taxonomic information for each taxid included in the genome metadata. To gather additional data for each assembly (e.g., sequencing technology), we used a custom web scraper script. Both this web scraper script and the scripts used to download and organize the metadata are available on this study's GitHub repository (https://github.com/pbfrandsen/metazoa_assemblies). We later supplemented this initial data set with a second round of metadata acquisition on 28 June 2021. For the full data set, we hand-refined the NCBI taxonomy classifications to subdivide our data set into three categories: species, subspecies, or hybrids (Table S1). If replicate assemblies for a taxon were present, we defined the "best available" assembly as the one with the highest contig N50 (the mid-point of the contig distribution where 50% of the genome is assembled into contigs of a given length or longer).

Figure 1. Variation in taxonomic richness and genome availability, quality, and assembly size across kingdom Animalia on GenBank (as of 28 June 2021). Taxonomic groups are clustered by phylogeny following (12). Only groups with 30 or more available assemblies as of January 2021 are shown with the exception of Hominidae ($n = 5$ genomes). In the tree, bold group names represent phyla and naming conventions follow those of the NCBI database. Of 34 recognized animal phyla, 10 do not have a representative genome sequence. (a) The total number of described species for each group following Zhang (9) and the references therein. (b) Genomic representation among animal groups for 3,278 species with available genome assemblies. Bars represent the magnitude of the observed minus the expected number of genomes given the proportion that each group comprises of described animal diversity. Significance was assessed with Fisher's exact tests and significantly under- or over-represented groups ($P < 0.05$) are denoted with asterisks. Gray numbers indicate the total number of species with available genome assemblies for each group. The number of available assemblies is not mutually exclusive with taxonomy; that is, a carnivore genome assembly would be counted in three categories (Order Carnivora, Class Mammalia, Phylum Chordata). (c) The percentage of described species within a group with an available genome sequence (bars) and the percentage of those assemblies that have corresponding annotations (red circles). For many groups (e.g., arthropods), only a fraction of a percent of all species have an available genome assembly, making their percentage appear near zero. (d) Assembly size for all animal genome assemblies, grouped by taxonomy. (e) Contig N50 by taxonomic group. The sequencing technology used for each assembly is denoted by circle fill color: short-read (blue), long-read (yellow), or not provided (gray). In (d) and (e), each circle represents one genome assembly and a few notable or outlier taxa are indicated with gray text.

We filtered our data in several ways: we removed subspecies (unless they were the only representative for a species) and hybrids, assemblies that were shorter than 15.3 Mb [the smallest confirmed assembly size for a metazoan to date, (13)], or had a contig N50 less than 1 kilobase (Kb). We also culled assemblies that were unusually short (i.e., 1-2.5 Mb) and had information in their descriptions that indicated they were not true nuclear genome assemblies (e.g., "exon capture"). In total, we culled 407 assemblies that did not fit the above criteria. The remaining assemblies were classified as "short-read", "long-read", or "not provided" based on whether only short-reads (e.g., Illumina) were used, any long-read sequences (e.g., PacBio) were used, or no information was available. We defined a species as having gene annotations available if any assembly for that taxon also had annotations on GenBank. When the best available assembly did not have annotations included or when multiple assemblies had annotations, we retained the annotations for the assembly with the highest contig N50. Finally, we used the submitting institution for each assembly as a surrogate for the institution that led the genome assembly effort. Using these data, we classified assemblies to a country, region (Africa, Asia, Europe, Middle East, North America, Oceania, South America, Southeast Asia), and the Global North (e.g., Australia, Canada, Europe, USA) or Global South (e.g., Africa, Asia including China, Mexico, Middle East, South America).

To test if clades were under- or overrepresented in terms of genome availability relative to their species richness, we compared the observed number of species with assemblies to the expected total for the group. We obtained totals for the number of described species overall and for each group from previous studies, primarily from Zhang (9) and the references therein. We assessed significance between observed and expected representation with Fisher's exact tests (alpha = 0.05). We tested for differences in distributions of contig N50 or assembly size between short- and long-read genomes with Welch's T-tests. For both display (i.e., Figure 1) and analysis, we subdivided the data set into the lowest taxonomic level that still contained 30 or more assemblies as of January 2021 (with the exception of hominids which were given their own category due to their exceptionally high genomic resource quality).

155
156
Figure 2. Genome availability for kingdom Animalia versus taxonomic descriptions and over time.
(a) The proportion of described taxonomic groups versus the number with sequenced genome
assemblies from phyla to species. The gray plot to the right is a zoomed in perspective of the higher
taxonomic level categories in the full plot to the left. For genus through phylum, the number of described
categories is based on the NCBI taxonomy. For species, the total number described is from Zhang (9). (b)
The timeline of genome contiguity versus availability for animals according to the GenBank publication
date (x-axis, see panel c). A rise in assembly contiguity has been precipitated by long-read sequencing.
Particularly contiguous assemblies for a given time period are labeled. (c) The number of animal genome
assemblies deposited on GenBank each month since February 2004. Several notable events are labeled.
When specific dates are indicated, those (and the assemblies referred to) are included within that month's
total. For (b) and (c), it is important to note that when a genome assembly is updated to a newer version,
its associated date is also updated. Thus, the date associated with many early animal assemblies [e.g.,
Caenorhabditis elegans, (2)] has shifted to be more recent with updates.
170
Results:
Taxonomic representation
Genome assemblies were available for 3,278 species representing 24 phyla, 64 classes, and
258 orders (Fig. 2a, Table S1). The data set was exceptionally enriched for the phylum
Chordata (which includes all vertebrates) with 1,770 assemblies for the group (54% of all
assemblies) despite chordates comprising just 3.9% of animal species (*P,* Fisher's < 1e-5; Fig.
1). Conversely, arthropods were underrepresented with 1,115 assemblies (34% of the data set)
for a group that comprises 78.5% of animal species (*P,* Fisher's < 1e-5; Fig. 1). However, not all
arthropods were underrepresented; five insect clades were overrepresented [Apidae (bees),
Culicidae (mosquitoes), *Drosophila* (fruit flies), Formicidae (ants), and Lepidoptera (butterflies
and moths); all *P,* Fisher's < 1e-3; Fig. 1]. Collectively, of the 59 animal taxonomic groups
included in our data set, 14 groups were underrepresented, 17 were represented as expected,
and 28 were overrepresented (primarily chordates; Fig. 1). Ten phyla had no publicly available
genome sequence (Fig. 1). Over the ~17-year GenBank genome assembly record, animal
assemblies have been deposited at a rate of 0.52 species' assemblies per day. Over the most
recent year, however, this rate increased eight-fold to 4.07 assemblies per day. If the most

recent rate were maintained, all currently described animals would have a genome assembly
available by 3136. To achieve this goal by 2031 instead, an average of 165,614 novel animal
genomes would need to be sequenced and assembled each year (~112x faster than the rate for
the most recent year).
191
Assembly size, contiguity, and annotations
The average animal genome assembly was 1.02 gigabases (Gb) in length (SD = 1.21 Gb) with
a contig N50 of 2.26 Mb (SD = 25.16 Mb; Fig. 1d,e). Two animal genome assemblies were 25
Gb longer than all other assemblies—the axolotl [32.4 Gb (14)] and Australian lungfish [34.6 Gb
(15); Fig. 1d]. The smallest genome assembly in the data set, the mite *Aculops lycopersici*, was
over 1000x smaller, spanning just 32.5 Mb (16). Still smaller is the 15.3 Mb assembly of the
marine parasite, *Intoshia variabili*, which has the smallest animal genome currently known (13).
But, since the *I. variabili* assembly was not available on GenBank as of June 2021, it was not
included in our data set.
201
Contiguity varied dramatically across groups. For instance, hominid assemblies (family
Hominidae, n = 5) were the most contiguous with an average contig N50 of 24.2 Mb. Bird
assemblies (class Aves, n = 515) were also highly contiguous (mean contig N50 = 1.4 Mb)
despite being so numerous (and accumulating for a long period of time). On the other end of the
spectrum, jellyfish and related species (phylum Cnidaria) exhibited some of the least contiguous
genome assemblies with a mean contig N50 of 0.18 Mb (n =65; Fig. 1e). Roughly 34% of
animals with genome assemblies had corresponding annotations on GenBank but annotation
rates differed substantially among groups (Fig. 1c). For example, the rate of arthropod
annotations (22.3%) lags behind that for chordates (41.3%); however much of this disparity
appeared to be driven by the low and high annotation rates of butterflies and moths (order
Lepidoptera) and birds (class Aves), respectively. Of 445 assemblies, just 6.5% of lepidopteran
assemblies on GenBank have corresponding annotations versus 72.8% of birds (n = 519
assemblies; Fig. 1c). Notably, since most gene models are based on sequence similarity to
known functional genes and not functional data, the true rate of annotation is likely even lower
than reported here.
217
Geographic representation
Animal genome assemblies have been contributed by researchers at institutions on every
continent with permanent inhabitants, including 52 countries. From a regional perspective,
institutions in North America (n = 1,331), Europe (n = 972), and Asia (n = 828) collectively
accounted for 95.5% of all assemblies (Fig. 3a). And, nearly 70% of all animal genome
assemblies have been submitted by researchers in just three countries: USA (n = 1,275), China
(n = 676), and Switzerland (n = 317; Fig. 3a). When countries were grouped by their inclusion in
the Global North or South, similarly stark patterns emerged. Researchers affiliated with
institutions in the Global North contributed roughly 75% of animal genome assemblies (Fig. 3b).
From a taxonomic perspective, researchers at North American institutions have contributed the
most insect and mammal assemblies, European researchers have contributed the most fish
assemblies, and Asian researchers the most bird assemblies (Fig. 3a). The first assembly in
GenBank from the Global North was deposited in 2004 and the first assembly from the Global
South was deposited in 2011 (Fig. 3c). Since then, the number of assemblies deposited each
year has steadily risen with the proportions from the Global North and South staying relatively
constant (Fig. 3c).
234

Figure 3. Where animal genome assemblies have been produced around the world according to the submitting institutions on GenBank. (a) For each geographic region, total numbers of genome assemblies are shown by dark circles with white lettering. This total is further broken down by country and taxon. For regions where more than four countries have contributed assemblies (e.g., Europe), an "Other" category represents all other countries. The same applies to all assemblies that are not insects, birds, fish, or mammals in the taxon plots. Countries are color-coded by assignment to the Global North or South. (b) The total number of genome assemblies contributed by countries in the Global North (e.g., USA, Europe, Australia) versus the Global South (e.g., Africa, South America, China, Mexico, Middle East). (c) The rate of genome assembly deposition by major sources in the Global North (Europe, USA) and Global South [China, Southeast (SE) Asia] as well as all other countries collectively in each (Other).

Use of long-reads in genome assemblies and availability of key metadata also differs with geography. For assemblies deposited since 2018, researchers from the Global South have used long-reads slightly more frequently than those from the Global North (25.7% versus 20.2%; Fig. 4a). However, researchers from the Global North were far less likely to report the types of sequence data used (19.9% of assemblies for the Global North versus 1.4% of assemblies for the Global South; Fig. 4a). Much of this difference appears to be driven by genome assemblies deposited by researchers at European institutions (Fig. 4b). This gap in metadata may reflect an issue with data mirroring between the European Nucleotide Archive (ENA) and GenBank. For instance, many new genome assemblies being generated by the United Kingdom, for example,

257 are part of the Wellcome Sanger Institute's Darwin Tree of Life project, which is generating
258 exceptionally high-quality assemblies using long-read sequencing and depositing them into the
259 ENA (Fig. 5). One region (Oceania) and three countries (Australia, Finland, India) reported long
260 reads being used in more than 50% of deposited assemblies (Fig. 4b,c).
261

262
263
264 **Figure 4. Sequencing technologies used around the world: (a) between the Global North versus**
265 **Global South, (b) among regions, and (c) among countries.** To limit bias due to the limited availability
266 of long-read sequencing technologies before ~2017 (see Fig. 2b), only assemblies deposited on or after 1
267 January 2018 were included in the analysis and in (c) only countries that deposited five or more
268 assemblies during the focal period (January 2018 – June 2021) are shown.
269
Discussion:
Taxonomic representation
Animal genome sequencing has dramatically progressed in the last 25 years. In that span, the
field has moved from sequencing the first nuclear genome for any animal (2)—a landmark
achievement—to targeting the generation of genome assemblies for *all* of Earth's eukaryotic
biodiversity (1). Here, we provided a contemporary perspective on progress towards this goal
for the 1.6-million species in the animal kingdom (9). We showed that while tremendous
progress has been made, major gaps and biases remain both in terms of taxonomic and
geographic representation, at least within the most commonly used database of genomic
resources, GenBank. For instance, a major bias exists in favor of vertebrates which are vastly
overrepresented relative to their total species diversity (Fig. 1a-c). From the perspectives of

biomedicine and human evolution, this bias is reasonable since humans are vertebrates.
However, from a basic research perspective, particularly as it relates to genomic natural history
and an overarching goal to sequence all animal genomes, there is a need to taxonomically
diversify sequencing efforts.
285
At the highest taxonomic levels, 10 animal phyla still have no genomic representation. To
illustrate the scale of this disparity versus other groups and the unique biology that is being
overlooked, genome assemblies are available for 685 ray-finned fishes (class Actinopterygii) but
none exists for phylum Nematomorpha, a ~2,000-species clade of parasitic worms whose
presence can dramatically alter energy budgets of entire stream ecosystems (17). Another
phylum without genomic representation—Loricifera—was first described in 1983 (18). This
group of small, sediment-dwelling animals includes the only examples of multicellular species
that spend their entire life cycles under permanently anoxic conditions (19). Loriciferans
accomplish this feat by foregoing the energy-producing mitochondria found in virtually all
animals in favor of hydrogenosome-like organelles akin to those found in prokaryotes inhabiting
anaerobic habitats (19). Clearly, there is much to discover in terms of genomic diversity and
functional biology in undersampled clades.
298
Global representation
A few select countries—primarily the USA, several European nations, and China—have led the
sequencing of the vast majority of animal genome assemblies (Fig. 3a). Aside from China, all of
these countries are within the Global North. This pattern of geographic bias raises two potential
issues for representation in animal genome science. First, the researcher population of animal
genome sequencing likely does not reflect the global population. Second, sampling biases may
exist towards the regions where most of the genome sequencing is occurring. Some of this
"bias" is intentional and reflects funding goals for a given region. For instance, the Darwin Tree
of Life project (https://www.darwintreeoflife.org/) seeks to sequence the genomes of all ~70,000
eukaryotic species living in Britain and Ireland. Still, however, similar to how sampling biases
can yield skewed understanding of the natural world in other disciplines (e.g., 20), so too could
bias towards specific ecoregions, habitats, or other classifications skew genomic insight.
311
Inherently linked with questions of representation in animal genome science is the specter of
parachute science (or helicopter research)—the practice where international scientists, typically
from wealthy nations, conduct studies in other countries that are often poorer without
meaningful communication nor collaborations with local people (21). Parachute science has a
long history in ecological research, and signatures of these practices have been observed for
genome sciences. For instance, Marks *et al.* (22) found that the majority of plant genome
assemblies for species that are native to South America and Africa were sequenced "off-
continent" by researchers at European, North American, or Asian institutions. Given the sheer
number of animal genome assemblies that have been submitted by a small number of countries
and institutions, a similar pattern likely exists for animal genome science. However, to properly
assess this issue, parsing authorship to quantify collaboration, at a minimum, would need to
occur and this approach would still overlook key aspects of representation that need to be
considered (e.g., if a researcher from the Global South is working at an institution in the Global
North).
326

327
328
Figure 5. Examples of major contributors of genome assemblies for (a) butterflies (order
Lepidoptera), (b) birds (class Aves), and fish (primarily class Actinopterygii). Major contributors
were defined as any consortium, organization, or project that has deposited more than 5% of all
assemblies for butterflies and birds or 2.5% of all assemblies for fish.

Assembly quality varies dramatically within and among clades
For the purpose of biological discovery, not all genome assemblies are created equal. As long-
read sequencing technologies have matured, so too have the quality of assemblies being
generated (5). In the last year alone, the largest ever animal genome assembly was deposited
[Australian lungfish, (15)] as well as the most complete human genome to date, a telomere-to-
telomere assembly (23). Still, many species on GenBank only have low-quality assemblies
available (i.e., contig N50 < 100 Kb with no corresponding gene annotations; Fig. 1). Since
fragmentation and/or poor or missing gene annotations reduces the research value of an
assembly, genome quality is an important consideration, particularly when the end goal is
resource development for a broader community. As of April 2021, the Earth BioGenome Project
sought assembly quality of 6.C.Q40 (https://www.earthbiogenome.org/assembly-standards) for
reference genomes where "6" refers to a 1e6 contig N50 (i.e., 1 Mb). In our data set, 568
assemblies (17.3%) reach this contiguity standard. And, that number drops to 271 assemblies
(8.3%) when contig N50 ≥ 1 Mb and deposited gene annotations are both required. For
reference, the "C" above refers to chromosomal scale scaffolding and "Q40" to a less than
1/10,000 error rate. Neither of these metrics were assessed in this study.

Independent research laboratories, institutions, and consortia have contributed genome
assemblies on both ends of the quality spectrum (Fig. 5). For example, among butterflies (order
Lepidoptera), a bimodal quality distribution is being primarily driven by contributions made in
2021 by two submitting institutions, the Florida Museum of Natural History (e.g., 24) and the
Wellcome Sanger Institute (Fig. 5a). When viewing genome assembly contributions holistically
across the animal Tree of Life, it is clear that two consortia—the Vertebrate Genomes Project
(6) and the Darwin Tree of Life (https://www.darwintreeoflife.org/), part of the Wellcome Sanger
Institute—warrant specific recognition for contributing exceptional genomic resources relative to
closely related species (Fig. 5a-c).

Going forward

While animal genome science has dramatically matured in recent years, the field still rests on
the cusp of massive change. Thousands of genome assemblies are now available for a wide
range of taxa, a resource that can empower unprecedented scales of genomic comparison.
Simultaneously, multiple consortia are building momentum towards their goals and generating
some of the highest quality genome assemblies ever produced. The field is also diversifying
with researchers around the world, particularly from the Global South, leading a rising number of
efforts. These ongoing advances will yield higher quality, more globally representative genomic
data for animals. As we collectively build towards this new genomic future, we offer
recommendations to improve assembly quality and accessibility while also continuing to
increase representation within the discipline.

The quality of a genome assembly is likely the most important factor dictating its long-term
value. Genome assembly "quality," however, is difficult to define. Here, we propose a holistic
view on genome assembly quality that generally echoes the guidelines proposed by the Earth
BioGenome Project and other consortia. Briefly, assemblies should reach minimum levels of
contiguity (e.g., contig N50 > 1 Mb) and accuracy in order to be considered a reference that will
likely not need to be updated for most applications. At a minimum, assemblies should also
include high-quality gene annotations that perhaps take advantage of standardized pipelines
[e.g., NCBI Eukaryotic Genome Annotation Pipeline, (25)] to maximize compatibility across taxa.
We recommend the field further improve the quality of genome assembly resources in two
ways. First, refining and expanding the coordinated deposition of genome assemblies will
improve the usability of the resources and reproducibility of analyses. It will also reduce
duplications of effort—i.e., when a group sequences a genome that has already been
produced—an issue that is likely to become increasingly common.

To refine and expand coordinated resource deposition, we recommend the continued use of
GenBank (10) or one of the other archives that are members of the International Nucleotide
Sequence Database Collaboration (INSDC)—the European Nucleotide Archive (ENA) and DNA
Database of Japan (DDBJ)—as the central repositories for genome assemblies and their
metadata given their tripartite data sharing agreement. Next, we call on genetic archive
administrators, consortia, and independent researchers to collectively improve the metadata
submitted with each assembly and the mirroring of data across repositories. Too many
assemblies lack basic information about the sequence data and methods used (e.g., Fig. 4),
and with the difficulty of linking assemblies to published studies (if available), it can be
challenging or impossible to find this information. Further, an expansion of the metadata
associated with each assembly—ideally to make more of the categories required and expand
demographic data—would make efforts to quantify geographic representation, for instance, far
more straightforward. Alternatively, the metadata associated with genome assembly accessions
could be integrated with existing efforts like the Genomic Observatories Metadatabase (GeOMe;
26). Furthermore, a set of minimum quality characteristics for a genome assembly may need to
be defined. A number of exceptionally low-quality "genome assemblies" (e.g., with contig N50
values shorter than 1 Kb) that often cover only a small fraction of the expected total genome
sequence length for a given group are present on GenBank. The presence of these assemblies
raises the question: where is the inflection point between resource quality and value to other
researchers versus diluting the resources of a shared repository?

For our second recommendation, we amplify and expand the message of Buckner *et al.* (27)
and Thompson *et al.* (28): genome science needs specimen vouchers. Vouchers serve as a key
physical link between taxonomy and molecular insight. Rarely, however, are vouchers
referenced in publications of genome assemblies; only 11% of vertebrate assemblies included

such a reference as of January 2020 (27). While vouchers represent a physical reference for
assessing taxonomic classification or morphological variation, a properly stored voucher could
also provide a long-term source of material for future resource improvement. If a physical
specimen cannot be deposited, photographs and/or genomic DNA should be deposited in its
place (e.g., 29). Tied to the metadata discussion above, additional fields should be added to
GenBank genome assembly accessions to directly link the assembly to a specimen, photos, or
genomic DNA that has been deposited elsewhere.
419
Though geographic representation in animal genome science has improved in recent years, the
discipline appears far from properly reflecting the global researcher pool. This issue is almost
certainly multi-faceted, likely stemming from a lack of infrastructure (e.g., fewer high-throughput
sequencing platforms in developing countries), fewer resources for expensive molecular
research, and a corresponding lack of training in genome data analysis. To bridge this gap and
to empower a more diverse discipline, the nations and institutions that are devoting large
amounts of resources to animal genome sequencing (e.g., China, United Kingdom, USA), and
the researchers within those countries, should develop meaningful collaborations with
researchers within the countries where their focal species reside (30). These meaningful
collaborations—where all parties are valued for their expertise and involved in decision-
making—improve the science through transfer of local knowledge, provide a means for local
researchers to expand their skillset and network while raising their scholarly profile, and most
importantly, can effectively end the practice of parachute science (30). Within-continent (or
country) initiatives also have transformative potential for people and genome research. For
instance, the African-led effort to sequence three million African genomes over the next 10
years (the "3MAG" project) will yield massive investment in African genomics, an incredible
resource for understanding the full scope of human genetic diversity, and a new generation of
African genome scientists (31). While focused on human genetics, the infrastructure and
expertise that arise from the 3MAG project will no doubt translate to other taxa in the coming
years.
440
A practical justification also exists for increasing representation in genome science, particularly
as we seek to generate genome assemblies for every animal on Earth. The Global South is
home to the bulk of the world's biodiversity (32) and as such, researchers in these regions have
greater access to key habitats and specimens. Thus, it behooves everyone, including
researchers in the Global North, to deepen collaborations with peers in the Global South while
also helping to build indigenous capacity for collection, storage, and sequencing of new
specimens.
448
Conclusion:
Animal genome science continues to grow and expand at an exceptional rate. The coming
years will surely see thousands, and perhaps tens of thousands, of new genome assemblies
from across the Tree of Life, massive technological and analytical improvements, and some of
the largest scale and most in-depth studies of animal genome biology conducted to date.
However, if we are to realize the ambitious goals of efforts like the Earth BioGenome Project—a
self-described biological "moonshot"—the rate and mean quality of animal genome assembly
production will have to increase by roughly two orders of magnitude. Regardless of rates and
timelines, however, perhaps the most important goal for the future of animal genome science is
that we empower a more diverse, representative researcher community in parallel with the
generation of new resources.
460

Acknowledgements:
S.H. and J.L.K. were supported by NSF award OPP-1906015. We would like to thank Guangfeng Song, Eric Cox, and Anne Ketter from the "datasets" development team at the NCBI for their responsiveness and receptiveness to improving this valuable tool for data science.

References:
1. H. A. Lewin *et al.*, Earth BioGenome Project: Sequencing life for the future of life. *Proceedings of the National Academy of Sciences* **115**, 4325-4333 (2018).
2. C. elegans Sequencing Consortium, Genome sequence of the nematode C. elegans: a platform for investigating biology. *Science* **282**, 2012-2018 (1998).
3. C. Hoencamp *et al.*, 3D genomics across the tree of life reveals condensin II as a determinant of architecture type. *Science* **372**, 984-989 (2021).
4. G. W. Thomas *et al.*, Gene content evolution in the arthropods. *Genome Biology* **21**, 1-14 (2020).
5. S. Hotaling *et al.*, Long-reads are revolutionizing 20 years of insect genome sequencing. *Genome Biology and Evolution* (2021).
6. A. Rhie *et al.*, Towards complete and error-free genome assemblies of all vertebrate species. *Nature* **592**, 737-746 (2021).
7. G. Zhang, Bird sequencing project takes off. *Nature* **522**, 34-34 (2015).
8. i5K Consortium, The i5K Initiative: advancing arthropod genomics for knowledge, human health, agriculture, and the environment. *Journal of Heredity* **104**, 595-600 (2013).
9. Z.-Q. Zhang, *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness* (Magnolia press, 2011).
10. E. W. Sayers *et al.*, GenBank. *Nucleic Acids Research* **48**, D84-D86 (2021).
11. W. Shen, J. Xiong, TaxonKit: a cross-platform and efficient NCBI taxonomy toolkit. *Biorxiv*, 513523 (2019).
12. C. E. Laumer *et al.*, Revisiting metazoan phylogeny with genomic sampling of all phyla. *Proceedings of the royal society B* **286**, 20190831 (2019).
13. G. S. Slyusarev, V. V. Starunov, A. S. Bondarenko, N. A. Zorina, N. I. Bondarenko, Extreme genome and nervous system streamlining in the invertebrate parasite Intoshia variabili. *Current Biology* **30**, 1292-1298. e1293 (2020).
14. S. Nowoshilow *et al.*, The axolotl genome and the evolution of key tissue formation regulators. *Nature* **554**, 50-55 (2018).
15. A. Meyer *et al.*, Giant lungfish genome elucidates the conquest of land by vertebrates. *Nature* **590**, 284-289 (2021).
16. R. Greenhalgh *et al.*, Genome streamlining in a minute herbivore that manipulates its host plant. *Elife* **9**, e56689 (2020).
17. T. Sato *et al.*, Nematomorph parasites drive energy flow through a riparian ecosystem. *Ecology* **92**, 201-207 (2011).
18. R. M. Kristensen, An introduction to loricifera, cycliophora, and micrognathozoa. *Integrative and Comparative Biology* **42**, 641-651 (2002).
19. R. Danovaro *et al.*, The first metazoa living in permanently anoxic conditions. *BMC biology* **8**, 1-10 (2010).
20. A. C. Hughes *et al.*, Sampling biases shape our view of the natural world. *Ecography* (2021).
21. P. V. Stefanoudis *et al.*, Turning the tide of parachute science. *Current Biology* **31**, R184-R185 (2021).
22. R. A. Marks, S. Hotaling, P. B. Frandsen, R. VanBuren, Lessons from 20 years of plant genome sequencing: an unprecedented resource in need of more diverse representation. *bioRxiv* (2021).
23. S. Nurk *et al.*, The complete sequence of a human genome. *bioRxiv* (2021).

24.   E. A. Ellis, C. G. Storer, A. Y. Kawahara, De novo genome assemblies of butterflies.
GigaScience **10**, giab041 (2021).
25.   F. Thibaud-Nissen *et al.*, The NCBI Eukaryotic Genome Annotation Pipeline. *Journal of*
Animal Science **94**, 184-184 (2016).
26.   J. Deck *et al.*, The Genomic Observatories Metadatabase (GeOMe): A new repository
for field and sampling event metadata associated with genetic samples. *PLoS biology*
15, e2002925 (2017).
27.   J. C. Buckner, R. C. Sanders, B. C. Faircloth, P. Chakrabarty, Science Forum: The
critical importance of vouchers in genomics. *Elife* **10**, e68264 (2021).
28.   C. W. Thompson *et al.*, Preserve a voucher specimen! The critical need for integrating
natural history collections in infectious disease studies. *Mbio* **12**, e02698-02620 (2021).
29.   S. Hotaling *et al.*, Species discovery and validation in a cryptic radiation of endangered
primates: coalescent-based species delimitation in Madagascar's mouse lemurs.
Molecular ecology **25**, 2029-2045 (2016).
30.   F. Adame, Meaningful collaborations can end'helicopter research'. *Nature* (2021).
31.   A. Wonkam (2021) Sequence three million genomes across Africa. (Nature Publishing
Group).
32.   R. Dirzo, P. H. Raven, Global state of biodiversity and loss. *Annual review of*
Environment and Resources **28**, 137-167 (2003).
531