Towards a genome sequence for every animal: where are we now?

Scott Hotaling[1], Joanna L. Kelley[1], and Paul B. Frandsen[2,3,4]

Affiliations:
[1] School of Biological Sciences, Washington State University, Pullman, WA, USA
[2] LOEWE Centre for Translational Biodiversity Genomics (LOEWE-TBG), Frankfurt, Germany
[3] Department of Plant and Wildlife Sciences, Brigham Young University, Provo, UT, USA
[4] Data Science Lab, Smithsonian Institution, Washington, DC, USA

Correspondence:
Scott Hotaling, School of Biological Sciences, Washington State University, Pullman, WA, 99164, USA; Email: scott.hotaling@wsu.edu: Phone: (828) 507-9950

Paul B. Frandsen, Department of Plant and Wildlife Sciences, Brigham Young University, Provo, UT, 84602, USA; Email: paul_frandsen@byu.edu; Phone: (804) 422-2283

Keywords: animal genomes, genome biology, Arthropoda, Mammalia, Metazoa, genomic natural history

Abstract:
In less than 25 years, the field of animal genome science has transformed from a discipline seeking its first glimpses into genome sequences across the Tree of Life to a global enterprise with ambitions to sequence genomes for all of Earth's eukaryotic diversity (1). As the field rapidly moves forward, it is important to take stock of the progress that has been made to best inform the discipline's future. In this perspective, we provide a contemporary, quantitative perspective on animal genome sequencing. We identified the best available genome assemblies on GenBank, the world's most extensive genetic database, for 3,278 unique animals across 24 phyla. We assessed taxonomic representation, assembly quality, and annotation status for major clades. We show that while tremendous taxonomic progress has occurred, stark disparities in genomic representation exist, highlighted by a systemic overrepresentation of vertebrates and underrepresentation of arthropods. In terms of assembly quality, long-read sequencing has dramatically improved contiguity and, on average, gene annotations are available for just 34.3% of taxa. Furthermore, we show that animal genome science has diversified in recent years with an ever-expanding pool of researchers participating. However, the field still appears to be dominated by institutions in the Global North, which have been listed as the submitting institution for 77% of all assemblies. We conclude by offering recommendations for how we can collectively improve genomic resource availability and value while also broadening representation worldwide.

Significance statement:
The field of animal genome science is rapidly developing, and efforts are underway to sequence genomes for all of Earth's eukaryotic biodiversity. Here, we provide a contemporary overview of animal genome sequencing, with emphases on taxonomic representation, assembly quality, and geographic representation. We show that while a staggering 3,278 unique animals have had their genomes sequenced, massive disparities exist in terms of the taxonomic groups receiving attention, the quality of the resources being produced, and the institutions driving the field. We highlight areas where improvements can be made, notably by continuing to increase the quality of genome assemblies, including by improving metadata and voucher specimen associations, and actively developing meaningful collaborations between researchers form the Global North and South.

52
Introduction:
The first animal genome sequence was published 23 years ago (2). The 97 million base pair (Mb) *C. elegans* genome assembly ushered in a new era of animal genome biology where genetic patterns and processes could be investigated at genome scales. As genome assemblies have accumulated for an increasingly diverse set of species, so too has our knowledge of how genomes vary and shape Earth's biodiversity (e.g., 3, 4). Major shifts in genome availability and quality have been driven by two key events. First, the invention of high-throughput, short-read sequencing provided an economical means to generate millions of reads for any species from which sufficient DNA could be obtained. These ~100-bp short reads could be assembled into useful, albeit fragmented, genome assemblies. Later, the rise of long-read sequencing allowed for similarly economical generation of reads that are commonly orders of magnitude longer than short reads, resulting in vastly more contiguous genome assemblies (5).

We have now entered an era of genomic natural history. Building on ~250 years of natural history efforts to describe and classify the morphological diversity of life on Earth, we are gaining a complementary genomic perspective of Earth's biodiversity. However, a baseline accounting of our progress towards a complete perspective of Earth's genomic natural history— where every species has a corresponding, reference-quality genome assembly available—has not been presented. This knowledge gap is particularly important given the momentum towards sequencing all animal genomes which is being driven by a host of sequencing consortia. For instance, the Vertebrates Genome Project seeks to generate high-quality assemblies for all vertebrates (6), the Bird10K project seeks to generate assemblies for all extant birds (7), the Earth BioGenome Project (EBP) aims to sequence all eukaryote genomes (1), and the Darwin Tree of Life project plans to sequence all eukaryotes in Britain and Ireland (https://www.darwintreeoflife.org/).

In this perspective, we curated, quantified, and summarized genomic progress for a major component of Earth's biodiversity: kingdom Animalia (Metazoa) and its roughly 1.66 million described species (8). We show that as of June 2021, 3,278 unique animals have had their nuclear genome sequenced and the assembly made publicly available on the National Center for Biotechnology Information (NCBI) GenBank database (9). This translates to 0.2% of all animal species, but when viewed through the lens of major clades, massive disparities exist. For instance, just 0.09% of arthropods have genome sequences available versus 2.7% of chordates, a difference of ~3,200% (Fig. 1).

Methods:
To construct a database of the best available genome assembly for all animals, we downloaded metadata from GenBank for all kingdom Animalia taxa using the "summary genome" function in v.10.9.0 of the NCBI data sets command line tool on 4 February 2021. Next, we used the TaxonKit (10) "lineage" function to retrieve taxonomic information for each taxid included in the genome metadata. To gather additional data for each assembly (e.g., sequencing technology), we used a custom web scraper script. Both this web scraper script and the scripts used to download and organize the metadata are available on this study's GitHub repository (https://github.com/pbfrandsen/metazoa_assemblies). We later supplemented this initial data set with a second round of metadata acquisition on 28 June 2021. For the full data set, we hand-refined the NCBI taxonomy classifications to subdivide our data set into three categories: species, subspecies, or hybrids (Table S1). If replicate assemblies for a taxon were present, we defined the "best available" assembly as the one with the highest contig N50 (the mid-point of the contig distribution where 50% of the genome is assembled into contigs of a given length or longer).

  Figure 1. Variation in taxonomic richness and genome availability, quality, and assembly size
across kingdom Animalia on GenBank (as of 28 June 2021). Taxonomic groups are clustered by
phylogeny following (11). Only groups with 30 or more available assemblies as of January 2021 are
shown with the exception of Hominidae (n = 5 genomes). In the tree, bold group names represent phyla
and naming conventions follow those of the NCBI database. Of 34 recognized animal phyla, 10 do not
have a representative genome sequence. (a) The total number of described species for each group
following Zhang (8) and the references therein. (b) Genomic representation among animal groups for
3,278 species with available genome assemblies. Bars represent the magnitude of the observed minus
the expected number of genomes given the proportion that each group comprises of described animal
diversity. Significance was assessed with Fisher's exact tests and significantly under- or over-represented
groups (P < 0.05) are denoted with asterisks. Gray numbers indicate the total number of unique species
with available genome assemblies for each group. The number of available assemblies is not mutually
exclusive with taxonomy; that is, a carnivore genome assembly would be counted in three categories
(Order Carnivora, Class Mammalia, Phylum Chordata). (c) The percentage of described species within a
group with an available genome sequence (bars) and the percentage of those assemblies that have
corresponding annotations (red circles). For many groups (e.g., arthropods), only a fraction of a percent
of all species have an available genome assembly, making their percentage appear near zero. (d)
Assembly size for all animal genome assemblies, grouped by taxonomy. (e) Contig N50 by taxonomic
group. The sequencing technology used for each assembly is denoted by circle fill color: short-read
(blue), long-read (yellow), or not provided (gray). In (d) and (e), each circle represents one genome
assembly and a few notable or outlier taxa are indicated with gray text.

We culled our data in several ways: we removed subspecies (unless they were the only
representative for a species) and hybrids, assemblies that were shorter than 15.3 Mb [the
smallest assembly size for a metazoan to date, (12)], or had a contig N50 less than 1 kilobase
(Kb). We also culled assemblies that were unusually short (i.e., 1-2.5 Mb) and had information
in their descriptions that indicated they were not true nuclear genome assemblies (e.g., "exon
capture"). In total, we culled 407 assemblies that did not fit the above criteria. The remaining
assemblies were classified as "short-read", "long-read", or "not provided" based on whether only
short-reads (e.g., Illumina) were used, any amount of long-read sequences (e.g., PacBio) were
used, or no information was available. We defined a species as having gene annotations
available if any assembly for that taxon also had annotations on GenBank. When the best
available assembly did not have annotations included or when multiple assemblies had
annotations, we retained the annotations for the assembly with the highest contig N50. Finally,
we used the submitting institution for each assembly as a surrogate for the institution that led
the genome assembly effort for each taxon. Using these data, we classified assemblies to a
country, region (Africa, Asia, Europe, Middle East, North America, Oceania, South America,
Southeast Asia), and the Global North (e.g., Australia, Canada, Europe, USA) or Global South
(e.g., Africa, Asia including China, Mexico, Middle East, South America).

To test if clades were under- or overrepresented in terms of genome availability relative to their
species richness, we compared the observed number of species with assemblies to the
expected total for the group. We obtained totals for the number of described species overall and
for each group from previous studies, primarily from Zhang (8) and the references therein. We
assessed significance between observed and expected representation with Fisher's exact tests
(alpha = 0.05). We tested for differences in distributions of contig N50 or assembly size between
short- and long-read genomes with Welch's T-tests. For both display (i.e., Figure 1) and
analysis, we subdivided the data set into the lowest taxonomic level that still contained 30 or
more assemblies as of January 2021 (with the exception of hominids which were given their
own category due to their exceptionally high genomic resource quality).

Figure 2. Genome availability for kingdom Animalia versus taxonomic descriptions and over time.
(a) The proportion of described taxonomic groups versus the number with sequenced genome
assemblies from phyla to species. The gray plot to the right is a zoomed in perspective of the lower
frequency (i.e., higher taxonomic level) categories in the full plot to the left. For genus-phylum, the
number of described categories is based on the NCBI taxonomy. For species, the total number described
is from Zhang (8). (b) The timeline of genome contiguity versus availability for animals according to the
GenBank publication date (x-axis, see panel c). A sudden rise in assembly contiguity has been
precipitated by long-read sequencing. Particularly contiguous assemblies for a given time period are
labeled. (c) The number of animal genome assemblies deposited on GenBank each month since
February 2004. Several notable events are labeled. When specific dates are indicated, those (and the
assemblies referred to) are included within that month's total. For (b) and (c), it is important to note that
when a genome assembly is updated to a newer version, its associated date is also updated. Thus, the
date associated with many early animal assemblies [e.g., *Caenorhabditis elegans*; (2)] has shifted to be
more recent with updates.
170

Results:
Taxonomic representation
Genome assemblies were available for 3,278 animal species representing 24 phyla, 64 classes,
and 258 orders (Fig. 2a, Table S1). The data set was exceptionally enriched for the phylum
Chordata (which includes all vertebrates) with 1,770 assemblies for the group (54% of all
assemblies) despite chordates comprising just 3.9% of animals (*P*, Fisher's < 1e-5; Fig. 1).
Conversely, arthropods were underrepresented with 1,115 assemblies (34% of the data set) for
a group that comprises 78.5% of animal species (*P*, Fisher's < 1e-5; Fig. 1). However, not all
arthropods were underrepresented; five insect clades were overrepresented [Apidae (bees),
Culicidae (mosquitoes), *Drosophila* (fruit flies), Formicidae (ants), and Lepidoptera (butterflies
and moths); all *P*, Fisher's < 1e-3; Fig. 1]. Collectively, of the 59 taxonomic groups included, 14
groups were underrepresented, 17 were represented as expected, and 28 were
overrepresented (primarily chordates; Fig. 1). Ten phyla had no publicly available genome
sequence (Fig. 1). Over the ~17-year GenBank genome assembly record, animal assemblies
have been deposited at a rate of 0.52 unique species' assemblies per day. Over the most recent

year, however, this rate increased eight-fold to 4.07 assemblies per day. If this most recent rate
was maintained, all currently described animals would have a genome assembly by 3136. To
achieve this goal by 2031 instead, an average of 165,614 animal genomes would need to be
sequenced each year (~112x faster than the historical rate).
190
Assembly size, contiguity, and annotations
The average animal genome assembly was 1.02 gigabases (Gb) in length (SD = 1.21 Gb) with
a contig N50 of 2.26 Mb (SD = 25.16 Mb; Fig. 1d,e). Two animal genome assemblies were 25
Gb longer than all other assemblies—the axolotl [32.4 Gb (13)] and Australian lungfish [34.6 Gb
(14); Fig. 1d]. The smallest genome assembly in the data set, the mite *Aculops lycopersici*, was
over 1000x smaller, spanning just 32.5 Mb (15). Still smaller is the 15.3 Mb assembly of the
marine parasite, *Intoshia variabili*, which has the smallest animal genome currently known (12).
But, since the *I. variabili* assembly was not available on GenBank as of June 2021, it was not
included in our data set.
200
Contiguity varied dramatically across groups. For instance, hominid assemblies (family
Hominidae, $n = 5$) were the most contiguous with an average contig N50 of 24.2 Mb. Bird
assemblies (class Aves, $n = 515$) were also highly contiguous (mean contig N50 = 1.4 Mb)
despite being so numerous (and accumulating for a long period of time). On the other end of the
spectrum, jellyfish and related species (phylum Cnidaria) exhibited some of the least contiguous
genome assemblies with a mean contig N50 of 0.18 Mb (n =65; Fig. 1e). Roughly 34% of
animals with genome assemblies had corresponding annotations on GenBank but annotation
rates differed substantially among groups (Fig. 1c). For example, the rate of arthropod
annotations (22.3%) lags behind chordates (41.3%), however much of this disparity appeared to
be driven by the low and high annotation rates of butterflies and moths (order Lepidoptera) and
birds (class Aves), respectively. Of 445 assemblies, just 6.5% of lepidopteran assemblies on
GenBank have corresponding annotations versus 72.8% of birds ($n = 519$ assemblies; Fig. 1c).
213
Geographic representation
Animal genome assemblies have been contributed by researchers at institutions on every
continent with permanent inhabitants, including 52 countries. From a regional perspective,
institutions in North America ($n = 1,331$), Europe ($n = 972$), and Asia ($n = 828$) collectively
accounted for 95.5% of all assemblies (Fig. 3a). And, nearly 70% of all animal genome
assemblies have been submitted by researchers in just three countries: USA ($n = 1,275$), China
($n = 676$), and Switzerland ($n = 317$; Fig. 3a). When countries were grouped by their inclusion in
the Global North or South, similarly stark patterns emerged. Researchers affiliated with
institutions in the Global North contributed roughly 75% of animal genome assemblies (Fig. 3b).
From a taxonomic perspective, researchers at North American institutions have contributed the
most insect and mammal assemblies, European researchers have contributed the most fish
assemblies, and Asian researchers the most bird assemblies (Fig. 3a). The first assembly in
GenBank from the Global North was deposited in 2004 and the first assembly from the Global
South was deposited in 2011 (Fig. 3c). Since then, the number of assemblies deposited each
year has steadily risen with the proportions from the Global North and South staying relatively
constant (Fig. 3c).
230

Figure 3. Where animal genome assemblies have been produced around the world according to the submitting institutions on GenBank. (a) For each geographic region, the total number of genome assemblies are shown by dark circles with white lettering. This total is further broken down by country and taxon. For regions where more than four countries have contributed assemblies (e.g., Europe), an "Other" category represents all other countries. The same applies to all assemblies that are not insects, birds, fish, or mammals in the taxon plots. Countries are color-coded by assignment to the Global North or South. (b) The total number of genome assemblies contributed by countries in the Global North (e.g., USA, Europe, Australia) versus the Global South (e.g., Africa, South America China, Mexico, Middle East). (c) The rate of genome assembly deposition by major sources in the Global North (Europe, USA) and Global South (China, Southeast Asia) as well as all other countries collectively in each.

Use of long-reads in genome assemblies and availability of key metadata also differs with geography. For assemblies deposited since 2018, researchers from the Global South have used long-reads slightly more frequently than those from the Global North (25.7% versus 20.2%; Fig. 4a). However, researchers from the Global North were far less likely to report the types of sequence data used (19.9% of assemblies for the Global North versus 1.4% of assemblies for the Global South; Fig. 4a). Much of this difference appears to be driven by genome assemblies deposited by researchers at European institutions (Fig. 4b). This gap in metadata may reflect an

issue with data mirroring between the European Nucleotide Archive (ENA) and GenBank. For
instance, many new genome assemblies being generated by the United Kingdom, for example,
are part of the Wellcome Sanger Institute's Darwin Tree of Life project, which is generating
extremely high-quality assemblies using long-read sequencing and were deposited into the ENA
(Fig. 5). One region (Oceania) and three countries (Australia, Finland, India) reported long reads
being used in more than 50% of deposited assemblies (Fig. 4b,c).
257

258
259 **Figure 4. Sequencing technologies used around the world: (a) between the Global North versus**
260 **Global South, (b) among regions, and (c) among countries.** To limit bias due to the limited availability
261 of long-read sequencing technologies before ~2017 (see Fig. 2b), only assemblies deposited on or after 1
262 January 2018 were included in the analysis and in (c) only countries that deposited five or more
263 assemblies during the focal period (January 2018 – June 2021) are shown.
264
Discussion:
Taxonomic representation
Animal genome sequencing has dramatically progressed in the last 25 years. In that span, the
field has moved from sequencing the first nuclear genome for any animal (2)—a landmark
achievement—to targeting the generation of genome assemblies for *all* of Earth's eukaryotic
biodiversity (1). Here, we provided a contemporary perspective on progress towards this goal
for the 1.6-million species in the animal kingdom (8). We showed that while tremendous
progress has been made, major gaps and biases remain both in terms of taxonomic and
geographic representation, at least within the most commonly used database of genomic
resources, GenBank. For instance, a major bias exists in favor of vertebrates which are vastly

overrepresented relative to their total species diversity (Fig. 1a-c). From the perspectives of
biomedicine and human evolution, this bias is reasonable since vertebrates are the closest
relatives of humans. However, from a basic research perspective, particularly as it relates to
genomic natural history and an overarching goal to sequence all animal genomes, there is a
need to taxonomically diversify sequencing efforts.
280
At the highest taxonomic levels, 10 animal phyla still have no genomic representation. To
illustrate the scale of this disparity versus other groups and the unique biology that is being
overlooked, genome assemblies are available for 685 ray-finned fishes (class Actinopterygii) but
none exists for phylum Nematomorpha, a ~2,000-species clade of parasitic worms whose
presence can dramatically alter energy budgets of entire stream ecosystems (16). Another
phylum without genomic representation—Loricifera—was first described in 1983 (17). This
group of small, sediment-dwelling animals includes the only examples of multicellular species
spending their entire life cycles under permanently anoxic conditions (18). Loriciferans
accomplish this feat by foregoing the energy-producing mitochondria found in virtually all
animals in favor of hydrogenosome-like organelles akin to those found in prokaryotes inhabiting
anaerobic habitats (18). Clearly, there is much to discover in terms of genomic diversity in
undersampled clades.
293
Global representation
A few select countries—primarily the USA, several European nations, and China—have led the
sequencing of the vast majority of animal genome assemblies (Fig. 3a). Aside from China, all of
these countries are within the Global North. This pattern of geographic bias raises two potential
issues for representation in animal genome science. First, the researcher population of animal
genome sequencing likely does not reflect the global population. Second, sampling biases may
exist towards the regions where most of the genome sequencing is occurring. Some of this
"bias" is intentional and reflects funding goals for a given region. For instance, the Darwin Tree
of Life project (https://www.darwintreeoflife.org/) seeks to sequence the genomes of all ~70,000
eukaryotic species living in Britain and Ireland. Still, however, similar to how sampling biases
can yield skewed understanding of the natural world in other disciplines (e.g., 19), so too could
bias towards specific ecoregions, habitats, or other classifications skew genomic insight.
306
Inherently linked with questions of representation in animal genome science is the specter of
parachute science (or helicopter research)—the practice where international scientists, typically
from wealthy nations, conduct studies in other countries that are often poorer without
meaningful communication nor collaborations with local people (20). Parachute science has a
long history in ecological research and signatures of these practices have been observed for
genome sciences. For instance, Marks *et al.* (21) found that the majority of plant genome
assemblies for species that are native to South America and Africa were sequenced "off-
continent" by researchers at European, North American, or Asian institutions. Given the sheer
number of animal genome assemblies that have been submitted by a small number of countries
and institutions, a similar pattern may exist for animal genomics. However, to properly assess
this issue, parsing authorship to quantify collaboration, at a minimum, would need to occur and
this approach would still overlook key aspects of representation that need to be considered
(e.g., if a researcher from the Global South is working at an institution in the Global North).
320

Figure 5. Examples of major contributors of genome assemblies for (a) butterflies (order Lepidoptera), (b) birds (class Aves), and fish (primarily class Actinopterygii). Major contributors were defined as any consortium, organization, or project that has deposited more than 5% of all assemblies for butterflies and birds or 2.5% of all assemblies for fish.

Assembly quality varies dramatically within and among clades
For the purpose of biological discovery, not all genome assemblies are created equal. As long-read sequencing technologies have matured, so too have the quality of assemblies being generated (5). In the last year alone, the largest ever animal genome assembly was deposited [Australian lungfish, (14)] as well as the most complete human genome to date, a telomere-to-telomere assembly (22). Still, many species on GenBank only have low-quality assemblies available (i.e., contig N50 < 100 Kb with no corresponding gene annotations; Fig. 1). Fragmentation and/or poor or missing gene annotations reduce the value of any assembly. Thus, genome assembly quality is an important consideration, particularly when the end goal is resource development for a broader community. As of April 2021, the EBP sought assembly quality of 6.C.Q40 (https://www.earthbiogenome.org/assembly-standards) for reference genomes where "6" refers to a 1e6 contig N50 (i.e., 1 Mb). In our data set, 568 assemblies (17.3%) reach this contiguity standard. And, that number drops to 271 assemblies (8.3%) when contig N50 ≥ 1 Mb and deposited gene annotations are both required. For reference, the "C" above refers to chromosomal scale scaffolding and "Q40" to a less than 1/10,000 error rate. Neither of these metrics were assessed in this study.

Independent research laboratories, institutions, and consortia have contributed genome assemblies on both ends of the quality spectrum (Fig. 5). For example, among butterflies (order Lepidoptera), a bimodal quality distribution is being primarily driven by contributions made in 2021 by two submitting institutions, the Florida Museum of Natural History (e.g., 23) and the Wellcome Sanger Institute (Fig. 5a). When viewing genome assembly contributions holistically across the animal Tree of Life, it is clear that two consortia—the Vertebrate Genomes Project (6) and the Darwin Tree of Life (https://www.darwintreeoflife.org/), part of the Wellcome Sanger Institute—warrant specific recognition for contributing exceptional genomic resources relative to closely related species (Fig. 5a-c).

354
Going forward
While animal genome science has dramatically matured in recent years, the field still rests on
the cusp of massive change. Thousands of genome assemblies are now available for a wide
range of taxa, a resource that can empower unprecedented scales of genomic comparison.
Simultaneously, multiple consortia are building momentum towards their goals and generating
some of the highest quality genome assemblies ever produced. The field is also diversifying
with researchers around the world, particularly from the Global South, leading a rising number of
efforts. These ongoing advances will yield higher quality, more globally representative genomic
data for animals. As we collectively build towards this new genomic future, we offer
recommendations to improve assembly quality and accessibility while also continuing to
increase representation within the discipline.
366
The quality of a genome assembly is likely the most important factor dictating its long-term
value. Genome assembly "quality," however, is difficult to define. Here, we propose a holistic
view on genome assembly quality that generally echoes the guidelines proposed by the EBP
and other consortia. Briefly, assemblies should reach minimum levels of contiguity (e.g., contig
N50 > 1 Mb) and accuracy in order to be considered a reference that will likely not need to be
updated for most applications. At a minimum, assemblies should also include high-quality gene
annotations that perhaps take advantage of standardized pipelines [e.g., NCBI Eukaryotic
Genome Annotation Pipeline, (24)] to maximize compatibility across taxa. We recommend the
field further improve the quality of genome assembly resources in two ways. First, refining and
expanding the coordinated deposition of genome assemblies will improve the usability of the
resources, reproducibility of analyses, and will reduce duplications of effort—i.e., when a group
sequences a genome that has already been produced—an issue that is likely to become
increasingly common.
380
To refine and expand coordinated resource deposition, we recommend the continued use of
GenBank (9) or one of the other archives that are members of the International Nucleotide
Sequence Database Collaboration (INSDC)—the European Nucleotide Archive (ENA) and DNA
Database of Japan (DDBJ)—as the central repositories for genome assemblies and their
metadata given their tripartite data sharing agreement. Next, we call on genetic archive
administrators, consortia, and independent researchers to collectively improve the metadata
submitted with each assembly and the mirroring of data across repositories. Too many
assemblies lack basic information about the sequence data and methods used (e.g., Fig. 4) and
with the difficulty of linking assemblies to published studies (if available), it can be difficult to find
this information. Further, an expansion of the metadata associated with each assembly—ideally
to make more of the categories required and expand demographic data—would make efforts to
quantify geographic representation, for instance, far more straightforward. In the same vein, a
set of minimum quality characteristics for a genome assembly may need to be defined. A
number of exceptionally low-quality "genome assemblies" (e.g., with contig N50 values shorter
than 1 Kb) that often cover only a small fraction of the expected total genome sequence length
for a given group are present on GenBank. The presence of these assemblies raise the
question: where is the inflection point between resource quality and value to other researchers
versus diluting the resources of a shared repository?
399
400 For our second recommendation, we amplify and expand the message of Buckner et al. (25):
401 genome science needs specimen vouchers. Vouchers serve as a key physical link between
402 taxonomy and molecular insight. Rarely, however, are vouchers referenced in publications of
403 genome assemblies; only 11% of vertebrate assemblies included such a reference as of
404 January 2020 (25). While vouchers represent a physical reference for assessing taxonomic

classification or morphological variation, a properly stored voucher could also provide a long-term source of material for future resource improvement. If a physical specimen cannot be deposited, photographs and/or genomic DNA should be deposited in its place (e.g., 26). Tied to the metadata discussion above, additional fields should be added to GenBank genome assembly accessions to directly link the assembly to a specimen, photos, or genomic DNA that has been deposited elsewhere.

Though geographic representation in animal genome science has improved in recent years, the discipline appears far from properly reflecting the global researcher pool. This issue is almost certainly multi-faceted, likely stemming from a lack of infrastructure (e.g., fewer high-throughput sequencing platforms in developing countries), fewer resources for expensive molecular research, and a corresponding lack of training in genome analysis. To bridge this gap and to empower a more diverse discipline, the nations and institutions that are devoting large amounts of resources to animal genome sequencing (e.g., China, United Kingdom, USA), and the researchers within those countries, should develop meaningful collaborations with researchers within the countries where their focal species reside (27). These meaningful collaborations—where all parties are valued for their expertise and involved in decision-making—improve the science through transfer of local knowledge, provide a means for local researchers to expand their skillset and network while raising their scholarly profile, and most importantly, can effectively end the practice of parachute science (27). Within-continent (or country) initiatives also have transformative potential for people and genome research. For instance, the African-led effort to sequence three million African genomes over the next 10 years (the "3MAG" project) will yield massive investment in African genomics, an incredible resource for understanding the full scope of human genetic diversity, and a new generation of African genome scientists (28). While focused on human genetics, the infrastructure and expertise that stems from the 3MAG project will no doubt translate to other taxa in the coming years.

Conclusion:
Animal genome science continues to grow and expand at an exceptional rate. The coming years will surely see thousands, and perhaps tens of thousands, of new genome assemblies from across the Tree of Life, massive technological and analytical improvements, and some of the largest scale and most in-depth studies of animal genome biology conducted to date. However, if we are to realize the ambitious goals of efforts like the Earth BioGenome Project (EBP)—a self-described biological "moonshot"—the rate and mean quality of animal genome assembly production will have to increase by roughly two orders of magnitude. Regardless of rates and timelines, however, perhaps the most important goal for the future of animal genome science is that we empower a more diverse, representative researcher community in parallel with these genomic resources.

Acknowledgements:
S.H. and J.L.K. were supported by NSF award OPP-1906015. We would like to thank Guangfeng Song, Eric Cox, and Anne Ketter from the "datasets" development team at the NCBI for their responsiveness and receptiveness to improving this valuable tool for data science.

References:
1.   H. A. Lewin *et al.*, Earth BioGenome Project: Sequencing life for the future of life. *Proceedings of the National Academy of Sciences* **115**, 4325-4333 (2018).
2.   C. elegans Sequencing Consortium, Genome sequence of the nematode C. elegans: a platform for investigating biology. *Science* **282**, 2012-2018 (1998).
3.   C. Hoencamp *et al.*, 3D genomics across the tree of life reveals condensin II as a determinant of architecture type. *Science* **372**, 984-989 (2021).

4. G. W. Thomas *et al.*, Gene content evolution in the arthropods. *Genome Biology* **21**, 1-
14 (2020).
5. S. Hotaling *et al.*, Long-reads are revolutionizing 20 years of insect genome sequencing.
Genome Biology and Evolution (2021).
6. A. Rhie *et al.*, Towards complete and error-free genome assemblies of all vertebrate
species. *Nature* **592**, 737-746 (2021).
7. G. Zhang, Bird sequencing project takes off. *Nature* **522**, 34-34 (2015).
8. Z.-Q. Zhang, *Animal biodiversity: An outline of higher-level classification and survey of*
taxonomic richness (Magnolia press, 2011).
9. E. W. Sayers *et al.*, GenBank. *Nucleic Acids Research* **48**, D84-D86 (2021).
10. W. Shen, J. Xiong, TaxonKit: a cross-platform and efficient NCBI taxonomy toolkit.
Biorxiv, 513523 (2019).
11. C. E. Laumer *et al.*, Revisiting metazoan phylogeny with genomic sampling of all phyla.
Proceedings of the royal society B **286**, 20190831 (2019).
12. G. S. Slyusarev, V. V. Starunov, A. S. Bondarenko, N. A. Zorina, N. I. Bondarenko,
Extreme genome and nervous system streamlining in the invertebrate parasite Intoshia
variabili. *Current Biology* **30**, 1292-1298. e1293 (2020).
13. S. Nowoshilow *et al.*, The axolotl genome and the evolution of key tissue formation
regulators. *Nature* **554**, 50-55 (2018).
14. A. Meyer *et al.*, Giant lungfish genome elucidates the conquest of land by vertebrates.
Nature **590**, 284-289 (2021).
15. R. Greenhalgh *et al.*, Genome streamlining in a minute herbivore that manipulates its
host plant. *Elife* **9**, e56689 (2020).
16. T. Sato *et al.*, Nematomorph parasites drive energy flow through a riparian ecosystem.
Ecology **92**, 201-207 (2011).
17. R. M. Kristensen, An introduction to loricifera, cycliophora, and micrognathozoa.
Integrative and Comparative Biology **42**, 641-651 (2002).
18. R. Danovaro *et al.*, The first metazoa living in permanently anoxic conditions. *BMC*
biology **8**, 1-10 (2010).
19. A. C. Hughes *et al.*, Sampling biases shape our view of the natural world. *Ecography*
(2021).
20. P. V. Stefanoudis *et al.*, Turning the tide of parachute science. *Current Biology* **31**,
R184-R185 (2021).
21. R. A. Marks, S. Hotaling, P. B. Frandsen, R. VanBuren, Lessons from 20 years of plant
genome sequencing: an unprecedented resource in need of more diverse
representation. *bioRxiv* (2021).
22. S. Nurk *et al.*, The complete sequence of a human genome. *bioRxiv* (2021).
23. E. A. Ellis, C. G. Storer, A. Y. Kawahara, De novo genome assemblies of butterflies.
GigaScience **10**, giab041 (2021).
24. F. Thibaud-Nissen *et al.*, The NCBI Eukaryotic Genome Annotation Pipeline. *Journal of*
Animal Science **94**, 184-184 (2016).
25. J. C. Buckner, R. C. Sanders, B. C. Faircloth, P. Chakrabarty, Science Forum: The
critical importance of vouchers in genomics. *Elife* **10**, e68264 (2021).
26. S. Hotaling *et al.*, Species discovery and validation in a cryptic radiation of endangered
primates: coalescent-based species delimitation in Madagascar's mouse lemurs.
Molecular ecology **25**, 2029-2045 (2016).
27. F. Adame, Meaningful collaborations can end'helicopter research'. *Nature* (2021).
28. A. Wonkam (2021) Sequence three million genomes across Africa. (Nature Publishing
Group).
505