

1 **Opposing roles of p38 $\alpha$ -mediated phosphorylation and arginine methylation in driving TDP-**  
2 **43 proteinopathy.**

3  
4 Mari Aikio<sup>1</sup>, Heike J. Wobst<sup>2§</sup>, Hana M. Odeh<sup>3</sup>, Bo Lim Lee<sup>3</sup>, Bradley Class<sup>2</sup>, Thomas A.  
5 Ollerhead<sup>1</sup>, Korrie L. Mack<sup>3,4</sup>, Alice F. Ford<sup>3,5</sup>, Edward M. Barbieri<sup>3</sup>, Ryan R. Cupo<sup>3,6</sup>, Lauren E.  
6 Drake<sup>3</sup>, Nicholas Castello<sup>7</sup>, Ashmita Baral<sup>7</sup>, John Dunlop<sup>2</sup>, Aaron D. Gitler<sup>8</sup>, Ashkan Javaherian<sup>7</sup>,  
7 Steven Finkbeiner<sup>7,9</sup>, Dean G. Brown<sup>10§</sup>, Stephen J. Moss<sup>1\*</sup>, Nicholas J. Brandon<sup>1,2\*</sup>, and James  
8 Shorter<sup>3,4,5,6\*</sup>.

9  
10 <sup>1</sup>AstraZeneca-Tufts Laboratory for Basic and Translational Neuroscience, Tufts University, Boston,  
11 MA, U.S.A.

12 <sup>2</sup>Neuroscience, BioPharmaceuticals R&D, AstraZeneca, Boston, MA, U.S.A.

13 <sup>3</sup>Department of Biochemistry and Biophysics, <sup>4</sup>Biochemistry and Molecular Biophysics Graduate  
14 Group, <sup>5</sup>Neuroscience Graduate Group, <sup>6</sup>Pharmacology Graduate Group, Perelman School of  
15 Medicine, University of Pennsylvania, Philadelphia, PA, U.S.A.

16 <sup>7</sup>Center for Systems and Therapeutics, Taube/Koret Center for Neurodegenerative Disease  
17 Research, Gladstone Institutes, San Francisco, CA, U.S.A.

18 <sup>8</sup>Department of Genetics, Stanford University School of Medicine, Stanford, CA, U.S.A.

19 <sup>9</sup>Departments of Neurology and Physiology, University of California, San Francisco, CA, U.S.A.

20 <sup>10</sup>Hit Discovery, Discovery Sciences, BioPharmaceuticals R&D, AstraZeneca, Boston, MA, U.S.A.

21 <sup>§</sup>Current address: Jnana Therapeutics, 6 Tide St., Boston, MA, U.S.A.

22 **Running Title: p38 $\alpha$  promotes TDP-43 proteinopathy**

23  
24 \*To whom correspondence should be addressed: [jshorter@pennmedicine.upenn.edu](mailto:jshorter@pennmedicine.upenn.edu),

25 [nick.j.brandon@gmail.com](mailto:nick.j.brandon@gmail.com), [stephen.moss@tufts.edu](mailto:stephen.moss@tufts.edu)

26 **Keywords:** ALS, TDP-43, p38 $\alpha$ , phosphorylation, PRMT1, arginine methylation

27 **Abstract**

28 Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disorder typically characterized by  
29 insoluble inclusions of hyperphosphorylated TDP-43. The mechanisms underlying toxic TDP-43  
30 accumulation are not understood. Persistent activation of p38 mitogen-activated protein kinase  
31 (MAPK) is implicated in ALS. However, it is unclear how p38 MAPK affects TDP-43 proteinopathy.  
32 Here, we demonstrate that inhibition of p38 $\alpha$  MAPK reduces pathological TDP-43 phosphorylation,  
33 aggregation, cytoplasmic mislocalization, and neurotoxicity. We establish that p38 $\alpha$  MAPK  
34 phosphorylates TDP-43 at pathological serine 409/410 (S409/S410) and serine 292 (S292), which  
35 reduces TDP-43 liquid-liquid phase separation (LLPS) but allows pathological TDP-43 aggregation.  
36 Moreover, we show that protein arginine methyltransferase 1 methylates TDP-43 at R293.  
37 Importantly, S292 phosphorylation reduces R293 methylation, and R293 methylation reduces  
38 S409/S410 phosphorylation. R293 methylation permits TDP-43 LLPS and reduces pathological  
39 TDP-43 aggregation. Thus, strategies to reduce p38 $\alpha$ -mediated TDP-43 phosphorylation and promote  
40 R293 methylation could have therapeutic utility for ALS and related TDP-43 proteinopathies.

41

42

## 43 **Introduction**

44 Amyotrophic lateral sclerosis (ALS) is a fatal disorder caused by degeneration of motor neurons  
45 (Wobst et al., 2020). While most ALS cases (~90–95%) are considered sporadic with unknown  
46 etiology (sALS), ~5–10% of cases are familial in nature (fALS), exhibiting a dominant pattern of  
47 inheritance (Rowland and Shneider, 2001; Valdmanis and Rouleau, 2008). ALS is linked to mutations  
48 in more than 25 genes, with the C9ORF72 hexanucleotide repeat expansion and mutations in the  
49 copper–zinc superoxide dismutase (SOD1) being the most common genetic causes (Nguyen et al.,  
50 2018). Although mutations in Transactive Response DNA binding protein 43 kDa (*TARDBP*), the  
51 gene encoding TDP-43, are a rare cause of ALS, ~97% of ALS cases and ~50% of patients with  
52 frontotemporal dementia (FTD) present with TDP-43 proteinopathy characterized by TDP-43-  
53 positive insoluble nuclear and cytoplasmic inclusions in affected neurons (Arai et al., 2006; Guo and  
54 Shorter, 2017; Neumann et al., 2006).

55  
56 TDP-43 is an essential, highly conserved, ubiquitously expressed, and predominantly nuclear protein  
57 with RNA/DNA-binding properties (Portz et al., 2021). It acts as a transcriptional repressor and is  
58 implicated in RNA transport and stability, alternative splicing, microRNA biogenesis, and formation  
59 of stress granules (SGs) (Alami et al., 2014; Aulas et al., 2012; Buratti and Baralle, 2008; Kawahara  
60 and Mieda-Sato, 2012; Khalfallah et al., 2018; Lalmansingh et al., 2011; Li et al., 2018; McDonald  
61 et al., 2011; Tollervey et al., 2011). TDP-43 is comprised of an N-terminal domain involved in  
62 dimerization and recruitment of other RNA-binding proteins, a bipartite nuclear localization sequence  
63 (NLS), two RNA-recognition motifs (RRM1 and RRM2) and a C-terminal glycine-rich, low  
64 complexity prion-like domain (PrLD) that mediates protein–protein interactions and formation of  
65 membraneless organelles, such as SGs, through liquid–liquid phase separation (LLPS) (Figure 1A)  
66 (Ayala et al., 2008; Buratti et al., 2005; Chang et al., 2012; Freibaum et al., 2010; Jiang et al., 2017;  
67 Johnson et al., 2009; Li et al., 2018; Molliex et al., 2015; Winton et al., 2008). The majority of ALS-

68 linked TDP-43 mutations reside in the PrLD, which can enhance aggregation propensity and also  
69 indicates that this region of TDP-43 may be prone to pathological protein modifications (Figure 1A)  
70 (Harrison and Shorter, 2017; Johnson et al., 2009; Prasad et al., 2019).

71  
72 Phosphorylation at serine residues 409/410 (S409/S410) is one of the major pathological markers for  
73 TDP-43 inclusions in human brains (Hasegawa et al., 2008; Neumann et al., 2009). In addition to  
74 phosphorylation, TDP-43 also undergoes other post-translational modifications (PTMs) in ALS  
75 patients and disease-mimicking models, including ubiquitination, generation of C-terminal domain  
76 fragments (CTFs), cysteine oxidation, sumoylation, and acetylation (Buratti, 2018). However, the  
77 functional and pathological significance of TDP-43 PTMs remains unknown. Thus, a clear  
78 understanding of the effects of phosphorylation and other PTMs on solubility, localization, and  
79 aggregation propensity of TDP-43 will enable new insights into the mechanisms of ALS  
80 pathogenesis.

81  
82 The mitogen-activated protein kinase (MAPK) signaling pathway plays a key role in the regulation  
83 of cellular differentiation, motility, growth, and survival (Brennan et al., 2021). MAPKs, which  
84 include extracellular-signal-regulated kinases (ERK), Jun amino-terminal kinases (JNK) and p38  
85 MAPKs, are activated in response to cytokines, growth factors and various stressors including  
86 oxidative stress and endoplasmic reticulum stress (Brennan et al., 2021; Cuadrado and Nebreda, 2010;  
87 Morrison, 2012). In mammals, p38 MAPKs comprise four isoforms ( $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$ ) (Zarubin and Han,  
88 2005). Of these,  $\alpha$  and  $\beta$  are expressed in most tissues, including the brain (Yasuda et al., 2011). p38 $\gamma$   
89 is most highly expressed in skeletal muscle, and p38 $\delta$  in testis, pancreas, kidney and small intestine  
90 (Cuenda et al., 1997). Aberrant p38 signaling has been linked to several neurodegenerative diseases,  
91 including ALS (Burton et al., 2021). Specifically, p38 MAPK inhibition can reduce motor neuron  
92 apoptosis and restore the physiological rate of axonal retrograde transport in SOD1-ALS models

93 (Dewil et al., 2007; Gibbs et al., 2018; Pickhardt et al., 2019). Furthermore, both genetic and  
94 pharmacological inhibition of p38 $\beta$  in a *Drosophila* model of TDP-43 toxicity rescued premature  
95 lethality (Zhan et al., 2015).

96

97 In this study, we establish a role for p38 $\alpha$  MAPK in promoting TDP-43 proteinopathy. We show that  
98 inhibition of p38 $\alpha$  MAPK reduces ALS-associated TDP-43 phenotypes, including TDP-43  
99 aggregation, S409/S410 phosphorylation, cytoplasmic accumulation, and toxicity. Furthermore, *in*  
100 *vitro* kinase assays combined with mass spectrometry revealed that p38 $\alpha$  directly phosphorylates  
101 TDP-43 at S292 and S409/S410. Subsequent cellular experiments identified S292, a residue  
102 previously found to be altered in a genetic risk allele for ALS (S292N) (Xiong et al., 2010; Zou et al.,  
103 2012), as an important site in regulating TDP-43 aggregation. We found that phosphorylation at S292  
104 induced phosphorylation at S409/S410 and promoted TDP-43 aggregation. Moreover, we found that  
105 the residue R293, adjacent to S292, is methylated by protein arginine methyltransferase 1 (PRMT1).  
106 Interestingly, loss or gain of function for other PRMTs is implicated in the pathogenesis of  
107 neurodegenerative diseases (Ratovitski et al., 2015; Simandi et al., 2018). We found that  
108 phosphorylation at S292 inhibited R293 methylation, whereas R293 methylation reduced  
109 phosphorylation at S409/S410, suggesting an interplay between phosphorylation at S292 and  
110 S409/S410 and methylation at R293. Biochemical studies indicated that S292 and S409/S410  
111 phosphorylation reduce TDP-43 LLPS but allow TDP-43 aggregation. By contrast, R293 methylation  
112 allows TDP-43 LLPS but reduces TDP-43 aggregation. Taken together, our results reveal additional  
113 regulatory mechanisms in TDP-43 homeostasis mediated by p38 $\alpha$  and PRMT1. We suggest that  
114 strategies aimed at reducing p38 $\alpha$ -mediated TDP-43 phosphorylation and promoting R293  
115 methylation could have therapeutic utility for ALS and related TDP-43 proteinopathies.

116

117

118 **Materials and methods**

119 **Plasmids**


120 cDNA sequences were based on the accession number NM\_007375.3 for human TARDBP,  
121 NM\_001315 for human p38 $\alpha$  (MAPK14) and NM\_001536 for human PRMT1. Plasmids harboring  
122 N-terminally myc-tagged wild-type or M337V-mutant TDP-43 sequences were generated as  
123 described (Wobst et al., 2017). Plasmids harboring C-terminally Flag-tagged wild-type or mutant  
124 human TDP-43 and p38 $\alpha$  sequences in the pcDNA3.1+/C-(K)-DYK mammalian expression vector  
125 were purchased from Genscript and PRMT1 plasmid was purchased from Origene. TDP-43 bacterial  
126 expression vector harboring a C-terminal MBP tag (pJ4M TDP-43-TEV-MBP-6xHis) was purchased  
127 from Addgene (Plasmid # 104480). TDP-43 mutations were generated by site-directed mutagenesis  
128 using QuikChange (Agilent) and confirmed by DNA sequencing. The presence of the S409:S410E,  
129 S409:S410A, S292E, S292N, S292A, S292:S409:S410A, S292:S409:S410E, and R293F mutations  
130 in TDP-43, as well as dominant negative (DN) T180A/Y182F (Winzen et al., 1999) and constitutively  
131 active (CA) D176A/F327S mutations (Diskin et al., 2004) in p38 $\alpha$  were confirmed by DNA  
132 sequencing performed by Tufts University Core Facilities (forward sequencing primer 5'-TAA-TAC-  
133 GAC-TCA-CTA-TAG-GG-3', reverse sequencing primer 5'-CAG-GAA-ACA-GCT-ATG-AC-3')  
134 and Eurofins Genomics (TDP-43 sequencing primer 1 5'-TAA-TAC-GAC-TCA-CTA-TAG-GGG-  
135 AAT-TG-3', sequencing primer 2 5'-CGG-TGA-GGT-GCT-GAT-GGT-CC-3', sequencing primer  
136 3 5'-GGC-TTT-GGC-AAT-TCG-CGT-GG-3')

137

138

139 **Synthesis of 6-(2,4-difluorophenoxy)-8-ethyl-2-(2-isopropoxyethylamino)pyrido[2,3-**  
140 **d]pyrimidin-7-one (compound 1)**

141



142

143

144 A stirred solution of 2-isopropoxyethan-1-amine (406 mg, 3.93 mmol) was supplemented with 6-  
145 (2,4-difluorophenoxy)-8-ethyl-2-(methylsulfonyl)pyrido[2,3-d]pyrimidin-7(8H)-one (300 mg, 0.79  
146 mmol) in DMF (3 mL). The resulting solution was stirred at 80 °C for 1 hour in a microwave reactor.  
147 The crude product was purified by preparative HPLC (Column: Xselect CSH OBD Column  
148 30\*150mm 5 μm n; Mobile Phase A: Water (0.1%FA), Mobile Phase B: ACN; Flow rate: 60 mL/min;  
149 Gradient: 40% B to 72% B in 8 min; 254/220 nm; Rt: 7.57 min). Fractions containing the desired  
150 compound were evaporated to dryness to afford 6-(2,4-difluorophenoxy)-8-ethyl-2-((2-  
151 isopropoxyethyl)amino)pyrido[2,3-d]pyrimidin-7(8H)-one (190 mg, 59.7 %) as a white solid. <sup>1</sup>H  
152 NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 8.56 (s, 1H), 7.76 (m, 1H), 7.52 - 7.41 (m, 2H), 7.26-7.18 (m, 1H),  
153 7.06-7.02 (m, 1H), 4.42-4.32 (m, 2H), 3.65-3.45 (m, 5H), 1.28-1.22 (m, 3H), 1.09-1.02 (m, 6H). *m/z*  
154 (ES<sup>+</sup>), [M+H]<sup>+</sup> = 405; TFA, HPLC t<sub>R</sub> = 1.822 min. The intermediate 6-(2,4-difluorophenoxy)-8-  
155 ethyl-2-(methylsulfonyl)pyrido[2,3-d]pyrimidin-7(8H)-one was prepared as described (Goldstein et  
156 al., 2011). Compound 1 was tested in a Z'-Lyte kinase assay for p38α (Thermo Fisher) and inhibited  
157 p38α activity with an IC<sub>50</sub> of 25nM.

158

159

160 **Cell culture, transfections and inhibition of p38 $\alpha$  MAPK and methyltransferase activity**

161 SH-SY5Y cells were cultured in minimum essential medium (MEM; Thermo Fisher Scientific)  
162 supplemented with L-glutamine, 10% fetal bovine serum and 1% penicillin/streptomycin in a  
163 humidified incubator at 37°C and 5% CO<sub>2</sub>. For immunocytochemical analysis, cells were grown on  
164 glass coverslips coated with 1 mg/mL poly-L-lysine (Sigma) in 24-well plates (Cellstar). Cells were  
165 transiently transfected using Fugene HD (Promega) following the manufacturer's instructions 24  
166 hours after seeding (Fugene HD:DNA ratio 3:1). Total transfection times were 8 hours - 48 hours.  
167 For pharmacological inhibition of p38 $\alpha$  MAPK, compound 1 (synthesized by AstraZeneca) or equal  
168 volume of DMSO vehicle, was added to cells for 24 or 48 hours at a final concentration of 0.1  $\mu$ M, 1  
169  $\mu$ M or 10  $\mu$ M. For pharmacological inhibition of arginine methyltransferase activity, adenosine-2',3'-  
170 dialdehyde (AdOx) (Sigma) or equal volume of DMSO vehicle was added to cells for 24 hours at a  
171 final concentration of 20  $\mu$ M.

172

173 **Antibodies**

174 The following antibodies were used for immunocytochemical staining: rabbit N-terminal TDP-43  
175 (1:500; 10782-2-AP, Proteintech), mouse anti FLAG-tag (1:500; A00187, GenScript). Anti-rabbit  
176 and anti-mouse secondary antibodies coupled to Alexa-488, and Alexa-647 were used for detection  
177 (1:1,000; Thermo Fisher Scientific). For western blot analysis, the following antibodies were used:  
178 rabbit N-terminal TDP-43 (1:7,000; 10782-2-AP, Proteintech), mouse S409/S410 phospho-TDP-43  
179 (1:3,000; CAC-TIP-PTD-M01, Cosmo), rabbit p38 MAPK (1:1,500; 9212S, Cell Signaling  
180 Technology), rabbit COX IV (3E11) (1:1,500; 4850, Cell Signaling Technology), mouse Histone H3  
181 (96C10) (1:1,500; 3638, Cell Signaling Technology), rabbit Mono-Methyl Arginine (R\*GG)  
182 (D5A12) (1:1,500; 8711S, Cell Signaling Technology), rabbit Asymmetric Di-Methyl Arginine Motif  
183 [adme-R] MultiMab (1:1,500; 13522S, Cell Signaling Technology), rabbit Symmetric Di-Methyl  
184 Arginine Motif [sdme-RG] MultiMab (1:1,500; 13222S, Cell Signaling Technology), rabbit PRMT1



185 (A33) (1:1,500; 2449, Cell Signaling Technology), mouse GAPDH (1:10,000; 60004-1-Ig,  
186 Proteintech). Anti-mouse and anti-rabbit horseradish peroxidase-coupled secondary antibodies were  
187 purchased from Jackson ImmunoResearch (1:10,000).

188

### 189 **siRNA knockdown of p38 $\alpha$ and PRMT1**

190 Small interfering RNAs (siRNA) were obtained from Thermo Fisher Scientific: p38 $\alpha$ : s3585 and  
191 s3586, PRMT1: s6917, s6919, Negative control: 4390846, 4390843. Reverse transfection was  
192 performed with Lipofectamine RNAiMAX-reagent (Invitrogen) following the manufacturer's  
193 instructions. In brief, 25 pmol of siRNA were mixed with 5  $\mu$ L of Lipofectamine RNAiMAX in 500  
194  $\mu$ L of Opti-MEM (Invitrogen) in a 6-well plate. The mixture was incubated for 20 min at room  
195 temperature and 3-3.5  $\times 10^5$  cells were added to mixture. Total knockdown times were 8 - 72 hours.

196

### 197 **Sequential extraction of insoluble protein aggregates**

198 Extraction of insoluble proteins was performed as previously described (Wobst et al., 2017). Briefly,  
199 transfected cells were lysed in 300  $\mu$ L radio-immunoprecipitation assay (RIPA) buffer [50 mM Tris-  
200 HCl, 150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, and 0.1% SDS] (Boston bioproducts)  
201 supplemented with 2 mM EDTA, protease inhibitors (cOmplete, Roche), and phosphatase inhibitors  
202 (PhosStop, Roche). Lysates were sonicated 2 x 15 s with 20% maximum amplitude and centrifuged  
203 for 30 min at 100,000  $\times g$  and 4°C. The supernatant was collected as the RIPA-soluble fraction. The  
204 pellet was washed in RIPA buffer and centrifuged as above. The supernatant was discarded and the  
205 urea-soluble fraction was generated by resuspending the pellet in 100  $\mu$ L urea buffer [7 M urea, 2 M  
206 thiourea, 30 mM Tris pH 8.5, 4% 3-[(3-Cholamidopropyl) dimethylammonio]-1-propanesulfonate  
207 (CHAPS, Sigma)], and sonicating the samples as above followed by centrifugation at room  
208 temperature for 30 min at 100,000  $\times g$ . The supernatant was collected as the urea-soluble fraction.

209

## 210 **Immunoblotting**

211 Cell extracts or RIPA and urea fractions obtained from sequential extractions were diluted with  
212 NuPAGE sample buffer. Proteins were separated on 4-12% NuPAGE Bis-Tris gels (Thermo Fisher  
213 Scientific) under denaturing conditions and transferred onto polyvinylidene difluoride (PVDF)  
214 membranes (Millipore). All blocking and antibody incubation steps were performed either in 5% milk  
215 in Tris-buffered saline (TBS) [25 mM Tris, 3 mM KCl, 140 mM NaCl, pH 7.4] supplemented with  
216 0.05% Tween-20 (TBS-T) (Thermo Fisher Scientific) or in 5% bovine serum albumin (BSA) in TBS-  
217 T. Western blots were developed with enhanced chemiluminescent substrates (ECL). Digital images  
218 were acquired with a ChemiDoc MP imaging system (BioRad). Where necessary, blots were stripped  
219 with stripping buffer for 15 min (Restore, Thermo Fisher Scientific) and re-probed with loading  
220 control antibodies.

221

## 222 **Lactate Dehydrogenase (LDH) assay to monitor cell death**

223 NSC-34 cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Thermo Fisher  
224 Scientific), supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin in a  
225 humidified incubator at 37°C and 5% CO<sub>2</sub>. Cells grown on 96-well plates coated with poly-*L*-lysine  
226 (BioCoat multiwell plates, Corning) were transiently transfected using Lipofectamine 3000 (Thermo  
227 Fisher Scientific) following the manufacturer's instructions 24 h after seeding (Lipofectamine  
228 3000:DNA ratio 3:1). Cells were treated with DMSO or p38α MAPK inhibitor (compound 1) at final  
229 concentrations of 1 or 10 μM for 4 hours post-transfection. LDH activity was measured from 50 μL  
230 of conditioned medium using LDH assay kit (Life Technologies) following the manufacturer's  
231 instructions.

232

233

## 234 **Longitudinal imaging and neuronal survival analysis**

235 Primary cortical neurons from E17 mouse embryos were cultured in 384-well plates. Neurons were  
236 co-transfected with plasmids expressing the fluorescent marker mApple and TDP-43<sup>M337V</sup>-EGFP.  
237 Neurons were treated with p38 $\alpha$  inhibitor, VX-745 (synthesized by AstraZeneca), at 0.3, 1, 3, or 9  
238  $\mu$ M, and imaged daily for 7 days on a custom-built highthroughput robotic microscopy system  
239 (Arrasate and Finkbeiner, 2005; Barmada et al., 2010). Images were montaged and neurons were  
240 segmented and tracked using custom-built algorithms. Cox proportional hazard analysis was used to  
241 determine hazard ratios. DMSO was used as a control.

242

## 243 **Cytoplasmic and nuclear protein extraction**

244 Cytoplasmic and nuclear protein extraction was performed using a commercial subcellular protein  
245 fractionation kit (Thermo Fisher Scientific). Briefly, ~80% confluent SH-SY5Y cells plated in a 100  
246 -mm dish were transfected as indicated for 24 hours. Cells were trypsinized and rinsed with cold  
247 phosphate-buffered saline (PBS), and cell suspensions were transferred to pre-chilled 1.5 mL  
248 microcentrifuge tubes. Cells were lysed in cytoplasmic extraction buffer at 4°C for 10 minutes with  
249 gentle mixing. After centrifugation at 500 x g at 4 °C for 5 min, the supernatant was collected and  
250 stored as the cytoplasmic fraction. After addition of membrane extraction buffer to the pellet, the tube  
251 was vortexed vigorously for 5 s and incubated at 4°C for 10 minutes with gentle mixing. After  
252 centrifugation at 3,000 x g at 4 °C for 5 min, the supernatant was collected and stored as the membrane  
253 fraction. The pellet was resuspended in nuclear extraction buffer followed by vigorous vortexing for  
254 15 s and incubation at 4°C for 30 minutes with gentle mixing. After centrifugation at 5,000 x g at  
255 4 °C for 5 min, the supernatant was collected and stored as the soluble nuclear fraction. The pellet  
256 was resuspended in chromatin-bound extraction buffer followed by vigorous vortexing for 15 s and  
257 incubation at room temperature for 15 s. The vortexing and incubation steps were repeated twice.  
258 After centrifugation at 16,000 x g at 4 °C for 5 min, the supernatant was collected and stored as the

259 chromatin-bound nuclear fraction. Bicinchoninic acid (BCA) assay was used to measure protein  
260 concentrations.

261

## 262 **Immunocytochemistry**

263 Cells grown on poly-*L*-lysine-coated glass coverslips were washed in PBS and fixed in 4%  
264 paraformaldehyde (in PBS) for 15 min followed by permeabilization in 0.25% Triton-X (in PBS) for  
265 10 min. Cells were blocked with 10% normal goat serum (in PBS, Abcam) for 1 h at room temperature  
266 and incubated overnight at 4°C in primary antibody diluted in blocking solution. The next day, cells  
267 were washed with PBS and incubated for 1 h in secondary antibody diluted in blocking solution.  
268 Coverslips were mounted with Prolong Gold Antifade Mountant with DAPI (Thermo Fisher  
269 Scientific). Image acquisition was performed using a Nikon A1 confocal/Eclipse Ti inverted  
270 microscope system and NIS Elements software (Nikon).

271

## 272 **Protein kinase assays with recombinant proteins**

273 5  $\mu$ L of 10x kinase assay buffer [25 mM Tris (pH 7.5), 5 mM  $\beta$ -glycerophosphate, 2 mM DTT, 0.1  
274 mM  $\text{Na}_2\text{VO}_4$ , 10 mM  $\text{MgCl}_2$ ] (Cell Signaling Technology), 200  $\mu$ M ATP (Cell Signaling  
275 Technology), 750 ng of recombinant human TDP-43 protein (Proteintech) and 300 ng of recombinant  
276 active p38 $\alpha$  kinase (SignalChem) were mixed in pre-chilled 1.5 mL tubes. Reactions were made up  
277 to a total volume of 50  $\mu$ L with ddH<sub>2</sub>O. Samples were mixed by flicking the tubes followed by brief  
278 centrifugation at 4°C and incubation at 30°C for 30 min. When indicated, kinase reactions were  
279 treated with 10  $\mu$ M the p38 $\alpha$  inhibitor compound 1. Reactions were stopped by adding 18  $\mu$ L of 4 $\times$   
280 NuPAGE sample buffer and boiling samples at 95°C for 5 minutes. Samples were subjected to  
281 immunoblotting or Coomassie staining followed by phosphorylation analysis by LC-MS/MS.

282

283

## 284 **Phosphorylation analysis by LC-MS/MS**

285 *In vitro* kinase reactions were separated on 4-12% NuPAGE Bis-Tris gels (Thermo Fisher Scientific)  
286 and gel bands were visualized with SimplyBlue™ SafeStain (Thermo Fisher Scientific). Gel bands  
287 were excised and cut into ~1 mm<sup>3</sup> pieces. The samples were reduced with 1 mM dithiothreitol (DTT)  
288 for 30 minutes at 60°C and then alkylated with 5 mM iodoacetamide for 15 minutes at room  
289 temperature. Gel pieces were then subjected to a modified in-gel trypsin digestion procedure  
290 (Shevchenko et al., 1996). Briefly, gel pieces were washed and dehydrated with acetonitrile for 10  
291 min followed by removal of acetonitrile. Pieces were then completely dried using a speed-vac. Gel  
292 pieces were rehydrated in 50 mM ammonium bicarbonate solution containing 12.5 ng/μL modified  
293 sequencing-grade trypsin (Promega) at 4°C before incubation overnight at 37°C. Peptides were later  
294 extracted by removing the ammonium bicarbonate solution, followed by one wash with a solution  
295 containing 50% acetonitrile and 1% formic acid. The extracts were then dried in a speed-vac for 1  
296 hour and stored at 4°C until analysis. On the day of analysis, samples were reconstituted in 5 - 10 μL  
297 of High Performance Liquid Chromatography (HPLC) solvent A (2.5% acetonitrile, 0.1% formic  
298 acid). A nano-scale reverse-phase HPLC capillary column was created by packing 2.6 μm C18  
299 spherical silica beads into a fused silica capillary (100 μm inner diameter, ~30 cm length) with a  
300 flame-drawn tip (Peng and Gygi, 2001). After equilibrating the column, each sample was loaded onto  
301 the column via a Famos auto sampler (LC Packings). A gradient was formed, and peptides were  
302 eluted with increasing concentrations of solvent B (97.5% acetonitrile, 0.1% formic acid). As each  
303 peptide was eluted it was subjected to electrospray ionization before entering an LTQ Orbitrap Velos  
304 Pro ion-trap mass spectrometer (Thermo Fisher Scientific). Eluting peptides were detected, isolated,  
305 and fragmented to produce a tandem mass spectrum of specific fragment ions for each peptide.  
306 Peptide sequences were determined by matching protein or translated nucleotide databases with the  
307 acquired fragmentation pattern using the software program Sequest (Thermo Finnigan) (Eng et al.,  
308 1994). The modification of 79.9663 mass units to serine, threonine, and tyrosine was included in the

309 database searches to determine phosphopeptides. Phosphorylation assignments were determined  
310 using the Ascore algorithm (Beausoleil et al., 2006). All databases include a reversed version of all  
311 sequences and the data was filtered to 1-2% peptide false discovery rate.

312

### 313 **Co-immunoprecipitation (Co-IP)**

314 Approximately 80% confluent SH-SY5Y cells plated in 6-well plates or 100-mm dishes were  
315 transfected as indicated. Cells were rinsed with cold PBS, and then lysed in cold lysis buffer [20 mM  
316 Tris-HCl (pH 7.5), 150 mM NaCl, 1 mM Na<sub>2</sub>EDTA, 1 mM EGTA, 1% Triton, 2.5 mM sodium  
317 pyrophosphate, 1 mM beta-glycerophosphate, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 1 µg/mL leupeptin] (Cell Signaling  
318 Technology), supplemented with protease inhibitors (cOmplete, Roche). Cells were incubated on ice  
319 for 5 minutes, collected in pre-chilled 1.5 mL tubes, sonicated briefly and cleared by centrifugation  
320 (14,000 x g at 4°C for 10 min). Lysate aliquots were stored as the input samples. Cell lysates were  
321 either incubated with Anti-FLAG® M2 Magnetic Beads (Millipore Sigma) overnight with continuous  
322 rotation at 4°C, or were subjected to pre-cleaning with protein A Dynabeads (Thermo Fisher  
323 Scientific) for 1 hour at 4°C followed by incubation with indicated primary antibodies overnight with  
324 continuous rotation at 4°C. Protein A Dynabeads were then added to pre-cleared antibody-containing  
325 samples, and the incubation was continued for an additional 2 hours at room temperature. Beads with  
326 immunoprecipitated proteins were washed 5x with either TBS [50 mM Tris HCl, 150 mM NaCl, pH  
327 7.4] (Anti-FLAG® M2 Magnetic Beads) or lysis buffer (Protein A Dynabeads). Immunoprecipitated  
328 proteins were eluted with 2x NuPAGE sample buffer by boiling for 3 min. Both input samples and  
329 immunoprecipitated proteins were analyzed by immunoblotting.

330

### 331 **Purification of recombinant TDP-43-MBP**

332 Wild-type TDP-43-MBP-6xHis and TDP-43 mutants S292E, S409:S410E, S292:S409:S410E, and  
333 R293F were expressed and purified as previously described (Wang et al., 2018). Briefly, TDP-43

334 variants were expressed in *E. coli* BL21-CodonPlus (DE3)-RIL cells (Agilent). Cell cultures were  
335 grown to an OD<sub>600</sub> of ~0.5-0.7 and then cooled down to 16°C. Protein expression was induced with  
336 1 mM IPTG overnight. Cells were harvested and resuspended in purification buffer (20 mM Tris-  
337 HCl, pH 8.0, 1 M NaCl, 10 mM imidazole, 10% (v/v) glycerol, 2 mM β-mercaptoethanol  
338 supplemented with cOmplete EDTA-free protease inhibitor cocktail) and lysed using 1 mg/mL  
339 lysozyme and sonication. Proteins were purified using Ni-NTA agarose (Qiagen) and eluted using  
340 300 mM imidazole in purification buffer. Proteins were then further purified over amylose resin  
341 (NEB) and eluted with elution buffer (20 mM Tris-HCl, pH 8.0, 1 M NaCl, 10 mM imidazole, 10  
342 mM maltose, 10% (v/v) glycerol, and 1 mM DTT). Purified proteins were concentrated, flash frozen  
343 and stored at -80°C.

344

#### 345 ***In vitro* TDP-43 aggregation assay**

346 Purified recombinant TDP-43-MBP-6xHis wild-type and TDP-43 mutants were first thawed and  
347 buffer exchanged into 20 mM HEPES-NaOH, pH 7.4, 150 mM NaCl and 1mM DTT using Micro  
348 Bio-Spin™ P-6 Gel Columns (Bio-Rad). Protein concentration was determined by NanoDrop, and  
349 the final concentration of TDP-43 was then adjusted to 5 μM in the same buffer. To measure  
350 aggregation kinetics, aggregation was initiated by cleavage of the MBP-6xHis tag using 1 μg/mL  
351 TEV protease (Cupo and Shorter, 2020) at t = 0, and turbidity was measured over 16 h at an  
352 absorbance of 395 nm using a TECAN M1000 plate reader. Values were normalized to wild-type  
353 TDP-43 + TEV protease to determine the extent of aggregation of TDP-43 mutants.

354

#### 355 ***In vitro* TDP-43 LLPS assay**

356 Purified recombinant TDP-43-MBP-6xHis wild-type and TDP-43 mutants were thawed and buffer  
357 exchanged as described above for the aggregation assay. Protein concentration was determined by  
358 NanoDrop, and reactions were prepared in phase separation buffer (20 mM HEPES-NaOH, pH 7.4,

359 150 mM NaCl, 1mM DTT, 100 mg/mL dextran from *Leuconostoc spp.* (Sigma)). Protein was always  
360 added last to each phase separation reaction, at a final concentration of 10  $\mu$ M. Reactions were  
361 incubated for 30 min at room temperature, and then 7.5  $\mu$ L of each reaction was mounted onto a glass  
362 slide and imaged by differential interference contrast (DIC) microscopy.  
363

#### 364 **Droplet image analysis**

365 DIC images of TDP-43 wild-type and its variants were analyzed using custom-written code in  
366 MATLAB. Each image was first converted into grayscale for processing. Then, Roberts gradient  
367 was used to filter out the noise. The pixel weight for each pixel in the image was determined based  
368 on the grayscale intensity differences before segmenting the images. The threshold for image  
369 segmentation was adjusted manually to ensure complete and accurate conversion to logical array.  
370 Droplets in the images were identified using circle Hough transform. The sensitivity was toggled  
371 either to detect missed droplets or to reduce the number of false positives. The detected circles were  
372 visualized on the original images. Various quantitative parameters, including the average area, total  
373 area, number of droplets and the lists of areas, were given as outputs to the code and were analyzed  
374 further through GraphPad Prism. The accuracy of the circle Hough transform was limited for  
375 droplets that were smaller than 5 pixels, where 10.8 pixels are equivalent to 1  $\mu$ m.

376

#### 377 **Quantification of immunocytochemistry and western blots, and statistical analysis**

378 Western blot band densities and immunocytochemical staining were quantified with ImageJ-Win64  
379 software. For statistical analysis, we used GraphPad Prism 7 and 8, and used unpaired t-test or one-  
380 way ANOVA followed by Sidak's or Dunnett's multiple comparison test, as indicated for each  
381 experiment. All assays were repeated at least three times. A *p*-value less than 0.05 was considered  
382 statistically significant.

383



## 384 **Results**

### 385 **Inhibition of p38 $\alpha$ reduces TDP-43 aggregation, phosphorylation, and toxicity**

386 Several lines of evidence point to a role for p38 MAPK in the development and progression of ALS  
387 (Bendotti et al., 2004; Corrêa and Eales, 2012; Gibbs et al., 2018; Tortarolo et al., 2003; Zhan et al.,  
388 2015). Thus, we first asked whether p38 $\alpha$  modulates the formation of insoluble TDP-43 aggregates  
389 in human neuronal SH-SY5Y cells. Increased expression of wild-type or ALS-linked TDP-43 variants  
390 elicits ALS-like phenotypes in various *in vitro* and *in vivo* models (Arnold et al., 2013; Johnson et  
391 al., 2009; Watkins et al., 2020; Wobst et al., 2017; Xu et al., 2011). Indeed, elevated expression of  
392 wild-type TDP-43 is connected with FTD (Gitcho et al., 2009) and disease-linked TDP-43  
393 aggregation is proposed to increase TDP-43 expression due to loss of TDP-43 autoregulation (Gasset-  
394 Rosa et al., 2019; Polymenidou et al., 2011). TDP-43<sup>M337V</sup>, a pathological mutant form of TDP-43  
395 associated with fALS, has been shown to be especially aggregation-prone and highly phosphorylated  
396 (Johnson et al., 2009; Sreedharan et al., 2008; Wobst et al., 2017). We therefore used the TDP-43<sup>M337V</sup>  
397 variant in our experiments to detect more prominent changes in TDP-43 solubility and  
398 phosphorylation, and to maximize our experimental signal window. Using siRNA-mediated  
399 knockdown of p38 $\alpha$  and TDP-43<sup>M337V</sup> expression, we monitored the accumulation of total and  
400 phosphorylated TDP-43 (pTDP-43) in RIPA- or urea-soluble fractions over time. We found that the  
401 depletion of p38 $\alpha$  significantly reduced the accumulation of insoluble TDP-43<sup>M337V</sup> in the urea  
402 fraction at all time points (Figure 1B-1C). Thus, reduction of p38 $\alpha$  reduces TDP-43 aggregation in  
403 human neuronal cells.

404

405 Phosphorylation of S409 and S410 of TDP-43 is a consistent feature in all sporadic and familial forms  
406 of TDP-43 proteinopathies (Neumann et al., 2009). TDP-43 is phosphorylated at S409/S410 in  
407 pathological inclusions, but S409/S410 phosphorylation is not observed under physiological  
408 conditions in the nucleus (Neumann et al., 2009). Interestingly, we observed a marked decrease in

409 pTDP-43 (i.e. TDP-43 phosphorylated at S409/S410) in the urea fraction when p38 $\alpha$  is knocked down  
410 (Figure 1B and 1D). Similarly, increased expression of TDP-43<sup>WT</sup>, in place of TDP-43<sup>M337V</sup>, or using  
411 a different siRNA to knockdown p38 $\alpha$  also decreased pTDP-43 in the urea fraction (Figure S1). Thus,  
412 reduction of p38 $\alpha$  reduces TDP-43 aggregation and pathological phosphorylation of S409/S410 in  
413 human neuronal cells.

414  
415 Next, we assessed whether pharmacological inhibition of p38 $\alpha$  activity affects the solubility and the  
416 phosphorylation status of TDP-43 in a similar manner to genetic ablation. Indeed, treatment of SH-  
417 SY5Y cells with the p38 $\alpha$  inhibitor, compound 1 (see Materials and Methods), significantly reduced  
418 TDP-43<sup>M337V</sup> phosphorylation and aggregation, in a concentration- and time-dependent manner  
419 (Figure 1E-1H). Note that we only observe TDP-43 phosphorylation at S409/S410 in the insoluble  
420 urea fraction and not in the soluble RIPA fraction (Figure 1E), indicating the pathological nature of  
421 S409/S410 phosphorylation. Our findings suggest that p38 $\alpha$  inhibition is an effective strategy to  
422 reduce pathological TDP-43 aggregation and phosphorylation.

423  
424 To test whether p38 $\alpha$  pharmacological inhibition affects toxicity induced by increased TDP-43  
425 expression, we performed an LDH cytotoxicity assay in mouse motor-neuron-like NSC-34 cells.  
426 Expression of TDP-43<sup>WT</sup> (Figure 1I) or TDP-43<sup>M337V</sup> in NSC-34 cells induces cytotoxicity.  
427 Importantly, inhibition of p38 $\alpha$  by compound 1 significantly rescued TDP-43<sup>WT</sup>-induced cytotoxicity  
428 (Figure 1I). Thus, pharmacological inhibition of p38 $\alpha$  mitigates TDP-43 toxicity in motor-neuron-  
429 like NSC-34 cells.

430  
431 To determine whether p38 $\alpha$  pharmacological inhibition rescues TDP-43-induced neurodegeneration,  
432 we employed a longitudinal imaging system to monitor neuronal survival (Barmada et al., 2010). We  
433 cultured primary mouse cortical neurons, uniformly labeled them with a fluorescent protein

434 (mApple), and expressed TDP43<sup>M337V</sup>. We imaged and tracked the neurons daily for 7 days and used  
435 Cox proportional hazard analysis to measure the cumulative risk of death and hazard ratios. We have  
436 previously shown that this assay is very sensitive to detecting TDP-43-induced neurodegeneration  
437 (Barmada et al., 2010). We found that treatment of neurons with a brain-penetrant p38 $\alpha$   
438 pharmacological inhibitor, VX-745 (also known as Neflamapimod) (Duffy et al., 2011), which is in  
439 phase 2 clinical trials for Alzheimer's disease, Huntington's disease, and dementia with Lewy bodies  
440 (Germann and Alam, 2020; Prins et al., 2021), significantly reduced the hazard ratio of TDP-43<sup>M337V</sup>-  
441 expressing neurons (Figure 1J). Thus, clinical stage, brain-penetrant p38 $\alpha$  inhibitors can also mitigate  
442 TDP-43 neurotoxicity. Taken together, our results demonstrate that p38 $\alpha$  inhibition reduces the  
443 pathological aggregation and phosphorylation of TDP-43, and mitigates TDP-43 toxicity in multiple  
444 settings.

445

#### 446 **Constitutively active p38 $\alpha$ promotes TDP-43 aggregation, S409/S410 phosphorylation, and** 447 **cytoplasmic accumulation**

448 The siRNA knockdown experiments demonstrated that p38 $\alpha$  depletion reduces TDP-43 aggregation  
449 and phosphorylation at S409/S410. Therefore, we hypothesized that p38 $\alpha$  overexpression would  
450 increase TDP-43 aggregation and phosphorylation. To test our hypothesis, we co-expressed TDP-  
451 43<sup>M337V</sup> with wild-type, constitutively active (CA), or dominant negative (DN) forms of p38 $\alpha$  in SH-  
452 SY5Y cells. Expression of WT or DN-p38 $\alpha$  did not have any significant effect on S409/S410  
453 phosphorylation or accumulation of TDP-43 in the urea fraction (Figure 2A-2D). By contrast, CA-  
454 p38 $\alpha$  promoted TDP-43<sup>M337V</sup> aggregation, as evidenced by the accumulation of TDP-43 in the urea  
455 fraction (Figure 2A-2D). CA-p38 $\alpha$  also significantly increased TDP-43 phosphorylation at  
456 S409/S410 (Figure 2A-2D). Thus, p38 $\alpha$  activation promotes pathological TDP-43 phosphorylation  
457 and aggregation.

458

459 Next, we assessed whether TDP-43<sup>M337V</sup> co-localizes with p38 $\alpha$  in cells. Previously, TDP-43<sup>M337V</sup>  
460 and p38 $\alpha$  have been detected in ubiquitinated inclusions (Bendotti et al., 2004; Wobst et al., 2017).  
461 Interestingly, we found a significant increase in the number of cells displaying intranuclear TDP-  
462 43<sup>M337V</sup> inclusions specifically when CA-p38 $\alpha$  was co-expressed (Figure 2E-2F). Furthermore, these  
463 intranuclear TDP-43 inclusions stained positive for p38 $\alpha$  (Figure 2E). We also found that expression  
464 of CA-p38 $\alpha$  promoted the cytoplasmic accumulation of TDP-43, as shown by an increase of TDP-43  
465 in the cytoplasmic fraction (Figure 2G-2H). Together, our data demonstrate that aberrant activation  
466 of p38 $\alpha$  promotes several hallmarks of ALS pathology, including TDP-43 aggregation,  
467 phosphorylation at S409/S410, and cytoplasmic accumulation.

#### 468 469 **TDP-43 is directly phosphorylated by p38 $\alpha$ at residues S292, S409, and S410**

470 Co-localization of TDP-43 and CA-p38 $\alpha$  in intranuclear aggregates might indicate a physical  
471 interaction between TDP-43 and p38 $\alpha$  (Figure 2E). To explore this possibility, we transfected SH-  
472 SY5Y with TDP-43<sup>WT</sup> and p38 $\alpha$ , and then immunoprecipitated p38 $\alpha$  or TDP-43 (Figure 3A). TDP-  
473 43<sup>WT</sup> and p38 $\alpha$  co-immunoprecipitated in both reciprocal immunoprecipitation experiments,  
474 suggesting a robust interaction between TDP-43 and p38 $\alpha$  (Figure 3A). Consistently, siRNA-  
475 mediated knockdown of p38 $\alpha$  reduced the amount of pulled down pTDP-43, and also diminished the  
476 amount of endogenous p38 $\alpha$  that co-immunoprecipitated with TDP-43<sup>M337V</sup> (Figure S2), further  
477 validating an interaction between p38 $\alpha$  and TDP-43.

478  
479 Next, we asked whether TDP-43 is directly phosphorylated by p38 $\alpha$ . Thus, we performed *in vitro*  
480 kinase assays. Samples containing recombinant wild-type TDP-43 with or without active p38 $\alpha$  were  
481 analyzed by western blot analysis after a 30-minute incubation period. In the absence of p38 $\alpha$ , no  
482 phosphorylation was detected at the S409/S410 site (Figure 3B). However, in the presence of active  
483 kinase, a robust band of phosphorylated TDP-43 was observed, indicating that TDP-43 is directly

484 phosphorylated by p38 $\alpha$  at S409/S410 (Figure 3B). As expected, the phosphorylation of TDP-43 was  
485 prevented by pharmacological inhibition of p38 $\alpha$  with compound 1 (Figure 3B).

486

487 While S409/S410 residues are thought to be the major pathological phosphorylation sites in TDP-43  
488 (Neumann et al., 2009), we sought to investigate whether p38 $\alpha$  phosphorylates TDP-43 at additional  
489 serine or threonine residues. Therefore, we performed LC-MS/MS analysis of trypsin-digested TDP-  
490 43 after incubation with p38 $\alpha$ , which revealed that p38 $\alpha$  also phosphorylates TDP-43 at serine residue  
491 292 (S292) (Figure 1A). In fact, TDP-43 is phosphorylated at S292 in the brains of ALS patients  
492 (Kametani et al., 2016). Furthermore, a serine-to-asparagine mutation at this site (S292N) is  
493 genetically linked to sALS and fALS (Harrison and Shorter, 2017; Xiong et al., 2010; Zou et al.,  
494 2012). To investigate the potential effects of phosphorylation at S292 on the aggregation propensity  
495 of TDP-43, we took a site-directed mutagenesis approach to generate phospho-mimetic (S292E; TDP-  
496 43<sup>S292E</sup>), phospho-dead (S292A; TDP-43<sup>S292A</sup>), and ALS-linked (S292N; TDP-43<sup>S292N</sup>) TDP-43  
497 mutants, which were transfected into SH-SY5Y cells. We compared the aggregation propensity of  
498 these S292 mutants to that of TDP-43<sup>WT</sup> and TDP-43 variants harboring S409:S410A (TDP-  
499 43<sup>S409:S410A</sup>), S409:S410E (TDP-43<sup>S409:S410E</sup>) and S292:S409:S410A (TDP-43<sup>S292:S409:S410A</sup>)  
500 mutations. Interestingly, overexpressing the phospho-mimetic mutant TDP-43<sup>S292E</sup> increased the  
501 aggregation propensity of TDP-43, as evidenced by the accumulation of TDP-43 in the urea fraction,  
502 and also enhanced its phosphorylation at serine residues 409/410 (Figure 3C and 3D). Moreover,  
503 TDP-43<sup>S292E</sup> enhanced the formation of 35 kDa C-terminal fragments of TDP-43 (CTF), which are  
504 associated with TDP-43 proteinopathy (Buratti, 2018) (Figure 3C and 3E). Neither TDP-43<sup>S292A</sup>,  
505 TDP-43<sup>S292N</sup>, TDP-43<sup>S409:S410A</sup>, TDP-43<sup>S409:S410E</sup> nor TDP-43<sup>S292:S409:S410A</sup> had any significant effect  
506 on TDP-43 aggregation propensity compared to TDP-43<sup>WT</sup> (Figure 3C-3D). These results suggest  
507 that aggregation of TDP-43 may be enhanced by S292 phosphorylation in neuronal cells. Moreover,  
508 S292 phosphorylation also likely stimulates phosphorylation at S409/410.

## 509 **TDP-43 undergoes arginine methylation catalyzed by PRMT1**

510 Our data suggest a role for S292 in regulating TDP-43 aggregation via p38 $\alpha$ -mediated  
511 phosphorylation. An alignment of TDP-43 amino acid sequences revealed that S292 is highly  
512 conserved across the phylogenic spectrum from *Homo sapiens* to *Gallus gallus* (Figure 4A).  
513 Intriguingly, an arginine-glycine-glycine (RGG) motif directly follows the S292 residue and is also  
514 highly conserved (Figure 4A). RGG/RG motifs are preferred substrates for methylation by members  
515 of the PRMT family (Thandapani et al., 2013). Given that several studies have shown that arginine  
516 methylation can attenuate phosphorylation at nearby residues on the same protein (Guo et al., 2010;  
517 Hsu et al., 2011; Lu et al., 2017; Yamagata et al., 2008), we hypothesized that arginine methylation  
518 at R293 could potentially interfere with p38 $\alpha$ -mediated phosphorylation at the adjacent residue S292.  
519 Recent proteomic studies have found that TDP-43 is methylated in HCT116 and HEK293 cells as  
520 well as in mouse embryos and human brain tissue (Guo et al., 2014; Larsen et al., 2016). To assess  
521 TDP-43 arginine methylation in SH-SY5Y cells, we immunoprecipitated endogenous TDP-43 and  
522 probed using antibodies against monomethyl-arginine (MMA), asymmetric dimethyl-arginine  
523 (ADMA) or symmetric dimethyl-arginine (SDMA). Western blot analysis showed that  
524 immunoprecipitated TDP-43 was indeed mono- and asymmetrically dimethylated and was efficiently  
525 recognized by the MMA and ADMA antibodies, but not a SDMA antibody (Figure 4B). Critically,  
526 when we depleted PRMT1, the most abundant PRMT in mammalian cells (Tang et al., 2000), using  
527 siRNA-mediated knockdown in SH-SY5Y cells, followed by TDP-43 immunoprecipitation and  
528 western blot analysis, we found a significant decrease in methylation of endogenous TDP-43 (Figure  
529 4C). Additionally, treating cells with AdOx, a global methyltransferase inhibitor, for 24 hours  
530 significantly decreased the methylation of overexpressed TDP-43<sup>WT</sup>, evident by a decrease in  
531 immunoprecipitated TDP-43 detected by anti-MMA and anti-ADMA (Figure 4D). Collectively, these  
532 data suggest that PRMT1 methylates TDP-43.

533

534 Next, we wanted to verify whether R293 is the targeted residue for arginine methylation. Using site-  
535 directed mutagenesis, we generated a methylation-dead mutant of TDP-43 by substituting the arginine  
536 residue at 293 to a lysine (TDP-43<sup>R293K</sup>). Unlike for TDP-43<sup>WT</sup>, immunoprecipitating TDP-43<sup>R293K</sup>  
537 and probing for monomethylated TDP-43 using anti-MMA antibody revealed an absence of  
538 monomethylated TDP-43 (Figure 4E). Overall, our results provide evidence that PRMT1 methylates  
539 TDP-43 at R293 in human neuronal cells.

540

### 541 **TDP-43 arginine methylation favors normal LLPS over aberrant aggregation**

542 Arginine methylation of TDP-43 has been largely unexplored. Thus, little is known about R293  
543 methylation and whether it affects TDP-43 LLPS or aggregation propensity at the pure protein level.  
544 Likewise, we have limited understanding of how specific phosphorylation events might affect TDP-  
545 43 LLPS and aggregation at the pure protein level. To address how phosphorylation and methylation  
546 can alter TDP-43 LLPS behavior, we first performed *in vitro* droplet formation assays. Purified  
547 recombinant maltose-binding protein (MBP)-tagged TDP-43<sup>WT</sup>, the phospho-mimicking mutants  
548 TDP-43<sup>S292E</sup>, TDP-43<sup>S409:S410E</sup>, TDP-43<sup>S292:S409:S410E</sup>, and the arginine methylation-mimic TDP-  
549 43<sup>R293F</sup> (Campbell et al., 2012; Huq et al., 2006; Liu et al., 2019), were separately incubated at  
550 physiological concentration (10 $\mu$ M) (Ling et al., 2010) in phase separation buffer containing  
551 physiological salt concentration and 10% (w/v) dextran to mimic the crowded cellular environment  
552 (Mann et al., 2019; McGurk et al., 2018). Formation of TDP-43 droplets was then visualized using  
553 DIC microscopy. TDP-43<sup>WT</sup> formed spherical droplets that were relatively large in size (average area  
554 of  $\sim 12.6\mu\text{m}^2 \pm 1.3$ ), and capable of fusion events indicating liquid-like properties (Figure 5A-C). In  
555 contrast, all three phosphomimetic TDP-43 mutants partitioned into droplets that were smaller in size  
556 (TDP-43<sup>S292E</sup>,  $\sim 5.8\mu\text{m}^2 \pm 0.6$ ; TDP-43<sup>S409:S410E</sup>,  $\sim 3.9\mu\text{m}^2 \pm 0.3$ ; and TDP-43<sup>S292:S409:S410E</sup>,  $\sim 3.2\mu\text{m}^2 \pm$   
557  $0.2$ ) compared to those of TDP-43<sup>WT</sup> (Figure 5A-C). There was no change in the number of droplets  
558 between mutants and TDP-43<sup>WT</sup>. Thus, these reductions in droplet sizes suggest that phosphorylation



559 of S292, S409, and S410 may limit the LLPS propensity of TDP-43. Interestingly, the arginine-  
560 methylation mimic, TDP-43<sup>R293F</sup>, formed large droplets that were comparable to TDP-43<sup>WT</sup>  
561 ( $\sim 12.6 \mu\text{m}^2 \pm 1.9$ ) (Figure 5A-C). These results indicated that phosphorylation and methylation likely  
562 have contrasting effects on LLPS of TDP-43. Thus, R293 methylation likely permits wild-type levels  
563 of TDP-43 LLPS, whereas S292, S409, and S410 phosphorylation likely reduce TDP-43 LLPS.

564  
565 Given these outcomes, we next compared the effects of phosphorylation and methylation mimics on  
566 TDP-43 aggregation propensity using an *in vitro* aggregation assay. Recombinant TDP-43-MBP  
567 protein constructs were incubated with TEV protease and their aggregation was monitored over time.  
568 Under these conditions, selective cleavage of the MBP tag by TEV protease results in the formation  
569 of solid-phase TDP-43 aggregates and fibrils (Cook et al., 2020), indicated by an increase in turbidity.  
570 Here, we found that there were only minor differences in aggregation between TDP-43<sup>WT</sup> and the  
571 phosphomimics TDP-43<sup>S292E</sup> and TDP-43<sup>S409E:S410E</sup> (Figure 5D). By contrast, the phosphomimic  
572 TDP-43<sup>S292E:S409E:S410E</sup> and the arginine methylation mimic, TDP-43<sup>R293F</sup>, exhibited modestly reduced  
573 aggregation (Figure 5D). However, compared to TDP-43<sup>WT</sup> this reduced aggregation was only  
574 significant for TDP-43<sup>R293F</sup> (Figure 5E). Strikingly, plotting the normalized aggregation against the  
575 LLPS propensities of TDP-43<sup>WT</sup> and its mutant forms, we found that the phosphomimetic mutants  
576 have a relatively higher tendency for aggregation over LLPS (Figure 5F). By contrast, TDP-43<sup>R293F</sup>  
577 has a relatively higher tendency to undergo LLPS over aggregation (Figure 5F). Taken together, our  
578 results imply that TDP-43 phosphorylation and methylation may have opposing effects on TDP-43  
579 (Figure 5F). Phosphorylation at S292, S409, and S410 reduces the propensity for TDP-43 to undergo  
580 LLPS, but has limited effects on TDP-43 aggregation (Figure 5G). Thus, S292, S409, and S410  
581 phosphorylation may divert TDP-43 toward aggregation and away from LLPS (Figure 5G). This  
582 finding could explain why phosphorylated TDP-43 accumulates as aggregates in the urea fraction in  
583 cells. In contrast, R293 methylation allows TDP-43 to undergo normal LLPS, but reduces TDP-43



584 aggregation (Figure 5G). Thus, R293 methylation may reduce the propensity of TDP-43 to aggregate  
585 and enter the urea fraction in cells.

586

### 587 **Arginine methylation regulates TDP-43 aggregation in human neuronal cells**

588 We next investigated the impact of arginine methylation on TDP-43 in a cellular context. SH-SY5Y  
589 cells were treated with AdOx, an arginine methyltransferase inhibitor, followed by fractionation and  
590 western blot analysis. We found that global methyltransferase inhibition promoted the accumulation  
591 of TDP-43 in the urea fraction (Figure 6A and B). Conversely, the overexpression of PRMT1 led to  
592 a decrease in TDP-43 aggregation, evident by a decrease of total and phosphorylated TDP-43 in the  
593 urea fraction (Figure 6C and D). To further investigate the effect of hypomethylation on TDP-43  
594 aggregation, we next mutated TDP-43 to have an additional RGG motif (TDP-43<sup>G308R</sup>) (Figure 4A).  
595 Based on previous studies, the introduction of an additional RGG motif has led to the  
596 hypomethylation and subsequent aggregation of another ALS-linked RBP, FUS (Qamar et al., 2018).  
597 Accordingly, we found that TDP-43<sup>G308R</sup> significantly accumulated in the urea fraction and was more  
598 highly phosphorylated at S409/S410, compared to TDP-43<sup>WT</sup> (Figure 6E and F). Together, these  
599 findings suggest that arginine methylation exerts a protective role on TDP-43, perhaps by decreasing  
600 its aggregation propensity via regulation of phosphorylation.

601

### 602 **Crosstalk between TDP-43 arginine methylation and p38 $\alpha$ -mediated phosphorylation**

603 The observation that TDP-43 undergoes PRMT1-mediated arginine methylation at the R293 site,  
604 coupled with our purified protein data suggesting contrasting outcomes between TDP-43  
605 phosphorylation and methylation, led us to ask whether phosphorylation at S292 interferes with  
606 arginine methylation at the adjacent residue, R293, or vice versa (Figure 4A). To answer this question,  
607 we expressed flag-tagged TDP-43<sup>WT</sup> as well as TDP-43<sup>S292E</sup>, TDP-43<sup>S292N</sup>, and TDP-43<sup>S292A</sup> mutants  
608 in SH-SY5Y cells and analyzed their methylation status. Immunoprecipitation followed by

609 immunoblotting revealed a striking reduction in the MMA-signal in the phospho-mimicking TDP-  
610 43<sup>S292E</sup> mutant (Figure 7A). Interestingly, when compared to TDP-43<sup>WT</sup>, the ALS-linked TDP-43<sup>S292N</sup>  
611 mutant showed a modest decrease in MMA-levels, whereas the TDP-43<sup>S292A</sup> mutant did not show  
612 any difference (Figure 7A). As shown previously (Figure 3C), the TDP-43<sup>S292E</sup> mutant also promoted  
613 phosphorylation of TDP-43 at S409/S410, further underlining the anti-correlative relationship  
614 between TDP-43 arginine methylation and S409/S410 phosphorylation (Figure 7A). These  
615 observations suggest that phosphorylation at S292, possibly combined with increased  
616 phosphorylation at S409/S410, could interfere with TDP-43 methylation. However, these findings do  
617 not rule out that a reduction in TDP-43 methylation could be due to steric hindrance caused by the  
618 glutamic acid residue at position 292.

619  
620 To further define the relationship between these two PTMs, we studied the methylation status of TDP-  
621 43 after genetic depletion of p38 $\alpha$ . As shown previously (Figure 1B), we found that p38 $\alpha$   
622 downregulation reduced the phosphorylation of TDP-43<sup>WT</sup> at S409/S410 (Figure 7B). Interestingly,  
623 western blot analysis also revealed that p38 $\alpha$  depletion significantly decreased the formation of the  
624 ~35 kDa TDP-43-CTF (Figure 7B and C), and resulted in elevated levels of mono-methylated TDP-  
625 43-CTF (Figure 7B and D). These observations suggest that there is an interplay between arginine  
626 methylation and p38 $\alpha$ -mediated phosphorylation, and that reduced p38 $\alpha$  activity could increase TDP-  
627 43 arginine methylation. Indeed, we found that overexpression of PRMT1 led to a strong increase in  
628 mono-methylated TDP-43-CTF and a striking reduction in TDP-43 phosphorylation at S409/S410  
629 (Figure 7E), further corroborating the hypothesis that there is crosstalk between TDP-43 arginine  
630 methylation and phosphorylation at disease-relevant residues.

631

632

## 633 **Discussion**

634 Since the discovery that ~97% of ALS cases and ~50% of FTD cases present with TDP-43  
635 proteinopathy (Arai et al., 2006; Harrison and Shorter, 2017; Neumann et al., 2006), TDP-43 has  
636 been subject to much investigation. However, the mechanisms leading to an accumulation of  
637 insoluble TDP-43 aggregates are not yet fully understood. Aberrant TDP-43 phosphorylation is one  
638 of the major distinguishing pathological features of TDP-43 inclusions in human brains (Hasegawa  
639 et al., 2008; Neumann et al., 2009). Although the consequences of these phosphorylation events have  
640 not been unequivocally established, aberrant phosphorylation of TDP-43 is associated with  
641 cytoplasmic mislocalization, decreased solubility, aberrant cleavage and cytotoxicity (Barmada et al.,  
642 2010; Brady et al., 2011; Kim et al., 2015; Liachko et al., 2010; Nonaka et al., 2009; Zhang et al.,  
643 2010). To date, casein kinases CK1 and CK2, CDC7 and TTBK1/2 have been shown to phosphorylate  
644 TDP-43 *in vitro* and *in vivo*, and promote its pathological aggregation and neurotoxicity (Carlomagno  
645 et al., 2014; Choksi et al., 2014; Goh et al., 2018; Kametani et al., 2009; Liachko et al., 2013, 2014;  
646 Liu et al., 2015; Meyerowitz et al., 2011; Nonaka et al., 2016; Sreedharan et al., 2015; Taylor et al.,  
647 2018). However, some studies have found that hyperphosphorylation of the TDP-43 PrLD can reduce  
648 TDP-43 LLPS and aggregation (Li et al., 2011; Silva et al., 2021), indicating a complex interplay  
649 between combinatorial TDP-43 PTMs and pathology. Importantly, increased activation of the MAP  
650 kinase p38 has been detected in human post-mortem ALS tissue, which is further substantiated by the  
651 findings that persistent activation of the p38 signaling pathways induce neurodegeneration (Bendotti  
652 et al., 2004; Dewil et al., 2007; Tortarolo et al., 2003). However, until now, the effect of p38 $\alpha$  on  
653 TDP-43 specifically has not been explored.

654

655 In this study, we elucidated the impact of p38 $\alpha$  on TDP-43 proteinopathy. Using neuronal cells, we  
656 first showed that genetic depletion and pharmacological inhibition of p38 $\alpha$  suppressed TDP-43  
657 phosphorylation, aggregation, and toxicity. We further established that TDP-43 is a substrate of p38 $\alpha$ -

658 mediated phosphorylation. Using *in vitro* biochemical assays followed by LC-MS/MS we established  
659 that TDP-43 is directly phosphorylated by p38 $\alpha$  at residues S292, and S409/S410. Intriguingly,  
660 mutations at S292 have been linked to pathogenicity in both sporadic and familial ALS. However, no  
661 biochemical mechanistic data have been reported on the effect of this mutation (Xiong et al., 2010;  
662 Zou et al., 2012). Here, we demonstrated that the phospho-mimetic mutant TDP-43<sup>S292E</sup> significantly  
663 promoted phosphorylation of TDP-43 at S409/S410 and enhanced the accumulation of insoluble  
664 TDP-43 aggregates. These findings identified S292 as a major site for TDP-43 phospho-regulation.  
665 Interestingly, we found that S292, and the RGG-motif immediately following this residue are highly  
666 conserved across the phylogenic spectrum. The RGG-motif presents a major site for methylation by  
667 members of the PRMT family (Chang et al., 2011; Huang et al., 2018; Thandapani et al., 2013; Wall  
668 and Lewis, 2017), and we show here that R293 is a major site for methylation by PRMT1 in human  
669 neuronal cells. This result supports earlier proteomic studies of mouse embryonic and brain tissue  
670 indicating that TDP-43 can be methylated at R293 (Guo et al., 2014; Larsen et al., 2016). Furthermore,  
671 recent evidence also corroborates the possible role of arginine methylation in neurodegeneration.  
672 Specifically, PRMT1 was identified as a significant modulator of toxicity in C9-ALS (Ortega et al.,  
673 2020).

674  
675 There is increasing evidence suggesting that phosphorylation and arginine methylation co-exist on  
676 the same protein, and that these PTMs can have opposing or potentiating effects on protein function  
677 (Basso and Pennuto, 2015; Lu et al., 2017). For example, the functions of p16 protein are regulated  
678 by the antagonistic crosstalk between arginine methylation at residue R138 and phosphorylation at  
679 residue S140 (Lu et al., 2017). Here, we provide similar evidence for the crosstalk between PRMT1-  
680 catalyzed arginine methylation at R293 and p38 $\alpha$ -mediated phosphorylation at the adjacent residue  
681 S292 in the regulation of TDP-43 LLPS and aggregation. Our *in vitro* and *in vivo* data elucidate a  
682 dichotomous relationship between methylation and phosphorylation of TDP-43. Perhaps as a

683 protective mechanism, TDP-43 methylation at R293 reduces phosphorylation at S292 and  
684 S409/S410, which in turn reduces TDP-43 aggregation. Conversely, the phosphomimetic S292E  
685 reduces the levels of TDP-43 methylation. Our biochemical studies suggest that S292 and S409/S410  
686 phosphorylation render TDP-43 less prone to undergo LLPS, which may divert TDP-43 along  
687 pathological aggregation trajectories (Conicella et al., 2016). Whether TDP-43 methylation at R293  
688 inhibits phosphorylation at S292, or vice versa, due to steric hindrance have yet to be further  
689 investigated. Nevertheless, our study suggests for the first time an intricate interplay between protein  
690 phosphorylation and arginine methylation in the regulation of TDP-43. While additional studies will  
691 be necessary to further clarify the dynamics of these regulatory processes, especially in the context  
692 of more complex, clinically-relevant models, our study provides a platform for developing novel  
693 therapeutic strategies to inhibit p38 $\alpha$  or promote PRMT1 activity to rescue TDP-43 pathologies  
694 associated with ALS/FTD. Indeed, it is interesting to note that the brain-penetrant p38 $\alpha$  inhibitor,  
695 VX-745, which mitigates TDP-43 toxicity in primary neurons (Figure 1J), has reached phase 2  
696 clinical trials for Alzheimer's disease, Huntington's disease, and dementia with Lewy bodies  
697 (Germann and Alam, 2020; Prins et al., 2021). Our data indicate that VX-745 might also be  
698 considered as a clinical candidate for ALS/FTD that presents with TDP-43 proteinopathy. Indeed, we  
699 suggest that strategies to reduce p38 $\alpha$ -mediated TDP-43 phosphorylation and promote R293  
700 methylation could have therapeutic utility for ALS/FTD and other TDP-43 proteinopathies.

701

## 702 **Acknowledgements**

703 We thank Charlotte Fare and Katie Copley for feedback on the manuscript, and Rebecca Jarvis for  
704 small-molecule curation. MA and HMO were supported by the AstraZeneca postdoctoral fellowships.  
705 BC was a member of the AstraZeneca graduate program. KLM was supported by supported by a NSF  
706 graduate research fellowship (DGE-1321851). AFF was supported by NIH grants T32AG00255 and  
707 F31NS087676. EMB was supported by a Milton Safenowitz Post-Doctoral Fellowship from the ALS

708 Association and NIH grant F32NS108598. RRC was supported by NIH grants T32GM008275,  
709 F31AG060672, and a Blavatnik Family Fellowship in Biomedical Research. JS was supported by  
710 Target ALS, The Association for Frontotemporal Degeneration, The Packard Foundation for ALS  
711 research, The ALS Association, The G. Harold and Leila Y. Mathers Charitable Foundation, the  
712 Office of the Assistant Secretary of Defense for Health Affairs (USA), through the Amyotrophic  
713 Lateral Sclerosis Research Program (W81XWH-20-1-0242), and NIH grants R01GM099836 and  
714 R21AG065854. This work was supported by a grant to NJB, DGB, JS, ADG, SF from the Target  
715 ALS Foundation and ALS Finding a Cure.

716

717 **Conflict of interest**

718 HJW, DGB, and NJB were all full-time employees and shareholders of AstraZeneca at the time these  
719 studies were conducted. SJM serves as a consultant for SAGE Therapeutics and AstraZeneca,  
720 relationships that are regulated by Tufts University. JS is a consultant for Dewpoint Therapeutics,  
721 Maze Therapeutics, Vivid Sciences, Korro Bio, and ADRx.

722

723 **References**

- 724 Alami, N.H., Smith, R.B., Carrasco, M.A., Williams, L.A., Winborn, C.S., Han, S.S.W., Kiskinis,  
725 E., Winborn, B., Freibaum, B.D., Kanagaraj, A., et al. (2014). Axonal Transport of TDP-43 mRNA  
726 Granules Is Impaired by ALS-Causing Mutations. *Neuron* 81, 536–543.
- 727 Arai, T., Hasegawa, M., Akiyama, H., Ikeda, K., Nonaka, T., Mori, H., Mann, D., Tsuchiya, K.,  
728 Yoshida, M., Hashizume, Y., et al. (2006). TDP-43 is a component of ubiquitin-positive tau-  
729 negative inclusions in frontotemporal lobar degeneration and amyotrophic lateral sclerosis.  
730 *Biochem Bioph Res Co* 351, 602–611.
- 731 Arnold, E.S., Ling, S.-C., Huelga, S.C., Lagier-Tourenne, C., Polymenidou, M., Ditsworth, D.,  
732 Kordasiewicz, H.B., McAlonis-Downes, M., Platoshyn, O., Parone, P.A., et al. (2013). ALS-linked  
733 TDP-43 mutations produce aberrant RNA splicing and adult-onset motor neuron disease without  
734 aggregation or loss of nuclear TDP-43. *Proc National Acad Sci* 110, E736–E745.
- 735 Arrasate, M., and Finkbeiner, S. (2005). Automated microscope system for determining factors that  
736 predict neuronal fate. *P Natl Acad Sci Usa* 102, 3840–3845.
- 737 Aulas, A., Stabile, S., and Velde, C.V. (2012). Endogenous TDP-43, but not FUS, contributes to  
738 stress granule assembly via G3BP. *Mol Neurodegener* 7, 1.
- 739 Ayala, Y.M., Zago, P., D’Ambrogio, A., Xu, Y.-F., Petrucelli, L., Buratti, E., and Baralle, F.E.  
740 (2008). Structural determinants of the cellular localization and shuttling of TDP-43. *J Cell Sci* 121,  
741 3778–3785.
- 742 Barmada, S.J., Skibinski, G., Korb, E., Rao, E.J., Wu, J.Y., and Finkbeiner, S. (2010). Cytoplasmic  
743 Mislocalization of TDP-43 Is Toxic to Neurons and Enhanced by a Mutation Associated with  
744 Familial Amyotrophic Lateral Sclerosis. *J Neurosci* 30, 639–649.
- 745 Basso, M., and Pennuto, M. (2015). Serine phosphorylation and arginine methylation at the  
746 crossroads to neurodegeneration. *Exp Neurol* 271, 77–83.
- 747 Beausoleil, S.A., Villén, J., Gerber, S.A., Rush, J., and Gygi, S.P. (2006). A probability-based  
748 approach for high-throughput protein phosphorylation analysis and site localization. *Nat Biotechnol*  
749 24, 1285–1292.
- 750 Bendotti, C., Atzori, C., Piva, R., Tortarolo, M., Strong, M.J., DeBiasi, S., and Migheli, A. (2004).  
751 Activated p38MAPK Is a Novel Component of the Intracellular Inclusions Found in Human  
752 Amyotrophic Lateral Sclerosis and Mutant SOD1 Transgenic Mice. *J Neuropathology Exp*  
753 *Neurology* 63, 113–119.
- 754 Brady, O.A., Meng, P., Zheng, Y., Mao, Y., and Hu, F. (2011). Regulation of TDP-43 aggregation  
755 by phosphorylation and p62/SQSTM1. *J Neurochem* 116, 248–259.
- 756 Brennan, C.M., Emerson, C.P., Owens, J., and Christoforou, N. (2021). p38 MAPKs — roles in  
757 skeletal muscle physiology, disease mechanisms, and as potential therapeutic targets. *Jci Insight* 6,  
758 e149915.



- 759 Buratti, E. (2018). TDP-43 post-translational modifications in health and disease. *Expert Opin Ther*  
760 *Tar* 22, 279–293.
- 761 Buratti, E., and Baralle, F.E. (2008). Multiple roles of TDP-43 in gene expression, splicing  
762 regulation, and human disease. *Front Biosci* 13, 867.
- 763 Buratti, E., Brindisi, A., Giombi, M., Tisminetzky, S., Ayala, Y.M., and Baralle, F.E. (2005). TDP-  
764 43 Binds Heterogeneous Nuclear Ribonucleoprotein A/B through Its C-terminal Tail AN  
765 IMPORTANT REGION FOR THE INHIBITION OF CYSTIC FIBROSIS TRANSMEMBRANE  
766 CONDUCTANCE REGULATOR EXON 9 SPLICING\*. *J Biol Chem* 280, 37572–37584.
- 767 Burton, J.C., Antoniadis, W., Okalova, J., Roos, M.M., and Grimsey, N.J. (2021). Atypical p38  
768 Signaling, Activation, and Implications for Disease. *Int J Mol Sci* 22, 4183.
- 769 Campbell, M., Chang, P.-C., Huerta, S., Izumiya, C., Davis, R., Tepper, C.G., Kim, K.Y.,  
770 Shevchenko, B., Wang, D.-H., Jung, J.U., et al. (2012). Protein Arginine Methyltransferase 1-  
771 directed Methylation of Kaposi Sarcoma-associated Herpesvirus Latency-associated Nuclear  
772 Antigen\*. *J Biol Chem* 287, 5806–5818.
- 773 Carlomagno, Y., Zhang, Y., Davis, M., Lin, W.-L., Cook, C., Dunmore, J., Tay, W., Menkosky, K.,  
774 Cao, X., Petrucelli, L., et al. (2014). Casein Kinase II Induced Polymerization of Soluble TDP-43  
775 into Filaments Is Inhibited by Heat Shock Proteins. *Plos One* 9, e90452.
- 776 Chang, C., Wu, T.-H., Wu, C.-Y., Chiang, M., Toh, E.K.-W., Hsu, Y.-C., Lin, K.-F., Liao, Y.,  
777 Huang, T., and Huang, J.J.-T. (2012). The N-terminus of TDP-43 promotes its oligomerization and  
778 enhances DNA binding affinity. *Biochem Bioph Res Co* 425, 219–224.
- 779 Chang, Y.-I., Hsu, S.-C., Chau, G.-Y., Huang, C.-Y.F., Sung, J.-S., Hua, W.-K., and Lin, W.-J.  
780 (2011). Identification of the methylation preference region in heterogeneous nuclear  
781 ribonucleoprotein K by protein arginine methyltransferase 1 and its implication in regulating  
782 nuclear/cytoplasmic distribution. *Biochem Bioph Res Co* 404, 865–869.
- 783 Choksi, D.K., Roy, B., Chatterjee, S., Yusuff, T., Bakhoun, M.F., Sengupta, U., Ambegaokar, S.,  
784 Kayed, R., and Jackson, G.R. (2014). TDP-43 Phosphorylation by casein kinase I $\epsilon$  promotes  
785 oligomerization and enhances toxicity in vivo. *Hum Mol Genet* 23, 1025–1035.
- 786 Conicella, A.E., Zerze, G.H., Mittal, J., and Fawzi, N.L. (2016). ALS Mutations Disrupt Phase  
787 Separation Mediated by  $\alpha$ -Helical Structure in the TDP-43 Low-Complexity C-Terminal Domain.  
788 *Structure* 24, 1537–1549.
- 789 Cook, C.N., Wu, Y., Odeh, H.M., Gendron, T.F., Jansen-West, K., Rosso, G. del, Yue, M., Jiang,  
790 P., Gomes, E., Tong, J., et al. (2020). C9orf72 poly(GR) aggregation induces TDP-43  
791 proteinopathy. *Sci Transl Med* 12, eabb3774.
- 792 Corrêa, S.A.L., and Eales, K.L. (2012). The Role of p38 MAPK and Its Substrates in Neuronal  
793 Plasticity and Neurodegenerative Disease. *J Signal Transduct* 2012, 649079.
- 794 Cuadrado, A., and Nebreda, A.R. (2010). Mechanisms and functions of p38 MAPK signalling.  
795 *Biochem J* 429, 403–417.



- 796 Cuenda, A., Cohen, P., Buée-Scherrer, V., and Goedert, M. (1997). Activation of stress-activated  
797 protein kinase-3 (SAPK3) by cytokines and cellular stresses is mediated via SAPKK3 (MKK6);  
798 comparison of the specificities of SAPK3 and SAPK2 (RK/p38). *Embo J* 16, 295–305.
- 799 Cupo, R., and Shorter, J. (2020). Expression and Purification of Recombinant Skd3 (Human ClpB)  
800 Protein and Tobacco Etch Virus (TEV) Protease from *Escherichia coli*. *Bio-Protocol* 10, e3858.
- 801 Dewil, M., Cruz, V.F. dela, Bosch, L.V.D., and Robberecht, W. (2007). Inhibition of p38 mitogen  
802 activated protein kinase activation and mutant SOD1G93A-induced motor neuron death. *Neurobiol*  
803 *Dis* 26, 332–341.
- 804 Diskin, R., Askari, N., Capone, R., Engelberg, D., and Livnah, O. (2004). Active Mutants of the  
805 Human p38 $\alpha$  Mitogen-activated Protein Kinase\*. *J Biol Chem* 279, 47040–47049.
- 806 Duffy, J.P., Harrington, E.M., Salituro, F.G., Cochran, J.E., Green, J., Gao, H., Bemis, G.W.,  
807 Evindar, G., Galullo, V.P., Ford, P.J., et al. (2011). The Discovery of VX-745: A Novel and  
808 Selective p38 $\alpha$  Kinase Inhibitor. *Acs Med Chem Lett* 2, 758–763.
- 809 Eng, J.K., McCormack, A.L., and Yates, J.R. (1994). An approach to correlate tandem mass  
810 spectral data of peptides with amino acid sequences in a protein database. *J Am Soc Mass Spectr* 5,  
811 976–989.
- 812 Freibaum, B.D., Chitta, R.K., High, A.A., and Taylor, J.P. (2010). Global Analysis of TDP-43  
813 Interacting Proteins Reveals Strong Association with RNA Splicing and Translation Machinery. *J*  
814 *Proteome Res* 9, 1104–1120.
- 815 Gasset-Rosa, F., Lu, S., Yu, H., Chen, C., Melamed, Z., Guo, L., Shorter, J., Cruz, S.D., and  
816 Cleveland, D.W. (2019). Cytoplasmic TDP-43 De-mixing Independent of Stress Granules Drives  
817 Inhibition of Nuclear Import, Loss of Nuclear TDP-43, and Cell Death. *Neuron* 102, 339-357.e7.
- 818 Germann, U.A., and Alam, J.J. (2020). P38 $\alpha$  MAPK Signaling—A Robust Therapeutic Target for  
819 Rab5-Mediated Neurodegenerative Disease. *Int J Mol Sci* 21, 5485.
- 820 Gibbs, K.L., Kalmar, B., Rhymes, E.R., Fellows, A.D., Ahmed, M., Whiting, P., Davies, C.H.,  
821 Greensmith, L., and Schiavo, G. (2018). Inhibiting p38 MAPK alpha rescues axonal retrograde  
822 transport defects in a mouse model of ALS. *Cell Death Dis* 9, 596.
- 823 Gitcho, M.A., Bigio, E.H., Mishra, M., Johnson, N., Weintraub, S., Mesulam, M., Rademakers, R.,  
824 Chakraverty, S., Cruchaga, C., Morris, J.C., et al. (2009). TARDBP 3'-UTR variant in autopsy-  
825 confirmed frontotemporal lobar degeneration with TDP-43 proteinopathy. *Acta Neuropathol* 118,  
826 633.
- 827 Goh, C.W., Lee, I.C., Sundaram, J.R., George, S.E., Yusoff, P., Brush, M.H., Sze, N.S.K., and  
828 Shenolikar, S. (2018). Chronic oxidative stress promotes GADD34-mediated phosphorylation of the  
829 TAR DNA-binding protein TDP-43, a modification linked to neurodegeneration. *J Biol Chem* 293,  
830 163–176.
- 831 Goldstein, D.M., Soth, M., Gabriel, T., Dewdney, N., Kuglstatter, A., Arzeno, H., Chen, J.,  
832 Bingenheimer, W., Dalrymple, S.A., Dunn, J., et al. (2011). Discovery of 6-(2,4-Difluorophenoxy)-  
833 2-[3-hydroxy-1-(2-hydroxyethyl)propylamino]-8-methyl-8 H -pyrido[2,3- d ]pyrimidin-7-one

- 834 (Pamapimod) and 6-(2,4-Difluorophenoxy)-8-methyl-2-(tetrahydro-2 H -pyran-4-  
835 ylamino)pyrido[2,3- d ]pyrimidin-7(8 H )-one (R1487) as Orally Bioavailable and Highly Selective  
836 Inhibitors of p38 $\alpha$  Mitogen-Activated Protein Kinase. *J Med Chem* 54, 2255–2265.
- 837 Guo, L., and Shorter, J. (2017). Biology and Pathobiology of TDP-43 and Emergent Therapeutic  
838 Strategies. *Csh Perspect Med* 7, a024554.
- 839 Guo, A., Gu, H., Zhou, J., Mulhern, D., Wang, Y., Lee, K.A., Yang, V., Aguiar, M., Kornhauser, J.,  
840 Jia, X., et al. (2014). Immunoaffinity Enrichment and Mass Spectrometry Analysis of Protein  
841 Methylation. *Mol Cell Proteomics* 13, 372–387.
- 842 Guo, Z., Zheng, L., Xu, H., Dai, H., Zhou, M., Pascua, M.R., Chen, Q.M., and Shen, B. (2010).  
843 Methylation of FEN1 suppresses nearby phosphorylation and facilitates PCNA binding. *Nat Chem*  
844 *Biol* 6, 766–773.
- 845 Harrison, A.F., and Shorter, J. (2017). RNA-binding proteins with prion-like domains in health and  
846 disease. *Biochem J* 474, 1417–1438.
- 847 Hasegawa, M., Arai, T., Nonaka, T., Kametani, F., Yoshida, M., Hashizume, Y., Beach, T.G.,  
848 Buratti, E., Baralle, F., Morita, M., et al. (2008). Phosphorylated TDP-43 in frontotemporal lobar  
849 degeneration and amyotrophic lateral sclerosis. *Ann Neurol* 64, 60–70.
- 850 Hsu, J.-M., Chen, C.-T., Chou, C.-K., Kuo, H.-P., Li, L.-Y., Lin, C.-Y., Lee, H.-J., Wang, Y.-N.,  
851 Liu, M., Liao, H.-W., et al. (2011). Crosstalk between Arg 1175 methylation and Tyr 1173  
852 phosphorylation negatively modulates EGFR-mediated ERK activation. *Nat Cell Biol* 13, 174–181.
- 853 Huang, L., Wang, Z., Narayanan, N., and Yang, Y. (2018). Arginine methylation of the C-terminus  
854 RGG motif promotes TOP3B topoisomerase activity and stress granule localization. *Nucleic Acids*  
855 *Res* 46, gky103-.
- 856 Huq, M.D.M., Gupta, P., Tsai, N., White, R., Parker, M.G., and Wei, L. (2006). Suppression of  
857 receptor interacting protein 140 repressive activity by protein arginine methylation. *Embo J* 25,  
858 5094–5104.
- 859 Jiang, L.-L., Xue, W., Hong, J.-Y., Zhang, J.-T., Li, M.-J., Yu, S.-N., He, J.-H., and Hu, H.-Y.  
860 (2017). The N-terminal dimerization is required for TDP-43 splicing activity. *Sci Rep-Uk* 7, 6196.
- 861 Johnson, B.S., Snead, D., Lee, J.J., McCaffery, J.M., Shorter, J., and Gitler, A.D. (2009). TDP-43 Is  
862 Intrinsically Aggregation-prone, and Amyotrophic Lateral Sclerosis-linked Mutations Accelerate  
863 Aggregation and Increase Toxicity\*. *J Biol Chem* 284, 20329–20339.
- 864 Kametani, F., Nonaka, T., Suzuki, T., Arai, T., Dohmae, N., Akiyama, H., and Hasegawa, M.  
865 (2009). Identification of casein kinase-1 phosphorylation sites on TDP-43. *Biochem Bioph Res Co*  
866 382, 405–409.
- 867 Kametani, F., Obi, T., Shishido, T., Akatsu, H., Murayama, S., Saito, Y., Yoshida, M., and  
868 Hasegawa, M. (2016). Mass spectrometric analysis of accumulated TDP-43 in amyotrophic lateral  
869 sclerosis brains. *Sci Rep-Uk* 6, 23281.

- 870 Kawahara, Y., and Mieda-Sato, A. (2012). TDP-43 promotes microRNA biogenesis as a component  
871 of the Drosha and Dicer complexes. *Proc National Acad Sci* *109*, 3347–3352.
- 872 Khalfallah, Y., Kuta, R., Grasmuck, C., Prat, A., Durham, H.D., and Velde, C.V. (2018). TDP-43  
873 regulation of stress granule dynamics in neurodegenerative disease-relevant cell types. *Sci Rep-Uk*  
874 *8*, 7551.
- 875 Kim, K.Y., Lee, H.-W., Shim, Y., Mook-Jung, I., Jeon, G.S., and Sung, J.-J. (2015). A  
876 phosphomimetic mutant TDP-43 (S409/410E) induces Drosha instability and cytotoxicity in Neuro  
877 2A cells. *Biochem Bioph Res Co* *464*, 236–243.
- 878 Lalmansingh, A.S., Urekar, C.J., and Reddi, P.P. (2011). TDP-43 Is a Transcriptional Repressor  
879 THE TESTIS-SPECIFIC MOUSE *acr1* GENE IS A TDP-43 TARGET IN VIVO\*. *J Biol Chem*  
880 *286*, 10970–10982.
- 881 Larsen, S.C., Sylvestersen, K.B., Mund, A., Lyon, D., Mullari, M., Madsen, M.V., Daniel, J.A.,  
882 Jensen, L.J., and Nielsen, M.L. (2016). Proteome-wide analysis of arginine monomethylation  
883 reveals widespread occurrence in human cells. *Sci Signal* *9*, rs9–rs9.
- 884 Li, H.-R., Chen, T.-C., Hsiao, C.-L., Shi, L., Chou, C.-Y., and Huang, J. (2018). The physical forces  
885 mediating self-association and phase-separation in the C-terminal domain of TDP-43. *Biochimica*  
886 *Et Biophysica Acta Bba - Proteins Proteom* *1866*, 214–223.
- 887 Li, H.-Y., Yeh, P.-A., Chiu, H.-C., Tang, C.-Y., and Tu, B.P. (2011). Hyperphosphorylation as a  
888 Defense Mechanism to Reduce TDP-43 Aggregation. *Plos One* *6*, e23075.
- 889 Liachko, N.F., Guthrie, C.R., and Kraemer, B.C. (2010). Phosphorylation Promotes Neurotoxicity  
890 in a *Caenorhabditis elegans* Model of TDP-43 Proteinopathy. *J Neurosci* *30*, 16208–16219.
- 891 Liachko, N.F., McMillan, P.J., Guthrie, C.R., Bird, T.D., Leverenz, J.B., and Kraemer, B.C. (2013).  
892 CDC7 inhibition blocks pathological TDP-43 phosphorylation and neurodegeneration. *Ann Neurol*  
893 *74*, 39–52.
- 894 Liachko, N.F., McMillan, P.J., Strovas, T.J., Loomis, E., Greenup, L., Murrell, J.R., Ghetti, B.,  
895 Raskind, M.A., Montine, T.J., Bird, T.D., et al. (2014). The Tau Tubulin Kinases TTBK1/2  
896 Promote Accumulation of Pathological TDP-43. *Plos Genet* *10*, e1004803.
- 897 Ling, S.-C., Albuquerque, C.P., Han, J.S., Lagier-Tourenne, C., Tokunaga, S., Zhou, H., and  
898 Cleveland, D.W. (2010). ALS-associated mutations in TDP-43 increase its stability and promote  
899 TDP-43 complexes with FUS/TLS. *Proc National Acad Sci* *107*, 13318–13323.
- 900 Liu, L., Sun, W., Fan, X., Xu, Y., Cheng, M., and Zhang, Y. (2019). Methylation of C/EBP $\alpha$  by  
901 PRMT1 inhibits its tumor suppressive function in breast cancer. *Cancer Res* *79*, canres.3211.2018.
- 902 Liu, Y.-J., Ju, T.-C., Chen, H.-M., Jang, Y.-S., Lee, L.-M., Lai, H.-L., Tai, H.-C., Fang, J.-M., Lin,  
903 Y.-L., Tu, P.-H., et al. (2015). Activation of AMP-activated protein kinase  $\alpha$ 1 mediates  
904 mislocalization of TDP-43 in amyotrophic lateral sclerosis. *Hum Mol Genet* *24*, 787–801.


- 905 Lu, Y., Ma, W., Li, Z., Lu, J., and Wang, X. (2017). The interplay between p16 serine  
906 phosphorylation and arginine methylation determines its function in modulating cellular apoptosis  
907 and senescence. *Sci Rep-Uk* 7, 41390.
- 908 Mann, J.R., Gleixner, A.M., Mauna, J.C., Gomes, E., DeChellis-Marks, M.R., Needham, P.G.,  
909 Copley, K.E., Hurtle, B., Portz, B., Pyles, N.J., et al. (2019). RNA Binding Antagonizes Neurotoxic  
910 Phase Transitions of TDP-43. *Neuron* 102, 321-338.e8.
- 911 McDonald, K.K., Aulas, A., Destroismaisons, L., Pickles, S., Beleac, E., Camu, W., Rouleau, G.A.,  
912 and Velde, C.V. (2011). TAR DNA-binding protein 43 (TDP-43) regulates stress granule dynamics  
913 via differential regulation of G3BP and TIA-1. *Hum Mol Genet* 20, 1400–1410.
- 914 McGurk, L., Gomes, E., Guo, L., Mojsilovic-Petrovic, J., Tran, V., Kalb, R.G., Shorter, J., and  
915 Bonini, N.M. (2018). Poly(ADP-Ribose) Prevents Pathological Phase Separation of TDP-43 by  
916 Promoting Liquid Demixing and Stress Granule Localization. *Mol Cell* 71, 703-717.e9.
- 917 Meyerowitz, J., Parker, S.J., Vella, L.J., Ng, D.C., Price, K.A., Liddell, J.R., Caragounis, A., Li, Q.-  
918 X., Masters, C.L., Nonaka, T., et al. (2011). C-Jun N-terminal kinase controls TDP-43 accumulation  
919 in stress granules induced by oxidative stress. *Mol Neurodegener* 6, 57.
- 920 Molliex, A., Temirov, J., Lee, J., Coughlin, M., Kanagaraj, A.P., Kim, H.J., Mittag, T., and Taylor,  
921 J.P. (2015). Phase Separation by Low Complexity Domains Promotes Stress Granule Assembly and  
922 Drives Pathological Fibrillization. *Cell* 163, 123–133.
- 923 Morrison, D.K. (2012). MAP Kinase Pathways. *Csh Perspect Biol* 4, a011254.
- 924 Neumann, M., Sampathu, D.M., Kwong, L.K., Truax, A.C., Micsenyi, M.C., Chou, T.T., Bruce, J.,  
925 Schuck, T., Grossman, M., Clark, C.M., et al. (2006). Ubiquitinated TDP-43 in Frontotemporal  
926 Lobar Degeneration and Amyotrophic Lateral Sclerosis. *Science* 314, 130–133.
- 927 Neumann, M., Kwong, L.K., Lee, E.B., Kremmer, E., Flatley, A., Xu, Y., Forman, M.S., Troost, D.,  
928 Kretzschmar, H.A., Trojanowski, J.Q., et al. (2009). Phosphorylation of S409/410 of TDP-43 is a  
929 consistent feature in all sporadic and familial forms of TDP-43 proteinopathies. *Acta Neuropathol*  
930 117, 137–149.
- 931 Nguyen, H.P., Broeckhoven, C.V., and Zee, J. van der (2018). ALS Genes in the Genomic Era and  
932 their Implications for FTD. *Trends Genet* 34, 404–423.
- 933 Nonaka, T., Arai, T., Buratti, E., Baralle, F.E., Akiyama, H., and Hasegawa, M. (2009).  
934 Phosphorylated and ubiquitinated TDP-43 pathological inclusions in ALS and FTL-DU are  
935 recapitulated in SH-SY5Y cells. *Febs Lett* 583, 394–400.
- 936 Nonaka, T., Suzuki, G., Tanaka, Y., Kametani, F., Hirai, S., Okado, H., Miyashita, T., Saitoe, M.,  
937 Akiyama, H., Masai, H., et al. (2016). Phosphorylation of TAR DNA-binding Protein of 43 kDa  
938 (TDP-43) by Truncated Casein Kinase 1 $\delta$  Triggers Mislocalization and Accumulation of TDP-43\*.  
939 *J Biol Chem* 291, 5473–5483.
- 940 Ortega, J.A., Daley, E.L., Kour, S., Samani, M., Tellez, L., Smith, H.S., Hall, E.A., Esengul, Y.T.,  
941 Tsai, Y.-H., Gendron, T.F., et al. (2020). Nucleocytoplasmic Proteomic Analysis Uncovers eRF1

- 942 and Nonsense-Mediated Decay as Modifiers of ALS/FTD C9orf72 Toxicity. *Neuron* *106*, 90-  
943 107.e13.
- 944 Peng, J., and Gygi, S.P. (2001). Proteomics: the move to mixtures. *J Mass Spectrom* *36*, 1083–  
945 1091.
- 946 Pickhardt, M., Tassoni, M., Denner, P., Kurkowsky, B., Kitanovic, A., Möhl, C., Fava, E., and  
947 Mandelkow, E. (2019). Screening of a neuronal cell model of tau pathology for therapeutic  
948 compounds. *Neurobiol Aging* *76*, 24–34.
- 949 Polymenidou, M., Lagier-Tourenne, C., Hutt, K.R., Huelga, S.C., Moran, J., Liang, T.Y., Ling, S.-  
950 C., Sun, E., Wancewicz, E., Mazur, C., et al. (2011). Long pre-mRNA depletion and RNA  
951 missplicing contribute to neuronal vulnerability from loss of TDP-43. *Nat Neurosci* *14*, 459–468.
- 952 Portz, B., Lee, B.L., and Shorter, J. (2021). FUS and TDP-43 Phases in Health and Disease. *Trends*  
953 *Biochem Sci* *46*, 550–563.
- 954 Prasad, A., Bharathi, V., Sivalingam, V., Girdhar, A., and Patel, B.K. (2019). Molecular  
955 Mechanisms of TDP-43 Misfolding and Pathology in Amyotrophic Lateral Sclerosis. *Front Mol*  
956 *Neurosci* *12*, 25.
- 957 Prins, N.D., Harrison, J.E., Chu, H.-M., Blackburn, K., Alam, J.J., Scheltens, P., Arnold, Coskinas,  
958 Gonzales, Joseph, et al. (2021). A phase 2 double-blind placebo-controlled 24-week treatment  
959 clinical study of the p38 alpha kinase inhibitor neflamapimod in mild Alzheimer’s disease.  
960 *Alzheimer’s Res Ther* *13*, 106.
- 961 Qamar, S., Wang, G., Randle, S.J., Ruggeri, F.S., Varela, J.A., Lin, J.Q., Phillips, E.C., Miyashita,  
962 A., Williams, D., Ströhl, F., et al. (2018). FUS Phase Separation Is Modulated by a Molecular  
963 Chaperone and Methylation of Arginine Cation- $\pi$  Interactions. *Cell* *173*, 720-734.e15.
- 964 Ratovitski, T., Arbez, N., Stewart, J.C., Chighladze, E., and Ross, C.A. (2015). PRMT5- mediated  
965 symmetric arginine dimethylation is attenuated by mutant huntingtin and is impaired in  
966 Huntington’s disease (HD). *Cell Cycle* *14*, 1716–1729.
- 967 Rowland, L.P., and Shneider, N.A. (2001). Amyotrophic Lateral Sclerosis. *New Engl J Medicine*  
968 *344*, 1688–1700.
- 969 Shevchenko, A., Wilm, M., Vorm, O., and Mann, M. (1996). Mass Spectrometric Sequencing of  
970 Proteins from Silver-Stained Polyacrylamide Gels. *Anal Chem* *68*, 850–858.
- 971 Silva, L.G. da, Simonetti, F., Hutten, S., Riemenschneider, H., Sternburg, E.L., Pietrek, L.M.,  
972 Gebel, J., Dötsch, V., Edbauer, D., Hummer, G., et al. (2021). Disease-linked TDP-43  
973 hyperphosphorylation suppresses TDP-43 condensation and aggregation. *Biorxiv*  
974 2021.04.30.442163.
- 975 Simandi, Z., Pajer, K., Karolyi, K., Sieler, T., Jiang, L.-L., Kolostyak, Z., Sari, Z., Fekecs, Z., Pap,  
976 A., Patsalos, A., et al. (2018). Arginine Methyltransferase PRMT8 Provides Cellular Stress  
977 Tolerance in Aging Motoneurons. *J Neurosci* *38*, 7683–7700.



- 978 Sreedharan, J., Blair, I.P., Tripathi, V.B., Hu, X., Vance, C., Rogelj, B., Ackerley, S., Durnall, J.C.,  
979 Williams, K.L., Buratti, E., et al. (2008). TDP-43 Mutations in Familial and Sporadic Amyotrophic  
980 Lateral Sclerosis. *Science* *319*, 1668–1672.
- 981 Sreedharan, J., Neukomm, L.J., Brown, R.H., and Freeman, M.R. (2015). Age-Dependent TDP-43-  
982 Mediated Motor Neuron Degeneration Requires GSK3, hat-trick, and xmas-2. *Curr Biol* *25*, 2130–  
983 2136.
- 984 Tang, J., Frankel, A., Cook, R.J., Kim, S., Paik, W.K., Williams, K.R., Clarke, S., and Herschman,  
985 H.R. (2000). PRMT1 Is the Predominant Type I Protein Arginine Methyltransferase in Mammalian  
986 Cells\*. *J Biol Chem* *275*, 7723–7730.
- 987 Taylor, L.M., McMillan, P.J., Liachko, N.F., Strovass, T.J., Ghetti, B., Bird, T.D., Keene, C.D., and  
988 Kraemer, B.C. (2018). Pathological phosphorylation of tau and TDP-43 by TTBK1 and TTBK2  
989 drives neurodegeneration. *Mol Neurodegener* *13*, 7.
- 990 Thandapani, P., O’Connor, T.R., Bailey, T.L., and Richard, S. (2013). Defining the RGG/RG Motif.  
991 *Mol Cell* *50*, 613–623.
- 992 Tollervey, J.R., Curk, T., Rogelj, B., Briese, M., Cereda, M., Kayikci, M., König, J., Hortobágyi,  
993 T., Nishimura, A.L., Župunski, V., et al. (2011). Characterizing the RNA targets and position-  
994 dependent splicing regulation by TDP-43. *Nat Neurosci* *14*, 452–458.
- 995 Tortarolo, M., Veglianese, P., Calvaresi, N., Botturi, A., Rossi, C., Giorgini, A., Migheli, A., and  
996 Bendotti, C. (2003). Persistent activation of p38 mitogen-activated protein kinase in a mouse model  
997 of familial amyotrophic lateral sclerosis correlates with disease progression. *Mol Cell Neurosci* *23*,  
998 180–192.
- 999 Valdmanis, P.N., and Rouleau, G.A. (2008). Genetics of familial amyotrophic lateral sclerosis.  
1000 *Neurology* *70*, 144–152.
- 1001 Wall, M.L., and Lewis, S.M. (2017). Methylarginines within the RGG-Motif Region of hnRNP A1  
1002 Affect Its IRES Trans-Acting Factor Activity and Are Required for hnRNP A1 Stress Granule  
1003 Localization and Formation. *J Mol Biol* *429*, 295–307.
- 1004 Wang, A., Conicella, A.E., Schmidt, H.B., Martin, E.W., Rhoads, S.N., Reeb, A.N., Nourse, A.,  
1005 Montero, D.R., Ryan, V.H., Rohatgi, R., et al. (2018). A single N-terminal phosphomimic disrupts  
1006 TDP-43 polymerization, phase separation, and RNA splicing. *Embo J* *37*.
- 1007 Watkins, J., Ghosh, A., Keerie, A.F.A., Alix, J.J.P., Mead, R.J., and Sreedharan, J. (2020). Female  
1008 sex mitigates motor and behavioural phenotypes in TDP-43Q331K knock-in mice. *Sci Rep-Uk* *10*,  
1009 19220.
- 1010 Winton, M.J., Igaz, L.M., Wong, M.M., Kwong, L.K., Trojanowski, J.Q., and Lee, V.M.-Y. (2008).  
1011 Disturbance of Nuclear and Cytoplasmic TAR DNA-binding Protein (TDP-43) Induces Disease-  
1012 like Redistribution, Sequestration, and Aggregate Formation\*. *J Biol Chem* *283*, 13302–13309.
- 1013 Winzen, R., Kracht, M., Ritter, B., Wilhelm, A., Chen, C.A., Shyu, A., Müller, M., Gaestel, M.,  
1014 Resch, K., and Holtmann, H. (1999). The p38 MAP kinase pathway signals for cytokine-induced


- 1015 mRNA stabilization via MAP kinase-activated protein kinase 2 and an AU-rich region-targeted  
1016 mechanism. *Embo J* *18*, 4969–4980.
- 1017 Wobst, H.J., Wesolowski, S.S., Chadchankar, J., Delsing, L., Jacobsen, S., Mukherjee, J., Deeb,  
1018 T.Z., Dunlop, J., Brandon, N.J., and Moss, S.J. (2017). Cytoplasmic Relocalization of TAR DNA-  
1019 Binding Protein 43 Is Not Sufficient to Reproduce Cellular Pathologies Associated with ALS In  
1020 vitro. *Frontiers in Molecular Neuroscience* *10*, 46.
- 1021 Wobst, H.J., Mack, K.L., Brown, D.G., Brandon, N.J., and Shorter, J. (2020). The clinical trial  
1022 landscape in amyotrophic lateral sclerosis—Past, present, and future. *Med Res Rev* *40*, 1352–1384.
- 1023 Xiong, H.-L., Wang, J.-Y., Sun, Y.-M., Wu, J.-J., Chen, Y., Qiao, K., Zheng, Q.-J., Zhao, G., and  
1024 Wu, Z.-Y. (2010). Association between novel TARDBP mutations and Chinese patients with  
1025 amyotrophic lateral sclerosis. *Bmc Med Genet* *11*, 8.
- 1026 Xu, Y.-F., Zhang, Y.-J., Lin, W.-L., Cao, X., Stetler, C., Dickson, D.W., Lewis, J., and Petrucelli,  
1027 L. (2011). Expression of mutant TDP-43 induces neuronal dysfunction in transgenic mice. *Mol*  
1028 *Neurodegener* *6*, 73.
- 1029 Yamagata, K., Daitoku, H., Takahashi, Y., Namiki, K., Hisatake, K., Kako, K., Mukai, H., Kasuya,  
1030 Y., and Fukamizu, A. (2008). Arginine Methylation of FOXO Transcription Factors Inhibits Their  
1031 Phosphorylation by Akt. *Mol Cell* *32*, 221–231.
- 1032 Yasuda, S., Sugiura, H., Tanaka, H., Takigami, S., and Yamagata, K. (2011). p38 MAP Kinase  
1033 Inhibitors as Potential Therapeutic Drugs for Neural Diseases. *Central Nerv Syst Agents Medicinal*  
1034 *Chem* *11*, 45–59.
- 1035 Zarubin, T., and Han, J. (2005). Activation and signaling of the p38 MAP kinase pathway. *Cell Res*  
1036 *15*, 11–18.
- 1037 Zhan, L., Xie, Q., and Tibbetts, R.S. (2015). Opposing roles of p38 and JNK in a *Drosophila* model  
1038 of TDP-43 proteinopathy reveal oxidative stress and innate immunity as pathogenic components of  
1039 neurodegeneration. *Hum Mol Genet* *24*, 757–772.
- 1040 Zhang, Y.-J., Gendron, T.F., Xu, Y.-F., Ko, L.-W., Yen, S.-H., and Petrucelli, L. (2010).  
1041 Phosphorylation regulates proteasomal-mediated degradation and solubility of TAR DNA binding  
1042 protein-43 C-terminal fragments. *Mol Neurodegener* *5*, 33.
- 1043 Zou, Z.-Y., Peng, Y., Wang, X.-N., Liu, M.-S., Li, X.-G., and Cui, L.-Y. (2012). Screening of the  
1044 TARDBP gene in familial and sporadic amyotrophic lateral sclerosis patients of Chinese origin.  
1045 *Neurobiol Aging* *33*, 2229.e11-2229.e18.
- 1046



1047


1048 **Figure 1. Genetic and pharmacological inhibition of p38α MAPK reduces TDP-43 aggregation**  
 1049 **and phosphorylation at S409/S410, and inhibits TDP-43-induced neurotoxicity.** **A)** Schematic of  
 1050 domain architecture of TDP-43 and location of ALS-linked mutations and phosphorylation sites (P)  
 1051 detected after p38α MAPK treatment *in vitro*. **B)** Western blot of total and phosphorylated TDP-  
 1052 43<sup>M337V</sup> in RIPA and urea fractions of SH-SY5Y cells with siRNA-induced p38α knockdown.  
 1053 GAPDH was used as a loading control. **C)** Quantification of urea/RIPA ratio of total TDP-43  
 1054 normalized to levels in scrambled siRNA (si-scr). **D)** Quantification of pTDP-43/total TDP-43 ratio  
 1055 in urea fraction normalized to levels in si-scr (mean band signal ± SD, two-way ANOVA with Sidak's  
 1056 multiple comparison test, n = 3). **E)** Western blot of total and pTDP-43<sup>M337V</sup> in RIPA and urea  
 1057 fractions of SH-SY5Y cells with pharmacological p38α inhibition with compound 1. GAPDH was  
 1058 used as a loading control. Quantification of urea/RIPA ratio of total TDP-43 at time points 24h (**F**)  
 1059 and 48h (**G**) post-transfection, and pTDP-43/total TDP-43 ratio in urea fraction at time point 48h  
 1060 post-transfection (**H**) normalized to levels in DMSO-treated cells (mean band signal ± SD, one-way  
 1061 ANOVA with Dunnett's multiple comparison test, n = 3). **I)** Quantification of LDH activity in  
 1062 conditioned medium normalized to levels in DMSO-treated TDP-43<sup>WT</sup>-transfected NSC-34 cells.  
 1063 (one-way ANOVA with Dunnett's multiple comparison test, n = 4 with 6 replicates in each). \*p <  
 1064 0.05, \*\*p < 0.01, \*\*\*p < 0.001, \*\*\*\*p < 0.0001. **J)** Hazard ratios of primary neurons expressing mApple  
 1065 and TDP-43<sup>M337V</sup>-EGFP treated with p38α inhibitor VX-745 at 0.3, 1, 3, and 9 μM compared with  
 1066 DMSO control (reference, set at 1.0) were 0.6296, 0.6378, 0.4596, and 0.6869 respectively.  
 1067 Reduction of hazard ratio was most significant at 3 μM (Cox proportional hazard, p < 0.01).  
 1068 See also Figure S1.  
 1069





1070

1071 **Figure 2. Constitutively active p38α induces aggregation, phosphorylation and cytoplasmic**  
 1072 **accumulation of TDP-43.** **A)** Western blot of total and pTDP-43 in RIPA and urea fractions of SH-  
 1073 SY5Y cells co-transfected with TDP-43<sup>M337V</sup> and empty control plasmid (Ctrl), WT-p38α, CA-p38α  
 1074 or DN-p38α. GAPDH was used as a loading control. Quantification of urea/RIPA ratio of total TDP-  
 1075 43 at time points 24h (**B**) or 48h (**C**) post-transfection, and pTDP-43/total TDP-43 ratio in urea  
 1076 fraction at time point 48h post-transfection (**D**), normalized to levels in ctrl-plasmid transfected cells  
 1077 (mean band signal ± SD, one-way ANOVA with Dunnett's multiple comparison test, n = 3). **E)**  
 1078 Representative confocal images of SH-SY5Y cells co-expressing TDP-43<sup>M337V</sup> (green) and WT-  
 1079 p38α, CA-p38α or DN-p38α (red) with quantification of the number of cells with nuclear TDP-43  
 1080 puncta/granules (**F**) (one-way ANOVA with Dunnett's multiple comparison test, n = 4 and 10 fields  
 1081 each, scale bar 20 μm). **G)** Western blot of TDP-43 in different cellular compartments of SH-SY5Y  
 1082 cells co-transfected with TDP-43<sup>M337V</sup> and ctrl-plasmid or CA-p38α. Exo denotes exogenous TDP-  
 1083 43 and endo denotes endogenous TDP-43. GAPDH was used as a loading control for cytoplasmic  
 1084 fraction (CP), COX IV for the membrane fraction (Mem), and histone H3 for the soluble nuclear (NS)  
 1085 and chromatin-bound (CB) fractions. **H)** Quantification of total TDP-43 in cytoplasmic, membrane,  
 1086 soluble nuclear and chromatin-bound fractions (mean band signal ± SD, two-way ANOVA with  
 1087 Sidak's multiple comparison test, n = 3). \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001, \*\*\*\*p < 0.0001.  
 1088




1089

1090 **Figure 3. TDP-43 directly interacts with and is phosphorylated by p38 $\alpha$  at S292 and S409/S410,**  
 1091 **with S292 regulating phosphorylation at S409/S410. A)** Western blot of immunoprecipitated TDP-  
 1092 43<sup>WT</sup> and co-immunoprecipitated WT-p38 $\alpha$  and immunoprecipitated WT-p38 $\alpha$  and co-  
 1093 immunoprecipitated TDP-43<sup>WT</sup> in SH-SY5Y cells. Inputs are shown on the right. GAPDH was used  
 1094 as a loading control. **B)** Western blot of *in vitro* kinase assay (30°C for 30 min) of recombinant human  
 1095 TDP-43 (750 ng) and recombinant active p38 $\alpha$  (300 ng) with and without the p38 $\alpha$  inhibitor  
 1096 compound 1. **C)** Western blot of total and pTDP-43 in RIPA and urea fractions of SH-SY5Y cells  
 1097 transfected with TDP-43<sup>WT</sup> and with S292 and S409/S410 mutant constructs. Quantification of  
 1098 urea/RIPA ratio of total TDP-43 (**D**), and CTF/full length ratio of TDP-43 in RIPA fraction (**E**)  
 1099 at time point 24h post-transfection normalized to levels in TDP-43<sup>WT</sup> (mean band signal  $\pm$  SD, one-way  
 1100 ANOVA with Dunnett's multiple comparison test, n = 3). \*p < 0.05.  
 1101 See also Figure S2.  
 1102


A)

*Homo sapiens* 287-GGFGN**SRGG**GAGLGNNQGSNMGGG-310  
*Pongo abelii* 287-GGFGN**SRGG**GAGLGNNQGSNMGGG-310  
*Bos taurus* 287-GGFGN**SRGG**GAGLGNNQGSNMGGG-310  
*Mus musculus* 287-GGFGN**SRGG**GAGLGNNQGSNMGGG-310  
*Gallus gallus* 287-GGFGN**SRGG**GGGLGNNQGSNMGGG-310


B)




C)



D)



E)




1103

1104 **Figure 4. TDP-43 is methylated by PRMT1 primarily at residue R293.** A) Sequence alignment of amino acids 287-310 of TDP-43 from diverse vertebrate species. Conserved S292 and R293 sites are bolded. S292 phosphorylation site and R293-G295 RGG-motif are highlighted in yellow and green, respectively. B) Western blot of immunoprecipitated endogenous TDP-43 from SH-SY5Y-cells probed with antibodies against TDP-43, ADMA, MMA and SDMA. TDP-43 with arginine methylation is marked by arrows. C) Western blot of immunoprecipitated endogenous TDP-43 from SH-SY5Y-cells with siRNA-induced PRMT1 knockdown probed with antibodies against TDP-43, MMA and PRMT1. D) Western blot of expressed and immunoprecipitated TDP-43<sup>WT</sup> using anti-Flag antibody from SH-SY5Y-cells treated with DMSO (D) or with methyltransferase inhibitor AdOx (A) at a final concentration of 20  $\mu$ M for 24h. FL denotes full-length TDP-43 and CTF<sub>35</sub> denotes C-terminal 35kDa fragment of TDP-43. E) Western blot of expressed and immunoprecipitated TDP-43<sup>WT</sup> and TDP-43<sup>R293K</sup> using anti-Flag antibody from SH-SY5Y cells probed with antibodies against TDP-43 and MMA.


1117

1118



1119  
 1120 **Figure 5. TDP-43 phosphorylation favors aggregation whereas arginine methylation favors**  
 1121 **LLPS *in vitro*.** **A)** Representative DIC microscopy images of liquid-like droplets of 10 $\mu\text{M}$  TDP-43-  
 1122 MBP wild-type protein and its mutant forms. Purified recombinant proteins were incubated for 30  
 1123 minutes with phase separation buffer prior to imaging. Scale bar = 10  $\mu\text{m}$  (n=3). **B)** Bar graph  
 1124 showing the total droplets area for each protein. Mean  $\pm$  SEM, one-way ANOVA with Dunnett's  
 1125 multiple comparison test (n=3, \*p<0.05). **C)** Vertical scatterplot displaying the size distribution of  
 1126 droplets for each protein. Each data point corresponds to a single droplet. Bolded bars represent the  
 1127 average droplet area for each variant. **D)** Turbidity measurements of 5  $\mu\text{M}$  TDP-43-MBP co-  
 1128 incubated with TEV protease (1  $\mu\text{g}/\text{mL}$ ). Turbidity was measured at an absorbance of 395 nm. Values  
 1129 represent the normalized mean  $\pm$  SEM (n = 4). **E)** Aggregation data from **D** was quantified by  
 1130 calculating the area under the curve. Values represent means  $\pm$  SEM (n = 4). One-way ANOVA with

1131 Tukey's multiple comparison test was performed (\* $p < 0.05$ ). **F)** LLPS versus aggregation plot shows  
1132 the phospho-mimetics cluster below the  $y=x$  dotted line, whereas the arginine methylation-mimic  
1133 appears above the  $y=x$  dotted line. Phospho-mimetics inhibit TDP-43 LLPS more severely than TDP-  
1134 43 aggregation, whereas the arginine-methylation mimic inhibits TDP-43 aggregation but not TDP-  
1135 43 LLPS. Y-axis represents normalized LLPS propensity relative to wild-type based on the average  
1136 area of droplets from analysis in **C**. X-axis represents normalized aggregation relative to wild-type  
1137 from **D**. Error bars represent SEM with  $n=3$  for their respective dimension. **G)** Schematic diagram  
1138 describing the dichotomy in outcomes between p38 $\alpha$ -mediated TDP-43 phosphorylation and  
1139 PRMT1-mediated arginine methylation. Phospho-mimicking mutants favor aberrant aggregation  
1140 (thick arrow) over LLPS (thin arrow), whereas the arginine methylation-mimic favors LLPS (thick  
1141 arrow) over aberrant aggregation (thin arrow). This panel was made with BioRender.  
1142




1143

1144 **Figure 6. Methylation of TDP-43 regulates its aggregation.** **A)** Western blot of total and pTDP-43  
 1145 in RIPA and urea fractions of TDP-43<sup>WT</sup>-transfected SH-SY5Y-cells treated with DMSO (D) or with  
 1146 methyltransferase inhibitor AdOx (A) at a final concentration of 20 μM for 24h. **B)** Quantification of  
 1147 urea/RIPA ratio of total TDP-43 normalized to levels in DMSO-treated cells (mean band signal ± SD,  
 1148 unpaired t test, n = 4). **C)** Western blot of total and pTDP-43 in RIPA and urea fractions of TDP-  
 1149 43<sup>WT</sup>-transfected SH-SY5Y-cells co-transfected with empty control plasmid (Ctrl) or PRMT1. **D)**  
 1150 Quantification of urea/RIPA ratio of total TDP-43 normalized to levels in Ctrl-plasmid co-transfected  
 1151 cells (mean band signal ± SD, unpaired t test, n = 5). **E)** Western blot of total and pTDP-43 in RIPA  
 1152 and urea fractions of TDP-43<sup>WT</sup> or TDP-43<sup>G308R</sup> transfected SH-SY5Y-cells at time point of 24h  
 1153 post-transfection. **F)** Quantification of urea/RIPA ratio of total TDP-43 normalized to levels in TDP-  
 1154 43<sup>WT</sup>-transfected cells (mean band signal ± SD, one-way ANOVA with Dunnett's multiple  
 1155 comparison test, n = 4). \*p < 0.05, \*\*\*p < 0.001.

1156






1157


1158 **Figure 7. Crosstalk between PRMT1-catalyzed arginine methylation and p38 $\alpha$ -mediated**  
 1159 **phosphorylation of TDP-43. A)** Western blot of immunoprecipitated flag-tagged TDP-43 from SH-  
 1160 SY5Y cells probed with antibodies against TDP-43, pTDP-43 and MMA. GAPDH serves as a loading  
 1161 control. **B)** Western blot of immunoprecipitated flag-tagged TDP-43<sup>WT</sup> from the SH-SY5Y cells with  
 1162 and without siRNA-induced p38 $\alpha$  knockdown probed with antibodies against TDP-43, p38 $\alpha$ , pTDP-  
 1163 43 and MMA. GAPDH serves as a loading control. **C)** Quantification of TDP-43-CTF/full length-  
 1164 ratio (mean band signal  $\pm$  SD, unpaired t test, n = 3). (\*p < 0.05, \*\*p < 0.01). **D)** Quantification of  
 1165 mono-methylated TDP-43-CTF normalized to total TDP-43-CTF (mean band signal  $\pm$  SD, unpaired  
 1166 t test, n = 3). (\*p < 0.05, \*\*p < 0.01). **E)** Western blot of immunoprecipitated flag-tagged TDP-43<sup>WT</sup>  
 1167 from the SH-SY5Y cells with and without PRMT1 overexpression probed with antibodies against  
 1168 TDP-43, pTDP-43 and MMA. GAPDH serves as a loading control.  
 1169





1170

1171 **Figure S1. Genetic ablation of p38α MAPK reduces the aggregation and phosphorylation at**  
 1172 **S409/S410 for both TDP-43<sup>WT</sup> and TDP-43<sup>M337V</sup>.** **A)** Western blot of total and pTDP-43<sup>WT</sup> in RIPA  
 1173 and urea fractions of cells with siRNA-induced p38α knockdown in SH-SY5Y cells after 48h.  
 1174 GAPDH was used as a loading control. **B)** Western blot of total and pTDP-43<sup>M337V</sup> in RIPA and urea  
 1175 fractions of cells with siRNA-induced p38α knockdown in SH-SY5Y cells after 48h. GAPDH was  
 1176 used as a loading control. **C)** Quantification of pTDP-43<sup>WT</sup>/total TDP-43<sup>WT</sup> -ratio in urea fraction  
 1177 normalized to levels in scrambled siRNA (si-scr). **D)** Quantification of pTDP-43<sup>M337V</sup>/total TDP-  
 1178 43<sup>M337V</sup> -ratio in urea fraction normalized to levels in si-scr (mean band signal ± SD, one-way  
 1179 ANOVA with Sidak's multiple comparison test, n = 3). \*p < 0.05, \*\*\*p < 0.001.  
 1180 Related to Figure 1.  
 1181



1182

1183 **Figure S2. Knockdown of p38 $\alpha$  reduces the co-immunoprecipitation of endogenous p38 $\alpha$  with**  
1184 **TDP-43.** Western blot of immunoprecipitated TDP-43<sup>M337V</sup> and co-immunoprecipitated endogenous  
1185 p38 $\alpha$  from the SH-SY5Y cells with and without siRNA-induced p38 $\alpha$  knockdown. GAPDH serves  
1186 as a loading control.

1187 Related to Figure 3.

1188

1189