Functional multi-omics reveals genetic and pharmacologic regulation of surface CD38 in multiple myeloma

Authors:

Priya Choudhry[1,+], Olivia Gugliemini[1], Huimin Geng[1], Vishesh Sarin[1], Letitia Sarah[2], Yu-Hsiu T. Lin[1], Neha Paranjape[1], Poornima Ramkumar[3], Makeba Marcoulis[1], Donghui Wang[4], Paul Phojanakong[4], Veronica Steri[4], Byron Hann[4], Martin Kampmann[3,5,6], Arun P. Wiita[1,4]*

Affiliations:

[1]Dept. of Laboratory Medicine, University of California, San Francisco, CA

[2]Dept. of Cellular and Molecular Pharmacology, University of California, San Francisco, CA

[3]Institute for Neurodegenerative Diseases, University of California, San Francisco, CA

[4]Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco, CA

[5]Dept. of Biochemistry and Biophysics, University of California, San Francisco, CA

[6]Chan Zuckerberg Biohub, San Francisco, CA

[+]current address: Genentech/Roche, South San Francisco, CA

*Correspondence:

Arun P. Wiita, MD, PhD

Dept. of Laboratory Medicine

University of California, San Francisco

San Francisco, CA 94107

Email: Arun.wiita@ucsf.edu

Running Title: CD38 regulation in myeloma

Number of Figures: 6

Number of Tables: 0

Number of Supplementary Figures: 5

Number of Supplementary Tables/Datasets: 5

Abstract/Manuscript Main Text Word Count: 192/4,293

Abstract

CD38 is a surface ectoenzyme expressed at high levels on myeloma plasma cells and is the target

for the monoclonal antibodies (mAbs) daratumumab and isatuximab. CD38 density on tumor cells

is an important determinant of mAb efficacy, and CD38 loss after mAb treatment may play a role in

resistance. Several small molecules have been found to increase tumor surface CD38, with the goal

of boosting mAb efficacy in a co-treatment strategy. Here we sought to extend our currently limited

insight into CD38 surface expression by using a multi-omics approach. Genome-wide CRISPR-

interference screens integrated with patient-centered epigenetic analysis confirmed known

regulators of *CD38*, such as RARA, while revealing XBP1 and SPI1 as other key transcription

factors governing surface CD38 levels. *CD38* knockdown followed by cell surface proteomics

demonstrated no significant remodeling of the myeloma "surfaceome" after genetically-induced

loss of this antigen. Integrated transcriptome and surface proteome data confirmed high specificity

of all-trans retinoic acid in upregulating CD38 in contrast to broader effects of azacytidine and

panobinostat. Finally, unbiased phosphoproteomics identified inhibition of MAP kinase pathway

signaling in tumor cells after daratumumab treatment. Our work provides a resource to design

strategies to enhance efficacy of CD38-targeting immunotherapies in myeloma.

1 Introduction

Harnessing the immune system to treat myeloma has rapidly become the most exciting therapeutic

frontier in this disease. The first such immunotherapy agent to achieve United States Food and Drug

Administration (FDA) approval was the monoclonal antibody (mAb) daratumumab[1]. Daratumumab

targets CD38, a cell surface ectoenzyme highly expressed on myeloma plasma cells. Daratumumab

is currently FDA-approved for use as either monotherapy or combination therapy in the

relapsed/refractory setting, or front-line therapy in combination with other small molecule agents[1].

A second mAb targeting CD38, isatuximab, was also recently approved for relapsed/refractory

myeloma; at least 15 additional CD38-targeting agents are in development[2]. Extensive and

encouraging clinical data has already been obtained with daratumumab, though unfortunately

resistance appears to inevitably occur[3,4]. Biologically, this process appears to be quite complex,

with determinants of resistance ranging from alteration of surface antigens on tumor cells[4-6] to

dysfunction of the tumor immune microenvironment[7]. While it remains unclear whether CD38

downregulation on tumor cells after mAb treatment is a marker of impending resistance[5,8], or,

instead, a sign of successful therapy[9], compelling preclinical and clinical data suggests that CD38

surface antigen density prior to treatment strongly correlates with mAb efficacy[5].

This latter observation has led to numerous efforts to identify small molecules which can

increase tumor surface antigen density of CD38, representing potential co-treatments with CD38-

targeting mAbs. The first such example of a CD38-boosting small molecule was all-trans retinoic

acid (ATRA)[10]. These experiments in myeloma were inspired by earlier literature, suggesting that

removal of retinoic acid receptor alpha (RARA) suppression can drive CD38 expression in myeloid

lineage cells[11]. Subsequent studies identified the pan-histone deacetylase (HDAC) inhibitor

panobinostat[12], the thalidomide analog lenalidomide[13], and the Janus kinase (JAK) inhibitor

ruxolitinib[14] as agents that could potentially lead to myeloma surface CD38 increase. Based on the

observation that the *CD38* promoter contains an annotated CpG island, possibly repressing

transcription via methylation, we recently demonstrated that the DNA methyltransferase inhibitor

azacytidine (Aza) could also increase surface CD38 and enhance daratumumab efficacy as part of a

co-treatment strategy[15]. We have now opened a clinical trial at our institution to evaluate whether

this combination may also overcome prior daratumumab resistance (NCT04407442). A clinical trial

combining ATRA with daratumumab has led to encouraging outcomes in patients previously

refractory to daratumumab[16].

While these published strategies are certainly cause for optimism about ways to improve

CD38 mAb outcomes, they also leave many questions unanswered. Most notably, we do not yet

have a broad global sense of the transcriptional or post-transcriptional networks that most strongly

impact CD38 expression. There may be alternate strategies to even more potently increase CD38

expression that have not yet been identified. Furthermore, prior studies showed that CD38

downregulation after daratumumab treatment was accompanied by increases in the complement

inhibitors CD55 and CD59[5]. Are there other features of myeloma surface remodeling driven by

CD38 downregulation? And, for the small molecules noted above, it is unknown how they more

generally impact the makeup of the myeloma cell surface proteome beyond CD38. The tumor cell

surface not only harbors opportunities for specific immunotherapeutic targeting but also serves as

the interface for communication between the tumor and its microenvironment, potentially leading to

other alterations in myeloma biology after changes in surface proteomic profile. To address these

questions, here we have taken advantage of CRISPR interference-based functional genomic screens,

cell surface proteomics, epigenetic analyses, and phosphoproteomics to provide a multi-omic

perspective on CD38 regulation and tumor cell consequences of targeting CD38 in myeloma.

Methods

CRISPR interference screening and hit validation

Genome-wide CRISPRi screening was performed as described previously[17]. Briefly, RPMI-8226

cells stably expressing dCas9-KRAB were transduced with a genome-wide library comprised 5

sgRNA/protein coding gene. After 14 days cells were stained for surface CD38 and flow-sorted to

enrich for populations of cells expressing low or high cell surface levels of CD38. Cell populations

were then processed for next-generation sequencing as previously described[18] and sequenced on a

HiSeq-4000 (Illumina). Reads were analyzed by using the MAGeCK pipeline as previously

described[19]. Further validation was performed by knockdown with individual sgRNA's extracted

from the genome-wide library with conformation by flow cytometry or Western blotting. Antibody-

dependent cytotoxicity assays were performed using NK92-CD16 cells as described previously[15].

Additional details in Supplementary Methods.

Epigenetic analysis and machine learning for CD38 *transcriptional regulation*

Publicly available ATAC-seq data from primary myeloma samples (ref.[20]) was analyzed with the

Homer tool findPeaks. Motif binding in the identified ATAC peak regions was called with PROMO

tool[21]. Newly-diagnosed patient tumor RNA-seq data in the Multiple Myeloma Research

Foundation CoMMpass trial (MMRF; research.themmrf.org) was used to correlate expression of

predicted transcription factors with *CD38* expression. To build a predictive model for *CD38*

expression as a function of transcription factor expression, we developed an XGBoost (Extreme

Gradient Boosting) model with randomized search with cross validation to find optimal parameters.

80% of CoMMpass data was used for training and the remainder for model testing. Additional

details in Supplementary Methods.

Cell surface proteomics and phosphoproteomics

Cell surface proteomics was performed using an adapted version of the *N*-glycoprotein Cell Surface

Capture[22] method, as we have described previously[23]. Unbiased phosphoproteomics was performed

using immobilized metal affinity (IMAC) chromatography using methods described previously[24].

All samples were analyzed on a Thermo Q-Exactive Plus mass spectrometer with data processing in

MaxQuant[25]. Additional details in Supplementary Methods.

Results

A CRISPR interference-based screen reveals regulators of CD38 surface expression

We first sought to use an unbiased approach to identify regulators of surface CD38 in myeloma

tumor cells. We specifically employed genome-wide screening with CRISPR interference

(CRISPRi), an approach which leads to much higher specificity of knockdown than shRNA while

avoiding potential toxicity of double-strand breakage with CRISPR deletion[26]. We recently used

this approach to characterize regulators of surface B-cell Maturation Antigen (BCMA) in

myeloma[17]. Here, we employed an RPMI-8226 cell line with the dCas9-KRAB machinery, required

for CRISPRi, as described previously[17]. We confirmed that this RPMI-8226 cell line robustly

expressed CD38 (**Supp. Fig. 1A**).

The genome-wide screen was performed as shown in **Fig. 1A**. Briefly, RPMI-8226 cells

were transduced with a pooled genome-wide sgRNA library. After 14 days the cells were then

stained with fluorescently-labeled anti-CD38 antibody and flow sorted into low- and high-CD38

populations. Frequencies of cells expressing each sgRNA was quantified using next generation

sequencing. As an important positive control, increasing confidence in the screen results, we first

noted that knockdown of *CD38* itself strongly decreased surface CD38 expression (**Fig. 1B**). On the

other hand, several dozen genes, when repressed, did indeed lead to increased surface CD38 (right

side of volcano plot in **Fig. 1B**; **Supp. Table 1**). As another positive control, one of these top hits

included *RARA*, whose degradation is catalyzed by ATRA treatment to drive CD38 increase[10].

To find pathways that may be useful for pharmacologic targeting, we first applied Gene

Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis to the list of

genes that, when inhibited, significantly increased CD38 (**Fig. 1C**). We were intrigued to find that

many of the strongest effects appeared to be driven by transcriptional or other epigenetic factors.

These specifically included pathways such as "DNA replication", "mRNA processing", "DNA-

templated transcription", and "spliceosome".

We considered whether any hits associated with these pathways may be "druggable", with

the goal of expanding our repertoire of small molecules that enhance surface CD38 in myeloma.

SS18, a component of the BAF (BRG1/BRM associated factor) chromatin remodeling complex,

scored highly as a hit. However, treatment with the proposed BAF inhibitor caffeic acid phenol

ester (CAPE)[27] did not lead to consistent increases in surface CD38 (**Supp. Fig. 1B**). Similarly, the

lysine demethylase *KDM4A* was a prominent hit, but treatment with the inhibitory metabolite (*R*)-2-

hydroxyglutarate[28] also had no effect (**Supp. Fig. 1B**).

The strongest hits for genes whose knockdown increased surface CD38 were two

transcription factors, *HEXIM1* and *TLE3*. While validation studies using individual sgRNA

knockdown confirmed increased surface CD38 (**Fig. 1D**, **Fig. 2A** and **Supp. Fig. 2A**), as well as

functional impact in NK cell antibody-dependent cellular cytoxicity (ADCC) assays with

daratumumab (**Fig. 2B**), these effects were modest. Furthermore, these proteins are known to be

widespread negative regulators of transcription[29,30], suggesting little scope for specific therapeutic

targeting at the *CD38* locus, and to our knowledge no small molecule inhibitors of these proteins are

under development. Overall, our screen did not reveal any straightforward alternatives to

pharmacologically increase CD38 beyond those already known.

We were surprised that other targets proposed to increase CD38 expression after

pharmacologic inhibition, such as HDACs[3], or catalyzed degradation, such as IKZF1/3 (ref.[25]), did

not appear as prominent hits (**Fig. 1B**). However, this result may reflect a limitation of functional

genomic screens. A pharmacologic agent may inhibit multiple members of a protein class to drive a

phenotype, whereas, with single gene knockdown, functional redundancy may prevent this

phenotype from appearing[31] (i.e., multiple HDACs may need to ablated at once, or both IKZF1 and

IKZF3 simultaneously, to drive increased CD38). We speculate this is the case with DNA

methyltransferases (DNMTs). We previously showed that treatment with the DNMT inhibitor

azacytidine, which promotes degradation of all cellular DNMTs[32], could robustly increase surface CD38[15]. Here, however, we found that knockdown of any individual DNMT only led to minor CD38 increase (**Fig. 1B**).

Given these findings, we therefore shifted our focus to genes that, when knocked down, led to CD38 decrease (left side of volcano plot in **Fig. 1B**). We reasoned this approach could still reveal important biological inputs that regulate the surface expression of CD38. Examining specific genes, we found that the transcription factor *SPI1* was the strongest hit besides *CD38* that, when knocked down, repressed surface CD38 expression. We also noted that *NFKB1* and *NFKB2* knockdown appeared to drive CD38 decrease. This finding was intriguing given the known importance of NF-κB signaling in myeloma proliferation and survival[33]. KEGG and GO analysis of genes whose knockdown significantly decreased CD38 showed enrichment for MAP kinase pathway and protein phosphorylation more broadly (**Supp. Fig. 1C**), suggesting key roles for intracellular signaling in regulating surface CD38.

Validation experiments with individual sgRNAs confirmed that *SPI1* knockdown strongly decreased CD38 surface expression by flow cytometry, with a lesser decrease in surface CD38 with *NFKB2* knockdown (**Fig. 1D, Fig. 2C, Supp. Fig. 2C-D**). These alterations also led to functional impacts. RPMI-8226 cells with knockdown of these genes showed significantly decreased NK cell lysis in ADCC assays (**Fig. 2D**). We further probed this dynamic *in vivo*, finding that RPMI-8226 cells with *SPI1* knockdown were relatively resistant to daratumumab in a murine model (**Fig. 2E**). We note that we attempted to expand these results to additional cell lines. However, our four other myeloma cell lines harboring the CRISPRi machinery[17] all express extremely low levels of *SPI1* (**Supp. Fig. 2B**) and attempted knockdown in two of them (AMO1, KMS12PE) did not elicit any phenotype (not shown). Therefore, this finding suggests that SPI1 may play an important role in regulating CD38 expression in some myeloma tumors, but it is less likely to be a universal regulator.

Epigenetic analysis suggests XBP1 *as a key determinant of* CD38 *in primary myeloma tumors*

Our CRISPRi results suggest that epigenetic and/or transcriptional regulation is a critical driver of surface CD38 levels. However, we do not know whether these specific hits in a myeloma cell line will extend to primary myeloma tumors. We therefore took a complementary approach to find potential transcriptional regulators of *CD38*. Using ATAC-seq data from 24 primary myeloma tumor samples[20], we extracted open chromatin motifs near the *CD38* promoter (**Fig. 3A**) to identify

a list of 46 transcription factors with potential binding sites at this locus (**Supp. Table 2**). We then correlated expression (via Pearson R) of these transcription factors with *CD38* expression across 664 primary patient tumors at diagnosis in the Multiple Myeloma Research Foundation CoMMpass database (research.themmrf.org; release IA13). In this analysis, we found the transcription factor most negatively correlated with *CD38* expression was *RARA* (**Fig. 3A**), consistent with our CRISPRi screen data, and underscoring the promise of ATRA as a co-treatment to increase *CD38*. Intriguingly, the transcription factor with strongest positive correlation was *XBP1* (**Fig. 3A-B**), a central driver of plasma cell identity[34]. *XBP1* also showed strong positive correlations with *CD38* in two other patient tumor gene expression datasets[35,36] (**Supp. Table 2**).

To further extend this analysis, we sought to build a predictive model which could estimate *CD38* transcript level as a function of transcription factor expression. We used an XGBoost method applied to CoMMpass mRNA-seq data to find weights of transcription factor expression that most-influence *CD38* levels in patient tumors. We first tested this analysis on 80% of patient data as a training set with 20% left out as test set. We found this model, solely based on transcription factor expression, could predict about half of the variance (coefficient of variation $R^2 = 0.49$ using 5-fold cross validation) in test set *CD38* levels (**Fig. 3C**). Using model weights and Shapley Additive Explanations (SHAP) analysis (see Supplementary Methods) to determine transcription factors that have the greatest impact, either positive or negative, on CD38 expression, we found that *XBP1* played the strongest role overall. Other strong hits from both of our analyses included *IRF2*, *ATF1*, and *STAT1*. *SPI1* also appeared in the top 10 most relevant transcriptional regulators (**Fig. 3D**), consistent with our CRISPRi results, suggesting that SPI1 may play a key role in regulating *CD38* in a subset of tumors.

We further evaluated XBP1 given its prominent role in these two complementary bioinformatic analyses. Consistent with the hypothesis that XBP1 regulates CD38 expression, we noted in our CRISPRi screen that *XBP1* knockdown indeed led to significant surface CD38 decrease (**Fig. 1B**). We further examined a prior dataset of shRNA knockdown of *XBP1* in myeloma plasma cells[37]. In this independent experiment, *CD38* mRNA was decreased ~3-fold after *XBP1* silencing (**Supp. Fig. 3C**). Taken together, these results nominate XBP1 as a particularly strong determinant of surface CD38 in myeloma plasma cells, while also supporting a role for SPI1 as highlighted in our functional genomic screen.

No consistent large-scale remodeling of the myeloma surface proteome after CD38 downregulation

We next evaluated CD38 surface regulation from the perspective of monoclonal antibody

therapy. It has previously been demonstrated that myeloma plasma cells rapidly lose surface CD38

after treatment with daratumumab[5]. In clinical samples, CD38 loss was accompanied by increases in

CD55 and CD59, which may inhibit complement-dependent cytotoxicity and contribute to

daratumumab resistance[5]. In preclinical studies, macrophage trogocytosis has been proposed as a

mechanism contributing to CD38 loss after mAb treatment, that also leads to alterations in other

surface antigens including CD138/SDC1[7]. However, we hypothesized that given its enzymatic

activity and role as a cellular differentiation marker[38], loss of CD38 on its own may influence

surface expression of other myeloma antigens. Such alterations may reveal new biology or

(immuno)therapeutic vulnerabilities of CD38 mAb-treated disease.

To test this hypothesis, we employed a method we recently developed termed "antigen

escape profiling"[23]. We used CRISPRi to transcriptionally repress *CD38* in RPMI-8226, AMO-1,

and KMS12-PE myeloma cell lines, using this genetic approach to partially mimic loss of surface

antigen seen after mAb therapy (**Fig. 4A**). We then performed Cell Surface Capture proteomics[22,23]

to uncover surface proteome alterations in a relatively unbiased fashion. Across cell lines, analyzed

in biological triplicate with *CD38* knockdown vs. non-targeting sgRNA, we quantified 897 proteins

annotated as membrane-spanning in Uniprot (minimum of two peptides per protein) (**Supp. Table**

3). As a positive control, in all lines we found that the strongest signature was decrease of CD38

itself (**Supp. Fig. 4A-C**). Initial analysis of the data suggested potential alterations across several

dozen other surface proteins (**Supp. Fig. 4A-C**). However, there was notable heterogeneity across

cell lines; when aggregating proteomic data, we found no significant alterations in any surface

antigens beyond CD38 (**Fig. 4B**). Integration with RNA-seq data on the same samples allowed us to

identify transcriptionally-regulated changes also present at the surface protein level (**Supp. Table**

3). This analysis revealed only THY-1/CD90 as upregulated >3-fold at both the mRNA and surface

proteomic level after *CD38* knockdown (**Fig. 4C**). Intriguingly, CD90 is known as a marker of

"stemness" in early hematopoietic lineage cells that is lost when CD38 expression is increased[39].

CoMMpass analysis also confirmed increased *THY1* expression in tumors with lower *CD38* (**Supp.**

Fig. 4D). However, further validation as to whether CD90 is truly altered after CD38 mAb will

require pre- and post-treatment clinical specimens, beyond the scope of our work here. Overall, we

conclude that loss of CD38 in isolation leads to only minor, if any, consistent remodeling of the

myeloma surface proteome.

Integrated surface proteomic and transcriptional analysis suggests ATRA is highly specific in CD38

upregulation

Data from our group[15] and others[10,12-14] have suggested that several small molecules can increase myeloma surface CD38. However, the broader impacts of these agents on membrane antigens beyond CD38 have not been directly compared. We performed integrated cell surface proteomics and transcriptional analysis of RPMI-8226 cells treated with 10 nM all-trans retinoic acid (ATRA), 2 μM azaciditine (Aza), and 10 nM Panobinostat (Pano), all treated for 72 hr, in comparison to DMSO (**Supp. Table 4**). These doses are chosen as they have been previously published to significantly increase myeloma surface CD38 by flow cytometry[10,12,15]. In this integrated analysis we found much broader impacts of azacytidine and panobinostat than ATRA on the "surfaceome" of plasma cells. For example, after ATRA, no genes showed greater than 4-fold (2-fold on $\log_2$ scale) expression change for membrane protein-coding genes, unlike prominent changes above this threshold for Aza and Pano (**Fig. 5A**). However, CD38 was one of the few genes detectably upregulated at both the proteomic and mRNA level after ATRA. This observation appears consistent with the known mechanism of action for these small molecules. ATRA is expected to only influence the small subset of genes, including CD38, that are transcriptionally regulated by the retinoic acid receptor[40]. In contrast, Aza and pano are expected to drive broad changes in epigenetic regulation across the genome[32,41]. These results suggest that, at doses driving CD38 upregulation, for Aza or pano altering CD38 is just a small component of their impact on myeloma tumor cells, whereas ATRA is much more specific in driving CD38 upregulation with minimal other cellular impacts.

Toward understanding how CD38 is modulated after drug treatment, in our previous work[15] we noted that the mechanism of *CD38* increase after Aza treatment was unclear. We thus further investigated the global transcriptional response (i.e. not limited to membrane protein-coding genes) after Aza (**Supp. Table 4**). Prior studies have suggested that the anti-tumor effect of Aza is largely mediated by reactivation of endogenous retroviruses, thus stimulating a tumor-autonomous interferon response ultimately leading to apoptosis[42,43]. Consistent with this work in other model systems, in this myeloma model we found a pronounced increase in interferon-responsive genes after Aza, but not ATRA, treatment including *IRF1*, *IFITM1*, *IFITM2*, and *IFITM3* (**Fig. 5B**). KEGG analysis also confirmed this effect, where pathways most significantly upregulated after Aza treatment were related to response to bacterial or viral infection (**Fig. 5C**). Given evidence across multiple systems that interferon upregulates *CD38* expression[44-46], our transcriptional profiling data

also supports an interferon-based mechanism driving surface CD38 increase in plasma cells after

Aza treatment. This proposed mechanism is also consistent with our prior observation that the

CD38 promoter exhibits minimal methylation at baseline in myeloma plasma cells[15], making it

unlikely that direct demethylation of the *CD38* CpG island is the primary driver of increased *CD38*

transcription after Aza.

Plasma cell proliferative signaling pathways are inhibited by mAb binding to CD38

In our final set of experiments related to targeting surface CD38, we were intrigued as to

whether binding of a therapeutic mAb leads to specific cellular phenotypes within myeloma plasma

cells. For example, isatuximab is known to directly lead to apoptosis of plasma cells[47], and

daratumumab can do so after crosslinking[48]. However, the mechanism underlying this transduction

of extracellular mAb binding to intracellular phenotype remains unclear. In addition, our CRISPRi

screen data (**Fig. 1B and Supp. Fig. 1C**) suggests that surface CD38 expression may be strongly

impacted by intracellular phospho-signaling pathways.

Therefore, we used unbiased phosphoproteomics by mass spectrometry to probe intracellular

signaling effects driven by CD38 mAb binding. In RPMI-8226 cells we compared 20 uM

daratumumab treatment vs. IgG1 isotype control. We chose a time point of 20 minutes of treatment

given known rapid alterations in signaling pathways in similar phosphoproteomic experiments[49]. In

total, across triplicate samples we quantified 5430 phosphopeptides (**Supp. Table 5; Supp. Fig.**

5A). Analyzing phosphopeptide changes by Kinase Substrate Enrichment Analysis (KSEA)[50], we

were intrigued to find downregulation of phosphorylation motifs consistent with both cyclin-

dependent kinases as well as several kinases of the MAP kinase pathway (**Fig. 6A**). Downregulation

of phosphorylation on several central nodes in the MAP kinase as well as AKT pathway was also

apparent via KEGG analysis (**Supp. Fig. 5B**). Across a time course we further confirmed effects on

MAP kinase pathway (reported by phosphorylation of MAPK (ERK1/2), a key node in this

response) and AKT signaling after daratumumab treatment via Western blotting in RPMI-8226 and

MM.1S cell lines, respectively (**Fig. 6B**). While the absolute value of changes in MAPK signaling

are modest, both by phosphoproteomics and Western blot, these results indicate that daratumumab

binding to CD38 can at least partially inhibit this central proliferative pathway within myeloma

tumor cells, and thus may form a component of daratumumab's anti-tumor effect.

Discussion

Our studies here present a "multi-omics" view of therapeutically targeting CD38 in multiple

myeloma. Our integrated functional genomics and epigenetic analysis point to the central role of

transcriptional regulators in governing CD38 surface expression at baseline. Using cell surface

proteomics, we further identify that loss of CD38 in isolation is unlikely to drive large changes in

the cell surface proteome, while known drug treatment strategies to increase CD38 have largely

divergent impacts on other surface antigens. Finally, unbiased phosphoproteomics reveals that

binding of anti-CD38 mAb can impair intracellular proliferative signaling within plasma cells.

Our initial CRISPRi screen illustrated the central role of numerous transcription factors,

such as SPI1, HEXIM1, and TLE3, in regulating surface CD38. This functional genomic study

suggests that regulation of surface CD38 largely occurs at the transcriptional, as opposed to protein

trafficking, level. This finding was in sharp contrast to our prior CRISPRi results with BCMA,

where we found that post-transcriptional mechanisms, such as proteolytic cleavage by γ-secretase

and protein trafficking via the SEC61 translocon, played some of the strongest roles in determining

surface BCMA levels[17].

Another recent study used genome-wide CRISPR deletion screening to find genes that,

when knocked out, could abrogate IL-6-mediated downregulation of surface CD38 (ref.[14]). The

strongest hits in this prior study included the transcription factors *STAT1* and *STAT3*, demonstrating

a role for JAK-STAT signaling in regulating tumor CD38 expression within the bone marrow

microenvironment[14]. In support of this notion, our integrated epigenetic and machine learning

analyses, extracted from bone marrow-derived patient tumor samples, also support a critical role for

STAT1 in governing surface CD38. However, in our CRISPRi screen in an *in vitro* monoculture

system, neither *STAT1* or *STAT3* affected CD38 surface expression (**Fig. 1B**). This result suggests

that JAK-STAT signaling may not play a major role in CD38 regulation in the absence of

exogenous tumor stimulation. This finding illustrates the complementary nature of our genome-

wide screen to that previously published under the context of IL-6 stimulation[14].

Toward the goal of finding key regulators of CD38 that were not previously known, our

epigenetic and machine learning approaches suggest that *XBP1* may be a critical regulator of

plasma cell *CD38*. While this initial analysis of primary samples only demonstrates correlation in

gene expression, our CRISPRi screen, as well as the independent experiment of Leung-Hagesteijn

et al[37], both suggest that genetic knockdown of *XBP1* does directly lead to CD38 decrease,

supporting a true functional relationship. Further exploring the connection between XBP1 and

CD38 will be an intriguing avenue for future studies. Dissecting this interaction may ultimately lead to new strategies to increase surface CD38 in myeloma plasma cells via modulating XBP1.

Given that it is now well-known that plasma cells demonstrate strong downregulation of CD38 after daratumumab treatment[5], a pressing clinical question is whether CD38-low, daratumumab-resistant cells have novel immunotherapeutic vulnerabilities. However, our recently-described strategy of "antigen escape profiling"[23] – CRISPRi knockdown followed by unbiased cell surface proteomics (**Fig. 4**) – suggests that other cell surface antigens on plasma cells do not exhibit consistent changes due to CD38 downregulation alone. This finding supports the notion that alterations in surface proteins found after mAb treatment on patent tumors, such as increases in CD55 and CD59 (ref.[5]), are caused by other therapy-induced selective pressure within the tumor microenvironment, not CD38 loss.

Also directly related to mAb therapeutic effects, our unbiased phosphoproteomic results suggest that daratumumab binding to CD38 can directly decrease signaling along the MAP kinase and PI3K-AKT pathways. While these findings were confirmed by Western blot in both RPMI-8226 and MM.1S cells, we note that both of these cell lines harbor mutations in $KRAS$[24]. It is unclear if similar intracellular signaling effects occur in myeloma tumors without MAPK-pathway mutations. Furthermore, it remains to be investigated whether this inhibition of central proliferative signaling pathways plays a role in the anti-tumor effect of daratumumab in patients. It will be interesting to address this question in future studies.

In terms of limitations of our work, the most prominent is that the many of our studies are derived from large-scale "omics" experiments in myeloma cell lines. There may be biological differences between our findings *in vitro* and primary tumors growing within the bone marrow microenvironment.

Taken together, our multi-omic studies comprise a resource that reveals new insight into the genetic, epigenetic, and pharmacologic regulation of surface CD38 in myeloma plasma cells. We anticipate these findings will have utility in deriving new strategies to enhance CD38 mAb efficacy or to overcome resistance to these agents.

Acknowledgements: We thank Ruilin Tian for assistance in modifying MAGeCK scripts for the analysis here. This work was supported by grants K08 CA184116, R01 CA226851, and the UCSF Stephen and Nancy Grand Multiple Myeloma Translational Initiative (to A.P.W.), NCI P30

1   CA082103 supporting the Preclinical Therapeutics Core facility (managed by V.Steri and B.H.),

2   and K99/R00 CA181494 and a Stand Up to Cancer Innovative Research Grant (to M.K.).

Author Contributions: P.C., P.R., M.K., and A.P.W. conceived and designed the study. P.C., O.G., L.S., Y-H.T.L., P.R., N.P., and M.M. performed experiments. P.C., N.P., Y-H.T.L., and P.R. analyzed data. D.W., P.P., V. Steri, and B.H. performed murine studies. H.G. analyzed patient epigenetic data. V. Sarin performed machine learning analysis. P.C. and A.P.W. wrote the manuscript with input from all authors.

Author Contributions: P.C., P.R., M.K., and A.P.W. conceived and designed the study. P.C.,

O.G., L.S., Y-H.T.L., P.R., N.P., and M.M. performed experiments. P.C., N.P., Y-H.T.L., and P.R.

analyzed data. D.W., P.P., V. Steri, and B.H. performed murine studies. H.G. analyzed patient

epigenetic data. V. Sarin performed machine learning analysis. P.C. and A.P.W. wrote the

manuscript with input from all authors.

Disclosures: P.C. is currently an employee and shareholder of Genentech/Roche, though during the

time of completing this project she was fully employed by the University of California, San

Francisco. P.R. is currently an employee and shareholder of Senti Biosciences, though during the

time of completing this project she was fully employed by the University of California, San

Francisco. A.P.W. is an equity holder and scientific advisory board member of Indapta

Therapeutics, LLC and Protocol Intelligence, LLC. M.K. has filed a patent application related to

CRISPRi screening (PCT/US15/40449); and serves on the Scientific Advisory Boards of Engine

Biosciences, Cajal Neuroscience and Casma Therapeutics. The other authors declare no relevant

conflicts of interest.

REFERENCES:

Syed, Y. Y. Daratumumab: A Review in Combination Therapy for Transplant-Ineligible Newly Diagnosed Multiple Myeloma. *Drugs* **79**, 447-454, doi:10.1007/s40265-019-01080-6 (2019).

Bonello, F. *et al.* CD38 as an immunotherapeutic target in multiple myeloma. *Expert Opin Biol Ther* **18**, 1209-1221, doi:10.1080/14712598.2018.1544240 (2018).

van de Donk, N. & Usmani, S. Z. CD38 Antibodies in Multiple Myeloma: Mechanisms of Action and Modes of Resistance. *Front Immunol* **9**, 2134, doi:10.3389/fimmu.2018.02134 (2018).

Saltarella, I. *et al.* Mechanisms of Resistance to Anti-CD38 Daratumumab in Multiple Myeloma. *Cells* **9**, doi:10.3390/cells9010167 (2020).

Nijhof, I. S. *et al.* CD38 expression and complement inhibitors affect response and resistance to daratumumab therapy in myeloma. *Blood* **128**, 959-970, doi:10.1182/blood-2016-03-703439 (2016).

Overdijk, M. B. *et al.* Antibody-mediated phagocytosis contributes to the anti-tumor activity of the therapeutic antibody daratumumab in lymphoma and multiple myeloma. *mAbs* **7**, 311-321, doi:10.1080/19420862.2015.1007813 (2015).

Krejcik, J. *et al.* Daratumumab depletes CD38+ immune regulatory cells, promotes T-cell expansion, and skews T-cell repertoire in multiple myeloma. *Blood* **128**, 384-394, doi:10.1182/blood-2015-12-687749 (2016).

Krejcik, J. *et al.* Monocytes and Granulocytes Reduce CD38 Expression Levels on Myeloma Cells in Patients Treated with Daratumumab. *Clin Cancer Res* **23**, 7498-7511, doi:10.1158/1078-0432.CCR-17-2027 (2017).

Plesner, T., van de Donk, N. & Richardson, P. G. Controversy in the Use of CD38 Antibody for Treatment of Myeloma: Is High CD38 Expression Good or Bad? *Cells* **9**, doi:10.3390/cells9020378 (2020).

Nijhof, I. S. *et al.* Upregulation of CD38 expression on multiple myeloma cells by all-trans retinoic acid improves the efficacy of daratumumab. *Leukemia* **29**, 2039-2049, doi:10.1038/leu.2015.123 (2015).

Drach, J., Zhao, S., Malavasi, F. & Mehta, K. Rapid induction of CD38 antigen on myeloid leukemia cells by all trans-retinoic acid. *Biochem Biophys Res Comm* **195**, 545-550, doi:10.1006/bbrc.1993.2080 (1993).

Garcia-Guerrero, E. *et al.* Panobinostat induces CD38 upregulation and augments the antimyeloma efficacy of daratumumab. *Blood* **129**, 3386-3388, doi:10.1182/blood-2017-03-770776 (2017).

Fedele, P. L. *et al.* IMiDs prime myeloma cells for daratumumab-mediated cytotoxicity through loss of Ikaros and Aiolos. *Blood* **132**, 2166-2178, doi:10.1182/blood-2018-05-850727 (2018).

Ogiya, D. *et al.* The JAK-STAT pathway regulates CD38 on myeloma cells in the bone marrow microenvironment: therapeutic implications. *Blood* **136**, 2334-2345, doi:10.1182/blood.2019004332 (2020).

Choudhry, P. *et al.* DNA methyltransferase inhibitors upregulate CD38 protein expression and enhance daratumumab efficacy in multiple myeloma. *Leukemia* **34**, 938-941, doi:10.1038/s41375-019-0587-5 (2020).

Frerichs, K. A. *et al.* Efficacy and Safety of Daratumumab Combined with All-Trans Retinoic Acid in Relapsed/Refractory Multiple Myeloma; Results of the Phase 1/2 Dara/ATRA Study. *Blood (ASH Abstract)* **134**, 1826 (2019).

Ramkumar, P. *et al.* CRISPR-based screens uncover determinants of immunotherapy response in multiple myeloma. *Blood Adv* **4**, 2899-2911, doi:10.1182/bloodadvances.2019001346 (2020).

Kampmann, M., Bassik, M. C. & Weissman, J. S. Functional genomics platform for pooled screening and generation of mammalian genetic interaction maps. *Nat Protoc* **9**, 1825-1847, doi:10.1038/nprot.2014.103 (2014).

Tian, R. *et al.* CRISPR Interference-Based Platform for Multimodal Genetic Screens in Human iPSC-Derived Neurons. *Neuron* **104**, 239-255 e212, doi:10.1016/j.neuron.2019.07.014 (2019).

Jin, Y. *et al.* Active enhancer and chromatin accessibility landscapes chart the regulatory network of primary multiple myeloma. *Blood* **131**, 2138-2150, doi:10.1182/blood-2017-09-808063 (2018).

Messeguer, X. *et al.* PROMO: detection of known transcription regulatory elements using species-tailored searches. *Bioinformatics* **18**, 333-334, doi:10.1093/bioinformatics/18.2.333 (2002).

Wollscheid, B. *et al.* Mass-spectrometric identification and relative quantification of N-linked cell surface glycoproteins. *Nat Biotechnol* **27**, 378-386, doi:10.1038/nbt.1532 (2009).

Nix, M. A. *et al.* Surface proteomics reveals CD72 as a target for in vitro-evolved nanobody-based CAR-T cells in KMT2A/MLL1-rearranged B-ALL. *Cancer Discov*, doi:10.1158/2159-8290.CD-20-0242 (2021).

Lin, Y. T. *et al.* Integrated phosphoproteomics and transcriptional classifiers reveal hidden RAS signaling dynamics in multiple myeloma. *Blood Adv* **3**, 3214-3227, doi:10.1182/bloodadvances.2019000303 (2019).

Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat Protoc* **11**, 2301-2319, doi:10.1038/nprot.2016.136 (2016).

Gilbert, L. A. *et al.* Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell* **159**, 647-661, doi:10.1016/j.cell.2014.09.029 (2014).

Dykhuizen, E. C., Carmody, L. C., Tolliday, N., Crabtree, G. R. & Palmer, M. A. Screening for inhibitors of an essential chromatin remodeler in mouse embryonic stem cells by monitoring transcriptional regulation. *J Biomol Screen* **17**, 1221-1230, doi:10.1177/1087057112455060 (2012).

Carbonneau, M. *et al.* The oncometabolite 2-hydroxyglutarate activates the mTOR signalling pathway. *Nat Commun* **7**, 12700, doi:10.1038/ncomms12700 (2016).

Michels, A. A. & Bensaude, O. Hexim1, an RNA-controlled protein hub. *Transcription* **9**, 262-271, doi:10.1080/21541264.2018.1429836 (2018).

Cinnamon, E. & Paroush, Z. Context-dependent regulation of Groucho/TLE-mediated repression. *Curr Opin Genet Dev* **18**, 435-440, doi:10.1016/j.gde.2008.07.010 (2008).

Agrotis, A. & Ketteler, R. A new age in functional genomics using CRISPR/Cas9 in arrayed library screening. *Front Genet* **6**, 300, doi:10.3389/fgene.2015.00300 (2015).

Gnyszka, A., Jastrzebski, Z. & Flis, S. DNA methyltransferase inhibitors and their emerging role in epigenetic therapy of cancer. *Anticancer Res* **33**, 2989-2996 (2013).

Matthews, G. M. *et al.* NF-kappaB dysregulation in multiple myeloma. *Semin Cancer Biol* **39**, 68-76, doi:10.1016/j.semcancer.2016.08.005 (2016).

Reimold, A. M. *et al.* Plasma cell differentiation requires the transcription factor XBP-1. *Nature* **412**, 300-307, doi:10.1038/35085509 (2001).

Mulligan, G. *et al.* Gene expression profiling and correlation with outcome in clinical trials of the proteasome inhibitor bortezomib. *Blood* **109**, 3177-3188, doi:10.1182/blood-2006-09-044974 (2007).

Shi, L. *et al.* The MicroArray Quality Control (MAQC)-II study of common practices for the development and validation of microarray-based predictive models. *Nat Biotechnol* **28**, 827-838, doi:10.1038/nbt.1665 (2010).

Leung-Hagesteijn, C. *et al.* Xbp1s-negative tumor B cells and pre-plasmablasts mediate therapeutic proteasome inhibitor resistance in multiple myeloma. *Cancer Cell* **24**, 289-304, doi:10.1016/j.ccr.2013.08.009 (2013).

Li, Y., Yang, R., Chen, L. & Wu, S. CD38 as an immunomodulator in cancer. *Future Oncol* **16**, 2853-2861, doi:10.2217/fon-2020-0401 (2020).

Majeti, R., Park, C. Y. & Weissman, I. L. Identification of a hierarchy of multipotent hematopoietic progenitors in human cord blood. *Cell Stem Cell* **1**, 635-645, doi:10.1016/j.stem.2007.10.001 (2007).

Giuli, M. V. *et al.* Current Trends in ATRA Delivery for Cancer Therapy. *Pharmaceutics* **12**, doi:10.3390/pharmaceutics12080707 (2020).

Ganai, S. A. Panobinostat: The Small Molecule Metalloenzyme Inhibitor with Marvelous Anticancer Activity. *Curr Top Med Chem* **16**, 427-434, doi:10.2174/1568026615666150813145800 (2016).

Chiappinelli, K. B. *et al.* Inhibiting DNA Methylation Causes an Interferon Response in Cancer via dsRNA Including Endogenous Retroviruses. *Cell* **162**, 974-986, doi:10.1016/j.cell.2015.07.011 (2015).

Roulois, D. *et al.* DNA-Demethylating Agents Target Colorectal Cancer Cells by Inducing Viral Mimicry by Endogenous Transcripts. *Cell* **162**, 961-973, doi:10.1016/j.cell.2015.07.056 (2015).

Mihara, K. *et al.* All-trans retinoic acid and interferon-alpha increase CD38 expression on adult T-cell leukemia cells and sensitize them to T cells bearing anti-CD38 chimeric antigen receptors. *Blood Cancer J* **6**, e421, doi:10.1038/bcj.2016.30 (2016).

Bauvois, B. *et al.* Upregulation of CD38 gene expression in leukemic B cells by interferon types I and II. *J Interferon Cytokine Res* **19**, 1059-1066, doi:10.1089/107999099313299 (1999).

Frasca, L. *et al.* IFN-gamma arms human dendritic cells to perform multiple effector functions. *J Immunol* **180**, 1471-1481, doi:10.4049/jimmunol.180.3.1471 (2008).

Moreno, L. *et al.* The Mechanism of Action of the Anti-CD38 Monoclonal Antibody Isatuximab in Multiple Myeloma. *Clin Cancer Res* **25**, 3176-3187, doi:10.1158/1078-0432.CCR-18-1597 (2019).

Overdijk, M. B. *et al.* The Therapeutic CD38 Monoclonal Antibody Daratumumab Induces Programmed Cell Death via Fcgamma Receptor-Mediated Cross-Linking. *J Immunol* **197**, 807-813, doi:10.4049/jimmunol.1501351 (2016).

Humphrey, S. J., Azimifar, S. B. & Mann, M. High-throughput phosphoproteomics reveals in vivo insulin signaling dynamics. *Nat Biotechnol* **33**, 990-995, doi:10.1038/nbt.3327 (2015).

Casado, P. *et al.* Kinase-substrate enrichment analysis provides insights into the heterogeneity of signaling pathway activation in leukemia cells. *Sci Signal* **6**, rs6, doi:10.1126/scisignal.2003573 (2013).

Li, W. *et al.* MAGeCK enables robust identification of essential genes from genome-scale CRISPR/Cas9 knockout screens. *Genome Biol* **15**, 554, doi:10.1186/s13059-014-0554-4 (2014).

FIGURE 1

Figure 1. CRISPR interference (CRISPRi) screening reveals genetic determinants of surface CD38 regulation. A. Schematic of CRISPRi screen design. B. Results of CRISPRi screen demonstrating genes that, when knocked down, regulate surface CD38 in RPMI-8226 cells. X-axis indicates phenotype (epsilon) from MAGeCK[51] statistical analysis. Dashed line indicates cutoff for significant change at False Discovery Rate (FDR) < 0.05. Genes of interest are specifically labeled. 4,000 negative control non-targeting sgRNAs are in grey. C. Gene Ontology (GO) Biological Process and Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis of all genes that when knocked down lead to significant CD38 upregulation. D. Follow-up flow cytometry validation of CRISPRi screen hits using two individual sgRNAs per gene demonstrates *TLE3* knockdown drives increased CD38, while *SPI1* knockdown leads to CD38 decrease.

FIGURE 2

Figure 2. Validation of CRISPRi screen hits as functionally impacting daratumumab efficacy.
A. Knockdown of HEXIM1 and TLE3 with two independent sgRNA's/gene (AMO1 myeloma
cells, $n = 3$) followed by flow cytometry shows significant surface CD38 increase with TLE3_i2
sgRNA and trend toward increased CD38 with HEXIM1_i1 sgRNA. Scri = non-targeting control
sgRNA. **B.** Results from ADCC assays with AMO1 cells stably expressing the noted sgRNA's and
incubated with the indicated concentration of daratumumab or isotype control antibody (1:20
myeloma:NK ratio, 20 hours, $n = 2$). The percent lysis by ADCC was calculated as outlined in the
Supplemental methods. . At 10 μM daratumumab, both HEXIM1 and TLE3 knockdown led to
significant increase in ADCC. **C.** Similar to A., sgRNA knockdown of *NFKB1*, *NFKB2*, and *SPI1*
with fold-change in CD38 by flow cytometry (RPMI-8226 cells, $n = 3$). **D.** Similar to B.,
knockdown with the most effective sgRNA for each gene show significant decreases in NK-cell
ADCC at 10 μM daratumumab in the RMPI-8266 cells ($n = 2$). **E.** *In vivo* validation of *SPI1*
knockdown driving daratumumab resistance. NOD *scid* gamma mice were I.V. implanted with
CRISPRi RPMI-8226 cells stably expressing both luciferase and noted sgRNA, then treated with
200 μg daratumumab on the noted schedule. Bioluminescence imaging measurement of tumor
burden demonstrates significantly increased fold-change in tumor burden (normalized to pre-
daratumumab intensity) with either CD38 or SPI1 knockdown compared to scramble sgRNA. For
A-**E**, *$p < 0.05$, **$p < 0.01$ by two-tailed *t*-test.

FIGURE 3

Figure 3. Patient-centered epigenetic analysis and machine learning predicts most potent transcriptional regulators of *CD38*. **A.** 46 transcription factors predicted to bind to the CD38 locus were derived from motif analysis of published ATAC-seq data (see Supplementary Figure 3). Gene expression of each transcription factor (TF) was correlated with *CD38* expression in the Multiple Myeloma Research Foundation (MMRF) CoMMpass database (release IA13), with RNA-seq data from CD138+ enriched tumor cells at diagnosis ($n = 664$ patients). Top predicted positive and negative regulators are shown based on Pearson correlation (R). **B.** CoMMpass RNA-seq data illustrates strong positive correlation between *XBP1* and *CD38* expression. **C.** XGBoost machine learning model was used to extract features of transcription factor gene expression that best-model CD38 expression in CoMMpass tumors (shown in log2 TPM (Transcripts per Million)). 80% of data was used as a test set with 20% left out as a training set. Coefficient of variation (R^2) for predictive model = 0.49 after five-fold cross validation. **D.** Shapley Additive Explanations (SHAP) analysis indicates transcription factors whose expression most strongly impacts CD38 expression levels in CoMMpass tumors.

FIGURE 4

1
2
Figure 4. Minimal alterations of the myeloma cell surface proteome after CD38 loss. A.
Schematic of "antigen escape profiling" approach to reveal new cell-surface therapeutic
vulnerabilities in the context of CD38 downregulation. **B.** Cell surface capture proteomics
comparing *CD38* knockdown vs. non-targeting sgRNA control, with aggregated data across three
cell lines (CRISPRi-expressing RPMI-8226, AMO1, and KMS12-PE; $n = 3$ replicates per cell line
per sgRNA) reveals minimal changes in the cell surface proteome beyond CD38 knockdown at
significance cutoff of $p < 0.05$ and $\log_2$ fold-change $>|1.5|$. **C.** Integrated analysis of cell surface
proteomics and mRNA-seq ($n = 2$ per cell line per guide) across three cell lines reveals the only
consistent change at both protein and transcript level after *CD38* knockdown is *THY1*/CD90
upregulation. $\log_2$ fold-change cutoff $= |1.5|$.

FIGURE 5

1
2
Figure 5. ATRA drives CD38 upregulation with limited additional cellular impact while Aza
leads to a broad interferon-mediated response. A. Integrated mRNA-seq (n = 2 per drug
treatment) and cell surface proteomics (n = 2 per drug treatment) across RPMI-8226 treatment with
10 nM all-trans retinoic acid (ATRA), 2 μM azacytidine (Aza), and 10 nM panobinostat (Pano).
All plots are in comparison to control replicates treated with 0.1% DMSO. Doses chosen are based
on those previously published to lead to CD38 upregulation for each agent. Data points shown are
for proteins and genes corresponding to Uniprot-annotated membrane-spanning proteins. Log$_2$-fold
change cutoffs shown at |0.5| for ATRA and |2.0| for Aza and Pano to increase clarity of plots given
many fewer changed genes with ATRA treatment. **B.** RNA-seq for same samples with ATRA or
Aza treatment vs. DMSO but here showing all mapped genes, not just those annotated as
membrane-spanning. Significance cutoff at $p < 0.05$ with log$_2$ fold-change cutoff set at |0.8| to
illustrate prominent differences above this level in transcriptome alteration after either ATRA or
Aza treatment. **C.** KEGG analysis of genes from RNA-seq dataset meeting cutoff criteria of $p <$
0.05 and log$_2$ fold-change >0.8 after Aza treatment.

FIGURE 6

A

B

Figure 6. Unbiased phosphoproteomics reveals downregulation of proliferative signaling after daratumumab treatment. A. RPMI-8226 cells were treated with 20 μM daratumumab or IgG1 isotype control for 20 minutes ($n = 3$ each) and then harvested for unbiased phosphoproteomics with immobilized metal affinity chromatography enrichment for phosphopeptide enrichment. Plot displays results of Kinase Substrate Enrichment Analysis, indicating modest decrease in phosphorylation of numerous predicted substrates of MAPK pathway kinases as well as cyclin-dependent kinases (cutoffs of $p < 0.05$, $\log_2$ fold-change $>|0.5|$). **B.** Western blot in RPMI-8226 of MAPK (ERK1/2) (Thr202/Tyr204) relative to total MAPK demonstrates modest decrease in MAPK phosphorylation after 5, 10, or 15 min daratumumab (Dara) treatment; magnitude of change normalized to IgG1 control at each time point (red) appears consistent with phosphoproteomic data. **C.** Western blot of MM.1S cells treated with daratumumab and blotted for p-AKT (Ser473) and total AKT, with quantification of p-AKT relative to total AKT and normalized to IgG1 at each time point. All images representative of two independent Western blots.