1    **NO-stressed *Y. pseudotuberculosis* have decreased cell division rates in the mouse spleen**

3    Running title: Fluorescence dilution detects slowed bacterial growth

8    Bessie Liu[1], Rezia Era D. Braza[1], Katherine L. Cotten[1], Robert K. Davidson[1], Kimberly M.

9                                    Davis[1,*]

13    [1]W. Harry Feinstone Department of Molecular Microbiology and Immunology

14        Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

17    *Corresponding author:

18    E-mail: kdavi140@jhu.edu

**Abstract**

Fluorescence dilution approaches can detect bacterial cell division events, and can

detect if there are differential rates of cell division across individual cells within a

population. This approach typically involves inducing expression of a fluorescent protein, and

then tracking partitioning of fluorescence into daughter cells. However, fluorescence can be

diluted very quickly within a rapidly replicating population, such as pathogenic bacterial

populations replicating within host tissues. To overcome this limitation, we have generated

two revTetR reporter constructs, where either mCherry or yellow fluorescent protein (YFP) is

constitutively expressed, and repressed by addition of tetracyclines, resulting in fluorescence

dilution within defined timeframes. We show that fluorescent signals are diluted in replicating

populations, and that signal accumulates in growth-inhibited populations, including during nitric

oxide exposure. Furthermore, we show that tetracyclines can be delivered to the mouse

spleen during *Yersinia pseudotuberculosis* infection, and defined a drug concentration that

results in even exposure of cells to tetracyclines. We then used this system to visualize bacterial

cell division within defined timeframes post-inoculation. We detected growth attenuation of the

revTetR-mCherry strains within mouse tissues, however data suggested heightened NO exposure

correlated with heightened mCherry signal. We were able to restore normal bacterial growth with

revTetR-YFP, and use this strain to show that heightened NO exposure correlated with

heightened YFP signal, indicating decreased cell division rates within this subpopulation *in vivo*.

This revTetR reporter will provide a critical tool for future studies to identify and isolate slowly

replicating bacterial subpopulations from host tissues.

**Introduction**

Bacterial growth and metabolism have been the focus of decades of research, in part because we need to learn more about the pathways utilized by bacteria for growth to be able to target these essential processes with antibiotics or novel therapeutics. It is also well appreciated that bacteria with slowed metabolic rates are less susceptible to antibiotics (1-4). Earlier studies focused on growth of bacteria and antibiotic susceptibility under different laboratory conditions (1, 5-7); more recently, there has been a focus on studying bacterial growth and metabolism within host tissues, to more closely approximate bacterial growth conditions at the point where antibiotics or other therapeutics would be administered (8-10). Importantly, bacterial pathogens may utilize a range of different nutrients for growth within host tissues, and individual bacteria, or subsets of bacteria, may concurrently utilize different metabolites to promote their growth (10-14). Some of these substrates may support faster bacterial replication rates, and others, slower replication rates. Nutrient availability and the presence of host-derived antimicrobials, in addition to interactions with different immune cells subsets, can also generate complex microenvironments within host tissues, which can result in differences in growth rates across bacterial populations (12, 15-18).

When treating a bacterial infection, differences in bacterial growth rates can dramatically impact antibiotic efficacy. Slow-growing subsets of bacteria are inherently less susceptible to antibiotic treatment, due to decreased metabolic rates (1, 3, 4). This is termed antibiotic tolerance, a transient, phenotypic change in a subset of bacterial cells within a population, which alters antibiotic susceptibility (19, 20). Many of these experiments have been performed in bacteriological media, so it still remains largely unclear which pathways promote decreased antibiotic susceptibility and slowed growth within host tissues. To identify these pathways,

investigators will need to isolate antibiotic tolerant cells from the host environment for

downstream analyses. The specific bacterial pathways linked to tolerance may depend in part on

the antibiotic and the mode of action, or the environment around the bacteria during the

antibiotic exposure (18, 21, 22).

Molecular tools currently exist to identify and isolate slow-growing bacterial cells (8, 23-

26), but there are also limitations with some of these approaches that need to be addressed to

apply them within the host environment. One of the main approaches is utilizing fluorescence

dilution to identify individual cells that have divided within a certain timeframe. Typically,

fluorescence dilution approaches involve inducing expression of a fluorescent protein then

removing the inducer, and detecting changes in the amount of fluorescent protein within cells as

a readout for cell division events and protein partitioning into daughter cells (27, 28). Many

research groups have successfully utilized this approach to identify dividing and non-dividing

bacterial cells, by inducing fluorescent expression prior to introducing bacteria into tissue culture

or infection models (27-30). For some bacterial infection models, this approach can easily be

applied, because the size of the colonizing population of bacteria is relatively large, and the

number of cell division events within host tissues is relatively small (24, 25, 27). However, in

other infection models, only a few bacterial cells establish infection, and these cells go on to

replicate many times, which would quickly dilute out fluorescent signals (17, 31-33).

Furthermore, it is difficult to deliver inducer compounds into host tissues at sufficient levels to

modulate gene expression, and more challenging to remove these compounds to allow for

subsequent fluorescence dilution.

*Yersinia pseudotuberculosis* is an enteric pathogen that spreads systemically from

intestinal tissues to colonize deep tissue sites, such as the spleen (34-36). There is a strong

population bottleneck within intestinal tissues, and few bacterial cells spread systemically to

colonize deep tissue sites (37). Within deep tissues, individual bacteria found clonal clusters of

replicating extracellular bacteria, termed microcolonies or pyogranulomas (17, 38, 39).

Microcolonies can grow to contain hundreds to thousands of bacteria within a single replication

site, despite recruitment of circumscribing layers of neutrophils and monocytes (17, 39, 40).

Recent studies have shown that subpopulations of bacteria are present within microcolony

structures, and that subsets of bacteria stressed by the presence of nitric oxide (NO)

preferentially survive treatment with doxycycline, suggesting they may represent a slow-growing

subset of cells (17, 22). However, it has been difficult to determine growth rates in this model

system, since fluorescence dilution approaches cannot typically be used when the bacterial

population contracts and expands so dramatically.

To overcome these limitations, we have generated revTetR reporter constructs, where

exposure to tetracyclines will inhibit additional fluorescent gene expression, resulting in

fluorescence dilution. These constructs will promote fluorescent signal accumulation in non-

dividing cells, and fluorescent protein will be diluted into dividing cells, thus allowing us to

differentiate between dividing and non-dividing subpopulations of bacteria. Tetracyclines can

also easily be delivered into our mouse model of *Yersinia pseudotuberculosis* systemic infection

(22), which allows us to visualize fluorescence dilution as a measure of cell division at specific

stages of infection and within specific timeframes. Here, revTetR constructs allow us to ask if

responses to host-derived stresses, specifically nitric oxide (NO), impacts *in vivo* cell division

rates. These reporter constructs will be a useful tool for future studies to isolate and characterize

bacterial subpopulations with differential growth rates.

**Results**

**The revTetR reporter can be used to visualize differential bacterial growth rates *in vitro***

**with mCherry fluorescence dilution.**

To detect differential cell division rates within host tissues, we generated a fluorescence

dilution reporter system by modifying our recently described TetON fluorescent reporter with

stable mCherry (*tetR::P_{tetA}::mCherry*) (22). The revTetR reporter (*revtetR::P_{tetA}::mCherry*) was

constructed by introducing two mutations (E15A and L17G) into the wild-type *tetR* gene. These

two mutations were predicted to cause a reversal of TetR activity resulting in repression of $P_{tetA}$

specifically in the presence of tetracycline and tetracycline derivatives (41, 42). Therefore,

revTetR is a tetracycline-responsive reporter that allows cells to constitutively express mCherry

in the absence of tetracycline derivatives, and subsequent addition of tetracyclines should inhibit

additional mCherry transcription **(Figure 1A)**. This reporter can be used to compare growth rates

of bacteria via a fluorescence dilution approach (27, 28). For example, in cultures treated with

tetracycline, fast-growing cells will continue to divide, and the existing mCherry signal will be

partitioned between daughter cells with every replication event **(Figure 1B)**. However, slow-

growing or nondividing cells will not go through the same number of replication events and

therefore will not undergo fluorescence dilution **(Figure 1B)**.

To confirm that the mCherry fluorescence from the revTetR reporter dilutes into actively

dividing cells, we generated a reporter strain containing the revTetR reporter alongside a plasmid

containing *dps::gfp-ssrA* (revTetR *dps*) (16). Dps, a ferritin-like iron-sequestering protein,

accumulates within bacteria in stationary phase in response to multiple different stresses and

protects DNA from oxidative damage (43-45). Slow-growing cells have increased levels of the

*dps* (GFP) reporter signal (16). If the revTetR reporter is functioning as predicted, we would

expect to detect heightened revTetR (mCherry) in cells expressing *dps,* identifying these cells as

slow-growing, non-dividing cells within the population. revTetR *dps* cells were grown into

exponential phase then treated with increasing doses of anhydrotetracycline (ATc), a tetracycline

derivative that binds TetR or revTetR but has limited ribosomal targeting (46, 47). Aliquots of

each sample were measured for optical density or absorbance ($A_{600nm}$) and fluorescence using a

microplate reader. The absorbance values increased across all treatment groups, and ATc did not

have a detrimental effect upon growth rate with either 0.1µg/ml or 1µg/ml treatment **(Figure**

**1C)**. However, 2µg/ml ATc treatment did slightly inhibit growth, resulting in significantly lower

OD values at 3h and 4h post-treatment (hours, h). mCherry fluorescence values were normalized

to cell number by dividing the background-subtracted fluorescence value from each well by the

corresponding OD value. The resulting value was further normalized to the untreated sample

value for each time point (set at 100%). Following 2h of ATc treatment, all treated groups had

significantly reduced mCherry fluorescence relative to untreated cells **(Figure 1D).** At 4h post-

treatment, the fluorescent signal for all treated groups decreased by approximately 50% relative

to the signal of the untreated group, showing that ATc successfully induces dilution of mCherry

with each replication event **(Figure 1D)**. A 50% dilution in fluorescent signal should correlate

with one cell division event in all the cells within the population, suggesting the doubling time

was approximately 4h. This is slow for *Y. pseudotuberculosis* strains, and was the first indication

that the high levels of mCherry expression associated with this revTetR construct may impact

bacterial growth. We noted that 2µg/ml ATc treatment resulted in significantly higher mCherry

**(Figure 1D)** and *dps* reporter signal **(Figure 1E)** relative to 0.1µg/ml and 1µg/ml treatment,

again suggesting the 2µg/ml dose may slightly inhibit growth, and providing evidence that

slowed growth can be detected by an increase in mCherry signal. Cells from the final time point

(4h) were fixed and imaged using fluorescence microscopy to quantify single cell fluorescence

using the sum single cell signal for each channel. The average signal of the untreated group

(0µg/ml) was significantly higher than the average signal of the group treated with the lowest

dose of ATc (0.1µg/ml), again showing that signal dilution was successful **(Figure 1F)**. 1µg/ml

ATc treated cells had mCherry fluorescence similar to the 0.1µg/ml treated group. However, at

the highest dose of ATc (2µg/ml), the average mCherry signal signficantly increased **(Figure

1F)**, and we found there were also significant increases in *dps* reporter expression with

increasing doses of ATc **(Figure 1G)**, suggesting that high concentrations of ATc may have an

inhibitory effect on bacterial cells. Slower growing cells with heightened *dps* reporter expression

should also exhibit heightened revTetR-mCherry signal, and we confirmed these signals were

correlated under all treatment conditions, based on significantly non-zero slopes of linear

regression best-fit lines **(Figure 1H)**. Collectively these data indicate that revTetR fluorescence

is accumulating in slow-growing cells, and that actively growing cells are partitioning mCherry

into daughter cells, resulting in significant decreases in mCherry signal.

**Growth inhibition with doxycycline promotes mCherry fluorescence accumulation.**

To further test whether growth-inhibited cells exhibit heightened revTetR-mCherry

fluorescence, cells of the revTetR *dps* strain were sub-cultured to exponential phase and treated

with increasing doses of the antibiotic, doxycycline (Dox). Aliquots of each sample were

measured for optical density (OD) by absorbance ($A_{600nm}$) and fluorescence at hourly timepoints

using a microplate reader. Cells exposed to 1µg/mL Dox had significantly decreased growth

rates, however the doses less than 1µg/ml (0.01-0.1µg/ml) were subinhibitory, and did not

impact growth **(Figure 2A)**. mCherry fluorescence was initially heightened with 1µg/ml Dox

treatment **(Figure 2B)**, and *dps* reporter signal was slightly, but not significantly, elevated

**(Figure 2C)**, consistent with growth inhibition at this treatment concentration. Three hours after

treatment, the mCherry signal in all treatment groups was significantly below untreated values,

including the 1µg/ml treatment group **(Figure 2B)**, suggesting that growth may resume after

initial growth inhibition. Single cell analysis was performed by fluorescence microscopy, and the

sum single cell fluorescence values were graphed for each treatment group. Dilution of mCherry

signal could be seen with treatment of 0.01µg/mL Dox, but as the dosage increased, the mCherry

signal within cells increased, indicating signal accumulation and stalled growth **(Figure 2D)**. The

mCherry fluorescent signal was significantly higher in cells treated with 1µg/ml Dox compared

to untreated cells (0µg/ml), suggesting that growth inhibition resulted in significant revTetR-

mCherry fluorescence accumulation. Again, we observed a correlation between *dps* reporter

signal and revTetR-mCherry signal under all treatment conditions, confirming that slower

growing cells have heightened revTetR fluorescence **(Figure 2E)**.

**Nitric oxide-stressed cells divide at a slower rate than unstressed cells.**

194            It has been shown recently that exposure to nitric oxide (NO) results in cell division

arrest, through collapse of the FtsZ cell division machinery (48). Our results have also suggested

NO-stressed *Yersinia pseudotuberculosis* preferentially survive doxycycline treatment during

growth in the mouse spleen (22), which is consistent with a hypothesis that NO causes cell

division arrest within host tissues, and subsequently promotes decreased antibiotic susceptibility.

To test this hypothesis, we generated a strain containing the revTetR-mCherry reporter and

*P_{hmp::}gfp*, which can be used to mark NO-stressed cells (16, 17).

The revTetR-mCherry $P_{hmp::}gfp$ strain was treated with the slow-releasing NO donor

compound DETA-NONOate for 2h, then further treated with ATc (1µg/ml). The $P_{hmp::}gfp$

reporter was used to detect NO exposure, and to mark the level of stress experienced by

individual cells, which can be variable across the population in response to DETA-NONOate

(16, 17). Growth inhibition during treatment with the NO donor, DETA-NONOate, was

confirmed based on a significantly lower absorbance value in treated cultures at the 4h timepoint

**(Figure 3A)**. Induction of *hmp* in response to the NO donor was also confirmed across all

treatment groups **(Supplemental Figure 1)**. In the absence of ATc treatment, NO-exposed cells

increased in mCherry signal over time, which is consistent with growth inhibition and decreased

protein turnover **(Figure 3B)**. In cells treated with NO and ATc, the revTetR-mCherry

fluorescence decreased between 0h and 2h, indicating some cell division within this timeframe,

but signal slightly increased between 2h and 4h ATc treatment indicating cell division arrest

**(Figure 3B)**. This timing was consistent with the growth inhibition observed at 4h in Figure 3A.

In contrast, cells grown in the absence of NO treatment had a significant drop in mCherry

fluorescence between 2h and 4h, suggesting significant levels of cell division events within this

timeframe **(Figure 3B)**.  Consistent with previous literature, these results indicate that NO

treatment results in growth arrest, and reduced levels of cell division events (48), as detected by

revTetR-mCherry.

To determine whether heightened NO stress correlated with a reduced level of cell

division events, single cell mCherry (revTetR) and GFP (Hmp) values from NO-treated cells

were compared in the absence and presence of ATc treatment using correlation plots and linear

regression analyses. In the absence of ATc treatment, the distributions of cells within the NO-

stressed bacterial populations were very similar in fluorescence across all timepoints **(Figure**

**3C)**. In the presence of ATc treatment, there was a clear shift in the NO-treated population

towards reduced mCherry signal over time **(Figure 3D)**. This suggested some cell division still

occurred in the presence of NO treatment, which was consistent with Figure 3B.  However, the

correlation between the two fluorescent signals strengthened over the timecourse based on

increased $R^2$ values **(Figure 3D)**. These data support the conclusion that cells experiencing

higher level of NO stress are dividing more slowly, and retaining higher levels of revTetR-

mCherry fluorescence.

**Low doses of doxycycline and anhydrotetracycline diffuse evenly across microcolonies and**

**do not inhibit bacterial growth.**

One of the major goals for generating the revTetR reporter was to set-up a system to

detect bacterial cell division events within host tissues over defined periods of time. To do this,

we would need to administer tetracyclines during infection, ensure that the dosage used would

not impact bacterial growth, and also diffuse evenly across microcolonies to allow for accurate

detection of signal dilution and cell division. We have previously developed a model of

doxycycline treatment of *Y. pseudotuberculosis* infection, and found that a single 40mg/kg

injection of doxycycline significantly reduces bacterial CFUs and also accumulates at a higher

concentration at the periphery of microcolonies (22). We hypothesized that lowering the

tetracycline dose may allow us to administer sub-inhibitory levels of antibiotic that would still

modulate the revTetR reporter, while allowing for more even diffusion of the antibiotic across

microcolonies.  Our rationale was that lower antibiotic concentrations would be more readily

eliminated from circulation, and so antibiotic movement into the spleen would occur within a

more defined timeframe, limiting continued accumulation.

To identify a tetracycline concentration that results in equal diffusion across

microcolonies, a previously characterized reporter, $P_{tetA}::mCherry\text{-}ssrA$ (TetON), was used for

tetracycline detection (22). The TetON reporter contains the wild-type *tetR* gene; addition of

tetracyclines induce the production of mCherry, and therefore levels of mCherry can be used as a

readout for antibiotic exposure **(Figure 4A)**. C57BL/6 mice were inoculated intravenously with

the TetON *gfp*$^+$ strain, which has constitutive expression of GFP alongside the TetON reporter,

allowing for visualization of relative reporter expression. A single dose of 4mg/kg Dox or

4mg/kg ATc was administered intraperitoneally at 48 hours post-inoculation. This timepoint was

chosen for antibiotic administration since we can detect subsets of bacteria responding to host-

derived nitric oxide at this timepoint, thus allowing us to ask whether stressed subpopulations of

bacteria are dividing more slowly than unstressed bacteria.

At 4h or 24h post-treatment, spleens were harvested to quantify CFUs and quantify

reporter expression by fluorescence microscopy. Bacterial numbers continued to increase

between 4h and 24h post-treatment with 4mg/kg doses of either ATc or Dox, confirming this

dosage does not interfere with bacterial growth **(Figure 4B)**. Microcolonies were then visualized

by fluorescence microscopy to determine if ATc and Dox diffuse evenly across microcolonies at

this dosage.  The centroid and periphery of each microcolony were quantified for mCherry and

GFP signal, and the ratio of mCherry/GFP was graphed. There were no significant differences in

mCherry/GFP ratios between the centroid and periphery of microcolonies treated with either

ATc or Dox, suggesting that each drug was diffusing evenly across microcolonies **(Figure 4C)**.

The mCherry/GFP ratios remained comparable at the centroid and peripheries even at 24 hours

after treatment, showing that throughout the infection time course, there was even diffusion of

the drugs **(Figure 4C, 4D)**. These data here suggest that lowering the tetracycline dosage 10-fold

results in even diffusion of a sub-inhibitory dose of tetracyclines, which will allow us to

modulate revTetR reporter expression in all cells across the microcolony without impacting

bacterial growth.

**revTetR signal appears to increase in both the centroid and periphery of microcolonies.**

To determine if there are slowly-dividing subpopulations of bacteria within

microcolonies, we infected mice with the revTetR-mCherry $gfp^+$ strain, which constitutively

expressed $gfp$, and treated mice with ATc at 48h post-inoculation or left mice untreated. Mice

were sacrificed at either 4h or 24h post-treatment, and spleens were harvested to quantify CFUs

and visualize reporter expression by fluorescence microscopy. Bacterial CFUs were comparable

between ATc-treated and untreated mice at both timepoints, suggesting that ATc treatment did

not impair bacterial growth (**Figure 5A**). However, we did not see an increase in CFUs between

4h and 24h, which suggests there was limited, or variable, bacterial growth within this selected

timeframe for this bacterial strain. To determine if there were differences in reporter expression,

mCherry signal was quantified relative to the constitutive GFP signal at the periphery and

centroid of individual microcolonies. Since we have previously shown that bacteria experience

heightened levels of NO stress and express higher levels of the T3SS at the periphery of

microcolonies, we were expecting to see heightened levels of mCherry signal relative to GFP at

the periphery of microcolonies. At 4h post-treatment, we did not observe any differences in

mCherry signal at the periphery of microcolonies compared to the centroid, in the absence or

presence of ATc treatment (**Figure 5B**). This suggested there may not have been many cell

division events within the first 4h of treatment, which is why tissue was also harvested 24h post-

treatment. At 24h post-treatment, untreated mice had significantly lower levels of mCherry

signal at the periphery of microcolonies, and ATc treated mice also had significantly lower

mCherry signal at the periphery, indicating we were not seeing increased mCherry signal at the

periphery of microcolonies as expected **(Figure 5C, 5D)**. It did appear that ATc treated

microcolonies had higher mCherry signals at both the centroid and periphery relative to

untreated microcolonies, although this was not statistically significant. These results suggest that

revTetR signal remained high after ATc treatment, indicating there may be slowly dividing cells

present within microcolonies. Without much bacterial growth across these timepoints in general,

it was difficult to assess whether slow-growing cells were present in a specific spatial location or

part of a subpopulation without an additional reporter alongside revTetR.

**NO stress correlates with heightened mCherry signal accumulation and slowed cell division**

**rates.**

Hmp$^+$ bacteria responding to NO stress preferentially survive doxycycline treatment,

suggesting they may represent a slow-growing subpopulation (22). Consistent with this, in

culture, NO-stressed cells accumulated heightened mCherry signal, suggesting they were

dividing more slowly than unstressed cells **(Figure 3)**. However, in our mouse model it was

difficult to determine if peripheral cells of the microcolony had heightened revTetR signal

without marking the peripheral subpopulation with the *hmp* reporter. To determine if NO-

stressed bacteria are dividing more slowly than unstressed cells within host tissues, we infected

mice with the revTetR-mCherry $P_{hmp}::gfp$ strain and treated mice with ATc at 48h post-

inoculation, or left mice untreated. Spleens were harvested at 4h and 24h post-treatment to

quantify CFUs and visualize reporter expression by fluorescence microscopy. CFUs within ATc-

treated mice did not differ significantly from CFUs within untreated mice; again, we noted that

there was not significant growth based on total CFUs between 4h and 24h, but anticipated that

there would be sufficient replication at the level of individual microcolonies to assess differences

in revTetR signal **(Figure 6A)**.

Periphery and centroid measurements were taken from each microcolony image and

combined into a single periphery/centroid ratio for each microcolony to allow for comparison

across treatment groups. At 4h post-ATc treatment, the average ratio for the *hmp* reporter signal

was above a value of 1, indicating higher expression in the peripheral cells **(Figure 6B)**. The

average ratio for revTetR-mCherry values was close to 1 in both ATc-treated and untreated mice,

suggesting equal expression in peripheral and centroid cells and very little cell division in this

time frame **(Figure 6B)**. At 24h post-ATc treatment, GFP ratios remained above 1, indicating

high peripheral expression of the *hmp* reporter **(Figure 6C)**. The revTetR-mCherry ratio

appeared to increase slightly, but remained below a value of 1, and was not significantly

different from untreated mice **(Figure 6C)**. This again suggested there was little cell division

within this timeframe, since we could not detect much impact of the ATc treatment. We then

used linear regressions to determine if there was a correlation in fluorescent signals at distinct

spatial locations within the microcolony, and overall saw little impact of the ATc treatment on

the fluorescence of the cells **(Figure 6D-H)**. The exception was the analysis at the periphery 24h

post-treatment, where ATc treatment resulted in a slight positive correlation suggesting $Hmp^+$

cells were less likely to divide **(Figure 6G, 6H)**. The limited levels of cell division with this

strain *in vivo* made these results difficult to interpret, and we then sought to modify the reporter

to generate a new revTetR construct with normal growth kinetics *in vivo*.

**Dilution of the revTetR-YFP signal coincides with the replication of fast-growing cells.**

To address the growth issues in the revTetR-mCherry strain evident in previous

experiments, we modified the revTetR reporter to express YFP instead of mCherry

(*revtetR::P_{tetA}::yfp)*. We surmised that the growth issues were due to the more toxic and

disruptive nature of constant mCherry overexpression in the absence of tetracycline derivatives,

and YFP is known to be less toxic for cells (49, 50).

To determine if revTetR-YFP strains exhibit normal growth kinetics, and confirm NO

stress is sufficient to slow growth and promote YFP signal accumulation, we generated a strain

containing the revTetR-YFP reporter and *P_{hmp}::mCherry* (*revtetR::P_{tetA}::yfp P_{hmp}::mCherry*).

Cells were grown in the presence and absence of the NO donor compound DETA-NONOate for

2h, then treated with ATc (1µg/ml) as described above. Aliquots were taken at the indicated

timepoints after ATc addition to quantify single cell fluorescence by microscopy. Growth

inhibition due to the NO donor was confirmed by negligible signal dilution between 0h and 2h

post-ATc treatment **(Supplemental Figure 2A)**, and all treated cultures responded to the NO

donor based on induction of *hmp* **(Supplemental Figure 2B)**. Cells exposed to NO and ATc had

a slight signal dilution between 0-2h and 2-4h, suggesting some continued cell division, however

cells grown in the absence of NO showed dramatic YFP signal dilution between 0-2h

**(Supplemental Figure 2A)**. This suggested that the revTetR-YFP construct behaves similarly to

revTetR-mCherry, and can be used to mark growth-arrested cells. However, unstressed cells (-

NO, +ATc) no longer had signal dilution between 2-4h post-ATc treatment, suggesting growth

was slowing in these cultures. Since cells were incubated with NO for 2h before ATc treatment,

this indicates that the cells were beginning to enter stationary phase by the 4h timepoint (6h of

growth). These changes in growth kinetics indicate that the revTetR-YFP strain may grow faster

than the revTetR-mCherry strain in culture. Additionally, the signal dilution between 2-4h in the

+NO +ATc group **(Supplemental Figure 2A)** suggests cells are recovering from NO exposure

and are starting to replicate after 6h exposure. This is consistent with our previous work (16), and

consistent with the presence of some Hmp$^{low}$/mCherry$^{low}$ cells in the +NO +ATc group at 4h

**(Supplemental Figure 2B)**.

To ensure we were capturing the timeframe of growth arrest, we modified the *in vitro* NO

donor experiment to shorten the NO exposure. Cells containing the revTetR-YFP reporter and

*P$_{hmp}$::mCherry* were treated with the NO donor compound for only 30 minutes before ATc

addition, and single cell fluorescence was quantified via microscopy. Cells grown in the absence

of ATc treatment continued to accumulate YFP signal, which was unsurprising **(Supplemental**

**Figure 3A)**. Heightened *hmp* induction in the +NO groups compared to the -NO groups

confirmed NO exposure throughout the timecourse **(Supplemental Figure 3B)**. With this

experimental protocol, cells exposed to the NO donor and ATc treatment increased in revTetR-

YFP signal between 0-2h, and only a slight decline between 2-4h, indicating overall growth

arrest **(Figure 7A)**. Cells grown in the absence of the NO donor with ATc treatment showed

continuous signal dilution, indicating continuous growth and replication **(Figure 7A)**. These

results suggest that in this faster growing strain, YFP signal dilution does correspond with

bacterial cell growth, and signal accumulation corresponds with growth arrest.

Since our results above indicate the growth kinetics are different between the revTetR-

YFP and revTetR-mCherry strains, we estimated the length of time it takes for 100% of the cells

in culture to undergo one full replication cycle, as indicated by a 50% dilution of the revTetR

signals. Calculations were completed using data from unstressed cells: -NO +ATc and -NO -ATc

groups. YFP signals from our single cell microscopy data were normalized to 100% at 0h, then

the ATc-treated group (-NO +ATc) data was normalized to the untreated control group (-NO -

ATc) to account for signal dilution over time. Using a linear regression equation, the revTetR-

YFP strain was estimated to undergo one replication cycle, or doubling, in 1.75h **(Figure 7B)**.

This is consistent with normal growth kinetics of *Y. pseudotuberculosis* at $37^{\circ}$ C (16). We then

confirmed that the revTetR-mCherry strain does indeed grow slower, with an estimated 3.31h for

one replication cycle **(Figure 7C)**. Together, these data indicate that the revTetR-YFP reporter

can be used to differentiate between dividing and non-dividing cells, and confirm the growth

issues seen in the revTetR-mCherry strains have been resolved with revTetR-YFP.

**Cells that lack NO stress rapidly dilute revTetR-YFP signal indicating faster cell division**

**rates.**

Our previous work indicates that cells at the periphery of the microcolony respond to NO

stress and may be slow-growing, since Hmp$^+$ cells had reduced susceptibility to a ribosome-

targeting antibiotic (22). While our *in vitro* data support this **(Figure 7)**, our previous study using

a modified TIMER protein (DsRed$_{42}$) to detect slow-growing bacteria suggested that NO stress is

transient within host tissues (16). At late timepoints during infection, Hmp$^-$ cells at the centroid

of microcolonies were instead slow-growing due to nutrient limitation (16).  In our mouse

infections with the revTetR-mCherry $P_{hmp}::gfp$ strain above, it appeared that cells responding to

NO stress (Hmp$^+$, GFP$^+$) may have higher mCherry levels **(Figure 6G)**. Due to growth issues

with the strain, it was difficult to interpret these results. To clarify if the subset of NO-exposed

cells are truly growing at a slower rate than unstressed cells within host tissues, we infected mice

with the revTetR-YFP $P_{hmp}::mCherry$ strain, then treated with ATc (+ATc) at 48h post-

inoculation, or left mice untreated (unt). We confirmed the revTetR-YFP $P_{hmp}::mCherry$ strain

was growing well within host tissues based on significant increases in CFUs in the spleen,

comparing between 4h and 24h post-ATc treatment **(Figure 8A)**. Spleens were processed for

fluorescence microscopy, periphery and centroid measurements were taken from each

microcolony, and combined into a single periphery/centroid ratio to allow for comparison across

treatment groups. At 4h post-treatment, the average ratio for the *hmp* (mCh) reporter signal was

above a value of 1 in both untreated (unt) and treated mice (+ATc), indicating higher expression

in the peripheral cells, as expected. The average ratio of revTetR-YFP signal was close to 1 in

untreated mice, suggesting equal expression in peripheral and centroid cells **(Figure 8B)**. In

contrast, ATc-treated mice had a ratio above 1, suggesting there is higher YFP expression in the

periphery relative to the centroid **(Figure 8B)**. Similar trends held true at 24h post-treatment,

although the *hmp* (mCh) reporter signal was higher in treated mice for unknown reasons **(Figure**

**8C)**.

An increased periphery/centroid ratio can mean either the periphery signal increased, or

conversely, the centroid signal decreased. Because of the nature of fluorescence dilution, we

would expect this means the centroid is diluting revTetR-YFP signal more rapidly than the

periphery, suggesting the centroid is more rapidly dividing. To assess this, raw YFP signal

values were compared across treatment groups and timepoints. At 4h post-ATc treatment, YFP

signals for untreated mice were not significantly different between the centroid and periphery

**(Figure 8D)**. The YFP signal in the centroid of ATc-treated mice was trending, but not

significantly lower than the periphery **(Figure 8D)**. However, the centroids of ATc-treated mice

were significantly lower than the centroids of untreated mice, suggesting cell division and signal

dilution at the centroid **(Figure 8D)**. At 24h post-ATc treatment, untreated mice had similar

levels of YFP signal at the centroid and periphery of microcolonies, while centroids of treated

mice had significantly lower levels of YFP when compared to treated peripheries or untreated

centroids **(Figure 8E)**. These results suggest significant cell division within the centroid of

microcolonies, and relative growth arrest at the periphery, since peripheral cells were not

changing in YFP signal **(Figure 8E)**. These conclusions are corroborated by the correlation of

revTetR-YFP and mCherry (Hmp) periphery/centroid ratios, where revTetR and Hmp exhibit a

weak correlation at 24h post-treatment, based on a significantly non-zero slope **(Figure 8F)**. It is

important to note that while cells in the periphery tend to have higher YFP signals in the treated

mice at 24h **(Figure 8F)**, some subsets of peripheral cells do not exhibit this high YFP signal

**(Figure 9)**. It is, however, striking, how much YFP dilution occurs at the centroid of

microcolonies, which strongly supports the conclusion that the centroid is rapidly dividing

**(Figure 8E, 9)**. Overall, this microcolony phenotype is consistent with our previous findings that

only small subsets of peripheral cells actively respond to NO stress at any given timepoint (16),

which appears to correlate with detection of growth arrest with revTetR-YFP.

**Discussion**

In this study, we have developed revTetR constructs that can be utilized to detect

bacterial cell division events within host tissues using a fluorescence dilution approach. We have

shown both revTetR-mCherry and revTetR-YFP constructs result in heightened fluorescent

signal in growth-arrested cells in culture, and have also shown that nitric oxide (NO) stress is

sufficient to promote increased fluorescent signal in culture **(Figures 3, 7)**. This was consistent

with previous studies showing NO exposure results in an arrest of cell division (48). Our *in vivo*

results in the mouse model of infection were initially less clear; our revTetR-mCherry strains had

notable slowed growth within host tissues **(Figures 5A, 6A)**, which made it difficult to

differentiate dividing and non-dividing cells. Experiments with revTetR-mCherry alongside a

*hmp* reporter hinted that Hmp$^+$ cells may be slowly dividing **(Figure 6G)**, but this conclusion

remained unclear until experiments were completed with the revTetR-YFP $P_{hmp}::mCherry$ strain.

Using revTetR-YFP, we detected clear differences in cell division rates across microcolonies.

Bacterial cells at the centroid are clearly replicating more quickly and diluting YFP signal, while

we saw YFP signal was maintained within peripheral cells **(Figures 7D, 7E)**. Some Hmp$^+$ cells

retained YFP signal while others did not, which we believe reflects transient NO exposures

within host tissues (16).

The Hmp$^+$ subpopulation was a focus of this study primarily because we knew this

subpopulation preferentially survives doxycycline treatment in the *Y. pseudotuberculosis* mouse

model of infection. We hypothesized this was due to increased levels of growth arrest as a result

of NO exposure (22, 48). However, it is important to note that the Hmp$^+$ subpopulation is a

complex mixture of cells, that also includes subsets expressing high levels of the type-III

secretion system (T3SS) due to direct neutrophil contact (17). Based on the pattern of revTetR-

YFP signal within host tissues **(Figure 9)**, revTetR-YFP more closely correlates with Hmp

reporter expression than T3SS, but additional experiments could directly assess this. We also

chose to mark the Hmp$^+$ subpopulation here with a stable fluorescent protein to detect NO

exposure. However, we know that only subsets of this population are actively responding to NO

at any given timepoint (16), which adds further complexity to this subpopulation. It will be very

interesting to utilize this revTetR-YFP construct in future experiments to isolate bacterial cells

with slowed division rates from the host environment, and further characterize the pathways that

are associated with bacterial slowed growth within host tissues.

Importantly, we believe the cell division rates within microcolonies vary over the course

of infection, and revTetR-YFP will allow us to probe this in more detail by selecting different

timepoints over the course of infection for ATc treatment and infection endpoints. Here, we

chose to compare between 52h and 72h post-inoculation, which equates to +ATc 4h and 24h,

respectively. These were timepoints where we expected rapid expansion within the bacterial

population, allowing us to differentiate between slowly-dividing and quickly-dividing cells.

These timepoints worked well for distinguishing differential growth rates with revTetR-YFP, and

showed the centroid of microcolonies were dividing rapidly and diluting fluorescent signal

**(Figure 9)**. However, this does seem to contrast with some of our other results that indicate the

centroid is slow-growing (16). In Patel P, *et al.*, slowed growth was detected in the centroid of

microcolonies with a slow-folding DsRed variant, DsRed$_{42}$, specifically at 72h post-inoculation

(16). We hypothesize the timepoint is key here, and with the timepoints we have chosen in this

study, we are capturing a period of rapid growth that precedes nutrient depletion at later

timepoints. By switching from revTetR-mCherry to revTetR-YFP, we have generated strains

without growth attenuation, and also generated a construct where revTetR-YFP fluorescence

could be directly compared with $DsRed_{42}$ to better understand bacterial growth dynamics over

the course of infection.

Experiments with the revTetR-mCherry constructs have shown us that to truly detect

differences in cell division rates *in vivo*, we need to sample timepoints where there is rapid

expansion within the bacterial population. Based on our estimates in Figure 7, revTetR-YFP

strains double approximately every 1.75h during culture at $37^{o}$ C, while revTetR-mCherry strains

double approximately every 3.3h, and may take as long as 4h to divide, based on Figure 1D. The

doubling time, or replication rate, thus essentially establishes the sensitivity of the system, and

needs to be taken into account when choosing the timepoints for experiments. Conversely, by

completing full timecourses, researchers would be able to utilize these reporters to approximate

the growth rate of strains grown under different conditions. We believe this application could be

particularly useful in approximating the bacterial cell division rates within host tissues, and plan

to explore this in future experiments.

In this study, we compared revTetR-mCherry reporter signal during treatment with two

tetracycline derivatives, the active antibiotic, doxycycline (Dox), and the inactive derivative,

anhydrotetracycline (ATc). Both tetracycline derivatives bind to TetR and revTetR, however

ATc is predicted to bind these proteins with higher affinity, and has poor affinity to bacterial

ribosomes (46, 47). This allowed us to modulate revTetR with ATc without significantly

inhibiting translation, however we would expect to see translational inhibition during Dox

treatment, and did see evidence of this when detecting *dps* reporter signal alongside revTetR-

mCherry. Because Dox inhibits translation, the amount of *dps* signal did not significantly

increase with Dox treatment due to translational inhibition **(Figure 2C)**. However, we were able

to conclude that slow-growing or growth-arrested cells had higher levels of revTetR mCherry

signal using ATc and NO treatments.

We recently showed with a slow-folding fluorescent protein derivative ($DsRed_{42}$) that

NO stress was sufficient to promote fluorescence accumulation, but also found there was limited

overlap in slow-growing and NO-stressed populations within host tissues (16).  Destabilized and

stable versions of fluorescent proteins were used to show that NO stress was transient within host

tissues, and that few bacteria were actively responding and detoxifying NO at any given

timepoint (16).  However, the fact that $Hmp^+$ bacteria preferentially survive doxycycline

treatment suggests that, while NO exposure may be transient, there could also be sustained

changes within cells exposed to high levels of NO (22).  Although *Yersinia* repairs iron-sulfur

cluster-containing proteins in response to this stress (51), there may be long-term damage, or

even genetic changes that occur following NO exposure.  It would be very interesting to

determine if any genetic changes occur within microcolonies over the course of infection,

especially given these bacterial populations are clonal, founded by a single bacterium, and

presumed to be genetically homogenous (17).

**Materials and Methods**

*Bacterial strains and growth conditions.* *Y. pseudotuberculosis* IP2666 was used as the wild-type strain throughout. For *in vitro* experiments, overnight cultures were grown in LB at 26°C with rotation for 16 hours. Exponential phase cells for *in vitro* experiments were generated by sub-culturing overnight cultures by diluting 1:100 into fresh LB, and rotating at 37°C 2 hours. All *in vitro* experiments were performed by culturing at 37°C with rotation. For all mouse infections, bacterial inocula were grown overnight (16 hours) to post-exponential phase in 2xYT broth (LB, with 2x concentration of yeast extract and tryptone) at 26°C with rotation.

*revTetR reporter construction.* The revTetR reporter was generated by introducing specific point mutations into our previously described *tetR::$P_{tetA}$::mCherry* reporter construct in pMMB67EH (22). *tetR* was mutated by site-directed mutagenesis at two sites, E15A (GAG→GCU) and L17G (CUU→GGU), using 4 point mutations to generate the revTetR construct. An additional L25V mutation could not be successfully introduced. Primers were designed that contained the desired mutations, and were used to PCR amplify around the pMMB67EH plasmid using PfuUltra II high-fidelity polymerase (Agilent). PCR products were purified (QIAGEN) and transformed into electrocompetent DH5αλpir *E. coli*. Colonies were screened based on mCherry fluorescence in the absence of tetracyclines, plasmids were then isolated using the PureYield Plasmid Midiprep kit (Promega), and sequenced to confirm mutations. *revtetR::$P_{tetA}$::mCherry* plasmids were transformed into *Y. pseudotuberculosis* strains as previously described (52), to generate the following reporter strains: *revtetR::$P_{tetA}$::mCherry $P_{dps}$::gfp-ssrA, revtetR::$P_{tetA}$::mCherry gfp$^+$,* and *revtetR::$P_{tetA}$::mCherry $P_{hmp}$::gfp*. *$P_{dps}$::gfp-ssrA, gfp$^+$,* and *$P_{hmp}$::gfp* reporter constructs have been previously described (16), and are all inserted into the pACYC184 plasmid.

*revtetR::P~tetA~::yfp* was constructed by PCR amplifying *revtetR::P~tetA~* and fusing this to *yfp* using

overlap extension PCR. *revtetR::P~tetA~::yfp* was inserted into the pACYC184 plasmid, and

transformed into a *Y. pseudotuberculosis* strain containing the previously described

*P~hmp~::mCherry* plasmid (17).

*In vitro reporter characterization.* Once subcultures reached exponential phase (2h growth), the

indicated doses of doxycycline (Dox; 0.01 µg/mL, 0.1 µg/mL, or 1 µg/mL) or

anhydrotetracycline (ATc; 0.1 µg/mL, 1 µg/mL, or 2 µg/mL) were added to samples to promote

mCherry dilution, with one sample left untreated as a control. Samples were incubated at 37°C

with rotation throughout the time course, and time points of 0, 0.5, 1, 2, 3, and 4h post-antibiotic

treatment were taken. For each time point, 600 µL from each sample was pelleted and

resuspended in 600 µL of PBS. 200 µL of each sample was then pipetted into 3 wells of a black

walled, clear bottom 96 well plate, and analyzed using a Synergy Microplate Reader (Biotek

Instruments) for $OD_{600}$ and fluorescence signal measurements. Optical density ($A_{600nm}$) was used

to approximate cell number, and mCherry fluorescence was detected with 560nm

excitation/610nm emission. All experiments were repeated in triplicate to generate a total of

three independent data sets, or nine replicates. In strains of Yptb also containing a GFP reporter,

GFP fluorescence was detected with 480nm excitation/520nm emission.

*Induction of hmp using nitric oxide donor compound.* Overnight cultures were diluted 1:50 into

fresh LB with the nitric oxide donor compound DETA-NONOate (2.5 mM dose) or were left

untreated. After either 2 hours or 30 min of incubation, each culture was evenly split into two

separate tubes. Samples were then further treated with either ATc (1 µg/mL dose) or left

untreated, for a total of four different treatment groups: +NO/+ATc, +NO/-ATc, -NO/+ATc, -

NO/-ATc. Aliquots were taken every 2h after ATc treatment to quantify absorbance and

fluorescence, and cells were prepared for fluorescence microscopy, as described below. Fixed

cells were imaged and quantified for single-cell fluorescence measurements.

*Fluorescence microscopy: bacteria.* To visualize individual bacterial cells, 500 µL from each

sample was pelleted, resuspended in 4% paraformaldehyde (PFA), and incubated overnight at 4$^{o}$

C for fixation. Agarose pads were prepared by placing two pieces of tape approximately 1 cm

apart on slides, pipetting 25 µL of 1% agarose in PBS, and placing a glass coverslip over the

area. After leaving the agarose to solidify for 20 min, the coverslips and tape were peeled off and

the agarose was trimmed to 1 cm$^{2}$ squares. PFA-fixed bacteria were pelleted, resuspended in 50

µL PBS, and 5 µL of sample was pipetted onto each agarose pad. Bacteria were imaged with the

63x oil immersion objective, using a Zeiss Axio Observer.Z1 (Zeiss) fluorescent microscope

with Colibri.2 LED light source and an ORCA-R$^{2}$ digital CCD camera (Hamamatsu). Five

images were taken of each sample in distinct fields of view, using DIC, GFP, and mCherry

channels. Volocity image analysis software was used to specifically select individual bacterial

cells and quantify the fluorescent signal associated with each cell.

*Murine model of systemic infection.* The following strains were used for mouse infections: *Y.*

*pseudotuberculosis tetR::P$_{tetA}$::mCherry-ssrA gfp$^{+}$ (22), revtetR::P$_{tetA}$::mCherry gfp$^{+}$,* and

*revtetR::P$_{tetA}$::mCherry P$_{hmp::}$gfp,* and *revtetR::P$_{tetA}$::yfp P$_{hmp}$*::mCherry. Six to eight week old

female C57BL/6 mice were obtained from Jackson Laboratories (Bar Harbor, ME). All animal

studies were approved by the Institutional Animal Care and Use Committee of Johns Hopkins

University. Mice were injected intravenously with $10^3$ bacteria for all experiments. After 48

hours, mice were treated with the indicated dose (4mg/kg, equates to approximately 72μg in

100μl sterile PBS) of Dox or ATc, or left untreated. At the indicated timepoints, either 4h or 24h

post-treatment, mice were euthanized via both lethal dose of isoflurane and cervical dislocation,

and spleens were removed, halved, and processed for histology, or homogenized to quantify

CFUs.

*Histology.* Harvested spleen halves were fixed in 4% PFA overnight at $4^o$C, and were embedded

using Sub Xero OCT compound (Mercedes Medical), frozen on dry ice, and stored at -80°C.

Spleens were cross-sectioned into 10 um sections using a cryostat microtome; one representative

section was imaged per mouse. To visualize reporters, sections were thawed in PBS, stained with

Hoechst at a 1:10,000 dilution, washed in PBS, and coverslips were mounted using ProLong

Gold (Life Technologies). Images were taken of all microcolonies within each section with a 63x

oil immersion objective, using a Zeiss Axio Observer.Z1 (Zeiss) fluorescent microscope with

Colibri.2 LED light source, an Apotome.2 (Zeiss) for optical sectioning, and an ORCA-R$^2$ digital

CCD camera (Hamamatsu).

*Image analysis.* For *in vitro* experiments, Volocity image analysis software was used to select

single bacterial cells in each image with the 'find objects' function. The mean and sum signal

intensity was calculated for each object (single cell) for mCherry, GFP, and YFP channels. Sum

signal intensity values were used for revTetR reporter quantification. For *in vivo* experiments,

either Volocity or Fiji image analysis software was used to select regions of interest at the

centroid and periphery of each microcolony and quantify the fluorescent signal intensity of each

channel to generate relative signal intensities of fluorescent reporters. A small representative

point (~0.01pixels$^2$) was selected at the centroid and periphery of each microcolony, which

generated a 'signal intensity' value at that point that correlates with the sum. Centroid bacteria

were defined as surrounded only by other bacteria. Peripheral bacteria were defined based on

contact with host cells (**Figure 4D**). All data was analyzed and graphed, and all statistical

analyses performed, using GraphPad Prism 9.

**Author Contributions.**  Conceptualization: BL, RKD, KMD; Formal Analysis: BL, REB,

KMD; Funding Acquisition and Supervision: KMD; Investigation: BL, REB, KLC, KMD;

Methodology: BL, REB, KMD; Writing – Original Draft Preparation: BL, REB, KMD; Writing

– Review & Editing: BL, REB, KLC, RKD, KMD.

**Acknowledgments.**  We thank the members of the Davis lab, who provided feedback and

suggestions during the final steps of manuscript preparation. We also thank the members of the

Pekosz and Klein lab for their helpful feedback throughout this project. This work was supported

by NIAID grants 1K22AI123465-01 and 1R21AI154116-01A1 to KMD. REB is also supported

by training grant 2T32AI007417-26 through NIAID. The authors of this manuscript declare no

conflicts of interest.

**References**

1.  Balaban N, Merrin J, Chait R, Kowalik L, Leibler S. 2004. Bacterial persistence as a phenotypic switch. Science 305:1622-5.

2.  Goerke C, Wolz C. 2010. Adaptation of *Staphylococcus aureus* to the cystic fibrosis lung. Int J Med Microbiol 300:520-5.

3.  Harms A, Maisonneuve E, Gerdes K. 2016. Mechanisms of bacterial persistence during stress and antibiotic exposure. Science 354.

4.  Lopatkin A, Stokes J, Zheng E, Yang J, Takahashi M, You L, Collins J. 2019. Bacterial metabolic state more accurately predicts antibiotic lethality than growth rate. Nat Microbiol 4:2109-2117.

5.  Arnoldini M, Vizcarra I, Peña-Miller R, Stocker N, Diard M, Vogel V, Beardmore R, Hardt W, Ackermann M. 2014. Bistable expression of virulence genes in *Salmonella* leads to formation of an antibiotic-tolerant subpopulation. PLoS Biol 12:e1001928.

6.  Nguyen D, Joshi-Datar A, Lepine F, Bauerle E, Olakanmi O, Beer K, McKay G, Siehnel R, Schafhauser J, Wang Y, Britigan B, Singh P. 2011. Active starvation responses mediate antibiotic tolerance in biofilms and nutrient-limited bacteria. Science 334:982-986.

7.  Sánchez-Romero M, Casadesús J. 2014. Contribution of phenotypic heterogeneity to adaptive antibiotic resistance. Proc Natl Acad Sci 111:355-60.

8.  Claudi B, Spröte P, Chirkova A, Personnic N, Zankl J, Schürmann N, Schmidt A, Bumann D. 2014. Phenotypic variation of *Salmonella* in host tissues delays eradication by antimicrobial chemotherapy. Cell 158:722-33.

9.  Kaiser P, Regoes R, Dolowschiak T, Wotzka S, Lengefeld J, Slack E, Grant A, Ackermann M, Hardt W. 2014. Cecum lymph node dendritic cells harbor slow-growing bacteria phenotypically tolerant to antibiotic treatment. PLoS Biol 12.

10. Manina G, Dhar N, McKinney J. 2015. Stress and host immunity amplify *Mycobacterium tuberculosis* phenotypic heterogeneity and induce nongrowing metabolically active forms. Cell Host Microbe 17:32-46.

11.  Perry W, Spraggins J, Sheldon J, Grunenwald C, Heinrichs D, Cassat J, Skaar E, Caprioli
R. 2019. *Staphylococcus aureus* exhibits heterogeneous siderophore production within
the vertebrate host. Proc Natl Acad Sci 116:21980-21982.

12.  Cunrath O, Bumann D. 2019. Host resistance factor SLC11A1 restricts *Salmonella*
growth through magnesium deprivation. Science 366:995-999.

13.  Rosenthal A, Qi Y, Hormoz S, Park J, Li S, Elowitz M. 2018. Metabolic interactions
between dynamic bacterial subpopulations. Elife 7.

14.  Stewart P, White B, Boegli L, Hamerly T, Williamson K, Franklin M, Bothner B, James
G, Fisher S, Vital-Lopez F, Wallqvist A. 2019. Conceptual model of biofilm antibiotic
tolerance that integrates phenomena of diffusion, metabolism, gene expression, and
physiology. J Bacteriol 201:e00307-19.

15.  Burton N, Schürmann N, Casse O, Steeb A, Claudi B, Zankl J, Schmidt A, Bumann D.
2014. Disparate impact of oxidative host defenses determines the fate of *Salmonella*
during systemic infection of mice. Cell Host Microbe 15:72-83.

16.  Patel P, O'Hara B, Aunins E, Davis K. 2021. Modifying TIMER to generate a slow-
folding DsRed derivative for optimal use in quickly-dividing bacteria. PLoS Pathog 17.

17.  Davis K, Mohammadi S, Isberg R. 2015. Community behavior and spatial regulation
within a bacterial microcolony in deep tissue sites serves to protect against host attack.
Cell Host Microbe 17:21-31.

18.  Rowe S, Wagner N, Li L, Beam J, Wilkinson A, Radlinski L, Zhang Q, Miao E, Conlon
B. 2020. Reactive oxygen species induce antibiotic tolerance during systemic
*Staphylococcus aureus* infection. Nat Microbiol 5:282-290.

19.  Brauner A, Fridman O, Gefen O, Balaban N. 2016. Distinguishing between resistance,
tolerance and persistence to antibiotic treatment. Nat Rev Microbiol 14:320-30.

20.  Fisher R, Gollan B, Helaine S. 2017. Persistent bacterial infections and persister cells.
Nat Rev Microbiol 15:453-64.

21.  Stapels D, Hill P, Westermann A, Fisher R, Thurston T, Saliba A, Blommestein I, Vogel
J, Helaine S. 2018. *Salmonella* persisters undermine host immune defenses during
antibiotic treatment. Science 362:1156-1160.

22.    Raneses J, Ellison A, Liu B, Davis K. 2020. Subpopulations of stressed *Y.*
695           *pseudotuberculosis* preferentially survive doxycycline treatment within host tissues.
696           mBio 11:e00901-20.

697    23.    Henry T, Brynildsen M. 2016. Development of persister-FACSeq: a method to massively
698           parallelize quantification of persister physiology and its heterogeneity. Sci Rep 6.

699    24.    Luk C, Valenzuela C, Gil M, Swistak L, Bomme P, Chang Y, Mallet A, Enninga J. 2021.
700           *Salmonella* enters a dormant state within human epithelial cells for persistent infection.
701           PLoS Pathog 17.

702    25.    Schulte M, Olschewski K, Hensel M. 2021. Fluorescent protein-based reporters reveal
703           stress response of intracellular *Salmonella enterica* at level of single bacterial cells. Cell
704           Microbiol 23.

705    26.    Zalis E, Nuxoll A, Manuse S, Clair G, Radlinski L, Conlon B, Adkins J, Lewis K. 2019.
706           Stochastic variation in expression of the tricarboxylic acid cycle produces persister cells.
707           MBio 10:e01930-19.

708    27.    Helaine S, Thompson J, Watson K, Liu M, Boyle C, Holden D. 2010. Dynamics of
709           intracellular bacterial replication at the single cell level. Proc Natl Acad Sci 107:3746-51.

710    28.    Roostalu J, Jöers A, Luidalepp H, Kaldalu N, Tenson T. 2008. Cell division in
711           *Escherichia coli* cultures monitored at single cell resolution. BMC Microbiol 8:68.

712    29.    Helaine S, Holden D. 2013. Heterogeneity of intracellular replication of bacterial
713           pathogens. Curr Opin Microbiol 16:184-91.

714    30.    Saliba A, Li L, Westermann A, Appenzeller S, Stapels D, Schulte L, Helaine S, Vogel J.
715           2016. Single-cell RNA-seq ties macrophage polarization to growth rate of intracellular
716           *Salmonella*. Nat Microbiol 2.

717    31.    Gonzalez R, Lane M, Wagner N, Weening E, Miller V. 2015. Dissemination of a highly
718           virulent pathogen: tracking the early events that define infection. PLoS Pathog
719           11:e1004587.

720    32.    McVicker G, Prajsnar T, Williams A, Wagner N, Boots M, Renshaw S, Foster S. 2014.
721           Clonal expansion during *Staphylococcus aureus* infection dynamics reveals the effect of
722           antibiotic intervention. PLoS Pathog 10:e1003959.

33.   Pollitt E, Szkuta P, Burns N, Foster S. 2018. *Staphylococcus aureus* infection dynamics.
PLoS Pathog 14.

34.   Capron J, Delamarre J, Delcenserie R, Gineston J, Dupas J, Lorriaux A. 1981. Liver
absess complicating *Yersinia pseudotuberculosis* ileitis. Gastroenterology 81:150-152.

35.   Hubbert W, Petenyi C, Glasgow L, Uyeda C, Creighton S. 1971. *Yersinia*
*pseudotuberculosis* infection in the United States. Septicemia, appendicitis, and
mesenteric lymphadenitis. Am J Trop Med Hyg 20:679-684.

36.   Paff J, Triplett D, Saari T. 1976. Clinical and laboratory aspects of *Yersinia*
*pseudotuberculosis* infections, with a report of two cases. Am J Clin Pathol 66:101-110.

37.   Barnes P, Bergman M, Mecsas J, Isberg R. 2006. *Yersinia pseudotuberculosis*
disseminates directly from a replicating bacterial pool in the intestine. J Exp Med
203:1591-1601.

38.   Peterson L, Philip N, DeLaney A, Wynosky-Dolfi M, Asklof K, Gray F, Choa R, Bjanes
E, Buza E, Hu B, Dillon C, Green D, Berger S, Gough P, Bertin J, Brodsky I. 2017.
RIPK1-dependent apoptosis bypasses pathogen blockage of innate signaling to promote
immune defense. J Exp Med 214:3171-3182.

39.   Zhang Y, Khairallah C, Sheridan B, van der Velden A, Bliska J. 2018. CCR2+
inflammatory monocytes are recruited to *Yersinia pseudotuberculosis* pyogranulomas and
dictate adaptive responses at the expense of innate immunity during oral infection. Infect
Immun 86:e00782-17.

40.   Green E, Clark S, Crimmins G, Mack M, Kumamoto C, Mecsas J. 2016. Fis is essential
for *Yersinia pseudotuberculosis* virulence and protects against reactive oxygen species
produced by phagocytic cells during infection. PLoS Pathog 12:e1005898.

41.   Resch M, Striegl H, Henssler E, Sevvana M, Egerer-Sieber C, Schiltz E, Hillen W,
Muller Y. 2008. A protein functional leap: how a single mutation reverses the function of
the transcription regulator TetR. Nucleic Acids Res 36:4390-401.

42.   Scholz O, Henssler E, Bail J, Schubert P, Bogdanska-Urbaniak J, Sopp S, Reich M,
Wisshak S, Köstner M, Bertram R, Hillen W. 2004. Activity reversal of Tet repressor
caused by single amino acid exchanges. Mol Microbiol 53:777-789.

43.   Almiron M, Link A, Furlong D, Kolter R. 1992. A novel DNA-binding protein with
regulatory and protective roles in starved *Escherichia coli*. Genes Dev 6:2646-54.

44.   Nair S, Finkel S. 2004. Dps protects cells against multiple stresses during stationary
phase. J Bacteriol 186:4192-8.

45.   Karas V, Westerlaken I, Meyer A. 2015. The DNA-Binding Protein from Starved Cells
(Dps) Utilizes Dual Functions To Defend Cells against Multiple Stresses. J Bacteriol
197:3206-15.

46.   Degenkolb J, Takahashi M, Ellestad G, Hillen W. 1991. Structural requirements of
tetracycline-Tet repressor interaction: determination of equilibrium binding constants for
tetracycline analogs with the Tet repressor. Antimicrob Agents Chemother 35:1591-5.

47.   Rasmussen B, Noller H, Daubresse G, Oliva B, Misulovin Z, Rothstein D, Ellestad G,
Gluzman Y, Tally F, Chopra I. 1991. Molecular basis of tetracycline action: identification
of analogs whose primary target is not the bacterial ribosome. Antimicrob Agents
Chemother 35:2306-11.

48.   Jones-Carson J, Yahashiri A, Kim J, Liu L, Fitzsimmons L, Weiss D, Vázquez-Torres A.
2020. Nitric oxide disrupts bacterial cytokinesis by poisoning purine metabolism. Sci
Adv 6.

49.   M B, Damron F. 2016. Rainbow Vectors for Broad-Range Bacterial Fluorescence
Labeling. PLoS One 11.

50.   Shen Y, Chen Y, Wu J, Shaner N, Campbell R. 2017. Engineering of mCherry variants
with long Stokes shift, red-shifted fluorescence, and low cytotoxicity. PLoS One 12.

51.   Davis K, Krupp J, Clark S, Isberg R. 2019. Iron-sulfur cluster repair contributes to
*Yersinia pseudotuberculosis* survival within deep tissues. Infect Immun 87.

52.   Davidson R, Davis K. 2020. *Yersinia pseudotuberculosis*: Cultivation, Storage, and
Methods for Introducing DNA. Curr Protoc Microbiol 59.
777
778
779

 **Figure Legends**

 **Figure 1. The revTetR reporter can be used to visualize differential bacterial growth rates**

 *in vitro* **with mCherry fluorescence dilution.** (**A**) The revTetR-mCherry reporter was created

 by introducing two mutations (E15A and L17G) in the wild-type tetR sequence, which results in

 constitutive mCherry expression in the absence of tetracyclines (-Tet). Tetracycline addition

 (+Tet) promotes revTetR binding and repression of mCherry transcription. (**B**) After tetracycline

 addition (+Dox/ATc), fast-growing/dividing bacteria (top row, red circles) will partition

 mCherry equally into daughter cells with each cell division event, resulting in dilution. Slow-

 growing/non-dividing cells will retain a high level of mCherry (bottom row). (**C-E**)**:** Exponential

 phase cultures of the revTetR-mCherry *dps* strain (2h subculture in LB, 37$^o$ C with rotation) were

 treated with the indicated ATc doses at 0h (hours, h), then further incubated at 37$^o$ C with

 rotation. Aliquots were taken at the indicated timepoints to detect (**C**) optical density by

 absorbance (A$_{600nm}$), (**D**) mCh fluorescence (normalized by dividing total fluorescence by OD

 values, untreated value set to 100%), dotted line depicts one doubling (50%), and (**E**) GFP

 fluorescence normalized to OD. (**C-E**): Data shown from 9 replicates. (**F**) Cells treated with the

 indicated doses of ATc for 4h were fixed for fluorescence microscopy. Sum single cell mCherry

 signal is shown. (**G**) Sum single cell GFP signal is shown. (**H**) Correlation plot depicting single

 cell sum mCherry and sum GFP signals. (**F-H**)**:** Data shown from three biological replicates.

 Mean and standard deviation are shown. Statistics: (**C-E**) Two-way ANOVA with Tukey's

 multiple comparison test, comparisons made between 2µg/ml and other groups; (**F-G**) Kruskal-

 Wallis one-way ANOVA with Dunn's post-test; (**H**) linear regression with R$^2$ value and slope of

 best-fit line, significantly non-zero slope indicates values are correlated. \*\*\*\*p<.0001, \*\*p<.01,

 \*p<.05, n.s.: not-significant.

**Figure 2. Growth inhibition with doxycycline promotes mCherry fluorescence accumulation.** Exponential phase cultures of the revTetR-mCherry *dps* strain were treated with the indicated doses of doxycycline (Dox) at 0h (hours, h), aliquots were taken at the indicated timepoints to detect **(A)** optical density by absorbance ($A_{600nm}$), **(B)** mCh fluorescence (normalized by dividing total fluorescence by OD values, untreated value set to 100%), and **(C)** GFP fluorescence normalized to OD. **(A-C)**: Data shown from 9 replicates. **(D)** Cells treated with the indicated doses of Dox for 4h were fixed for fluorescence microscopy. Sum single cell mCherry signal is shown. **(E)** Correlation plot depicting single cell sum mCherry and sum GFP signals. **(D-E):** Data shown from three biological replicates. Mean and standard deviation are shown. Statistics: **(A-C)** Two-way ANOVA, Tukey's multiple comparison test, comparisons shown between untreated (0µg/ml) and 1µg/ml Dox; **(D)** Kruskal-Wallis one-way ANOVA with Dunn's post-test; **(E)** linear regression with $R^2$ value and slope of best-fit line, significantly non-zero slope indicates values are correlated. \*\*\*\*p<.0001, \*\*\*p<.001, \*\*p<.01, \*p<.05, n.s.: not-significant.

**Figure 3. Nitric oxide-stressed cells divide at a slower rate than unstressed cells.** Exponential phase cultures of the revTetR-mCherry *$P_{hmp}$::gfp* strain were cultured two hours in the presence (+NO) or absence (-NO) of NO, then split, and either treated with 1µg/ml ATc (+ATc) or left untreated (-ATc). Timepoints are shown relative to ATc addition. **(A)** Aliquots were taken at the indicated timepoints to detect optical density by absorbance ($A_{600nm}$). Mean and standard deviation are shown. **(B)** Aliquots of cells were fixed at the indicated timepoints for fluorescence microscopy. Sum single cell mCherry signal is shown. Median value/group is highlighted. **(C-E)**

Correlation plots depicting single cell sum mCherry and mean GFP signals. Comparisons were

made for NO-treated cells (+NO) in the presence (+ATc) and absence (-ATc) of ATc, cells were

imaged at the indicated timepoints. Values were normalized to the max signal intensity detected

in samples, represented by a value of 1.0. All data shown represents three biological replicates.

Number of cells analyzed is shown. Statistics: **(A)** Two-way ANOVA, Tukey's multiple

comparison test, comparisons shown between untreated (-NO) and NO donor treated (+NO)

samples; **(B)** Kruskal-Wallis one-way ANOVA with Dunn's post-test; **(C-E)** linear regression

with $R^2$ value and slope of best-fit line, significantly non-zero slope indicates values are

correlated. ****p<.0001, *p<.05, n.s.: not-significant.

**Figure 4. Low doses of doxycycline and anhydrotetracycline diffuse evenly across**

**microcolonies and do not inhibit bacterial growth.** C57BL/6 mice were infected intravenously

with the TetON $gfp^+$ strain and a single dose of Dox (4mg/kg) or ATc (4 mg/kg) was

administered intraperitoneally at 48h post-inoculation. Spleens were harvested at the indicated

timepoints after treatment. **(A)** Schematic of the TetON reporter: addition of tetracycline relieves

TetR repression and allows for *mCherry* transcription. **(B)** CFU/spleen quantification at the

indicated timepoints post-treatment. Dotted line: limit of detection, dots: individual mice. **(C)**

Reporter signal was detected with fluorescence microscopy and is shown as the ratio of TetON

(mCherry)/GFP signal at the centroid and periphery of microcolonies. Timepoints indicate time

post-treatment with ATc (left panel) or Dox (right panel). Dotted line: equivalent levels of

mCherry and GFP signal. Each dot: individual microcolony. Mean and standard deviation are

shown. **(D)** Representative images. Merged images and individual channels are shown. Cells

outside the dotted line are defined as peripheral. Dataset represents 4-5 mice/group, collected

from two independent experiments; 2-8 microcolonies were analyzed/spleen from a single cross-

section of tissue. Statistics: **(B)** Kruskal-Wallis one-way ANOVA with Dunn's post-test; **(C)**

Wilcoxon matched-pairs; *$p<.05$, n.s.: not-significant.

**Figure 5. revTetR-mCherry signal appears to increase in both the centroid and periphery**

**of microcolonies.** C57BL/6 mice were infected intravenously with the revTetR-mCherry $gfp^+$

strain and a single dose of ATc (4 mg/kg) was administered intraperitoneally at 48h post-

inoculation. Spleens were harvested at the indicated timepoints after treatment to quantify CFUs

and prepare tissue for fluorescence microscopy. **(A)** CFU/spleen quantification at the indicated

timepoints post-treatment. Dotted line: limit of detection, dots: individual mice, horizontal lines:

median values. **(B)** Reporter signal at 4h quantified using the ratio of revTetR (mCherry)/GFP

signal at the centroid and periphery of microcolonies. Dotted line: equivalent levels of mCherry

and GFP signal. Each dot: individual microcolony. Mean and standard deviation are shown. **(C)**

Reporter signal at 24h quantified using ratio of revTetR (mCherry)/GFP signal at the centroid

and periphery of microcolonies. Mean and standard deviation are shown. **(D)** Representative

images. Dataset represents 4-6 mice/group, collected from three independent experiments; 2-13

microcolonies were analyzed/spleen from a single cross-section of tissue. Statistics: **(A)** Kruskal-

Wallis one-way ANOVA with Dunn's post-test; **(B-C)** Wilcoxon matched-pairs; **$p<.01$,

*$p<.05$, n.s.: not-significant.

**Figure 6. NO stress correlates with heightened mCherry signal accumulation and slowed**

**cell division rates.** Mice were infected with the revTetR-mCherry $P_{hmp}::gfp$ strain and a single

dose of ATc (4 mg/kg) was administered intraperitoneally at 48h post-inoculation. Spleens were

harvested at the indicated timepoints after treatment and prepared for fluorescence microscopy.

**(A)** CFU/spleen quantification. Dotted line: limit of detection, dots: individual mice. **(B)**

Reporter signal at 4h quantified using ratio of periphery/centroid signal for either mCherry

(mCh) or GFP. Dotted line: equivalent levels of mCherry and GFP signal. Each dot: individual

microcolony. **(C)** Reporter signal at 24h quantified using ratio of periphery/centroid signal for

either mCh or GFP. **(D-G)** Correlation plots depicting single cell revTetR signal (mCherry) and

Hmp signal (GFP) at the indicated timepoints, comparing +/- ATc treatment. Graphs represent

either centroid or peripheral values, dots: individual microcolonies. **(H)** Representative images

with the green and red channels shown alongside merged images for 24h post-treatment images.

Dataset represents 5-6 mice/group, collected from three independent experiments; 3-8

microcolonies were analyzed/spleen from a single cross-section of tissue. Median values shown

with horizontal lines. Statistics: (A-C) Kruskal-Wallis one-way ANOVA with Dunn's post-test

**(D-G)** linear regression with $R^2$ value and slope of best-fit line, significantly non-zero slope

indicates values are correlated. \*\*p<.01, n.s.: not-significant.

**Figure 7. Dilution of the revTetR-YFP signal coincides with replication of fast-growing**

**cells.** The revTetR-YFP *P~hmp~::mCherry* strain was grown in the presence (+NO) or absence (-

NO) of NO for 30 min. Cultures were then treated with 1µg/ml ATc (+ATc) or left untreated.

Aliquots were taken at timepoints after ATc addition. **(A)** Sum single cell YFP signal determined

by microscopy. Bars: mean. At least 1,500 individual cells were quantified per group. **(B-C)**

Replication rates of unstressed (-NO) cells were calculated by first normalizing to fluorescent

signal at 0h, then values of treated cells (+ATc) were further normalized to the untreated values

(-ATc, set to 100%) to depict signal dilution over time. Linear regressions were used to calculate

the time it takes for 100% of the cells in culture to undergo one full replication (50% dilution of

the signal). This was done for **(B)** revTetR-YFP (yellow tick, 1.75h for full replication cycle) and

**(C)** revTetR-mCherry (red tick, 3.31h). Statistics: **(A)** Kruskal-Wallis one-way ANOVA with

Dunn's post-test; ****p<.0001, **p<.001.

**Figure 8. Cells that lack NO stress rapidly dilute revTetR-YFP signal indicating faster cell**

**division rates.** Mice were infected with the revTetR-YFP $P_{hmp}::mCherry$ strain, then either

injected with 4 mg/kg ATc (+ATc) or left untreated (unt) at 48h post-inoculation. **(A)**

CFU/spleen was quantified for each mouse at the indicated timepoints post-ATc treatment.

Dotted line indicates limit of detection (LOD), and each dot represents one mouse. **(B-C)** The

ratio of periphery/centroid signals were quantified with revTetR (YFP) and Hmp reporter

(mCherry) signals at **(B)** 4h and **(C)** 24h post-ATc treatment, along with the raw YFP signals at

**(D)** 4h and **(E)** 24h post-treatment. **(F)** Correlation plots are shown for periphery/centroid ratios

for revTetR (YFP) and Hmp (mCherry) reporter signals, comparing treated groups at each

timepoint. **(B-F)** Each dot represents a microcolony. Mean values are highlighted. Data represent

4-5 mice per group per timepoint, with 11-71 microcolonies analyzed for each group and

timepoint. Statistics: **(A-E)** Kruskal-Wallis one-way ANOVA with Dunn's post-test; **(F)** linear

regression was used to determine correlation between values, a significant deviation from zero

indicates the values are correlated; ****p<.0001, *p<.05.

**Figure 9. Representative images of spleens during revTetR-YFP $P_{hmp}::mCherry$ infection.**

Representative images are shown from spleens of mice infected with the revTetR-YFP

*P_{hmp}::mCherry* strain, as described and quantified in Figure 8. Merged images are shown

alongside red and yellow single channels images at the indicated timepoints after ATc treatment.