Mathematical Relationships between Spinal Motoneuron Properties

Arnault Caillet[1], Andrew T.M. Phillips[1], Dario Farina[2],*, Luca Modenese[1],*

[1]Department of Civil and Environmental Engineering, Imperial College London, SW7 2AZ, UK
[2]Department of Bioengineering, Imperial College London, SW7 2AZ, UK
Email: arnault.caillet17@imperial.ac.uk

* These authors have equal contributions and share the senior authorship

NUMBER OF PAGES: 43

NUMBER OF FIGURES: 7

NUMBER OF TABLES: 8

NUMBER OF WORDS FOR ABSTRACT: 194

CONFLICT OF INTEREST STATEMENT: The authors declare no competing financial interests.

ACKNOWLEDGEMENTS: Imperial College Skempton Scholarship to A.C. and Imperial College Research Fellowship to L.M.

Abstract

Our understanding of the behaviour of alpha-motoneurons (MNs) in mammals partly relies on our knowledge of the relationships between MN membrane properties, such as MN size, resistance, rheobase, capacitance, time constant, axonal conduction velocity and afterhyperpolarization period. Based on scattered but converging evidence, current experimental studies and review papers qualitatively assumed that some of these MN properties are related. In this systematic meta-analysis, we reprocessed the data from 40 experimental studies in adult cat, rat and mouse MN preparations *in vivo* to empirically derive mathematical relations between experimentally measured MN electrophysiological and anatomical properties. These mathematical relationships provide the first validated framework of MN-specific morphometric and electrophysiological properties that reproduce published experimental mammal data *in vivo*. These relationships moreover support the classic description of a MN as a membrane equivalent electrical circuit and describe for the first time the association between MN size and MN membrane capacitance and time constant. The obtained relationships finally indicate that motor units are recruited in order of increasing MN size, muscle unit size, MN rheobase, unit force recruitment thresholds and tetanic forces, but underlines that MN size and recruitment order may not be related to motor unit type.

Keywords: motor neuron, motoneuron, motor neuron size, mathematical relationships, motor unit, Henneman's size principle

Introduction

Our understanding of the behaviour of alpha-motoneurons (MNs) in mammals partly relies on the
knowledge of the relationships between the MN morphometric and electrophysiological properties
reported in Table 1. Direct measurement of some of these properties has been performed in animal
studies and yielded some significant correlations, such as the MN input resistance being strongly
inversely associated to rheobase current (Krawitz et al., 2001; Button et al., 2008; Krutki et al., 2015;
Huh et al., 2021). Further investigating such associations in mammals is important to improve both our
understanding of MN pool behaviour and the computational models of MNs. Indeed, due to the
current lack of mathematical descriptions of the relationships between MN morphometric and/or
electrophysiological quantities, MN property profiles in computational models are typically built with
scattered data from a relatively small number of experimental studies. Such generic MN profiles
indicate reasonable orders of magnitude but result in a lack of inter-consistency between the MN
properties. This approach yields obvious limitations when the sets of physiological and MN-specific
characteristics must be known, as in numerical models of a MN membrane equivalent electrical circuit
(Izhikevich, 2004; Dong et al., 2011; Negro et al., 2016; Teeter et al., 2018).

To tackle these current limitations in our understanding of the working principles of MNs and in MN
modelling, algebraic relationships must be derived between the main MN properties listed in Table 1.
This could be done by concurrently measuring all properties in single MNs. However, because of
technical difficulties in measuring MN morphometric and electrophysiological properties *in vivo*, it is
currently challenging to experimentally measure a complete and reliable set of MN-specific properties
for a large sample of MNs. Specialised experimental set-ups could measure a maximum of six MN
properties concurrently in animal preparations, and for limited numbers of MNs (Gustafsson & Pinter,
1984a; Gustafsson & Pinter, 1984b).

Our knowledge of MN properties and their associations is therefore currently limited to a series of
datasets that individually provide a crude picture of MN physiological and biophysical features but not
a detailed understanding of the working principles of MNs. For this reason, the statements reported
in review papers on the associations between MN properties rely on converging conclusions from
independent studies and remain speculative (Henneman, 1981; Burke, 1981; Binder et al., 1996;
Powers & Binder, 2001; Kernell, 2006; Heckman & Enoka, 2012). While qualitative associations, such
as between MN size and resistance (Kernell & Zwaagstra, 1981; Burke et al., 1982), have led to
extensions of Henneman's size principle (Henneman, 1957; Wuerker et al., 1965; Henneman et al.,
1965a; Henneman et al., 1965b; Henneman et al., 1974; Henneman, 1981; Henneman, 1985), other
relationships, such as between MN size and membrane capacitance or time constant, are not known.

Therefore, algebraic relationships between MN properties must be extrapolated from existing data,

taking all data into account in a single mathematical framework.

Here, we derived and validated currently unknown mathematical relationships between any pair of

the MN morphometric and electrophysiological cat properties listed in Table 1 by reprocessing and

merging the published data from 18 available experimental studies in adult cat preparations *in vivo*. In

this way, we first demonstrate that the electrophysiological properties reported in Table 1 are all

associated to each other and can be precisely predicted by MN size in cats. We then assess whether

the cat mathematical relationships apply to other mammals by validating them with data from 9 adult

rat and mouse studies *in vivo*. These empirical relationships provide for the first time a mathematical

framework for the association between any pair of spinal MN properties and for the derivation of MN

profiles of MN-specific morphometric and electrophysiological properties that reproduce published

experimental mammal data *in vivo*. Finally, using additional correlations obtained between some MN

and muscle unit (mU) properties from 14 mammal studies, we discuss the empirical relationships

obtained between MN properties in the context of Henneman's size principle.

		67	**Table 1. The motoneuron (MN) and muscle unit (mU) properties investigated in this study with their notations**
	68	**and SI base units.** S_{MN} is the size of the MN. As reproduced in Table 2, the MN size S_{MN} is adequately described
	69	by measures of the MN surface area S_{neuron} and the soma diameter D_{soma}. R and R_m define the MN-specific
	70	electrical resistance properties of the MN and set the value of the MN-specific current threshold I_{th} (Binder et
	71	al., 1996; Powers & Binder, 2001; Heckman & Enoka, 2012). C and C_m (constant among the MN pool) define the
	72	capacitance properties of the MN and contribute to the definition of the MN membrane time constant τ
	73	(Gustafsson & Pinter, 1984b; Zengel et al., 1985). ΔV_{th} is the constant amplitude of the membrane voltage
	74	depolarization threshold relative to resting state required to elicit an action potential. I_{th} is the corresponding
	75	electrical current causing a membrane depolarization of ΔV_{th}. AHP is defined in most studies as the duration
	76	between the action potential onset and the time at which the MN membrane potential meets the resting state
	77	after being hyperpolarized. ACV is the axonal conduction velocity of the elicited action potentials on the MN
	78	membrane. S_{mU} is the size of the mU. As indicated in Table 2, the mU size S_{mU} is adequately described by
	79	measures of (1) the sum of the cross-sectional areas (CSAs) of the fibres composing the mU CSA^{tot}, (2) the mean
	80	fibre CSA CSA^{mean}, (3) the innervation ratio IR, i.e. the number of innervated fibres constituting the mU, and (4)
	81	the mU tetanic force F^{tet}. The muscle force at which a mU starts producing mU force is called mU force
	82	recruitment threshold F^{th}.

	Properties	**Notation**	**Unit**
	Size:	S_{MN}	
	Neuron surface area	S_{neuron}	$[m^2]$
	Soma diameter	D_{soma}	$[m]$
	Resistance	R	$[\Omega]$
	Specific resistance per unit area	R_m	$[\Omega \cdot m^2]$
	Capacitance	C	$[F]$
Motoneuron (MN) properties	Specific capacitance per unit area	C_m	$[F \cdot m^{-2}]$
	Time constant	τ	$[s]$
	Rheobase (current recruitment threshold)	I_{th}	$[A]$
	Voltage threshold	ΔV_{th}	$[V]$
	Afterhyperpolarization period	AHP	$[s]$
	Conduction velocity	ACV	$[m \cdot s^{-1}]$
	Size:	S_{mU}	
	Total fibre cross-sectional area (CSA)	CSA^{tot}	$[m^2]$
Muscle unit (mU) properties	Mean fibre CSA	CSA^{mean}	$[m^2]$
	Innervation ratio	IR	$[\ ]$
	Tetanic force	F^{tet}	$[N]$
	Force recruitment threshold	F^{th}	$[N]$

Methods

Eligible studies

To optimize inter-study analysis, this meta-analysis focuses exclusively on studies, named "eligible" in the following, that concurrently measured at least two of the morphometric and/or electrophysiological properties reported in Table 1 in individual spinal alpha-motoneurons of healthy adult cats, rats and mice observed *in vivo*. Larval and postnatal specimens were therefore disregarded because of the pronounced age-related variance (Highlander et al., 2020) in the mean values of the electrophysiological properties listed in Table 1, which corroborates the non-extrapolability of neonatal neuronal circuitry to older ages (Song et al., 2006; Carp et al., 2008; Nakanishi & Whelan, 2010; Mitra & Brownstone, 2012). Due to the nonlinear inter-species scalability of the spinal alpha-MN electrophysiological and morphometric properties (Manuel et al., 2019) and a pronounced variance in the inter-species mean values (Highlander et al., 2020), non-mammalian species were disregarded, while the retained data from cats, rats and mice were processed separately. Highlander et al. (2020) also report an 'unexplained' variance in the mean property values reported between studies investigating specimens of the same species, sex and age, which may be explained by differences between *in vivo* and *in vitro* protocols (Carp et al., 2008). For this reason, only *in vivo* studies were considered in this meta-analysis, while most *in vitro* studies had already been disregarded at this stage as dominantly performed on neonatal specimens due to experimental constraints (Manuel et al., 2009a; Mitra & Brownstone, 2012).

To build an extensive set of eligible studies, the output of the systematic websearch provided in Highlander et al. (2020) was screened and used for a further search by reference lists. For convenience, in the following, the notation $\{A; B\}$ refers to the pair of morphometric or electrophysiological MN properties A and B, with $A, B \in \{S_{MN}; ACV; AHP; R; I_{th}; C; \tau\}$, defined in Table 1. The eligible studies generally provided clouds of data points for pairs $\{A; B\}$ of concurrently measured MN properties through scatter graphs. These plots were digitized using the online tool WebPlotDigitizer (Ankit, 2020). When an eligible study did not provide such processable data – most reported the mean $\pm SD$ property values of the cohort of measured MNs – the corresponding author was contacted to obtain the raw data of the measured MN properties, following approval of data sharing. Upon reception of data, datasets of all possible $\{A; B\}$ pairs were created and included into the metanalysis. As all but two eligible studies focused on lumbosacral MNs, the final set of eligible studies was constrained to MNs innervating hindlimb muscles to improve inter-study consistency.

116 From a preliminary screening of the retrieved studies, it was found that relatively more cat studies
117 were obtained than rat and mouse studies, while the cat studies also investigated more $\{A; B\}$ pairs of
118 morphometric and electrophysiological properties. The mathematical relationships sought in this
119 paper between MN properties were thus derived from cat data, and the validated for extrapolation to
120 rat and mouse data.

Relationships between MN properties

<u>Normalized space</u>

The sets of data retrieved from different cat studies for each property pair $\{A; B\}$ were merged into a
'global' dataset dedicated to that property pair. The $\{A; B\}$ data was also normalized for each study
and transformed as a percentage of the maximum property value measured in the same study and
normalized 'global' datasets were similarly created. A least squares linear regression analysis was
performed for the $\ln(A) - \ln(B)$ transformation of each global dataset yielding relationships of the
type $\ln(A) = a \cdot \ln(B) + k$, which were then converted into power relationships of the type $A = k \cdot$
B^a (eq.1). Power fitting was chosen for flexibility and simplicity. The adequacy of these global power
trendlines and the statistical significance of the correlations were assessed with the coefficient of
determination r^2 (squared value of the Pearson's correlation coefficient) and a threshold of 0.01 on
the p-value of the regression analysis, respectively. For each $\{A; B\}$ pair, the normalized global
datasets systematically returned both a higher r^2 and a lower p-value than the datasets of absolute
$\{A; B\}$ values, in agreement with the 'unexplained' inter-experimental variance reported in
(Highlander et al., 2020). It was therefore decided to investigate the $\{A; B\}$ pairs of MN properties in
the normalized space using the normalized global datasets to improve the cross-study analysis.

<u>Size-dependent normalized relationships</u>

From inspection of the considered cat studies, most of the investigated MN property pairs comprised
either a direct measurement of MN size, noted as S_{MN} in this study, or another variable strongly
associated to size, such as axonal conduction velocity (ACV) or afterhyperpolarization period (AHP).
Accordingly, and consistently with Henneman's size principle, we identified S_{MN} as the reference MN
property with respect to which relationships with the electrophysiological MN properties in Table 1
were investigated.

To integrate the data from all global normalized datasets into the final relationships, the MN properties
in Table 1 were processed in a step-by-step manner in the order $ACV, AHP, R, I_{th}, C, \tau$, as reproduced
in Figure 1 for two arbitrary properties, to seek a 'final' power relationship of the type eq.1 between
each of them and S_{MN}. For each electrophysiological property A and each $\{A; B\}$ dataset, we

considered two cases. If $B = S_{MN}$, the global dataset was not further processed as the

electrophysiological property A was already related to MN size S_{MN} from experimental

measurements. If $B \neq S_{MN}$, the global $\{A; B\}$ dataset was transformed into a new $\{A; S_{MN}\}$ dataset by

converting the discrete values of B with the trendline regression $S_{MN} = k_d \cdot B^d$, which was obtained

at a previous step of the data processing. With this dual approach, as many MN size-dependent

$\{A; S_{MN}\}$ intermediary datasets as available $\{A; B\}$ global datasets were obtained for each property A.

These size-dependent datasets were merged into a 'final' $\{A; S_{MN}\}$ dataset to which a least squares

linear regression analysis was performed for the $\ln(A) - \ln(S_{MN})$ transformation, yielding

relationships of the type $\ln(A) = c \cdot \ln(S_{MN}) + k$, which were converted into the power relationships:

$$A = k_c \cdot S_{MN}^c$$

The adequacy of these final power trendlines and the statistical significance of the correlations were

assessed identically to the $\{A; B\}$ relationships derived directly from normalized experimental data,

using the coefficient of determination r^2 (squared value of the Pearson's correlation coefficient) and

a threshold of 0.01 on the p-value of the regression analysis, respectively. If $r^2 > 0.3$ (i.e. $r > 0.55$)

and $p < 0.01$, another power trendline $S_{MN} = k_d \cdot A^d$ was fitted to the final dataset to describe the

inverse relationship for $\{S_{MN}; A\}$ and used in the processing of the next-in-line property.

Figure 1. Two steps of the step-by-step approach to create the final MN size-related datasets. Here, the final
$\{A; S_{MN}\}$ and $\{C; S_{MN}\}$ MN size-related datasets are created (right-side solid contour boxes), to which the final
relationships $A, C = k_c \cdot S_{MN}^c$ were fitted. If $r^2 > 0.3$ and $p < 0.01$, power relationships $S_{MN} = f_A(A)$ and
$S_{MN} = f_C(C)$ (oval boxes) were also fitted to the final $\{A; S_{MN}\}$ and $\{C; S_{MN}\}$ datasets. Here, $S_{MN} = f_B(B)$ was
obtained from a previous unshown step. The normalized global datasets of experimental data (dashed-contour
grey-filled boxes) were either directly included into the final datasets if already obtained as MN size-related, or
first transformed into an MN size-related intermediary datasets (dotted-contour boxes) by converting the A
and/or B properties into S_{MN} values with the $S_{MN} = f_A(A)$ and $S_{MN} = f_B(B)$ relationships. The final $\{A; S_{MN}\}$
and $\{C; S_{MN}\}$ MN size-related datasets were retained for further processing in this study.

 <u>Normalized mathematical relationships between electrophysiological properties</u>

175 When a final relationship with S_{MN} was obtained for two MN properties A and B by the procedure

176 described above, i.e., $A = k_{cA} \cdot S_{MN}^{c_A}$ and $B = k_{cB} \cdot S_{MN}^{c_B}$, one of the expressions was mathematically

177 inverted and a third empirical relationship was derived for the property pair $\{A; B\}$:

178
$$A = \frac{k_{cA}}{(k_{cB})^{\frac{c_A}{c_B}}} \cdot (B)^{\frac{c_A}{c_B}} = k_e \cdot B^e$$

This procedure was applied to all possible $\{A; B\}$ pairs in Table 1.

<u>Validation of the normalized relationships</u>

The data in each $\{A; B\}$ global normalized dataset was initially randomized and therefore made

independent from the studies constituting it. Each global dataset was then randomly split so that 70%

of the data was processed to build the relationships, while the remaining 30% was used as a test

dataset. The final size-related relationships $A = k_c \cdot S_{MN}^c$ and relationships between

electrophysiological properties $A = k_e \cdot B^e$ were obtained as described previously using the 70%-

datasets. These mathematical relationships were applied to the S_{MN} or B data in each 30%-dataset,

yielding predicted values A. It was then assessed to what extent the mathematical relationships

predicted the test data A by calculating the normalized maximum error (nME), the normalized root

mean squared error ($nRMSE$) and the coefficient of determination r_{pred}^2 between predicted and

control values of A in each 30%-dataset. This process was repeated for a total of 5 times, randomly

creating novel 70-30 partitions of the global datasets. The nME, $nRMSE$ and r_{pred}^2 values were finally

averaged across the five repetitions and the average r_{pred}^2 hence obtained was compared to the

average of the r_{exp}^2 values obtained from the power trendlines least-squared fitted to the control data.

This approach validated the relationships. Once validated with this cross-validation method, the final

normalized size-dependent relationships $A = k_c \cdot S_{MN}^c$ and relationships between electrophysiological

properties $A = k_e \cdot B^e$ were provided for the complete global datasets.

<u>Scaling of the normalized relationships</u>

The $A = k_c \cdot S_{MN}^c$ and $A = k_e \cdot B^e$ normalized relationships were finally scaled to typical cat property

values in three steps. First, it was assessed from the literature the fold-range over which each MN

property was to vary. S_{MN} varied over a q_S^E-fold range, taken as the average across cat studies of the

ratios of minimum and maximum values measured for S_{MN}:

202
$$q_S^E = \frac{1}{N_{studies}} \sum_{i=1}^{N_{studies}} \left[\frac{(S_{MN_i})_{max}}{(S_{MN_i})_{min}} \right]$$

An experimental ratio q_A^E was similarly obtained for each electrophysiological property A. As $A = k_c \cdot$

S_{MN}^c, any electrophysiological property A could theoretically vary over a $q_A^T = (q_S^E)^{|c|}$-fold range,

which was compared to q_A^E for consistency in the results. Then, a theoretical range $[A_{min}; A_{max}]$ of

values were derived for each electrophysiological property A. Defining A_{min}^E and A_{max}^E as the

respective average of the minimum and maximum A-values across studies, it was enforced $\frac{A_{max}}{A_{min}} = q_A^T$

and $\frac{A_{min}+A_{max}}{2} = \frac{A_{min}^E + A_{max}^E}{2}$ so that the $[A_{min}; A_{max}]$ theoretical ranges were consistent with the

normalized size-dependent relationships while best reproducing the experimental data from cat

literature. A theoretical range for S_{MN} $[(S_{MN})_{min}; (S_{MN})_{max}]$ was similarly built over the q_S^E-fold

range previously derived. Finally, the intercept k_c in the size-dependent relationships $A = k_c \cdot S_{MN}^c$

was scaled as:

213
$$
\begin{cases}
k_c = \dfrac{A_{min}}{(S_{MN})_{min}^c} \;\; ; \;\; c > 0 \\
k_c = \dfrac{A_{max}}{(S_{MN})_{min}^c} \;\; ; \;\; c < 0
\end{cases}
$$

A similar approach was used to mathematically derive the intercept k_e and scale the relationships $A =$

$k_e \cdot B^e$ between electrophysiological properties.

<u>Extrapolation to rats and mice</u>

It was finally assessed whether the $A = k_c \cdot S_{MN}^c$ and $A = k_e \cdot B^e$ scaled relationships derived from cat

data accurately predicted rat and mouse quantities reported in other experimental studies. These

mathematical relationships were applied to the S_{MN} or B data in each $\{A; B\}$ global dataset of

absolute values obtained for rats and mice independently. The accuracy in predicting the quantity A

was assessed for each available dataset with the same three validation metrics used to validate the

normalized power trend lines: the normalized maximum error (nME), the normalized root-mean-

square error ($nRMSE$) and the coefficient of determination r^2 between predicted and experimental

values of A.

Definitions of MN Size S_{MN} and mU size S_{mU}

<u>MN Size S_{MN}</u>

The eligible papers that performed both electrophysiological and morphometric measurements on the

same MNs dominantly measured the MN soma diameter D_{soma} as an index of MN size. Therefore, we

choose $S_{MN} = D_{soma}$ in the described methodology. In the literature on spinal MNs in adult mammals

in vivo, the size of a MN S_{MN} is also related to the measures of the somal cross-sectional area CSA_{soma}

(Gao et al., 2009; Friese et al., 2009; Deardorff et al., 2013; Mierzejewska-Krzyżowska et al., 2014;

Dukkipati et al., 2018), soma surface area S_{soma} (Ulfhake, B. & Kellerth, 1984; Brandenburg et al., 2020), axonal diameter D_{axon} (Cullheim, 1978), individual dendrite diameter $D_{dendrite}$ (Amendola & Durand, 2008; Carrascal et al., 2009; Mantilla et al., 2018), individual dendritic surface area $S_{dendrite}$ (Li et al., 2005; Obregón et al., 2009; Carrascal et al., 2009; Filipchuk & Durand, 2012; Kanjhan et al., 2015), total dendritic surface area (Ulfhake, Brun & Cullheim, 1988; Amendola & Durand, 2008; Brandenburg et al., 2020) or total MN surface area S_{neuron} (Burke et al., 1982) defined as the summed soma and dendritic surface areas.

In the references (Zwaagstra & Kernell, 1981; Ulfhake & Kellerth, 1981), a linear correlation between D_{soma} and the average diameter of the stem dendrites $D_{dendrite_{mean}}$ has been reported ($r > 0.62$, population size in $\{40; 82\}$ cells). Moreover, a linear or quasi-linear correlation has been found (Cullheim et al., 1987; Ulfhake, Brun & Cullheim, 1988; Moschovakis et al., 1991; Prakash et al., 2000; Li et al., 2005; Obregón et al., 2009; Mantilla et al., 2018) between the stem dendrite diameter $D_{dendrite}$ and the membrane surface area of the corresponding dendritic tree $S_{dendrite}$ ($r > 0.78$, population size in $[33; 342]$ dendrites). Therefore, from these studies, we can assume an approximate linear correlation between D_{soma} and the average dendritic surface area $S_{dendrite_{mean}}$. Moreover, the number of dendritic trees $N_{dendrites}$ per cell increases with increasing soma surface S_{soma} (Brandenburg et al., 2018), and a linear correlation between D_{soma} or $S_{dendrites}$ with $N_{dendrites}$ has been observed (Zwaagstra & Kernell, 1981; Ulfhake, Brun & Cullheim, 1988; Amendola & Durand, 2008) ($r > 0.40$, population size in $\{14; 32; 87\}$ cells). It was therefore assumed that D_{soma} and total dendritic surface area $S_{dendrite}^{tot}$ are linearly correlated. This assumption/approximation is consistent other conclusions of a linear correlations between D_{soma} and $S_{dendrite}^{tot}$ (Amendola & Durand, 2008; Brandenburg et al., 2020). Then, according to typical values of D_{soma}, S_{soma} and $S_{dendrites}^{tot}$ obtained from the studies previously cited, yielding $S_{dendrites}^{tot} \gg S_{soma}$, it was also assumed $S_{dendrites}^{tot} \approx S_{neuron}$. It is thus concluded that D_{soma} is linearly related to total neuron membrane area S_{neuron}, consistently with results by Burke et al. (1982) ($r = 0.61$, 57 cells).

For the above reasons and assumptions, the mathematical relationships $A = k_c \cdot S_{MN}^c$ derived previously, with $S_{MN} = D_{soma}$, were extrapolated with a gain to relationships between A and $S_{MN} = S_{neuron}$, notably permitting the definition of surface specific resistance R_m and capacitance C_m. Following the same method as described above, theoretical ranges $[(S_{neuron})_{min}; (S_{neuron})_{max}]$ were derived from additional morphometric studies on adult cat spinal alpha-motoneurons. The new relationships were extrapolated as $A = k_c \cdot D_{soma}^c = \frac{(S_{neuron})_{min}}{(D_{neuron})_{min}} \cdot S_{neuron}^c$.

It must however be highlighted that the conclusion of a linear correlation between D_{soma} and S_{neuron}, while plausible, is crude. Morphometric measurements of individual MNs are indeed difficult and

suffer many limitations. MN staining, slice preparations and MN reconstruction and identification are complex experimental procedures requiring a large amount of work; the results from most of the cited studies thus rely on relatively small pools of investigated MNs. Most cited studies did not account for tissue shrinkage after dehydration, while some may have failed to assess the full dendritic trees from MN staining techniques (Brandenburg et al., 2020), thus underestimating dendritic membrane measurements. Moreover, before automated tools for image segmentation, landmark mapping and surface tracking (Amendola & Durand, 2008; Obregón et al., 2009; Mierzejewska-Krzyżowska et al., 2014) were available, morphometric measurements were performed from the manual reproduction of the cell outline under microscope, yielding important operator errors reported to be of the order of $\sim 0.5 \mu m$, i.e. $\sim 20\%$ stem diameter. In these studies, morphologic quantities were besides obtained from geometrical approximations of the 2D MN shape, such as modelling the individual dendritic branches as one (Cullheim et al., 1987) or more (Prakash et al., 2000) equivalent cylinders for the derivation of $S_{dendrite}$. Morphometric measurement approaches also varied between studies; for example, oval or circle shapes were best fitted by sight onto the soma outline and equivalent D_{soma} quantities were derived either as the mean of measured maximum and minimum oval diameters or as the equivalent diameter of the fitted circle. In these studies, CSA_{soma} and S_{soma} were therefore derived by classical equations of circle and spheric surface areas, which contradicts the results by Mierzejewska-Krzyżowska et al. (2014) obtained from surface segmentation of a linear relationship between D_{soma} and CSA_{soma} ($r = 0.94$, 527 MNs). Finally, no correlation between soma size and $N_{dendrites}$ was found (Ulfhake & Kellerth, 1981; Cullheim et al., 1987). Therefore, the linear correlation between D_{soma} and S_{neuron} is plausible but crude and requires awareness of several important experimental limitations and inaccuracies. Conclusions and predictions involving S_{neuron} should be treated with care in this study as these morphometric measurements lack the precision of the measures performed for the electrophysiological properties listed in Table 1.

<u>Muscle unit size S_{mU}</u>

To enable future comparisons between MN and mU properties, we here assess potential indices of the mU size S_{mU} suggested in the literature. The size of a mU (S_{mU}) can be defined as the sum CSA^{tot} of the cross-sectional areas CSAs of the innervated fibres composing the mU. CSA^{tot} depends on the mU innervation ratio (IR) and on the mean CSA (CSA^{mean}) of the innervated fibres: $S_{mU} = CSA^{tot} = IR \cdot CSA^{mean}$. CSA^{tot} was measured in a few studies on cat and rat muscles, either indirectly by histochemical fibre profiling (Burke & Tsairis, 1973; Dum & Kennedy, 1980; Burke, 1981), or directly by glycogen depletion, periodic acid Schiff (PAS) staining and fibre counting (Burke et al., 1982; Rafuse et al., 1997). The mU tetanic force F^{tet} is however more commonly measured in animals. As the fibre mean specific force σ is considered constant among the mUs of one muscle in animals (Lucas et al.,

1987; Enoka, 1995), the popular equation $F^{tet} = \sigma \cdot IR \cdot CSA^{mean}$ (Burke, 1981; Enoka, 1995)

returns a linear correlation between F^{tet} and $IR \cdot CSA^{mean} = S_{mU}$ in mammals. Experimental results

(Burke & Tsairis, 1973; Bodine et al., 1987; Chamberlain & Lewis, 1989; Kanda & Hashizume, 1992;

Rafuse et al., 1997; Hegedus et al., 2007) further show a linear correlation between F^{tet} and IR and

between F^{tet} and CSA^{mean}. Consequently, F^{tet}, IR and CSA^{mean} are measurable, consistent, and

linearly related measures of S_{mU} in animals, as summarized in Table 2.

Table 2. Measurable indices of MN and mU sizes in mammals. S_{MN} and S_{mU} are conceptual parameters which
are adequately described by the measurable and linearly inter-related quantities reported in this table.

MN size (S_{MN})	mU size (S_{mU})
S_{neuron}	F^{tet}
D_{soma}	IR
	CSA^{mean}
	CSA^{tot}

Relationships between MN and mU properties

To assess whether the size-dependent relationships $A = k_c \cdot S_{MN}^c$ derived in this study were in

accordance with Henneman's size principle of motor unit recruitment, we identified a set of

experimental studies that concurrently measured a MN property B_{MN} (Table 1) and a muscle unit (mU)

property A_{mU} for the same MU. The normalized global datasets obtained for the pairs $\{A_{mU}; B_{MN}\}$

were fitted with power trendlines, as previously described for MN properties, yielding $A_{mU} = k \cdot B_{MN}^b$

relationships. Using both the definition of S_{mU} (Table 2) and the $A = k_c \cdot S_{MN}^c$ relationships derived

previously, the $A_{mU} = k \cdot B_{MN}^b$ relationships were then mathematically transformed into $S_{mU} = k \cdot$

S_{MN}^c relationships. If all c-values obtained from different $\{A_{mU}; B_{MN}\}$ pairs were of the same sign, it

was concluded that mU and MN sizes were monotonically related. Considering the limited data

available obtained from the literature, data obtained for MNs innervating different hindlimb muscles

were merged. Also, cat, rat and mouse studies were processed independently but the resulting c-

values were compared without regards to the related species.

Results

We respectively identified 18, 6 and 4 eligible studies on cats, rats and mice that reported processable experimental data for the morphometric and electrophysiological properties listed in Table 1. Additional eligible studies including some from the past 10 years were identified but could not be included in the meta-analysis as no processable data could be recovered. An additional 14 studies were found to perform concurrent MN and mU measurements on individual motor units.

From the cat eligible studies, the 17 pairs of MN properties and the 8 pairs of one MN and one mU property represented in the bubble diagram of Figure 2(A) were investigated.

Figure 2. Experimental (A) and unknown (B) relations between MN and mU properties. (A) Bubble diagram representing the pairs of MN and/or mU properties that could be investigated in this study from the results provided by the 40 eligible studies. MN and mU properties are represented in circle and square bubbles, respectively. Relationships between MN properties are represented by coloured connecting lines; the colours red, blue, green, yellow, and purple are consistent with the order $ACV, AHP, R, I_{th}, C, \tau$ in which the pairs were investigated (see Table 3 for mathematical relationships). Relationships between one MN and one mU property are represented by black dashed lines. (B) Bubble diagram representing the mathematical relationships proposed in this study between pairs of MN properties for which no concurrent experimental data has been measured to date.

Relationships between MN properties

<u>Cat eligible studies</u>

All 18 eligible studies on cats included in this meta-analysis were published between 1966 and 2001. They applied similar experimental protocols. Briefly, animals were anesthetized, immobilized, and kept under artificial breathing. Hindlimb muscles of interest were dissected free and their nerves were mounted onto stimulating electrodes, while maintaining body temperature between 35°C and 38°C. After a laminectomy was performed over the lumbosacral region of the spinal cord, the MNs were

identified by antidromic invasion following electrical stimulation of the corresponding muscle nerves.

All studies reported the use of electrodes having stable resistances and being able to pass currents up

to 10 nA without evident polarization or rectification. Most studies performed intracellular recordings

using similar experimental approaches for the measurements of the electrophysiological parameters.

For instance, all studies measured the MN membrane input resistance using the spike height method

(Frank & Fuortes, 1956) by measuring the effects of steady currents, injected through the intracellular

microelectrode, on the amplitude of antidromic spikes. Some aspects of the experimental protocol

however diverged between studies, such as the age and the sex of the group of adult cats, the size

population of cats and recorded MNs, the muscles innervated by the recorded MNs, the means of

anaesthesia, the level of oxygen, and whether the spinal dorsal roots were severed (Kernell, 1966;

Burke, 1968; Barrett & Crill, 1974; Fleshman et al., 1981; Gustafsson & Pinter, 1984a; Gustafsson &

Pinter, 1984b; Zengel et al., 1985; Foehring et al., 1987), the complete surrounding hindlimbs were

denervated (Burke, 1968; Kernell & Zwaagstra, 1981) or the spinal cord was transected (Burke, 1968;

Krawitz et al., 2001). All the eligible studies performing morphometric measurements injected the

recorded MNs intracellularly with horseradish peroxidase (HRP) tracer by applying a continuous train

of anodal current pulses. The spinal cord was eventually sliced frozen or at room temperature with a

microtome or a vibratome and the MN morphometry was investigated. All morphometric

measurements were manually performed, and the MN compartments (soma, dendrites, axon)

approached by simple geometrical shapes, yielding the limitations previously listed.

<u>Normalized relationships</u>

The experimental data retrieved for the 17 pairs of MN properties drawn in Figure 2(A) were

successfully digitized and merged into 17 normalized global datasets as reproduced in Figure 3. For

information, power trendlines were fitted to these normalized datasets and reported in Table 3. All 17

trendlines were statistically significant, showing inter-study consistency in normalized measurements,

and adequately represented by a power model $A = k_a \cdot B^a$ (p-value$< 10^{-5}$ and $r^2 \in [0.34; 0.72]$ for

15 datasets; p-value $< 10^{-5}$ and $r^2 \in \{0.24; 0.17\}$ for the $\{C; AHP\}$ and $\{C; CV\}$ datasets

respectively).

Figure 3. Normalized global datasets. These are obtained from the 18 eligible studies on cats that measured and investigated the 17 pairs of MN properties reported in Figure 2(A). For each $\{A; B\}$ pair, the property A is read on the y axis and B on the x axis. For information, power trendlines $A = k \cdot B^a$ (red dotted curves) are fitted to the data of each dataset and reported in Table 3. The studies are identified with the following symbols: • (Gustafsson, B., 1979; Gustafsson & Pinter, 1984a; Gustafsson & Pinter, 1984b), ○ (Munson et al., 1986), ▲ (Zengel et al., 1985), Δ (Foehring et al., 1987), ■ (Cullheim, 1978), □ (Burke, 1968; Burke & Ten Bruggencate, 1971; Burke et al., 1982), ◆ (Krawitz et al., 2001), ◇ (Fleshman et al., 1981), + (Kernell, 1966), X (Kernell & Zwaagstra, 1981; Kernell & Monster, 1981), - (Zwaagstra, B. & Kernell, 1980), − (Sasaki, 1991), ✻ (Pinter & Vanden Noven, 1989). The axes are given in % of the maximum retrieved values in the studies consistently with the Methods section.

The 17 normalized global datasets were processed according to the Methods to derive normalized

mathematical relationships between the electrophysiological properties listed in Table 1 and MN size

S_{MN}. First, the normalized size-dependent relationship $ACV = 4.0 \cdot S_{MN}^{0.7}$, reported in Table 3, was

derived from the trendline fitting of the normalized $\{ACV; S_{MN}\}$ dataset, which was obtained from 3

studies (Cullheim, 1978; Kernell & Zwaagstra, 1981; Burke et al., 1982) and is represented in the upper-

left panel in Figure 3. This resulted in a statistically significant relationship ($r^2 = 0.58$, p-value$< 10^{-5}$).

A statistically significant ($r^2 = 0.59$, p-value$< 10^{-5}$) inverse relationship $S_{MN} = k_d \cdot ACV^d$ was also

fitted to this dataset. Then, the $\{AHP; ACV\}$ dataset, represented in the second panel upper row in

Figure 3 and obtained from 3 studies (Zwaagstra, B. & Kernell, 1980; Gustafsson & Pinter, 1984a;

Foehring et al., 1987) was transformed into a new $\{AHP; S_{MN}\}_1$ dataset by applying the priory derived

$S_{MN} = k_d \cdot ACV^d$ relationship to the list of ACV-values. This transformed $\{AHP; S_{MN}\}_1$ dataset was

then merged with the $\{AHP; S_{MN}\}_2$ dataset (third panel upper row in Figure 2) obtained from

(Zwaagstra, B. & Kernell, 1980), yielding the final $\{AHP; S_{MN}\}_f$ dataset of $N = 492$ data points shown

in Figure 4, second panel. A statistically significant ($r^2 = 0.58$, p-value$< 10^{-5}$) power trendline

$AHP = 2.5 \cdot 10^4 \cdot S_{MN}^{-1.5}$ was fitted to the $\{AHP; S_{MN}\}_f$ dataset and is reported in Table 3. As before,

a statistically significant ($r^2 = 0.38$, p-value$< 10^{-5}$) inverse relationship $S_{MN} = k_d \cdot AHP^d$ was also

fitted to this dataset for future use. A similar procedure was applied to derive the normalized final

relationships between $\{R, I_{th}, C, \tau\}$ and S_{MN} reported in the last column of Table 3. A statistically

significant (p-value$< 10^{-5}$) correlation was obtained between each electrophysiological property

$\{ACV, AHP, R, I_{th}, C, \tau\}$ and S_{MN} as reported in Table 3. With $r^2 \in [0.28; 0.58]$, it was obtained that

$\{I_{th}, C, CV\}$ and $\{R, AHP, \tau\}$ respectively increased and decreased with increasing MN sizes S_{MN}, with

slopes reported in Table 3 and visually displayed in Figure 5.

Table 3. Fitted experimental data of pairs of MN properties and subsequent normalized final size-related relationships. For information, the r^2, p-value and the equation
$A = k_a \cdot B^a$ are reported for each fitted global dataset. The normalized MN-size dependent relationships $A = k_c \cdot S_{MN}^c$ are mathematically derived from the transformation
of the global datasets and from the power trendline fitting of the final datasets (N data points) as described in the Methods.

MN property		$A = k_a \cdot B^a$ (normalized global datasets)					$A = k_c \cdot S_{MN}^c$ (final MN-size dependent datasets)				
A	Relationship	k_a	a	r^2	p-value	Reference studies	k_c	c	r^2	p-value	N points
ACV	$ACV = k_a \cdot S_{MN}^a$	4.0	0.7	0.58	$p < 10^{-5}$	(Cullheim, 1978; Kernell & Zwaagstra, 1981; Burke et al., 1982)	4.0	0.7	0.58	$p < 10^{-5}$	109
AHP	$AHP = k_a \cdot S_{MN}^a$	$6.1 \cdot 10^3$	-1.2	0.34	$p < 10^{-5}$	(Zwaagstra, B. & Kernell, 1980)	$2.5 \cdot 10^4$	-1.5	0.41	$p < 10^{-5}$	492
	$AHP = k_a \cdot ACV^a$	$1.5 \cdot 10^4$	-1.4	0.43	$p < 10^{-5}$	(Zwaagstra, B. & Kernell, 1980; Gustafsson & Pinter, 1984a; Foehring et al., 1987)					
R	$R = k_a \cdot S_{MN}^a$	$1.5 \cdot 10^5$	-2.1	0.61	$p < 10^{-5}$	(Kernell & Zwaagstra, 1981; Burke et al., 1982)	$9.6 \cdot 10^5$	-2.4	0.37	$p < 10^{-5}$	745
	$R = k_a \cdot ACV^a$	$6.3 \cdot 10^4$	-2.3	0.40	$p < 10^{-5}$	(Kernell, 1966; Burke, 1968; Barrett & Crill, 1974; Kernell & Zwaagstra, 1981; Fleshman et al., 1981; Gustafsson & Pinter, 1984a; Sasaki, 1991)					
	$R = k_a \cdot AHP^a$	$6.2 \cdot 10^{-1}$	1.1	0.65	$p < 10^{-5}$	(Gustafsson, B., 1979; Gustafsson & Pinter, 1984a; Foehring et al., 1987; Pinter & Vanden Noven, 1989; Sasaki, 1991)					
I_{th}	$I_{th} = k_a \cdot R^a$	$1.1 \cdot 10^3$	-1.0	0.37	$p < 10^{-5}$	(Kernell, 1966; Fleshman et al., 1981; Gustafsson & Pinter, 1984b; Zengel et al., 1985; Munson et al., 1986; Foehring et al., 1987; Krawitz et al., 2001)	$9.0 \cdot 10^{-4}$	2.5	0.37	$p < 10^{-5}$	722
	$I_{th} = k_a \cdot ACV^a$	$3.2 \cdot 10^{-6}$	3.7	0.37	$p < 10^{-5}$	(Kernell & Monster, 1981; Gustafsson & Pinter, 1984b)					
	$I_{th} = k_a \cdot AHP^a$	$2.5 \cdot 10^4$	-1.7	0.60	$p < 10^{-5}$	(Gustafsson & Pinter, 1984b)					
C	$C = k_a \cdot R^a$	$2.4 \cdot 10^2$	-0.4	0.57	$p < 10^{-5}$	(Gustafsson & Pinter, 1984a)	1.2	1.0	0.28	$p < 10^{-5}$	444
	$C = k_a \cdot I_{th}^a$	$2.9 \cdot 10^1$	0.26	0.51	$p < 10^{-5}$	(Gustafsson & Pinter, 1984b)					
	$C = k_a \cdot AHP^a$	$2.8 \cdot 10^2$	-0.4	0.24	$p < 10^{-5}$	(Gustafsson & Pinter, 1984a)					
	$C = k_a \cdot ACV^a$	2.5	0.8	0.17	$p < 10^{-5}$	(Gustafsson & Pinter, 1984a)					
τ	$\tau = k_a \cdot R^a$	8.7	0.5	0.52	$p < 10^{-5}$	(Burke & Ten Bruggencate, 1971; Barrett & Crill, 1974; Gustafsson, B., 1979; Gustafsson & Pinter, 1984a; Zengel et al., 1985; Pinter & Vanden Noven, 1989; Sasaki, 1991)	$2.6 \cdot 10^4$	-1.5	0.46	$p < 10^{-5}$	649
	$\tau = k_a \cdot AHP^a$	2.2	0.8	0.63	$p < 10^{-5}$	(Gustafsson & Pinter, 1984a)					
	$\tau = k_a \cdot I_{th}^a$	$2.3 \cdot 10^2$	-0.45	0.72	$p < 10^{-5}$	(Gustafsson & Pinter, 1984b)					
	$\tau = k_a \cdot ACV^a$	$1.2 \cdot 10^4$	-1.3	0.30	$p < 10^{-5}$	(Gustafsson & Pinter, 1984a)					

Figure 4. Normalized MN size-related final datasets. These are obtained from the 18 eligible studies on cats that concurrently measured at least two of the morphometric and electrophysiological properties listed in Table 1. For each $\{A; S_{MN}\}$ pair, the property A is read on the y axis and S_{MN} on the x axis. The power trendlines $A = k_c \cdot S_{MN}^c$ (red dotted curves) are fitted to each dataset and are reported in Table 3.

Figure 5. Normalized size-dependent behaviour of the MN properties ACV, AHP, R, I_{th}, C and τ. For displaying purposes, the MN properties are plotted in arbitrary units as power functions (intercept $k = 1$) of S_{MN}: $A = S_{MN}^c$ according to Table 3. The larger the MN size, the larger ACV, C and I_{th} in the order of increasing slopes, and the lower AHP, τ and R in the order of increasing slopes.

<u>Normalized mathematical relationships between electrophysiological properties</u>

The final size-dependent relationships reported in Table 3 were mathematically combined to yield normalized relationships between all electrophysiological properties listed in Table 1. As an example, $I_{th} = 9.4 \cdot 10^{-4} \cdot S_{MN}^{2.5}$ and $R = 9.6 \cdot 10^5 \cdot S_{MN}^{-2.4}$ were obtained in Table 3. When mathematically combined (with the latter mathematically inverted as $S_{MN} = 2.9 \cdot 10^2 \cdot R^{-0.4}$), these relationships yielded the normalized relationship between rheobase and input resistance $I_{th} = 1.5 \cdot 10^3 \cdot R^{-1}$. All normalized relationships between MN electrophysiological properties hence obtained are provided in Supplementary Material.

<u>Validation of the normalized relationships</u>

The procedure and normalized relationships reported in Table 3 were obtained from the complete global datasets represented in Figure 3. To provide a validation of this approach and of the normalized relationships, the global datasets were split into two random partitions respectively containing 70% and 30% of the data, after the data points order was randomized within each dataset. Normalized relationships between $\{CV, AHP, R, I_{th}, C, \tau\}$ and S_{MN}, and then between all electrophysiological properties were derived from the 70%-datasets following the same procedure as previously described, and their predictions compared to the ground truth 30%-datasets with calculations of the $nME, nRMSE$ and r_{pred}^2 validation metrics. This validation was performed for five random 70%-30% partitions of the global datasets. Figure 6 reports the average $nME, nRMSE$ and r_{pred}^2 values across the five trials. The coefficient of determination between experimental and predicted quantities was

besides compared to the r_{exp}^2-value of the power trendline best fitted to the experimental data directly, which provides the highest r^2 value possibly reachable by our model. As displayed in Figure 6, the normalized relationships obtained from the 17 70%-datasets predicted the experimental data from the 30%-datasets with average nME between 48% and 412%, 12% to 26% $nRMSE$ and coefficients of determination r_{pred}^2 that diverged from the higher limit r_{exp}^2 by a maximum of 7% (for the $\{R; ACV\}$ pair in Figure 6). These results confirm the validity of the obtained relationships.

Figure 6. Validation of the normalized mathematical relationships obtained from the 70%-datasets against the corresponding 30%-datasets. For each 30%-dataset are reported the average values across the five trials of (A) the normalized maximum error (nME), (B) the normalized root-mean-square error ($nRMSE$), and (C) coefficient of determination (r_{pred}^2, grey bars) which is compared with the coefficient of determination (r_{exp}^2, black bars) of the power trendline fitted to the experimental data directly.

The normalized relationships reported in Supplementary material (and in Table 6 after scaling) were also strongly consistent with other experimental data in cats that were not included in the global datasets displayed in Figure 3 and used for deriving the normalized relationships. The normalized relationships $I^{th} = k \cdot \tau^{-1.7}$, $\frac{1}{C} = k \cdot \tau^{0.6}$, $\frac{1}{S_{MN}} = k \cdot CV^{-1.4}$ and $\frac{1}{R} = k \cdot CV^{3.5}$ yielded, when combined, $\frac{I^{th}}{C} = k \cdot \tau^{-1.1}$ and $\frac{1}{R \cdot S_{MN}} = k \cdot CV^{2.1}$, which are consistent with the relationship $\frac{I^{th}}{C} = k \cdot \tau^{-1}$ experimentally observed by Gustafsson & Pinter (1985) and $\frac{1}{R \cdot S_{MN}} = k \cdot CV^{1.9}$ measured by Kernell & Zwaagstra (1981). The stronger-than-linear inverse relationship $R = k \cdot S_{MN}^{-2.4}$ is also consistent with the phenomenological conclusions by Kernell & Zwaagstra (1981).

<u>Scaling of the normalized relationships</u>

The normalized relationships provided in Supplementary Material were finally scaled to typical cat data obtained from the literature following the procedure described in the Methods section. D_{soma} and S_{neuron} were found to vary in cats over an average $q_S^E = 2.4$-fold range according to a review of the morphometric studies reported in the two first lines of Table 4. The similar ranges obtained for these two morphometric quantities support the findings of the linear inter-relationship existing between them and justified in the Methods.

It must however be noted that the minimum and maximum values of S_{neuron} and D_{soma} reported in Table 4 may underestimate the true distribution of MN sizes in a MN pool for several reasons. First, the studies reported in Table 4 investigated MN populations of small size, so that the largest and smallest MNs of the investigated MN pools may not have been systematically identified and measured. Also, dendritic labelling by staining may only identify dendrites up to the 3rd dendritic branch (Issa et al., 2010; Fogarty et al., 2018) and full dendritic trees may not be identified, thus underestimating S_{neuron} for the largest MNs that exhibit the highest dendritic tree complexity (Fogarty et al., 2020). Finally, the distribution of gamma- and alpha-MN sizes is typically bimodal and the size distributions of the two populations are reported to overlap (Moschovakis et al., 1991; von Steyern et al., 1999; Friese et al., 2009; Deardorff et al., 2013). However, most studies reported in Table 4 identify the alpha MNs from the gamma MNs by visually splitting the bimodal distributions of MN sizes in two independent domains (Donselaar et al., 1986; Ishihara et al., 2001), thus neglecting the overlap. Similarly, some studies identify as gamma MNs any MN showing an axonal conduction velocity typically less than $\sim 60 m \cdot s^{-1}$ (Zwaagstra, B. & Kernell, 1980; Ulfhake, B. & Kellerth, 1984). These approaches lead to overestimating the lower limit for the reported D_{soma} and S_{neuron} values. Consequently, true ranges in MN sizes may be larger than the theoretical ones reported in Table 4. A similar comment can be made for the electrophysiological properties reported in Table 5.

However, the $q_S^E = 2.4$-fold range obtained for cats is consistent with the fold-ranges obtained from rat and mouse data for D_{soma} as reported in Table 4, while it underestimates the 4.4-fold range in S_{neuron} values observed in mice. Table 4 finally shows that typical cat MN sizes are larger than in rats and mice respectively, as reviewed by Manuel et al. (2019).

	Property	Unit	Absolute {min;max}	Average {min; max}	q_S^E	Reference studies	Theoretical range
Cat	D_{soma}	$[\mu m]$	{25.0; 90.0}	{36.7; 75.5}	2.2	(Kernell, 1966; Cullheim, 1978; Zwaagstra, B. & Kernell, 1980; Kernell & Zwaagstra, 1981; Ulfhake & Kellerth, 1981; Zwaagstra & Kernell, 1981; Burke et al., 1982; Donselaar et al., 1986; Destombes et al., 1992)	[33; 79]
	S_{neuron}	$[mm^2]$	{0.08; 0.64}	{0.17; 0.45}	2.7	(Barrett & Crill, 1974; Ulfhake & Kellerth, 1981; Burke et al., 1982; Ulfhake, B. & Kellerth, 1984; Ulfhake, Brun & Cullheim, 1988; Moschovakis et al., 1991)	[0.18; 0.44]
Rat	D_{soma}	$[\mu m]$	{17.5; 66.0}	{23.9; 55.7}	2.4	(Swett et al., 1986; von Steyern et al., 1999; Copray & Kernell, 2000; Ishihara et al., 2001; Deardorff et al., 2013; Mierzejewska-Krzyżowska et al., 2014)	
Mouse	D_{soma}	$[\mu m]$	{14.0; 35.0}	{14.0; 35.0}	2.5	(von Steyern et al., 1999)	
	S_{neuron}	$[mm^2]$	{0.01; 0.08}	{0.01; 0.06}	4.4	(Amendola & Durand, 2008; Brandenburg et al., 2020)	

Then, the empirical q_A^E and theoretical q_A^T ratios, defined in the Methods, were calculated for each MN

electrophysiological property A and are reported in Table 5. For example, MN resistance R was found

to vary over an average $q_R^E = 10.9$-fold range in a MN pool according to the literature, while the

theoretical fold range $q_R^T = (q_S^E)^{|c_R|} = 2.4^{2.4} = 8.4$ was obtained from the $R = 9.6 \cdot 10^5 \cdot S_{MN}^{-2.4}$

normalized relationship previously derived when a 2.4-fold range is set for S_{MN}. As shown in Table 5,

$\frac{q_A^E}{q_A^T} \in [0.8; 1.3]$ for the MN properties C, τ, AHP and ACV, while the theoretical ranges for R and I_{th}

span over a narrower domain than expected from experimental results. The latter either suggests an

underestimation of the MN q_S^E-fold range as discussed previously or of the power c in the relationships

$I_{th}, R = k_c \cdot S_{MN}^c$.

Table 5: Typical ranges of physiological values for the MN properties $R, R_m, C, \tau, I_{th}, AHP$ and ACV. As described in the Methods section, q_A^E is the average among reference studies of the ratios of minimum and maximum values; the properties experimentally vary over a q_A^E-fold range. This ratio compares with the theoretical ratio $q_A^T = (q_S^E)^{|c|}$ (with c taken from Table 3), which sets the amplitude of the theoretical ranges. Absolute and average {min; max} are obtained as described in Table 4.

MN property	Unit	Absolute exp {min;max}	Average exp {min; max}	q_A^E	Reference studies	q_A^T	$\dfrac{q_A^E}{q_A^T}$	Theoretical range
R	$M\Omega$	{0.2; 8.1}	{0.4; 4.0}	10.9	(Kernell, 1966; Burke, 1968; Burke & Ten Bruggencate, 1971; Barrett & Crill, 1974; Gustafsson, B., 1979; Kernell & Zwaagstra, 1981; Fleshman et al., 1981; Burke et al., 1982; Ulfhake, B. & Kellerth, 1984; Gustafsson & Pinter, 1984a; Zengel et al., 1985; Foehring et al., 1986; Munson et al., 1986; Foehring et al., 1987; Sasaki, 1991; Krawitz et al., 2001)	8.4	1.3	[0.5; 4.0]
C	nF	{2.2; 8.5}	{2.2; 8.5}	3.9	(Gustafsson & Pinter, 1984a; Gustafsson & Pinter, 1985)	2.3	1.6	[3.2; 7.5]
τ	ms	{2.0; 14.2}	{2.9; 10.2}	3.5	(Burke & Ten Bruggencate, 1971; Barrett & Crill, 1974; Gustafsson, B., 1979; Ulfhake, B. & Kellerth, 1984; Gustafsson & Pinter, 1984a; Gustafsson & Pinter, 1985; Sasaki, 1991)	3.7	0.9	[2.8; 10.3]
I_{th}	nA	{1.7; 52.7}	{2.3; 36.6}	16.3	(Fleshman et al., 1981; Kernell & Monster, 1981; Ulfhake, B. & Kellerth, 1984; Gustafsson & Pinter, 1984b; Zengel et al., 1985; Foehring et al., 1986; Munson et al., 1986; Foehring et al., 1987; Krawitz et al., 2001)	9.1	1.8	[3.9; 35.0]
AHP	ms	{10.6; 266.6}	{44.2; 158.7}	4.1	(Eccles et al., 1957; Gustafsson, B., 1979; Dum & Kennedy, 1980; Zwaagstra, B. & Kernell, 1980; Ulfhake, B. & Kellerth, 1984; Gustafsson & Pinter, 1984a; Zengel et al., 1985; Foehring et al., 1987; Sasaki, 1991)	3.8	1.1	[42.6; 160.2]
ACV	$m \cdot s^{-1}$	{51.1; 127.2}	{65.3; 114.8}	1.8	(Eccles et al., 1957; McPhedran et al., 1965; Kernell, 1966; Appelberg & Emonet-Dénand, 1967; Burke, 1968; Barrett & Crill, 1974; Proske & Waite, 1974; Bagust, 1974; Stephens & Stuart, 1975; Cullheim, 1978; Dum & Kennedy, 1980; Zwaagstra, B. & Kernell, 1980; Kernell & Zwaagstra, 1981; Fleshman et al., 1981; Glenn & Dement, 1981; Burke et al., 1982; Gustafsson & Pinter, 1984a; Zengel et al., 1985; Foehring et al., 1986; Foehring et al., 1987)	2.3	0.8	[63.5; 116.6]

The normalized relationships were finally scaled using the theoretical ranges reported in the last column of Table 5. Taking the $\{I_{th}; R\}$ pair as example, as $I_{th} = 1.4 \cdot 10^3 \cdot R^{-1}$ (normalized relationship derived previously and provided in Supplementary Material), $R \in [0.5; 4.0] \cdot 10^6 \Omega$ and

$I_{th} \in [3.9; 35.0] \cdot 10^{-9} A$ (Table 5), it was directly obtained from the Methods that $I_{th} = \frac{2.7 \cdot 10^{-2}}{R} =$

$\frac{\Delta V^{th}}{I^{th}}$ in SI base units. A similar approach yielded the mathematical relationships reported in Table 6

between all MN electrophysiological and morphometric properties $ACV, AHP, R, I_{th}, C, \tau$ and S_{MN}. All

constants and relationships are given in SI base units. The specific capacitance C_m and the membrane

voltage threshold ΔV_{th} were found to be constant among MNs, a property discussed in the Discussion

section.

<u>Specific resistance R_m</u>

A specific membrane input resistance is generally defined in the literature (Barrett & Crill, 1974; Burke

et al., 1982; Kernell, 2006) as the surface input resistance $R = \frac{R_m}{S_{neuron}}$. According to the relationships

in Table 6, we obtain $R = \frac{R_m}{S_{neuron}} = \frac{1.7 \cdot 10^{-10}}{S_{neuron}^{2.4}}$, which yields $R_m = \frac{1.7 \cdot 10^{-10}}{S_{neuron}^{1.4}}$. This finding is consistent with

the conclusions from Barrett & Crill (1974), who reported a weak but significant linear relationship

between R_m and $1/S_{MN}$. Also, $R_m = \frac{1.7 \cdot 10^{-10}}{S_{MN}^{1.4}}$ yields $R_m = 4.1 \cdot AHP^{0.9}$ from the relationships in

Table 6, which is consistent with the indirect conclusion of a positive correlation between R_m and AHP

in Gustafsson & Pinter (1984b). Finally, considering that $q_S^E = 2.4$ and $S_{neuron} \in [0.18; 0.44] \, mm^2$

(Table 4), $R_m = \frac{1.7 \cdot 10^{-10}}{S_{MN}^{1.4}}$ yields that R_m varies over the $q_A^E =$3.5-fold theoretical range $[0.21; 0.73] \, \Omega \cdot$

m^2. This is both consistent with the $q_A^E = 3.2$-fold range and the $[A_{min}^E; A_{max}^E] = [0.16; 0.62] \, \Omega \cdot m^2$

mean range previously speculated in the literature (Albuquerque & Thesleff, 1968; Barrett & Crill,

1974; Burke et al., 1982; Gustafsson & Pinter, 1984a; Kernell & Zwaagstra, 1989). The MN property R_m

was therefore added to Table 6.

Table 6. Mathematical empirical cat relationships between the MN properties S_{neuron}, D_{soma}, R, R_m, C, τ, I^{th}, AHP and ACV. Each column provides the relationships between one and the
nine other MN properties. If one property is known, the complete MN profile can be reconstructed by using the pertinent line in this table. All constants and properties are provided in SI base
units (meters, seconds, ohms, farads and ampers). These relationships were obtained for alpha-MNs innervating the hindlimbs of adult cats in vivo.

	$S_{neuron}[m^2]$	$D_{soma}[m]$	$R[\Omega]$	$R_m[\Omega \cdot m^2]$	$C[F]$	$\tau[s]$	$I_{th}[A]$	$AHP[s]$	$ACV[m \cdot s^{-1}]$
$S_{neuron}[m^2]$		$D_{soma} = 1.8 \cdot 10^2 \cdot S_{neuron}$	$R = \frac{1.7 \cdot 10^{-10}}{S_{neuron}^{2.4}}$	$R_m = \frac{1.7 \cdot 10^{-10}}{S_{neuron}^{1.4}}$	$C = 1.0 \cdot 10^{-2} \cdot S_{neuron}$	$\tau = \frac{1.0 \cdot 10^{-12}}{S_{neuron}^{1.5}}$	$I_{th} = 3.8 \cdot 10^8 \cdot S_{neuron}^{2.5}$	$AHP = \frac{1.0 \cdot 10^{-11}}{S_{neuron}^{1.5}}$	$ACV = 3.0 \cdot 10^6 \cdot S_{neuron}^{0.7}$
$D_{soma}[m]$	$S_{neuron} = 5.5 \cdot 10^{-3} \cdot D_{soma}$		$R = \frac{5.1 \cdot 10^{-5}}{D_{soma}^{2.4}}$	$R_m = \frac{2.8 \cdot 10^{-7}}{D_{soma}^{1.4}}$	$C = 6.7 \cdot 10^{-5} \cdot D_{soma}$	$\tau = \frac{2.3 \cdot 10^{-9}}{D_{soma}^{1.5}}$	$I_{th} = 7.8 \cdot 10^2 \cdot D_{soma}^{2.5}$	$AHP = \frac{2.7 \cdot 10^{-8}}{D_{soma}^{1.5}}$	$ACV = 8.1 \cdot 10^4 \cdot D_{soma}^{0.7}$
$R[\Omega]$	$S_{neuron} = \frac{9.5 \cdot 10^{-5}}{R^{0.4}}$	$D_{soma} = \frac{1.7 \cdot 10^{-2}}{R^{0.4}}$		$R_m = 9.5 \cdot 10^{-5} \cdot R^{0.6}$	$C = \frac{1.3 \cdot 10^{-6}}{R^{0.4}}$	$\tau = 9.6 \cdot 10^{-7} \cdot R^{0.6}$	$I_{th} = \frac{2.7 \cdot 10^{-2}}{R}$	$AHP = 1.3 \cdot 10^{-5} \cdot R^{0.6}$	$ACV = \frac{4.9 \cdot 10^3}{R^{0.3}}$
$R_m[\Omega \cdot m^2]$	$S_{neuron} = \frac{1.5 \cdot 10^{-7}}{R_m^{0.7}}$	$D_{soma} = \frac{2.6 \cdot 10^{-5}}{R_m^{0.7}}$	$R = 6.8 \cdot 10^6 \cdot R_m^{1.7}$		$C = \frac{2.6 \cdot 10^{-9}}{R_m^{0.7}}$	$\tau = 1.4 \cdot 10^{-2} \cdot R_m$	$I_{th} = \frac{2.2 \cdot 10^{-9}}{R_m^{1.8}}$	$AHP = 2.2 \cdot 10^{-1} \cdot R_m^{1.1}$	$ACV = \frac{5.4 \cdot 10^1}{R_m^{0.5}}$
$C[F]$	$S_{neuron} = 1.2 \cdot 10^2 \cdot C$	$D_{soma} = 2.1 \cdot 10^4 \cdot C$	$R = \frac{1.5 \cdot 10^{-15}}{C^{2.5}}$	$R_m = \frac{1.8 \cdot 10^{-13}}{C^{1.5}}$		$\tau = \frac{8.6 \cdot 10^{-16}}{C^{1.5}}$	$I_{th} = 6.5 \cdot 10^{13} \cdot C^{2.6}$	$AHP = \frac{7.7 \cdot 10^{-15}}{C^{1.6}}$	$ACV = 8.2 \cdot 10^7 \cdot C^{0.7}$
$\tau[s]$	$S_{neuron} = \frac{8.4 \cdot 10^{-9}}{\tau^{0.7}}$	$D_{soma} = \frac{1.5 \cdot 10^{-6}}{\tau^{0.7}}$	$R = 7.1 \cdot 10^9 \cdot \tau^{1.6}$	$R_m = 6.0 \cdot 10^1 \cdot \tau$	$C = \frac{1.6 \cdot 10^{-10}}{\tau^{0.6}}$		$I_{th} = \frac{1.6 \cdot 10^{-12}}{\tau^{1.7}}$	$AHP = 1.7 \cdot 10^1 \cdot \tau$	$ACV = \frac{7.5}{\tau^{0.5}}$
$I_{th}[A]$	$S_{neuron} = 4.0 \cdot 10^{-4} \cdot I_{th}^{0.4}$	$D_{soma} = 7.1 \cdot 10^{-2} \cdot I_{th}^{0.4}$	$R = \frac{3.1 \cdot 10^{-2}}{I_{th}}$	$R_m = \frac{1.2 \cdot 10^{-5}}{I_{th}^{0.6}}$	$C = 5.3 \cdot 10^{-6} \cdot I_{th}^{0.4}$	$\tau = \frac{1.2 \cdot 10^{-7}}{I_{th}^{0.6}}$		$AHP = \frac{1.5 \cdot 10^{-6}}{I_{th}^{0.6}}$	$ACV = 1.3 \cdot 10^4 \cdot I_{th}^{0.3}$
$AHP[s]$	$S_{neuron} = \frac{5.5 \cdot 10^{-8}}{AHP^{0.7}}$	$D_{soma} = \frac{9.8 \cdot 10^{-6}}{AHP^{0.7}}$	$R = 7.5 \cdot 10^7 \cdot AHP^{1.6}$	$R_m = 4.1 \cdot AHP^{0.9}$	$C = \frac{1.0 * 10^{-9}}{AHP^{0.6}}$	$\tau = 6.2 \cdot 10^{-2} \cdot AHP$	$I_{th} = \frac{1.8 * 10^{-10}}{AHP^{1.7}}$		$ACV = \frac{2.7 \cdot 10^1}{AHP^{0.5}}$
$ACV[m \cdot s^{-1}]$	$S_{neuron} = 4.6 \cdot 10^{-10} \cdot ACV^{1.4}$	$D_{soma} = 8.3 \cdot 10^{-8} \cdot ACV^{1.4}$	$R = \frac{8.3 \cdot 10^{12}}{ACV^{3.5}}$	$R_m = \frac{3.8 \cdot 10^3}{ACV^{2.1}}$	$C = 1.0 \cdot 10^{-11} \cdot ACV^{1.4}$	$\tau = \frac{7.4 \cdot 10^1}{ACV^{2.1}}$	$I_{th} = 1.1 \cdot 10^{-15} \cdot ACV^{3.6}$	$AHP = \frac{1.4 \cdot 10^3}{ACV^{2.2}}$	

 Predicting correlations between MN properties

This study also predicts the correlations between MN properties that were either never reported in
past review studies, such as the positive $S_{MN} - \tau$ relationship, or never concurrently measured in the
literature, as displayed in Figure 2(B). Such unknown relationships were indirectly extracted from the
combination of known relationships (Table 3) and typical ranges of values obtained from the literature
for these properties. Due to the prior validation of the relationships in Table 6, these findings are
reliable as indirectly consistent with the literature data processed in this study and provide new
insights on the size-dependency of the MN recruitment mechanisms.

Extrapolation to rats and mice

It was assessed whether the scaled relationships obtained from cat data and reported in Table 6 could
be successfully applied to rat and mouse data. A global $\{I_{th}; R\}$ dataset was obtained from 5 eligible
rat studies reported in Table 7, while four mice studies reported in Table 7 provided processable data
for the pairs $\{I_{th}; R\}$, $\{\tau; R\}$ and $\{C; R\}$. Figure 7 reports the global datasets for all 4 pairs of properties
and the predictions obtained with the cat relationships reported in Table 6. As displayed in Table 7,
r_{pred}^2 and r_{exp}^2 values were close for all 4 datasets, with a maximum 12.5% difference for $\{\tau; R\}$ in
mice. $nRMSE$ values remained low in both rat and mouse $\{I_{th}; R\}$ datasets, while being substantially
higher in both $\{\tau; R\}$ and $\{C; R\}$ datasets in mice because of an inter-species-specific offset in the
relationships.

Table 7. Validation of the cat relationships (Table 6) against rat and mouse data. nME: normalized maximum
error. nRMSE: normalized root-mean-square error. r_{pred}^2: coefficient of determination between experimental
and predicted quantities. r_{exp}^2: coefficient of determination of the power trendline directly fitted to the
experimental data.

Animal	Dataset	Reference studies	nME (%)	$nRMSE$ (%)	r_{pred}^2	r_{exp}^2
Rat	$\{I_{th}; R\}$	(Gardiner, 1993; Bakels & Kernell, 1993; Lee & Heckman, 1998; Button et al., 2008; Turkin et al., 2010; Krutki et al., 2015)	591	16	0.53	0.54
Mouse	$\{I_{th}; R\}$	(Delestrée et al., 2014; De Lourdes Martínez-Silva et al., 2018; Huh et al., 2021)	621	17	0.37	0.41
	$\{\tau; R\}$	(Manuel et al., 2009b)	1183	155	0.35	0.40
	$\{C; R\}$	(Manuel et al., 2009b)	1802	87	0.38	0.38

Figure 7. Global datasets for rat and mouse species and predictions of the MN properties with the cat mathematical relationships (Table 6). They were obtained from the 5 eligible studies on rats and the 4 eligible studies on mice reported in Table 7 that measured the $\{I_{th}; R\}$, $\{\tau; R\}$ and $\{C; R\}$ pairs of MN electrophysiological properties. I_{th}, R, τ, C are given in $nA, M\Omega, ms$ and nF respectively. The experimental data ($\circ$ symbol) is fitted with a power trendline (red dotted curve) and compared to the predicted quantities obtained with the scaled cat relationships in Table 6 ($\blacksquare$ symbol).

Relationships between MN and mU properties

As shown in Figure 2(A), eight pairs of one MN and one mU property were investigated in 14 studies

in the literature in cats and rats. As remarked by Heckman & Enoka (2012) and Manuel et al. (2019),

most recent studies were performed on mice, while most dated were dominantly performed on cats.

One study on the rat gastrocnemius muscle (Kanda & Hashizume, 1992) indicated no correlation

between IR and CV. However, after removing from the dataset 2 outliers that fell outside two

standard deviations of the mean IR data, a statistically significant correlation ($p < 0.05$) between IR

and CV was successfully fitted with a power trendline ($r^2 = 0.43$). Eight studies, dominantly focusing

on the cat soleus and medial gastrocnemius muscles, found a strong correlation between F^{tet} and CV,

while one study showed a significant correlation for the pair $\{F^{tet}; R\}$ in both the cat tibialis anterior

and extensor digitorum longus muscles. One study (Burke *et al.*, 1982) on the cat soleus, medial and

lateral gastrocnemius muscles inferred a statistically significant correlation for $\{F^{tet}; S_{MN}\}$. In mice,

both R and I_{th} were significantly related to muscle tetanic and twitch forces. As F^{tet}, F^{tw} and IR are

reliable indices of S_{mU} as discussed in the Methods, the MN size-dependent relationships in Table 3

return eight $S_{mU} = k \cdot S_{MN}^c$ relationships between mU and MN properties (Table 8). A 2.8-fold range

in positive c-values $c \in [2.4; 6.6]$ (mean 3.8 ± 1.5 s.d.) was obtained between studies. This result

infers than mU and MN size are positively related in cats, mice and potentially in rats.

Table 8. Fitted experimental data of pairs of one mU and one MN property and subsequent $S_{mU} \propto S_{MN}^c$
relationships. The r^2, p-value and the equation $A \propto B^c$ of the trendlines are reported for each pair of properties.
In the last column, prior knowledge is used to derive $S_{mU} \propto S_{MN}^c$ relationships.

Species	$A = k \cdot B^c$ (fitted relationships)					$S_{mU} = k \cdot S_{MN}^c$ (final relationships)
	Relationship	c	r^2	p-value	Reference studies	c
Rat	$IR = k \cdot CV^c$	3.4	0.45	$6 \cdot 10^{-3}$	(Kanda & Hashizume, 1992)	2.4
	$F^{tw} = k \cdot CV^c$	9.4	0.43	$p < 10^{-5}$	(Knott et al., 1971)	6.6
Cat	$F^{tet} = k \cdot CV^c$	7.2	0.37	$p < 10^{-5}$	(McPhedran et al., 1965; Wuerker et al., 1965; Appelberg & Emonet-Dénand, 1967; Proske & Waite, 1974; Bagust, 1974; Jami & Petit, 1975; Stephens & Stuart, 1975; Burke et al., 1982; Emonet-Dénand et al., 1988)	5.0
	$F^{tet} = k \cdot R^c$	−1.3	0.27	$6 \cdot 10^{-5}$	(Dum & Kennedy, 1980)	3.2
	$F^{tet} = k \cdot S_{MN}^c$	2.0	0.21	$2 \cdot 10^{-2}$	(Burke *et al.*, 1982)	2.0
Mouse	$F^{tw} = k \cdot R^c$	−2.1	0.42	$p < 10^{-5}$	(Manuel & Heckman, 2011; De Lourdes Martínez-Silva et al., 2018)	5.1
	$F^{tw} = k \cdot I_{th}^c$	1.3	0.64	$2 \cdot 10^{-4}$	(Manuel & Heckman, 2011)	3.3
	$F^{tet} = k \cdot I_{th}^c$	1.0	0.80	$6 \cdot 10^{-2}$	(Manuel & Heckman, 2011)	2.5
Mean $\pm$**sd**						3.8 ± 1.5

Discussion

In this systematic meta-analysis, we processed the data (Figure 3) from previous cat experimental studies to extract mathematical relationships between several MN electrophysiological properties and MN size. First, this allowed us to demonstrate that all investigated MN properties are predicted by MN size (Table 3, Figure 4, Figure 5) and that electrophysiological properties at the level of individual MNs are interrelated. Second, we established (Table 6) and validated (Figure 6) mathematical relationships linking all the pairs of MN properties, providing the first mathematical framework for building virtual MN pools of MN-specific morphometric and electrophysiological properties that are consistent with published experimental cat data *in vivo*.

The findings are consistent with considerations from previous literature reviews, either drawn from direct but isolated measurements or speculated. For example, Binder et al. (1996) concluded that the AHP duration was inversely related to MN size S_{MN} based on measures of a single experimental study. Conversely, the size-dependencies of R and I_{th} were qualitatively assumed by Powers & Binder (2001) from the equations obtained from a variant of Rall's model of MN membrane equivalent electrical circuit (Rall, 1957; Rall, 1959; Rall, 1960), and not from experimental data. The negative correlation between S_{MN} and R was predicted by Binder et al. (1996) from the findings of Henneman's studies on the correlation between extracellular spike amplitude and S_{MN} (Henneman, 1957; Henneman et al., 1965a). The fact that R defines the MN rheobase and thus dictates the size order of MN recruitment was similarly justified in Binder et al. (1996) and Powers & Binder (2001) from a combination of Henneman's findings and Rall's model equations. Moreover, all previous papers used the debated association between motor unit type and S_{MN}, mainly obtained from measures in the cat gastrocnemius muscle (Fleshman et al., 1981; Burke et al., 1982; Zengel et al., 1985; Bakels & Kernell, 1993), to indirectly speculate on correlations between S_{MN} and AHP (Heckman & Enoka, 2012), R (Powers & Binder, 2001; Heckman & Enoka, 2012), R_m (Binder et al., 1996; Powers & Binder, 2001) or I_{th} (Binder et al., 1996). Finally, we found a single paper that inferred a positive correlation between S_{MN} and C (Heckman & Enoka, 2012) but without reference to direct experimental results.

Relevance for MN modelling

The empirical equations in Table 6 provide a mathematical framework that supports the common approach of modelling the MN membrane behaviour with an equivalent resistance-capacitance electrical circuit as variants of Rall's cable model (Rall, 1957; Rall, 1959; Rall, 1960). The relationship $C = 1.0 \cdot 10^{-2} \cdot S_{MN}^{1.0}$ reported in Table 6 validates the definition $C = C_m \cdot S_{MN}$, as well as emphasizing that the specific capacitance C_m per unit area is constant among the MN pool, and indirectly yields, from typical ranges of C and S_{MN} in the literature, the relation $C_m = 1.0 \cdot 10^{-2} F \cdot m^{-2}$, which is highly

consistent with the typical values reported in the literature (Lux & Pollen, 1966; Albuquerque &

Thesleff, 1968; Barrett & Crill, 1974; Adrian & Hodgkin, 1975; Sukhorukov et al., 1993; Major et al.,

1994; Solsona et al., 1998; Thurbon et al., 1998; Gentet et al., 2000). Similarly, the empirical

relationship $I_{th} = \frac{2.7 \cdot 10^{-2}}{R^{1.0}}$ (Table 6) yields $\Delta V_{th} = 27mV$, consistently with typical reported values

(Brock et al., 1952; Eccles et al., 1958), despite uncertainties in the value of the membrane resting

potential (Heckman & Enoka, 2012). This supports the conclusions that the relative voltage threshold

ΔV_{th} is constant within the MN pool (Coombs et al., 1955; Gustafsson & Pinter, 1984a; Gustafsson &

Pinter, 1984b; Powers & Binder, 2001), and that Ohm's law is followed in MNs (Glenn & Dement, 1981;

Spruston & Johnston, 1992; Kernell, 2006). Also, no correlation between $\frac{\Delta V_{th}}{R \cdot I_{th}}$ and C is reported in

Table 6, consistent with measurements performed by Gustafsson & Pinter (1984b), substantiating that

the dynamics of MN recruitment dominantly rely on R, I^{th} and ΔV_{th} (Heckman & Enoka, 2012). Finally,

the findings $R = 7.1 \cdot 10^9 \cdot \tau^{1.6}, C = 1.6 \cdot 10^{-10} \cdot \frac{1}{\tau^{0.6}}$ and $R_m = 6.0 \cdot 10^1 \cdot \tau, C_m = 1.0 \cdot 10^{-2} F \cdot m^{-2}$

in Table 6 yield $R \cdot C = 1.1 \cdot \tau$ and $R_m \cdot C_m = 0.6 \cdot \tau$, which is consistent with the classic empirical

relationship $\tau = RC = R_m C_m$ (Gustafsson & Pinter, 1984a; Zengel et al., 1985).

Relevance for other mammals

It is demonstrated in Figure 7 and with the calculation of the r^2 values in Table 7 that the mathematical

equations in Table 6 obtained from cat data adequately predict the normalized associations between

pairs of MN properties in rats and mice. However, the scaling factor K applied to the intercept k_e to

scale the normalized relationships $A = K \cdot k_e \cdot B^e$ is species-specific, being for example around 3

times less in mice than in cat for the $\{\tau; R\}$ and $\{C; R\}$ pairs in mice (Figure 7) and explaining the large

obtained $nRMSE$ values in these cases. Absolute values of the $\{I_{th}; R\}$ pair were nevertheless

accurately predicted in both rats and mice from the cat relationships, as explained by the larger values

of R which counterbalances the respectively lower I_{th} values in mice and rats than in cats (Manuel et

al., 2019; Highlander et al., 2020). Therefore, the mathematical equations in Table 6 should be used

for the modelling of cat MN pools only, while the normalized equations in Supplementary material

should be used and scaled with species-specific data if investigating other mammals (mice, rats,

humans). It was not assessed whether these relationships applied to non-mammal species.

Henneman's size principle of motor unit (MU) recruitment

Table 8 reports statistically significant power relationships of positive power values between MN and

mU indices of size. These results substantiate the concept that S_{MN} and S_{mU} are positively correlated

in a MU pool and that large MNs innervate large mUs (Henneman, 1981; Heckman & Enoka, 2012), a

statement that has never been demonstrated from the concurrent direct measurement of S_{MN} and

S_{mU}. Besides, considering the positive $I_{th} - S_{MN}$ correlation (Table 6), and that the mU force

recruitment threshold F^{th} is positively correlated to F^{tet} (Heckman & Enoka, 2012) and thus to S_{mU}

(Table 2), larger MUs have both larger current and force recruitment thresholds I_{th} and F^{th} than

relatively smaller MUs, which are thus recruited first, consistently with the Henneman's size principle

of MU recruitment (Henneman, 1957; Wuerker et al., 1965; Henneman et al., 1965a; Henneman et al.,

1965b; Henneman et al., 1974; Henneman, 1981; Henneman, 1985). The terminologies 'small MU',

'low-force MU' and 'low-threshold MU' are thus equivalent. Henneman's size principle thus relies on

the amplitude of the MN membrane resistance (Binder et al., 1996; Powers & Binder, 2001; Heckman

& Enoka, 2012). Finally, the relationships $S_{MN} \propto CV^{1.4} \propto \tau^{-0.7} \propto AHP^{-0.7}$ (Table 6) suggest that high-

threshold MUs rely on relatively faster MN dynamics, which might partially explain why large MNs can

attain relatively larger firing rates than low-thresholds MNs.

It has been repeatedly attempted to extend Henneman's size principle and the correlations between

the MU properties in Table 1 to the concept of 'MU type' (Burke & Ten Bruggencate, 1971; Burke,

1981; Bakels & Kernell, 1993; Powers & Binder, 2001). While a significant association between 'MU

type' and indices of MU size has been observed in some animal (Fleshman et al., 1981; Burke et al.,

1982; Zengel et al., 1985) and a few human (Milner-Brown et al., 1973; Stephens & Usherwood, 1977;

Garnett et al., 1979; Andreassen & Arendt-Nielsen, 1987) studies, it has however not been observed

in other animal studies ((Bigland-Ritchie et al., 1998) for a review) and in the majority of human

investigations (Sica & McComas, 1971; Goldberg & Derfler, 1977; Yemm, 1977; Young & Mayer, 1982;

Thomas et al., 1990; Nordstrom & Miles, 1990; Elek et al., 1992; Macefield et al., 1996; Cutsem et al.,

1997; Mateika et al., 1998; Fuglevand et al., 1999; Keen & Fuglevand, 2004). Moreover, the reliability

of these results is weakened by the strong limitations of the typical MU type identification protocols.

Sag experiments are irrelevant in humans (Buchthal & Schmalbruch, 1970; Thomas et al., 1991; Bakels

& Kernell, 1993; Macefield et al., 1996; Bigland-Ritchie et al., 1998; Fuglevand et al., 1999), and lack

consistency with other identification methods (Nordstrom & Miles, 1990). MU type identification by

twitch contraction time measurements is limited by the strong sources of inaccuracy involved in the

transcutaneous stimulation, intramuscular microstimulation, intraneural stimulation, and spike-

triggered averaging techniques (Taylor et al., 2002; Keen & Fuglevand, 2004; McNulty & Macefield,

2005; Negro et al., 2014; Dideriksen & Negro, 2018). Finally, as muscle fibres show a continuous

distribution of contractile properties among the MU pool, some MUs fail to be categorized in discrete

MU types in some animal studies by histochemical approaches (Reinking et al., 1975; Totosy de

Zepetnek et al., 1992). Owing to these conflicting results and technical limitations, MU type may not

be related to MN size and the basis for MU recruitment during voluntary contractions (McNulty &

Macefield, 2005; Duchateau & Enoka, 2011).

Limitations

The mathematical relationships derived Table 6 and the conclusions drawn in the Discussion present some limitations, which were partly discussed in the Methods and Result sections and are summarised here. A first limitation is related to the normalization of all datasets to the highest measured value retrieved in each study; this approach is only valid if the experimental studies identified the same largest MN relatively to the MN populations under investigation. This cannot be verified and is unlikely considering the small MN populations investigated in most eligible studies. A second limitation is due to the limited processable experimental data available in the literature, especially in rats and mice, as also discussed by Heckman & Enoka (2012). Therefore, some pairs of MN properties were investigated in only one study, especially those involving C, preventing inter-study comparisons. Also for this reason, measurements obtained from MNs innervating different muscles from cats of different sex were merged into unique datasets, implicitly assuming no sex- and muscle-specific distributions of MN properties within the MN pools. Besides, only MNs innervating the hindlimbs were investigated. The extrapolation of the relationships in Table 6 to rat and mouse species could only be assessed against a small set of studies and for three associations only. Finally, the conclusions on the associations between S_{mU} and S_{MN} relied on few studies providing few measurements, due to the challenging experimental protocols and the considerable amount of work required to measure both MN and mU properties in single MUs. Thirdly, the available data comes from relatively dated cat studies, as observed in Manuel et al. (2019). Thus, recent computer-assisted techniques of MN morphometric measurements were notably not applicable in those studies, yielding important sources of inaccuracies, as discussed. Therefore, the relationships involving S_{neuron} are subjected to a higher level of inaccuracy. Consequently, the mathematical relationships in Table 6 best reproduce published cat data but might remain a crude description of the association of MN properties in real MN pools. They however meet the level of inaccuracy that would be obtained from direct experimental approaches.

Conclusion

This study provides the first empirical and algebraical proof that the MN size S_{MN} and the MN electrophysiological properties (ACV, AHP, R, I_{th}, C and τ) are related. The derived mathematical relationships between any of these MN properties and/or S_{MN} are provided in Table 6. They accurately describe the experimental cat data available in the literature and provide for the first time a method for building virtual MN profiles of inter-consistent MN-specific properties in mammals and a framework to generate hypothesis for experimental studies aiming at investigating currently unreported relationships.

References

Adrian RH, Hodgkin AL (1975) Conduction velocity and gating current in the squid giant axon. Proc. Royal Soc. 189:81-86.

Albuquerque EX, Thesleff S (1968) A comparative study of membrane properties of innervated and chronically denervated fast and slow skeletal muscles of the rat. Acta Physiol. Scand. 73:471-480.

Amendola J, Durand J (2008) Morphological differences between wild-type and transgenic superoxide dismutase 1 lumbar motoneurons in postnatal mice. Journal of comparative neurology (1911) 511:329-341.

Andreassen S, Arendt-Nielsen L (1987) Muscle fibre conduction velocity in motor units of the human anterior tibial muscle: A new size principle parameter. J. Physiol. 391:561-571.

Ankit, R. (2020) WebPlotDigitizer. **4.4** https://automeris.io/WebPlotDigitizer.

Appelberg B, Emonet-Dénand F (1967) Motor units of the first superficial lumbrical muscle of the cat. J. Neurophysiol. 30:154-160.

Bagust J (1974) Relationships between motor nerve conduction velocities and motor unit contraction characteristics in a slow twitch muscle of the cat. J. Physiol. 238:269-278.

Bakels R, Kernell D (1993) Matching between motoneurone and muscle unit properties in rat medial gastrocnemius. J. Physiol. 463:307-324.

Barrett JN, Crill WE (1974) Specific membrane properties of cat motoneurones. J. Physiol. 239:301-324.

Bigland-Ritchie B, Fuglevand AJ, Thomas CK (1998) Contractile properties of human motor units: Is man a cat? The Neuroscientist 4:240-249.

Binder, M. D., Heckman, C. J. & Powers, R. K. (1996) The physiological control of motoneuron activity. In Rowell LB, S. J. (ed), *Handbook of Physiology: Exercise, Regulation and Integration of Multiple Systems.* New York: Oxford Univ Press, pp. 3-53.

Bodine SC, Roy RR, Eldred E, Edgerton VR (1987) Maximal force as a function of anatomical features of motor units in the cat tibialis anterior. J. Neurophysiol. 57:1730-1745.

Brandenburg, J. E.,Fogarty, M.J.,Brown, A.D.,Sieck, G.C.(2020) Phrenic motor neuron loss in an animal model of early onset hypertonia. Journal of Neurophysiology, **123**, 1682.

Brandenburg, J. E.,Gransee, H.M.,Fogarty, M.J.,Sieck, G.C.(2018) Differences in lumbar motor neuron pruning in an animal model of early onset spasticity. Journal of Neurophysiology, **120**, 601.

Brock LG, Coombs JS, Eccles JC (1952) The recording of potentials from motoneurones with an intracellular electrode. J.Physiol.(Lond.) 117:431-460.

Buchthal F, Schmalbruch H (1970) Contraction times and fibre types in intact human muscle. Acta Physiol. Scand. 79:435-452.

Burke RE, Ten Bruggencate G (1971) Electrotonic characteristics of alpha motoneurones of varying
size. J. Physiol. 212:1-20.

Burke RE, Dum RP, Fleshman JW, Glenn LL, Lev-Tov A, O'Donovan MJ, Pinter MJ (1982) An HRP study
of the relation between cell size and motor unit type in cat ankle extensor motoneurons. J. Comp.
Neurol. 209:17-28.

Burke RE, Tsairis P (1973) Anatomy and innervation ratios in motor units of cat gastrocnemius. J.
Physiol. 234:749-765.

Burke RE (1968) Firing patterns of gastrocnemius motor units in the decerebrate cat. J. Physiol.
196:631-654.

Burke, R. E. (1981) Motor units: Anatomy, physiology, and functional organization. In Brooks, V. B.
(ed), *Handbook of Physiology, the Nervous System, Motor Control.* American Physiological Society,
pp. 345-422.

Button DC, Kalmar JM, Gardiner K, Marqueste T, Zhong H, Roy RR, Edgerton VR, Gardiner PF (2008)
Does elimination of afferent input modify the changes in rat motoneurone properties that occur
following chronic spinal cord transection? The Journal of physiology 586:529-544.

Carp JS, Tennissen AM, Mongeluzi DL, Dudek CJ, Chen XY, Wolpaw JR (2008) An in vitro protocol for
recording from spinal motoneurons of adult rats. Journal of Neurophysiology 100:474-481.

Carrascal L, Nieto-Gonzalez JL, Torres B, Nunez-Abades P (2009) Changes in somatodendritic
morphometry of rat oculomotor nucleus motoneurons during postnatal development. Journal of
comparative neurology (1911) 514:189-202.

Chamberlain S, Lewis DM (1989) Contractile characteristics and innervation ratio of rat soleus motor
units. J. Physiol. 412:1-21.

Coombs JS, Eccles JC, Fatt P (1955) The electrical properties of the motoneurone membrane.
J.Physiol.(Lond.) 130:291-325.

Copray, S.,Kernell, D.(2000) Neurotrophins and trk-receptors in adult rat spinal motoneurons:
Differences related to cell size but not to 'slow/fast' specialization. Neuroscience Letters, **289**, 217.

Cullheim S, Fleshman JW, Glenn LL, Burke RE (1987) Membrane area and dendritic structure in type-
identified triceps surae alpha motoneurons. Journal of comparative neurology (1911) 255:68-81.

Cullheim S (1978) Relations between cell body size, axon diameter and axon conduction velocity of
cat sciatic α-motoneurons stained with horseradish peroxidase. Neurosci. Lett. 8:17-20.

Cutsem MV, Feiereisen P, Duchateau J, Hainaut K (1997) Mechanical properties and behaviour of
motor units in the tibialis anterior during voluntary contractions. Can. J. Appl. Physiol. 22:585-597.

De Lourdes Martínez-Silva M, Imhoff-Manuel RD, Sharma A, Heckman CJ, Shneider NA, Roselli F,
Zytnicki D, Manuel M (2018) Hypoexcitability precedes denervation in the large fast-contracting
motor units in two unrelated mouse models of ALS. ELife 7:e30955.

Deardorff AS, Romer SH, Deng Z, Bullinger KL, Nardelli P, Cope TC, Fyffe REW (2013) Expression of
postsynaptic Ca2+-activated K+ (SK) channels at C-bouton synapses in mammalian lumbar -
motoneurons. The Journal of physiology 591:875-897.

Delestrée, N.,Manuel, M.,Iglesias, C.,Elbasiouny, S.M.,Heckman, C.J.,Zytnicki, D.(2014) Adult spinal
motoneurones are not hyperexcitable in a mouse model of inherited amyotrophic lateral sclerosis.
The Journal of Physiology, **592**, 1687.

Destombes J, Horcholle-Bossavit G, Thiesson D, Jami L (1992) Alpha and gamma motoneurons in the
peroneal nuclei of the cat spinal cord: An ultrastructural study. Journal of comparative neurology
(1911) 317:79-90.

Dideriksen JL, Negro F (2018) Spike-triggered averaging provides inaccurate estimates of motor unit
twitch properties under optimal conditions. J Electromyogr Kinesiol 43:104-110.

Dong Y, Mihalas S, Russell A, Etienne-Cummings R, Niebur E (2011) Estimating parameters of
generalized integrate-and-fire neurons from the maximum likelihood of spike trains. Neural Comput.
23:2833-2867.

Donselaar Y, Kernell D, Eerbeek O (1986) Soma size and oxidative enzyme activity in normal and
chronically stimulated motoneurones of the cat's spinal cord. Brain Research 385:.

Duchateau J, Enoka RM (2011) Human motor unit recordings: Origins and insight into the integrated
motor system. Brain Res. 1409:42-61.

Dukkipati, S. S.,Garrett, T.L.,Elbasiouny, S.M.(2018) The vulnerability of spinal motoneurons and
soma size plasticity in a mouse model of amyotrophic lateral sclerosis. J Physiol, **596**, 1723.

Dum RP, Kennedy TT (1980) Physiological and histochemical characteristics of motor units in cat
tibialis anterior and extensor digitorum longus muscles. J. Neurophysiol. 43:1615-1630.

Eccles JC, Libet B, Young RR (1958) The behaviour of chromatolysed motoneurones studied by
intracellular recording. J.Physiol.(Lond.) 143:11-40.

Eccles JC, Eccles RM, Lundberg A (1957) Durations of after-hyperpolarization of motoneurones
supplying fast and slow muscles. Nature 179:866-868.

Elek JM, Kossev A, Dengler R, Schubert M, Wohlfahrt K, Wolf W (1992) Parameters of human motor
unit twitches obtained by intramuscular microstimulation. Neuromuscul. Disord. 2:261-267.

Emonet-Dénand F, Hunt CC, Petit J, Pollin B (1988) Proportion of fatigue-resistant motor units in
hindlimb muscles of cat and their relation to axonal conduction velocity. J. Physiol. 400:135-158.

Enoka RM (1995) Morphological features and activation patterns of motor units. J Clin Neurophysiol
12:538-559.

Filipchuk AA, Durand J (2012) Postnatal dendritic development in lumbar motoneurons in mutant
superoxide dismutase 1 mouse model of amyotrophic lateral sclerosis. Neuroscience 209:144-154.

Fleshman JW, Munson JB, Sypert GW, Friedman WA (1981) Rheobase, input resistance, and motor-
unit type in medial gastrocnemius motoneurons in the cat. J Neurophysiol 46:1326-1338.

Foehring RC, Sypert GW, Munson JB (1987) Motor-unit properties following cross-reinnervation of
cat lateral gastrocnemius and soleus muscles with medial gastrocnemius nerve. II. influence of
muscle on motoneurons. J Neurophysiol 57:1227-1245.

Foehring RC, Sypert GW, Munson JB (1986) Properties of self-reinnervated motor units of medial
gastrocnemius of cat. II. axotomized motoneurons and time course of recovery. J Neurophysiol
55:947-965.

Fogarty, M. J.,Mu, E.W.H.,Lavidis, N.A.,Noakes, P.G.,Bellingham, M.C.(2020) Size-Dependent
vulnerability of lumbar motor neuron dendritic degeneration in SOD1 G93A mice. Anat. Rec., **303**,
1455.

Fogarty MJ, Omar TS, Zhan W, Mantilla CB, Sieck GC (2018) Phrenic motor neuron loss in aged rats.
J.Neurophysiol. 119:1852-1862.

Frank K, Fuortes M (1956) Stimulation of spinal motoneurones with intracellular electrodes.
J.Physiol.(Lond.) 134:451-470.

Friese A, Kaltschmidt JA, Ladle DR, Sigrist M, Jessell TM, Arber S (2009) Gamma and alpha motor
neurons distinguished by expression of transcription factor Err3. Proceedings of the National
Academy of Sciences - PNAS 106:13588-13593.

Fuglevand AJ, Macefield VG, Bigland-Ritchie B (1999) Force-frequency and fatigue properties of
motor units in muscles that control digits of the human hand. J Neurophysiol 81:1718-1729.

Gao H, Glatzer NR, Williams KW, Derbenev AV, Liu D, Smith BN (2009) Morphological and
electrophysiological features of motor neurons and putative interneurons in the dorsal vagal
complex of rats and mice. Brain research 1291:40-52.

Gardiner PF (1993) Physiological properties of motoneurons innervating different muscle unit types
in rat gastrocnemius. J Neurophysiol 69:1160-1170.

Garnett RAF, O'Donovan MJ, Stephens JA, Taylor A (1979) Motor unit organization of human medial
gastrocnemius. J. Physiol. 287:33-43.

Gentet L, Stuart GJ, Clements JD (2000) Direct measurement of specific membrane capacitance in
neurons. Biophys. J. 79:314-320.

Glenn LL, Dement WC (1981) Membrane resistance and rheobase of hindlimb motoneurons during
wakefulness and sleep. J Neurophysiol 46:1076-1088.

Goldberg LJ, Derfler B (1977) Relationship among recruitment order, spike amplitude, and twitch
tension of single motor units in human masseter muscle. J Neurophysiol 40:879-890.

Gustafsson, Pinter MJ (1985) On factors determining orderly recruitment of motor units: A role for
intrinsic membrane properties. Trends Neurosci. 431-433.

Gustafsson, Pinter MJ (1984a) Relations among passive electrical properties of lumbar alpha-
motoneurones of the cat. J. Physiol. 356:401-431.

Gustafsson, Pinter MJ (1984b) An investigation of threshold properties among cat spinal alpha-
motoneurones. J. Physiol. 357:453-483.

Gustafsson B (1979) Changes in motoneurone electrical properties following axotomy. J. Physiol.
293:197-215.

Heckman CJ, Enoka RM (2012) Motor unit. Compr. Physiol. 2:2629-2682.

Hegedus J, Putman CT, Gordon T (2007) Time course of preferential motor unit loss in the SOD1
G93A mouse model of amyotrophic lateral sclerosis. Neurobiology of disease 28:154-164.

Henneman E (1985) The size-principle: A deterministic output emerges from a set of probabilistic
connections. J. Exp. Biol. 115:105-112.

Henneman E (1981) Recruitment of motoneurons : The size principle. Motor Unit Types, Recruitment
and Plasticity in Health and Disease 26-60.

Henneman E, Clamann HP, Gillies JD, Skinner RD (1974) Rank order of motoneurons within a pool:
Law of combination. J Neurophysiol 37:1338-1349.

Henneman E, Somjen G, Carpenter DO (1965a) Functional significance of cell size in spinal
motoneurons. J Neurophysiol 28:560-580.

Henneman E, Somjen G, Carpenter DO (1965b) Excitability and inhibitibility of motoneurons of
different sizes. J Neurophysiol 28:599-620.

Henneman E (1957) Relation between size of neurons and their susceptibility to discharge. Science
126:1345-1347.

Highlander, M. M.,Allen, J.M.,Elbasiouny, S.M.(2020) Meta-analysis of biological variables' impact on
spinal motoneuron electrophysiology data. Journal of Neurophysiology, **123**, 1380.

Huh, S.,Heckman, C.J.,Manuel, M.(2021) Time course of alterations in adult spinal motoneuron
properties in the SOD1(G93A) mouse model of ALS. eNeuro, **8**, ENEURO.

Ishihara, A.,Kawano, F.,Ishioka, N.,Oishi, H.,Higashibata, A.,Shimazu, T.,Ohira, Y.(2001) Growth-
related changes in cell body size and succinate dehydrogenase activity of spinal motoneurons
innervating the rat soleus muscle. Int. j. dev. neurosci., **21**, 461.

Issa AN, Zhan W, Sieck GC, Mantilla CB (2010) Neuregulin-1 at synapses on phrenic motoneurons.
J.Comp.Neurol. 518:4213-4225.

Izhikevich EM (2004) Which model to use for cortical spiking neurons? IEEE Trans.Neural Networks
15:1063-1070.

Jami L, Petit J (1975) Correlation between axonal conduction velocity and tetanic tension of motor
units in four muscles of the cat hind limb. Brain Res. 96:114-118.

Kanda K, Hashizume K (1992) Factors causing difference in force output among motor units in the rat
medial gastrocnemius muscle. J. Physiol. 448:677-695.

Kanjhan, R.,Fogarty, M.J.,Noakes, P.G.,Bellingham, M.C.(2015) Developmental changes in the
morphology of mouse hypoglossal motor neurons. Brain Struct Funct, **221**, 3755.

Keen DA, Fuglevand AJ (2004) Distribution of motor unit force in human extensor digitorum assessed
by spike-triggered averaging and intraneural microstimulation. J Neurophysiol 91:2515-2523.

Kernell D (2006) *The Motoneurone and its Muscle Fibres.* Oxford University Press UK, .

Kernell D, Zwaagstra B (1989) Dendrites of cat's spinal motoneurones: Relationship between stem
diameter and predicted input conductance. J. Physiol. 413:255-269.

Kernell D, Monster AW (1981) Threshold current for repetitive impulse firing in motoneurones
innervating muscle fibres of different fatigue sensitivity in the cat. Brain Res. 229:193-196.

Kernell D, Zwaagstra B (1981) Input conductance, axonal conduction velocity and cell size among
hindlimb motoneurones of the cat. Brain Res. 204:311-326.

Kernell D (1966) Input resistance, electrical excitability, and size of ventral horn cells in cat spinal
cord. Science 152:1637-1640.

Knott S, Lewis DM, Luck JC (1971) Motor unit areas in a cat limb muscle. Exp.Neurol. 30:475-483.

Krawitz S, Fedirchuk B, Dai Y, Jordan LM, McCrea DA (2001) State-dependent hyperpolarization of
voltage threshold enhances motoneurone excitability during fictive locomotion in the cat. The
Journal of physiology 532:271-281.

Krutki P, Hałuszka A, Mrówczyński W, Gardiner PF, Celichowski J (2015) Adaptations of motoneuron
properties to chronic compensatory muscle overload. Journal of neurophysiology 113:2769-2777.

Lee RH, Heckman CJ (1998) Bistability in spinal motoneurons in vivo: Systematic variations in
persistent inward currents. Journal of neurophysiology 80:583-593.

Li Y, Brewer D, Burke RE, Ascoli GA (2005) Developmental changes in spinal motoneuron dendrites in
neonatal mice. Journal of comparative neurology (1911) 483:304-317.

Lucas SM, Ruff RL, Binder MD (1987) Specific tension measurements in single soleus and medial
gastrocnemius muscle fibers of the cat. Exp.Neurol. 95:142-154.

Lux HD, Pollen DA (1966) Electrical constants of neurons in the motor cortex of the cat.
J.Neurophysiol. 29:207-220.

Macefield VG, Fuglevand AJ, Bigland-Ritchie B (1996) Contractile properties of single motor units in
human toe extensors assessed by intraneural motor axon stimulation. J Neurophysiol 75:2509-2519.

Major G, Larkman AU, Jonas P, Sakmann B, Jack JJ (1994) Detailed passive cable models of whole-cell
recorded CA3 pyramidal neurons in rat hippocampal slices. J.Neurosci. 14:4613-4638.

Mantilla, C. B.,Zhan, W.,Gransee, H.M.,Prakash, Y.S.,Sieck, G.C.(2018) Phrenic motoneuron structural
plasticity across models of diaphragm muscle paralysis. J Comp Neurol, **526**, 2973.

Manuel M, Chardon M, Tysseling V, Heckman CJ (2019) Scaling of motor output, from mouse to
humans. Physiology (Bethesda, Md.) 34:5-13.

Manuel M, Heckman CJ (2011) Adult mouse motor units develop almost all of their force in the
subprimary range: A new all-or-none strategy for force recruitment? Journal of Neuroscience
31:15188-15194.

Manuel M, Iglesias C, Donnet M, Leroy F, Heckman CJ, Zytnicki D (2009a) Fast kinetics, high-
frequency oscillations, and subprimary firing range in adult mouse spinal motoneurons. The Journal
of neuroscience 29:11246-11256.

Manuel M, Iglesias C, Donnet M, Leroy F, Heckman CJ, Zytnicki D (2009b) Fast kinetics, high-
frequency oscillations, and subprimary firing range in adult mouse spinal motoneurons. The Journal
of neuroscience 29:11246-11256.

Mateika JH, Essif EG, Dellorusso C, Fregosi RF (1998) Contractile properties of human nasal dilator
motor units. J Neurophysiol 79:371-378.

McNulty PA, Macefield VG (2005) Intraneural microstimulation of motor axons in the study of human
single motor units. Muscle & Nerve: Official Journal of the American Association of Electrodiagnostic
Medicine 32:119-139.

McPhedran AM, Wuerker RB, Henneman E (1965) Properties of motor units in a homogeneous red
muscle (soleus) of the cat. J Neurophysiol 28:71-84.

Mierzejewska-Krzyżowska B, Bukowska D, Taborowska M, Celichowski J (2014) Sex differences in the
number and size of motoneurons innervating rat medial gastrocnemius muscle. Anatomia, histologia,
embryologia 43:182-189.

Milner-Brown HS, Stein RB, Yemm R (1973) The orderly recruitment of human motor units during
voluntary isometric contractions. J. Physiol. 230:359-370.

Mitra P, Brownstone RM (2012) An in vitro spinal cord slice preparation for recording from lumbar
motoneurons of the adult mouse. Journal of Neurophysiology 107:728-741.

Moschovakis AK, Burke RE, Fyffe REW (1991) The size and dendritic structure of HRP-labeled gamma
motoneurons in the cat spinal cord. Journal of comparative neurology (1911) 311:531-545.

Munson JB, Foehring RC, Lofton SA, Zengel JE, Sypert GW (1986) Plasticity of medial gastrocnemius
motor units following cordotomy in the cat. Journal of neurophysiology 55:1454.

Nakanishi ST, Whelan PJ (2010) Diversification of intrinsic motoneuron electrical properties during
normal development and botulinum Toxin–Induced muscle paralysis in early postnatal mice.
J.Neurophysiol. 103:2833-2845.

Negro F, Yavuz U, Farina D (2016) The human motor neuron pools receive a dominant slow-varying
common synaptic input. J. Physiol. 594:5491-5505.

Negro F, Yavuz U, Farina D (2014) Limitations of the spike-triggered averaging for estimating motor
unit twitch force: A theoretical analysis. PloS one 9:e92390.

Nordstrom MA, Miles TS (1990) Fatigue of single motor units in human masseter. J. Appl. Physiol.
68:26-34.

Obregón G, Ermilov LG, Wen-Zhi Z, Sieck GC, Mantilla CB (2009) Modeling dendritic arborization
based on 3D-reconstructions of adult rat phrenic motoneurons. Revista Ingeniería Biomédica 3:47-
54.

Pinter MJ, Vanden Noven S (1989) Effects of preventing reinnervation on axotomized spinal
motoneurons in the cat. I. motoneuron electrical properties. Journal of neurophysiology 62:311-324.

Powers RK, Binder MD (2001) Input-output functions of mammalian motoneurons. Rev. Physiol.
Biochem. Pharmacol. 143:137-263.

Prakash YS, Mantilla CB, Zhan W, Smithson KG, Sieck GC (2000) Phrenic motoneuron morphology
during rapid diaphragm muscle growth. Journal of Applied Physiology 89:563-572.

Proske U, Waite PME (1974) Properties of types of motor units in the medial gastrocnemius muscle
of the cat. Brain Res. 67:89-101.

Rafuse VF, Pattullo MC, Gordon T (1997) Innervation ratio and motor unit force in large muscles: A
study of chronically stimulated cat medial gastrocnemius. J. Physiol. 499:809-823.

Rall W (1960) Membrane potential transients and membrane time constant of motoneurons.
Exp.Neurol. 2:503-532.

Rall W (1959) Branching dendritic trees and motoneuron membrane resistivity. Exp.Neurol. 1:491-
527.

Rall W (1957) Membrane time constant of motoneurons. Science 126:454.

Reinking RM, Stephens JA, Stuart DG (1975) The motor units of cat medial gastrocnemius: Problem of
their categorisation on the basis of mechanical properties. Exp. Brain Res. 23:301-313.

Sasaki M (1991) Membrane properties of external urethral and external anal sphincter
motoneurones in the cat. The Journal of physiology 440:345-366.

Sica REP, McComas AJ (1971) Fast and slow twitch units in a human muscle. J. Neurol. Neurosurg.
Psychiatry 34:113-120.

Solsona C, Innocenti B, Fernández JM (1998) Regulation of exocytotic fusion by cell inflation.
Biophys.J. 74:1061-1073.

Song Z, Hu J, Rudy B, Redman SJ (2006) Developmental changes in the expression of calbindin and
potassium-channel subunits Kv3. 1b and Kv3. 2 in mouse renshaw cells. Neuroscience 139:531-538.

Spruston N, Johnston D (1992) Perforated patch-clamp analysis of the passive membrane properties
of three classes of hippocampal neurons. J Neurophysiol 67:508-529.

Stephens JA, Stuart DG (1975) The motor units of cat medial gastrocnemius: Speed-size relations and
their significance for the recruitment order of motor units. Brain. Res. 91:177-195.

Stephens JA, Usherwood TP (1977) The mechanical properties of human motor units with special
reference to their fatiguability and recruitment threshold. Brain Res. 125:91-97.

Sukhorukov VL, Arnold WM, Zimmermann U (1993) Hypotonically induced changes in the plasma
membrane of cultured mammalian cells. J.Membr.Biol. 132:27-40.

Swett, Wikholm, Blanks, Conley (1986) Motoneurons of the rat sciatic nerve. Exp. Neurol. 93:227-
252.

Taylor AM, Steege JW, Enoka RM (2002) Motor-unit synchronization alters spike-triggered average
force in simulated contractions. J Neurophysiol 88:265-276.

Teeter C, Iyer R, Menon V, Gouwens N, Feng D, Berg J, Szafer A, Cain N, Zeng H, Hawrylycz M, Koch C,
Mihalas S (2018) Generalized leaky integrate-and-fire models classify multiple neuron types. Nat
Commun 9:1-15.

Thomas CK, Bigland-Richie B, Johansson RS (1991) Force-frequency relationships of human thenar
motor units. J. Neurophysiol. 65:1509-1516.

Thomas CK, Johansson RS, Westling G, Bigland-Ritchie B (1990) Twitch properties of human thenar
motor units measured in response to intraneural motor-axon stimulation. J Neurophysiol 64:1339-
1346.

Thurbon D, Hans-R. Lüscher, Hofstetter T, Redman SJ (1998) Passive electrical properties of ventral
horn neurons in rat spinal cord slices. J.Neurophysiol. 79:2485-2502.

Totosy de Zepetnek, Zung HV, Erdebil S, Gordon T (1992) Innervation ratio is an important
determinant of force in normal and reinnervated rat tibialis anterior muscles. J Neurophysiol
67:1385-1403.

Turkin VV, O'Neill D, Jung R, Iarkov A, Hamm TM (2010) Characteristics and organization of
discharge properties in rat hindlimb motoneurons. Journal of Neurophysiology 104:1549-1565.

Ulfhake, Kellerth JO (1981) A quantitative light microscopic study of the dendrites of cat spinal α-
motoneurons after intracellular staining with horseradish peroxidase. J. Comp. Neurol. 202:571-583.

Ulfhake B, Kellerth JO (1984) Electrophysiological and morphological measurements in cat
gastrocnemius and soleus α-motoneurones. Brain Res. 307:167-179.

Ulfhake B, Cullheim S (1988) Postnatal development of cat hind limb motoneurons. III: Changes in
size of motoneurons supplying the triceps surae muscle. Journal of comparative neurology (1911)
278:103-120.

von Steyern FV, Martinov V, Rabben I, Njå A, de Lapeyrière O, Lømo T (1999) The homeodomain
transcription factors islet 1 and HB9 are expressed in adult alpha and gamma motoneurons identified
by selective retrograde tracing. The European journal of neuroscience 11:2093-2102.

Wuerker RB, McPhedran AM, Henneman E (1965) Properties of motor units in a heterogeneous pale
muscle (m. gastrocnemius) of the cat. J.Neurophysiol. 28:85-99.

Yemm R (1977) The orderly recruitment of motor units of the masseter and temporal muscles during
voluntary isometric contraction in man. J. Physiol. 265:163-174.

Young JL, Mayer RF (1982) Physiological alterations of motor units in hemiplegia. J. Neurol. Sci.
54:401-412.

Zengel JE, Reid SA, Sypert GW, Munson JB (1985) Membrane electrical properties and prediction of
motor-unit type of medial gastrocnemius motoneurons in the cat. J Neurophysiol 53:1323-1344.

Zwaagstra, Kernell (1981) Sizes of soma and stem dendrites in intracellularly labelled α-
motoneurones of the cat. Brain Research 204:295-309.

Zwaagstra B, Kernell D (1980) The duration of after-hyperpolarization in hindlimb alpha
motoneurones of different sizes in the cat. Neurosci. Lett. 19:303-307.

1041