Nontypeable *Haemophilus influenzae* infection impedes *Pseduomonas aeruginosa*

colonization and persistence in mouse respiratory tract

Natalie Lindgren [1,2], Lea Novak[3], Benjamin C. Hunt [1,2], Melissa S. McDaniel [1,2], and W. Edward

Swords [1,2#]

[1] Department of Medicine, Division of Pulmonary, Allergy, and Critical Care Medicine

[2] Gregory Fleming James Center for Cystic Fibrosis Research

[3] Department of Pathology, Division of Anatomic Pathology

University of Alabama at Birmingham

Running title: Competitive infections in CF related opportunists

Key words: Cystic fibrosis, bacteria, *Haemophilus*, *Pseudomonas*, biofilm

Communicating author:

1918 University Boulevard, MCLM 818

Birmingham, AL 35294

wswords@uabmc.edu

ABSTRACT

Patients with cystic fibrosis (CF) experience lifelong respiratory infections which are a significant

cause of morbidity and mortality. These infections are polymicrobial in nature wherein the

predominant bacterial species changes as patients age. Young patients have populations

dominated by pathobionts such as nontypeable *Haemophilus influenzae* (NTHi), that are

eventually supplanted by pathogens such as *Pseudomonas aeruginosa* (*Pa*), which are more

typical of late-stage CF disease. In this study, we investigated how initial colonization with NTHi

impacts colonization and persistence of *Pa* in the respiratory tract. Analysis of polymicrobial

biofilms *in vitro* by confocal microscopy revealed that NTHi promoted greater levels of *Pa* biofilm

volume and diffusion. However, sequential infection of mice with NTHi followed by *Pa* showed

significant reduction in *Pa* in the lungs as compared to infection with *Pa* alone. Coinfected mice

also had reduced severity of airway tissue damage and lower levels of inflammatory cytokines

as compared mice infected with *Pa* alone. Similar results were observed using heat-inactivated

NTHi bacteria prior to *Pa* introduction. Based on these results, we conclude that NTHi can

significantly reduce susceptibility to subsequent *Pa* infection, most likely due to immune priming

rather than a direct competitive interaction between species. These findings have potential

significance with regard to therapeutic management of early life infections in patients with CF.

Word count: 214 words

INTRODUCTION

Cystic fibrosis (CF) is an autosomal recessive genetic illness that causes dysfunctional ionic transport at epithelial surfaces, with concomitant impacts on mucociliary defenses and clearance in the lungs. As a consequence, patients with CF experience lifelong infections of the airway mucosa that cause chronic inflammation and epithelial damage, among other sequelae (1-3). Despite therapeutic gains that have significantly increased average lifespan for CF patients, airway infections and associated respiratory complications remain a leading cause of morbidity and mortality in patients with CF disease (4). The microbial populations within the CF lung are complex, polymicrobial communities that undergo dynamic changes that correlate with patients' age and overall health, and can be a determinant of disease severity (3, 5-7). Despite years of study of opportunistic infections in the context of CF, there remains much to be learned about how specific pathogens and opportunists impact the course and severity of disease (6-9). Nontypeable *Haemophilus influenzae* (NTHi) is a non-encapsulated, Gram-negative pathobiont that normally resides in the human nasopharynx and upper airways as part of the normal flora. (10-12). Although typically benign in healthy individuals, NTHi is among the most common bacterial species isolated from CF patients during the first year of life and is thus thought of as a common early-stage pathogen (13-15). As with other mucosal airway infections, NTHi persists in the lung within biofilm communities (16, 17).

As CF patients age, the bacterial population diversity typically declines, and patients become chronically colonized with late-stage pathogens such as *Pseudomonas aeruginosa* (*Pa*) (8, 9, 18, 19). *Pa* is a Gram-negative opportunistic pathogen that has a high level of inherent resistance to antimicrobials and host immune effectors and is therefore difficult to treat (20). To adapt to the CF airway environment, *Pa* populations typically undergo a genetic conversion to a mucoid phenotype that overexpresses the exopolysaccharide alginate, which is a component of *Pa* biofilm (21, 22). Mucoid conversion of *Pa* is associated with increased antibiotic resistance, persistent inflammation, and overall worse clinical outcomes (23). Initial *Pa* infections, which

may take place within the first year of life, frequently become chronic after mucoid conversion

(19, 20). While the role of *Pa* in end-stage CF disease has been well established, less has been

done to establish the role of *Pa* in early-stage CF disease, particularly the ages in which NTHi

colonization peaks in frequency (4, 9). We hypothesize that the NTHi infections that are

common in early childhood in CF, can potentially influence subsequent infections, with impacts

on progression of CF disease.

Bacteria reside in the CF lung as a complex, evolving polymicrobial community in which

individual organisms may interact with one another in a synergistic, antagonistic, or null fashion

(11, 24-26). Changes in these communities as different species, or strains within species,

interact are thought to greatly influence antibiotic susceptibility, immune evasion, and mutations

for chronic colonization, all influencing patient outcomes (8, 27). In order to better model the

polymicrobial airway infections associated with CF and other conditions, we have carried out

multispecies experimental approaches (5, 8, 28). While we know that *Pa* and NTHi clearly

reside in the airways at the same time, there is still a need for better understanding of how each

species is impacted during polymicrobial infection.

RESULTS

In this study, we aimed to assess the polymicrobial interaction between NTHi and *Pa*

both *in vitro* and *in vivo*, using clinically isolated bacterial strains from patients with CF. Mice

were intratracheally infected with NTHi and/or mucoid *Pa* (mPA) to produce localized, acute

respiratory infections. To model the temporal acquisition of NTHi and *Pa* in CF patients, we

tested whether pre-colonization with NTHi influences colonization of *Pa* and began to

characterize how this interaction may influence the innate immune response.

Nontypeable *H. influenzae* and *P. aeruginosa* form polymicrobial biofilms *in vitro* that

support *Pa* growth.

In any environment that contains more than one bacterial species, including the CF lungs,

organisms may interact with one another synergistically, competitively, or null (5)(6). To begin to

characterize the polymicrobial interaction between NTHi and *Pa*, we first wanted to evaluate

how these two organisms form a polymicrobial biofilm *in vitro*. Since NTHi and *Pa* are typically

acquired sequentially over a patient's lifespan, dual-species static biofilms were sequentially

seeded with NTHi 86028NP-gfp+, followed by mPA 08-31-mCherry+ 12 hours later.

Polymicrobial biofilms were grown for a total of 18 hours, while single-species biofilms of NTHi

and *Pa* were grown for 12 hours and 6 hours, respectively, at 37°C + 5% CO_2. These timepoints

for growth were selected based on the peak viability of NTHi and *Pa*, as determined by counts

of viability over time (Fig. S1A in supplemental material). Due to limitations of differential plating

that do not allow for quantification of both organisms in an NTHi/*Pa* co-culture, viable colony

counts of *Pa* from sequential dual biofilms were compared to single-species *Pa* biofilms only to

assess how pre-introduction of NTHi may influence *Pa* growth. *In vitro*, it appears that pre-

colonization with NTHi has no significant impact on *Pa* growth when both organisms are

inoculated at roughly the same amount (~10^8 CFU/mL) (Fig. S1B).

We then visualized NTHi 86028NP-gfp+ and mPA 08-31-mCherry+ single- and dual-species

fluorescent biofilms with confocal laser scanning microscopy (CLSM) after 18 hours of growth.

In single-species biofilms, both organisms formed lawn structures with some towers (Fig. 1A

and B). Sequentially seeded dual-species biofilms showed a complex lawn with more diffuse *Pa*

colony formation that appeared to support *Pa* growth. As expected, the spatial orientation of

dual biofilms were stratified, reflecting the sequential introduction of NTHi followed by *Pa*, with a

significant lawn structure of NTHi on the bottom (Fig. 1C). Quantification of *Pa* in both single-

and dual-species biofilms by BiofilmQ software revealed that the total volume of *Pa* biofilm is

greater in dual-species biofilms than in single-species *Pa* biofilms (Fig. 2A). Segmentation within

BiofilmQ software allowed for fluorescent images to be divided by 1 μm cubes with each cube

being individually tracked. Following segmentation, the mean cube surface is calculated as the

number of points between occupied and unoccupied volume in cube. The calculated mean cube

surface showed that dual biofilms have more *Pa* than single-species *Pa* biofilms, recapitulating

the results seen with total biofilm volume and verifying the accuracy of cubing (Fig. 2B). This

segmentation technique was used for further 3D visualization of single- and dual-species

biofilms of *Pa.* Segmentation shows that there is more diffuse local density of *Pa* associated

with dual-species (Fig. 2C) rather than single-species (Fig. 2D) growth. Overall, this

quantification suggests that *in vitro*, NTHi may positively influence *Pa* growth, producing greater

biofilm volume and more dispersed spatial distribution when grown together compared to a

single-species *Pa* biofilm.

Nontypeable *H. influenzae* infection decreases bacterial burden of *P. aeruginosa* in

mouse lung

While our *in vitro* results suggest that NTHi and *Pa* may positively interact in polymicrobial

biofilms, we aimed to evaluate how each species is impacted during polymicrobial infection

within the host. Through serial passage, we induced antibiotic resistance in a CF sputum-

derived clinical isolate of NTHi, HI-1, developing a model strain for acute infection (NTHi HI-1r)

that can be differentially plated in the presence of *Pa*. To model the sequential acquisition of an

early life pathogen (NTHi) followed by a later-stage pathogen (*Pa*) that typically ensues as CF

patients age, BALB/cJ mice were intratracheally infected with NTHi HI-1r (inoculum ~10^8

CFU/mouse) followed by a mucoid *P. aeruginosa* isolate, mPA 08-31 (inoculum ~10^7

CFU/mouse) 24 hours later. Both organisms colonized the lungs and were detectable by viable

plate counting of lung homogenate 48 hours post-infection. However, colonization of *Pa* was

significantly impeded after sequential introduction with NTHi compared to a *Pa* single-infection

(**$P<0.01$) (Fig. 3A). To determine the importance of temporal spacing in this interaction, we

also infected mice with both organisms concurrently (inoculum ~10^7 CFU/mL of each). When we

compared the bacterial burden of *Pa* in the lungs after a single-infection, a sequential dual-

infection, and a concurrent dual-infection, we saw that mice sequentially infected with NTHi

followed by *Pa* showed significant reduction in *Pa* abundance compared to a *Pa* only infection

(*$P<0.05$). Additionally, sequential dual-infection resulted in less *Pa* in lung homogenate than

concurrent dual-infection (****$P<0.0001$) (Fig. 3B). This suggests that the competitive interaction

between NTHi and *Pa* is dependent on time of introduction and *Pa* abundance in the lungs is

only impeded during sequential infection. Histopathological analysis of hematoxylin and eosin

(H&E) stained lung sections indicated immune cell infiltration and inflammation 48 hours after

infection of both NTHi and *Pa* infection. (Fig. 3C). H&E stained slides were graded for severity

of lung injury as indicated by neutrophil infiltration of alveoli, peri-vascular inflammation and peri-

bronchial inflammation. At 48-hours post-infection, where a high burden of both NTHi and *Pa*

were present in the airways, there was moderate tissue damage for all infection groups with no

difference between groups. All infections resulted in significantly more severe tissue damage

than a PBS control infection (****$P<0.0001$) (Fig. 3D). Immune cell infiltration was expressed by

total cell infiltrates counted in the bronchoalveolar lavage fluid (BALF) after cytospin preparation

and differential staining. As expected from histological results, all infection groups showed

significantly elevated total cell counts, as well as PMN infiltrates in those counts, than uninfected

controls with no significant difference between infection groups (*P<0.05, **P<0.01, ***P<0.001,

****P<0.0001) (Fig. 3E-F). Weight loss per animal was evaluated over the course of infection

and weight loss increased compared to uninfected controls (Fig. S2A). Taken together, these

results suggests that *in vivo*, NTHi impedes the bacterial burden of *Pa* in the airways when

introduced sequentially, but not concurrently, and infection with either organism results in

respiratory tissue damage with significant inflammatory response.

Impacts of nontypeable *H. influenzae* on *P. aeruginosa* are not strain-specific.

Our coinfection results indicate that a clinical isolate of NTHi, HI-1r, significantly inhibits the

bacterial burden of mucoid *P. aeruginosa* isolate mPA 08-31 in the airways. To determine if the

impact of NTHi on *Pa in vivo* is strain specific, we induced antibiotic resistance in laboratory

strain NTHi 86-028NP to develop strain NTHi 86r, allowing for differential plating of NTHi and

Pa. Following the sequential infection schematic previously described, we intratracheally

introduced NTHi 86-028NPr (inoculum ~10^8 CFU/mouse) into the airways of BALBc/J mice

followed by introduction of mPA 08-31 (inoculum ~10^7 CFU/mouse) 24 hours later. Unlike the

positive interaction between NTHi and *Pa* seen in our polymicrobial biofilms with the parent

strain of NTHi 86r (NTHi 86-028NP), in the airways, viable colony counts of *Pa* were

significantly decreased after pre-introduction with NTHi 86r compared to a *Pa* single-infection

(*P<0.05) (Fig. 4A). Mucoid conversion of *Pa* is a common consequence of long-term

colonization (19, 24, 29). For most patients, the initial colonizing *Pa* strain is nonmucoid, and

can remain so for relatively long periods of persistent infection (19). To extend the impact of this

phenotype to patients colonized with *Pa* before and after mucoid conversion, we then

determined if NTHi pre-introduction impedes the abundance of non-mucoid *Pa.* To do this, we

performed sequential infections with the clinical NTHi HI-1r followed by a non-mucoid *Pa* (PA

FRD1*mucA+).* Viable colony counts from lung homogenate revealed that, as in preceding

infection studies, *Pa* counts were significantly diminished after infection with NTHi compared to

Pa alone (*P<0.05) (Fig. 4B). Weight loss per animal was evaluated over the course of infection

and weight loss increased compared to uninfected controls (Fig. S2B-C). These results show

that NTHi infection significantly reduces the colonization/persistence of *Pa*, regardless of the

specific strain of NTHi or the mucoidy status of *Pa*.

Infection with nontypeable *H. influenzae* followed by *P. aeruginosa* results in less severe

airway tissue damage and less inflammatory cytokine infiltration than *P. aeruginosa*

alone.

While clinical isolate NTHi Hi-1r persists longer than previously established NTHi strains, such

as NTHi 86-028NP (data not shown), this strain still begins to be cleared from the airways of

mice around 72 hours post-infection, with viable colony counts below the limit of detection for

some animals. We wanted to test if *Pa* abundance is inhibited in the lungs once NTHi clearance

begins. BALBc/J mice were intratracheally infected with NTHi HI-1r, followed by mPA 08-31 24

hours later and viable colony counts from lung homogenate were assessed after 72 hours post-

dual infection, extending the previously established 48 hour timepoint. As expected, sequential

introduction of NTHi prior to *Pa* significantly reduced colonization of *Pa* compared to a *Pa* alone

(***P<0.001) (Fig. 5A). Reflecting the inhibition of *Pa* abundance in viable colony counts during

dual-infection, histological grading of tissue damage revealed a significantly decreased damage

score in dual-infected animals compared to a *Pa* single-infection (**P<0.01) (Fig. 5B). H&E

staining of lung sections of each infection group revealed more severe immune cell infiltration,

pleuritis, perivascular inflammation, and peri-bronchial inflammation 72 hours post-infection than

was seen at the 48 hour post-infection timepoint (Fig. 5C). Finally, total cell counts in BALF and

the percentage of PMN infiltrates in those counts was not different among single or dual-

infection groups, but was significantly increased compared to uninfected controls (*P<0.05,

**P<0.01, ****P<0.0001) (Fig. 5D-E). Weight loss per animal was evaluated over the course of

infection and weight loss increased compared to uninfected controls (Fig. S2D).

To evaluate the host immune response following NTHi and *Pa* single or dual-infections, we

measured cytokine and chemokine levels in BALF with DuoSet ELISA kits designed to detect

inflammatory markers including MCP-1, TNF-α, IL-10, IL-1α, IL-1β, or IL-6. Several of these

well-established inflammatory markers, including MCP-1, TNF-α, and the anti-inflammatory

marker IL-10 were significantly higher after *Pa* single-infection compared to a dual-infection with

both NTHi and *Pa* (****P<0.0001). (Fig. 6A-C). While IL-1α, IL-1β and IL-6 levels were not

different between single and dual-infected groups at 72 hours post-infection, the temporal

regulation of these acute response markers indicate that they may be associated with earlier

timepoints after bacterial introduction (Fig. 6D-F). Taken together, these data suggest that at 72

hours post-infection, after bacterial clearance has begun, NTHi still inhibits *Pa* bacterial burden

in the lungs. In addition to impeding abundace of *Pa*, sequential infection reduces the severity of

damage to airway tissues, as measured by histological scoring, and reduces inflammatory

cytokine levels in BALF.

Inhibitory effect of nontypeable *H. influenzae* on *P. aeruginosa* infection is independent

of bacterial viability.

Previous polymicrobial studies suggest that NTHi may impact subsequent infections directly via

interspecies interaction, or by priming the host response (30-33). To determine if the impact of

preceding infection on *Pa* requires viable NTHi bacteria, we sequentially infected mice with

heat-killed NTHi HI-1r (HK NTHi HI-1r) (inoculum equvialent to ~10^8 CFU/mL) followed by

mucoid *P. aeruginosa* mPA 08-31 (inoculum ~10^7 CFU/mL) for 48 hours. The total infection time

of this experiment was 72 hours, repeating our previously established timepoint in which live

NTHi infection reduced *Pa* infection. Viable colony counts from lung homogenate showed *Pa*

counts were still significantly reduced by pre-treatment with heat-killed NTHi (*P<0.05) (Fig. 7A).

Additionally, *Pa* infection resulted in a significantly higher histolopathological score than HK

NTHi/*Pa* dual-infection, suggesting that non-viable NTHi is sufficient to protect the lung

parenchyma from the severe effects of *Pa* infection, indicating the protection from tissue

damage previously seen at this time point during dual-infection is independent of NTHi viability

(**P<0.01) (Fig. 7B). H&E stained lung sections harvested from this experiment showed more

neutrophilic infiltration and more severe tissue damage after *Pa* single-infection compared to

dual-infection (Fig. 7C). As previously seen, the total cell counts in BALF and the percentage of

PMN infiltration was not different among single or dual-infection groups, but was significantly

increased compared to uninfected controls (*P<0.05, ***P<0.001, ****P<0.0001) (Fig. 5D-E).

Overall, we found that heat-killed NTHi is sufficient to significantly impede the abundance of *Pa*

in the lungs at 72 hours post-infection, suggesting that the interaction between NTHi and *Pa*

may be entirely mediated by the immune response of the host.

DISCUSSION

With the advent and widespread use of culture independent technologies for profiling microbial consortia, it has become clear that the populations within the lungs of patients with CF are diverse and subject to dynamic change. These shifts are likely of importance in determining whether species will persist or be cleared, as well as in the initiation of the inflammatory events that are a hallmark of CF disease (6-8, 24, 34, 35). The consequences of respiratory exacerbations include an intense neutrophilic inflammatory response that can cause significant lung damage and decreased respiratory function, contributing significantly to the morbidity and mortality associated with CF disease (8, 35). There is still much to learn regarding how specific organisms interact within the compromised respiratory tract of CF airways and how these interactions influence disease progression (8).

The complex microbial communities in the lung often change in dominant species and overall population diversity as patients age (19, 24). Decreased microbial diversity and the emergence of mucoid *Pa* as the dominant microbial species are significantly associated with detrimental host immune response, lung functional decline, and eventual patient mortality (19, 24). Previous studies have focused on characterizing key CF pathogens individually, however, these infections are undoubtedly polymicrobial in nature, influencing the pathogenesis of disease as organisms interact (3, 19). Nontypeable *Haemophilus influenzae*, a typically benign colonizer of nasopharyngeal region in healthy individuals, has long been recognized as an opportunistic pathogen of the airway mucosa, including in young children with CF (10, 36). While it is clear that NTHi and *Pa* share a common niche within the CF lung, including patients with coinfections, there remains much to be learned regarding how these organisms influence one another (14, 26-27). Our work considers the interaction between and disease consequences of two key CF pathogens, NTHi, as early-life pathogen, and *Pa*, a late-stage pathogen that can be acquired early in life.

Polymicrobial interactions in static biofilms provide basic information about interspecies

bacterial interactions. Our results indicate that NTHi and *Pa* form polymicrobial biofilms that may

support *Pa* growth (Fig. 1-2). However, to understand polymicrobial infections in patients, we

must how these two organisms interact within the lung environment. In a mouse model, we

tested how NTHi impacted *Pa* abundance in the lungs following sequential introduction,

representing how patients typically acquire these organisms over time, contrasted with

concurrent acquisition. We found that both NTHi and *Pa* colonize and persist in murine lungs for

at least 48 hours in single-infections, but viable colony counts of *Pa* from lung homogenate were

significantly reduced only following sequential introduction with NTHi (Fig. 3). This competitive

inhibition contrasts with the cooperative interaction seen in dual-species biofilms, possibly

because of the additional factor of the host immune response.

Bacterial infections in the CF lungs result in chronic inflammation as microbial and

phagocytic debris buildup in the airways, contributing to the reduced lung function and

overactive immune response associated with exacerbations (12). The inherently diminished

airways defenses associated with CFTR dysfunction are thought to predispose CF patients to

frequent microbial infections followed by an inadequate, yet overactive, innate immune

response that further initiate tissue damage by airway remodeling (12, 29). Histopathological

analysis of lung sections can evaluate immune cell infiltration within alveoli, pleura, perivascular

and peri-bronchial spaces, and grade the severity of tissue damage. In CF, the ensuing tissue

damage following chronic bacterial infection is a strong indicator of respiratory failure, the

leading cause of morbidity and mortality (14). Pre-introduction with NTHi not only reduces the

bacterial burden of *Pa* in the lungs, but also significantly reduces the visible epithelial damage

compared to a *Pa* single-infection, an effect that may reduce detrimental airway remodeling

following infection (Fig. 5). While dual-infection reduces the severity of airway parenchymal

damage, we wanted to know if inflammatory cytokine levels were reduced after dual-infection as

well. The BALF collected from single- and dual-infected mice showed trending decreases in

early response cytokines, such as IL-1-α, IL-1-β, and IL-6 in dual-infected mice compared to a

Pa single-infection (Fig. 6A-C). Since the interleukins are some of the first cytokines to respond

to bacterial stimuli, we suspect that significant differences between dual- and single-infected

groups may appear at earlier timepoints, prior to 72 hours post-infection (13, 30). We saw a

significant decrease in inflammatory cytokine levels of MCP-1, TNF-α, and IL-10 in sequentially

infected duals compared to a *Pa* single-infection, suggesting that the reduction in bacterial

burden also diminishes a potentially harmful innate immune response (Fig. 6D-F). Most notably,

this reduction in TNF-α could extend beneficial effects for CF patients, as the continual

production of this early response cytokine indicates an ongoing and overactive innate immune

response in the CF lungs (13).

While we suspect that NTHi and *Pa* are interacting in vivo, understanding the

mechanism of their interaction would provide a valuable therapeutic target. Previous studies

have noted that NTHi is capable of interacting with other bacterial species via interspecies

quorum signaling, a form of interbacterial communication that can influence the establishment of

polymicrobial biofilms and the development of chronic infection. This mechanism of interaction

commonly   utilizes   highly   conserved   interspecies   signaling   molecules   that   facilitate

communication across species, commonly through a canonical signal, autoinducer-2 (16).

Alternatively, NTHi has also been known to participate in a host-pathogen interactions that

overtly exclude the successful colonization of other pathogens, allowing for niche competition by

activation of opsonophagocytic killing (17). Therefore, possible mechanisms of interaction

between NTHi and *Pa* could be broadly categorized as active, requiring live bacterial cells, or

passive, solely reliant on cellular components. To determine if NTHi and *Pa* are engaging in a

pathogen-pathogen or host-pathogen interaction, we sequentially infected animals with heat-

killed NTHi HI-1r, followed by *Pa* 24 hours later. The introduction of heat-inactivated bacterial

components of NTHi provided the pathogen-associated-molecular-patterns (PAMPs) necessary

to mount an innate immune response against NTHi components, broadly including the

lipooligosaccharide (LOS) and other protein components, prior to *Pa* introduction (13). We found

that even in the absence of live bacteria, pre-introduction of NTHi is capable of diminishing *Pa*

bacterial burden in the lungs (Fig. 7A). This indicates that the host inflammatory response is key

to *Pa* bacterial clearance, suggesting the interaction between NTHi and *Pa* is host-pathogen

mediated. It is also important to note that our infection studies were carried out in mice with

normal CFTR function; extending these studies into genetically altered CF mutant animals will

be an important avenue for future work.

Defining how different microbes interact within the CF lung to impact bacterial

colonization, persistence, and host response has the potential to guide rational treatment

strategies to lengthen periods of asymptomatic or mildly symptomatic infection that are typically

associated with younger patients. The results of our infection studies indicate that colonization

of the lung with NTHi may provide inherent resistance to subsequent colonization with *Pa*. A

likely mechanism for such an effect would be stimulation of innate immune defenses, which can

prime host phagocyte responses to respond to incoming pathogens (31, 33, 37, 38). These

results raise interesting possibilities regarding whether aggressive treatment against early-stage

opportunists, such as NTHi, may be contraindicated or even harmful by providing a

susceptibility window to other infections. It is also notable that for opportunistic airway infections

associated with COPD, an extensive body of recent work indicates that treatment with

inflammatory agonists that evoke innate responses can confer protection and are under active

investigation as therapeutics (39-43). The concept of therapeutically inducing a low level of

airway inflammation to enhance resistance may merit further study in CF related infections.

MATERIALS AND METHODS

Bacterial strains and growth conditions

NTHi 86-028NP is well characterized patient isolate for which a fully annotated genome sequence data base is available (44). NTHi HI-1 is a sputum-derived isolate from a pediatric patient with CF and was provided by Timothy Starner (University of Iowa Children's Hospital). NTHi 86-028NPr and NTHi HI-1r were derived from NTHi 86-028NP and NTHi HI-1 by serial passage on plates containing spectinomycin (18). NTHi 86-028NP-gfp+ was constructed by electroporation with plasmid PGM1.1+gfp (provided by K. Mason, Nationwide Childrens' Research Hospital) (45, 46). For heat-killed NTHi experiments, bacteria were scraped from a plate into sterile PBS, resuspended to $\sim10^9$ CFU/ml, and incubated at 65°C for 1 h; bacterial killing was confirmed by lack of growth. All NTHi strains were routinely cultured on supplemented brain-heart infusion (sBHI) agar (Difco), containing 10 μg/mL of hemin (ICN Biochemical) and 1μg/mL of NAD (Sigma).

P. aeruginosa mPA08-31 is a mucoid clinical isolate from a CF patient at UAB Hospital, and was provided by S. Birket (University of Alabama at Birmingham) (47, 48). *A* nonmucoid derivative of *P. aeruginosa*, (*P. aeruginosa* FRD1*mucA*+) was provided by J. Scoffield (University of Alabama at Birmingham) (49). *P. aeruginosa* mPA08-31-mCherry+ was derived by electroporation with plasmid pUCP19+mCherry (provided by D. Wozniak, Ohio State University). All *P. aeruginosa* strains were routinely cultured on Luria-Bertani (LB) agar (Difco). Strains were streaked for colony isolation before inoculation into LB broth and shaking overnight at 37°C and 200 rpm.

Confocal microscopy of static biofilms

In vitro biofilms were prepared by resuspending NTHi 86028NP-gfp+ from an overnight plate culture to$\sim10^8$ CFU/ml in 2x sBHI, or overnight broth culture of mPA08-31-mCherry+ diluted to $\sim10^8$ CFU/ml in LB broth. Dual-species biofilms were sequentially seeded with NTHi 86028NP-

gfp+ preceding mPA08-31-mCherry+ by 12 hours into a 35-mm glass-bottom confocal dish

(MatTek) in 2-ml increments such that the final density in each well was ~10^8 CFU of one or of

both organisms. NTHi 86028NP-gfp+ single-species biofilms grew for 12 hours. mPA08-31-

mCherry+ single-species biofilms grew for 6 hours. Sequentially seeded dual biofilms grew for

18 hours. All dishes were incubated at 37°C with 5% CO_2 before being washed with sterile PBS

and fixed with 4% paraformaldehyde (Alfa Aesar, Tewksbury, MA). Viable colony counts of

mPA08-31-mCherry+ from single and dual-species biofilms were obtained by plating on LB

(antibiotic selection). Confocal laser scanning microscopy (CLSM) was performed using a

Nikon-A1R HD25 confocal laser microscope (Nikon, Tokyo, Japan) at the University of Alabama

at Birmingham. All images were processed using the NIS-elements 5.0 software.

Fluorescent biofilm quantification

Fluorescent biofilms were quantified using BiofilmQ software (50). Single fluorescence channels

were automatically segmented using the Otsu algorithm. Background was removed with semi-

manually thresholding denoising. Global biofilm parameters of the fluorescence channel

representing *Pa* were quantified for both single-species and dual-species biofilms to assess

biovolume. 3D spatial representation of cubing (representing 1 μm per cube) was calculated by

BiofilmQ and visualized with ParaView according to manufacturer's instructions.

Mouse model of respiratory infections

BALB/cJ mice (8 to 10 weeks old) were obtained from Jackson Laboratories (Bar Harbor, ME).

For single-species infections, mice were anesthetized with isoflurane and intratracheally

infected with NTHi (~10^9 CFU in 100 μl PBS) or *Pa* (~10^8 CFU in 100 μL PBS). For dual

infections, mice were anesthetized with isoflurane and infected intratracheally either

concurrently with NTHi and *Pa* (~10^9 CFU NTHi and ~10^8 CFU *Pa* in 100 μL PBS), or

sequentially, in which infection with NTHi occurred 24 hours prior to *Pa* introduction. For viable

plate counting, the left lung of each mouse was harvested and homogenized in 500 μL of sterile

PBS at 30 Hz/s. Lung homogenate was serially diluted in PBS and plated LB agar to obtain

viable colony counts of *Pa* and sBHI agar containing spectinomycin (2 mg/mL) for selection of

NTHi HI-1r. All samples from polymicrobial infections were also plated on sBHI without antibiotic

for total bacterial counts of both organisms. For histological analysis, the right lung of each

animal was inflated with 10% buffered formalin and stored at 4°C until processing. All mouse

infection protocols were approved by the University of Alabama at Birmingham (UAB)

Institutional Animal Care and Use Committees.

Bronchoalveolar lavage fluid collection

Bronchoalveolar lavage was performed by flushing lungs with 6 ml of cold PBS in 1-mL

increments as previously described (19). Collected BALF was stored on ice until processing and

the first 1mL isolated was centrifuged (1350 × *g*, 5 min) to separate the supernatant from

immune cells. The cell pellet was used for total and differential cell counts. The remaining 5 mL

of collected BALF was stored at -20°C until use for cytokine analyses.

Cytokine analyses and differential cell counting

Cytokine analyses were performed on the 5 mL of collected BALF via DuoSet ELISA kits (R&D

Systems, Minnesota, MN). For total and differential cell counts, cell pellets from BALF were

resuspended in fresh PBS and mounted on a slide by cytospin at 500 rpm for 5 minutes. Cells

were stained with a Kwik-Diff differential cell stain (ThermoFisher Scientific, Waltham, MA).

Three representative fields from each spot were counted to determine the composition of

immune cells.

Histological analysis

Inflated right lungs from infected animals were stored in 10% neutral buffered formalin (Fisher

Scientific, Waltham, MA) at 4°C until processing. Sections from each lobe of the right lung were

trimmed and sent to the UAB Comparative Pathology Laboratory to be paraffin embedded,

sectioned and stained with hematoxylin and eosin (H&E). Images of lung sections (200 x

magnification) were taken with Olympus DP25 camera (Tokyo, Japan) using an Olympus BX40

microscope. Semiquantitative grading of all lung sections was performed by a board-certified

surgical pathologist (L.N.). Semiquantitative histopathological scores were assigned using a

scoring matrix based primarily on neutrophilic influx. Severity was rated on a scale of 0 to 3,

where 0 represents no observable neutrophils,1 represents dispersed acute inflammation (mild

damage), 2 represents dense infiltration (moderate damage), and 3 represents solid infiltrate

with necrosis (severe damage). of the section affected of independently viewed fields of view.

The final histopathological score represents the maximum grade assigned to each replicate.

Statistical analyses

All bar graphs represent sample means ± standard error of the mean (SEM). All mouse

experiments were repeated at least two times and data from independent experiments were

combined for analyses. For nonparametric analyses, differences between groups were analyzed

by Kruskal-Wallis test with the uncorrected Dunn's test for multiple comparisons. For normally

distributed data sets, as determined by Shapiro-Wilk normality test, a one-way analysis of

variance (ANOVA) was used with Tukey's multiple-comparison test. Outliers were detected via

the ROUT method (Q, 1%) and excluded from the analysis. All statistical tests were performed

using GraphPad Prism 9 (San Diego, CA).

ACKNOWLEDGMENTS

The authors gratefully acknowledge helpful input and discussions with colleagues in the UAB

Center for Cystic Fibrosis Research. This work was supported by grants to W.E.S. from the

Cystic Fibrosis Foundation (CFFSWORDS1810 and CFFSWORDS20G0). N.L. was a

predoctoral trainee supported by the UAB Cystic Fibrosis Foundation Basic Research Center

(Steven Rowe, PI). B.C.H. was supported by a fellowship from the UAB Predoctoral Training

Program in Lung Biology (T32 HL13640). We also thank Timothy Starner (University of Iowa

Children's Hospital), Daniel Wozniak (Ohio State University), Jessica Scoffield (UAB), and

Susan Birket (UAB) for providing the bacterial strains for this study.

Figure Legends

Fig. 1: Nontypeable _H. influenzae_ and _P. aeruginosa_ form polymicrobial biofilms _in vitro_.

Sequentially seeded biofilms on abiotic surface of A) single-species _P. aeruginosa_ mPA 08-31 (pUCP19+mCherry+) B) Nontypeable _H. influenzae_ 86-028NP (PGM1.1+gfp) and C) dual-species NTHi and _Pa_ in both single channels and merged. Inoculation of NTHi preceded _Pa_ by 12 hours and both organisms grew for additional 6 hours at 37 °C + 5% CO_2 (18 hours total). Imaged at 40X magnification. Scale bar represents 50 μm.

Fig. 2 Quantification of nontypeable _H. influenzae_ and _P. aeruginosa_ polymicrobial biofilms.

Bacterial density within fluorescent biofilms was quantified using BiofilmQ software to measure A) biofilm volume of _Pa_ in dual and single-species biofilms and B) mean cube surface of _Pa_ quantified as 1 cube = 1μm. Cubing can be visualized as local density of _Pa_ in both C) a sequentially seeded dual-species biofilm and D) a single-species _Pa_ biofilm. Mean ± SEM. (N = 4). Mann-Whitney test, two-tailed, *P<0.05.

Fig. 3: Nontypeable H. influenzae diminishes _P. aeruginosa_ abundance in murine lungs.

BALBc/J mice were intratracheally infected with nontypeable _H. influenzae_ (NTHi Hi-1r) either 24 hours prior to introduction of mucoid _P. aeruginosa_ (mPA 08-31) in a sequential fashion, or were introduced at the same time (concurrently) and assessed 48 post-infection. A) Bacterial counts of NTHi and _Pa_ in the lung homogenate enumerated by viable colony counting. B) Bacterial counts of _Pa_ in the lung homogenate comparing sequential and concurrent introduction. Severity of disease was indicated by C) H&E stained lung sections (200x

magnification) and D) semi-quantitative grading by a pathologist (L.N.). E) Total cell counts in

BALF quantified by differential cell counts. F) Percentage of PMNs in the BALF. Mean ± SEM. (N

= 10). LOD = limit of detection. Kruskal-Wallis with Dunn's multiple comparison test, * P<0.05,

** P<0.01, *** P<0.001, **** P<0.0001.

Fig. 4: Nontypeable *H. influenzae* impact on subsequent *P. aeruginosa* infection is strain

independent.

BALBc/J mice were intratracheally infected with nontypeable *H. influenzae* (NTHi 86r) followed

by mucoid *P. aeruginosa* (mPA 08-31). A) Bacterial counts in the lung homogenate were

enumerated by viable colony counting 72 hours post- dual infection. Intratracheal infections

were also performed with nontypeable *H. influenzae* (NTHi HI-1r) and nonmucoid *P. aeruginosa*

(PA FRD1*mucA+)* and B) bacterial counts in the lung homogenate were enumerated by viable

colony counting 72 hours post- dual infection. Mean ± SEM. (N = 10). LOD = limit of detection.

Kruskal-Wallis with Dunn's multiple comparison test, * P<0.05, ** P<0.01, *** P<0.001, ****

P<0.0001.

Fig. 5: Nontypeable *H. influenzae* and *P. aeruginosa* dual-infection reduces the severity of

airway tissue damage.

BALBc/J mice were intratracheally infected with nontypeable *H. influenzae* (NTHi Hi-1r) 24

hours prior to introduction of mucoid *P. aeruginosa* (mPA 08-31) and assessed 72 hours post-

dual infection. A) Bacterial counts of NTHi and *Pa* in the lung homogenate enumerated by

viable colony counting. (N = 13-15). Severity of disease was indicated by B) semi-quantitative

grading by a pathologist (L.N.) and C) H&E stained lung sections (200x magnification) (N = 6-10).

D) Total cell counts in BALF quantified by differential counting. (N = 6-8). E) Percentage of PMNs

in the BALF (N = 6-9). Mean ± SEM. LOD = limit of detection. Kruskal-Wallis with Dunn's multiple

comparison test, * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Fig. 6: Inflammatory cytokine levels after dual-infection are reduced compared to *P.*

***aeruginosa* infection alone.**

Cytokine and chemokine analyses were performed on BALF supernatant from infected or mock-

infected mice 72 hours post-infection using a DuoSet ELISA kits for A) MCP-1, B) TNF-α, C) IL-10,

D) IL-1α, E) IL-1β, and F) IL-6. Mean ± SEM, (N = 7-10). One-way ANOVA with Tukey's multiple

comparisons test for post-hoc analysis. **** P<0.0001.

Fig. 7: Impact of nontypeable *H. influenzae* on *P. aeruginosa* infection is independent of

bacterial viability.

BALBc/J mice were intratracheally infected with heat-inactivated nontypeable *H. influenzae*

(NTHi Hi-1r) 24 hours prior to introduction of mucoid *P. aeruginosa* (mPA 08-31) and assessed

72 hours post- dual infection. A) Bacterial counts of NTHi and *Pa* in the lung homogenate were

enumerated by viable colony counting. (N = 10). Severity of disease was indicated by B) semi-

quantitative grading by a pathologist (L.N.) and C) H&E stained lung sections (200x

magnification) (N = 8-10). D) Total cell counts in BALF quantified by differential counting. E)

Percentage of PMNs in the BALF. (N = 9-10). Mean ± SEM. LOD = limit of detection. Kruskal-

Wallis with Dunn's multiple comparison test, * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Fig. S1: Quantification of nontypeable *H. influenzae* and *P. aeruginosa* polymicrobial

fluorescent biofilms.

Viable colony counts of A) single -species NTHI 86-028NP-gfp+ and mPA 08-31-mCherry+ static

biofilms seeded at ~10^8 and grown over time. B) Bacterial counts of single- and dual-species

static biofilms sequentially seeded at ~10^8 and grown in sBHI broth at 37°C + 5% CO_2 for 18

hours total. Dual counts depicts Pa growth. Mean ± SEM. (N = 4-5). Dotted line = inoculum.

Kruskal-Wallis with Dunn's multiple comparison test.

Fig. S2: Weight change following nontypeable *H. influenzae* and *P. aeruginosa* infection.

Percentage of initial weight was calculated for all animal infections with weight change

compared to uninfected controls. A) Weight change at 48 hours post-infection with NTHi HI-1r

and mPA 08-31. B) Weight change at 72 hours post-infection with NTHi 86-028NPr and mPA 08-

31. C) Weight change at 72 hours post-infection with NTHi HI-1r and Pa FRD1*mucA+*. D) Weight

change at 72 hours post-infection with NTHi HI-1r and mPA 08-31. E) Weigh change at 72 hours

post-infection with heat-killed NTHi HI-1r and mPA 08-31. (N = 10-15). Mean ± S1

SEM. Kruskal-Wallis with Dunn's multiple comparison test, * P<0.05, ** P<0.01, *** P<0.001,

**** P<0.0001.

References:

1.   Lyczak JB, Cannon CL, Pier GB. 2002. Lung infections associated with cystic fibrosis. Clin
Microbiol Rev 15:194-222.

2.   Ratjen F, Bell SC, Rowe SM, Goss CH, Quittner AL, Bush A. 2015. Cystic fibrosis. Nat Rev
Dis Primers 1:15010.

3.   Magalhaes AP, Lopes SP, Pereira MO. 2016. Insights into Cystic Fibrosis Polymicrobial
Consortia: The Role of Species Interactions in Biofilm Development, Phenotype, and
Response to In-Use Antibiotics. Front Microbiol 7:2146.

4.   Foundation CF. 2019. Cystic fibrosis patient registry annual data report.

5.   O'Toole GA. 2018. Cystic Fibrosis Airway Microbiome: Overturning the Old, Opening the
Way for the New. J Bacteriol 200.

6.   Huang YJ, LiPuma JJ. 2016. The Microbiome in Cystic Fibrosis. Clin Chest Med 37:59-67.

7.   Caverly LJ, Zhao J, LiPuma JJ. 2015. Cystic fibrosis lung microbiome: opportunities to
reconsider management of airway infection. Pediatr Pulmonol 50 Suppl 40:S31-8.

8.   Filkins LM, O'Toole GA. 2015. Cystic Fibrosis Lung Infections: Polymicrobial, Complex,
and Hard to Treat. PLoS Pathog 11:e1005258.

9.   Zemanick ET, Sagel SD, Harris JK. 2011. The airway microbiome in cystic fibrosis and
implications for treatment. Curr Opin Pediatr 23:319-24.

10.   Erwin AL, Smith AL. 2007. Nontypeable *Haemophilus influenzae*: understanding
virulence and commensal behavior. **Trends Microbiol** 15:355-362.

11.   Dickson RP, Erb-Downward JR, Martinez FJ, Huffnagle GB. 2016. The Microbiome and
the Respiratory Tract. Annu Rev Physiol 78:481-504.

12.   Weinberg F, Dickson RP, Nagrath D, Ramnath N. 2020. The Lung Microbiome: A Central
Mediator of Host Inflammation and Metabolism in Lung Cancer Patients? Cancers
(Basel) 13.

13.   Cardines R, Giufre M, Pompilio A, Fiscarelli E, Ricciotti G, Di Bonaventura G, Cerquetti M.
2012. Haemophilus influenzae in children with cystic fibrosis: antimicrobial
susceptibility, molecular epidemiology, distribution of adhesins and biofilm formation.
Int J Med Microbiol 302:45-52.

14.   Rayner RJ, Hiller EJ, Ispahani P, Baker M. 1990. Haemophilus infection in cystic fibrosis.
559   Arch Dis Child 65:255-8.

560   15.   Saliu F, Rizzo G, Bragonzi A, Cariani L, Cirillo DM, Colombo C, Dacco V, Girelli D, Rizzetto
561   S, Sipione B, Cigana C, Lore NI. 2021. Chronic infection by nontypeable Haemophilus
562   influenzae fuels airway inflammation. ERJ Open Res 7.

563   16.   Starner TD, Zhang N, Kim G, Apicella MA, McCray PB, Jr. 2006. *Haemophilus influenzae*
564   forms biofilms on airway epithelia: implications in cystic fibrosis. **Am J Respir Crit Care
565   Med** 174:213-220.

566   17.   Swords WE. 2012. Nontypeable Haemophilus influenzae biofilms: role in chronic airway
567   infections. Front Cell Infect Microbiol 2:97.

568   18.   Grahl N, Dolben EL, Filkins LM, Crocker AW, Willger SD, Morrison HG, Sogin ML, Ashare
569   A, Gifford AH, Jacobs NJ, Schwartzman JD, Hogan DA. 2018. Profiling of Bacterial and
570   Fungal Microbial Communities in Cystic Fibrosis Sputum Using RNA. mSphere 3.

571   19.   Burns JL, Gibson RL, McNamara S, Yim D, Emerson J, Rosenfeld M, Hiatt P, McCoy K,
572   Castile R, Smith AL, Ramsey BW. 2001. Longitudinal assessment of Pseudomonas
573   aeruginosa in young children with cystic fibrosis. J Infect Dis 183:444-52.

574   20.   Gellatly SL, Hancock RE. 2013. Pseudomonas aeruginosa: new insights into pathogenesis
575   and host defenses. Pathog Dis 67:159-73.

576   21.   Rossi E, La Rosa R, Bartell JA, Marvig RL, Haagensen JAJ, Sommer LM, Molin S, Johansen
577   HK. 2021. Pseudomonas aeruginosa adaptation and evolution in patients with cystic
578   fibrosis. Nat Rev Microbiol 19:331-342.

579   22.   Gill JF, Deretic V, Chakrabarty AM. 1987. Alginate production by the mucoid
580   Pseudomonas aeruginosa associated with cystic fibrosis. Microbiol Sci 4:296-9.

581   23.   Mayer-Hamblett N, Aitken ML, Accurso FJ, Kronmal RA, Konstan MW, Burns JL, Sagel SD,
582   Ramsey BW. 2007. Association between pulmonary function and sputum biomarkers in
583   cystic fibrosis. Am J Respir Crit Care Med 175:822-8.

584   24.   Burns JL, Emerson J, Stapp JR, Yim DL, Krzewinski J, Louden L, Ramsey BW, Clausen CR.
585   1998. Microbiology of sputum from patients at cystic fibrosis centers in the United
586   States. Clin Infect Dis 27:158-63.

587   25.   Dickson RP, Erb-Downward JR, Huffnagle GB. 2013. The role of the bacterial microbiome
588   in lung disease. Expert Rev Respir Med 7:245-57.

26.　Dickson RP, Erb-Downward JR, Prescott HC, Martinez FJ, Curtis JL, Lama VN, Huffnagle
GB. 2014. Cell-associated bacteria in the human lung microbiome. Microbiome 2:28.

27.　Dickson RP, Huffnagle GB. 2015. The Lung Microbiome: New Principles for Respiratory
Bacteriology in Health and Disease. PLoS Pathog 11:e1004923.

28.　Welp AL, Bomberger JM. 2020. Bacterial Community Interactions During Chronic
Respiratory Disease. Front Cell Infect Microbiol 10:213.

29.　Burns JL, Ramsey BW, Smith AL. 1993. Clinical manifestations and treatment of
pulmonary infections in cystic fibrosis. Adv Pediatr Infect Dis 8:53-66.

30.　Lysenko ES, Clarke TB, Shchepetov M, Ratner AJ, Roper DI, Dowson CG, Weiser JN. 2007.
Nod1 signaling overcomes resistance of S. pneumoniae to opsonophagocytic killing.
PLoS Pathog 3:e118.

31.　Lysenko ES, Ratner AJ, Nelson AL, Weiser JN. 2005. The role of innate immune responses
in the outcome of interspecies competition for colonization of mucosal surfaces. **PLoS**
Pathog 1:e1.

32.　Armbruster CE, Hong W, Pang B, Weimer KE, Juneau RA, Turner J, Swords WE. 2010.
Indirect pathogenicity of Haemophilus influenzae and Moraxella catarrhalis in
polymicrobial otitis media occurs via interspecies quorum signaling. mBio 1.

33.　Weimer KE, Armbruster CE, Juneau RA, Hong W, Pang B, Swords WE. 2010. Coinfection
with Haemophilus influenzae promotes pneumococcal biofilm formation during
experimental otitis media and impedes the progression of pneumococcal disease. J
Infect Dis 202:1068-75.

34.　Hampton TH, Green DM, Cutting GR, Morrison HG, Sogin ML, Gifford AH, Stanton BA,
O'Toole GA. 2014. The microbiome in pediatric cystic fibrosis patients: the role of shared
environment suggests a window of intervention. Microbiome 2:14.

35.　Zemanick ET, Harris JK, Wagner BD, Robertson CE, Sagel SD, Stevens MJ, Accurso FJ,
Laguna TA. 2013. Inflammation and airway microbiota during cystic fibrosis pulmonary
exacerbations. PLoS One 8:e62917.

36.　Starner TD, McCray PB, Jr., American College of P, American Physiological S. 2005.
Pathogenesis of early lung disease in cystic fibrosis: a window of opportunity to
eradicate bacteria. Ann Intern Med 143:816-22.

37.　Ratner AJ, Lysenko ES, Paul MN, Weiser JN. 2005. Synergistic proinflammatory
responses induced by polymicrobial colonization of epithelial surfaces. **Proc Natl Acad**
Sci U S A 102:3429-3434.

38.   Mukherjee S, Weimer KE, Seok SC, Ray WC, Jayaprakash C, Vieland VJ, Swords WE, Das J.
2014. Host-to-host variation of ecological interactions in polymicrobial infections. Phys
Biol 12:016003.

39.   Cleaver JO, You D, Michaud DR, Pruneda FA, Juarez MM, Zhang J, Weill PM, Adachi R,
Gong L, Moghaddam SJ, Poynter ME, Tuvim MJ, Evans SE. 2014. Lung epithelial cells are
essential effectors of inducible resistance to pneumonia. Mucosal Immunol 7:78-88.

40.   Duggan JM, You D, Cleaver JO, Larson DT, Garza RJ, Guzman Pruneda FA, Tuvim MJ,
Zhang J, Dickey BF, Evans SE. 2011. Synergistic interactions of TLR2/6 and TLR9 induce a
high level of resistance to lung infection in mice. J Immunol 186:5916-26.

41.   Evans SE, Tuvim MJ, Fox CJ, Sachdev N, Gibiansky L, Dickey BF. 2011. Inhaled innate
immune ligands to prevent pneumonia. Br J Pharmacol 163:195-206.

42.   Evans SE, Xu Y, Tuvim MJ, Dickey BF. 2010. Inducible innate resistance of lung
epithelium to infection. Annu Rev Physiol 72:413-35.

43.   Moghaddam SJ, Clement CG, De la Garza MM, Zou X, Travis EL, Young HW, Evans CM,
Tuvim MJ, Dickey BF. 2008. Haemophilus influenzae lysate induces aspects of the
chronic obstructive pulmonary disease phenotype. Am J Respir Cell Mol Biol 38:629-38.

44.   Harrison A, Dyer DW, Gillaspy A, Ray WC, Mungur R, Carson MB, Zhong H, Gipson J,
Gipson M, Johnson LS, Lewis L, Bakaletz LO, Munson RS, Jr. 2005. Genomic sequence of
an otitis media isolate of nontypeable Haemophilus influenzae: comparative study with
H. influenzae serotype d, strain KW20. J Bacteriol 187:4627-36.

45.   Vogel AR, Szelestey BR, Raffel FK, Sharpe SW, Gearinger RL, Justice SS, Mason KM. SapF-
mediated heme-iron utilization enhances persistence and coordinates biofilm
architecture of Haemophilus. Front Cell Infect Microbiol 2:42.

46.   Mitchell MA, Skowronek K, Kauc L, Goodgal SH. 1991. Electroporation of *Haemophilus*
influenzae is effective for transformation of plasmid but not chromosomal DNA. Nucl
Acids Res 19:3625-3628.

47.   Das S, Howlader DR, Zheng Q, Ratnakaram SSK, Whittier SK, Lu T, Keith JD, Picking WD,
Birket SE, Picking WL. 2020. Development of a Broadly Protective, Self-Adjuvanting
Subunit Vaccine to Prevent Infections by Pseudomonas aeruginosa. Front Immunol
11:583008.

48.   McDaniel MS, Schoeb T, Swords WE. 2020. Cooperativity between Stenotrophomonas
maltophilia and Pseudomonas aeruginosa during polymicrobial airway infections. Infect
Immun doi:10.1128/IAI.00855-19.

49.   Silo-Suh LA, Suh SJ, Ohman DE, Wozniak DJ, Pridgeon JW. 2015. Complete Genome
656   Sequence of Pseudomonas aeruginosa Mucoid Strain FRD1, Isolated from a Cystic
657   Fibrosis Patient. Genome Announc 3.

658   50.   Hartmann R, Jeckel H, Jelli E, Singh PK, Vaidya S, Bayer M, Rode DKH, Vidakovic L, Diaz-
659   Pascual F, Fong JCN, Dragos A, Lamprecht O, Thoming JG, Netter N, Haussler S, Nadell
660   CD, Sourjik V, Kovacs AT, Yildiz FH, Drescher K. 2021. Quantitative image analysis of
661   microbial communities with BiofilmQ. Nat Microbiol 6:151-156.

Fig. 1: Nontypeable *H. influenzae* and *P. aeruginosa* form polymicrobial biofilms *in vitro*.
Sequentially seeded biofilms on abiotic surface of A) single-species *P. aeruginosa* mPA 08-31 (pUCP19+mCherry) B) Nontypeable *H. influenzae* 86-028NP (PGM1.1+gfp) and C) dual-species NTHi and *Pa* in both single channels and merged. Inoculation of NTHi preceded *Pa* by 12 hours and both organisms grew for an additional 6 hours at 37 °C + 5% CO_2 (18 hours total). Imaged at 40X magnification. Scale bar represents 50 μm.

Fig. 2: Quantification of nontypeable *H. influenzae* and *P. aeruginosa* polymicrobial biofilms.
Bacterial density within fluorescent biofilms was quantified using BiofilmQ software to measure A) biofilm volume of mPA 08-31(mCherry+) in dual and single-species biofilms and B) mean cube surface of *Pa* quantified as 1 cube = 1µm. Cubing can be visualized as local density of *Pa* in both C) a sequentially seeded dual-species biofilm and D) a single-species *Pa* biofilm. Mean ± SEM. (N = 4). Mann-Whitney test, two-tailed, *P<0.05.

Fig. 3: Nontypeable *H. influenzae* diminishes *P. aeruginosa* abundance in murine lungs.
BALBc/J mice were intratracheally infected with nontypeable *H. influenzae* (NTHi Hi-1r) either 24 hours prior to introduction of mucoid *P. aeruginosa* (mPA 08-31) in a sequential fashion, or were introduced at the same time (concurrently) and assessed 48 post-infection. A) Bacterial counts of NTHi and *Pa* in the lung homogenate enumerated by viable colony counting. B) Bacterial counts of *Pa* in the lung homogenate comparing sequential and concurrent introduction. Severity of disease was indicated by C) H&E stained lung sections (200x magnification) and D) semi-quantitative grading by a pathologist (L.N.). E) Total cell counts in BALF quantified by differential cell counts. F) Percentage of PMNs in the BALF. Mean ± SEM. (N = 10). LOD = limit of detection. Kruskal-Wallis with Dunn's multiple comparison test, * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Fig. 4: Nontypeable *H. influenzae* impact on subsequent *P. aeruginosa* infection is strain independent.
BALBc/J mice were intratracheally infected with nontypeable *H. influenzae* (NTHi 86r) followed by mucoid *P. aeruginosa* (mPA 08-31). A) Bacterial counts in the lung homogenate were enumerated by viable colony counting 72 hours post- dual infection. Intratracheal infections were also performed with nontypeable *H. influenzae* (NTHi HI-1r) and nonmucoid *P. aeruginosa* (PA FRD1*mucA+*) and B) bacterial counts in the lung homogenate were enumerated by viable colony counting 72 hours post- dual infection. Mean ± SEM. (N = 10). LOD = limit of detection. Kruskal-Wallis with Dunn's multiple comparison test, * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Fig. 5: Nontypeable *H. influenzae* and *P. aeruginosa* dual-infection reduces the severity of airway tissue damage.

BALBc/J mice were intratracheally infected with nontypeable *H. influenzae* (NTHi Hi-1r) 24 hours prior to introduction of mucoid *P. aeruginosa* (mPA 08-31) and assessed 72 hours post- dual infection. A) Bacterial counts of NTHi and *Pa* in the lung homogenate enumerated by viable colony counting. (N = 13-15). Severity of disease was indicated by B) semi-quantitative grading by a pathologist (L.N.) and C) H&E stained lung sections (200x magnification). (N = 6-10). D) Total cell counts in BALF quantified by differential counting. (N = 6-8). E) Percentage of PMNs in the BALF (N = 6-9). Mean ± SEM. LOD = limit of detection. Kruskal-Wallis with Dunn's multiple comparison test, * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Fig. 6: Inflammatory cytokine levels after dual-infection are decreased compared to *P. aeruginosa* infection alone.
Cytokine and chemokine analyses were performed on BALF supernatant from infected or mock-infected mice 72 hours post-infection using a DuoSet ELISA kits for A) MCP-1, B) TNF-α, C) IL-10, D) IL-1α, E) IL-1β, and F) IL-6 Mean ± SEM, (N = 7-10). One-way ANOVA with Tukey's multiple comparisons test for post-hoc analysis.
**** P<0.0001.

Fig. 7: Impact of nontypeable *H. influenzae* on *P. aeruginosa* infection is independent of bacterial viability. BALBc/J mice were intratracheally infected with heat-inactivated nontypeable *H. influenzae* (NTHi Hi-1r) 24 hours prior to introduction of mucoid *P. aeruginosa* (mPA 08-31) and assessed 72 hours post- dual infection. A) Bacterial counts of NTHi and *Pa* in the lung homogenate were enumerated by viable colony counting. (N = 10). Severity of disease was indicated by B) semi-quantitative grading by a pathologist (L.N.) and C) H&E stained lung sections (200x magnification). (N = 8-10). D) Total cell counts in BALF quantified by differential counting. E) Percentage of PMNs in the BALF. (N = 9-10). Mean ± SEM. LOD = limit of detection. Kruskal-Wallis with Dunn's multiple comparison test, * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

A.

Viable Biofilm Counts Over Time

- ■ NTHi 86-028NP (gfp+)
- ● mPA 08-31 (mCherry+)

B.

Polymicrobial Biofilm Counts (18hr)

ns

Fig. S1: Quantification of nontypeable *H. influenzae* and *P. aeruginosa* polymicrobial fluorescent biofilms.
Viable colony counts of A) single -species NTHI 86-028NP-gfp+ and mPA 08-31-mCherry+ static biofilms seeded at ~10^8 and grown over time. B) Bacterial counts of single- and dual-species static biofilms sequentially seeded at ~10^8 and grown in sBHI broth at 37°C + 5% CO_2 for 18 hours total. Dual counts depicts Pa growth. Mean ± SEM. (N = 4-5). Dotted line = inoculum. Kruskal-Wallis with Dunn's multiple comparison test.

Fig. S2: Weight change following nontypeable *H. influenzae* and *P. aeruginosa* infection.
Percentage of initial weight was calculated for all animal infections with weight change compared to uninfected controls. A) Weight change at 48 hours post-infection with NTHi HI-1r and mPA 08-31. B) Weight change at 72 hours post-infection with NTHi 86-028NPr and mPA 08-31. C) Weight change at 72 hours post-infection with NTHi HI-1r and *Pa* FRD1*mucA+*. D) Weight change at 72 hours post-infection with NTHi HI-1r and mPA 08-31. E) Weigh change at 72 hours post-infection with heat-killed NTHi HI-1r and mPA 08-31. (N = 10-15). Mean ± SEM. Kruskal-Wallis with Dunn's multiple comparison test, * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.