Disc and Actin-Associated Protein 1 Influence Attachment in the Intestinal Parasite

Giardia lamblia

Melissa C. Steele-Ogus[1], Ava M. Obenaus[2], Nathan J. Sniadecki[2], and Alexander R. Paredez[1]

[1]Department of Biology, University of Washington, Seattle, United States

[2]Department of Mechanical Engineering, University of Washington, Seattle, United States

Abstract

The deep-branching eukaryote *Giardia lamblia* is an extracellular parasite that attaches to

the host intestine via a microtubule-based structure called the ventral disc. Control of attachment

is mediated in part by the movement of two regions of the ventral disc that either permit or

exclude the passage of fluid under the disc. Several known disc-associated proteins (DAPs)

contribute to disc structure and function, but no force-generating protein has been identified

among them. We recently identified several *Giardia* actin (*Gl*Actin) interacting proteins at the

ventral disc, which could potentially employ actin polymerization for force generation and disc

conformational changes. One of these proteins, Disc and Actin Associated Protein 1 (DAAP1), is

highly enriched at the two regions of the disc previously shown to be important for fluid flow

during attachment. In this study, we investigate the role of both *Gl*Actin and DAAP1 in ventral

disc morphology and function. We confirmed interaction between *Gl*Actin and DAAP1 through

coimmunoprecipitation, and used immunofluorescence to localize both proteins throughout the

cell cycle and during trophozoite attachment. Similar to other DAPs, the association of DAAP1

with the disc is stable, except during cell division when the disc disassembles. Depletion of

*Gl*Actin by translation-blocking antisense morpholinos resulted in both impaired attachment and

defects in the ventral disc, indicating that *Gl*Actin contributes to disc-mediated attachment.

Depletion of DAAP1 through CRISPR interference resulted in intact discs but impaired

attachment, gating, and flow under the disc. As attachment is essential for infection, elucidation

of these and other molecular mediators is a promising area for development of new therapeutics

against a ubiquitous parasite.

Author Summary

Giardia lamblia is a single-celled organism and one of the most common gastrointestinal

parasites worldwide. In developing countries, recurrent *Giardia* infections are common, due to

lack of access to clean water. *Giardia* infections can lead to diarrhea, vomiting, dehydration,

disruption of the intestinal microbiome, and chronic infections can lead to irritable bowel

syndrome. Because existing drug treatments have side effects and *Giardia's* resistance to drugs

is increasing, new treatment strategies are needed. The parasite's attachment to the host's

intestine is mediated by a *Giardia*-specific structure that resembles a suction cup and is called

the ventral adhesive disc. We previously identified DAAP1, a protein which interacts with

Giardia actin and localizes to the ventral disc. Here, we explore the relationship between these

two proteins and investigate their role in disc-based attachment. Most disc proteins, including

DAAP1, are unrelated to any human proteins, making them appealing drug targets to inhibit

parasite attachment and infection.

Introduction

Giardia lamblia, a single-celled parasite, is the cause of over 280 million annual cases of

the gastrointestinal disease giardiasis worldwide [1,2] This extracellular parasite colonizes the

lumen of the intestine, where its ability to maintain infection requires tight attachment to

microvilli, in order to withstand the peristaltic flow of the gastrointestinal tract [3]. The primary

means of attachment is via a specialized microtubule-based organelle known as the ventral

adhesive disc [4–6]. A well-functioning disc is critical for cell survival and disease transmission

[7], but the molecular mechanisms controlling conformational dynamics and function remain

poorly understood.

Imaging of fluid flow at the disc margin identified two areas postulated to be important

for tight attachment: one at the anterior and the other at the posterior of the disc [8–10] (Figure

1A). The former is the marginal region of the overlap zone, the region of the disc where the sheet

of microtubules overlaps itself. The other is the ventral groove, a hump in the posterior region of

the disc where the ventral flagella exit between the disc and cell body. Interruption of contact in

these areas results in disruption of attachment and in fluid moving underneath the disc [9,11,12].

Cryo-electron tomography imaging of the disc reveals different protein densities in these areas,

indicating specialized architecture within the disc itself, which may indicate that specific effector

proteins control disc conformation and fluid flow [13]. In the current model, both hydrodynamic

flow and suction contribute to the mechanism of attachment [9]. Attachment occurs in several

distinct stages [8,9]. In early attachment, the anterior ventrolateral flange, lateral crest and

ventral groove make contact with the surface while the ventral flagella beat to generate fluid

flows. During this stage cells can either skim the surface or remain stationary. In the later stages

of attachment, the cell body becomes more intimately associated with the surface so that the

lateral shield and bare area make contact with the surface and the cells remain in place. Flagella-

beating is required for initiation, but not maintenance, of attachment, indicating that fluid must

be moved out from underneath the disc during the earlier stages of attachment [9,14].

These conformational changes in the disc which control attachment are not generated by

microtubule turnover, as microtubule depolymerizing or stabilizing drugs do not affect

attachment [15]. Though microtubules are the foundation of the disc, it is also composed of over

100 other proteins, known as disc-associated proteins (DAPs). At least five of these DAPs have

been shown to be important for disc biogenesis and function [9,12]. Only three DAPs identified

to date contain known microtubule-binding motifs; most are ankyrin-repeat proteins, NEK

kinases, or the annexin-like alpha-Giardins [15,16]. At least 25, however, lack identifiable

domains and are annotated in the *Giardia* genome as "hypothetical proteins" [17,18]. So far, no

DAPs have been identified as motor or force-generating proteins. Thus, the proteins that mediate

changes in disc conformation are unknown.

Actin has a conserved role in force generation and in controlling cell and organelle shape

in eukaryotic cells[19]. Previously, localization studies using heterologous antibodies

demonstrated that *Giardia* actin (*Gl*Actin) and actin regulators associate with the ventral disc

[6,20]. This observation led to multiple experiments attempting to test the role of *Gl*Actin in

attachment using actin inhibitors. These studies yielded conflicting results because reagents that

typically recognize actin or actin-binding proteins in other organisms were presumed to act

similarly in *Giardia* [6]. Later, *Gl*Actin was determined to be one of the most divergent actins

found among eukaryotes and its genome lacked genes encoding conventional actin-binding

proteins that were reportedly localized to the disc [21–23]. Further research revealed that

*Gl*Actin has non-conservative amino acid substitutions, predicted to disrupt actin inhibitor

binding, and anti-*Gl*Actin antibodies revealed localization throughout the cell [22]. Thus,

*Gl*Actin's potential role in disc-based attachment is controversial.

We recently identified eight *Giardia* actin (*Gl*Actin) interactors that localize to the

ventral disc [18], suggesting a means to resolve the role of actin and actin-binding proteins in

ventral disc function. One such actin-interacting protein, GL50803_16844, which we denote

Disc and Actin Associated Protein 1 (DAAP1), appears highly enriched in the two regions of the

disc important for regulating fluid flow during attachment.

In this study, we use morpholinos and CRISPR interference (CRISPRi) to deplete

*Gl*Actin and DAAP1 respectively and investigate the role of both proteins in the ventral disc.

Our results show that *Gl*Actin plays a role in normal ventral disc morphology and both *Gl*Actin

and DAAP1 are required for parasite attachment.

Results

***Gl*Actin affects attachment and disc morphology**

In vitro, Giardia normally grows attached to the inside of a culture tube. We used an

antisense translation-blocking morpholino to deplete *Gl*Actin for 24 hours, [22] after which we

assessed their attachment. We compared the numbers of cells attached to the tube wall to cells

unattached in the media for each condition. With an approximately 50% knockdown (Figure S1),

we found attachment to plastic culture tube walls to be significantly decreased, with an average

of 23% of cells attached compared to 75% of cells treated with a control morpholino (Figure

1B). Dead *Giardia* cells lose their ability to adhere to culture tubes. To account for the possibility

that the majority of the unattached cells were dead, we stained the unattached cells with

propidium iodide to mark dead cells, and fluorescein diacetate to mark live cells. While slightly

more unattached *Gl*Actin knockdown cells were dead compared to that of the control population,

this difference was not statistically significant (Figure 1C), indicating that the defect in

attachment in *Gl*Actin-depleted cells was not due to a preponderance of dead cells.

We asked if the role of *Gl*Actin in attachment could be attributed to an effect on the

ventral disc. Attached and unattached cells were treated with either control or anti-*Gl*Actin

morpholino, fixed after 24 h, and examined for distribution of a ventral disc marker using

immunofluorescence microscopy. We used a HALO-tagged delta-giardin which is widely

distributed in the disc to monitor whether the disc appeared normal or defective. Defects were

then further classified as 1) mild, with misshapen/broken discs or small gaps in the disc or 2)

severe, including unwound discs or severely misshapen discs (Figure S2, 1D, 1E). More *Gl*Actin

knockdown cells had defective discs compared to those in the control; however, this difference

was only significant in unattached cells (Figure 1D, 1E). While both attached and unattached

*Gl*Actin-depleted cells had more disc defects than their counterparts treated with the control

morpholino, the unattached *Gl*Actin-depleted cells had the most defective discs among all the

categories. Yet, there were more unattached cells than cells with disc defects, suggesting that

*Gl*Actin has a role in attachment beyond its contribution to disc morphology.

GlActin interacts with disc-associated protein DAAP1

We next asked if GL50803_16844, previously identified by mass spectrometry as a

potential actin filament interactor at the ventral disc [18], could be involved in attachment.

GL50803_16844 is conserved in all *Giardia lamblia* isolates and *Giardia muris*, but is absent

from the genome of *Spironucleus*, a close relative of *Giardia* which lacks a ventral disc [24].

Hereafter, we will refer to this protein that lacks identifiable domains and homologues in other

organisms as Disc- and Actin-Associated Protein 1 (DAAP1). We tagged DAAP1 with a C-

terminal 3xHA tag and expressed it episomally in *Giardia* (Figure S2). Immunoprecipitation of

this construct revealed it in complex with *Gl*Actin (Figure 2A). We also endogenously tagged

DAAP1 with a C-terminal mNeonGreen tag then used live and fixed cell fluorescent imaging.

Notably, this fusion protein localizes to the entirety of the disc, but it is highly enriched in the

ventral groove and the overlap zone, two areas implicated in attachment (Figure 1A, 2B).

Flexion of the overlap zone can be seen during detachment from the surface, consistent with the

notion that conformational changes of specific regions of the disc can regulate attachment

(Supplemental Movie 1).

To investigate co-localization between *Gl*Actin and DAAP1, we calculated Pearson's

correlation coefficients for these two proteins using the JACoP plugin for ImageJ [25,26]. While

DAAP1-mNeonGreen does not appear to co-localize with *Gl*Actin during interphase, in mitosis

DAAP1 leaves the ventral disc and was observed to co-localize with *Gl*Actin (Figure 2C).

We also measured the intensity of DAAP1-mNeonGreen fluorescence in *Gl*Actin-

depleted cells, selecting a region in the ventral groove, as that is where DAAP1 expression in the

highest (Figure 3). While there was no significant difference between fluorescence levels in

attached cells (Figure 3A, 3B), this difference was significant in unattached cells (Figure 3C,

3D). Furthermore, in some *Gl*Actin-depleted cells, we observed DAAP1 loss of enrichment in

the entire disc (Figure 3C). These results indicate that *Gl*Actin is responsible for DAAP1 disc

localization.

The microtubules in the ventral disc are stable and do not undergo dynamic instability;

they rely on a set of core DAPs to maintain this stability [15]. During mitosis, the disc is

dismantled as the cells must detach to "swim" away from each other during cytokinesis [27,28].

New discs in the daughter cells appear during early cytokinesis and are built from microtubules

originating from the mother disc and the median body [27,29]. Notably, DAAP1-mNeonGreen

remains on the disc during telophase, late into disassembly, and begins to coalesce again on the

daughter discs in mid cytokinesis. In newly-divided cells, DAAP1-mNeonGreen has returned to

its disc localization and is once more enriched in the ventral groove (Figure 2C). Fluorescence

recovery after photobleaching (FRAP) confirmed the stability of DAAP1-mNeonGreen for disc

localization, as fluorescence levels did not recover for 12 minutes after photobleaching (Figure

4) in either of the two regions of the disc where DAAP1 is enriched. Viability was confirmed by

cell movement as seen in Supplemental Movie 2. This result is similar to the fluorescence

recovery of other DAPs. Taken together, these data indicate that DAAP1 is a *Gl*Actin interactor

and stably localized to the ventral disc in interphase.

DAAP1-depleted cells have intact ventral discs

Since *Gl*Actin is involved in both attachment and disc morphology, we asked if DAAP1

could have a similar role. We attempted depletion of DAAP1 with translation-blocking

morpholinos, however, the sequence was design-resistant and the knockdown level inconsistent.

We then turned to CRISRPi to investigate DAAP1's potential role in attachment and disc

morphology. To monitor disc morphology as well as deplete DAAP1, we inserted the HALO-

tagged delta-giardin marker into the dead Cas9- (dCas9) based CRISPRi plasmid and introduced

this into a DAAP1-mNeonGreen expressing cell line. This allowed us to evaluate knockdown on

both a whole population (Figure 5A) and an individual cell basis. We analyzed the fluorescence

of endogenously-tagged DAAP1-mNeonGreen in control cells expressing dCas9 with a non-

specific guide RNA to those expressing a DAAP1 specific small guide RNA resulting in an

average decrease in expression of 40% for the population. We once again separated cells that

were attached to culture tubes from unattached cells to probe for disc defects in each

subpopulation. We did observe a higher proportion of disc defects in the DAAP1-depleted cells

than in the control, but this difference was not significant (Figure 5B and 5C). These results

indicate that a 40% depletion of DAAP1 did not impact gross ventral disc morphology, as

assessed by light microscopy.

Inability to Maintain Attachment in DAAP1-Depleted Cells

DAAP1 knockdown cells had no visible defects, so DAAP1 may not be needed for disc

assembly. To investigate a possible role in attachment, we used the aforementioned attachment

assay on DAAP1-depleted cells and observed a small but significant decrease in attachment for

two of three replicates relative to control cells (Supplemental Table 2). As CRISPRi is known to

have variable penetrance in *Giardia* [30], we analyzed the fluorescence in unattached and

attached cells separately. Here we found reduced levels of DAAP1-mNeonGreen in unattached

cells compared to attached ones (Figure 6A). This result prompted us to further explore the

possible role of DAAP1 in attachment using a shear force attachment assay. In their normal

intestinal habitat, *Giardia* must maintain attachment even when subjected to the flow of chyme.

With this in mind, we challenged DAAP1-depleted cells with a fluid shear force. DAAP1-

depleted and control cells were allowed to attach inside a flow chamber and then subjected to a

flow of growth medium. Cell position and velocity were tracked using Trackmate (Figure 6B,

[31]). DAAP1-depleted cells slid significantly faster under flow than their control counterparts

(Figure 6C, 6D). Furthermore, more control cells remained in the field of view than DAAP1-

depleted cells over time (Figure 6E), indicating a better ability to maintain attachment.

Impaired Seal Formation in DAAP1-Depleted Cells

We then hypothesized that the impaired attachment in DAAP1-depleted cells may be due to impairments of the ventral groove or overlap zone "gates" that control fluid flow and buildup of negative pressure differentials under the disc. We used fluorescent microspheres to investigate this fluid flow in live, attached cells imaged in normal growth media. We found that spheres entered under the disc more frequently in DAAP1-depleted than dCas9 control cells, consistent with a defect in gating flow under the disc (Figure 7A). Furthermore, we found more beads per cell in the DAAP1-depleted cells (Supplemental Table 4).

Since the ability of the gates to close affects disc seal formation [8,14], we used plasma membrane stain, CellMask Orange and Total Internal Fluorescence (TIRF) microscopy to evaluate seal formation for DAAP1-depleted and dCas9-Control cells. This approach allowed us to assess the different levels of attachment based on which portions of the cell were in contact with the surface. Since stage 1 of attachment is unattached skimming, we omitted these cells from our analysis, and only included cells in stage 2 (lateral crest contact), stage 3 (lateral shield contact), and stage 4 (bare area contact). We found more cells at stages 2 and 3 and fewer in stage 4 for DAAP1-depleted cells than in the dCas9-Control (Figure 7B), indicating impaired seal formation around the ventral disc. Taken together, these data demonstrate that DAAP1 functions at the disc gates that regulate fluid flow and ventral disc seal formation.

Discussion

Here, we have confirmed that *Gl*Actin has a role in ventral disc morphology and disc-based attachment (Figure 1). We also show that *Gl*Actin interactor DAAP1 is a stable component

of the ventral disc (Figure 4). While perturbation of DAAP1 levels was not associated with

visible effects on disc morphology, it resulted in decreased attachment (Figures 5-7).

The ventral disc is well known as a *Giardia*-specific microtubule based structure [17,32]

and our data show that the actin cytoskeleton has an important role in ventral disc assembly and

function. A functional relationship between actin and microtubules has been well-documented

for other structures. In *Chlamydomonas*, actin is required for intraflagellar transport [33,34].

Actin monomers are a key component of the flagella through interaction with the inner dynein

arm [35–38] and recent evidence indicates actin is a component of the gamma-tubulin ring

complex [39]. Actomyosin interactions influence the orientation of the mitotic spindle [40,41]

and F-actin associates directly with mitotic spindles in *Xenopus* and starfish embryos [42,43].

Actin is also involved in microtubule- or tubulin-based host interaction in other parasites. For

instance, one of *Toxoplasma gondii's* formins localize exclusively to the conoid [44], a

latticelike organelle composed of tubulin which initiates host invasion [45]. There is evidence of

coordination between the microtubules of *Toxoplasma*'s glideosome, which is used for actin-

based motility, during host invasion [46]. Furthermore, actin polymerization is believed to

contribute to conoid extrusion [47,48], and the conoid-localized Myosin H is essential for entry

and exit from host cells [49,50]. *Gl*Actin polymerization could have an analogous role in

controlling the movement of the ventral disc to facilitate attachment.

We also observed defects in the ventral disc in our *Gl*Actin-depleted cells (Figure 1),

which we characterized as either mild (misshapen, broken, or with small gaps) or severe

(unwound, severely misshapen, often associated with cytokinesis defects). There are several

possible explanations for these phenotypes. During mitosis, the ventral disc is dismantled and

two daughter discs are rebuilt during cytokinesis [27]. One possibility is that *Gl*Actin has a role

in loading proteins on the assembling disc. *Gl*Actin also likely contributes to ventral disc

positioning, since actin has a role in positioning basal bodies that nucleate the disc [51,52].

However, many of the cells with defective discs have normally-positioned flagella (Figure 1,

Figure S2), indicating defective discs are not entirely due to basal body mispositioning. While

the resolution of light microscopy limits our ability to see details in the microtubule array of the

disc [53], over 25% of our unattached *Gl*Actin-depleted cells lacked noticeable defects (Figure

1). Therefore, a portion of the attachment defect cannot be attributed to disc defects alone.

Another possibility is that *Gl*Actin helps maintain the structure of the disc, consistent

with its role in maintaining cell and organelle shape in other organisms [19,22]. The unusual

stability of disc microtubules can be at least partially attributed to certain DAPs [8,15,17], which

lock the microtubules into place, preventing their turnover. *Gl*Actin interacts with at least one of

these DAPs, GL50803_5188 [18], and could be working in conjunction with it to stabilize the

disc microtubules themselves. Another explanation is that *Gl*Actin works to stabilize one of the

non-microtubule components of the ventral disc. The microtubules of the disc are overlaid by the

giardin-composed microribbons [54], which are connected by crossbridges [13,55]. Since the

spiral shape is often maintained, but unwound, in *Gl*Actin-depleted cells (Figure 1, Figure S2), it

may be that *Gl*Actin contributes to establishing and/or maintaining disc overlap, as the overlap

zone also functions as the anterior gate of the ventral disc.

A previous study from our lab identified eight *Gl*Actin interactors which localize to the

ventral disc, including DAAP1 [18]; here we confirmed complex formation between *Gl*Actin and

DAAP1 through co-immunoprecipitation (Figure 2). We did not detect co-localization between

DAAP1 and *Gl*Actin in interphase cells, but present evidence that indicates co-localization

during mitosis when new discs are assembled (Figure 2). Polymerized *Gl*Actin may be enriched

in the ventral disc at these stages, but detection may be obscured due to technical limitations.

Unfortunately, use of the anti-*Gl*Actin antibody does not differentiate between filaments and

pools of monomeric *Gl*Actin which likely obscures some filaments. Currently, there are no

methods to specifically detect F-*Gl*Actin, as Rhodamine-Phalloidin does not bind to *Gl*Actin

filaments [22]. Actin in *Toxoplasma gondii* was believed to be primarily monomeric and only

capable of forming short filaments [56,57]. However, expression of an anti-actin nanobody

revealed a previously unseen complex network of F-actin [58]. Poor detection of *Gl*Actin using

the anti-*Gl*Actin antibody could also be due to steric hindrance. Actin-binding proteins could

sterically block access of anti-*Gl*Actin antibodies to disc-localized *Gl*Actin, similar to low anti-

tubulin antibody staining compared to our ability to see microtubules of the disc with

mNeonGreen-Tubulin [27]. Another possibility is that actin only transiently associates with the

disc for specific conformational changes which would be challenging to capture in fixed cells. A

live marker specific for *Gl*Actin is needed to determine the extent of actin association with

conformational dynamics of the ventral disc including disc doming, bare area protrusion and disc

flexion as seen in Movie S1.

Similar to other disc-associated proteins [44], DAAP1's localization to the disc is stable,

as we did not see any recovery of this protein for 12 minutes after photobleaching (Figure 4).

While DAAP1 was previously identified as a potential disc interactor, it was not found in high

enough abundance in the detergent-extracted disc proteome to qualify for further study as a disc-

associated protein [14]. These results possibly indicating that this protein is a peripheral

component of the disc, which would be expected in order for it to engage with a dynamic subset

of *Gl*Actin.

Using CRISPR-interference, we depleted DAAP1 from cells, and found a mild defect in

attachment, but no obvious impact on disc morphology (Figures 5-7). While a DAAP1 knockout

would give us better insight into how DAAP1 loss affects the disc, the ability to readily knockout

genes of interest remains under development. CRISPRi and Morpholino knockdown both have

variable penetrance, and thus the level of depletion fluctuates among cells, and we observed

normal discs even in cells with very little DAAP1 fluorescence (Figure 5). In contrast, depletion

of median body protein with morpholinos results in broken discs [12]. Although an inability to

attach can be attributed to disc defects [12,30], the depletion of gamma-giardin, another disc-

localized protein that interacts with *Gl*Actin [18], resulted in malformed discs with normal

attachment [59]. These findings indicate that some disc proteins may be involved in maintaining

the structural integrity of the disc, while others are involved in controlling conformational

changes of the disc required for modulating attachment. Our data is consistent with DAAP1

being in the latter category. [12]

The possibility that some disc proteins contribute to attachment but not structure may

also explain the disparity between the severity of attachment defects in *Gl*Actin-depleted cells

and DAAP1-depleted cells. Since many of the discs in *Gl*Actin-depleted cells were unwound or

misshapen, this suggests roles in building the disc or collaboration with DAPs that maintain the

disc structure. However, since over 25% of our unattached *Gl*Actin-depleted cells had apparently

normal discs, the attachment defect cannot be attributed to disc defects alone. Our previous study

identified eight other putative disc-localized F-*Gl*Actin interactors which could also contribute to

attachment and/or disc morphology [18].

An additional explanation for the severity of attachment defects in *Gl*Actin-depleted cells

is due to its involvements in supplemental mechanisms of attachment. *Gl*Actin has been

demonstrated to function in building the ventrolateral flange, *Giardia's* membrane lamella which

also contributes to attachment [60,61]. Further, the marginal plate, a rigid structure associated

with the anterior axonemes, is also hypothesized to be involved in attachment [11,32]. *Gl*Actin is

stably enriched in this structure [22,62] and we have described six interactors which localize

there as well [18]. Given *Gl*Actin's involvement in all three structures hypothesized to augment

attachment, depletion of *Gl*Actin likely has multivariant effects on attachment, contributing to a

more severe phenotype.

The biology of parasites is often unlike anything seen in other organisms, shaped by the

unusual environments in which they live. *Giardia's* ventral disc exemplifies parasitic innovation

and makes this *Giardia* specific structure a target for drug development. Since most DAPs lack

homologs, they are in theory potential drug targets; however, the lack of active sites reduces their

potential [63]. Further understanding of the mechanisms of attachment and its regulation will

increase the viability of developing drugs that could specifically function through regulating

attachment. Actin, however, is a druggable protein and conventional actin drugs do not bind to

*Gl*Actin or inhibit its function, rendering it an appealing therapeutic target.

Materials and Methods

Parasite Strain and Growth Conditions

G. lamblia strain WB clone 6 (ATCC 50803; American Type Culture Collection) was cultured as

described previously [22].

Vector Construction

All constructs in this study were C-terminal fusions, made using Gibson reactions with linearized

vectors and PCR products [64]. See Supplemental Table 3 for primer sequences and workflow.

Live Cell Imaging

Cells were chilled with ice for 15 minutes to detach from the culture tube and then placed into an

Attofluor cell chamber (Molecular Probes) and incubated in a GasPak EZ anaerobic pouch (BD)

or a Tri-gas incubator (Panasonic) set to 2.5% O2, 5% CO2 for 90 minutes at 37°C. Cells were

then washed four times with HEPES-buffered saline (HBS: 137mM NaCl, 5mM KCl, 0.91mM

Na2HpO4-heptahydrate, 5.55mM Glucose, 20mM HEPES, pH7). Live cell imaging was

performed on a DeltaVision Elite microscope (GE) equipped with DIC optics, using a 100×1.4

NA or 60×1.42 NA objective, and a sCMOS 5.4 PCle air-cooled camera (PCO-TECH).

Coimmunoprecipitation

Cell cultures of 500 mL each of *Giardia* wildtype and DAAP1-3xHA were grown for 3 days,

then iced for 2 hours to detach, and spun for 1500xg at 4°C for 20 minutes. Cells were then

washed twice in 1X HBS with 2X HALT protease inhibitors and 10 µM chymostatin, 1 µM

leupeptin, and 1 µM E64. Each pellet was resuspended to a final volume of 1.2 mL and 100 mM

DSP in DMSO was added to a final concentration of 1 mM and incubated at room temperature

for 30 minutes. The reaction was quenched for 15 minutes with an addition of Tris pH 7.4 final

concentration 20 mM. Cells were then pelleted by spinning for 7 minutes at 700xg and

resuspended in 350 µL lysis buffer (80 mM KCl, 10 mM imidazole, 1 mM MgCl2, 1 mM

EGTA, 5% Glycerol, 20 mM HEPES, 0.2 mM, CaCl2, 10 mM ATP, 0.1% Triton X-100, 500

mM NaCl, pH 7.2, 2XHALT protease inhibitors and 10 µM chymostatin, 1µM leupeptin, and 1

µM E64). Cells were then lysed by sonication and cleared with a 10 minute spin at 10,000xg. A

volume of 17.5 µL of equilibrated EZview Red Anti-HA Affinity gel (Sigma) was added to each

tube of lysate, then incubated at 4 ºC with end-over-end mixing for 1 hour. Beads were then spun

at 8,200xg for 30 seconds and the supernatant was discarded, followed by a total of three washes

with 750 µL wash buffer (80 mM KCl, 10 mM imidazole, 1 mM MgCl2, 1 mM EGTA, 5%

Glycerol, 20 mM HEPES, 0.2 mM CaCl2, 10 mM ATP, 0.5% Tween, 500 mM NaCl, pH 7.2).

Each wash consisted of end-over-end rotation for 5 minutes followed by a 30 second spin at

8,200 x g. Protein was then incubated with 50 µL of 8 M Urea at RT for 20 minutes to elute. The

beads were pelleted and then sample buffer was added to the supernatant, the sample was boiled

for 5 mins at 98 ºC, and run on a 12% SDS-PAGE gel, followed by western blot protocol

described previously [22].

Immunofluorescence Microscopy

Fixation and imaging were performed as described in previous work [65], with the addition of a

JaneliaFluor 646 HaloTag ligand (Promega, GA1120) or JaneliaFluor 549 HaloTag ligand

(Promega, GA1110) at 1:100 in the second immunostaining step. Image analysis was performed

in FIJI [66]. DAAP1 intensity was measured by generating a circular region of interest

surrounding the ventral groove and measuring fluorescence intensity at the brightest point in

each image stack.

Morpholino Knockdown

Morpholino knockdown was performed as described previously [67]. Cells were lysed and run

on a 10% SDS-PAGE gel, followed by western blot protocol as described previously [18,22].

For each of three biological replicates, samples were divided equally into three lanes and

intensity of each lane measured using FIJI and averaged. Cell counts were performed with a

MoxiZ coulter counter to ensure the same number of cells were included in each sample. For

morpholino sequences, see Supplemental Table 3.

Culture Tube Attachment Assay

*Gl*Actin: Cells were treated with either a standard control or anti-*Gl*Actin morpholino and

allowed to recover for 24 hours in a total volume of 13 mL of TYDK media at 37 °C. At 24

hours, each culture tube was inverted six times; detached cells in media were decanted and

counted with a MoxiZ coulter counter (Orflo), then replaced with 13 mL ice cold media. Culture

tubes were placed on ice for 30 minutes to detach the remaining cells, after which these were

also counted. Cells were then imaged live or fixed as described previously [27]. Three

independent replicates of each experimental and control were analyzed. DAAP1 assays were

performed similarly, except knockdown was accomplished with CRISPRi rather than

morpholinos (see below).

Assessment for Live/Dead Cells

Stock solutions of dye were created by dissolving 5 mg fluorescein diacetate (Sigma-Aldrich, F-

7378) in 1 ml acetone and 2 mg propidium iodide (Sigma-Aldrich, P4170) in 1 ml HBS. A 2X

stain solution was prepared by diluting the stock solutions of fluorescein diacetate and propidium

iodide by 1:100 in 1X HBS. Cells were pelleted by centrifugation at 500 x g and resuspended in

1 mL 1X HBS. 1mL of 2x stain solution was then added to the cell suspension, washed twice in

1X HBS by centrifugation and resuspended in 1X HBS for imaging.

Fluorescence Recovery after Photobleaching.

FRAP was performed on DAAP1-mNeonGreen in the ventral disc to measure protein turnover.

Cells were chilled with ice for 20 minutes to detach from the culture tube and then placed into an

Attofluor cell chamber (Molecular Probes) and incubated in a O_2/CO_2 incubator (Panasonic) for

1-2 hours at 37° C. Media was replaced with 37 ºC 1X HBS before imaging. All experiments

were conducted using a FRAP-enabled DeltaVision Spectris with 405 nm solid-state laser, 100%

laser power, 50 mW laser, and 25 ms stationary pulse. Three images were taken until the bleach

event for 0.5 second intervals, followed by image acquisition every 10 sec for 10 minutes.

Normalized mNeonGreen fluorescence recovery was calculated by subtracting the background

noise from the ROI intensity measurement; to normalize for photobleaching due to imaging, this

measurement was then divided by a fluorescent control ROI intensity measurement with the

background noise subtracted as well.

CRISPR Interference Knockdown

Constructs were designed as described previously [68]. For primer design and workflow, see

Supplemental Table 3. Cells containing an integrated C-terminally mNeonGreen-tagged DAAP1

were transfected with a dead Cas9 vector which contained either a non-specific small guide RNA

(dCas9-CTRL) or a small guide RNA targeting base pairs 392 through 411 of the coding region

of DAAP1 (dCas9+sgRNA392-411). Knockdown efficiency was measured using a Tecan Spark

plate reader. Cells were pelleted and washed with 1X PBS and then resuspended in 1X PBS. An

equal number of dCas9-CTRL or dCas9+sgRNA392-411cells were loaded into Tecan black flat

96-well plates and the volume was adjusted to a total of 150 µL. Three measurements per sample

were averaged and three independent transformations were averaged for total knockdown

efficiency, normalized to the control and blanked with 1X PBS.

Shear Force Attachment Assay

Microfluidic devices were fabricated using a silicon master, which was created using deep

reactive ion etching. The silicon master contained a raised channel that had a rectangular cross-

sectional area, with a length of 1 cm, a width of 1 mm, and a height of 115 µm. A

polydimethylsiloxane (PDMS) (Sylgard 184, Dow Corning) mold was made using a 10:1 base to

curing agent ratio, which was mixed for 5 minutes, degassed for 20 minutes, poured over the

master, and placed in an oven for 10 minutes at 110 ºC to cure. The mold was peeled away from

the silicon master and holes were punched for the inlet and outlet of the channel. Afterwards,

inlet and outlet ports were created using a custom aluminum mold with pegs to insert the silicone

tubing (0.040 inch ID/ 0.085 inch OD, HelixMark). Degassed 10:1 PDMS was poured into the

preheated aluminum mold and cured at 110 ◦C for 1 hour. The inlet and outlet ports were aligned

with holes on the top face of the microfluidic device and stuck together using uncured PDMS

and 5 minutes of oven heat. Finally, the bottom face of the microfluidic device and a clean glass

slide were plasma treated for 10 seconds and aligned together to create a water-tight seal

between the two layers.

Control dCas9 or DAAP1 targeting dCas9+sgRNA392-411 trophozoites were chilled for 20

minutes on ice and then loaded into microfluidic devices placed in a GasPak EZ anaerobic pouch

(BD Scientific) at 37 °C for 1 h to allow re-attachment. Growth media was drawn into a

disposable syringe (BD Scientific) and then loaded onto a syringe pump (Harvard Apparatus).

For experiments, the flow rate was set to 20 µl/min for 30 s to clear swimming trophozoites and

then ramped to 100 µl/min over 5 s and challenged for 60 s. The challenge assays were recorded

at 2 FPS using a Nikon Eclipse Ti microscope with a 40X objective, using phase illumination,

Clara DR-1357 CCD camera (Andor), and a live-cell incubation chamber (In Vivo Scientific,

Inc) to maintain a temperature of 37 °C. Trophozoite mean speed was quantified starting at 35s

using the TrackMate ImageJ plugin via manual tracking over the first 10 seconds of 100µL/min

flow [31].

TIRF Microscopy

Cells were prepared for live imaging as described above, with the addition of CellMask™

Orange Plasma membrane Stain (ThermoFisher) according to manufacturer's instructions after

media replacement with 1X HBS. Cells were imaged in a chamber with 2% CO2 and 5% O2

with a DeltaVision OMX (GE) using 60X TIRF objective. Images were taken from randomly

selected fields of view. Each dish was imaged for a maximum of 30 minutes to minimize stress

on the cells. Three dishes from two independent experiments were used for each condition.

Live Imaging with Microspheres

Cells expressing DAAP1-mNeonGreen were transfected with a plasmid containing Halo-tagged

delta-giardin and either dCas9 control or dCas9+sgRNA392-411 to deplete DAAP1. Cells were

grown in parallel and upon confluency were chilled with ice for 15 minutes to detach from the

culture tube and then placed into an Attofluor cell chamber (Molecular Probes), then incubated

in a GasPak EZ anaerobic pouch (BD) or a Tri-gas incubator (Panasonic) set to 2.5% O2, 5%

CO_2 for 90 minutes at 37 °C. 646 HaloTag ligand (Promega, GA1120) was added to cells 45

minutes after incubation at 1:1000 concentration, incubated for 15 minutes, and then replaced

with TYDK for 30 minutes prior to imaging. 2.5 µL of fluorescent microspheres (0.2 µm,

Polysciences Fluoresbrite® YO Carboxylate Microspheres catalog number 19391) were added to

0.5 mL 1xHBS and sonicated for 10 minutes using a bath sonicator to disrupt aggregates. This

diluted solution was then added to cells at a 1:10 ratio to each chamber (30 µL diluted bead

solution in 300 µL cells). Images of single focal planes in which the disc was visible were taken

every 1 second for 15 seconds for each condition. Cells with beads which were collected under

their ventral disc and cells with no beads were counted. Cells which detached or moved were

excluded from analysis. Live cell imaging was performed on a DeltaVision Elite microscope

(GE) equipped with DIC optics, using a 60×1.42 NA objective, and a sCMOS 5.4 PCle air-

cooled camera (PCO-TECH).

Acknowledgements

We thank Kelly Hennessey, Kelli Hvorecny, Han-wei Shih, Elizabeth Thomas, Barbara

Wakimoto, Susan Parkhurst and Germain Alas for manuscript editing. We thank Pang Chan for

assistance with TIRF microscopy at the Biology Imaging Facility.

This material is based upon work supported by the National Science Foundation Graduate

Research Fellowship under Grant No. DGE-1762114 to MSO. Funding for the OMX was

provided by grant proposal S10 OD021490. Research reported in this publication was supported

by the National Heart, Lung, And Blood Institute of the National Institutes of Health under

Award Number F31HL156697. The content is solely the responsibility of the authors and does

not necessarily represent the official views of the National Institutes of Health."

References

1. Lane S, Lloyd D. Current Trends in Research into the Waterborne Parasite Giardia. Critical Reviews in Microbiology. 2002;28. doi:10.1080/1040-840291046713
2. Savioli L, Smith H, Thompson A. Giardia and Cryptosporidium join the 'Neglected Diseases Initiative.' Trends in Parasitology. 2006;22. doi:10.1016/j.pt.2006.02.015
3. Adam RD. Biology of Giardia lamblia. Clinical Microbiology Reviews. 2001;14. doi:10.1128/CMR.14.3.447-475.2001
4. Dawson SC. An insider's guide to the microtubule cytoskeleton of *Giardia*. Cellular Microbiology. 2010;12. doi:10.1111/j.1462-5822.2010.01458.x
5. Schwartz CL, Heumann JM, Dawson SC, Hoenger A. A Detailed, Hierarchical Study of Giardia lamblia's Ventral Disc Reveals Novel Microtubule-Associated Protein Complexes. PLoS ONE. 2012;7. doi:10.1371/journal.pone.0043783
6. Elmendorf HG, Dawson SC, McCaffery JM. The cytoskeleton of Giardia lamblia. International Journal for Parasitology. 2003;33. doi:10.1016/S0020-7519(02)00228-X
7. Nosala C, Dawson SC. The Critical Role of the Cytoskeleton in the Pathogenesis of Giardia. Current Clinical Microbiology Reports. 2015;2. doi:10.1007/s40588-015-0026-y
8. Nosala C, Dawson SC. The ventral disc is a flexible microtubule organelle that depends on domed ultrastructure for functional attachment of Giardia lamblia. bioRxiv. bioRxiv; 2017. doi:10.1101/213421
9. Nosala C, Hagen KD, Dawson SC. 'Disc-o-Fever': Getting Down with Giardia's Groovy Microtubule Organelle. Trends in Cell Biology. 2018;28. doi:10.1016/j.tcb.2017.10.007
10. Lenaghan SC, Davis CA, Henson WR, Zhang Z, Zhang M. High-speed microscopic imaging of flagella motility and swimming in Giardia lamblia trophozoites. Proceedings of the National Academy of Sciences. 2011;108: E550–E558. doi:10.1073/pnas.1106904108
11. Hagen KD, McInally SG, Hilton ND, Dawson SC. Microtubule organelles in Giardia. 2020. doi:10.1016/bs.apar.2019.11.001
12. Woessner DJ, Dawson SC. The Giardia Median Body Protein Is a Ventral Disc Protein That Is Critical for Maintaining a Domed Disc Conformation during Attachment. Eukaryotic Cell. 2012;11. doi:10.1128/EC.05262-11
13. Brown JR, Schwartz CL, Heumann JM, Dawson SC, Hoenger A. A detailed look at the cytoskeletal architecture of the Giardia lamblia ventral disc. Journal of Structural Biology. 2016;194. doi:10.1016/j.jsb.2016.01.011
14. House SA, Richter DJ, Pham JK, Dawson SC. Giardia Flagellar Motility Is Not Directly Required to Maintain Attachment to Surfaces. PLoS Pathogens. 2011;7. doi:10.1371/journal.ppat.1002167
15. Nosala C, Hagen KD, Hilton N, Chase TM, Jones K, Loudermilk R, et al. Disc-associated proteins mediate the unusual hyperstability of the ventral disc in *Giardia lamblia*. Journal of Cell Science. 2020;133. doi:10.1242/jcs.227355

16. Hagen KD, Hirakawa MP, House SA, Schwartz CL, Pham JK, Cipriano MJ, et al. Novel
Structural Components of the Ventral Disc and Lateral Crest in Giardia intestinalis. PLoS
Neglected Tropical Diseases. 2011;5. doi:10.1371/journal.pntd.0001442

17. Nosala C, Hagen KD, Dawson SC. 'Disc-o-Fever': Getting Down with Giardia's Groovy
Microtubule Organelle. Trends in Cell Biology. 2018;28. doi:10.1016/j.tcb.2017.10.007

18. Steele-Ogus MC, Johnson R, MacCoss M, Paredez AR. Identification of Actin Filament
Interactors in Giardia lamblia. [cited 7 Apr 2021].
doi:https://doi.org/10.1128/Spectrum.00558-21

19. Pollard TD. Actin and Actin-Binding Proteins. Cold Spring Harbor Perspectives in
Biology. 2016;8. doi:10.1101/cshperspect.a018226

20. Feely DE, Schollmeyer J v., Erlandsen SL. Giardia spp.: Distribution of contractile
proteins in the attachment organelle. Experimental Parasitology. 1982;53.
doi:10.1016/0014-4894(82)90100-X

21. Drouin G, de Sá MM, Zuker M. The Giardia lamblia actin gene and the phylogeny of
eukaryotes. Journal of Molecular Evolution. 1995;41. doi:10.1007/BF00173163

22. Paredez AR, Assaf ZJ, Sept D, Timofejeva L, Dawson SC, Wang C-JR, et al. An actin
cytoskeleton with evolutionarily conserved functions in the absence of canonical actin-
binding proteins. Proceedings of the National Academy of Sciences. 2011;108.
doi:10.1073/pnas.1018593108

23. Morrison HG, McArthur AG, Gillin FD, Aley SB, Adam RD, Olsen GJ, et al. Genomic
Minimalism in the Early Diverging Intestinal Parasite Giardia lamblia. Science. 2007;317.
doi:10.1126/science.1143837

24. Aurrecoechea C, Brestelli J, Brunk BP, Carlton JM, Dommer J, Fischer S, et al.
GiardiaDB and TrichDB: integrated genomic resources for the eukaryotic protist
pathogens Giardia lamblia and Trichomonas vaginalis. Nucleic Acids Research. 2009;37.
doi:10.1093/nar/gkn631

25. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
analysis. Nature Methods. 2012;9. doi:10.1038/nmeth.2089

26. BOLTE S, CORDELIÈRES FP. A guided tour into subcellular colocalization analysis in
light microscopy. Journal of Microscopy. 2006;224. doi:10.1111/j.1365-
2818.2006.01706.x

27. Hardin WR, Li R, Xu J, Shelton AM, Alas GCM, Minin VN, et al. Myosin-independent
cytokinesis in *Giardia* utilizes flagella to coordinate force generation and direct membrane
trafficking. Proceedings of the National Academy of Sciences. 2017;114.
doi:10.1073/pnas.1705096114

28. Tůmová P, Kulda J, Nohýnková E. Cell division ofGiardia intestinalis: Assembly and
disassembly of the adhesive disc, and the cytokinesis. Cell Motility and the Cytoskeleton.
2007;64. doi:10.1002/cm.20183

29. Nohýnková E, Tu□mová P, Kulda J. Cell Division of Giardia intestinalis: Flagellar
Developmental Cycle Involves Transformation and Exchange of Flagella between
Mastigonts of a Diplomonad Cell. Eukaryotic Cell. 2006;5. doi:10.1128/EC.5.4.753-
761.2006

30. McInally SG, Hagen KD, Nosala C, Williams J, Nguyen K, Booker J, et al. Robust and
stable transcriptional repression in *Giardia* using CRISPRi. Molecular Biology of the
Cell. 2019;30. doi:10.1091/mbc.E18-09-0605

31.   Tinevez J-Y, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E, et al.
588   TrackMate: An open and extensible platform for single-particle tracking. Methods.
589   2017;115. doi:10.1016/j.ymeth.2016.09.016
590   32.   Dawson SC, Paredez AR. Alternative cytoskeletal landscapes: cytoskeletal novelty and
591   evolution in basal excavate protists. Current Opinion in Cell Biology. 2013;25.
592   doi:10.1016/j.ceb.2012.11.005
593   33.   Avasthi P, Onishi M, Karpiak J, Yamamoto R, Mackinder L, Jonikas MC, et al. Actin Is
594   Required for IFT Regulation in Chlamydomonas reinhardtii. Current Biology. 2014;24.
595   doi:10.1016/j.cub.2014.07.038
596   34.   Jack B, Mueller DM, Fee AC, Tetlow AL, Avasthi P. Partially Redundant Actin Genes in
597   Chlamydomonas Control Transition Zone Organization and Flagellum-Directed Traffic.
598   Cell Reports. 2019;27. doi:10.1016/j.celrep.2019.04.087
599   35.   Gervasi MG, Xu X, Carbajal-Gonzalez B, Buffone MG, Visconti PE, Krapf D. The actin
600   cytoskeleton of the mouse sperm flagellum is organized in a helical structure. Journal of
601   Cell Science. 2018;131. doi:10.1242/jcs.215897
602   36.   Yanagisawa H, Kamiya R. Association between Actin and Light Chains in
603   Chlamydomonas Flagellar Inner-Arm Dyneins. Biochemical and Biophysical Research
604   Communications. 2001;288. doi:10.1006/bbrc.2001.5776
605   37.   Kato-Minoura T, Hirono M, Kamiya R. Chlamydomonas Inner-Arm Dynein Mutant, ida5,
606   Has a Mutation in an Actin-encoding Gene. Journal of Cell Biology. 1997;137.
607   doi:10.1083/jcb.137.3.649
608   38.   Gui M, Ma M, Sze-Tu E, Wang X, Koh F, Zhong ED, et al. Structures of radial spokes
609   and associated complexes important for ciliary motility. Nature Structural & Molecular
610   Biology. 2021;28. doi:10.1038/s41594-020-00530-0
611   39.   Wieczorek M, Urnavicius L, Ti S-C, Molloy KR, Chait BT, Kapoor TM. Asymmetric
612   Molecular Architecture of the Human γ-Tubulin Ring Complex. Cell. 2020;180.
613   doi:10.1016/j.cell.2019.12.007
614   40.   Sugioka K, Bowerman B. Combinatorial Contact Cues Specify Cell Division Orientation
615   by Directing Cortical Myosin Flows. Developmental Cell. 2018;46.
616   doi:10.1016/j.devcel.2018.06.020
617   41.   Scarpa E, Finet C, Blanchard GB, Sanson B. Actomyosin-Driven Tension at
618   Compartmental Boundaries Orients Cell Division Independently of Cell Geometry
619   In Vivo. Developmental Cell. 2018;47. doi:10.1016/j.devcel.2018.10.029
620   42.   Kita AM, Swider ZT, Erofeev I, Halloran MC, Goryachev AB, Bement WM. Spindle–F-
621   actin interactions in mitotic spindles in an intact vertebrate epithelium. Molecular Biology
622   of the Cell. 2019;30. doi:10.1091/mbc.E19-02-0126
623   43.   Sugizaki A, Sato K, Chiba K, Saito K, Kawagishi M, Tomabechi Y, et al. POLArIS, a
624   versatile probe for molecular orientation, revealed actin filaments associated with
625   microtubule asters in early embryos. Proceedings of the National Academy of Sciences.
626   2021;118. doi:10.1073/pnas.2019071118
627   44.   Tosetti N, Dos N, Pacheco S, Soldati-Favre D, Jacot D. Three F-actin assembly centers
628   regulate organelle inheritance, cell-cell communication and motility in Toxoplasma
629   gondii. 2019. doi:10.7554/eLife.42669.001
630   45.   Hu K, Roos DS, Murray JM. A novel polymer of tubulin forms the conoid of Toxoplasma
631   gondii. Journal of Cell Biology. 2002;156. doi:10.1083/jcb.200112086

46. Das S, Stortz JF, Meissner M, Periz J. The multiple functions of actin in apicomplexan
parasites. Cellular Microbiology. 2021. doi:10.1111/cmi.13345

47. MONDRAGON R, FRIXIONE E. Ca $^{2+}$ -Dependence of Conoid Extrusion in Toxoplasma
gondii Tachyzoites. Journal of Eukaryotic Microbiology. 1996;43. doi:10.1111/j.1550-
7408.1996.tb04491.x

48. del Carmen MG, Mondragón M, González S, Mondragón R. Induction and regulation of
conoid extrusion in *Toxoplasma gondii*. Cellular Microbiology. 2009;11.
doi:10.1111/j.1462-5822.2009.01304.x

49. Graindorge A, Frénal K, Jacot D, Salamun J, Marq JB, Soldati-Favre D. The Conoid
Associated Motor MyoH Is Indispensable for Toxoplasma gondii Entry and Exit from
Host Cells. PLOS Pathogens. 2016;12. doi:10.1371/journal.ppat.1005388

50. dos Santos Pacheco N, Tosetti N, Koreny L, Waller RF, Soldati-Favre D. Evolution,
Composition, Assembly, and Function of the Conoid in Apicomplexa. Trends in
Parasitology. 2020;36. doi:10.1016/j.pt.2020.05.001

51. Roman A-C, Garrido-Jimenez S, Diaz-Chamorro S, Centeno F, Carvajal-Gonzalez JM.
Centriole Positioning: Not Just a Little Dot in the Cell. 2019. doi:10.1007/978-3-030-
23173-6_8

52. Antoniades I, Stylianou P, Skourides PA. Making the Connection: Ciliary Adhesion
Complexes Anchor Basal Bodies to the Actin Cytoskeleton. Developmental Cell. 2014;28.
doi:10.1016/j.devcel.2013.12.003

53. Halpern AR, Alas GCM, Chozinski TJ, Paredez AR, Vaughan JC. Hybrid Structured
Illumination Expansion Microscopy Reveals Microbial Cytoskeleton Organization. ACS
Nano. 2017;11. doi:10.1021/acsnano.7b07200

54. Holberton DV. Arrangement of subunits in microribbons from Giardia. Journal of Cell
Science. 1981;47. doi:10.1242/jcs.47.1.167

55. Gadelha APR, Benchimol M, de Souza W. Helium ion microscopy and ultra-high-
resolution scanning electron microscopy analysis of membrane-extracted cells reveals
novel characteristics of the cytoskeleton of Giardia intestinalis. Journal of Structural
Biology. 2015;190. doi:10.1016/j.jsb.2015.04.017

56. Skillman KM, Ma CI, Fremont DH, Diraviyam K, Cooper JA, Sept D, et al. The unusual
dynamics of parasite actin result from isodesmic polymerization. Nature Communications.
2013;4. doi:10.1038/ncomms3285

57. Angrisano F, Delves MJ, Sturm A, Mollard V, McFadden GI, Sinden RE, et al. A GFP-
Actin reporter line to explore microfilament dynamics across the malaria parasite
lifecycle. Molecular and Biochemical Parasitology. 2012;182.
doi:10.1016/j.molbiopara.2011.11.008

58. Periz J, Whitelaw J, Harding C, Gras S, del Rosario Minina MI, Latorre-Barragan F, et al.
Toxoplasma gondii F-actin forms an extensive filamentous network required for material
exchange and parasite maturation. eLife. 2017;6. doi:10.7554/eLife.24119

59. Kim J, Park S-J. Role of gamma-giardin in ventral disc formation of Giardia lamblia.
Parasites & Vectors. 2019;12. doi:10.1186/s13071-019-3478-8

60. Hardin WR, M Alas GC, Taparia N, Thomas EB, Hvorecny KL, Halpern AR, et al. The
Giardia lamellipodium-like ventrolateral flange supports attachment and rapid cytokinesis.
doi:10.1101/2021.01.31.429041

61.   ERLANDSEN SL, RUSSO AP, TURNER JN. Evidence for Adhesive Activity of the
Ventrolateral Flange in Giardia lamblia. The Journal of Eukaryotic Microbiology.
2004;51. doi:10.1111/j.1550-7408.2004.tb00165.x

62.   Paredez AR, Nayeri A, Xu JW, Krtková J, Cande WZ. Identification of Obscure yet
Conserved Actin-Associated Proteins in Giardia lamblia. Eukaryotic Cell. 2014;13.
doi:10.1128/EC.00041-14

63.   Makley LN, Gestwicki JE. Expanding the Number of 'Druggable' Targets: Non-Enzymes
and Protein-Protein Interactions. Chemical Biology & Drug Design. 2013;81.
doi:10.1111/cbdd.12066

64.   Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. Enzymatic
assembly of DNA molecules up to several hundred kilobases. Nature Methods. 2009;6.
doi:10.1038/nmeth.1318

65.   Krtková J, Thomas EB, Alas GCM, Schraner EM, Behjatnia HR, Hehl AB, et al. Rac
Regulates *Giardia lamblia* Encystation by Coordinating Cyst Wall Protein Trafficking and
Secretion. mBio. 2016;7. doi:10.1128/mBio.01003-16

66.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
analysis. Nature Methods. 2012;9. doi:10.1038/nmeth.2089

67.   Krtková J, Paredez AR. Use of Translation Blocking Morpholinos for Gene Knockdown
in Giardia lamblia. 2017. doi:10.1007/978-1-4939-6817-6_11

68.   Hennessey KM, Alas GCM, Rogiers I, Li R, Merritt EA, Paredez AR. Nek8445, a protein
kinase required for microtubule regulation and cytokinesis in *Giardia lamblia*. Molecular
Biology of the Cell. 2020;31. doi:10.1091/mbc.E19-07-0406

Figure 1: Depletion of *Gl*Actin affects attachment and ventral disc morphology

A) Left: Diagram of a *Giardia* trophozoite showing *Gl*Actin (magenta), disc marker delta-giardin

(cyan), and DAPI/nuclei (blue). Anterior (Af), posteriolateral (Plf), ventral (Vf), and caudal

flagella (Cf) are shown. Right: Enlarged diagram of ventral disc and areas: overlap zone (OZ),

ventral groove (VG), and bare area (BA). Arrows indicate the origin of the microtubule sheet and

orientation of the microtubule plus ends. **B)** Quantification of non-challenged attachment assays.

Attached and unattached cells were counted 24 hours after treatment with either a standard

control or anti-*Gl*Actin morpholino. Graph is representative of an average of three replicates,

mean ± sem. For control, 74% ± 6 were attached and 25% ± 6 were detached. For *Gl*Actin-

depleted cells 23% ± 2 were attached and 77% ± 2 detached. Raw numbers and detailed statistics

reported in Supplemental Table 1. Each replicate was analyzed separately with a two-sided chi-

square test with Yates' correction, P<0.0001 for each. **C)** Unattached cells from B were stained

with propidium iodide and fluorescein diacetate to distinguish dead from living cells

respectively. N=198 control cells and 222 *Gl*Actin-depleted cells. There was no significant

difference between the two groups based on a two-sided Fisher's exact test P= 0.7307. **D)** Both

attached and unattached *Gl*Actin-depleted and control cells were fixed and immunostained for

delta-giardin to assess ventral disc morphology using fluorescence microscopy. N=126 cells for

each condition. There was a significant difference in unattached cells based on a two-sided chi-

square test with Yates' correlation, P<0.0001. No significant difference was detected in attached

cells based on the same test, P<0.0903 **E)** Representative images of cells from D, with examples

of normal discs (teal), mild defects (yellow), and severe defects (pink). Scale bar denotes 5 µm.

Figure 2: DAAP1 is a *Gl*Actin interactor and a ventral disc-associated protein

A) Immunoprecipitation from extracts of cells expressing DAAP1-3XHA and wild type cells,

followed by western blots probed with anti-*Gl*Actin and anti-HA antibodies, revealed interaction

between the two proteins. **B)** Image of DAAP1-mNeonGreen in a live cell. The fusion protein

localizes to the disc and is enriched in the ventral groove (asterisk) and overlap zone (arrow).

Scale bar denotes 5 µm. **C)** Immunofluorescence of *Gl*Actin (magenta), tubulin (cyan), DAAP1-

mNeonGreen (green), and DAPI/nuclei (blue) during the cell cycle. DAAP1-

mNeonGreenremains on the disc into telophase, is incorporated into the nascent discs by mid

735   cytokinesis (see arrows), and localization is fully restored in newly divided cells. Pearson's

736   correlation coefficients using Costes randomization were calculated for colocalization between

737   DAAP1 and *Gl*Actin in each image, shown in the lower righthand corner. The Costes P-value

738   was 1 for all images. Scale bar denotes 5 µm.

Figure 3: DAAP1 loading onto the ventral disc requires *Gl*Actin.

A) Representative images of attached cells 24 hours after treatment with either an anti-*Gl*Actin

or control morpholino. Fluorescence intensity of DAAP1 at the ventral groove (see green circle)

was measured at the brightest section in each stack. *Gl*Actin shown in magenta, mNeonGreen-

DAAP1 in green, DAPI in blue. Scale bar denotes 5 µm. **B)** A two-tailed Mann-Whitney test

indicated that this difference was not statistically significant, ($N_{Control}$=57, $N_{GlActin-depletd}$=55),

P=0.227. Median and 95% confidence interval shown. Median of control=3131445. Median of

*Gl*Actin-depleted=3298353. Two and four outliers were detected by Tukey Fence, k=1.5 in the

control and *Gl*Actin-depleted condition respectively and were not included in the analysis. **C)**

Representative images of unattached cells 24 hours after treatment with either an anti-*Gl*Actin or

control morpholino. Fluorescence intensity of DAAP1 at the ventral groove (see green circle)

was measured at the brightest optical section in each stack. **D)** A two-tailed Mann-Whitney test

indicated that this difference was statistically significant, ($N_{Control}$=55, $N_{GlActin\text{-}depletd}$=54), P<0.0001.

Median and 95% confidence interval shown. Median of control= 3508064. Median of *Gl*Actin-

depleted=1609904. Four outliers were detected by Tukey Fence, k=1.5 in the *Gl*Actin-depleted

condition and were not included in the analysis.

Figure 4: DAAP1 is stably associated with the ventral disc

Fluorescence recovery after-photobleaching (FRAP) in DAAP1-mNeonGreen-cells showed no

recovery after 12 minutes for bleached areas in both the ventral groove **(A)** and overlap zone **(B).**

Three points of reference were taken, including bleached (ROI1, blue), unbleached (RO2, red),

764   and background (green), and normalized accordingly. Scale bar denotes 5 µm. N=10 for each

765   ROI in both A and B.

Figure 5: DAAP1-depleted discs do not have defects detectable by light microscopy A.)

Fluorescence of DAAP1-mNeonGreen cells transfected with a dead Cas9 vector containing

either a non-specific small guide RNA (dCas9-CTRL) or a small guide RNA targeting DAAP1

(dCas9+sgRNA392-411). Normalized to the control, the DAAP1-depleted cells demonstrated an

average knockdown of 40% ± 3 (three independent transformations). Attached and unattached

control and DAAP1-depleted cells were fixed and stained separately before assessing their disc

morphology. Based on a chi-square test, there was no significant difference in disc defects

775   observed between either attached or unattached groups. (Control attached N=129 cells, control

776   unattached N=126, DAAP1-depleted attached N=123, DAAP1-depleted unattached N=128.)

777   Attached cells, P=0.3483.Unattached cells:, P=0.1288. **C.)** Representative images from B:

778   *Gl*Actin (magenta), delta-giardin (cyan), DAAP1 (green), and DAPI/nuclei (blue). Scale bar

779   denotes 5 µm.

Figure 6: DAAP1 depletion results in reduced attachment

A) DAAP1 fluorescence levels from control (dCas9-CTRL) and DAAP1-depleted

(dCas9+sgRNA392-411). Unattached cells had a greater decrease in DAAP1 expression (38% ±

6) than attached cells (23% ± 2). Average of two independent transformations with three

technical replicates for each is shown. **B)** In this flow chamber assay, attached cells were

challenged with a flow rate of 100 µL/min. Cells were tracked for 10 seconds using TrackMate

software in ImageJ, colored lines indicating resulting paths. **C)** Distribution of mean flow-

induced speed. **D)** Mean flow-induced speed of each cell for 10 seconds after challenge, with

median and 95% confidence intervals. A two-tailed Mann-Whitney test indicated that this

difference was statistically significant, ($N_{dCas9-CTRL}$=124, $N_{dCas9+sgRNA392-411}$=111), P=0.001. Median of

control=10.56, upper 95% CI= 13.33, lower 95% CI=10.15. Median of DAAP1-depleted is

793   14.62, upper 95% CI=18.62, lower 95% CI=14.41. **E)** Kaplan-Meier curve of cells displaced

794   from the field of view over time, P= 0.0077 N=124 for control, N=111 for DAAP1-depleted

795   cells. Median and standard error shown.

Figure 7: DAAP1 has a role in ventral disc seal formation

A.) Fluorescent microspheres were added to live attached control (dCas) and DAAP1-depleted

(dCas9+sgRNA392-411) and imaged. Left: significantly more cells with microspheres under the

disc were observed in DAAP1-depleted cells based on a chi-square test exact test, P<0.001.

N=175 for control, 180 for DAAP1-depleted cells. Right, top: control cell. Right, bottom:

DAAP1-depleted cell with microsphere under disc (yellow arrow). Merged images shown with

DIC, microspheres (magenta), delta-giardin (cyan). Two-sided Fisher's exact test, P=0.0156 was

used for analysis. **B.)** TIRF microscopy assay to monitor the quality of attachment. Control and

DAAP1-depleted cells were stained with the plasma membrane stain CellMask Orange and

imaged with TIRF microscopy. Left: significantly more DAAP1-depleted cells were partially

attached than control cells based on a chi-square test, N=317 for each condition, P<0.005. Right:

images of cells in initial attachment (stage 2) with lateral crest (LC) contact, partial attachment

(stage 3), with lateral crest and lateral shield (LS) contact, or fully attached (stage 4), with lateral

crest, lateral shield, and bare area (BA) contact.

Supplemental figure 1: *Gl*Actin Depletion by antisense translation-blocking Morpholinos

A) Representative western blot of cells treated with either a standard control morpholino (left) or

morpholino targeting *GlActin* (right) for 24-hours and probed for tubulin (cyan) and *Gl*Actin

(magenta), resulting in a ~61% reduction of *Gl*Actin levels. **B)** Average of three morpholino

knockdowns results in an average of 50.3% ± 5.6 reduction of protein expression, as determined

by western blot. Values shown are mean ± SEM.

Supplemental Figure 2: Range of Disc Defect Phenotypes in *Gl*Actin-Depleted Cells

A) Representative *Giardia* cell from the control (attached) condition. **(B-E**) are *Gl*Actin depleted

cells with disc defects. Mild defects were categorized as discs which were broken **(B)** or

misshapen **(C)**. Severe defects included unwound discs **(D**) or disc defects associated with failed

cytokinesis **(E)**. All cells pictured in B-E were unattached, with the exception of the bottom row

of B. Scale bar denotes 5 µm.

Supplemental Table 1: Raw number of *Gl*Actin-depleted attachment assay cell counts and

statistics

Supplemental Table 2: DAAP1-depleted attachment assay cell counts and statistics

Supplemental Table 3: Primer and morpholino sequences and workflow

Supplemental Table 4: Number of microspheres per cell and statistics

836