TEMPORALLY AUTO-CORRELATED PREDATOR ATTACKS STRUCTURE ECOLOGICAL COMMUNITIES

Sebastian J. Schreiber*

Department of Evolution and Ecology

and Center for Population Biology

University of California

Davis, CA 95616, USA

sschreiber@ucdavis.edu

Abstract: For species regulated by a common predator, the P^* rule predicts that the prey species that supports the highest mean predator density (P^*) excludes the other prey species. This prediction is re-examined in the presence of temporal fluctuations in the vital rates of the interacting species including predator attack rates. When the fluctuations in predator attack rates are temporally uncorrelated, the P^* rule still holds even when the other vital rates are temporally auto-correlated. However, when temporal auto-correlations in attack rates are positive but not too strongly, the prey species can coexist due to the emergence of a positive covariance between predator density and prey vulnerability. This coexistence mechanism is similar to the storage effect for species regulated by a common resource. Strongly positive or negative auto-correlations in attack rates generate a negative covariance between predator density and prey vulnerability and a stochastic priority effect can emerge: with non-zero probability either prey species is excluded. These results highlight how temporally auto-correlated species' interaction rates impact the structure and dynamics of ecological communities.

Manuscript information: 2,412 words, 2 figures + online appendix.

INTRODUCTION

Predation or resource limitation can regulate populations. When multiple species are regulated by the same limiting factor, long-term coexistence is not expected under equilibrium conditions. Regulation due to a common, limiting resource can result in the R^* rule: the species suppressing the resource to the lower equilibrium level excludes other competitors [Volterra, 1928, Tilman, 1990, Wilson et al., 2007]. Regulation due to a common predator can result in the P^* rule: the prey species supporting the higher equilibrium predator density excludes the other prey species [Holt and Lawton, 1993, Schreiber, 2004, Holt and Bonsall, 2017]. Yet, many coexisting species share a common resource or a common predator. Understanding mechanisms permitting this coexistence is central to community ecology. One of these coexistence mechanisms, the storage effect, relies on temporal fluctuations in environmental conditions [Chesson and Warner, 1981, Chesson, 1994, Ellner et al., 2016]. Whether an analogous, fluctuation-dependent mechanism exists for species sharing a common predator is studied here.

Similar to species competing for a common resource, species sharing a common predator can exhibit mutually antagonistic interactions Holt [1977]: increasing the density of one prey species leads to an increase in predator density and a resulting increase in predation pressure on the other prey species. Thus, to the uninformed observer, the prey appear to be competing. Empirical support for apparent competition is extensive [Morris et al., 2004, Chaneton and Bonsall, 2000, Holt and Bonsall, 2017] and has significant implications for conservation biology [Gibson, 2006]. When the shared predator is the primary regulating factor, Holt and Lawton [1993] demonstrated that one prey excludes the other via the P^* rule. Yet in nature, coexisting species often share common predators. Holt and Lawton [1993] found that spatial refuges, resource limitation, and donor-controlled predation could help mediate coexistence. However, environmentally driven fluctuations in demographic rates did not promote coexistence [Holt and Lawton, 1993, Schreiber, 2021b]. These studies, however, assumed environmental fluctuations are temporally uncorrelated.

In contrast, environmental fluctuations are known, both theoretically and empirically, to mediate coexistence for species competing for a common resource. Intuitively, Hutchinson [1961] argued that environmental fluctuations favoring different competitors at different times may promote coexistence.

In a series of influential papers [Chesson and Warner, 1981, Chesson, 1983, 1988, 1994], Chesson re-

fined Hutchinson's idea and identified two fluctuation-dependent coexistence mechanisms: nonlinear

averaging and the storage effect. Empirical support for these mechanisms exist in a diversity of sys-

tems [Cáceres, 1997, Adler et al., 2006, Angert et al., 2009, Usinowicz et al., 2012, Chu and Adler,

2015, Ellner et al., 2016, Letten et al., 2018]. A key ingredient for the storage effect is a positive covari-

ance between favorable environmental conditions and species' densities. Temporal auto-correlations,

which are commonly observed in environmental factors [Vasseur and Yodzis, 2004, Sun et al., 2018],

can generate this positive covariance [Schreiber, 2021a].

Here, temporally auto-correlated fluctuations in predator attack rates are shown to generate non-

zero covariances between predator densities and their prey-specific attack rates. These covariances

are shown to alter ecological outcomes by mediating prey coexistence or generating stochastic priority

effects. To derive these conclusions, stochastic models of predator-prey interactions are studied using

a mixture of analytic and numerical methods.

Model and Methods

57 Following Holt and Lawton [1993], the model considers two prey species with densities N_1, N_2 that

58 are regulated by a common predator with density P. In the absence of predation, the density of prey

59 i increases by a factor, its intrinsic fitness $R_i(t)$, in generation t. The predator attack rate on prey i is

60 $a_i(t)$ in generation t. Predator immigration equals $I(t)$ in generation t and ensures the prey populations

61 are regulated [Holt and Lawton, 1993, Schreiber, 2021b]. The community dynamics are

$$N_i(t+1) = N_i(t)R_i(t)\exp(-a_i(t)P(t)) \text{ with } i = 1, 2$$

(1)

$$P(t+1) = \sum_{i=1}^{2} c_i(t)N_i(t)(1 - \exp(-a_i(t)P(t))) + I(t).$$

62 Consistent with meteorological models [Wilks and Wilby, 1999, Semenov, 2008], fluctuations in the

63 demographic rates, $R_i(t), a_i(t), I(t)$, are modeled by first-order auto-regressive processes. For example,

64 fluctuations in the predator's attack rate are modeled as $a_i(t+1) = \overline{a_i} + \rho(a_i(t) - \overline{a_i}) + \sqrt{1-\rho^2}Z_i(t)$

65 where $\overline{a_i}$ is the mean attack rate, $\rho = \text{Cor}[a_i(t), a_i(t+1)]$ is the auto-correlation in the attack rate, and

FIGURE 1. Auto-correlated attack rates have a nonlinear effect on mean predator densities. In (A), the mean predator density in the single prey and predator system is plotted a function of the auto-correlation $\rho = \text{Cor}[a_j(t), a_j(t+1)]$ in the predator attack rate. Shading of lines corresponds to the temporal variation in the attack rates. Dashed lines corresponds to where the small variance approximation (4) equals zero. In (B), the short-term versus long-term effects of an increase in the predator attack rate on the predator (top panel) and prey (bottom panel) densities.

$Z_i(t)$ are independent and identically distributed random variables with mean zero and variance σ_i^2.

The stationary distribution of this auto-regressive process has mean $\overline{a_i}$ and variance $\text{Var}[a_i] = \sigma_i^2$.

The dynamics of (1) are explored using analytical and numerical methods. The analytical meth-

ods are based invasion growth rates when rare of the prey species [Chesson, 1994, Schreiber, 2000,

Schreiber et al., 2011, Grainger et al., 2019]. These invasion growth rates in conjunction with results

from Benaïm and Schreiber [2019] allow one to determine whether each prey, in isolation, coexists with

the predator, whether the two prey species can coexist, whether one prey excludes the other prey via

apparent competition, or whether the system exhibits a stochastic priority effect i.e. with non-zero

probability either prey species is excluded. Analytical approximations for these invasion growth rates

are derived when the environmental fluctuations are small. To verify the analytical approximations

provide the qualitatively correct predictions, the invasion growth estimated numerically. The details of

the mathematical analysis are presented in the Appendix.

<center>RESULTS</center>

79 **Prey persistence.** In the absence of the other prey species, prey species j coexists with the predator

80 if its mean log fitness $\overline{\log R_j}$ exceeds the mean of the net attack rate $\overline{a_j I}$ of the immigrating predators.

81 Equivalently,

$$(2) \qquad \overline{\log R_j} > \overline{a_j} \times \overline{I} + \mathrm{Cov}[a_j, I]$$

Hence, negative covariances between prey vulnerability and predator immigration facilitate prey estab-

lishment, while positive covariances inhibit prey establishment. When this condition (2) is reversed,

the prey is driven extinct by the immigrating predators.

When the predator and prey j coexist, the mean log fitness $\overline{\log R_j}$ of the prey is balanced by the

mean of the net attack rate $\overline{a_j P_j}$ of the predator, where P_j denotes the predator's densities at the

stationary coexistence state. When the attack rates are uncorrelated in time, one has $\overline{a_j P_j} = \overline{a_j}\,\overline{P_j}$

and the mean predator density, $\overline{P_j} = \overline{\log R_j}/\overline{a_j}$, decreases with the mean attack rate. Intuitively, with

higher mean attack rates, the long-term balance between the log fitness of the prey and net predation

is only maintained by lower mean predator densities.

Auto-correlated predator attack rates generate a covariance $\mathrm{Cov}[a_j, P_j]$ between the attack rates and

the predator densities. Call this covariance, the prey vulnerability-predator density covariance. When

this covariance is positive, it increases the mean net predation, otherwise it decreases this mean i.e.

$\overline{a_j P_j} = \overline{a_j}\,\overline{P_j} + \mathrm{Cov}[a_j, P_j]$. Hence, a positive covariance $\mathrm{Cov}[a_j, P_j]$ reduces the mean predator density,

while a negative covariance increases the mean predator density. Specifically,

$$(3) \qquad \overline{P_j} = \frac{1}{\overline{a_j}}\left(\overline{\log R_j} - \mathrm{Cov}[a_j, P_j]\right).$$

96 The prey vulnerability-predator density covariance depends on the auto-correlation ρ of the attack

97 rate in a nonlinear fashion (Fig. 1A). The analytical approximation implies this covariance is propor-

98 tional to the variance of the attack rate and a quadratic function of the auto-correlation ρ:

$$(4) \qquad \mathrm{Cov}[a_j, P_j] \propto \mathrm{Var}[a_j]\rho(1 - \rho\overline{R_j})$$

FIGURE 2. Auto-correlated attack rates alter ecological outcomes. In (A), the dynamics of coexisting prey species and (B) two realizations of the dynamics prey species exhibiting a stochastic priority effect. In (C), the invasion growth rates r_i for both prey species when rare as a function of the temporal auto-correlation ρ in attack rates. Different lines correspond to different cross-correlations $\tau = -1, 0, 1$ in the attack rates on the two prey species. Positive values correspond to the prey coexisting (A) and negative values correspond to a stochastic priority effect (B). Dashed lines correspond to where the prey vulnerability-predator density covariance is zero.

where $\overline{R_j}$ is the mean fitness of prey j . The prey vulnerability-predator density covariance is positive

only if the auto-correlation is positive but not too positive, i.e. $0 < \rho < 1/\overline{R_j}$.

Figure 1B provide a graphical intuition for this non-linear relationship. When the predator's attack

rate increases, there is a short-term increase in the predator's density, but in the long-term there is a

decrease in the predator's density. The attack rates change on shorter time scales for weaker, positive

auto-correlations and longer time scales for stronger auto-correlations. Weaker auto-correlations gen-

erate a positive prey vulnerability-predator density covariance. Stronger auto-correlations, which let

the long-term dynamics play out, generate a negative covariance.

Uncorrelated attack rates and the $\overline{P_j}$-rule. When the attack rates are uncorrelated in time,

extending arguments in [Schreiber, 2021a] demonstrates that the prey species that can support the

higher mean predator density $\overline{P_j}$ excludes the other prey species.

Auto-correlated attack rates alter ecological outcomes. Auto-correlated fluctuations in attack

rates can alter the outcome of exclusion. Consider the informative case where prey only differ in

their vulnerability. In this case, the invasion growth rate of prey i equals the difference in the prey

vulnerability-predator density covariances:

$$(5) \qquad\qquad r_i = \text{Cov}[a_j, P_j] - \text{Cov}[a_i, P_j]$$

where prey $j \neq i$ is common and prey i is rare. If the common prey's vulnerability covaries more
strongly with the predator's density than the rare prey's vulnerability, then the rare species increases
from low densities and the prey species coexist (Fig. 2A). Alternatively, if rare species' vulnerability
covaries more strongly with the predator's density, then the rare species tends to decrease leading to a
stochastic priority effect: either prey species is lost with non-zero probability (Fig.2B).

A small variance approximation demonstrates that the difference in these covariances is proportional
the variance between the attack rates and a quadratic function in ρ (Fig.2C):

$$(6) \qquad\qquad \text{Var}[a_i - a_j]\rho(1 - \rho\overline{R})$$

where $\overline{R}$ is intrinsic fitness of both prey species. Provided there is differential vulnerability to the
predator (i.e. $\text{Var}[a_i - a_j]$), auto-correlations in these attack rates alter the ecological outcomes. When
this auto-correlation is positive but not too strong (i.e. less than $1/\overline{R}$ – see dashed line in Fig.2C),
the common prey's vulnerability has a greater covariance with the predator's densities, than the rare
prey. This results in a lower mean predator density and allows the rare prey to increase. When the
auto-correlation is too strong (i.e. greater than $1/\overline{R}$), the negative covariance between the common
prey's vulnerability and predator density leads to a higher predator density and, thereby, causes a
stochastic priority effect.

More generally, when the prey species differ in their intrinsic fitness R_i and differ in their mean
vulnerability, the invasion growth rate of the prey i depends on these differences:

$$(7) \qquad\qquad r_i = \overline{\log R_i} - \overline{\log R_j} + (\overline{a_j} - \overline{a_i})\overline{P_j} + \text{Cov}[a_j - a_i, P_j].$$

The first term $\overline{\log R_i} - \overline{\log R_j}$ corresponds the difference in the mean log fitness of the two prey species.
When prey species i has a larger mean log fitness, this term is positive, otherwise it is negative. The
second term, $(\overline{a_j} - \overline{a_i})\overline{P_j}$, is proportional to the difference in the mean attack rates. When the common
prey species is less vulnerable on average (i.e. $\overline{a_i} < \overline{a_j}$), this term is positive, otherwise it is negative.

The final term, $\text{Cov}[a_j - a_i, P_j]$, is equivalent to (5). However, differences in the variances leads to a more

complicated analytical approximation of this covariance term. Specifically, the term $\text{Var}[a_i - a_j]$ in (6) is

replaced by $\text{Var}[a_1 - a_2] + \text{Var}[a_j] - \text{Var}[a_i]$. For weak, positive auto-correlations, greater variation in the

common prey's vulnerability increases its invasion growth rate. Intuitively, greater variation increases

the covariance $\text{Cov}[a_j, P_j]$ that reduces the mean predator density (see (3)) and, thereby, increases the

invasion growth rate of the rare species.

141   DISCUSSION

For species regulated by a common predator, coexistence isn't expected under equilibrium conditions:

the prey supporting the higher predator density excludes the other via apparent competition [Holt and

Lawton, 1993, Schreiber, 2021b]. Similar to the storage effect for competing species [Chesson, 1994,

2018], we found that environmental fluctuations impacting prey specific attack rates can modify this

ecological outcome. Two conditions are necessary for alternative outcomes. First, fluctuating envi-

ronmental conditions must differentially impact the vulnerability of each prey species to the predator.

This condition is equivalent to "species-specific responses to environmental conditions" required for the

storage effect [Chesson, 1994]. The second condition requires a non-zero, within-generation covariance

between prey vulnerability and predator density. When positive, the prey vulnerability-predator den-

sity covariance results in a reduction of the density of the predator and facilitates the recovery of the

rarer prey species. Hence, coexistence is more likely. When the prey vulnerability-predator density

covariance is negative, each prey species tends toward extinction whenever they become rare. This re-

sults in a stochastic priority effect. This second condition is analogous to the "environment-competition

covariance" condition of the storage effect [Chesson, 1994].

The sign and magnitude of the temporal auto-correlations in predator attack rates determine the sign

of the prey vulnerability-predator density covariance. For positive auto-correlations, the sign of this co-

variance depends on the time-scale at which the fluctuations occur. When temporal auto-correlation is

weak, fluctuations occur on shorter time scales, generate a positive prey vulnerability-predator density

covariance, and promote coexistence. In contrast, when temporal auto-correlations are strong, fluctua-

tions occur over longer-time scales, generate a negative prey vulnerability-predator density covariance,

and promote stochastic priority effects. Similarly, for continuous-time models of species competing for a common resource, Li and Chesson [2016] found that fast resource depletion generates a positive environment-competition covariance and, thereby, can promote coexistence. Although not stated by Li and Chesson [2016], this positive environment-competition covariance arises from consumer attack rates being positively auto-correlated at the time scale of the resource depletion.

Taken together, the work presented here and earlier work [Li and Chesson, 2016, Schreiber, 2021a] highlight that covariances between species densities and per-capita species interaction rates can fundamentally alter the composition and dynamics of ecological communities. These covariances can be driven by the sign and magnitude of auto-correlated fluctuations in environmental conditions. Importantly, the magnitude of these auto-correlations can lead to different ecological outcomes due to differences in transient versus long-term responses of species to changing per-capita interaction rates [Bender et al., 1984, Hastings et al., 2018]. Understanding how these effects combine across multiple species to influence the composition and dynamics of ecological communities provides significant challenges for future work.

REFERENCES

P.B. Adler, J. HilleRisLambers, P.C. Kyriakidis, Q. Guan, and J.M. Levine. Climate variability has a stabilizing effect on the coexistence of prairie grasses. *Proceedings of the National Academy of Sciences*, 103:12793–12798, 2006.

A.L. Angert, T.E. Huxman, P. Chesson, and D.L. Venable. Functional tradeoffs determine species coexistence via the storage effect. *Proceedings of the National Academy of Sciences*, 106:11641–11645, 2009.

M. Benaïm and S.J. Schreiber. Persistence and extinction for stochastic ecological models with internal and external variables. *Journal of Mathematical Biology*, 79:393–431, 2019.

E. A. Bender, T. J. Case, and M. E. Gilpin. Perturbation experiments in community ecology: Theory and practice. *Ecology*, 65:1–13, 1984.

C.E. Cáceres. Temporal variation, dormancy, and coexistence: a field test of the storage effect. *Proceedings of the National Academy of Sciences*, 94:9171–9175, 1997.

E. J. Chaneton and M. B. Bonsall. Enemy-mediated apparent competition: empirical patterns and the evidence. *Oikos*, 88:380–394, 2000.

P. Chesson. Multispecies competition in variable environments. *Theoretical Population Biology*, 45: 227–276, 1994.

P. Chesson. Updates on mechanisms of maintenance of species diversity. *Journal of Ecology*, 106: 1773–1794, 2018.

P.L. Chesson. Coexistence of competitors in a stochastic environment: the storage effect. In *Population Biology*, pages 188–198. Springer, 1983.

P.L. Chesson. Interactions between environment and competition: How fluctuations mediate coexistence and competitive exclusion. In A. Hastings, editor, *Community Ecology*, pages 51–71, Berlin, Heidelberg, 1988. Springer Berlin Heidelberg.

P.L. Chesson and R.R. Warner. Environmental variability promotes coexistence in lottery competitive systems. *The American Naturalist*, 117:923–943, 1981.

C. Chu and P.B. Adler. Large niche differences emerge at the recruitment stage to stabilize grassland coexistence. *Ecological Monographs*, 85:373–392, 2015.

S.P. Ellner, R.E. Snyder, and P.B. Adler. How to quantify the temporal storage effect using simulations instead of math. *Ecology Letters*, 19:1333–1342, 2016.

L. Gibson. The role of lethal control in managing the effects of apparent competition on endangered prey species. *Wildlife Society Bulletin*, 34:1220–1224, 2006.

T.N. Grainger, J.M. Levine, and B. Gilbert. The invasion criterion: A common currency for ecological research. *Trends in ecology & evolution*, 34:925–935, 2019.

A. Hastings, K.C. Abbott, K. Cuddington, T. Francis, G. Gellner, Y.C. Lai, A. Morozov, S. Petrovskii, K. Scranton, and M.L. Zeeman. Transient phenomena in ecology. *Science*, 361:eaat6412, 2018.

R. D. Holt. Predation, apparent competition, and the structure of prey communities. *Theoretical Population Biology*, 12:197–229, 1977.

R. D. Holt and M. B. Bonsall. Apparent competition. *Annual Review of Ecology, Evolution, and Systematics*, 48:447–471, 2017.

R. D. Holt and J. H. Lawton. Apparent competition and enemy-free space in insect host-parasitoid
communities. *American Naturalist*, 142:623–645, 1993.

G.E. Hutchinson. The paradox of the plankton. *The American Naturalist*, 95:137–145, 1961.

A.D. Letten, M.K. Dhami, P. Ke, and T. Fukami. Species coexistence through simultaneous fluctuation-
dependent mechanisms. *Proceedings of the National Academy of Sciences*, 115:6745–6750, 2018.

L. Li and P. Chesson. The effects of dynamical rates on species coexistence in a variable environment:
the paradox of the plankton revisited. *The American Naturalist*, 188:E46–E58, 2016.

R.J. Morris, O.T. Lewis, and H.C.J. Godfray. Experimental evidence for apparent competition in a
tropical forest food web. *Nature*, 428:310–313, 2004.

S. J. Schreiber. Criteria for C^r robust permanence. *Journal of Differential Equations*, 162:400–426,
2000.

S. J. Schreiber. Coexistence for species sharing a predator. *Journal of Differential Equations*, 196:
209–225, 2004.

S.J. Schreiber. Positively and negatively autocorrelated environmental fluctuations have opposing
effects on species coexistence. *The American Naturalist*, 197:405–414, 2021a.

S.J. Schreiber. The P^* rule in the stochastic Holt-Lawton model of apparent competition. *Discrete &*
Continuous Dynamical Systems-B, 26:633–644, 2021b.

S.J. Schreiber, M. Benaïm, and K.A.S. Atchadé. Persistence in fluctuating environments. *Journal of*
Mathematical Biology, 62:655–683, 2011.

M.A. Semenov. Simulation of extreme weather events by a stochastic weather generator. *Climate*
Research, 35:203–212, 2008.

F. Sun, M.L. Roderick, and G.D. Farquhar. Rainfall statistics, stationarity, and climate change. *Pro-*
ceedings of the National Academy of Sciences, 115:2305–2310, 2018.

D. Tilman. Mechanisms of plant competition for nutrients: the elements of a predicitve theory of
competition. In *Perspectives on plant competition* (J. B. Grace and D. Tilman, eds.), pages 117–142,
San Diego, CA, 1990. Academic Press.

J. Usinowicz, S.J. Wright, and A.R. Ives. Coexistence in tropical forests through asynchronous variation
in annual seed production. *Ecology*, 93:2073–2084, 2012.

D. A. Vasseur and P. Yodzis. The color of environmental noise. *Ecology*, 85:1146–1152, 2004.

V. Volterra. Variations and fluctuations of the numbers of individuals in animal species living together. *ICES Journal of Marine Science*, 3:3–51, 1928.

D.S. Wilks and R.L. Wilby. The weather generation game: a review of stochastic weather models. *Progress in Physical Geography*, 23:329–357, 1999.

J. B. Wilson, E. Spijkerman, and J. Huisman. Is there really insufficient support for Tilman's R^* concept? A comment on Miller et al. *The American Naturalist*, 169:700–706, 2007.

₁ # APPENDIX FOR "TEMPORALLY AUTO-CORRELATED PREDATOR ATTACKS
₂ # STRUCTURE ECOLOGICAL COMMUNITIES"

Sebastian J. Schreiber*

Department of Evolution and Ecology

and Center for Population Biology

University of California

Davis, CA 95616, USA

₃ This appendix provides the mathematical details for the analysis reported in the main text. With

₄ the demographics rates modeled as a first-order auto-regressive processes, the full model is

$$N_i(t+1) = N_i(t)R_i(t)\exp(-a_i(t)P(t)) \text{ with } i = 1, 2$$

$$P(t+1) = \sum_{i=1}^{2} N_i(t)(1 - \exp(-a_i(t)P(t))) + I(t)$$

(A1)

$$a_i(t+1) = \overline{a}_i + \rho(a_i(t) - \overline{a}_i) + \sqrt{1-\rho^2}X_i(t)$$

$$R_i(t+1) = \overline{R}_i + \rho_R(R_i(t) - \overline{R}_i) + \sqrt{1-\rho_R^2}Y_i(t)$$

$$I(t+1) = \overline{I}_i + \rho_I(I(t) - \overline{I}) + \sqrt{1-\rho_I^2}Z(t)$$

₅ where $X_i(1), X_i(2), \ldots$ are a sequence of i.i.d. random variables with mean 0, variance σ_i^2 and cross-

₆ correlation τ, $Y_i(1), Y_i(2), \ldots$ are a sequence of i.i.d. random variables with mean 0 and variance $\sigma_{R,i}^2$,

₇ and $Z(1), Z(2), \ldots$ are a sequence of i.i.d. random variables with mean 0 and variance σ_I^2. These

₈ sequences of i.i.d. random variables are assumed to take values in a compact set. Provided the auto-

₉ correlation coefficients are strictly less than one, the processes $a_i(t), R_i(t), I(t)$ converge to a compactly

₁₀ supported stationary distribution with means $\overline{a}_i, \overline{R}_i, \overline{I}$ and variances $\sigma_i^2, \sigma_{R,i}^2, \sigma_I^2$.

₁₁ The dynamics of (A1) are analyzed in four steps. First, using results from Benaïm and Schreiber

₁₂ [2019], I determine under what conditions a single prey species can coexist with the predator. When this

₁₃ condition doesn't hold, the prey asymptotically goes extinct with probability one. For the remainder of

₁₄ the stochastic analysis, I assume that the condition for each prey species coexisting with the predator

holds. Second, I consider the dynamics of the three species model when the predator attack rates are

uncorrelated in time; the other demographic rates are allowed to be auto-correlated. Extending the

results of Schreiber [2021], I show that the prey species supporting the higher mean predator density

excludes the other prey species. Third, I use results from Benaïm and Schreiber [2019] to characterize

the general ecological outcomes using the invasion growth rates when rare. Finally, I use a diffusion

approximation [Karlin and Taylor, 1981] to derive the approximations for the invasion growth rates r_i

presented in the main text.

Persistence of a single prey species. The proof of the persistence of the single prey species, say

species j, in the absence of the other prey species closely follows the proof of Theorem 3.1 in [Schreiber,

2021]. The main difference is that [Schreiber, 2021] assumed that the demographic parameters are

independent in time i.e. $\rho = \rho_I = \rho_R = 0$. Relaxing this assumption changes the stochastic persis-

tence condition and the expression of the mean predator density. Assume $\widehat{a}_j$, $\widehat{I}$, and $\widehat{R}_j$ are random

variables whose distributions correspond to the stationary distributions of the $a_j(t)$, $R_j(t)$, and $I(t)$

auto-regressive processes. The invasion growth rate of prey j is given by

$$\mathbb{E}[\ln \widehat{\mathbb{R}}_j] - \mathbb{E}[\widehat{a}_j \widehat{I}]$$

29 where $\mathbb{E}[\cdot]$ denotes the expectation of a random variable.

30   Assume that $\mathbb{E}[\ln \widehat{\mathbb{R}}_j] - \mathbb{E}[\widehat{a}_j \widehat{I}] < 0$. As $P(t) \geq I(t-1)$ for all $t \geq 1$, it follows that for all $t \geq 1$

$$N_j(t+1) \leq R_j(t)N_j(t)\exp(-a_j(t)I(t-1))$$

$$\leq N_j(0)\prod_{s=0}^{t} R_j(s)\exp(-a_j(s)I(s-1)).$$

31 The ergodic theorem for auto-regressive processes implies that with probability one

$$\limsup_{t\to\infty} \frac{\ln N_j(t+1)}{t+1} \leq \lim_{t\to\infty} \frac{1}{t+1}\left(\ln x_1(0) + \sum_{s=0}^{t}(\ln R_1(s) - a_j(s)I(s-1))\right) = \mathbb{E}[\ln \widehat{\mathbb{R}}_j] - \mathbb{E}[\widehat{a}_j \widehat{I}] < 0.$$

Now assume $\mathbb{E}[\ln \widehat{\mathbb{R}}_j] - \mathbb{E}[\widehat{a}_j \widehat{I}] > 0$. [Benaïm and Schreiber, 2019, Theorem 1] with N_j as the only

species density and the remaining variables P, a_j, I, R_j as auxiliary variables implies that prey j coexists

with the predator in the sense of stochastic persistence.

Still assuming that $\mathbb{E}[\ln \widehat{\mathbb{R}}_j] - \mathbb{E}[\widehat{a}_j \widehat{I}] > 0$, let $(\widehat{N}_j, \widehat{P}_j, \widehat{a}_j, \widehat{I}, \widehat{R}_j)$ be a random vector that corresponds to

a stationary distribution of the model with $\widehat{N}_j > 0$ with probability one. Then, [Benaïm and Schreiber,

2019, Proposition 1] implies that the average per-capita growth rate of prey j is zero at this stationary

distribution i.e.

(A2) $$0 = \mathbb{E}[\ln \widehat{R}_j] - \mathbb{E}[\widehat{a}_j \widehat{P}_j].$$

The expressions for the mean predator density, $\overline{P}_j = \mathbb{E}[\widehat{P}_j]$, presented in the main text follow from (A2).

P^* rule when attack rates are serially uncorrelated. Assume that the attack rates are temporally

uncorrelated and both prey species can coexist with the predator i.e. $\mathbb{E}[\ln \widehat{\mathbb{R}}_j] > \mathbb{E}[\widehat{a}_j \widehat{I}]$ for $j = 1, 2$.

Without loss of generality, assume that prey 1 can maintain the higher mean predator density i.e.

$\mathbb{E}[\widehat{P}_1] = \mathbb{E}[\ln \widehat{R}_1]/\mathbb{E}[\widehat{a}_1] > \mathbb{E}[\ln \widehat{R}_2]/\mathbb{E}[\widehat{a}_2] = \mathbb{E}[\widehat{P}_2]$. The proof of the exclusion of prey 2 is similar to the

proof of Theorem 3.2 in [Schreiber, 2021]. However, [Schreiber, 2021] assumed all the demographic

parameters, not only the attack rates, are serially uncorrelated.

First, I show that prey species 1 is stochastically persists despite the presence of prey species 2.

Consider any random vector $(P_1 = 0, N_2 = \widehat{U}, P = \widehat{V}, \widehat{a}_1, \widehat{a}_2, \widehat{R}_1, \widehat{R}_2, \widehat{I})$ that corresponds to an ergodic,

stationary distribution of the model with prey 1 absent. By ergodicity, there are two cases to consider:

either $\widehat{U} = 0$ with probability one or $\widehat{U} > 0$ with probability one. In the first case, $\widehat{V} = \widehat{I}$ and the

invasion growth rate of prey 1 is given by

$$\mathbb{E}[\ln \widehat{R}_1] - \mathbb{E}[\widehat{a}_1 \widehat{I}] > 0.$$

51 In the second case, stationarity and $\widehat{U} > 0$ implies the average per-capita growth rate of prey species

52 2 equals zero i.e.

$$0 = \mathbb{E}[\ln \widehat{R}_2] - \mathbb{E}[\widehat{a}_2 \widehat{V}] = \mathbb{E}[\ln \widehat{R}_2] - \mathbb{E}[\widehat{a}_2]\mathbb{E}[\widehat{V}]$$

53 where the second equality follows from $a_2(t)$ being i.i.d. in time. Hence,

$$\mathbb{E}[\widehat{V}] = \mathbb{E}[\ln \widehat{R}_2]/\mathbb{E}[\widehat{a}_2].$$

54   On the other hand, the invasion growth rate of species 1 at this distribution equals

$$\mathbb{E}[\ln \widehat{R}_1] - \mathbb{E}[\widehat{a}_1 \widehat{V}] = \mathbb{E}[\ln \widehat{R}_1] - \mathbb{E}[\widehat{a}_1]\mathbb{E}[\ln \widehat{R}_2]/\mathbb{E}[\widehat{a}_2]$$

$$= \mathbb{E}[\widehat{a}_1]\left(\mathbb{E}[\ln \widehat{R}_1]/\mathbb{E}[\widehat{a}_1] - \mathbb{E}[\ln \widehat{R}_2]/\mathbb{E}[\widehat{a}_2]\right) > 0$$

where the last equality follows from the assumption that prey 1 can support the higher mean predator

density. [Benaïm and Schreiber, 2019, Theorem 1] treating all but N_1 as auxiliary variables implies

that prey 1 persists whether or not prey 2 is in the system.

Next I show that prey 2 is excluded whenever prey 1 is in the system. Assume $N_1(0) > 0$ and let

$W(t) = (N_1(t), N_2(t), P(t), a_1(t), a_2(t), I(t), R_1(t), R_2(t))$. Let $\Pi_t(dw)$ be the occupational measure at

time t i.e. the random measure defined by

$$\int h(w)\Pi_t(dw) = \frac{1}{t}\sum_{s=1}^{t} h(W(t))$$

61   for any continuous bounded function h. With probability one, the weak* limit points of Π_t as $t \to \infty$

62   are an invariant probability measure. Let $\widehat{W} = (\widehat{N}_1, \widehat{N}_2, \widehat{P}, \widehat{a}_1, \widehat{a}_2, \widehat{I}, \widehat{R}_1, \widehat{R}_2)$ be a random vector whose

63   distribution is given by one of these invariant probability measures. By stochastic persistence of prey

64   1, $\widehat{N}_1 > 0$ with probability one. Furthermore, by stationarity, the average per-capita growth rate of

65   prey 1 at this distribution equals zero:

$$0 = \mathbb{E}[\ln \widehat{R}_1] - \mathbb{E}[\ln \widehat{a}_1]\mathbb{E}[\ln \widehat{P}].$$

66   On the other hand, the average per-capita growth rate of prey 2 at this distribution equals

$$\mathbb{E}[\ln \widehat{R}_2] - \mathbb{E}[\widehat{a}_2\widehat{P}] = \mathbb{E}[\ln \widehat{R}_2] - \mathbb{E}[\widehat{a}_2]\mathbb{E}[\ln \widehat{R}_1]/\mathbb{E}[\widehat{a}_1]$$

$$= \mathbb{E}[\widehat{a}_2]\left(\mathbb{E}[\ln \widehat{R}_2]/\mathbb{E}[\widehat{a}_2] - \mathbb{E}[\ln \widehat{R}_1]/\mathbb{E}[\widehat{a}_1]\right) =: \beta < 0$$

67   As this holds for all weak* limit points of Π_t, it follows that

$$\lim_{t\to\infty} \frac{1}{t}\log\frac{N_2(t)}{N_2(1)} = \lim_{t\to\infty}\frac{1}{t}\sum_{s=1}^{t}(\log R_2(s) - a_2(s)P(s))$$

$$= \beta < 0$$

68 with probability one. Hence, prey species 2 goes extinct asymptotically an exponential rate.

General conditions for coexistence, bistability, and exclusion

When the attack rate is auto-correlated ecological outcomes are characterized the invasion growth

rates of both prey species. As before, assume that each prey species can persist with the predator

in isolation of the other prey species i.e. $\mathbb{E}[\ln \widehat{R}_j] > \mathbb{E}[\widehat{a}_j \widehat{I}]$ for $j = 1, 2$. Under this assumption, the

predator-prey j subsystem supports at least one stationary distribution for which $N_j > 0$ and $N_i = 0$

with probability one. Let $(\widehat{N}_j, \widehat{P}_j, \widehat{a}_1, \widehat{a}_2, \widehat{R}_1, \widehat{R}_2, \widehat{I})$ be a random vector whose distribution corresponds to

this stationary distribution. Assume this stationary distribution is unique – this happens for example,

when the environmental fluctuations are sufficiently large or when the environmental fluctuations are

sufficiently small and the predator-prey system without fluctuations has a globally stable equilibrium.

At this stationary distribution, the invasion growth rate of prey i is

$$r_i = \mathbb{E}[\ln \widehat{R}_i] - \mathbb{E}[\widehat{a}_i \widehat{P}_j]$$

79 Furthermore, at this stationary distribution, the resident prey j growth rate is zero

$$0 = \mathbb{E}[\ln \widehat{R}_j] - \mathbb{E}[\widehat{a}_j \widehat{P}_i]$$

80 Subtracting this expression from r_j and recalling that $\mathbb{E}[\widehat{a}_i \widehat{P}_i] = \mathbb{E}[\widehat{a}_i]\mathbb{E}[\widehat{P}_j] + \mathrm{Cov}[\widehat{a}_i, \widehat{P}_j]$ yields

$$r_i = \mathbb{E}[\ln \widehat{R}_i] - \mathbb{E}[\ln \widehat{R}_j] + (\mathbb{E}[\widehat{a}_j] - \mathbb{E}[\widehat{a}_i])\mathbb{E}[\widehat{P}_j] + \mathrm{Cov}[\widehat{a}_j, \widehat{P}_j] - \mathrm{Cov}[\widehat{a}_i, \widehat{P}_j]$$

as claimed in the main text.

Benaïm and Schreiber [2019, Theorem 1] implies that if $r_1 > 0, r_2 > 0$, then the prey coexist in

the sense of stochastic persistence. When $r_i < 0$ and $r_j > 0$, Benaïm and Schreiber [2019, Corollary

3] implies that species i goes exponentially quickly toward extinction when prey j is initially present.

Finally, when $r_1 < 0, r_2 < 0$, the same corollary implies that each species is excluded with positive

probability whenever the other species is present.

DIFFUSION APPROXIMATIONS FOR PREY INVASION GROWTH RATES

To approximate the invasion growth rates r_i from the previous section, consider the case of small

fluctuations in the predator attack rates. Namely, $X_i(t) = \varepsilon \eta_i(t)$ in equation (A1) where $\varepsilon > 0$ is

small, and $\eta_i(t)$ are i.i.d. with mean zero, variance v_i, and $\mathbb{E}[\eta_1(t)\eta_2(t)] = \tau$ and $\bar{a}_i = \bar{a} + \varepsilon^2 \alpha_i$.

These choices correspond to the classical diffusion scaling [see, e.g., Karlin and Taylor, 1981, Chapter

15]. Furthermore, for simplicity, assume that there are no fluctuations in the intrinsic prey fitness or

predator immigration i.e. $Y_i(t) = Z(t) = 0$ for all time t in equation (A1). As in the previous two

sections, assume each prey species can persist in the absence of the other i.e. $\ln \overline{R}_i > \bar{a}_i \overline{I}$ for $i = 1, 2$.

In the absence of the fluctuating attack rates, the dynamics of the predator with only prey j are

given by the deterministic difference equations:

(A3)
$$N_j(t+1) = N_j(t)\overline{R}_j \exp(-\bar{a}_j P(t))$$

$$P(t+1) = N_j(t)(1 - \exp(-\bar{a}_j(t)P(t))) + \overline{I}$$

97 for which the two species coexist about the equilibrium (N_j^*, P_j^*) given by

$$P_j^* = \frac{\ln \overline{R}_j}{\bar{a}_j} \text{ and } N_j^* = (P_j^* - \overline{I})/(1 - 1/\overline{R}_j).$$

98 The Jacobian matrix of (A3) at this equilibrium equals

$$\begin{pmatrix} 1 & -\bar{a}_j N_j^* \\ 1 - 1/\overline{R}_j & \bar{a}_j N_j^*/\overline{R}_j \end{pmatrix}.$$

99   For small fluctuations (i.e. small $\varepsilon > 0$), one can approximate the fluctuations

$$\begin{pmatrix} u(t) \\ v(t) \\ w(t) \end{pmatrix} = \begin{pmatrix} N_j(t) - N_j^* \\ P(t) - P_j^* \\ a_j(t) - \bar{a}_j \end{pmatrix}$$

100   around the equilibrium values as a first-order auto-regressive process:

$$
\begin{pmatrix} u(t+1) \\ v(t+1) \\ w(t+1) \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & -\bar{a}_j N_j^* & -N_j^* P_j^* \\ 1 - 1/\overline{R}_j & \bar{a}_j N_j^*/\overline{R}_j & N_j^* P_j^*/\overline{R}_j \\ 0 & 0 & \rho \end{pmatrix}}_{=:A} \begin{pmatrix} u(t) \\ v(t) \\ w(t) \end{pmatrix} + \underbrace{\begin{pmatrix} 0 \\ 0 \\ \sqrt{1-\rho^2}\xi_j(t) \end{pmatrix}}_{=:b(t)}.
$$

101   The covariance matrix C of $\begin{pmatrix} u(t) \\ v(t) \\ w(t) \end{pmatrix}$ at stationarity [see, e.g., Schreiber and Moore, 2018] satisfies

$$
\mathrm{vec(C)} = (\mathrm{Id} - A \otimes A)^{-1}\mathrm{vec(Cov[b(t)])}
$$

102   where $\mathrm{vec}(*)$ is a column vector given by concatenating the columns of its argument $*$ and $\otimes$ denotes

103   the Kroenker product. Carrying out this calculation yields the approximation

$$
\mathrm{Cov[a_j(t), P_j(t)]} \approx \mathrm{Cov[w(t), v(t)]} = \underbrace{\frac{N_j^* P_j^*}{\bar{a}_j \rho N_j^*(\rho\overline{R}_j - 1) + (1-\rho)\overline{R}_j}}_{=:\Psi} \times \sigma^2 \rho\left(1 - \rho\overline{R}_j\right).
$$

104   Next I show that $\Psi > 0$ whenever $\overline{R}_j > 1$. The definition of N_j^* implies that

$$
\Psi/\overline{R}_j = \rho\frac{\ln\overline{R}_j - \bar{a}_j\overline{I}}{\overline{R}_j - 1}(\rho\overline{R}_j - 1) + (1-\rho).
$$

105   Note that $\ln\overline{R}_j - \bar{a}_j\overline{I} > 0$ by assumption. For $\rho \le 0$, the first term of $\Psi/\overline{R}_j$ is non-negative and the

106   second term is positive. Hence, $\Psi > 0$ for $\rho \le 0$. Assume $0 < \rho < 1$ and define $f(x) = \ln x/(x-1)$. I

107   claim that $f(x) < 1$ for $x > 1$ in which case

$$
\Psi/\overline{R}_j > \rho(\rho - 1) + (1-\rho) = (1-\rho)^2 > 0.
$$

108   To prove the claim about f, l'Hopital's rule implies that

$$
\lim_{x\downarrow 1} f(x) = \lim_{x\downarrow 1} \frac{1/x}{1} = 1.
$$

109 Furthermore,

$$f'(x) = \frac{\overbrace{1 - 1/x - \ln x}^{=:g(x)}}{(x-1)^2}.$$

110 For $x > 1$, the sign of $f'(x)$ is determined by the sign of $g(x)$. As $g(1) = 0$ and $g'(x) = (1/x)(1/x-1) < 0$

111 for $x > 1$, $f'(x) < 0$ for $x > 1$. Hence, as claimed, $f(x) < 1$ for $x > 1$. Thus, as claimed in the main

112 text, $\mathrm{Cov}[a_j(t), P_j(t)]$ is proportional to $\sigma^2 \rho(1 - \rho \overline{R}_j)$.

113   REFERENCES

M. Benaïm and S.J. Schreiber. Persistence and extinction for stochastic ecological models with internal

and external variables. *Journal of Mathematical Biology*, 79:393–431, 2019.

S. Karlin and H.M. Taylor. *A second course in stochastic processes*. Academic Press, 1981.

S.J. Schreiber. The P^* rule in the stochastic Holt-Lawton model of apparent competition. *Discrete &*

Continuous Dynamical Systems-B, 26:633–644, 2021.

S.J. Schreiber and J.L. Moore. The structured demography of open populations in fluctuating environ-

ments. *Methods in Ecology and Evolution*, 9:1569–1580, 2018.