The initial response of females towards congeneric males matches the propensity to hybridise

in *Ophthalmotilapia*.

Maarten Van Steenberge[1,2], Noémie Jublier[3,4], Loïc Kéver[3], Sophie Gresham[1,5], Sofie

Derycke[1,6], Jos Snoeks[2,7], Eric Parmentier[3], Pascal Poncin[4], Erik Verheyen[1,5]

1.   Operational Directorate Taxonomy and phylogeny, Royal Belgian Institute of Natural

Sciences, 1000 Brussels, Belgium

2.   Laboratory of Biodiversity and Evolutionary Genomics, KU Leuven, Charles

Deberiotstraat 32, 3000 Leuven, Belgium

3.   Laboratory of Functional and Evolutionary Biology, FOCUS, Liège University, 4000

Liège, Belgium

4.   Laboratory of Fish and Amphibian Ethology, Behavioural Biology Unit, FOCUS, Liège

University, 4000 Liège, Belgium

5.   Department Biology, Evolutionary Ecology, University of Antwerp, 2610 Antwerp,

Belgium

6.   Flanders research institute for agriculture, fisheries and food, Animal Sciences Unit –

Aquatic Environment and Quality, Ankerstraat 1, 8400 Oostende, Belgium.

7.   Section Vertebrates, Ichthyology, Royal Museum for Central Africa, 3080 Tervuren,

Belgium

Corresponding author: Maarten Van Steenberge (mvansteenberge@naturalsciences.be)

Abstract:

Cichlid radiations often harbour closely related species with overlapping niches and

distribution ranges. Such species sometimes hybridise in nature, which raises the question

how can they coexist. This also holds for the Tanganyika mouthbrooders *Ophthalmotilapia*

ventralis and *O. nasuta*. Earlier studies found indications of asymmetrical hybridisation with

females of *O. ventralis* accepting males of *O. nasuta*, but not the other way around. We

hypothesised that this was due to differences in the capacity for species recognition. Given the

higher propensity of *O. ventralis* females towards hybridisation, we expect a reduced ability

for species recognition in *O. ventralis* females, compared to *O. nasuta* females. We staged

two experiments, one focusing on 22 female *O. nasuta* and one on 21 female *O. ventralis*.

These fish were placed in one half of a tank and briefly exposed to a conspecific or a

heterospecific male, a conspecific female, or nothing (control). Female response was

evaluated by scoring six tracking parameters and by noting the occurrence of ten discrete

behaviours before and during the encounter. Females always responded to the presence of

another fish by approaching it. Remarkably, for both *O. nasuta* and *O. ventralis*, we did not

find a different response between encounters with conspecific males and females. However,

in agreement with our hypothesis, *O. nasuta* females behaved differently towards conspecific

or heterospecific males, whereas *O. ventralis* females did not. When presented with a

heterospecific male, *O. nasuta* females performed a lower number of 'ram' behaviours.

Additionally, they never displayed the 'flee' behaviour, a component of the species' mating

repertoire that was seen in all but one of the presentations with a conspecific male. Our

findings show that differences in species recognition at first encounter predict to a large

degree the outcome of the mating process, even in the absence of mating behaviour.

Introduction

Speciation is traditionally seen as a gradual build-up of reproductive isolation between diverging populations (Mayr 1982; Coyne and Orr 2004). The classical view that this is a slow process that occurs between allopatric populations has recently been challenged by genomic findings (Marques et al. 2019) that showed how hybridisation can drive rapid speciation (Seehausen 2004; Jiggins et al. 2008). However, as unrestrained gene flow inevitably homogenizes the genomes of diverging lineages (Roux et al. 2016), the question remains what mechanisms keep incipient species separated. In scenarios of sympatric, closely related species, the ability to correctly distinguish between conspecific and heterospecific mates is probably crucial to maintain the integrity of the species (Sullivan 2009).

Mating is the end point of a complex decision-making process in which several potential mates are evaluated (Luttbeg et al. 2001). An encounter with a potential mate can be seen as the first step in this process. The outcome of this initial contact can have profound implications on fitness, either through the refusal of suitable mates, or through the acceptance of suboptimal partners. Therefore, we expect intra- and interspecific differences in individual responses when confronted with a choice of partners. Additionally, the preference for a given mate can depend on environmental, social and intrinsic parameters, explaining variation in preference both between and within species, and between and within individuals (Pfennig 2008; Sommer-Trembo et al. 2017). Assortative mate selection was traditionally seen as a sequential process in which individuals first assess whether 'the other' is a conspecific and then assess its quality as a mate (Mayr 1982). However, empirical data and theory suggest that assessment of species and quality are not independent steps (Sullivan 2009; Mendelson and Shaw 2012) as the specific status of an individual could be judged using the same cues as its quality. Additionally,

adaptive hybridisation has been observed in several taxa (overview in Mendelson and Shaw

2012), indicating that preferred mates are not necessarily always conspecific.

Cichlids of the large East African Lakes form endemic, species-rich radiations (Salzburger

2018). Several suggested "key adaptations" of cichlids, such as their pharyngeal jaws, help to

explain their evolution into numerous trophic niches (Kocher 2004). However, a large

proportion of these closely related species coexist without apparent eco-morphological

differences (Van Oppen et al. 1998). Since most cichlid assemblages are relatively young,

several taxa may be classified as incipient species and they retain the potential to hybridise.

The oldest East African lake, Lake Tanganyika, however, contains a mature cichlid radiation

(Salzburger 2018) in which most species are well-delineated (Ronco et al. 2020). However,

even between well-delineated biological species, boundaries can be permeable and molecular

studies identified several instances of inter-specific hybridisation (Rüber et al. 2001;

Koblmüller et al. 2007; Nevado et al. 2011). Such examples allowed us to select a case to study

the importance of prezygotic, behavioural isolation after, or at the last stages, of the speciation

process.

Ophthalmotilapia Pellegrin, 1904 species are maternal mouth brooders that occur on the rocky

and intermediate (rocky patches separated by sand) shores of Lake Tanganyika. The genus

contains four currently accepted valid species: *O. ventralis* (Boulenger 1898), *O. boops*

(Boulenger 1901), *O. heterodonta* (Poll and Matthes 1962) and *O. nasuta* (Poll and Matthes

1962) (Hanssens et al. 1999). They are sexually dimorphic maternal mouthbrooders with

territorial males that protect a spawning site, and females that aggregate in feeding schools

when they are not breeding. *Ophthalmotilapia* males possess egg-shaped lappets at the distal

ends of their elongated pelvic fins that are unique among Great Lake cichlids (Poll 1986). These

lappets function as egg dummies during the species' mating behaviour in a similar way as the

egg spots on the anal fins of the so-called 'modern' haplochromines (sensu Salzburger et al.

2007; Theis et al. 2012). During the mating process, the female deposits the eggs and almost

immediately takes them into her mouth. By snapping at the egg dummies, which are situated

close to the genital opening of the male, the intake of sperm is thought to be facilitated,

increasing the fertilisation rate of the eggs within the female's mouth (Salzburger et al. 2007).

The four species of *Ophthalmotilapia* have different but partially overlapping, distribution

ranges. *Ophthalmotilapia nasuta* is the sole species in the genus with a patchy but lake-wide

distribution. The sister species *O. heterodonta* and *O. ventralis* have non-overlapping ranges

with the former occurring in the northern half and the latter in the southern third of the Lake.

The fourth species, *O. boops* only occurs along a rather limited stretch of Lake Tanganyika's

south-eastern shoreline. There, it prefers sites where large stones are available (Konings 2019).

This is the only part of the lake where up to three species of *Ophthalmotilapia* occur in sympatry

(Hanssens et al. 1999).

Although specimens of *Ophthalmotilapia* can be easily assigned to one of the valid species

(with the possible exception of *O. heterodonta* and *O. ventralis*, see Hanssens et al. 1999), a

phylogeographic study discovered gene flow among these species. Nevado et al. (2011)

observed that specimens of *O. nasuta* often carried mitochondrial DNA of the other species,

whereas the opposite was much less often the case. They suggested that this pattern either has

a postzygotic, (*e.g.* by cyto-nuclear incompatibilities that affects mutual crossbreedings

differently) or a prezygotic (*e.g.* by an asymmetry in reproductive behaviour that results in a

different resistance towards hybridisation) cause. The latter scenario implies that females of all

species would occasionally mate with *O. nasuta* males, while *O. nasuta* females would be much

less inclined to mate with heterospecific males. It also implies that the female hybrid offspring

would backcross into *O. nasuta*. This scenario agrees with the recent description of a successful

mating between a female *O. ventralis* and a male *O. nasuta* (Kéver et al. 2018). Reproductive

isolation in closely related species of East African cichlids is mostly maintained through

prezygotic isolation (Turner et al. 2001). Hence, models that describe speciation in cichlids

emphasize the importance of female mate choice in the initial stages of the speciation process

(Danley and Kocher 2001).

The Lake Tanganyika cichlids assemblage contains species with profoundly different mating

strategies (Ronco et al. 2020). *Ophthalmotilapia* stands out by its extreme sexual dimorphism

and female-biased reproductive investment. Although the correlation between reproductive

investment, sexual selection and choosiness is well-established, it still remains debated whether

choosiness is an evolutionary outcome (sensu Trivers 1972), or rather a determinant of

differences between the sexes in parental investment (Thomas and Szekely 2005). Using Lake

Tanganyika cichlids, Gonzales-Voyer et al. (2008) showed support for the latter hypothesis.

Regardless of the evolutionary mechanism, females should be considered the choosy sex in

Ophthalmotilapia (sensu Wirtz, 1999). Therefore, if a prezygotic mechanism explains the

asymmetric pattern observed in nature (Nevado et al. 2011), it would be caused by differences

between females of the different species in accepting matings with heterospecific males. As

increased capacity for species recognition leads to increased preference in the choosier sex

(Kozak & Boughman, 2009), we predict to see an interspecific difference in female response

to conspecific and heterospecific males. As males of the different *Ophthalmotilapia* species

have very similar courtship behaviours, in which the few species-specific elements are

insufficient to prevent hybridisation (Kéver et al. 2018), females would mainly rely on other

cues like colour patterns, body size and pheromones. Although the reproductive behaviour of

Ophthalmotilapia species is well documented (Haesler et al. 2011; Immler and Taborsky 2009;

Kéver et al. 2018), little is known on how *Ophthalmotilapia* species recognize conspecifics and

select potential mates.

In this study, we will investigate species recognition, which was, in spite of its shortcomings

(Mendelson and Shaw 2012), defined as ''a measurable difference in behavioural response

toward conspecifics as compared to heterospecifics''. We will study this by comparing the

ability of female *O. nasuta* and *O. ventralis* to distinguish con- from heterospecific males. Often,

dichotomous mate choice trails, in which a female can choose between two males, are used for

such studies. Such a setup includes an aspect of male competition, whereas we are interested in

the response of females in the absence of such behaviour. Additionally, with dichotomous trails

one cannot discriminate mate preference from species recognition. Although we expect species

recognition to be a good predictor of female mate preference, we did not aim to quantify

preference directly. Instead, we choose to investigate differences in species recognition by

presenting a set of *O. ventralis* and *O. nasuta* females to males of both species separately and

by measuring a range of behavioural variables.

This study described and compared the initial behavioural response of *O. ventralis* and *O.*

nasuta females towards males of both species in an aquarium setting. In view of the pattern of

asymmetric hybridisation discovered among these species (Nevado et al. 2011), we expected a

difference in the capacity for species recognition between females of *O. ventralis* and *O. nasuta*.

Specifically, we hypothesized that *O. nasuta* females would be able to differentiate between

conspecific and heterospecific males at the initial stages of an encounter. For *O. ventralis*

females, however, we expected that this capacity would be less pronounced or absent.

Materials and Methods

Experimental setting

We performed two independent experiments using females and males of *O. ventralis* and *O. nasuta*. The first focused on the behaviour of focal *O. nasuta* females (ON experiment), the second on that of focal *O. ventralis* females (OV experiment). All individuals were wild caught off the coast of Ulwile Island or nearby Kala, on the mainland (Tanzania). Specimens were acquired as juveniles and hence had no prior mating experience. They were kept in monospecific aquaria per species until they reached maturity. Hence, all females used had similar ages. Males were older and most had prior mating experience (Kéver et al. 2018). We verified the origin of a random selection of these fishes by sequencing the mitochondrial control region, and by comparing these sequences with the data collected by Nevado et al. (2011). The experiments were performed at the aquarium facilities of the University of Liège. Prior to the onset of the experimental trials, the sex of the specimens was checked by visually inspecting their genital papillae. Female specimens were kept jointly but isolated from males and heterospecific specimens in a separate tank for at least two weeks. This tank was devoid of hiding places, in order to prevent the development of territoriality. During that period, males were kept in monospecific tanks in which they were visually isolated from each other using opaque partitions. We kept all specimens in the same condition for at least two weeks with photoperiod: 12:12 h L:D, water temperature: 26±1°C, carbonate hardness: >8 dKH. Fishes were fed daily ad libitum with 'Tropical Spirulina forte' mini-granules.

We used three identical experimental aquaria (88cm*50cm*40cm with water level ca. 40cm), which we divided into two equal parts by a perforated transparent partition (separation wall), through which fishes could not pass, and by an opaque wall (visual barrier) that could be

removed (Fig. 1A). A flower pot was placed on each side of the separation to allow the fish to

take refuge. We kept the fishes in these aquaria for at least twelve hours before they were used

in the experimental trials. During the trial, the visual barrier was removed.

Figure 1. Experimental setup. A: aquarium setup: A focal female of either *O. nasuta* or *O. ventralis* (here *O.*

nasuta) (A) was placed in one half of the experimental tank whereas no fish, a conspecific female or a hetero- or

conspecific male was placed in the other half (here an *O. ventralis* male) (B). The tank was divided in two by a

transparent wall (D) and a visual barrier (C), a microphone (E) was placed on the side of the non-focal specimen

and an empty flowering pot was placed in both halves of the tank, allowing fishes to take refuge. Video and audio

recordings were made 15 minutes prior and 45 minutes after the visual barriers were removed. B: Contrasts tested

using permanova: A: focal females presented with another fish *vs*. with no fish, B: focal females presented with a

conspecific female *vs*. a conspecific male, C: focal females presented with a conspecific *vs*. a heterospecific male,

D: conspecific females and males presented to a focal female and E: conspecific and heterospecific presented to a

focal female. Black and grey fishes represent different species, the white fish represents all possible non-focal

specimens used.

We recorded the behaviour of focal specimens (*O. nasuta* females or *O. ventralis* females) in

four different experimental conditions. They were either exposed to (i) no other specimen (Co),

(ii) a conspecific female (CF), (iii) a conspecific male (CM) or (iv) a heterospecific male (HM)

(Supplement 1). For each experiment (ON and OV) and for each experimental condition (i to

iv), we conducted a minimum of five replicates. We filmed (using a CANON Legria HF R606)

the entire aquarium (i.e. focal and non-focal fishes) during one hour: from 15 minutes before to

45 minutes after the visual barrier was removed. Experimenters were only briefly present in the

room to remove the visual barrier. As *O. ventralis* males are known to produce weak-pulsed

sounds during the inviting behaviour (Kéver et al. 2018), we used a HTI Min-96 hydrophone

(−164.4 dB re. 1 V μPa−1; bandwidth 2 Hz and 30 kHz, MS, USA), connected to a Tascam

DR-05 recording (TEAC, Wiesbaden, Germany) at a 44.1 kHz sampling rate to record sounds

during the whole experiment. The hydrophone was positioned near the separation wall, at half

the height of the water column, on the side of the non-focal specimen. At the start of each trial,

we switched off the aeration of the tank so that sounds could be recorded. However, these

recordings were not analysed, as we detected no communication sounds. After each trial, the

focal female was euthanized. Both the focal and the non-focal specimens were weighed. Focal

specimens were measured, dissected and the stage of gonad development was scored following

Panfili et al. (2006).

We performed a total of 28 ON and 21 OV experimental trials, with a maximum of three trials

per day (Supplement 1). However, after the dissections (see below), we observed that six focal

O. nasuta females from the first set of trials possessed male or ambiguous gonads (Supplement

2). These specimens were referred to as floater males and the recordings for these trials were

not analysed. As we suspected that these specimens had changed sex, we photographed the

genital papillae of the focal females that were to be used subsequently, two weeks before the

onset of the experimental trials. A comparison between papillae of the same individuals after

two weeks revealed that some could be identified as males. This either confirmed that a sex

change did indeed take place in several specimens, or showed that the examination of genital

papillae was insufficient for sex determination in subadult *Ophthalmotilapia*. These specimens

were not studied. After each trial, the aquarium was cleaned and the water fully renewed.

Collection of tracking and qualitative behavioural data

Video files were converted into JPG images using Adapter v2.1.6 (available at

https://www.macroplant.com), capturing one frame per second and saving it as an 8-bit, gray-

scale JPG file. Images taken within ten seconds before or after an experimenter was performing

an action (i.e. removing the wall) were discarded from analyses. We chose to analyse the same

number of frames for all trials within the ON and the OV experiment, respectively. For the ON

experiment, this resulted in a minimum of 721 and 2186 frames collected before and after the

removal of the separation wall, respectively. For the OV experiment, 871 and 2685 frames were

available for analyses. Both focal and non-focal specimens were tracked using the ImageJ v1.49

(Schneider et al. 2012) software package.

Given the presence of both light- and dark- coloured backgrounds in the aquarium setting, the

set of frames was studied twice. Specimens that were present before a light coloured

background were tracked by inverting black and white values whereas specimens present before

a dark-coloured background were tracked using non-inverted images. For computational

reasons, analyses were performed on subsets of the data containing a maximum of 1,000 frames.

For each set of images, a subset of 30 frames was used to create a background using the plug-

in '*Image stack merger plus*'. Backgrounds were removed using the image calculator and the

resulting frames were transformed into black-and-white images using the threshold function

with '*MaxEntropy*' as the methodology. Images were adjusted using the '*erode*' and '*dilate*'

functions to remove noise and to obtain a better representation of the fishes. The resulting image

series was then used for tracking using the plug-in '*MTrack2*', in which tracks were summarized

as the x- and y- coordinates of the centroids of the tracked object. The quality of the automated

tracking was checked by visually inspecting each of the frames. When the software failed to248 *Collection of tracking and qualitative behavioural data*

Video files were converted into JPG images using Adapter v2.1.6 (available at

https://www.macroplant.com), capturing one frame per second and saving it as an 8-bit, gray-

scale JPG file. Images taken within ten seconds before or after an experimenter was performing

an action (i.e. removing the wall) were discarded from analyses. We chose to analyse the same

number of frames for all trials within the ON and the OV experiment, respectively. For the ON

experiment, this resulted in a minimum of 721 and 2186 frames collected before and after the

removal of the separation wall, respectively. For the OV experiment, 871 and 2685 frames were

available for analyses. Both focal and non-focal specimens were tracked using the ImageJ v1.49

(Schneider et al. 2012) software package.

Given the presence of both light- and dark- coloured backgrounds in the aquarium setting, the

set of frames was studied twice. Specimens that were present before a light coloured

background were tracked by inverting black and white values whereas specimens present before

a dark-coloured background were tracked using non-inverted images. For computational

reasons, analyses were performed on subsets of the data containing a maximum of 1,000 frames.

For each set of images, a subset of 30 frames was used to create a background using the plug-

in '*Image stack merger plus*'. Backgrounds were removed using the image calculator and the

resulting frames were transformed into black-and-white images using the threshold function

with '*MaxEntropy*' as the methodology. Images were adjusted using the '*erode*' and '*dilate*'

functions to remove noise and to obtain a better representation of the fishes. The resulting image

series was then used for tracking using the plug-in '*MTrack2*', in which tracks were summarized

as the x- and y- coordinates of the centroids of the tracked object. The quality of the automated

tracking was checked by visually inspecting each of the frames. When the software failed to

track a specimen that was clearly present in the final images, coordinates were added manually.

Finally, tracks obtained from both datasets, inverted and non-inverted, were combined. When

a specimen was recognized by both methods, *e.g.* when the fish was partially before a light-

and partially before a dark-coloured background, the average of the coordinates was used.

When tracking data was missing, the average value of the coordinates of the previous and the

next positions were used. This is justified, as missing data either corresponded to fish that

remained stationary for many frames, and could hence not be distinguished from the

background, or to fish that hid behind the terracotta flowerpots (Fig. 1). Frames collected before

and after the removal of the visual barrier were analysed separately. Coordinates were shifted

using the lower- and anterior-most point of the separation wall as the origin, and rotated by

setting the anterior water level as a reference for the horizontal plane. Finally, all coordinates

were transformed from pixels to centimetres using the dimensions of the aquaria. Tracks were

visualized by plotting all individual positions as well as the shift in average position of a

specimen before and after the removal of the barrier (Supplement 3).

For each specimen, six tracking parameters were calculated from the coordinates (Table 1).

Each parameter was calculated three times: once using coordinates obtained for 721/871

(OV/ON experiment respectively) seconds before (before) the removal of the visual barrier,

once using coordinates obtained during 721/871 seconds (after1) after the visual barrier was

removed and finally using coordinates obtained during 2186/2685 seconds (after2) after the

visual barrier was removed, hence including the after1. Additionally, ten specific behaviours

were defined based on Baerends and Baerends-Van Roon (1950). These were encoded and

recorded as point events in Boris v. 2.72 open source software (Friard and Gamba 2016) (Table

1). This data was collected during the same three periods: before, after1 and after2. We choose

these time periods, as we aimed to study the immediate response following presentation to

another fish. Teleosts respond to changing social situations at two scales, by an immediate

change in behaviour, and by a modification of neural circuits (Oliveira 2012). The latter is

driven by the regulation of immediate early genes (IEG). Following Oliveira (2012), we

assumed that the transcription of IEG occurs roughly 15 min (after1), and their translation 45

min (after2) after the stimulus. Behaviours displayed within ten seconds before or after an

experimenter was performing an action (removing the wall) were discarded from the analyses.

Table 1. Tracking parameters and point events recorded for the focal and non-focal individuals during the

experimental trials. For point events, interpretation of the behaviour was added.

Tracking parameters	Description	
Dist.wall	Distance to the visual barrier (% of length of compartment).	
Dist.fish	Distance to the fish on the other side of the transparent wall (cm).	
Sp	Average speed (cm/s).	
SpX	Horizontal speed (cm/s).	
SpY	Vertical speed (cm/s).	
Height	Mean height (% of height of water column).	
Point events	**Description**	**Interpretation**
Chase	The fish suddenly swims very fast towards the other fish and rams (or almost rams) the separation wall.	Contextual: agonistic and male courtship behaviour (Kéver et al. 2018)
Flee	The fish suddenly swims away from the other fish.	Contextual: agonistic and female courtship behaviour (Kéver et al. 2018)
Lateral	Lateral display: The fish positions itself perpendicular to the other fish, keeping its head slightly downwards, erects its fins and bends its body.	Signal movement (Baerends and Baerends-Van Roon 1950), agonistic behaviour (Kéver et al. 2017)
Frontal	Frontal display: The fish faces the other fish head up and erects its fins.	Signal movement (Baerends and Baerends-Van Roon 1950), agonistic behaviour (Kéver et al. 2018)
Bite	Biting the wall: The fish bites the separation wall (possibly trying to bite the other fish).	Signal movement (Baerends and Baerends-Van Roon 1950), agonistic behaviour (Kéver et al. 2018)
Ram	Ram into the wall: The fish tries to enter the other part of the aquarium and rams (not very fast) the separation wall.	
Sand	Sand picking: The fish takes sand in its mouth.	Courtship behaviour when linked to construction of bower (Kéver et al. 2018), signal movement when nipping off a substrate (Baerends and Baerends-Van Roon 1950).
Spasm	A quick, strong, and unilateral contraction of the trunk musculature that results in a displacement of the head and the caudal fin in the same direction.	Contextual: comfort behaviour or signal movement. Observed in courtship behaviour or inter-territorial fights depending on the genus (Baerends and Baerends-Van Roon 1950).

Tail	Tail-wagging: Exaggerated movements of the caudal fin (+ caudal part of the dorsal fin). At its zenith, the movement of the caudal fin is completely counterbalanced by backpedalling.	Signal movement (Baerends and Baerends-Van Roon 1950), courtship behaviour (Kéver et al. 2018).
Flicker	Pelvics flickering: The fish quickly and alternatively moves its right and left pelvic fins.	Comfort behaviour (Baerends and Baerends-Van Roon 1950)

Statistical analyses

Prior to testing differences in species recognition, we visually explored the combined datasets of tracking parameters and point events. We did this by performing principal component analysis (PCA) and canonical variate analyses (CVA) in Past 3.14 (Hammer et al. 2001). The former allowed for an unbiased visualization of the variation in the data and was performed on the correlation matrices. The latter was conducted to maximize the differentiation between the different groups. Separate analyses were performed for each period of time recorded (before, after1, after2) and for each experiment (ON or OV). Point events that were not recorded during one of these periods were disregarded and missing values (i.e. for Dist.fish in the control condition Co) were treated using mean value imputation. Prior to the analyses, each of the tracking parameters and point events was normalised. This was done for each experiment (OV and ON), and for each of the time periods ('before', 'after1', 'after2' and 'shift') separately.

We used permanova to compare the behaviour of both focal and non-focal specimens in five different comparisons (Fig. 1B). With this non-parametric test, we could include all recorded behaviours. This allowed us to measure species recognition via differences in behaviour (sensu Mendelson and Shaw 2012) without defining *a priori* in what variable specimens would differ. In order to reduce the number of comparisons, we restricted ourselves to only biologically relevant contrasts. For focal specimens, we compared the behaviour between (A) females that were presented with another fish *vs*. with no fish, (B) focal females that were presented with a conspecific female *vs*. a conspecific male, and (C) females that were presented with a

conspecific *vs*. a heterospecific male. Two additional comparisons were tested for the non-focal

individuals. We tested (D) whether conspecific females and males respond differently to a focal

female and (E) whether conspecific and heterospecific males respond differently to a focal

female. Even though our main goal was to test whether females of *O. nasuta* and *O. ventralis*

differed in behavioural response towards conspecific and heterospecific males (C), we tested

the four other comparisons as well, following the recommendations of Moran (2003) and

Nakagawa (2004).

Tests were performed using non-parametric permanova, using the *pairwise.adonis* function, of

the R package *vegan* on the combined data of tracking parameters and point events. This

approach was chosen as the conditions for multivariate normality were violated. When

permanova revealed significant differences, we verified whether this could be due to dispersion

effects (Anderson 2006). For this, in view of the size of the dataset, non-parametric Mann-

Whitney U tests were performed on the within group dispersions from the mean, calculated

using the *betadisper* function implemented in the R package *vegan* (Oksanen et al. 2017). A

difference in dispersion across groups does, however, not contradict a difference in their

average values. Hence, the data was visually explored, using PCA, to assess whether dispersion

could be present together with a difference in average values. For each comparison revealed

significant by permanova, Mann Whitney U non-parametrical tests were performed on each of

the variables separately in order to detect which of these caused the difference between the

treatments. We choose this non-parametric approach as the assumptions of normality were often

not met. When significant, the effect sizes of these variables were estimated using Hedge's g,

which was calculated using the estimationstats.com web application (Ho et al. 2019).

In order to test whether the observed differences in behaviour depended on the visual presence

of another specimen, and whether these differences were already visible at the first stages of

the encounter, permanova tests were performed on data collected before the removal of the

visual separation (before) as well as on data collected over a short (after1) and a long (after2)

period of time after this separation was removed. Finally, an additional test was conducted

which removed individual variation between the different treatments. For this, behavioural

shifts were calculated for each tracking parameter and point event by subtracting the values of

the 'before' period before from those of the 'after1' period (shift). All tests were performed

separately for the ON and the OV experiment. As behaviour can be influenced by gonad

development and weight of the focal and non-focal specimens, Mann Whitney U tests were

performed to check whether these differed between the treatments. Such tests were also

performed on the amount of frames in which fishes could not be tracked. All statistical analyses

were performed using Past 3.14 (Hammer et al. 2001) and R (R core team 2017).

Results

We separately analysed two experiments, one focusing on *O. nasuta* and one on *O. ventralis*

females (ON and OV experiment). In the ON experiment, two males of *O. ventralis* performed

advanced courtship behaviours. After the encounter, these males started to swim in circles, in

fast and erratic movements. This was accompanied by tail wagging, generally displayed close

to the partition wall. These males often bit the hydrophone and picked up and moved around

sand (49 and 45 times within 45 min *vs.* 0 for the other males). One of these two males (*O.*

ventralis male presented to ON38) also tried to chase the female (79 times) and presented the

egg dummies of its pelvic fins (5 times). This behaviour stopped immediately when the

experimenter removed the female fish. During the encounter, these males turned dark grey, to

almost black, which was swiftly reversed after the experimental trial. As we designed our experiment to study behavioural response in the absence of courtship behaviour, we removed these outliers from all analyses. All ten point events were observed in at least one of the fishes in the ON experiment, whereas 'tail' (*i.e.* tail wagging) was never observed in the OV experiment (Supplement 4).

Visualization of the behavioural data

We visually explored the data using PCA and CVA to compare the behaviour of all specimens used in each of the two experiments. In the PCAs conducted on the behavioural data collected before the removal of the barrier, values of all females as well as of conspecific males overlapped (Fig. 2A,B), suggesting a highly similar behaviour. However, heterospecific males were (somewhat) separated from all other specimens by their higher values for PC1 (ON experiment) or PC2 (OV experiment). The loadings of the main PCs (Supplement 5.1) suggest that this difference was due to a more active swimming behaviour (Sp, SpX, SpY) higher up in the water column (height) for *O. ventralis* males (ON experiment) and a higher number of point events (ram, sand, bite) performed at the floor of the aquarium (height) for the *O. nasuta* males (OV experiment), prior to their presentation to a heterospecific female. The fact that *O. nasuta* males build true bowers could explain the higher number of point events, especially 'sand'. Additionally, *O. nasuta* might be less at ease close to the water surface as it generally occurs in deeper waters than *O. ventralis* (Konings 2019). The CVAs also reflected the behavioural differences of heterospecific males (Fig. 3A,B), as they had higher values for the first CVs. The behaviours that contributed strongly to the separating PCs, also contributed to the main CVs (Supplement 5.2).

The PCAs performed on the data collected 15 minutes after the removal of the barrier foremost

showed how the focal females that were used as controls behaved differently than the specimens

that were presented with another specimen (Fig. 2C,D). For both experiments, this can be

explained by control females spending less time closer to the wall (Dist.wall), and performing

less agonistic behaviour (ram, lateral, flee). We did not observe any additional separation in the

PCA of the OV experiment (Fig. 2D). In the ON experiment, values for (heterospecific) *O.*

ventralis males stood out by their high values for PC1, whereas those of (conspecific) *O. nasuta*

males had mostly low values for this axis (Fig. 2C). The tracking parameters Sp, SpX, SpY all

had high, positive contributions to PC1 (Supplement 5), reflecting that *O. ventralis* males kept

swimming actively when the barrier was removed in the ON experiment.

We carried out CVAs on the same datasets (Fig. 3C,D). For both experiments, the control

females stood out by their high values for CV1. This could again be explained by their higher

values for Dist.Wall. In the ON experiment, (heterospecific) *O. ventralis* males stood out by

their low values for CV1, which would be attributed to their more active swimming behaviour

(Sp, SpX, SpY). Conspecific *O. nasuta* males stood out by their low values for CV2, which

could be due to the higher occurrence of 'sand' and 'bite' behaviour. Values for *O. nasuta*

females that were presented with another fish had more intermediate values for CV1 and CV2.

However, values of *O. nasuta* females that were presented to a conspecific male clustered

between values of those males and of those of the other females. Similarly, females that were

presented to a heterospecific male had values that were intermediate between those of these

males and those of the other females (Fig. 3C). This suggests that the behaviour of focal females

shares characteristics with the behaviour of the non-focal fishes presented to them. In the CVA

of the OV experiment, (heterospecific) *O. nasuta* males stood out by their low values for CV1

and high values for CV2. This was most influenced by the higher occurrence of point

behaviours (ram, spasm, sand, flicker). Values of *O. ventralis* males overlapped with those of

female specimens that were presented with another fish (Fig. 3D). These patterns remained

present when performing similar analyses on the data collected 45 minutes after the removal of

the visual barrier (Supplement 6). Plotting the shift in average position before and after the

removal of the barrier revealed how almost all specimens moved towards the wall when

presented with another specimen. Additionally, this showed that *O. nasuta* specimens, on

average, spent more time closer to the bottom whereas *O. ventralis* specimens were more often

found higher up in the water column (Supplement 7).

Figure 2. Principal component analyses performed on the behavioural data collected 15 min before (A, B) and 15 min after (C, D) the visual barrier was removed in the ON (*O. nasuta*, left) and the OV (*O. ventralis*, right) experiments. Symbols on the scatter plots for the ON and OV experiment as in E and F, respectively, with full circles denoting focal females presented with no fish (red), a conspecific female (blue) an *O. ventralis* male (purple), and an *O. nasuta* male (green), empty circles denote non-focal conspecific females and full squares *O. ventralis* (purple) and *O. nasuta* (green) males. Explained variances are added to the axes.

Figure 3. Canonical variate analyses on the behavioural data collected 15 min before and 15 min after the visual barrier was removed in the ON (*O. nasuta*, left) and the OV (*O. ventralis*, right) experiment. Symbols on the scatter plots for the ON and OV experiments as in E and F, respectively, with full circles denoting focal females presented with no fish (red), a conspecific female (blue) an *O. ventralis* male (purple), and an *O. nasuta* male (green), empty circles denote non-focal conspecific females and full squares *O. ventralis* (purple) and *O. nasuta* (green) males. Explained variances are added to the axes.

Behaviour of focal females

We tested whether the behavioural responses of females of *O. nasuta* and *O. ventralis* differed

in three different comparisons (Fig. 1B), i.e. when they were presented to (A) another fish *vs.*

no fish, (B) a conspecific female *vs.* a conspecific male, and (C) a conspecific *vs.* a

heterospecific male. Within each of the comparisons in the ON experiment, we detected no

significant difference in the gonad development of the focal females, in their weights, in the

weights of the non-focal fishes, and in the percentage of missing frames. The same applies for

the OV experiment, although here, non-focal females had lower body weights than conspecific

males. Prior to the removal of the visual barrier, no significant difference in behaviour was

recorded for focal females from the different treatments in both experiments, and for all three

comparisons (Table 2).

For the three comparisons, and for both experiments (ON and OV), we performed permanova

on the behavioural data recorded during the 15 minutes after the removal of the visual barrier

(Table 2). For both the OV and the ON experiment, this revealed a significant difference in the

behaviour of focal females that were not presented to another fish (controls) and focal females

that were presented with another fish (comparison A). For the ON experiment, we could not

exclude that these differences stem from dispersion effects. However, these dispersion effects

can be explained by the (much) lower variation in values for control females compared to those

of other focal females. As visualisation of the data showed a clear separation of both groups,

we can assume that a difference in dispersion is present together with a difference in average

values (Fig. 2C). Mann-Whitney U tests revealed that controls differed from other focal females

by their higher values for the variable Dist.wall (ON: 46.3 +-18.2 *vs.* 20.1 +-8.8, p=0.005,

g=1.32; OV: 55.2 +-13.3 *vs*. 20.8 +-9.7, p=0.002, g=1.13) and their lower values for 'ram' (ON:

8.6+-12.1 *vs*. 74.5+-54.0, p=0.005, g=-2.14; OV: 8.0 +-11.9 *vs*. 48.8 +-38.4, p=0.014, g=-3.12).

Table 2. PERMANOVA performed on the behavioural parameters of the ON (*O. nasuta*) and OV (*O.

***ventralis*) experiment.** Tests were performed on the data collected during 15 minutes before (B), 15 minutes after

(A1) and 45 minutes after (A2) the removal of the opaque wall as well as on behavioural shifts (S) calculated by

subtracting data recorded during 15 minutes before from that of 15 min after the removal of the wall (A1-B). We

tested five comparisons (Fig. 1B): by comparing the behaviour of focal females that were presented with (A)

another fish *vs*. with nothing, (B) with a conspecific female *vs*. a conspecific male, and (C) with a conspecific *vs*.

a heterospecific male. We further compared (D) the behaviour of non-focal conspecific males vs. females, and (E)

conspecific *vs*. heterospecific males. Behaviours that were neither recorded before or after the removal of the wall,

'tail' and 'sand' for the ON, and 'flicker' and 'sand' for the OV experiment, were excluded. For the first

comparison, Dist.fish was excluded, as it could not be calculated. Values in bold are significant at the 0.05 level.

For these comparisons, ° and † denote that the assumption of equal dispersion was violated at the 0.05 and 0.01

levels.

	C	A	B	C	D	E
ON	B	F:0.31; p: 0.789	F:0.95; p: 0.423	F:0.91; p: 0.474	F:1.17; p: 0.339	F:1.15; p: 0.24
	A1	**F:7.28; p: 0.012** †	F:0.42; p: 0.768	**F:11.33; p: 0.016** °	F:0.45; p: 0.651	F:0.55; p: 0.73
	A2	**F:4.71; p: 0.027** †	F:0.49; p: 0.553	**F:8.13; p: 0.040**	F:0.66; p: 0.545	F:0.85; p: 0.47
	S	**F:8.40; p: 0.006** †	F:1.03; p: 0.416	**F:8.75; p: 0.018** †	F:1.18; p: 0.332	F:0.25; p: 0.93
OV	B	F:0.23; p: 0.762	F:1.62; p: 0.203	F:0.85; p: 0.387	F:3.19; p: 0.061	**F:5.13; p: 0.013**
	A1	**F:8.00; p: 0.005**	F:0.91; p: 0.386	F:0.40; p: 0.546	F:0.28; p: 0.633	F:0.48; p: 0.524
	A2	**F:4.28; p: 0.026**	F:0.78; p: 0.438	F:0.48; p: 0.610	F:0.09; p: 0.837	F:0.49; p: 0.525
	S	**F:7.61; p: 0.004**	F:0.89; p: 0.418	F:0.87; p: 0.407	F:0.42; p: 0.679	F:0.73; p: 0.447

Unexpectedly, in both experiments, we did not observe a difference in behaviour between focal

females that were presented with a conspecific female or a conspecific male (comparison B).

However, when comparing the behaviour of focal females presented with conspecific and

heterospecific males (comparison C), a difference became evident between the ON and the OV

experiment. In support of our hypothesis, females of *O. nasuta* responded differently towards

conspecific and heterospecific males, whereas females of *O. ventralis* did not (Table 2). Mann-

Whitney U tests revealed that this was due to the lower number of observed 'ram' behaviours

(43.8+-17.9 *vs*. 138+-52.9; p=0.04, g=2.25) in *O. nasuta* females that were presented to *O.*

nasuta males compared to those presented to *O. ventralis* males. Additionally, *O. nasuta*

females never performed a 'flee' behaviour when presented to an *O. ventralis* male, whereas

this was observed in all but one of the *O. nasuta* females presented to a *O. nasuta* male (0 *vs.*

2.6+-2.3; p=0.04, g=-1.33). We obtained highly similar results when analysing the data

collected 45 minutes after the visual barrier was removed, or when analysing the shift data

(Table 2).

Behaviour of non-focal specimens

We tested whether the behavioural response of conspecific females and males differed when

presented to a focal female (D) and whether the behavioural response of conspecific and

heterospecific males differed when presented to a focal female (E). Unexpectedly, permanova

revealed a difference in the behaviour of *O. nasuta* and *O. ventralis* males in the OV experiment,

prior to the removal of the barrier. Mann-Whitney U tests revealed that this was due to the

higher average vertical swimming speed of *O. ventralis* males compared to *O. nasuta* males

(SpY 1.7+-0.5 *vs.* 0.7+-0.4, p= 0.014, g=2.05). None of the other comparisons was shown to be

significantly different, neither before, nor after the removal of the barrier (Table 2).

Discussion

Summary

We tested the initial response of *O. nasuta* and *O. ventralis* females towards conspecific and

heterospecific males. In support of our hypothesis, *O. nasuta* females differentiated between

conspecific and heterospecific males, whereas *O. ventralis* females did not. Visualisation of the

data revealed that *O. nasuta* females mirrored the behaviour of the males to which they were

presented. We also presented females of both species with a conspecific female or with nothing

(control). Although females always responded to the presence of another fish, their behaviour

did not differ when presented with conspecific males and females. Comparisons of non-focal

specimens did not reveal any differences in behaviour after presentation to a focal female.

However, before the removal of the wall, males of *O. ventralis* and *O. nasuta* behaved

differently in the OV experiment.

Responses of Ophthalmotilapia *females*

Nevado et al. (2011) discovered signatures of unidirectional hybridisation in *Ophthalmotilapia*,

which could either be explained by cyto-nuclear incompatibilities, or by asymmetric mate

choice. The latter explanation implies that *O. nasuta* females would discriminate more strongly

against heterospecific males, than females of *O. ventralis*. This is supported by our experiments.

As we did not present focal females with heterospecific females, we cannot say that the

observed species recognition in *O. nasuta* females was due to a different response towards

males, or towards all specimens of the other species. However, as females in *Ophthalmotilapia*

are non-territorial and therefore often encounter heterospecific congeners, we expect that the

female response is specific to heterospecific males. Unexpectedly, females of both species

behaved similarly towards conspecific males and females. Hence, we cannot say whether we

observed sexually motivated behaviour, or just the routine behaviour of a (isolated) female that

encounters a conspecific individual. In the wild, non-breeding females of both species

aggregate in large feeding groups (Konings 2019). Hence, being isolated for 12 hours, as was

the case prior to the start of the experiment, represents an unnatural situation for

Ophthalmotilapia females. It would, therefore, not be unlikely if *Ophthalmotilapia* females are

behaviourally hardwired to reunite immediately with conspecifics, regardless of whether these

are female or male.

Females of *O. nasuta* only performed the 'flee' behaviour towards conspecific males and

displayed the 'ram' behaviour less frequently. Although we found no evidence that this

behaviour is sexually motivated, male chasing and female fleeing (i.e. 'flee') form the first steps

in the mating process of *Ophthalmotilapia* (Kéver et al. 2018). The ram behaviour, on the other

hand, was seen in all experimental trials in which a focal female was presented to another fish.

Indeed, the main difference in behaviour between focal females of both species that were, or

were not, presented with another fish was the amount of time spent close to the wall, and in the

display of the 'ram' behaviour.

We discovered that several of the presumed females that we planned to use in the experiments

had been, or changed into, males. This was also observed for one of the non-focal females in

the ON experiment, which was kept isolated from males after the experiment (Supplement 2).

Although there are several reports of sex changes occurring in cichlids (Peters 1975; Naish and

Ribbink 1990), evidence hereof remained, until now, limited. As no reproductions were

recorded when these specimens still had a female morphology, we cannot rule out that they

were males that delayed the development of the conspicuous male colouration and the elongated

pelvic fins. Hence, we did not provide solid evidence of sex change in *Ophthalmotilapia*.

Interpretation

Although we uncovered a significant difference in the behaviour of *O. nasuta* females that were

presented with a conspecific and a heterospecific male, permanova did not reveal a significant

interspecific difference in the behaviour of the males. This could indicate that *O. nasuta* females interpreted behaviour differently when displayed by *O. nasuta* or by *O. ventralis* males. Such species- or sex-dependent interpretation of behaviour is known for several cichlid species, in which territorial males present themselves identically towards both visiting females and intruding males (Baerends and Baerends-Van Roon 1950). Visual exploration of the data, however (Fig. 3C, Supplement 6), revealed a potential difference in male behaviour, which was mirrored by female response. As female *O. ventralis* did not appear to differentiate between conspecific and heterospecific males, one could ask why hybridisation is not even more prevalent. However, we only examined the very first stage in a potential mating process, so other differences that are present in the mating behaviour between both species could be responsible for this. For example, *O. ventralis* males display a specific late mating behaviour, called 'invite', which *O. nasuta* males never display (Kéver et al. 2018). Additionally, hybrids might have a lower fitness. In order to reach mitochrondrial introgression, female hybrids would also need to mate with *O. nasuta* males. This is not unlikely given that, in cichlids, female mate choice is influenced, via imprinting, by the maternal phenotype (Verzijden and ten Cate 2007; Verzijden et al. 2008).

The role of males

Asymmetric propensities towards hybridisation are expected in the intermediate stages of reproductive isolation (Arnold et al. 1996). This has been observed in a variety of animal taxa including lungless salamanders (Verrell 1990), spadefoot toads (Pfennig 2007), swordtails (Crapon de Caprona and Ryan 1990), pupfishes (Strecker and Kodric-Brown 1999), and several cichlids (Egger et al. 2008; Nevado et al. 2011). Although most examples are related to female mate choice, these patterns can also be caused by asymmetries in male choosiness (Svensson

et al. 2007). Although male mate choice is common in fishes (Wong and Jennions 2003; Werner

and Lotem 2003), we choose to focus on the role of females (see introduction, Seehausen et al.

2008; Sefc et al. 2017). However, the mode of fertilization in *Ophthalmotilapia* could also have

an influence on male choosiness. Haesler et al. (2011) studied the reproductive behaviour of *O.*

ventralis, but it can be assumed that the behaviour of its congeners is highly similar. In *O.*

ventralis, a ripe female will visit the territories of several males, either to spawn, or just to

collect additional ejaculates. Subsequently, sperm competition will take place within her mouth,

resulting in clutches with multiple sires (Haesler et al. 2011). Given that this dilutes the effect

of a 'wrong' choice, a female can afford to be less choosy. Differences in both male and female

courtship effort towards genetically distant or similar mates have been documented in another

mouth brooding cichlid: *Tropheus* Boulenger, 1898 (Zoppoth et al. 2013). However, *Tropheus*

species are sexually monomorphic and both sexes are territorial. Additionally, *Tropheus* males

invest significantly more in raising the clutch, by providing the female access to their feeding

territories. As males of *Ophthalmotilapia* do not share their resources, we can expect these

males to be less choosy than those of *Tropheus*. Additionally, a substantial role of male mate

choice is not supported by our data, as we did not observe a difference in behaviour between

non-focal males of *O. nasuta* and *O. ventralis* when presented with females of the two species.

It should be noted, however, that two *O. ventralis* males that displayed mating behaviour

towards *O. nasuta* females were excluded from the analyses.

Whereas our experiments only revealed the capacity for species recognition in females of *O.*

nasuta, we cannot conclude that males cannot distinguish between females of the two species.

Whereas the males of *O. nasuta* and *O. ventralis* behaved differently when a visual barrier was

present, no significant difference was found after its removal. This could imply that males of

the two species behave in a similar way when presented with a conspecific or a heterospecific

female. However, an alternative explanation would be that males recognise conspecific and

heterospecifics, and use this knowledge to court females using a repertoire appropriate to the

species. Although this was found in sister species pairs of freshwater sticklebacks (*Gasterosteus*

spp. L. 1758) (Kozak et al. 2009), our experimental design did not allow us to test this in

Ophthalmotilapia.

We cannot exclude that morphological, physiological and behavioural features that distinguish

O. nasuta males from males of congeners could have caused the asymmetric pattern of

introgression. Foremost, *O. nasuta* males become larger and possess longer pelvic fins. This

feature could render them more attractive as *O. ventralis* females have a preference towards

males with strongly elongated pelvic fins (Haesler et al. 2011). As a change in the feature

associated with attractiveness can alter species recognition in the mating process (Phelps et al.

2006), the extra-long pelvic fins of *O. nasuta* males could serve as a super-natural stimulus

(sensu Tinbergen 1948). Additionally, even though Haesler et al. (2011) found no correlation

between female choice and male body length in *O. ventralis*, they did observe that larger males

outcompeted their rivals in sperm competition within the females' mouth. Additionally, sperm

of *O. nasuta* remains viable for a significantly longer amount of time than that of *O. ventralis*

(Morita et al. 2014). Lastly, *O. nasuta* males construct true bowers (elaborate, crater-shaped

sand mounts), whereas the nests of males of the other species of *Ophthalmotilapia* only consist

of a small area of cleaned rock, or of a small pit in the sand (Konings 2019).

The importance of visual cues

Although animals can use multiple kinds of cues to assess the quality of a potential mate, their

final assessment depends on the overall information available. This is exemplified by female

mate choice in the allopatric swordtail species *Xiphophorus nigrensis* Rosen 1960 and *X. pygmaeus* Hubbs and Gordon 1943. Here, mating preferences differed depending on whether visual, olfactory or a combination of both cues were available (Crapon de Caprona and Ryan 1990). Different responses to visual and olfactory cues were also shown for females of sympatric *Cyprinodon* Lacipède, 1803 pupfish species from Lake Chichancanab (Mexico). Here, different degrees of asymmetric discrimination of males were observed depending on whether females had access to visual or olfactory information (Strecker and Kodric-Brown 1999). In species-rich systems and in species that form leks, such as *Ophthalmotilapia* spp., females must be able to rapidly assess the quality of a potential mate (Barlow 2002). *Ophthalmotilapia* females could use the morphological characteristics, conspicuous bowers and/or stereotypical displays of males to distinguish them from sympatric congeners. However, even though multiple cues can be involved, mate choice decisions in radiations are often based on just a small amount of (combinations of) these traits (Hohenlohe and Arnold 2010).

The separation wall used in our experiments contained holes that allowed for the exchange of water between both compartments. Hence, besides visual clues, the fishes most likely also received olfactory and acoustic information. Although visual cues were suggested to be the primary factor in species-isolating, female mate choice in other cichlids (Jordan et al. 2003; Kidd et al. 2006), we cannot rule out the importance of other types of information. Studies have shown that olfactory (Blais et al. 2009; Plenderleith et al. 2005), acoustic (Nelissen 1978; Amorim et al. 2004; Kéver et al. 2018) and behavioural (Barlow 2002) information can also influence the mating process. Although Seehausen and van Alphen (1998) showed a certain hierarchy of information, where other cues are taken into account when visual information is absent or masked, other experiments showed that female cichlids are more likely to select the right male when both olfactory and visual cues are available (Plenderleith et al. 2005; Blais et

al. 2009). When visual information suffices for mate recognition, the behaviour throughout the

mating process, i.e. potentially leading toward spawning, does not need to diverge between

closely-related species (Barlow 2002). This may explain why spawning behaviour of

Ophthalmotilapia is remarkably similar across the genus (Kéver et al. 2018) and why

differently-coloured, sympatric mbuna cichlids from Lake Malawi have identical courtship

behaviours (McElroy and Kornfeld 1990).

Ecological reasons for asymmetric hybridisation

Although they can be found in sympatry, *O. ventralis* is more associated with the rocky shores

of Lake Tanganyika, whereas *O. nasuta* has a wider ecological range. At rocky shores, *O.*

ventralis can be one of the most abundant cichlid species (Sturmbauer et al. 2008). Hence, for

an *O. ventralis* female, a random encounter with another *Ophthalmotilapia* male is much more

likely to result in a conspecific than a heterospecific encounter. In contrast, for an *O. nasuta*

female venturing into the preferred *O. ventralis* habitat, a conspecific encounter would be less

often the case. Therefore, the ability to discriminate between conspecific and heterospecifics

would be less important for females of *O. ventralis* than for those of *O. nasuta*. A similar

interpretation was given to explain asymmetries in female discrimination of sympatric

Cyprinodon species, where the species with the highest abundance had the lowest choosiness

(Strecker and Kodric-Brown 1999). Although a species' propensity for discrimination could be

a consequence of its distribution range, the opposite could also hold. Species that are better in

recognizing conspecifics are more likely to maintain the integrity of their gene pool. Hence,

they could be better in colonising habitats that have already been occupied by related species.

Finally, we showed that substantial behavioural differences can be observed between closely-

related species. This should be a warning to be cautious when assuming similarities in the

behaviour of certain (model) organisms and related taxa.

Acknowledgments

We are grateful to Siegfried Loose for his advice on the rearing of *Ophthalmotilapia* and to

Nedim Tüzün for his advice on the tracking software. We thank Ad Konings (Cichlid press) to

provide us with pictures of *Ophthalmotilapia*. This study is part of the GENBAS project, which

was funded by the BELSPO (Belgian Science Policy) BRAIN-be program.

Ethical note

All specimens were obtained from a certified commercial supplier (Cichlidenstadl, Alerheim,

Germany). Specimens were housed in the aquarium facilities of the University of Liège.

Experimental procedures were performed in accordance with Belgian law, and approved by the

University of Liège Institutional Animal Care and Use Committee (protocol #1759) in

accordance with the regulations of the ethical committee of the University of Liège. All

manipulations were performed by a FELASA-certified experimenter. The data and script are

deposited on Zenodo under: 10.5281/zenodo.5534485.

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content

of this article. MVS is one of the PCI Zool recommenders.

References

Amorim MCP, Knight ME, Stratoudakis Y, Turner GF. 2004. Differences in sounds made by

courting males of three closely related Lake Malawi cichlids. J Fish Biol. 65(5):1358-1371.

Anderson MJ. 2006. Distance-based tests for homogeneity of multivariate dispersions.

Biometrics. 62(1):245–253.

Arnold SJ, Verrell PA, Tilley SG. 1996. 1996 The evolution of asymmetry in sexual isolation:

A model and a test case. Evolution 50:1024-1033

Barlow GW. 2002. How behavioural studies contribute to the species problem: a piscine

perspective. Fish Fish. 3(3):197-212.

Baerends GP, Baerends-Van Roon JM. 1950. An introduction to the study of the ethology of

the cichlid fishes, Leiden, The Netherlands, E. J. Brill.

Blais J, Plenderleith M, Rico C, Taylor MI, Seehausen O, van Oosterhout C, Turner GF. 2009.

Assortative mating among Lake Malawi cichlid fish populations is not simply predictable from

male nuptial colour. BMC Evol Biol. 9(53):53.

Coyne JA, Orr HA. 2004. Speciation. Sunderland, MA: Sinauer Associates.

Crapon de Caprona MD, Ryan MJ. 1990. Conspecific mate recognition in swordtails,

Xiphophorus nigrensis and *X. pygmaeus* (Poeciliidae): olfactory and visual cues. Anim Behav.

39(2):290-296.

Danley PD, Kocher TD. 2001. Speciation in rapidly diverging systems: lessons from Lake

Malawi. Mol Ecol. 10:1075-1086.

Egger B, Obermüller B, Eigner E, Sturmbauer C, Sefc KM. 2008. Assortative mating

preferences between colour morphs of the endemic Lake Tanganyika cichlid genus *Tropheus*.

Hydrobiologia. 615:37-48.

Friard O, Gamba M. 2016. BORIS: A Free, Versatile Open-Source Event-Logging Software

for Video / Audio Coding and Live Observations. Methods Ecol Evol. 7(11):1–6.

Gonzalez-Voyer A, Fitzpatrick JL, Kolm N. 2008. Sexual selection determines parental care

patterns in cichlid fishes. Evolution 62(8): 2015-2026.

Haesler MP, Lindeyer CM, Otti O, Bonfils D, Heg D, Taborsky M. 2011. Female

mouthbrooders in control of pre- and postmating sexual selection. Behav Ecol. 22(5):1033–

1041.

Hammer Ø, Harper DA, Ryan PD. 2001. PAST: Paleontological statistics software package for

education and data analysis. Palaeontologia Electronica. 4:9pp.

Hanssens M, Snoeks J, Verheyen E. 1999. A Morphometric Revision of the Genus

Ophthalmotilapia (Teleostei, Cichlidae) from Lake Tanganyika (East Africa). Zool J Linn Soc-

Lon. 125(4):487–512.

Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A. 2019. Moving beyond P values: data

analysis with estimation graphics. Nat Met. 16(7):565-566.

Hohenlohe PA, Arnold SJ. 2010. Dimensionality of mate choice, sexual isolation, and

speciation. PNAS. 107(38):16583-16588.

Immler S, Taborsky M. 2009. Sequential polyandry affords post-mating sexual selection in the

mouths of cichlid females. Behav Ecol Sociobiol. 63(8):1219–1230.

Jiggins CD, Salazar C, Linares M, Mavarez J. 2008. Hybrid trait speciation and *Heliconius*

butterflies. Philos T R Soc B. 363(1506):3047-3054.

Jordan R, Kellogg K, Juanes F, Stauffer J. 2003. Evaluation of female mate choice cues in a

group of Lake Malawi mbuna (Cichlidae). Copeia. 2003(1):181-186.

Kéver L, Parmentier E, Derycke S, Verheyen E, Snoeks J, Van Steenberge M, Poncin P. 2018

Limited possibilities for prezygotic barriers in the reproductive behaviour of sympatric

Ophthalmotilapia species (Teleostei, Cichlidae). Zoology. 126:71-81.

Kidd MR, Danley PD, Kocher TD. 2006. A direct assay of female choice in cichlids: all the

eggs in one basket. J Fish Biol. 68(2):373–384.

Kocher TD. 2004. Adaptive evolution and explosive speciation: the cichlid fish model. Nat Rev

Genet. 5(4):288-298.

Koblmüller S, Duftner N, Sefc KM, Aibara M, Stipacek M, Blanc M, Egger B, Sturmbauer C.

2007. Reticulate phylogeny of gastropod-shell-breeding cichlids from Lake Tanganyika - the

result of repeated introgressive hybridisation. BMC Evol Biol. 7:7.

Konings A. 2019. Tanganyika cichlids in their natural habitat. 4th edition. El Paso, TX: Cichlid

press.

Kozak GM, Boughman JW. 2009. Learned conspecific mate preference in a species pair of

sticklebacks. Behav Ecol. 20: 1282–1288.

Kozak GM, Reisland M, Boughmann JW. 2009. Sex differences in mate recognition and

conspecific preference in species with mutual mate choice. Evolution. 63: 353-365.

Luttbeg B, Towner MC, Wandesforde-Smith A, Mangel M, Foster SA. 2001. State-dependent

mate-assessment and mate-selection behavior in female threespine sticklebacks (*Gasterosteus*

aculeatus, Gasterosteiformes: Gasterosteidae). Ethology. 107(6):545-558.

Mayr E. 1982. The growth of biological thought: Diversity, evolution and inheritance.

Cambridge, MA: Harvard University Belknap Press.

Marques DA, Meier JI, Seehausen O. 2019. A combinatorial view on speciation and adaptive

radiation. Trends Ecol Evol. 34(6):531-544.

Mendelson TC, Shaw KL. 2012. The (mis)concept of species recognition. Trends Ecol Evol.

27(8):421-427.

McElroy DM, Kornfeld I. 1990. Sexual selection, reproductive behaviour, and speciation in the

mbuna species fock of Lake Nyasa (Pisces: Cichlidae). Environ Biol Fish. 28:273-284.

Moran MD. 2003. Arguments for rejecting the sequential Bonferroni in ecological studies.

Oicos. 100(2):403-405.

Morita M, Awata S, Yorifuji M, Ota K, Kohda M, Ochi H. 2014. Bower-building behaviour is

associated with increased sperm longevity in Tanganyikan cichlids. J Evolution Biol.

27(12):2629–2643.

Naish KA, Ribbink AJ. 1990. A preliminary investigation of sex change in *Pseudotropheus*

lombardoi (Pisces: Cichlidae). Environ Biol Fish. 28:285–294.

Nakagawa S. 2004. A farewell to Bonferroni: the problem if low statistical power and

publication bias. Behav. Ecol. 15:1044-1045.

Nelissen MHJ. 1978. Sound production by some Tanganyikan cichlid fishes and a hypothesis

for the evolution of their communication mechanisms. Behaviour. 64(1-2):137–147.

Nevado B, Fazalova V, Backeljau T, Hanssens M, Verheyen E. 2011. Repeated unidirectional

introgression of nuclear and mitochondrial DNA between four congeneric Tanganyikan

cichlids. Mol Biol Evol. 28(8):2253-2267.

Oksanen JF, Blanchet G, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara

RB, Simpson GL, Solymos PM, Stevens HH, Szoecs E, Wagner H. 2017. vegan: Community

Ecology Package. R package version 2.4-5.

Panfili J, Thior D, Ecoutin J-M, Ndiaye P, Albaret J-J. 2006. Influence of salinity on the size at

maturity for fish species reproducing in contrasting West African estuaries. J Fish Biol.

69(1):95-113.

Peters HM. 1975. Hermaphroditism in cichlid fishes. pp. 228–235.In: R. Reinboth (ed.)

Intersexuality in the Animal Kingdom. Springer-Verlag, Berlin.

Pfennig KS. 2007. Facultative mate choice drives adaptive hybridization. Science, 318(5852):

965.

Phelps SM, Rand AS, Ryan MJ. 2006. A cognitive framework for mate choice and species

recognition. Am Nat. 167(1):28–42.

Plenderleith M, van Oosterhout C, Robinson RL, Turner GF. 2005. Female preference for

conspecific males based on olfactory cues in a Lake Malawi cichlid fish. Biol Lett. 1(4):411-

414.

Poll M. 1986. Classification des cichlidae du Lac Tanganyika: Tribus, genres et espèces. Bull

Acad R Belg. 8:1-163.

R core team. 2017. R: A Language and environment for statistical computing. R foundation for

statistical computing, Vienna, Austria.

Ronco F., Büscher H. H., Indermauer A., Salzburger W. 2020. The taxonomic diversity of the

cichlid fish fauna of ancient Lake Tanganyika, East Africa. J. Great Lakes Res. 46: 1067-1078.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image

analysis. Nat Met. 9(7):671-675.

Roux C, Fraïsse C, Romiguier J, Anciaux Y, Galtier N, Bierne N. 2016. Shedding light on the

grey zone of speciation along a continuum of genomic divergence. PLOS Biol.

14(12):e2000234.

Rüber L, Meyer A, Sturmbauer C, Verheyen E. 2001. Population structure in two sympatric

species of the Lake Tanganyika cichlid tribe Eretmodini: Evidence for introgression. Mol Ecol.

10(5):1207–1225.

Salzburger W. 2018. Understanding explosive diversification through cichlid fish genomics.

Nat Rev Genet. 19:705-717.

Salzburger W, Braasch I, Meyer A. 2007. Adaptive sequence divergence in a colour gene

involved in the formation of the characteristic egg-dummies of male *Haplochromis* cichlid

fishes. BMC Biol. 5(51):13pp.

Seehausen O. 2004. Hybridisation and adaptive radiation. Trends Ecol Evol. 19(4):198-207.

Seehausen O, van Alphen JJM. 1998. The effect of male coloration on female mate choice in

closely related Lake Victoria cichlids (*Haplochromis nyererei* complex). Behav Ecol Sociobiol.

42:1–8.

Seehausen O, Terai Y, Magalhaes IS, Carleton KL, Mrosso HDJ, Miyagi R, van der Sluijs I,

Schneider MV, Maan ME, Tachida H, Imai H, Okada N. 2008. Speciation through sensory

drive in cichlid fish. Nature. 455:620-626.

Sefc KM, Mattersdorfer K, Ziegelbecker A, Neuhüttler N, Steiner O, Goessler W, Koblmüller

S. 2017. Shifting barriers and phenotypic diversification by hybridisation. Ecol Lett. 20(5):651-

662.

Sommer-Trembo C, Plath M, Gismann J, Helfrich C, Bierbach D. 2017. Context-dependent

female mate choice maintains variation in male sexual activity. R Soc Open Sci. 4: 170303.

Strecker U, Kodric-Brown A. 1999. Mate recognition systems in a species flock of Mexican

pupfish. J Evolution Biol. 12(5):927- 935.

Sturmbauer C, Fuchs C, Harb G, Damm E, Duftner N, Maderbacher M, Koch M, Koblmüller

S. 2008. Abundance, distribution, and territory areas of rock-dwelling Lake Tanganyika cichlid

fish species. Hydrobiologia. 615(1):57-68.

Sullivan BK. 2009. Mate recognition, species boundaries and the fallacy of "species

recognition". Open Zool. 2:86-90.

Svensson EI, Karlsson K, Friberg M, Eroukhmanoff F. 2007. Gender differences in species

recognition and the evolution of asymmetric sexual isolation. Curr Biol. 17(22):1943–1947.

Tinbergen N. 1948. Social releasers and the experimental method required for their study.

Wilson Bull. 60(1):6-51.

Turner GF, Seehausen O, Knight ME, Allender CJ, Robinson RL. 2001. How many species of

cichlid fishes are there in African lakes. Mol Ecol. 10(3):793-806.

Theis A, Salzburger W, Egger B, 2012. The function of anal fin egg-spots in the cichlid fish

Astatotilapia burtoni. PLoS ONE 7(1):e29878.

Thomas GH, Székely T. 2005. Evolutionary pathways in shorebird breeding systems: sexual

conflict, parental care, and chick development. Evolution 59(10): 2222–2230.

Trivers RL. 1972. Parental investment and sexual selection. in Sexual selection and the descent

of man, EBG. Campbell, ed. (136-179). Aldine, Chicago.

Van Oppen MJH, Turner GF, Rico C, Robinson RL, Deutsch JC, Genner MJ, Hewitt GM. 1998.

Assortative mating among rock-dwelling cichlid fishes supports high estimates of species

richness from Lake Malawi. Mol Ecol. 7(8):991-1001.

Verrell PA. 1990. Frequency of interspecific mating in salamanders of the plethodontid genus

Desmognathus: different experimental designs may yield different results. J Zool. 221(3):441-

451.

Verzijden MN, Korthof REM, ten Cate C. 2008. Females learn from mothers and males learn

from others. The effect of mother and siblings on the development of female mate preferences

and male aggression biases in Lake Victoria cichlids, genus *Mbipia*. Behav Ecol Sociobiol. 62:

1359–1368.

Verzijden MN, ten Cate, C. 2007. Early learning influences species assortative mating

preferences in Lake Victoria cichlid fish. Biol lett 3(2): 134-136.

Werner NY, Lotem A. 2003. Choosy males in a haplochromine cichlid: first experimental

evidence for male mate choice in a lekking species. Anim Behav. 66(2):293-298.

Wirtz, P. 1999. Mother species-father species: unidirectional hybridization in animals with

female choice. Animal Behaviour. 58:1–12.

Wong BMB, Jennions MD. 2003. Costs influence male mate choice in a freshwater fish. P Roy

Soc B-Biol Sci. 270(Suppl 1):S36-S38.

Zoppoth P, Koblmüller S, Sefc KM. 2013. Male courtship preferences demonstrate

discrimination against allopatric colour morphs in a cichlid fish. J Evolution Biol. 26(3): 577-

586.

Additional material

Supplement 1. Summary of the experiments.

Supplement 2. Indications for sex change observed in *O. nasuta* females. Top row: gonads of

some of the specimens that were female in external phenotype but had male or ambiguous

gonads (ON33, 35, 26, 43). Vertical left: female gonads in developmental stage 3 (ON34), 4

(ON24) and 5 (ON37) respectively (Panfili et al. 2006). Bottom: two presumed non-focal *O.*

nasuta females used in the ON experiment, one of which could have undergone transition after

the experiments, and horizontal right, the ventral area of the same specimens, with A: anal pore

and UG: urogentital pore.

Supplement 3. Visualization of tracking data. The position of each fish is plotted for each

second in which specimens were tracked with the positions recorded before and after the

removal of the opaque wall (grey) coloured differently. Ellipses denote the area in which 90%

of tracks are situated, large dots denote the average positions before and after the removal of

the barrier and arrows shows the change in mean position. The separation wall is visualized as

a meshed partition. The average speed before (v0) and after (v1) the removal of the barrier is

plotted for each tracked specimen with data for focal specimens given in bold. Abbreviations

(ON: *O. nasuta*, OV, *O. ventralis*, F: female, M; male)

Supplement 4. Summary of the data.

Supplement 5. Loadings and variance of the main axes of the PCAs and CVAs conducted in

the study.

Supplement 6. Principal component analyses and Canonical variate analyses performed on the

behaviours recorded 45 min after the visual barrier was removed of the ON (*O. nasuta*, left)

and the OV (*O. ventralis,* right) experiment. Symbols on the scatter plots for the ON and OV

experiment as in E and F, respectively, with full circles denoting focal females presented with

no fish (red), a conspecific female (blue) an *O. ventralis* male (purple), and an *O. nasuta* male

(green), empty circles denote non-focal conspecific females and full squares *O. ventralis*

(purple) and *O. nasuta* (green) males. Explained variances are added to the axes.

Supplement 7. Shift in average position of the specimens analysed 15min before and 15min

after the removal of the separation wall for the ON (above) and OV (below) experiment. Focal

specimens are all visualised on the left, and non-focal specimens on the right. Dimensions in

cm, with the vertical bar representing the separation wall. Dashed arrows represent individual

fishes, bold arrows the average per treatment. Colours, for focal females (ON and OV)

presented with no fish (red), a conspecific female (turquoise) an *O. ventralis* male (pink), and

an *O. nasuta* male (light green), and non-focal specimens (right): conspecific females (blue),

O. ventralis males (purple) and *O. nasuta* males (green).