Title: Clenbuterol Attenuates Immune Reaction to Lipopolysaccharide and Its Relationship to Anhedonia in Adolescents

Running Title: Immune System Inhibition by Clenbuterol and Anhedonia in Adolescents

Authors: Tram N. B. Nguyen, B.S.[1]; Benjamin A. Ely, Ph.D.[1]; Danielle Pick[1]; Manishkumar Patel, M.S.[2]; Hui Xie, M.S.[2]; Seunghee Kim-Schulze, Ph.D.[2]; Vilma Gabbay, M.D., M.S.[1, 3]

1. Department of Psychiatry and Behavioral Sciences, Albert Einstein College of Medicine, Bronx, NY

2. Human Immune Monitoring Center, Icahn School of Medicine at Mount Sinai, New York, NY

3. Nathan S. Kline Institute for Psychiatric Research, Orangeburg, NY

Corresponding Author:

Vilma Gabbay, M.D., M.S.

Department of Psychiatry and Behavioral Sciences, Albert Einstein College of Medicine

Address: Van Etten 4A-44, 1225 Morris Park Ave, Bronx, NY 10461

Phone: + 1-718-839-7549

Fax: + 1-718-839-7575

Email: vilma.gabbay@einsteinmed.edu

Keywords: Clenbuterol; Lipopolysaccharide; Immune Challenge; Cytokines; Anhedonia; Adolescence; RDoC

Abstract

While inflammation has been implicated in depression, little is known of immune-inhibitory agents for individuals with depression. This study sought to assess whether β_2-agonist clenbuterol (CBL) would attenuate increased cytokine secretion in adolescents with mood and anxiety symptoms following *ex vivo* exposure of whole blood to lipopolysaccharide (LPS). Our focus on adolescents aimed to target a critical developmental period when psychiatric conditions often emerge and prior to chronicity effects. Participants (15.25 ± 2.16 years old, 59% female) were 97 psychotropic-medication free adolescents with mood and anxiety symptoms and 33 healthy controls. All had comprehensive evaluations and dimensional assessments of psychiatric symptoms. Fasting whole-blood samples were collected and stimulated with LPS in the presence and absence of CBL for 6 hours and analyzed for 41 cytokines. Comparison analyses used Bonferroni-corrected nonparametric tests. Exploratory factor analysis reduced 41 cytokines into 5 cytokine factors in each experimental condition, and their relationships with psychiatric symptoms were examined. Levels of nine cytokines were significantly reduced by CBL treatment compared to LPS alone. A cytokine factor in the LPS+CBL condition significantly correlated with anticipatory (*rho* = -0.39, *p* = 7.4 × 10^{-5}) and consummatory anhedonia (*rho* = -0.36, *p* = 3.3 × 10^{-4}), which remained significant when controlling for depression. This study supports the possible inhibitory effect of CBL on immune activation. Using a data-driven method, distinctive relationships between CBL-affected cytokines and dimensional anhedonia were reported, further elucidating the role of β_2-agonism in adolescent affective symptomatology.

1. Introduction

Inflammation has been implicated in numerous neuropsychiatric conditions (1, 2). Work by our group and others have consistently reported a link between peripheral inflammatory processes across psychiatric conditions and ages (3-11). The relationships between inflammation and psychopathologies are thought to reflect similar neurobiological mechanisms to those underlying "sickness behavior" in model animals (8). Sickness behavior is characterized by fatigue, malaise, and reductions in appetite, ability to concentrate, mood, social interactions, and pleasure-seeking behaviors (12-15). While sickness behavior is an adaptive process involving complex brain-body interactions, dysregulated immune activation is postulated to induce maladaptive changes in the central nervous system, which ultimately manifest as psychiatric disturbances (8). Since psychiatric conditions are heterogenous, our group and others have utilized the Research Domain Criteria (RDoC) framework, which focuses on behavioral constructs rather than categorical psychiatric diagnoses (16). We found that only anhedonia—the decreased capacity to experience pleasure—was associated with peripheral inflammation among adolescents (5, 6, 17). As anhedonia clinically reflects reward dysfunction, our findings suggest that inflammation may induce alterations in neurocircuits underlying reward function. Indeed, our neuroimaging work supported this investigative direction, providing evidence for the relationships between clusters of cytokines and alterations of the neural function across reward processes (18-20).

As inflammation involves many intertwined and tightly orchestrated systems, it is crucial to examine processes modulating immune activity to delineate the brain-immune relationships. One prominent immune regulator is the adrenergic nervous system, which

has been shown to halt overactivation of peripheral and central immune cells through β_2-adrenergic receptor transduction pathways (21). Attempts to study the role of adrenergic regulation in depressive pathophysiology have yielded promising results. Earlier post-mortem studies showed differential adrenoreceptor density in reward-related cortical regions between antidepressant-free suicide victims and matched controls (22, 23). More recently, catecholamine depletion has been linked to alterations in reward processing among adults with major depressive disorder (24). Nevertheless, the anti-inflammatory effects of the adrenergic system on reward function remains largely unexamined in depression and across the lifespan. Among immune-inhibitory β_2 receptor agonists, clenbuterol (CBL) is a promising candidate for use in modeling adrenergic regulation of immune activation *ex vivo*. Especially relevant to psychiatric contexts, CBL readily crosses the blood brain barrier to act on central β_2 receptors, with preclinical evidence indicating that CBL reduces neuroinflammation associated with strokes (25) and exhibits antidepressant-like effects (26) in rodent models. However, investigation of β_2 agonists adrenergic system regulating immune CBL's anti-inflammatory property using human blood samples is scarce, and available studies have only analyzed a limited number of inflammatory markers.

Building upon the above observations, here we sought to investigate whether adrenergic immune modulation could be achieved in a functional immune response model using lipopolysaccharide (LPS) and CBL in whole blood. In alignment with the RDoC framework, we examined behavioral constructs to capture the full range of symptom severity rather than stratifying participants by categorical diagnoses. Therefore, participants were psychotropic-medication free adolescents with diverse mood and

anxiety symptoms as well as healthy adolescents. Our focus on adolescence aimed to capture a critical period with increased vulnerability to psychopathology development. We used a comprehensive profile of 41 immune biomarkers including cytokines, chemokines, and growth factors (referred to throughout as "cytokines" for convenience). We hypothesized that CBL would attenuate immune activation following exposure to LPS in adolescents. There were no *a priori* hypotheses regarding which specific LPS-stimulated cytokines would be downregulated by CBL, as we expected a general attenuating effect on cytokine production. Further, based on our prior finding implicating inflammation in anhedonia, we also explored the relationships between CBL-induced attenuation of inflammatory effects and severity of depression, anxiety, and anhedonia symptoms.

2. Methods

2.1. Study Participants

Adolescents in the New York City area were recruited via community advertisements and clinician referrals. On the first visit, adolescents underwent a comprehensive medical and psychiatric assessment to determine eligibility for study participation. Eligible adolescents were instructed to return to the study site within approximately 2 weeks to undergo study procedures, including vital sign measurements, a fasting blood draw, a urine toxicology screen, a pregnancy test if female, and completion of clinical questionnaires.

This study was approved by the Institutional Review Boards at Icahn School of Medicine at Mount Sinai and Albert Einstein College of Medicine. Participants ages 18 and older (n=24) provided written informed consent; participants younger than 18 years old provided assent, and a legal guardian provided signed consent.

2.2. Inclusion and Exclusion Criteria

The study included medically healthy adolescents ages 12 to 20 years. Inclusion criteria for the clinical group: primary presentation of mood and anxiety symptoms regardless of meeting diagnostic criteria per the Diagnostic and Statistical Manual of Mental Disorders (DSM), Fourth or Fifth Edition (27, 28). Exclusion criteria for the clinical group: a) current or past DSM diagnosis of eating disorders, schizophrenia, pervasive developmental disorder, or substance use disorder, b) intake of any psychotropic medication within 30-90 days (depending on drug half-life) prior to the blood draw; c) history of immunological or hematological disorder; d) intake of any immune-affecting medication and supplements (e.g., steroids, non-steroidal anti-inflammatory drugs,

omega-3 fatty acids) within 2 weeks prior to study enrollment; c) history of chronic fatigue syndrome; d) any infection during the month prior to the blood draw (including the common cold); e) history of significant medical or neurological disorders; f) an estimated IQ below 80 based on the Kaufmann Brief Intelligence Test (KBIT; 29); g) positive urine toxicology screens; h) in females, a positive pregnancy test.

Healthy controls (HC) were required to meet all above exclusion criteria in addition to be without a lifetime history of any psychiatric diagnoses, including subthreshold symptoms as determined by clinical evaluations (detailed in section *2.3.* below).

2.3. Clinical Evaluations

Each participant received a thorough evaluation consisting of a comprehensive psychiatric evaluation, medical history as well as laboratory tests including complete blood count, metabolic panel, liver and thyroid function tests, and a urine toxicology test. Both the participating adolescent and their accompanying parent were interviewed by a clinician to assess for possible presence of an infectious illness (including the common cold) within the month immediately prior to enrollment.

Psychiatric diagnostic evaluation: The semi-structured Schedule for Affective Disorders and Schizophrenia – Present and Lifetime version (K-SADS-PL; 30), an interview-based assessment tool commonly used in pediatric research settings, was conducted by a trained, licensed psychiatrist or clinical psychologist to assess presence of any psychiatric symptoms and lifetime history of psychiatric diagnoses per the DSM. Evaluations were discussed between the interviewing clinician and the Principal Investigator, a board-certified child and adolescent psychiatrist, to enhance diagnostic

reliability. The K-SADS-PL was utilized to ensure participants meet study eligibility and determine if they belonged to the clinical or HC group.

Dimensional symptom assessment: The self-rated, 39-item Multidimensional Anxiety Scale for Children (MASC; 31) was employed to quantify anxiety severity (scores ranging from 0 to 117). Participants self-reported their depression severity with the 21-item Beck Depression Inventory, 2nd edition (BDI-II; scores ranging 0 to 63; 32). Higher scores on the MASC and BDI indicate more severe anxiety and depression, respectively. Additionally, we utilized the Temporal Experience of Pleasure Scale (TEPS; 33) to separately capture anticipatory (TEPS-A; scores ranging 10 to 60) and consummatory (TEPS-C; scores ranging 8 to 48) components of anhedonia. Higher TEPS scores suggest lower levels of anhedonia. The TEPS' internal consistency is comparable in both adult and adolescent samples (33, 34), and its breakdown of anticipatory and consummatory subscales has been supported by exploratory factory analysis using data collected in two independent samples (35). We note that the BDI, MASC, and TEPS are designed to dimensionally probe the full range of symptoms severity, and thus HCs in our sample were expected to report non-zero, though non-pathological, levels of depression, anxiety, and anhedonia (see **Table 1**).

2.4. Ex Vivo *Immune Stimulation with Lipopolysaccharide and Clenbuterol*

Blood samples were collected between 9 and 10 A.M. after an overnight fast lasting at least 12 hours. Functional immune responses were assessed *in vitro* using a well-established LPS challenge protocol that reliably induces pro-inflammatory cytokine production (36). Whole blood samples were cultured on standard growth medium for 6 hours in a tissue culture incubator under three conditions: in the presence of LPS (0.1

μg/ml) alone, in the presence of LPS along with the β_2-agonist CBL (10^{-6} M), and in the absence of both LPS and CBL (i.e. Control condition). After 6 hours of incubation, the cultured whole blood samples were centrifuged; supernatant was collected and stored at -20°C.

Levels of 41 immune biomarker analytes (cytokines, chemokines, and growth factors; see **Supplementary Table 1**) were measured using a bead-based Luminex-200 system and the XMap Platform (Luminex Corporation, Austin, TX), as described in **Supplementary Methods** – *SM1*. Assays were performed at the Mount Sinai Human Immune Monitoring Center by laboratory staff blinded to participants' clinical status. The combination of assays has been previously validated using secondary measurements (37-41). Duplicates of median fluorescent intensity (MFI) values were measured for each of the 3 peripheral whole blood culture growth conditions (Control, LPS, LPS+CBL), and mean MFI values were generated. While absolute concentration values (**Supplementary Table 1**) were computed from mean MFI values, we opted to use MFI values in our analyses as these allow for increased statistical power (42, 43). To enhance the reproducibility and transparency of our findings, we reported further details on the detection limits, methods to account for potential assay drifts, and reasons for missing MFI values for each analyte in **Supplementary Methods** – *SM1* and *SM2*.

2.5. Statistical Analyses

Primary analyses were conducted in MATLAB R2020b (MathWorks, Natick, MA). An overview of our analytic protocol can be found in **Supplementary Figure 2**. Shapiro-Wilk tests indicated a non-normal distribution for most cytokines in our sample, necessitating a nonparametric approach. We first utilized Friedman tests, controlling for

family-wise error rate using Bonferroni correction, to test for significant differences in cytokine levels across the three conditions (Control, LPS, and LPS+CBL). For any cytokines found to differ significantly, *post-hoc* Wilcoxon signed-rank tests with Bonferroni correction were used to examine pairwise differences between conditions. Effect sizes for Wilcoxon signed-rank tests were computed as the correlation coefficient r ($r = \frac{z}{\sqrt{N}}$) (44-47).

In secondary analyses, given the unknown relations between CBL-attenuated inflammatory markers and subjective psychiatric symptoms, we utilized an exploratory factor analysis (EFA) model to identify sources of shared variance within subjects' detailed immune profiles in a data-driven manner. Cytokine data were examined using 3 discrete EFA models corresponding to the 3 conditions (Control, LPS, LPS+CBL). EFA was performed in SPSS 27.0 (IBM, Armonk, NY) using principal component analysis (PCA) dimensionality reduction and varimax orthogonal rotation. Finally, Spearman partial correlations controlling for age, sex, and body mass index (BMI) were used to assess the relationships between each cytokine factor scores and BDI (depression), MASC (anxiety), TEPS-A (anticipatory anhedonia), and TEPS-C (consummatory anhedonia) scores. Spearman correlations between cytokine factors and anhedonia measures were repeated while controlling for depression severity.

Following our primary analyses outlined above, we performed a supplementary factor analysis to assess relative changes in cytokine levels between LPS and LPS+CBL conditions and related the resulting factors to anhedonia scores (see **Supplementary Results** – *SR1*). Further *post hoc* analyses were conducted to contrast cytokine relationships between LPS+CBL and LPS conditions. Spearman correlations (*rho*)

between pairs of cytokines in the 2 conditions were computed, *z*-transformed, subtracted from each other, then transformed back to *rho* coefficients. This yielded a total of 820 pairwise correlation differences (Δ*rho*) per subject. The significance of each Δ*rho* was assessed using a bias-corrected and accelerated bootstrap (48-50) with 10^5 resamples. We additionally utilized this bootstrapping method to determine whether observed relationships between symptom measures and immune factors are significantly stronger than other comparisons (details in **Supplementary Results** – *SR2*).

3. Results

3.1. Participants' Characteristics

This study enrolled a total of 130 adolescents, including 97 in the clinical group (85 endorsed mood and anxiety symptoms; 12 with other externalizing behavioral symptoms without comorbid mood and anxiety symptoms) and 33 HC. As summarized in **Table 1**, participants were racially and ethnically diverse. Demographically, participants with psychiatric symptoms and HC were similar ($p > 0.1$), except for a significantly higher proportion of Hispanic individuals among the clinical sample ($p = 0.012$). More than half of participants in the clinical group had at least two concurrent DSM diagnoses. As expected, adolescents in the clinical group exhibited higher depression and anxiety severity compared to HC (see **Table 1**). Depression and anhedonia scores were skewed, while anxiety scores were normally distributed (see **Supplementary Figure 3**). Across all participants, anticipatory and consummatory anhedonia scores were significantly correlated with each other and with depression, but not with anxiety (see **Supplementary Table 2**).

[**Table 1** here]

3.2. Lipopolysaccharide and Clenbuterol Effects on Cytokines

Cytokine levels were not significantly different between HC and participants with psychiatric symptoms for any condition (see **Supplementary Table 3**). Consequently, all analyses used combined data from the whole sample. Friedman tests (detailed in **Supplementary Table 4**) indicated significant differences in 21 cytokines across the three conditions after Bonferroni correction ($\alpha = \frac{0.05}{41} \approx 1.2 \times 10^{-3}$).

Pairwise follow-up analyses (i.e., Control vs. LPS, LPS vs. LPS+CBL, Control vs. LPS+CBL) of these 21 cytokines using Wilcoxon signed-rank tests are detailed in **Table 2**, with results considered significant at the Bonferroni corrected $\alpha = \frac{0.05}{21 \times 3} \approx 7.9 \times 10^{-4}$ level. Relative to the Control condition, LPS increased levels of 19 out of the 21 cytokines, with only PDGF-AB/BB and sCD40L not meeting significance. Levels of 17/21 cytokines (all except Fractalkine, G-CSF, IL-12P40, and sCD40L) were also elevated in the LPS+CBL condition relative to Control. For all but 2/21 cytokines (PDGF-AB/BB and sCD40L), effect sizes were smaller for LPS+CBL vs. Control than for LPS vs. Control, consistent with the immunosuppressant activity of CBL. Compared to LPS, levels of 9/21 cytokines were significantly reduced in the LPS+CLB condition, while only 1/21 (sCD40L) was found to be increased in the presence of CBL. Cytokines with the greatest differences in expression between LPS+CBL and LPS conditions included MIP-1β (r = -0.73), TNF-α (r = -0.70), MIP-1α (r = -0.65), and IL-6 (r = -0.62). These findings remained consistent when stratified by group membership (see **Supplementary Tables 5–6**).

[**Table 2** here]

3.3. Cytokine Dimensionality Reduction Across Experimental Conditions

In each of the 3 examined conditions (see **Methods –** *2.5.*), EFA yielded an initial model containing 8 orthogonal cytokine factors with eigenvalues greater than 1, of which 5 factors were retained for our analyses on the basis of variance explained (≥5%) and scree plots. The factors explained 81.1%, 73.5%, and 74.1% variance of the original cytokine data for the Control, LPS, and LPS+CBL conditions, respectively. Across all conditions, similar cytokines consistently loaded on the same factors. Factor loading scores can be found in **Table 3.** For ease of interpretation, factors in the LPS and

LPS+CBL conditions were listed using the same numbering scheme as that of Control factors throughout the study.

[**Table 3** here]

3.4. Relationships between Adrenergic Immune Modulation and Psychiatric Measures

Across the three experimental conditions, a similar cytokine factor (major loadings: PDGF-AB/BB, sCD40L, PDGF-AA, GRO, EGF, + RANTES for Control and LPS conditions, + G-CSF for LPS condition; see **Supplementary Table 7**) was identified to be significantly correlated with anticipatory anhedonia after Bonferroni correction ($\alpha = \frac{0.05}{5 \times 4 \times 3} \approx 8.3 \times 10^{-4}$) and controlling for age, sex, and BMI. Specifically, correlations were found between anticipatory anhedonia and a) Control Factor (F) 4 (**Figure 1A**, *rho* = -0.35, *p* = 3.6 × 10⁻⁴), b) LPS F4 (**Figure 1B**, *rho* = -0.36, *p* = 2.6 × 10⁻⁴), and c) LPS+CBL F4 (**Figure 1C**, *rho* = -0.39, *p* = 7.4 × 10⁻⁵). In addition, LPS+CBL F4 significantly correlated with consummatory anhedonia (**Figure 1D**, *rho* = -0.36, *p* = 3.3 × 10⁻⁴).

[**Figure 1** here]

When depression severity was also included as a covariate, LPS+CBL F4 remained significantly correlated with anticipatory anhedonia (*rho* = -0.34, *p* = 6.3 × 10⁻⁴) and consummatory anhedonia (*rho* = -0.34, *p* = 9.9 × 10⁻⁴), while relationships between anticipatory anhedonia and Control F4 and LPS F4 did not meet the Bonferroni-corrected significant threshold $\alpha = \frac{0.05}{5 \times 2 \times 3} \approx 1.7 \times 10^{-3}$ (i.e. adjusting for number of factors, TEPS subscales, and conditions). When these analyses were repeated with total TEPS score (sum of TEPS-A and TEPS-C), these negative correlation trends were similarly observed in cytokine factors with anhedonia relationships, with LPS+CBL F4 correlating more strongly with anhedonia compared to Control F4 and LPS F4 (see **Supplementary**

Results – *SR3*). No significant relationships were found between any immune factor and depression or anxiety. All correlation results are reported in **Supplementary Table 8**.

Across all three conditions, the observed relationships between F4 and anhedonia subconstructs were significantly different from associations between anhedonia and F1, F2, and F3. Additionally, consistent across LPS and LPS+CBL conditions was the significant difference between F4 and F5 in their associations with anticipatory anhedonia. Details of these results can be found in **Supplementary Tables 10 – 11**.

3.5. Exploring Clenbuterol-Induced Differences in Immune Profiles

Despite the lack of significant associations between anhedonia scores and cytokine ratios between LPS+CBL and LPS conditions (see **Supplementary Results – *SR1*** and **Supplementary Table 9**), our primary results showing differential relationships between anhedonia and factors in these two conditions alone called for further dissection of their immune profiles. Using stringent Bonferroni correction ($\alpha = \frac{0.05}{820} \approx 6.1 \times 10^{-5}$), we found that administration of CBL significantly affected 3 inter-cytokine correlations: RANTES and PDGF-AA (Δrho = -0.18, *p* = 3.36 × 10⁻⁶); MIP-1α and MCP-1 (Δrho = -0.29, *p* = 3.5 × 10⁻⁶); and MIP-1β and MCP-1 (Δrho = -0.33, *p* = 4.74 × 10⁻⁶). At a more relaxed exploratory significance threshold using false discovery rate (FDR) correction (51), we identified CBL-induced changes in correlations between 5 additional cytokine pairs: MIP-1β and IL-2 (Δrho = 0.19, p_{FDR} = 0.030); MCP-1 and IL-6 (Δrho = 0.29, p_{FDR} = 0.036); MCP-1 and IL-8 (Δrho = -0.22, p_{FDR} = 0.021); MCP-1 and IP-10 (Δrho = -0.26, p_{FDR} = 0.028); and RANTES and TNF-α (Δrho = -0.20, p_{FDR} = 0.018). **Figure 2** visualizes the pairwise differences in *rho* (lower triangle) and corresponding significance (upper triangle) values.

[**Figure 2** here]

4. Discussion

To our knowledge, this study is the first to examine adrenergic modulation of *ex vivo* inflammatory responses using comprehensive immune biomarker assays. As hypothesized, our results indicated that CBL had a general attenuating effect on LPS-induced immune activation in a large sample of medically healthy adolescents with diverse psychiatric profiles. Further, utilizing data-driven factor analysis, we documented relationships between adrenergic mediation of cytokine production and anhedonia in youth. We discuss our findings below.

4.1. Clenbuterol Functionally Modeled Adrenergic Immune Modulation

Using a conservative statistical approach with stringent correction for multiple comparisons, we identified nine immune analytes for which LPS-induced inflammatory responses were significantly attenuated by simultaneous administration of the β_2-adrenergic receptor agonist CBL, as well as one (sCD40L) for which CBL potentiated release compared to LPS alone. Notably, IL-1β, IL-6, and TNF-α, which have been consistently implicated in psychopathology (2, 52, 53), were among the cytokines most substantially attenuated by CBL in our study. While not previously reported in clinical studies, the observed post-CBL reduced levels of IL-1β, IL-6, and TNF-α mirror results from preclinical studies examining immune activity following CBL induction (54, 55).

Another important finding from our study is the pronounced changes in levels of MIP-1α and MIP-1β chemokines across *ex vivo* conditions. MIP-1α and MIP-1β have been implicated in multiple immune-related pathologies, including those affecting the central nervous system (56). The detected increase of MIP-1α and MIP-1β levels following LPS stimulation in our study is consistent with previous reports (56); however,

their attenuated levels following CBL addition have never been studied and are novel findings. Though infrequently investigated compared to other immune biomarkers in psychopathologic contexts, MIP chemokines have been assayed in a few studies recruiting adult psychiatric participants (56-58). One postmortem study found that MIP mRNA was expressed in brain tissue samples of 4 patients with schizophrenia and 1 patient with manic depressive illness but not in those of 8 patients with dementia, neurodegenerative disease, and brain tumor, suggesting a specific role in psychiatric vs. neurological illness (59). The handful of clinical studies to examine MIP levels in psychiatric cohorts have yielded inconsistent findings, with some reporting elevated serum MIP-1β concentrations in patients with more severe depression (57, 58), but others reporting lower or no difference in MIP levels in patients with major depression compared to healthy controls (60, 61). Our study presents new evidence supporting MIP chemokines as part of the general immune response to adrenergic stimulation in youth with diverse mood and anxiety symptoms.

While CBL attenuated the release of multiple pro-inflammatory cytokines, it had minimal impact on anti-inflammatory biomarkers, including IL-4 and IL-10 (62, 63). Animal literature has provided conflicting evidence regarding adrenergic modulation of IL-10, reporting both suppression (25, 64) and induction of IL-10 following administration of β_2 receptor agonists, including CBL (65, 66). As IL-4 and IL-10 were elevated in both LPS and LPS+CBL conditions compared to Control in our study, one interpretation is that these cytokines were already being produced in response to LPS to limit the predominantly inflammatory effects of other induced cytokines and thus were not sensitive to the coadministration of CBL. Furthermore, CBL exposure led to increased

levels of sCD40L, which is a signaling molecule typically characterized as pro-inflammatory but promotes the secretion of IL-10 and exhibits immunosuppressive activity in HIV infection and certain cancers (67, 68). In section *4.2* below, we discuss alternative mechanisms potentially underlying CBL's immune-attenuation effects given the observed minimal change in IL-10 levels between LPS and LPS+CBL conditions in this study. We note that these type of complex interactions between pro- and anti-inflammatory mechanisms are characteristic of the immune system, resulting in varied observations using available measurement and analytic methods.

Similarly, several hematopoietic growth factors (i.e. G-CSF, GM-CSF, PDGF-AA/BB, TGF-α) responsive to LPS stimulus did not appear to be affected by CBL. Few studies have examined if and how hematopoietic growth factors expression react to CBL (69, 70), and virtually none addressed such questions in post-endotoxin exposure settings. The lack of substantial change observed peripherally in levels of hematopoietic growth factor following CBL induction could be explained by hematopoietic growth factors acting more centrally in psychopathologic contexts (71).

4.2. Mechanisms underlying anti-inflammatory effects by β$_2$-adrenergic signaling

Though β$_2$ agonism induces a generalized anti-inflammatory effect, responses to β$_2$ signaling by subtypes of immune cells are nuanced (21, 72). Innate immune cells with well-differentiated functions appear to be consistent in their anti-inflammatory responses to β$_2$-adrenergic receptors (AR) activation. Human monocytes primarily exert an immune-suppressive effect through reducing levels of post-LPS MIP-1α, IL-18, and IL-12 production (73, 74). Dendritic cells, though less commonly studied, have also been shown to primarily engage in anti-inflammatory activities following adrenergic modulation (21).

However, in the absence of inflammatory triggers, β_2-AR activation might signal monocytes to upregulate pro-inflammatory activities, with increased secretion of IL-18, IL-6, and free radicals (75, 76). Adaptive immune cells express β_2-AR differentially, which was thought to influence their responses to post-inflammatory adrenergic stimulation (72, 77). Naïve CD4+ T cells can differentiate into Th1 or Th2 cells in response to β_2-AR engagement depending on their cellular environment, and downstream increased IFN-γ production through mature Th1 cells has been reported (78). B cells, known for their antibody producing capabilities, can initiate a downstream pro-inflammatory effect upon adrenergic stimulation (79).

Beyond the initial responses to adrenergic stimulation, cytokines released from these activated cells trigger feedback pathways to regulate the overall production of immune markers. Cross-regulations between innate and adaptive branches of the immune system add further granularity to cellular responses following β_2-agonism (21, 72, 77). Conflicting reports of various cytokine levels being promoted or suppressed after β_2-AR stimulation emphasize the complexity of various immune modulating systems and the importance of more global approaches to examine a wide spectrum of immune markers. In our study in which immune marker levels were quantified following *in vitro* experimentation on whole blood samples, the reported adrenergic attenuation of immune activation likely reflected the net effect of these mechanisms.

Although we cannot make direct inferences about precise molecular mechanisms of CBL's downregulating effect on LPS-induced immune dysregulation from the results of this study, intracellular cyclic adenosine monophosphate (cAMP) level elevation might mediate such observed effect, as supported by an *in vitro* study of cellular events in

human peripheral blood mononuclear cells and cumulative preclinical studies on mechanisms of β_2-AR signaling (55, 80). CBL has also been shown to suppress LPS-induced nuclear factor kappa-light-chain-enhancer of activated B cells (NF-κB) activity through prevention of nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor (IκB) phosphorylation in rodent brains (81). Given that transcription factor NF-κB is ubiquitous in human cells (82), CBL possibly exerts its inflammatory-attenuating property through this modulator, similar to other β_2-AR agonists (83).

4.3. New Insights into Adrenergic-Modulated Immunologic Correlates of Anhedonia

Our exploratory analytic strategy revealed associations between anhedonia sub-components and immune measures at baseline as well as in response to endotoxin and adrenergic challenge. This finding emphasizes the advantages of employing the RDoC approach focusing on behavioral constructs and neuronal circuits rather than limiting investigations to categorical classification of psychiatric disorders. Additionally, this finding fits our prior research reporting relationships between levels of various inflammatory mediators in the blood and anhedonia, as well as their associations with brain function during a reward fMRI task (6, 18-20). Complementing our works focusing on adolescence, others have shown a correlation between an LPS-induced immune cluster and anticipatory anhedonia, indexed by TEPS-A and a reward fMRI task, in adults (84). The varied cytokines contributing to different immune factors linked to anhedonia across studies is expected from the data-driven nature of factor analyses.

Interestingly, specific to post-CBL exposure, a cluster composed predominantly of hematopoietic growth factors was linked to higher degrees of both anticipatory and consummatory anhedonia. These relationships persisted after adjusting for depression

severity. Previously, several growth factors, including G-CSF, GM-CSF, sCD40L, and EGF, have been implicated in depression (85-89). Our findings corroborated this growing literature and encouraged the utilization of dimensional measures of reward constructs in immunopsychiatric investigations beyond conventional disorder categories. While anticipatory and consummatory anhedonia quantified with clinical self-reports might share quite some variance, our study supported their divergent immunological correlates across various experimental conditions. Perhaps the distinction between anhedonia sub-components is more apparent biologically, as suggested by works from our group and others (18, 84, 90).

Although there was a lack of association between anhedonia and relative changes of immune marker levels across LPS and LPS+CBL conditions, in dissecting the inter-cytokine relationships within their immune profiles, we showed that the complex interplay between pro-inflammatory, chemokines, and hematopoietic growth factors was potentially responsible for the newly identified factor-anhedonia associations. Taken together, our findings lent support to the developing notion that immunological involvement in psychopathology might not be specific to a group of selected few markers (91, 92).

4.4. Clinical Implications of Adrenergic Modulating Interventions in Depression

Our results suggest that targeting β_2-adrenergic receptors might be a promising anti-inflammatory approach for interventions in a subset of patients suffering from depression, as β_2-agonism could dampen the effects of peripheral inflammation modulating psychiatric symptoms. Particularly, we speculate that adrenergic-targeting interventions might be favorable for patients experiencing anhedonia. This approach

should be cautiously considered as attempts to develop adrenergic-specific psychopharmacologic therapy targeting depression have been met with cardiovascular and electrolytic challenges (93), and data to date have yet to suggest a clinically meaningful effect of beta-blockers in treating anxiety disorders (94). One previous trial of CBL as a potential anti-depressant treatment was conducted in 5 female patients with major depression, and all patients experienced systemic side effects, prompting the authors to conclude CBL was not a viable treatment for depression (95). Alternatively, previous reports have linked improvement in anhedonia to existing interventions that engage sympathetic and parasympathetic systems, including deep breathing exercises, meditation, and physical exercises (96-99). This study thus provides early neurobiological evidence to support the effectiveness of such interventions. Relatedly, while TNF-α antagonist infliximab, an anti-inflammatory biologic, failed to produce significant antidepressant effects in patients with major depressive episodes, meaningful improvements in anhedonia were observed (100, 101). In line with these reports, our findings advocate for anti-inflammatory trials specifically targeting anhedonia.

4.5. Limitations and Future Directions

This study has several limitations. First, as cytokine levels in our experiment were measured at only one time point (6 hours post-exposure), temporal cytokine behaviors following treatment of LPS with and without CBL remain unclear. Future research might measure cytokine levels at several post-exposure time points to gain more comprehensive insights in CBL-affected cytokine behaviors. Second, correlations were found between adrenergic-induced immune suppression and anhedonia, thus we cannot make further inferences regarding underlying neural mechanisms that might modulate

this relationship. Since we also collect neuroimaging data on many of these participants as part of our on-going research program, we aim to incorporate measurements of reward dysfunctions in our future studies to examine CBL-affected immune response and reward-related neural activity. Third, our study design is cross-sectional, limiting our ability to infer stability of these associations over time. Finally, our analysis did not account for several factors influencing the immune system and processes that modulate it, including menstrual cycle stage, exercise, diet, stress, and sleep. These data should be collected in future studies to further control for latent confounders and parse out more specific relationships between immune attenuation and symptomatology in adolescents.

4.6. Conclusions

The results of this study further inform the research and clinical communities about CBL's anti-inflammatory effects that can potentially target neuroinflammation implicated in adolescent psychopathology. More broadly, the results of this study can pave the way for further understanding of the role of β_2-agonism underlying reward-related constructs that contribute to mood and anxiety symptomatology.

Funding and Disclosure:

This study was supported by the National Institutes of Health (NIH) under Award Numbers R21MH126501 to B.E.; RM1DA055437, R01DA054885, R21MH126501, R01MH120601, R21MH121920, and R01MH126821 to V.G. (Principal Investigator). T.N. was additionally supported by NIH grants UL1TR001073 (Clinical Research Training Program; PIs: Harry Shamoon and Marla J. Keller) and T32GM007288 (Medical Scientist Training Program; PI: Myles H. Akabas). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Conflict of Interest:

The authors declare no conflict of interest.

Author Contributions:

Conceptualization: T.N, B.E., S.K, and V.G.

Data Curation: T.N., B.E., D.P, M.P, H.X., S.K., and V.G.

Formal Analysis: T.N.

Funding Acquisition: T.N., B.E, and V.G.

Investigation: T.N., B.E., M.P, H.X., S.K., and V.G.

Methodology: T.N., B.E., M.P, H.X, S.K, and V.G.

Project Administration: T.N., B.E., S.K., and V.G.

Resources: S.K and V.G.

Software: T.N. and B.E.

Supervision: V.G.

Validation: T.N, B.E., S.K., and V.G.

Visualization: T.N.

Writing – Original Draft: T.N., B.E., S.K., and V.G.

Writing – Review & Editing: T.N, B.E., D.P, M.P, H.X., S.K., and V.G.

References

1. O'Connor TG, Moynihan JA, Caserta MT. Annual research review: The neuroinflammation hypothesis for stress and psychopathology in children--developmental psychoneuroimmunology. J Child Psychol Psychiatry. 2014;55(6):615-31. doi:10.1111/jcpp.12187

2. Najjar S, Pearlman DM, Alper K, Najjar A, Devinsky O. Neuroinflammation and psychiatric illness. J Neuroinflammation. 2013;10:43. doi:10.1186/1742-2094-10-43

3. Gabbay V, Klein RG, Alonso CM, Babb JS, Nishawala M, De Jesus G, et al. Immune system dysregulation in adolescent major depressive disorder. J Affect Disord. 2009;115(1-2):177-82. doi:10.1016/j.jad.2008.07.022

4. Gabbay V, Klein RG, Guttman LE, Babb JS, Alonso CM, Nishawala M, et al. A preliminary study of cytokines in suicidal and nonsuicidal adolescents with major depression. J Child Adolesc Psychopharmacol. 2009;19(4):423-30. doi:10.1089/cap.2008.0140

5. Gabbay V, Liebes L, Katz Y, Liu S, Mendoza S, Babb JS, et al. The kynurenine pathway in adolescent depression: preliminary findings from a proton MR spectroscopy study. Prog Neuropsychopharmacol Biol Psychiatry. 2010;34(1):37-44. doi:10.1016/j.pnpbp.2009.09.015

6. Gabbay V, Ely BA, Babb J, Liebes L. The possible role of the kynurenine pathway in anhedonia in adolescents. J Neural Transm (Vienna). 2012;119(2):253-60. doi:10.1007/s00702-011-0685-7

7. D'Acunto G, Nageye F, Zhang J, Masi G, Cortese S. Inflammatory Cytokines in Children and Adolescents with Depressive Disorders: A Systematic Review and Meta-Analysis. J Child Adolesc Psychopharmacol. 2019;29(5):362-9. doi:10.1089/cap.2019.0015

8. Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. Nat Rev Neurosci. 2008;9(1):46-56. doi:10.1038/nrn2297

9. Miller AH, Haroon E, Raison CL, Felger JC. Cytokine targets in the brain: impact on neurotransmitters and neurocircuits. Depress Anxiety. 2013;30(4):297-306. doi:10.1002/da.22084

10. Ely BA, Nguyen TNB, Tobe RH, Walker AM, Gabbay V. Multimodal Investigations of Reward Circuitry and Anhedonia in Adolescent Depression. Frontiers in Psychiatry. 2021;12(1136). doi:10.3389/fpsyt.2021.678709

11. Gabbay V, Coffey BJ, Guttman LE, Gottlieb L, Katz Y, Babb JS, et al. A cytokine study in children and adolescents with Tourette's disorder. Prog Neuropsychopharmacol Biol Psychiatry. 2009;33(6):967-71. doi:10.1016/j.pnpbp.2009.05.001

12. Dantzer R, Kelley KW. Twenty years of research on cytokine-induced sickness behavior. Brain Behav Immun. 2007;21(2):153-60. doi:10.1016/j.bbi.2006.09.006

13. De La Garza R, 2nd. Endotoxin- or pro-inflammatory cytokine-induced sickness behavior as an animal model of depression: focus on anhedonia. Neurosci Biobehav Rev. 2005;29(4-5):761-70. doi:10.1016/j.neubiorev.2005.03.016

14. Konsman JP, Parnet P, Dantzer R. Cytokine-induced sickness behaviour: mechanisms and implications. Trends Neurosci. 2002;25(3):154-9. doi:10.1016/s0166-2236(00)02088-9

15. Dantzer R. Cytokine-induced sickness behavior: mechanisms and implications. Ann N Y Acad Sci. 2001;933:222-34. doi:10.1111/j.1749-6632.2001.tb05827.x

16. Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, et al. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. Am J Psychiatry. 2010;167(7):748-51. doi:10.1176/appi.ajp.2010.09091379

17. Freed RD, Mehra LM, Laor D, Patel M, Alonso CM, Kim-Schulze S, et al. Anhedonia as a clinical correlate of inflammation in adolescents across psychiatric conditions. World J Biol Psychiatry. 2019;20(9):712-22. doi:10.1080/15622975.2018.1482000

18. Bradley KA, Stern ER, Alonso CM, Xie H, Kim-Schulze S, Gabbay V. Relationships between neural activation during a reward task and peripheral cytokine levels in youth with diverse psychiatric symptoms. Brain Behav Immun. 2019;80:374-83. doi:10.1016/j.bbi.2019.04.014

19. Liu Q, Ely B, Simkovic S, Tao A, Wolchok R, Alonso CM, et al. Correlates of C-reactive protein with neural reward circuitry in adolescents with psychiatric symptoms. Brain Behav Immun Health. 2020;9. doi:10.1016/j.bbih.2020.100153

20. DeWitt SJ, Bradley KA, Lin N, Yu C, Gabbay V. A pilot resting-state functional connectivity study of the kynurenine pathway in adolescents with depression and healthy controls. J Affect Disord. 2018;227:752-8. doi:10.1016/j.jad.2017.11.040

21. Scanzano A, Cosentino M. Adrenergic regulation of innate immunity: a review. Front Pharmacol. 2015;6:171. doi:10.3389/fphar.2015.00171

22. De Paermentier F, Cheetham SC, Crompton MR, Katona CL, Horton RW. Brain beta-adrenoceptor binding sites in antidepressant-free depressed suicide victims. Brain Res. 1990;525(1):71-7. doi:10.1016/0006-8993(90)91321-7

23. Biegon A, Israeli M. Regionally selective increases in β-adrenergic receptor density in the brains of suicide victims. Brain research. 1988;442(1):199-203. doi:10.1016/0006-8993(88)91453-9

24. Hasler G, Luckenbaugh DA, Snow J, Meyers N, Waldeck T, Geraci M, et al. Reward processing after catecholamine depletion in unmedicated, remitted subjects with major depressive disorder. Biol Psychiatry. 2009;66(3):201-5. doi:10.1016/j.biopsych.2009.02.029

25. Lechtenberg KJ, Meyer ST, Doyle JB, Peterson TC, Buckwalter MS. Augmented beta2-adrenergic signaling dampens the neuroinflammatory response following ischemic stroke and increases stroke size. J Neuroinflammation. 2019;16(1):112. doi:10.1186/s12974-019-1506-4

26. Zhang HT, Huang Y, O'Donnell JM. Antagonism of the antidepressant-like effects of clenbuterol by central administration of beta-adrenergic antagonists in rats. Psychopharmacology (Berl). 2003;170(1):102-7. doi:10.1007/s00213-003-1512-0

27. American Psychiatric Association A. Diagnostic and statistical manual of mental disorders (DSM-IV). 1994.

28. American Psychiatric Association A. Diagnostic and statistical manual of mental disorders (DSM-5®). 2013.

29. Kaufman AS. Manual for the Kaufman brief intelligence test. Circle Pines, MN: American Guidance Service. 1990.

30. Kaufman J, Birmaher B, Brent D, Rao U, Flynn C, Moreci P, et al. Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime

Version (K-SADS-PL): initial reliability and validity data. J Am Acad Child Adolesc Psychiatry. 1997;36(7):980-8. doi:10.1097/00004583-199707000-00021

31. March JS, Parker JD, Sullivan K, Stallings P, Conners CK. The Multidimensional Anxiety Scale for Children (MASC): factor structure, reliability, and validity. J Am Acad Child Adolesc Psychiatry. 1997;36(4):554-65. doi:10.1097/00004583-199704000-00019

32. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. Arch Gen Psychiatry. 1961;4:561-71. doi:10.1001/archpsyc.1961.01710120031004

33. Gard DE, Gard MG, Kring AM, John OP. Anticipatory and consummatory components of the experience of pleasure: a scale development study. Journal of research in personality. 2006;40(6):1086-102.

34. Gooding DC, Pflum MJ, Fonseca-Pedero E, Paino M. Assessing social anhedonia in adolescence: The ACIPS-A in a community sample. Eur Psychiatry. 2016;37:49-55. doi:10.1016/j.eurpsy.2016.05.012

35. Ho PM, Cooper AJ, Hall PJ, Smillie LD. Factor structure and construct validity of the temporal experience of pleasure scales. J Pers Assess. 2015;97(2):200-8. doi:10.1080/00223891.2014.940625

36. Rossol M, Heine H, Meusch U, Quandt D, Klein C, Sweet MJ, et al. LPS-induced cytokine production in human monocytes and macrophages. Crit Rev Immunol. 2011;31(5):379-446. doi:10.1615/critrevimmunol.v31.i5.20

37. dupont NC, Wang K, Wadhwa PD, Culhane JF, Nelson EL. Validation and comparison of luminex multiplex cytokine analysis kits with ELISA: determinations of a panel of nine cytokines in clinical sample culture supernatants. J Reprod Immunol. 2005;66(2):175-91. doi:10.1016/j.jri.2005.03.005

38. Elshal MF, McCoy JP. Multiplex bead array assays: performance evaluation and comparison of sensitivity to ELISA. Methods. 2006;38(4):317-23. doi:10.1016/j.ymeth.2005.11.010

39. Leng SX, McElhaney JE, Walston JD, Xie D, Fedarko NS, Kuchel GA. ELISA and multiplex technologies for cytokine measurement in inflammation and aging research. J Gerontol A Biol Sci Med Sci. 2008;63(8):879-84. doi:10.1093/gerona/63.8.879

40. Tighe PJ, Ryder RR, Todd I, Fairclough LC. ELISA in the multiplex era: potentials and pitfalls. Proteomics Clin Appl. 2015;9(3-4):406-22. doi:10.1002/prca.201400130

41. Richens JL, Urbanowicz RA, Metcalf R, Corne J, O'Shea P, Fairclough L. Quantitative validation and comparison of multiplex cytokine kits. J Biomol Screen. 2010;15(5):562-8. doi:10.1177/1087057110362099

42. Breen EJ, Polaskova V, Khan A. Bead-based multiplex immuno-assays for cytokines, chemokines, growth factors and other analytes: median fluorescence intensities versus their derived absolute concentration values for statistical analysis. Cytokine. 2015;71(2):188-98. doi:10.1016/j.cyto.2014.10.030

43. Breen EJ, Tan W, Khan A. The Statistical Value of Raw Fluorescence Signal in Luminex xMAP Based Multiplex Immunoassays. Sci Rep. 2016;6:26996. doi:10.1038/srep26996

44. Fritz CO, Morris PE, Richler JJ. Effect size estimates: current use, calculations, and interpretation. J Exp Psychol Gen. 2012;141(1):2-18. doi:10.1037/a0024338

45. Rosenthal R, DiMatteo MR. Meta-analysis: recent developments in quantitative methods for literature reviews. Annu Rev Psychol. 2001;52:59-82. doi:10.1146/annurev.psych.52.1.59

46. Ellis PD. The essential guide to effect sizes : statistical power, meta-analysis, and the interpretation of research results. Cambridge ; New York: Cambridge University Press; 2010.

47. Tomczak M, Tomczak E. The need to report effect size estimates revisited. An overview of some recommended measures of effect size. 2014.

48. Efron B, Tibshirani RJ. An introduction to the bootstrap: CRC press; 1994.

49. Van Snellenberg JX, de Candia T. Meta-analytic evidence for familial coaggregation of schizophrenia and bipolar disorder. Arch Gen Psychiatry. 2009;66(7):748-55. doi:10.1001/archgenpsychiatry.2009.64

50. Van Snellenberg JX. BCa_bootstrap. MATLAB, MATLAB Central File Exchange. 2018. https://www.mathworks.com/matlabcentral/fileexchange/69119. Accessed April 25 2021.

51. Benjamini Y, Krieger AM, Yekutieli D. Adaptive linear step-up procedures that control the false discovery rate. Biometrika. 2006;93(3):491-507.

52. Miller AH, Maletic V, Raison CL. Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. Biol Psychiatry. 2009;65(9):732-41. doi:10.1016/j.biopsych.2008.11.029

53. Mitchell RH, Goldstein BI. Inflammation in children and adolescents with neuropsychiatric disorders: a systematic review. J Am Acad Child Adolesc Psychiatry. 2014;53(3):274-96. doi:10.1016/j.jaac.2013.11.013

54. Izeboud CA, Monshouwer M, van Miert AS, Witkamp RF. The beta-adrenoceptor agonist clenbuterol is a potent inhibitor of the LPS-induced production of TNF-alpha and IL-6 in vitro and in vivo. Inflamm Res. 1999;48(9):497-502. doi:10.1007/s000110050493

55. Yoshimura T, Kurita C, Nagao T, Usami E, Nakao T, Watanabe S, et al. Inhibition of tumor necrosis factor-alpha and interleukin-1-beta production by beta-adrenoceptor agonists from lipopolysaccharide-stimulated human peripheral blood mononuclear cells. Pharmacology. 1997;54(3):144-52. doi:10.1159/000139481

56. Menten P, Wuyts A, Van Damme J. Macrophage inflammatory protein-1. Cytokine Growth Factor Rev. 2002;13(6):455-81. doi:10.1016/s1359-6101(02)00045-x

57. Hoge EA, Brandstetter K, Moshier S, Pollack MH, Wong KK, Simon NM. Broad spectrum of cytokine abnormalities in panic disorder and posttraumatic stress disorder. Depress Anxiety. 2009;26(5):447-55. doi:10.1002/da.20564

58. Oglodek EA, Just MJ. The association between inflammatory markers (iNOS, HO-1, IL-33, MIP-1beta) and depression with and without posttraumatic stress disorder. Pharmacol Rep. 2018;70(6):1065-72. doi:10.1016/j.pharep.2018.06.001

59. Ishizuka K, Igata-Yi R, Kimura T, Hieshima K, Kukita T, Kin Y, et al. Expression and distribution of CC chemokine macrophage inflammatory protein-1 alpha/LD78 in the human brain. Neuroreport. 1997;8(5):1215-8. doi:10.1097/00001756-199703240-00031

60. Lehto SM, Niskanen L, Herzig KH, Tolmunen T, Huotari A, Viinamaki H, et al. Serum chemokine levels in major depressive disorder. Psychoneuroendocrinology. 2010;35(2):226-32. doi:10.1016/j.psyneuen.2009.06.007

61. Cassano P, Bui E, Rogers AH, Walton ZE, Ross R, Zeng M, et al. Inflammatory cytokines in major depressive disorder: A case-control study. Aust N Z J Psychiatry. 2017;51(1):23-31. doi:10.1177/0004867416652736

62. Moore KW, de Waal Malefyt R, Coffman RL, O'Garra A. Interleukin-10 and the interleukin-10 receptor. Annu Rev Immunol. 2001;19:683-765. doi:10.1146/annurev.immunol.19.1.683

63. Opal SM, DePalo VA. Anti-inflammatory cytokines. Chest. 2000;117(4):1162-72. doi:10.1378/chest.117.4.1162

64. van den Hoven R, Duvigneau JC, Hartl RT, Gemeiner M. Clenbuterol affects the expression of messenger RNA for interleukin 10 in peripheral leukocytes from horses challenged intrabronchially with lipopolysaccharides. Vet Res Commun. 2006;30(8):921-8. doi:10.1007/s11259-006-3383-4

65. McNamee EN, Ryan KM, Griffin EW, Gonzalez-Reyes RE, Ryan KJ, Harkin A, et al. Noradrenaline acting at central beta-adrenoceptors induces interleukin-10 and suppressor of cytokine signaling-3 expression in rat brain: implications for neurodegeneration. Brain Behav Immun. 2010;24(4):660-71. doi:10.1016/j.bbi.2010.02.005

66. Agac D, Estrada LD, Maples R, Hooper LV, Farrar JD. The beta2-adrenergic receptor controls inflammation by driving rapid IL-10 secretion. Brain Behav Immun. 2018;74:176-85. doi:10.1016/j.bbi.2018.09.004

67. Jenabian MA, Patel M, Kema I, Vyboh K, Kanagaratham C, Radzioch D, et al. Soluble CD40-ligand (sCD40L, sCD154) plays an immunosuppressive role via regulatory T cell expansion in HIV infection. Clin Exp Immunol. 2014;178(1):102-11. doi:10.1111/cei.12396

68. Schlom J, Jochems C, Gulley JL, Huang J. The role of soluble CD40L in immunosuppression. Oncoimmunology. 2013;2(1):e22546. doi:10.4161/onci.22546

69. Culmsee C, Stumm RK, Schafer MK, Weihe E, Krieglstein J. Clenbuterol induces growth factor mRNA, activates astrocytes, and protects rat brain tissue against ischemic damage. Eur J Pharmacol. 1999;379(1):33-45. doi:10.1016/s0014-2999(99)00452-5

70. Sharma AN, da Costa e Silva BF, Soares JC, Carvalho AF, Quevedo J. Role of trophic factors GDNF, IGF-1 and VEGF in major depressive disorder: A comprehensive review of human studies. J Affect Disord. 2016;197:9-20. doi:10.1016/j.jad.2016.02.067

71. Diederich K, Schabitz WR, Minnerup J. Seeing old friends from a different angle: novel properties of hematopoietic growth factors in the healthy and diseased brain. Hippocampus. 2012;22(5):1051-7. doi:10.1002/hipo.20904

72. Kohm AP, Sanders VM. Norepinephrine and beta 2-adrenergic receptor stimulation regulate CD4+ T and B lymphocyte function in vitro and in vivo. Pharmacol Rev. 2001;53(4):487-525.

73. Mizuno K, Takahashi HK, Iwagaki H, Katsuno G, Kamurul HA, Ohtani S, et al. Beta2-adrenergic receptor stimulation inhibits LPS-induced IL-18 and IL-12 production in monocytes. Immunol Lett. 2005;101(2):168-72. doi:10.1016/j.imlet.2005.05.008

74. Li CY, Chou TC, Lee CH, Tsai CS, Loh SH, Wong CS. Adrenaline inhibits lipopolysaccharide-induced macrophage inflammatory protein-1 alpha in human monocytes: the role of beta-adrenergic receptors. Anesth Analg. 2003;96(2):518-23, table of contents. doi:10.1097/00000539-200302000-00040

75. Paul-Eugene N, Kolb JP, Damais C, Abadie A, Mencia-Huerta JM, Braquet P, et al. Beta 2-adrenoceptor agonists regulate the IL-4-induced phenotypical changes and IgE-dependent functions in normal human monocytes. J Leukoc Biol. 1994;55(3):313-20. doi:10.1002/jlb.55.3.313

76. Takahashi HK, Morichika T, Iwagaki H, Yoshino T, Tamura R, Saito S, et al. Effect of beta 2-adrenergic receptor stimulation on interleukin-18-induced intercellular adhesion molecule-1 expression and cytokine production. J Pharmacol Exp Ther. 2003;304(2):634-42. doi:10.1124/jpet.102.042622

77. Sanders VM. The beta2-adrenergic receptor on T and B lymphocytes: do we understand it yet? Brain Behav Immun. 2012;26(2):195-200. doi:10.1016/j.bbi.2011.08.001

78. Swanson MA, Lee WT, Sanders VM. IFN-gamma production by Th1 cells generated from naive CD4+ T cells exposed to norepinephrine. J Immunol. 2001;166(1):232-40. doi:10.4049/jimmunol.166.1.232

79. Wu L, Tai Y, Hu S, Zhang M, Wang R, Zhou W, et al. Bidirectional Role of beta2-Adrenergic Receptor in Autoimmune Diseases. Front Pharmacol. 2018;9:1313. doi:10.3389/fphar.2018.01313

80. Lefkowitz RJ, Stadel JM, Caron MG. Adenylate cyclase-coupled beta-adrenergic receptors: structure and mechanisms of activation and desensitization. Annu Rev Biochem. 1983;52:159-86. doi:10.1146/annurev.bi.52.070183.001111

81. Ryan KJ, Griffin E, Yssel JD, Ryan KM, McNamee EN, Harkin A, et al. Stimulation of central beta2-adrenoceptors suppresses NFkappaB activity in rat brain: a role for IkappaB. Neurochem Int. 2013;63(5):368-78. doi:10.1016/j.neuint.2013.07.006

82. Pahl HL. Activators and target genes of Rel/NF-kappaB transcription factors. Oncogene. 1999;18(49):6853-66. doi:10.1038/sj.onc.1203239

83. Kolmus K, Tavernier J, Gerlo S. beta2-Adrenergic receptors in immunity and inflammation: stressing NF-kappaB. Brain Behav Immun. 2015;45:297-310. doi:10.1016/j.bbi.2014.10.007

84. Costi S, Morris LS, Collins A, Fernandez NF, Patel M, Xie H, et al. Peripheral immune cell reactivity and neural response to reward in patients with depression and anhedonia. Transl Psychiatry. 2021;11(1):565. doi:10.1038/s41398-021-01668-1

85. Becerril-Villanueva E, Perez-Sanchez G, Alvarez-Herrera S, Giron-Perez MI, Arreola R, Cruz-Fuentes C, et al. Alterations in the Levels of Growth Factors in Adolescents with Major Depressive Disorder: A Longitudinal Study during the Treatment with Fluoxetine. Mediators Inflamm. 2019;2019:9130868. doi:10.1155/2019/9130868

86. Dahl J, Ormstad H, Aass HC, Malt UF, Bendz LT, Sandvik L, et al. The plasma levels of various cytokines are increased during ongoing depression and are reduced to normal levels after recovery. Psychoneuroendocrinology. 2014;45:77-86. doi:10.1016/j.psyneuen.2014.03.019

87. Idemoto K, Ishima T, Niitsu T, Hata T, Yoshida S, Hattori K, et al. Platelet-derived growth factor BB: A potential diagnostic blood biomarker for differentiating bipolar disorder from major depressive disorder. J Psychiatr Res. 2021;134:48-56. doi:10.1016/j.jpsychires.2020.12.051

88. Leo R, Di Lorenzo G, Tesauro M, Razzini C, Forleo GB, Chiricolo G, et al. Association between enhanced soluble CD40 ligand and proinflammatory and prothrombotic states in major depressive disorder: pilot observations on the effects of

selective serotonin reuptake inhibitor therapy. J Clin Psychiatry. 2006;67(11):1760-6. doi:10.4088/jcp.v67n1114

89. Memon AA, Sundquist K, Ahmad A, Wang X, Hedelius A, Sundquist J. Role of IL-8, CRP and epidermal growth factor in depression and anxiety patients treated with mindfulness-based therapy or cognitive behavioral therapy in primary health care. Psychiatry Res. 2017;254:311-6. doi:10.1016/j.psychres.2017.05.012

90. Ely BA, Liu Q, DeWitt SJ, Mehra LM, Alonso CM, Gabbay V. Data-driven parcellation and graph theory analyses to study adolescent mood and anxiety symptoms. Transl Psychiatry. 2021;11(1):266. doi:10.1038/s41398-021-01321-x

91. Himmerich H, Patsalos O, Lichtblau N, Ibrahim MAA, Dalton B. Cytokine Research in Depression: Principles, Challenges, and Open Questions. Front Psychiatry. 2019;10:30. doi:10.3389/fpsyt.2019.00030

92. Audet MC, Anisman H. Interplay between pro-inflammatory cytokines and growth factors in depressive illnesses. Front Cell Neurosci. 2013;7:68. doi:10.3389/fncel.2013.00068

93. Millan MJ. Multi-target strategies for the improved treatment of depressive states: Conceptual foundations and neuronal substrates, drug discovery and therapeutic application. Pharmacol Ther. 2006;110(2):135-370. doi:10.1016/j.pharmthera.2005.11.006

94. Steenen SA, van Wijk AJ, van der Heijden GJ, van Westrhenen R, de Lange J, de Jongh A. Propranolol for the treatment of anxiety disorders: Systematic review and meta-analysis. J Psychopharmacol. 2016;30(2):128-39. doi:10.1177/0269881115612236

95. Wiegand M, Schreiber W, Lauer C, Berger M. The action of clenbuterol on sleep and symptomatology in depressives. Pharmacopsychiatry. 1991;24(3):89-92. doi:10.1055/s-2007-1014446

96. Muzik O, Diwadkar VA. Hierarchical control systems for the regulation of physiological homeostasis and affect: Can their interactions modulate mood and anhedonia? Neurosci Biobehav Rev. 2019;105:251-61. doi:10.1016/j.neubiorev.2019.08.015

97. Johnson DP, Penn DL, Fredrickson BL, Meyer PS, Kring AM, Brantley M. Loving-kindness meditation to enhance recovery from negative symptoms of schizophrenia. J Clin Psychol. 2009;65(5):499-509. doi:10.1002/jclp.20591

98. Toups M, Carmody T, Greer T, Rethorst C, Grannemann B, Trivedi MH. Exercise is an effective treatment for positive valence symptoms in major depression. J Affect Disord. 2017;209:188-94. doi:10.1016/j.jad.2016.08.058

99. Leventhal AM. Relations between anhedonia and physical activity. Am J Health Behav. 2012;36(6):860-72. doi:10.5993/AJHB.36.6.12

100. Lee Y, Mansur RB, Brietzke E, Carmona NE, Subramaniapillai M, Pan Z, et al. Efficacy of adjunctive infliximab vs. placebo in the treatment of anhedonia in bipolar I/II depression. Brain Behav Immun. 2020;88:631-9. doi:10.1016/j.bbi.2020.04.063

101. Raison CL, Rutherford RE, Woolwine BJ, Shuo C, Schettler P, Drake DF, et al. A randomized controlled trial of the tumor necrosis factor antagonist infliximab for treatment-resistant depression: the role of baseline inflammatory biomarkers. JAMA Psychiatry. 2013;70(1):31-41. doi:10.1001/2013.jamapsychiatry.4

Table 1

Measure	HC (n = 33)	Clinical (n = 97)	Whole Sample (N = 130)	p value
Demographics				
Age (Mean ± SD)	15.06 ± 2.41	15.31 ± 2.08	15.25 ± 2.16	0.48
Sex	F = 17, M = 16	F = 60, M = 37	F = 77, M = 53	0.30
Race [a]	W = 15, B = 13, A = 0, O = 5	W = 41, B = 28, A = 5, O = 23	W = 56, B = 41, A = 5, O = 28	0.40
Ethnicity [b]	H = 9, N = 24	H = 51, N = 46	H = 60, N = 70	**0.012***
Clinical Profile				
BDI (Mean ± SD)	2.06 ± 2.54 (n = 33)	14.67 ± 12.31 (n = 94)	11.39 ± 12.01 (N = 127)	**<0.01***
MASC (Mean ± SD)	27.81 ± 12.89 (n = 31)	44.29 ± 17.58 (n = 92)	40.14 ± 17.98 (N = 123)	**<0.01***
TEPS-A (Mean ± SD)	48.20 ± 8.52 (n = 30)	44.24 ± 9.50 (n = 74)	45.38 ± 9.36 (N = 104)	0.10
TEPS-C (Mean ± SD)	36.76 ± 8.10 (n = 29)	33.54 ± 7.28 (n = 72)	34.47 ± 7.62 (N = 101)	0.58
BMI (Mean ± SD)	24.27 ± 6.02 (n = 32)	25.60 ± 7.19 (n = 97)	25.27 ± 6.92 (N = 129)	0.50
Most Common DSM Diagnoses				
Mood Disorders [c]	0	58	58	–
Anxiety Disorders [c]	0	54	54	–
ADHD [c]	0	35	35	–
ODD [c]	0	12	12	–
PTSD [c]	0	9	9	–
Comorbidity [c, d]	N = 33, Y = 0	N = 43, Y = 54	N = 76, Y = 54	–

Table 1. Demographic and clinical characteristics of study participants. Reported p values are of appropriate bivariate tests comparing the clinical and healthy control groups.
[a] A = Asian, B = Black, O = Other/Mixed Race, W = White
[b] H = Hispanic, N = Non-Hispanic
[c] Includes past and/or subthreshold symptoms
[d] N = 0-1 diagnosis, Y = ≥ 2 diagnoses
*p < 0.05
Abbreviations: ADHD: Attention-deficit/hyperactivity disorder; BDI: Beck Depression Inventory; BMI: body mass index (kg/m²); Clinical: participants with psychiatric symptoms and/or diagnoses; DSM: Diagnostic and Statistical Manual of Mental Disorders; HC: healthy control participants; MASC: Multidimensional Anxiety Scale for Children; ODD: Oppositional Defiant Disorder; PTSD: Post Traumatic Stress Disorder; SD: standard deviation; TEPS-A: Temporal Experience of Pleasure Scale – Anticipatory; TEPS-C: Temporal Experience of Pleasure Scale – Consummatory.

Table 2

Cytokine	LPS vs. Control		LPS+CBL vs. LPS		LPS+CBL vs. Control	
	r	*p* value	r	*p* value	r	*p* value
Fractalkine	0.40	**3.5×10^{-6} ***	- 0.21	1.7×10^{-2}	0.24	5.3×10^{-3}
G-CSF	0.34	**9.5×10^{-5} ***	- 0.10	2.3×10^{-1}	0.23	1.0×10^{-2}
GM-CSF	0.40	**3.5×10^{-6} ***	- 0.15	6.9×10^{-2}	0.30	**6.3×10^{-4} ***
GRO	0.42	**1.0×10^{-6} ***	0.22	1.1×10^{-2}	0.42	**6.9×10^{-7} ***
IL-10	0.54	**1.2×10^{-10} ***	- 0.01	7.7×10^{-1}	0.54	**1.1×10^{-10} ***
IL-12P40	0.36	**3.7×10^{-5} ***	- 0.22	1.0×10^{-2}	0.23	9.5×10^{-3}
IL-1RA	0.73	**5.2×10^{-21} ****	- 0.47	**3.7×10^{-8} ***	0.57	**8.2×10^{-12} ****
IL-1α	0.41	**2.2×10^{-6} ***	- 0.21	1.2×10^{-2}	0.31	**3.1×10^{-4} ***
IL-1β	0.82	**7.8×10^{-29} ****	- 0.51	**1.0×10^{-9} ***	0.77	**3.4×10^{-24} ****
IL-4	0.43	**4.4×10^{-7} ***	0.039	6.6×10^{-1}	0.39	**6.8×10^{-6} ***
IL-6	0.82	**8.2×10^{-29} ****	- 0.62	**2.6×10^{-14} ****	0.76	**7.0×10^{-23} ****
IL-7	0.42	**1.6×10^{-6} ***	- 0.12	1.8×10^{-1}	0.36	**3.5×10^{-5} ***
IL-8	0.82	**3.4×10^{-28} ****	- 0.26	2.8×10^{-3}	0.79	**3.0×10^{-25} ****
IP-10	0.71	**3.5×10^{-19} ****	- 0.56	**2.4×10^{-11} ****	0.35	**4.8×10^{-5} ***
MCP-1	0.71	**2.6×10^{-19} ****	- 0.60	**4.8×10^{-13} ****	0.37	**2.3×10^{-5} ***
MIP-1α	0.77	**3.7×10^{-23} ****	- 0.65	**8.3×10^{-16} ****	0.64	**3.0×10^{-15} ****
MIP-1β	0.82	**1.2×10^{-28} ****	- 0.73	**2.4×10^{-20} ****	0.72	**8.7×10^{-20} ****
PDGF-AB/BB	0.23	8.0×10^{-3}	0.21	1.5×10^{-2}	0.33	**1.8×10^{-4} ***
sCD40L	0.21	1.5×10^{-2}	0.33	**2.0×10^{-4} ***	0.28	1.4×10^{-3}
TGF-α	0.65	**8.6×10^{-16} ****	- 0.36	**2.7×10^{-5} ***	0.54	**1.1×10^{-10} ***
TNF-α	0.84	**8.6×10^{-32} ****	- 0.70	**1.0×10^{-18} ****	0.78	**1.7×10^{-24} ****

Table 2. *Post hoc* pairwise comparisons of cytokine levels across 3 *ex vivo* conditions in whole sample (N=130).

*: significant at $\alpha = \frac{0.05}{21 \times 3} \approx 7.9 \times 10^{-4}$ (Bonferroni correction)

**: significant at $\alpha = 1.0 \times 10^{-10}$

r: effect size computed from nonparametric test statistics

Table 3

Cytokine	Control					LPS					LPS + CBL				
	F1	F2	F3	F4	F5	F1	F2	F3	F4	F5	F1	F2	F3	F4	F5
IL-3	**0.956**	0.071	0.021	-0.004	0.047	**0.871**	0.084	0.093	0.084	0.029	**0.882**	0.101	0.082	0.108	0.002
IL-12P70	**0.945**	0.193	0.104	-0.049	0.056	**0.778**	0.466	0.071	0.061	-0.004	**0.754**	0.447	0.203	0.108	0.041
Flt-3L	**0.943**	0.083	0.092	-0.012	0.058	**0.737**	0.277	0.035	0.144	-0.021	**0.663**	0.312	0.179	0.214	0.016
FGF-2	**0.942**	0.137	0.095	0.077	0.045	**0.747**	0.401	0.013	0.285	-0.032	**0.710**	0.366	0.142	0.372	0.015
IFN-α2	**0.919**	0.192	0.152	0.061	0.103	**0.735**	0.367	0.087	0.234	0.076	**0.704**	0.330	0.130	0.275	0.146
IL-7	**0.916**	0.189	0.135	0.165	0.111	**0.653**	0.425	0.192	0.370	0.093	**0.537**	0.520	0.134	0.333	0.148
VEGF	**0.913**	0.332	0.124	0.070	0.093	**0.761**	0.537	0.087	0.164	0.093	**0.724**	0.540	0.163	0.220	0.113
IL-15	**0.877**	0.277	-0.024	-0.068	0.150	**0.687**	0.364	-0.022	-0.028	0.245	**0.697**	0.411	-0.106	-0.017	0.191
IL-12P40	**0.870**	0.306	0.043	-0.097	0.146	**0.764**	0.297	0.101	-0.127	0.204	**0.747**	0.386	-0.041	-0.063	0.178
IL-2	**0.863**	0.249	0.039	0.098	-0.005	**0.859**	0.139	-0.024	0.099	0.013	**0.882**	0.122	0.073	0.143	0.004
GM-CSF	**0.856**	0.353	0.264	0.020	0.117	0.562	**0.718**	0.202	0.082	0.051	0.541	**0.729**	0.179	0.129	0.094
G-CSF	**0.825**	0.097	0.070	0.317	0.133	-0.047	0.106	-0.022	**0.269**	0.028	0.551	0.513	0.119	0.469	0.099
IFN-γ	**0.820**	0.185	0.222	0.006	0.051	**0.734**	0.247	0.182	0.017	0.029	**0.734**	0.223	0.295	0.062	0.047
IL-10	**0.809**	0.173	0.046	-0.093	0.093	**0.622**	0.183	0.366	-0.175	0.163	**0.625**	0.316	0.252	-0.109	0.092
IL-17A	**0.762**	0.399	0.120	-0.038	-0.025	**0.621**	0.531	0.116	-0.005	-0.058	**0.648**	0.425	0.254	0.045	-0.036
Fractalkine	**0.694**	0.152	0.049	0.163	0.045	**0.476**	0.230	0.015	0.300	-0.004	**0.391**	0.197	0.050	0.333	0.200
MDC	**0.202**	0.055	0.049	0.100	-0.053	0.186	**0.644**	-0.008	0.079	-0.098	0.103	**0.474**	-0.071	0.056	-0.086
MCP-3	0.269	**0.938**	-0.011	0.025	0.100	0.426	**0.816**	-0.032	0.053	0.221	0.483	**0.794**	-0.083	0.016	0.211
IL-13	0.293	**0.930**	-0.033	0.048	0.172	0.419	**0.797**	-0.097	0.080	0.337	0.486	**0.756**	-0.121	0.036	0.317
TNF-β	0.249	**0.912**	-0.035	0.036	0.294	0.392	**0.759**	-0.110	0.085	0.442	0.456	**0.717**	-0.125	0.031	0.439
IL-1RA	0.311	**0.910**	0.056	0.067	0.153	0.489	**0.722**	0.082	0.052	0.257	0.540	**0.709**	0.000	0.012	0.261
IL-5	0.315	**0.891**	-0.053	0.072	0.145	0.433	**0.748**	-0.104	0.113	0.299	0.496	**0.697**	-0.138	0.051	0.287
TGF-α	0.467	**0.761**	0.089	-0.051	-0.012	0.424	**0.845**	0.070	-0.006	0.015	0.428	**0.840**	0.110	0.025	0.024
Eotaxin	0.304	**0.718**	0.227	0.079	-0.016	0.500	**0.517**	-0.072	0.090	0.107	**0.497**	0.494	-0.072	0.016	0.078
IL-8	0.125	**0.692**	0.398	-0.039	-0.052	0.192	**0.695**	0.237	-0.034	-0.090	0.143	**0.762**	0.242	0.009	-0.057
MIP-1β	0.019	0.070	**0.960**	0.010	0.045	0.069	-0.003	**0.860**	-0.122	0.014	-0.008	0.120	**0.750**	-0.058	0.049
TNF-α	0.164	0.005	**0.948**	0.010	0.005	0.231	0.006	**0.859**	0.085	-0.047	0.280	0.000	**0.837**	0.021	-0.052
MIP-1α	0.010	0.021	**0.911**	0.054	0.041	0.086	0.019	**0.905**	-0.067	0.033	0.053	0.106	**0.792**	-0.008	0.042
IL-1β	0.392	0.131	**0.836**	-0.058	0.034	0.287	0.072	**0.762**	-0.170	-0.043	0.202	0.067	**0.751**	-0.131	-0.041
IL-6	0.099	0.305	**0.762**	-0.049	0.489	0.291	**0.747**	0.450	-0.011	0.046	0.308	**0.793**	0.352	0.017	0.069
MCP-1	0.097	0.617	**0.527**	-0.032	-0.014	0.064	**0.638**	0.620	-0.026	-0.069	0.084	**0.752**	0.449	0.048	-0.037
IP-10	0.118	0.005	**0.362**	-0.079	-0.092	-0.169	0.087	**0.677**	-0.028	-0.024	0.093	-0.034	**0.768**	-0.046	-0.037
PDGF-AA	0.114	0.021	0.018	**0.942**	-0.032	0.262	-0.116	-0.051	**0.881**	-0.059	0.202	-0.099	-0.020	**0.896**	-0.018
sCD40L	0.089	-0.066	-0.058	**0.928**	-0.032	0.165	-0.168	-0.065	**0.893**	-0.054	0.097	-0.157	-0.060	**0.915**	-0.040
PDGF-AB/BB	0.065	-0.003	-0.045	**0.928**	-0.032	0.142	-0.080	-0.081	**0.913**	-0.051	0.031	-0.048	-0.059	**0.920**	-0.049
GRO	0.071	0.030	-0.007	**0.902**	-0.026	0.160	0.192	-0.039	**0.826**	-0.070	0.074	0.263	-0.021	**0.851**	-0.029
EGF	0.156	0.430	-0.057	**0.731**	0.022	0.140	0.237	-0.112	**0.700**	0.103	0.173	0.259	-0.145	**0.718**	0.075
RANTES	-0.200	-0.100	-0.033	**0.387**	0.326	-0.250	-0.067	0.040	**0.457**	0.294	**-0.397**	0.034	-0.044	0.371	0.294
IL-1α	0.090	0.147	-0.012	-0.034	**0.962**	0.023	0.089	-0.007	0.005	**0.951**	0.032	0.059	-0.015	-0.018	**0.965**
IL-9	0.213	0.219	-0.004	-0.047	**0.931**	0.099	0.119	-0.021	-0.020	**0.951**	0.109	0.110	-0.040	-0.021	**0.950**
IL-4	0.317	0.155	0.129	0.008	**0.861**	0.179	0.107	-0.026	0.042	**0.882**	0.188	0.082	0.059	0.037	**0.917**

Table 3. Factor loadings for 5 factors (F1 – F5) derived from cytokine levels in the control, LPS, and LPS + CBL conditions using principal component analysis (Rotation method: Varimax with Kaiser Normalization). These 5 factors explained 81.1%, 73.5%, and 74.1% data variance, respectively. Major loadings (>0.3) for each factor are bolded.
*For ease of interpretation, all factors in LPS and LPS+CBL conditions were listed following the same numbering scheme as Control factors.

Figure 1

Figure 1. Partial correlations controlling for age, sex, and body mass index (BMI) between **(A)** Control Factor 4 and Anticipatory Anhedonia; **(B)** LPS Factor 4 and Anticipatory Anhedonia; **(C)** LPS+CBL Factor 4 and Anticipatory Anhedonia; **(D)** LPS+CBL Factor 4 and Consummatory Anhedonia. All are significant at Bonferroni-corrected significant threshold $\alpha = \frac{0.05}{5 \times 2 \times 3} \approx 1.7 \times 10^{-3}$.

Abbreviations: CBL: clenbuterol; HC: healthy control participants; LPS: lipopolysaccharide; Clinical: participants with psychiatric symptoms and/or diagnoses. TEPS-A: Temporal Experience of Pleasure Scale – Anticipatory; TEPS-C: Temporal Experience of Pleasure Scale – Consummatory.

Figure 2

Figure 2. Change in cytokine correlations in the LPS+CBL condition relative to the LPS condition (LPS+CBL – LPS; lower half triangle), and their corresponding log-transformed *p* values (upper half triangle). With false discovery rate correction, significant differences were detected in the following pairs: *(a)* RANTES and PDGF-AA, *(b)* MIP-1β and IL-2, *(c)* MCP-1 and IL-6, *(d)* MCP-1 and IL-8, *(e)* MCP-1 and IP-10, *(f)* MIP-1α and MCP-1, *(g)* MIP-1β and MCP-1, *(h)* RANTES and TNF-α. *Abbreviations*: CBL: clenbuterol; LPS: lipopolysaccharide