# Powerful eQTL mapping through low coverage RNA sequencing

Tommer Schwarz [1,2], Toni Boltz [3], Kangcheng Hou [1], Merel Bot [4], Chenda Duan [5], Loes Olde Loohuis [7,8], Marco P. Boks [9], René S. Kahn [9,10], Roel A. Ophoff [1,3,11], Bogdan Pasaniuc [1,2,3,6]

1. Bioinformatics Interdepartmental Program, University of California Los Angeles, Los Angeles, CA, USA
2. Department of Computational Medicine, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, USA
3. Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, USA
4. Center for Neurobehavioral Genetics, Semel Institute for Neuroscience and Human Behavior, University of California, Los Angeles, California, USA
5. Department of Computer Science, University of California, Los Angeles, Los Angeles, CA USA
6. Department of Pathology and Laboratory Medicine, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, USA
7. Department of Psychiatry, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, USA
8. Program in Neurobehavioral Genetics, Semel Institute, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, USA
9. Department of Psychiatry, Brain Center University Medical Center Utrecht, University Utrecht, Utrecht, The Netherlands
10. Department of Psychiatry, Icahn School of Medicine, Mount Sinai, NY, the United States of America
11. Department of Psychiatry, Erasmus University Medical Center, Rotterdam, The Netherlands

Corresponding authors: Tommer Schwarz (tomschwarz@g.ucla.edu), Bogdan Pasaniuc (pasaniuc@ucla.edu)

**ABSTRACT**

Mapping genetic variants that regulate gene expression (eQTL mapping) in large-scale RNA

sequencing (RNA-seq) studies is often employed to understand functional consequences of

regulatory variants. However, the high cost of RNA-Seq limits sample size, sequencing depth,

and therefore, discovery power. In this work, we demonstrate that, given a fixed budget, eQTL

discovery power can be increased by lowering the sequencing depth per sample and increasing

the number of individuals sequenced in the assay. We perform RNA-Seq of whole blood tissue

across 1490 individuals at low-coverage (5.9 million reads/sample) and show that the effective

power is higher than that of an RNA-Seq study of 570 individuals at high-coverage (13.9 million

reads/sample). Next, we leverage synthetic datasets derived from real RNA-Seq data to explore

the interplay of coverage and number individuals in eQTL studies, and show that a 10-fold

reduction in coverage leads to only a 2.5-fold reduction in statistical power. Our study suggests

that lowering coverage while increasing the number of individuals is an effective approach to

increase discovery power in RNA-Seq studies.

**KEYWORDS**

RNA-Seq, Gene expression, Association testing

**BACKGROUND**

The vast majority of risk loci identified in genome-wide association studies (GWAS) are difficult to

interpret as they lie in noncoding regions of the genome. Variants that regulate gene expression

abundance, as measured through expression quantitative trait locus (eQTL) studies, provide

insightful information about the functional interpretation of GWAS signals [1-2]. By integrating eQTL

associations with GWAS, we can hope to identify target genes that are driving the GWAS signal

at a locus [3-6]. RNA sequencing (RNA-Seq) is the state-of-the-art assay for measuring gene

expression in bulk tissue and is therefore the assay of choice for eQTL mapping [7-8]. However, the

high cost of RNA-Seq often limits the sample size and therefore reduces the discovery power of

eQTL studies based on RNA-Seq [2,6,9].

Traditional RNA-Seq study design prioritizes sequencing depth per individual (targeted levels of

coverage in the range of 30-50 million reads) over the number of individuals (samples) included

in the study [10-12]. However, given that high levels of coverage per individual limits the sample size

of a study, this results in a loss of statistical power in eQTL mapping. Previous studies have

established that the low-coverage whole genome sequencing of a larger number of individuals

attains increased power of association compared to higher-coverage studies of smaller sample

sizes in GWAS [13-17]. This raises the hypothesis that, similarly as for whole genome sequencing

and GWAS, lower coverage RNA-seq with a considerable increase in the number of individuals

sequenced could increase power of discovery in eQTL studies [18-21]. Currently, there is no

systematic approach for determining the optimal sample size (in terms of number of sequenced

individuals) and coverage to maximize eQTL discovery power.

In this work, we perform RNA-Seq in 1490 individuals at a lower coverage (average mapped read

depth of 5.9 million reads/sample) and find that eQTL discovery power is better than that of an

experiment with a similar budget, but with fewer individuals and higher coverage. Compared to

high-coverage RNA-Seq, we find a high degree of consistency in both the gene expression as

well as eQTL effects. We assess the interplay of coverage per sample and accuracy of expression

estimates using synthetic RNA-Seq datasets generated by the down-sampling of real high-

coverage data. Our analyses show that a sequencing experiment conducted with a target

coverage of 10 million reads/sample has an average correlation per-gene of 0.40, when compared

to an experiment conducted with a target coverage of 50 million reads/sample. We provide

evidence to show that under a fixed budget, sequencing at lower coverage levels (< 10 million

reads/sample) and increased sample size can boost the effective sample size per unit of cost

compared to standard approaches of eQTL study design.

**RESULTS**

***Low-coverage RNA-sequencing is accurate for eQTL mapping***

To validate the utility of low-coverage RNA-sequencing, we sequenced whole blood tissue from

N = 1490 unrelated individuals (**Methods**) (**Supplementary Figure 1A** and **Supplementary**

**Figure 1B**). We target a sequencing coverage of 9.5 million reads per sample, yielding M = 5.9

million reads mapped to RefSeq genes on average (sd across samples of 1.96 million,

**Supplementary Figure 2**). We refer to this dataset as the lower-coverage RNA-Seq, or the M=5.9

million reads/sample dataset. We contrast this dataset with an RNA-Seq dataset obtained with a

similar budget, but with 2.4-fold higher coverage (M = 13.9 reads) across N = 570 individuals

(**Supplementary Figure 1C** and **Supplementary Figure 1D**) [22]. We refer to this as the higher-

coverage whole blood RNA-Seq, or the M = 13.9 million reads/sample dataset (Table 1).

First, we assess the number of genes quantified in the two datasets. We observe 40459 genes

with at least one mapped read on average across samples in the whole blood high-coverage

dataset, and 27308 genes with at least one mapped read on average across samples in the whole

blood low-coverage dataset. Notably, when restricting to protein coding genes with at least one

mapped read in both the high-coverage and low-coverage datasets, we find more similar numbers

between the data sets, with 18329 and 15605 genes quantified, respectively. This is likely due to

the very sparse abundance of the non-protein coding genes, making them less likely to be

detected in a lower coverage dataset. Indeed, we observe similar effects across the high vs low

coverage datasets when assessing the genes with sufficient expression to be included in eQTL

analysis (TPM > 0.1 in 20% of individuals, see **Methods**): 26566 genes (15496 protein coding

genes) in high coverage data versus 19039 (13339 protein coding genes) in low coverage data.

Most importantly we observe a high correlation in the abundance levels across the two data sets

thus demonstrating that high and low coverage recover similar expression ($R^2$= 0.91, **Figure 1A**).

Next, we investigate the power of low-coverage RNA-Seq for eQTL mapping. We conducted cis-

eQTL mapping with a 1 Mb window using FastQTL, restricting to the 1490 unrelated individuals

in the low-coverage RNA-Seq data (**Methods**), to identify 7587 genes (eGenes) with a significant

association at FDR correction level of 5%. As expected, eQTL distribution is concentrated at

transcription start sites (TSS), with 73% of eGenes TSS within 250kb of the associated SNP

(eSNP). Repeating this approach using the high-coverage whole blood data in 570 individuals,

we only find 5971 genes with a significant association at FDR correction level of 5%. 4969 of the

7587 eGenes found using the low-coverage data are also significant in the high-coverage data.

Of these, 2151 of the eGenes are protein coding eGenes that share the same associated eSNP,

and we see an extremely high level of concordance between effect sizes for these eGenes across

the two datasets ($R^2$ = 0.93, **Figure 1B**). This further indicates that low-coverage RNA-Seq is

robust in capturing eQTL effect sizes. 1002 genes were found to be eGenes in the high-coverage

eQTL analysis but not in the low-coverage analysis, with 573 (of the 1002) not passing expression

levels (TPM >0.1 in 20% individuals) to be included in the low-coverage eQTL analysis; only 234

of the 573 were protein coding genes. Similar concordance is observed at the level of p-values

for the associations in both datasets (**Figure 1C**). Comparing the p-values for eGenes detected

in both eQTL analyses, the corresponding regression line has a slope of 0.39, consistent with

lower-coverage dataset having superior statistical power to detect associations over lower-

coverage dataset, and consistent with overall number of significant eQTL discoveries. We report

the results from using typed SNPs in these eQTL analyses (**Methods**), but observe similar

patterns when using the full set of imputed SNPs.

To further validate the performance of eQTL analysis using lower coverage RNA-Seq (coverage

5.9M, n = 1490), we compared the resulting eQTLs to the ones found by GTEx consortium in

whole blood [12] (**Supplementary Figure 3**). Restricting to the 12247 protein coding genes with

sufficient expression to be included in both studies (> 0.1 TPM in 20% of samples) we find that

4529 out of the 5957 protein coding genes (76%) with a significant association using the lower-

coverage data also had a significant association in GTEx, correcting at an FDR level of 5%.

Further restricting to eGenes with the same leading SNP in both of these datasets (140 genes)

(**Figure 1D**), we observe a correlation ($R^2$) of 0.85 between their effect sizes. Looking into the

1428 protein coding genes with a significant association in eQTL analysis using the lower-

coverage RNA-Seq but not in GTEx using an FDR cutoff of 5%, we observe that 372 have

significant association in GTEx using an FDR cutoff of 10%. To further ensure that these eGenes

are not false positives, we compare the set of 1428 genes with eQTL analysis conducted by the

eQTLGen Consortium [23] and find that all but 190 of these genes have been found to have a

significant association in eQTLGen. This suggests that the additional associations found using

lower-coverage data that are not found in GTEx are not false positives, but fall just below the

significance threshold in the GTEx analysis.

Finally, we explore the impact of RNA-Seq at lower coverages for cell type expression

estimation. We use Cibersort [24] to compare cell-type proportion estimates between the lower-

coverage data and higher-coverage data (**Methods**). We find that the median estimated cell

type proportions are conserved across both datasets, suggesting that deconvolution of cell type

specific signal from gene expression profiles of whole blood samples is not impacted when

coverage is reduced by half (**Supplementary Figure 7**).

***Impact of RNA-seq coverage on eQTL power***

Having demonstrated the accuracy of low-coverage RNA-Seq in eQTL mapping in real data, we

next focused on exploring the interplay of number of individuals and coverage for optimizing

power for discovery. As simulating RNA-Seq data is challenging [25-26], we down-sample reads from

higher-coverage RNA-Seq data to create synthetic datasets at various coverages (**Methods**). We

observe that with just a fraction of the reads, it is still possible to estimate gene expression (**Figure**

**2A**). For example, using just 10% of the data (5.0 million reads/sample) retains a per gene $R^2$ of

0.40, on average. The results from our analyses using these synthetic lower-coverage RNA-Seq

datasets suggest that under simplified settings of no per-sample library preparation cost, the

statistical power in an association study can be increased up to fourfold by spending more

resources on increasing sample size and fewer resources on increasing coverage. In practice,

increasing the number of samples in an RNA-Seq study leads to increased library preparation

costs, making the increase in obtainable statistical association power less obvious.

It has been established that statistical power in association studies is a function of sample size,

phenotype measurement accuracy, and genotype measurement accuracy [13,19,29]. This means that

the power of a study with sample size N and estimated gene expression is approximately the

same as the power of a study with sample size N, using the true gene expression measurements

(**Methods**). In this scenario, $R^2$ is the correlation between the true expression and the expression

estimates. We therefore report the squared correlation ($R^2$) between synthetic datasets at various

coverages and the full data at an average of 50 million reads/sample (which is assumed to be the

true gene expression). While these results show the mean $R^2$ for all genes obtained under one

synthetic dataset (one draw) per coverage level, we find that the synthetic datasets are consistent

across multiple draws at the same coverage level (**Supplementary Figure 4A**) and each show

similar correlations with the ground truth gene expression (**Supplementary Figure 4B**).

Next, we quantified how well lower-coverage RNA-Seq can be used to detect eGenes [30]. We

explore the number of genes with significant associations after FDR correction at 5% under

various levels of simulated coverage (**Figure 2B**). Using synthetic data, as the number of reads

per sample decreases, we find that many eGenes are still detectable. For example, at 10 million

reads per sample, just 20% of the full coverage, 60% of the eGenes are still detected. In the

context of eQTL studies, synthetic RNA-Seq supports the idea that sequencing at lower

coverages over a higher number of individuals is a promising approach to boosting statistical

power.

Finally, we explore the estimation accuracy in the synthetic data as a function of relative gene

expression abundance, since less abundant genes may not be captured altogether at lower

sequencing coverages. We stratify genes into five groups based on their relative expression in

the full dataset (M=50.3 million reads/sample) and report the $R^2$ for genes in each of these groups

in synthetic data (**Figure 2C**). We observe that in the synthetic RNA-Seq dataset at 10 million

reads/sample, we capture expression of highly expressed genes better than lower expressed

genes. Specifically, for genes in the lowest through the highest quintiles of relative gene

abundance, we find the average correlation ($R^2$) to the ground truth of expression to be 0.36, 0.44,

0.61, 0.73, 0.86, respectively. We observe the same effect for synthetic datasets at coverages of

1 million reads/sample and 25 million reads/sample (**Supplementary Figure 5A** and

**Supplementary Figure 5B**). These results suggest that the ability to achieve similar power in

eQTL analysis studies will differ per gene, and is a function of relative expression. We further

investigate the properties of genes with quantification accuracy influenced by coverage levels of

sequencing and find that that protein coding genes are more accurately quantified at lower

coverage levels (**Supplementary Figure 6A**). Conversely, the number of transcripts per gene,

gene length, and GC content do not appear to be factors that broadly influence the gene

quantification accuracy when sequencing coverage is reduced (**Supplementary Figure 6B**,

**Supplementary Figure 6C,** and **Supplementary Figure 6D**).

***Optimal association power for eQTLs is attained at lower coverage with a larger number***

***of samples***

In the context of reducing experimental costs, we explored the trade-off between the number of

samples sequenced and the average coverage per sample. We evaluated the expected effective

sample size obtained with lower coverage per sample and compared this with a conventional

approach of 50 million reads/sample. We down-sample reads (as described in Section 1 and

**Methods**) from a high-coverage RNA-Seq experiment derived from Fibroblast tissue in order to

create lower-coverage RNA-Seq synthetic data. This is done to match actual low coverage

sequencing as closely as possible. To evaluate the relationship between cost, coverage, and

sample size, we use the following equation to model the budget: $B = n * a + \frac{n*b*c}{d}$

Where $B$ is the cost/budget (in US dollars), $a$ is the library preparation cost per sample, $b$ is the

target coverage of each sample (in millions of reads), $c$ is the cost per lane (which contains $d$

million reads), and $d$ is the number of reads per sequencing lane (in millions). We compute the

effective sample size of an eQTL study as a function of average coverage, which determines the

number of samples sequenced under a fixed budget (**Figure 3A**). As an example, at a fixed

budget of $300,000, the highest effective sample size is achieved by sequencing 2045 individuals

using 10 million reads per sample, which leads to a corresponding effective sample size of 1107.

An experiment achieving the sample effective sample size, using 50 million reads per sample,

would cost $426,564 (N = 1107, $R^2$ = 1.0). Therefore, by lowering the coverage of each sample

and increasing sample size, we achieve the same effective sample size at just 70.3% of the cost.

In practice, it is common to observe a considerable discrepancy between the target number of

reads in an experiment and the number of reads that successfully map to genes. This can be

attributed to different library prep techniques, quality of samples, or tissue type. To show how

mapping rate can influence the effective sample size of an experiment, we model effective sample

size with varying levels of mapping rates (**Methods**). As expected, we observe that as the

mapping rate increases, there is a corresponding increase in effective sample size (**Figure 3C**).

We provide a webtool as a practical approach for selecting cost-effective designs for maximizing

eQTL power: https://tomschwarz.shinyapps.io/RNASeqCoverageCalculator/.

With a budget of ~$300k and an expected mapping rate of 0.60 (chosen based on mapping rate

of similar experiments using TruSeq Stranded plus rRNA and GlobinZero in whole blood tissue),

we see the maximum effective sample size would be achieved at a target coverage of 7 million

reads per sample, including 2227 individuals in the study. We estimate that achieving the same

effective sample size using data with 50 million reads per sample would cost ~$500k (N = 1328),

or 1.78x the cost of sequencing 2227 individuals at a coverage of 7 million reads/sample. To

explore other cost scenarios we provide a webtool we created a webtool where one can enter

budget, costs, and other details about the experiment, in order to see how to achieve optimal

effective sample size (https://tomschwarz.shinyapps.io/RNASeqCoverageCalculator/).

**DISCUSSION**

In this work, we generate RNA-Seq data at a lower coverage than typically used in eQTL studies

(5.9M reads/sample) and demonstrate how this approach boosts effective sample size per unit

cost in an association study. To further validate this approach, we use synthetic RNA-Seq data to

show that the optimal level of coverage in an RNA-Seq project for the purpose of identifying eQTL

associations is lower than is commonly practiced [10-12]. Based on our findings, we recommend

increasing sample size while lowering sequencing depth per sample in order to achieve optimal

statistical power in association studies.

We conclude with some notes, caveats, and future directions. First, synthetic RNA-Seq via

downsampling reads is potentially limited in several ways. These synthetic datasets of lower

coverage RNA-Seq are created by uniformly sampling from real RNA-Seq data with an average

of 50 million reads mapped per sample. However, in practice, it is possible that sequencing biases

are not captured by uniform sampling due to the different experimental setup compared to the

dataset from which we sample [27]. Additionally, these synthetic datasets are based on data

obtained from fibroblast tissue with different transcriptomic profiles from whole blood, potentially

influencing the sequencing depth required to detect associations with gene expression. Finally,

this approach is optimized for eQTL discovery. Other mechanisms that are detected using RNA-

Seq, such as RNA splicing, have different mechanisms and will likely have different optimal

coverages for detection. The fact that we identify different sets of eGenes depending on which

gene expression measurements we consider (GTEX vs eQTLGen vs lower-coverage RNA-Seq),

shows that we need to increase cohort sizes in order to fully understand the connection between

genetics and gene expression in blood. Furthermore, the results in **Figure 3A** (figure showing

effective sample size at various coverages) indicate that even including 1490 individuals under

this fixed budget is not enough to achieve the optimal effective sample size. Current approaches

are not sufficient to understand the full landscape of eQTLs in whole blood tissue, even while only

considering a single genetic ancestry group. We compare the eGenes identified by GTEx,

eQTLGen, and the lower-coverage RNA-Seq (**Supplementary Figure 8**) and find that no single

study is sufficient in capturing all of the associations in whole blood. As observed in GWAS, much

larger sample sizes including far more ancestral diversity in these samples will enable discovery

of novel associations in transcriptomics. Including non-European populations and considering the

temporal aspect of gene expression will help us gain a more complete understanding of the blood

transcriptome landscape in the entire population.

**CONCLUSIONS**

In summary, we show that reducing coverage and increasing the number of samples in an eQTL

study is a valid approach for increasing effective sample size of the association study. We use

both real and synthetic RNA-Seq data to confirm the benefit of increased sample sizes in eQTL

studies. This approach can be applied to any dataset for which genotypes are available and will

help scientists optimize resources when measuring gene expression for the purpose of integration

with genetics. We provide an online tool to assist with improved design of eQTL studies at

https://tomschwarz.shinyapps.io/RNASeqCoverageCalculator/.

**METHODS**

**Cohort Description**

The samples included are from a study with individuals ascertained for bipolar disorder (BP). The

cohort consists of 916 individuals with BP, 358 controls, and 216 relatives of the individuals with

BP.

**Connection between effect size and $R^2$**

If $g$ is the genotype at the SNP that we are testing for associations, and is the effect size of that

SNP when regressing on the true gene expression, y, and is the effect size of that SNP when

regressing on the estimated gene expression, $\tilde{y}$. The relationship between $y$ and $\hat{y}$ is as follows

that $r^2 = corr(y, \hat{y})$. It follows that the estimates of effect size for a SNP on the true gene

expression, $\hat{\beta}$, are related to the estimate of effect size for a SNP on the estimated gene

expression, $\tilde{\hat{\beta}}$ as $\tilde{\hat{\beta}} = cov(g, \tilde{y}) = cov(g, ry + \varepsilon) = cov(g, ry) + cov(g, \varepsilon) = r\hat{\beta}$ where $\varepsilon$ is a

random variable with mean 0 and variance 1. The association test statistics at low-coverage is

$x_{ground} = N cor^2(g, y)$ thus implying that the association statistic at low coverage

is $x_{low-coverage} = N cor^2(g, \tilde{y}) = N\tilde{\hat{\beta}}^2 = N(r\hat{\beta})^2 = r^2 * N cor^2(g, y) = r^2 x_{ground}$

**Budget model**

We modeled the cost of a large-scale bulk RNA-Seq experiment based on parameters from two

different library prep techniques: (1) TruSeq Stranded plus rRNA and GlobinZero and (2) TruSeq

Stranded polyA selected, both from the UCLA Neuroscience Genomics core. Cost, or *B,* is a

function of the following: *a,* the library preparation cost per sample, *b,* which is the target coverage

of each sample (in millions of reads). *c,* the cost per lane (which contains *d* million reads), and *d*

is the number of reads per sequencing lane (in millions). Altogether, we model the budget as $B =$

$n * a + \frac{n*b*c}{d}$

**Genotyping pipeline**

Genotypes for the low-coverage whole blood samples were obtained from the following platforms:

OmniExpressExome (N = 810), PSYCH (N = 523), and COEX (N = 163). Given that the SNP-

genotype data for both the fibroblast and whole blood samples came from numerous studies using

various genotyping platforms (including GSA, Illumina550, OmniExpress Exome, COEX, and

PsychChip) the number of overlapping SNPs across all platforms was < 80k, prompting us to

perform imputation separately for each genotyping platform. Genotypes were first filtered for

Hardy-Weinberg equilibrium p value < 1.0e-6 for controls and p value < 1.0e-10 for cases, with

minor allele frequency (MAF) > 0.01, leaving 148613 typed SNPs.

Genotypes were imputed using the 1000 Genomes Project phase 3 reference panel [33] by

chromosome using RICOPILI v.1 [34] separately per genotyping platform, then subsequently

merged. Imputation quality was assessed by filtering variants where genotype probability > 0.8

and INFO score > 0.1, resulting in 2289732 autosomal SNPs. We restricted to only autosomal

due to sex chromosome dosage, as commonly done [12].

**Synthetic low coverage RNA-Seq**

We use high-coverage RNA-Seq (average of 50 million reads/sample, TruSeq Stranded plus

rRNA and GlobinZero library prep) from a set of 152 cell lines derived from human fibroblast cells.

We assume this to be the ground truth of gene expression. We used seqtk

(https://github.com/lh3/seqtk) to randomly sample reads at various coverages, uniformly. We

performed five iterations of downsampling at each level of coverage in order to account for

potential variability in the sampling and sequencing errors.

**RNA-Seq processing pipeline**

We used FASTQC to visually inspect the read quality from the lower-coverage whole blood RNA-

Seq (5.9M reads/sample) and the higher-coverage fibroblast RNA-Seq (13.9M reads/sample).

We then used kallisto to pseudoalign reads to the GRCh37 transcriptome and quantify estimates

for transcript expression. We aggregated transcript counts using scripts from the GTEX

consortium (https://github.com/broadinstitute/gtex-pipeline) [12].

**344**  **cis-eQTL mapping**

**345**  Excluding related individuals (pi_hat > 0.2) from the analysis, we perform cis-eQTL analysis

**346**  mapping using FastQTL [30], using a defined window of 1 Mb both up and downstream of every

**347**  gene's TSS, for sufficiently expressed genes.  We run the eQTL analysis in permutation pass

**348**  mode (1000 permutations, and perform multiple testing corrections using the q-value FDR

**349**  procedure, correcting at 5% unless otherwise specified. We then restrict our associations to the

**350**  top (or leading) SNP per eGene.

**351**  **Cell type proportion estimation**

**352**  We estimate the proportion of cell types of both the lower-coverage and higher-coverage bulk

**353**  whole blood RNA-seq datasets using CIBERSORTx [35] with batch correction applied and LM22

**354**  signature matrix as the reference gene expression profile. The LM22 signature matrix uses 547

**355**  genes to distinguish between 22 human hematopoietic cell phenotypes.

**356**  $R^2$ **adjustment**

**357**  To account for the variability in mapping rate across different library prep techniques [37] and

**358**  different tissue types, we look at the mean $R^2$ at the expected coverage, which is calculated as

**359**  *expected coverage = target coverage * estimated mapping rate.* Using mean $R^2$ values from

**360**  comparing lower-coverage synthetic RNA-Seq to higher-coverage RNA-Seq real data, we fit a

**361**  log curve to estimate the adjusted mean $R^2$ ($R^2_{adj}$) at the expected coverage.

**362**  **Effective Sample Size**

**363**  Under a fixed-budget setting, we calculate effective sample size ($N_{eff}$) for a given coverage using

**364**  the adjusted mean $R^2$ ($R^2_{adj}$) and the number of samples included at a given coverage level (N)

**365**  $N_{eff} = R^2_{adj} * N$

**366**  **DECLARATIONS**

**367**  **Ethics approval and consent to participate**

The authors assert that all procedures contributing to this work comply with the ethical standards

of the relevant national and institutional committees of human experimentation and with the

Helsinki Declaration of 1975, as revised in 2008.

**Consent for publication**

Not applicable.

**Availability of data and materials**

Gene expression data will be made available upon publication

**Competing interests**

All authors declare they have no competing interest

**Funding**

T.S. was supported by the National Institute of Neurological Disorders and Stroke of the National

Institutes of Health under Award Number T32NS048004. T.B. was supported by the NIH (grant

number 5T32HG002536-19). This research was supported by the National Institute of Mental

Health of the National Institutes of Health under Award number 5R01MH115676-04. The content

is solely the responsibility of the authors and does not necessarily represent the official views of

the National Institutes of Health.

**Author Contributions**

T.S., B.P., and R.O. initialized the study. B.P. and R.O. directed and supervised the project. R.O.,

R.K., and M.P.B. collected samples. M.B. prepared samples for sequencing. Bioinformatics

analysis was conducted by T.S., T.B., K.H., C.D., and L.O.L. . The first draft of the manuscript

was drafted by T.S. and all authors contributed to editing, revisions, and approval.

**Acknowledgements**

We thank the study subjects for their willingness to provide specimens and clinical data. We thank Yi Ding, Kathryn Burch, Ruthie Johnson, Arjun Bhattacharya, and Malika Freund for meaningful discussion in helping make this work possible.

**SUPPLEMENTAL INFORMATION**

**Supplementary Figure 1:** *Distribution of ancestry among samples.* **(S1A)** MDS plot of 2000 samples in 5.9M read/sample cohort. **(S1B)** Distribution of ancestry among sample in 5.9M read/sample cohort. **(S1C)** MDS plot of 759 samples in 13.9M read/sample cohort: Genotype PC1 and PC2 are projected onto PCs from 1000 Genomes Project. **(S1D)** Distribution of ancestry among sample in 13.9M read/sample cohort.

**Supplementary Figure 2**: *Number of pseudoaligned reads per sample.* **(S2A)** Number of pseudoaligned reads per sample in low-coverage RNA-Seq. **(S2B)** Number of pseudoaligned reads per sample in high-coverage RNA-Seq.

**Supplementary Figure 3:** *Real data p-value comparison scatterplot with GTEX*

**Supplementary Figure 4:** *Variability in correlations in synthetic data.* **(S4A)** Scatterplot of log TPM of different draws in synthetic data. **(S4B)** Distribution of correlations observed between synthetic lower-coverage RNA-Seq and ground truth.

**Supplementary Figure 5:** *Using synthetic data, how well do we capture expression as a function of average expression in a given gene*. **(S5A)** Correlation as a function of relative expression, at 25 million reads/sample. (**S5B)** Correlation as a function of relative expression, at 1 million reads/sample.

**Supplementary Figure 6:** *Using synthetic data, how well do we capture expression in different gene categories*. (**S6A)** Using synthetic data, how well do we capture expression as a function of whether a gene is protein coding or not. **(S6B)** Using synthetic data, how well do we capture expression as a function of number of isoforms in a given gene. **(S6C)** Using synthetic RNA-Seq, how well do we capture expression as a function of gene length in a given gene Gene expression estimation accuracy simulated at 10 million reads/sample as a function of relative gene length. **(S6D)** Using synthetic RNA-Seq, how well do we capture expression as a function of GC content.

**Supplementary Figure 7:** *Estimation of cell-type proportions*.

**Supplementary Figure 8:** *Overlap of significant eGenes using RNA-Seq from three different datasets*

**REFERENCES**

1.    Montgomery, S. B. *et al.* Transcriptome genetics using second generation sequencing in
a Caucasian population. *Nature* vol. 464 773–777 (2010).

2.    Pickrell, J. K. *et al.* Understanding mechanisms underlying human gene expression
variation with RNA sequencing. *Nature* **464**, 768–772 (2010).

3.    Mancuso, N. *et al.* Integrating Gene Expression with Summary Association Statistics to
Identify Genes Associated with 30 Complex Traits. *Am. J. Hum. Genet.* **100**, 473–487 (2017).

4.    Gusev, A. *et al.* Integrative approaches for large-scale transcriptome-wide association
studies. *Nat. Genet.* **48**, 245–252 (2016).

5.    Gamazon, E. R. *et al.* A gene-based association method for mapping traits using
reference transcriptome data. *Nat. Genet.* **47**, 1091–1098 (2015).

6.    Zhernakova, D. V. *et al.* Identification of context-dependent expression quantitative trait
loci in whole blood. *Nat. Genet.* **49**, 139–145 (2017).

7.    Wang, Z., Gerstein, M. & Snyder, M. RNA-Seq: a revolutionary tool for transcriptomics.
*Nat. Rev. Genet.* **10**, 57–63 (2009).

8.    Rockman, M. V. & Kruglyak, L. Genetics of global gene expression. *Nat. Rev. Genet.* **7**,
862–872 (2006).

9.    Gallagher, M. D. & Chen-Plotkin, A. S. The Post-GWAS Era: From Association to
Function. *Am. J. Hum. Genet.* **102**, 717–730 (2018).

10.    Hoffman, G. E. *et al.* CommonMind Consortium provides transcriptomic and epigenomic
data for Schizophrenia and Bipolar Disorder. *Sci Data* **6**, 180 (2019).

11.    Franzén, O. *et al.* Cardiometabolic risk loci share downstream cis- and trans-gene
regulation across tissues and diseases. *Science* **353**, 827–830 (2016).

12.    GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across
human tissues. *Science* **369**, 1318–1330 (2020).

13.    Pasaniuc, B. *et al.* Extremely low-coverage sequencing and imputation increases power
for genome-wide association studies. *Nat. Genet.* **44**, 631–635 (2012).

14.    Sims, D., Sudbery, I., Ilott, N. E., Heger, A. & Ponting, C. P. Sequencing depth and
coverage: key considerations in genomic analyses. *Nat. Rev. Genet.* **15**, 121–132 (2014).

15.  CONVERGE consortium. Sparse whole-genome sequencing identifies two loci for major
depressive disorder. *Nature* **523**, 588–591 (2015).

16.  Homburger, J. R. *et al.* Low coverage whole genome sequencing enables accurate
assessment of common variants and calculation of genome-wide polygenic scores. *Genome
Med.* **11**, 74 (2019).

17.  Rubinacci, S., Ribeiro, D. M., Hofmeister, R. J. & Delaneau, O. Efficient phasing and
imputation of low-coverage sequencing data using large reference panels. *Nat. Genet.* **53**, 412
(2021).
18.  Baccarella, A., Williams, C. R., Parrish, J. Z. & Kim, C. C. Empirical assessment of the
impact of sample number and read depth on RNA-Seq analysis workflow performance. *BMC*
*Bioinformatics* **19**, 423 (2018).
19.  Pritchard, J. K. & Przeworski, M. Linkage disequilibrium in humans: models and data.
*Am. J. Hum. Genet.* **69**, 1–14 (2001).
20.  Tarazona, S., García-Alcalde, F., Dopazo, J., Ferrer, A. & Conesa, A. Differential
expression in RNA-seq: a matter of depth. *Genome Res.* **21**, 2213–2223 (2011).
21.  Robinson, D. G. & Storey, J. D. subSeq: determining appropriate sequencing depth
through efficient read subsampling. *Bioinformatics* **30**, 3424–3426 (2014).
22.  Krebs, C. E. *et al.* Whole blood transcriptome analysis in bipolar disorder reveals strong
lithium effect. *Psychol. Med.* **50**, 2575–2586 (2020).
23.  Võsa U, Claringbould A, Westra H-J, Bonder MJ, Deelen P, Zeng B, et al. Unraveling the
polygenic architecture of complex traits using blood eQTL meta-analysis. *bioRxiv*. 447367
(2018)
24.  Chen, B., Khodadoust, M. S., Liu, C. L., Newman, A. M. & Alizadeh, A. A. Profiling
Tumor Infiltrating Immune Cells with CIBERSORT. *Methods Mol. Biol.* **1711**, 243–259 (2018).
25.  Williams, A. G., Thomas, S., Wyman, S. K. & Holloway, A. K. RNA-seq Data: Challenges
in and Recommendations for Experimental Design and Analysis. *Curr. Protoc. Hum. Genet.* **83**,
11.13.1–20 (2014).
26.  Frazee, A. C., Jaffe, A. E., Langmead, B. & Leek, J. T. Polyester: simulating RNA-seq
datasets with differential transcript expression. *Bioinformatics* **31**, 2778–2784 (2015).
27.  Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).
28.  Liu, Y. *et al.* Evaluating the impact of sequencing depth on transcriptome profiling in
human adipose. *PLoS One* **8**, e66883 (2013).
29.  Mandric, I. *et al.* Optimized design of single-cell RNA sequencing experiments for cell-
type-specific eQTL analysis. *Nat. Commun.* **11**, 5504 (2020).
30.  Ongen, H., Buil, A., Brown, A. A., Dermitzakis, E. T. & Delaneau, O. Fast and efficient
QTL mapper for thousands of molecular phenotypes. *Bioinformatics* **32**, 1479–1485 (2016).

31.    Marioni, J. C., Mason, C. E., Mane, S. M., Stephens, M. & Gilad, Y. RNA-seq: an

assessment of technical reproducibility and comparison with gene expression arrays. *Genome*

*Res.* **18**, 1509–1517 (2008).

32.    Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-

Based Linkage Analyses. *The American Journal of Human Genetics* vol. 81 559–575 (2007).

33.  Consortium, T. 1000 G. P. & The 1000 Genomes Project Consortium. A global reference for

human genetic variation. *Nature* vol. 526 68–74 (2015).

34.    Lam, M. *et al*. RICOPILI: Rapid Imputation for COnsortias PIpeLIne. *Bioinformatics* **36**,

930-933 (2019).

35.    Newman, A.M. *et al*. Determining cell type abundance and expression from bulk tissues

with digital cytometry. *Nature Biotechnology* **37**, 773-782 (2019).

36. Hoffman, G. E. et al. CommonMind Consortium provides transcriptomic and epigenomic data

for Schizophrenia and Bipolar Disorder. Sci. Data 6, 180 (2019).

37.  Aguet, F. et al. Genetic effects on gene expression across human tissues. Nature 550, 204–

213 (2017)

519
520 **DISPLAY ITEMS**
521
522

| Referred to as: | Coverage (million reads per sample) | Tissue | Number of samples | Library prep method |
| --- | --- | --- | --- | --- |
| Lower-Coverage or M=5.9M reads/sample (Whole Blood) | 5.9 | Whole blood | 1490 | TruSeq Stranded plus rRNA and GlobinZero |
| Higher-Coverage or M=13.9M reads/sample (Whole Blood) [19] | 13.9 | Whole blood | 570 | Meta-analysis of (1) TruSeq Stranded plus rRNA and GlobinZero and (2) TruSeq Stranded polyA selected |
| High- | 50.3 | Fibroblast | 155 | TruSeq Stranded polyA selected |

| | | | | |
|---|---|---|---|---|
| coverage (Fibroblast) | | | | |
| GTEX[12] | 82 | Whole blood | 670 | TruSeq Non-stranded polyA selected |
| eQTLGen[13] | N/A | Whole blood | 31684 | Meta-analysis consisting of RNA-Seq and microarray |

523
524 **Table 1:** *RNA-Seq datasets discussed in this paper*
525 The coverage refers to the average number of reads that successfully map to the transcriptome,
526 except for GTEX, which refers to the median number of total reads per sample (average
527 mapped not available). Further description of sample overlaps among cohorts in
528 **Supplementary Note**.
529
530
531

| | Cost per lane | Cost per sample |
|---|---|---|
| Scenario 1 | $1790 | $87 |
| Scenario 2 | $1790 | $30 |
| Scenario 3 | $1790 | $150 |
| Scenario 4 | $1000 | $150 |

532
533 **Table 2:** *Sequencing cost scenarios (Figure 3)*
534 The cost parameters corresponding to the effective sample size scenarios in Figure 3. Cost per
535 sample reflects the cost of library prep to include an additional sample. Cost per lane reflects the
536 cost per sequencing lane, which allows for 300 million reads.
537
538
539

**Figure 1:** *Concordance of eQTL discovery when using lower-coverage RNA-Seq vs higher-coverage RNA-Seq* **(1A):** Restricting to the 20735 genes with sufficient expression levels to be included in eQTL analysis in both the 5.9M read/sample and 13.9M read/sample dataset, comparison of the median expression (log TPM) across samples, of every gene. $R^2$ = 0.91.
**(1B):** In real data, scatterplot of effect sizes of most significant eQTL hits for the 2151 protein coding genes with the same eQTL hit in both eQTL analyses performed (low-coverage and high-coverage). On the x-axis, we show the effect sizes for these genes using low-coverage RNA-Seq, on the y-axis we show the effect sizes for these genes using high-coverage RNA-Seq. **(1C):** Real data p-value comparison scatterplot: In real data, scatterplot of -log p-values of most significant eQTL hit for 13950 genes included in both eQTL analyses performed (low-coverage and high-coverage). On the x-axis, we show the -log p-values for these genes using low-coverage RNA-Seq, on the y-axis we show the -log p-values for these genes using high-coverage RNA-Seq. The dotted line shows $y = x$, while the solid line shows the line of best fit for the 3985 protein-coding eGenes with a significant eQTL hit in both datasets. **(1D):** In real data, scatterplot of effect sizes of the most significant eQTL hit for the 140 eGenes with the same leading SNP identified in both eQTL analyses performed (lower-coverage RNA-Seq with 5.9M reads/sample and GTEX). On the x-axis, we show the effect size for these eGenes from eQTL analysis conducted using the 1490 individuals of EUR ancestry and imputed genotypes, and on the y-axis we show the effect sizes for these eGenes from eQTL analysis published by the GTEX Consortium.

564
565
**Figure 2:** *Synthetic lower-coverage RNA-Seq captures expression signal* **(2A):** On the x-axis, we
show the level of simulated coverage, and on the y-axis we show the mean Pearson correlation
of every gene. We calculate this value by finding the $R^2$ values for the TPM values of each of
45,910 genes across 155 samples between the high coverage data (average of 50 million reads
per sample) and the simulated data, and reporting the mean $R^2$ value per gene. **(2B):** For a fixed
number of individuals, absolute number and percentage of eGenes captured at 5% FDR, for
synthetic RNA-Seq at varying levels of coverage. **(2C):** Gene expression accuracy as a function
of relative gene expression observed in actual RNA-Seq data with 50 million reads/sample.
23,540 genes (with average expression < 0.1 TPM) are divided into five ascending quintiles of
expression based on their average expression in 155 samples.
576
577

578
**Figure 3:** *Effective sample size under various budget parameters*. **(3A):** Effective sample size
in RNA-Seq under a fixed budget ($300,000) as a function of the number of samples and the
resulting coverage. Cost assumptions: $87 per library prep per sample, $1790 per lane of
sequencing (300 million reads). **(3B):** Effective sample size in RNA-Seq under a fixed budget
($300,000) as a function of the number of samples and the resulting coverage. Cost assumptions
vary and are reflected in Table 2. **(3C):** Effective sample size under a fixed budget ($300,000) as
a function of the number of samples and the results coverage. A global mapping rate parameter
is used to simulate actual experimental conditions (Methods).