A warmer growing season triggers earlier following spring phenology

Hongshuang Gu[1†], Yuxin Qiao[1†], Zhenxiang Xi[1†], Sergio Rossi[2,3], Nicholas G. Smith[4], Jianquan

Liu[1*], Lei Chen[1, 4*]

[1]Key Laboratory of Bio-Resource and Eco-Environment of Ministry of Education, College of

Life Sciences, Sichuan University, Chengdu, China

[2]Département des Sciences Fondamentales, Université du Québec à Chicoutimi, Chicoutimi

(QC), G7H SB1, Canada

[3]Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems,

Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical Garden,

Chinese Academy of Sciences, Guangzhou, China

[4]Department of Biological Sciences, Texas Tech University, Lubbock, USA

[†]These authors contribute equally to this work.

[*]Corresponding author: liujq@lzu.edu.cn, chen_lei1029@163.com

Authors' ORCID iDs:

HG, 0000-0003-0609-7862; YQ, 0000-0002-7944-6165; ZX, 0000-0002-2851-5474; NGS,

0000-0001-7048-4387; JL, 0000-0002-4237-7418; LC, 0000-0001-7011-8782

Abstract: Under global warming, advances in spring phenology due to the rising temperature
have been widely reported. However, the mechanisms underlying the warming-induced earlier
spring phenology remain poorly understood. Here, using multiple long-term and large-scale
phenological datasets between 1951 and 2018, we show that warmer temperatures during the
previous growing season between May and September led to earlier spring phenology in the
Northern Hemisphere. We also found that warming-induced increases in maximum
photosynthetic rate in the previous year advanced spring phenology, with an average of 2.50
days $°C^{-1}$. Furthermore, we found a significant decline in the advancing effect of warming
during previous growing season on spring phenology from cold to warm periods over the past
decades. Our results suggest that observed warming-induced earlier spring phenology may be
driven by increased photosynthetic carbon assimilation in the previous season, while the
slowdown in the advanced spring phenology is likely due to decreased carbon assimilation
when warming exceeding the optimal temperatures for photosynthesis. Our study suggested the
vital role of photosynthetic carbon assimilation during growing season in spring phenology
under global warming.

Introduction

Plant phenology influences the fitness of individual plants and the functioning of terrestrial ecosystems, including the fluxes of water and energy and food webs[1-6]. Since phenological events are highly sensitive to climate variations, monitoring changes in plant phenology can provide the first clear visible signals of the impact of climate change on terrestrial ecosystems[6,7]. Under global warming, advanced spring phenology due to rising temperature has been widely reported[8-12]. However, important questions regarding the mechanisms underlying this response remain unanswered[13-17]. This largely hinders the prediction of spring phenology and global carbon cycling under future warming conditions.

Generally, spring phenology is considered to be driven by temperatures in winter and spring because plants need to accumulate sufficient winter chilling to end endodormancy and spring forcing units to break ecodormancy before the budbreak [18-22]. Recent studies show that the response of earlier spring phenology to climate warming is declining[17]. However, there continues to be debate about the slowdown in the warming-induced spring phenology. In fact, plants need to assimilate and store sufficient carbohydrates in the preceding growing season to resist to the frost temperatures in winter and support growth reactivation in spring[23-26]. In temperate regions, nonstructural carbohydrates (NSC; soluble sugar and starch) often reach the maximum levels in autumn before winter dormancy, but become depleted by early summer after spring growth[27-29]. Girdling experiments have demonstrated that a later budbreak is often associated with a lower NSC availability[30,31]. The timing of spring phenology is therefore likely to depend on the photosynthetic carbon assimilation during the previous growing season.

Under global warming, increasing temperatures may influence the photosynthetic carbon assimilation and delaying spring phenology in the following year [32]. Photosynthetic carbon uptake tends to show a peaked response to temperature at the leaf and canopy scale[12,33-36]. As such, an increase in temperature might increase photosynthesis in cold and temperate regions, and advance spring phenology[37,38]. When temperatures increase above the optimal threshold for photosynthesis, this could explain the slowdown in warming-induced advancement in spring phenology. However, previous researches have largely overlooked the effect of previous growing season climate on spring phenology[39-42].

Using long-term phenological observations and remote-sensing chronologies collected in the Northern Hemisphere (Fig. 1), we analyzed the effect of warming during the previous growing season on spring phenology. We hypothesized that timing of spring phenology may depend on the photosynthetic carbon assimilation during the growing season prior to leaf senescence. According to this carbon-driven theory, warmer temperatures during the previous growing season are expected to increase photosynthetic carbon uptake and trigger earlier spring phenology.

Materials and Methods

PEP725 phenological network

Data were provided by the European phenology database PEP725 (http://www.pep725.eu/), which contains phenological observations of temperate species across central Europe since 1951[43]. We selected the date when the first leaf stalks are visible (BBCH 11 in PEP725) to represent the start of spring phenology (SOS) and date when 50% leaves had their autumnal color (BBCH94 in PEP725) to represent the end of autumn phenology (EOS). Data exceeding

2.5 times of median absolute deviation (MAD) were considered outliers and were removed[44].

We selected 466,988 records of nine temperate tree species (Table S1) at 2,300 sites, for a total

of 171,202 species-site combinations with at least 30-year observations.

PhenoCam network

The PhenoCam network (https://phenocam.sr.unh.edu/) is a cooperative database of digital

phenocamera imagery which provided the dates of phenological transition between 2000 and

2018 worldwide[45,46]. In the PhenoCam network, the 50%, and 90% of the Green Chromatic

Coordinate (G_{CC}) were calculated daily to extract the date of greenness rising and falling

According to the following formula:

90
$$G_{CC} = \frac{G_{DN}}{R_{DN} + G_{DN} + B_{DN}}, (1)$$

where R_{DN}, G_{DN} and B_{DN} are the average red, green and blue digital numbers (DN), respectively.

We selected 50% threshold of G_{CC}_90 (G_{CC} reaches 90^{th} quantiles of its seasonal amplitude)

as SOS[47]. We removed outliers according to the abovementioned procedure, and we selected

sites with at least 8-year observations between 2000 and 2018. We also excluded agricultural

ecosystem to avoid human influence. The final dataset had a total of 738 records at 78 sites

from three vegetation types: deciduous broadleaf forests, evergreen forests and grassland.

GIMMS NDVI$_{3g}$ phenological product

The Normalized Difference Vegetation Index (NDVI), a proxy of vegetation greenness and

photosynthetic activity, is commonly used to derive phenological metrics[48]. We derived SOS

from the third generation GIMMS NDVI$_{3g}$ dataset (http://ecocast.arc.nasa.gov) from Advanced

Very High Resolution Radiometer (AVHRR) instruments for the period 1982-2014 with a

spatial resolution of 8 km and a temporal resolution of 15 days[49].

We only kept areas outside tropics (latitudes >30 °N), which have a clear seasonal

phenology[50] and excluded bare lands with annual average NDVI < 0.1 to reduce bias. We

applied a Savizky-Golay filter[51] to smooth the time series and eliminate noise of atmospheric

interference and satellite sensor, and used a Double Logistic 1^{st} to extract phenology dates[50]

according to the formula:

109
$$y(t) = a\left(\frac{1}{1+e^{k(t-m)}} + \frac{1}{1+e^{e(t-n)}}\right) + b, (2)$$

where a, k, m, and n are parameters of logistic function and a is the initial background

NDVI value, a + b represents the maximum NDVI value, t is time in days, and y(t) is the NDVI

value at time t. The second-order derivative of the function (Eq. (2)) was calculated to extract

SOS and EOS at the first and second local maximum point, respectively[52,53].

FLUXNET dataset

The flux dataset was downloaded from FLUXNET (https://fluxnet.org/data/). The data were

released in November 2016 (total 212 sites) worldwide[54]. The dataset was processed with a

processing pipeline to reduce uncertainty by improving the data quality control. The pipeline

generates uniform and high-quality derived data products suitable for studies that compare

multiple sites[54]. We selected 39 sites with at least 5-year observations and daily records > 300

for each year between 1992 and 2014. The Singular Spectrum Analysis (SSA) filter method[55]

was used to smooth the time series of gross primary productivity (GPP) to minimize the noise.

GPP_{max}, daily maximum GPP in a year, is considered as an important index to evaluate the

carbon fixation of terrestrial ecosystems and the feedback of vegetation climate[56–58]. We

extracted the GPP_{max} from the smoothed GPP curve by the SSA-based de-nosing smoothing

method[59]. SOS and EOS were extracted from smoothed daily GPP curve based on the threshold

method[51]. The spring and autumn threshold were defined as 15% of the multi-year daily GPP

maximum following previous studies[60,61], and SOS and EOS were defined as the turning point

when the smoothed GPP was higher or lower than spring or autumn threshold, respectively.

Climate data

Gridded daily mean temperature, precipitation, solar radiation and air humidity during 1950-

2015 in Europe was downloaded from the database E-OBS (http://www.ecad.eu/)[62] at 0.25°

spatial resolution. Gridded monthly soil moisture during 1979-2015 was downloaded from

World   Meteorological   Organization   (http://climexp.knmi.nl/select.cgi?id=someone@

somewhere&field=clm_wfdei_soil01) at 0.5° spatial resolution and banded with PEP725

dataset. Global monthly mean temperature during 1981-2017 was downloaded from Climate

Research Unit (https://crudata.uea.ac.uk/cru/data/hrg/cru_ts_4.04/cruts.2004151 855.v4.04/) at

0.5° spatial resolution and matched the PhenoCam and GIMMS $NDVI_{3g}$ datasets. Bilinear

interpolation method was used to extract climate data of each site or pixel using the *raster*

package[63] in R version 4.0.3[64]. Environmental variables, including daily mean temperature (°C),

shortwave radiation (Wm^{-2}), CO_2 (ppm), and precipitation (mm) of FLUXNET dataset were

also extracted.

Statistical analysis

To tested our hypotheses, we primarily used the observations from the PEP725 network

corresponding E-OBS climate dataset. We did this because PEP725 data was relatively more

reliable than the extracted phenological metrics from imagines of PhenoCam network and

GIMMS $NDVI_{3g}$ product because its phenological records were taken manually in situ. In

addition, the PEP725 network covered a longer period (between 1951 and 2015) than

PhenoCam (between 2000 and 2018) and GIMMS $NDVI_{3g}$ dataset (between 1982 and 2014).

The PhenoCam and GIMMS $NDVI_{3g}$ phenology products were used to test the robustness and

generality of the results obtained from the PEP725 network. Specifically, we calculated the

temperature sensitivity (S_T, change in days per degree Celsius) based on mean temperatures

during previous growing season (from May to September) and timing of spring phenology using

three complementary large-scale datasets (PEP725, PhenoCam, GIMMS $NDVI_{3g}$) in the

Northern Hemisphere. To clarify the underlying mechanisms, we further examined the

relationships between GPP_{max} of previous growing season and SOS between 1992 and 2014

using FLUXNET data.

Temperature sensitivities

Temperature sensitivity (S_T, change in days per degree Celsius), defined as the slope of a linear

regression between the dates of phenological stages and the temperature[21,65,66], was used to

investigate the effects of temperature of the previous growing season on leaf unfolding dates in the PEP725 network. The length of growing season was defined as the period between SOS and EOS. The mean dates of SOS and EOS from the PEP725 network were DOY 120 and DOY 280. Therefore, the period between May and September was selected to represent the growing season. Linear regression models were used to calculate S_T of leaf unfolding for each species at each site. In the model, the response variable was the leaf unfolding date, the predictor was the mean temperature of previous growing season.

In addition, a linear mixed-effects model was used to exclude the covariate effects of other climate factors and autumn phenology, and further examine the overall effect of temperature of previous growing season on leaf unfolding dates by pooling all the records across species and study sites. In the model, the response variable was leaf unfolding dates, the predictors were temperature, radiation, precipitation, soil moisture, humidity during previous growing season between May and September and leaf senescence dates of previous year, with random intercepts among species and sites. In addition, we quantified and compared the effects of climate variables of previous growing season on leaf unfolding dates using boosted regression tree, an ensemble statistical learning method[67], which has been widely applied in ecological modeling and prediction[68,69]. Because radiation and soil moisture data were only available since 1980, we selected phenology and climate datasets between 1984 and 2015 to perform the linear mixed-effects model and boosted regression tree. Linear mixed-effects model fitting was conducted using the *lme4* package[70] of R[64]. Significance testing of the fixed effects terms was done using the Satterthwaite method incorporated into the *lmerTest* package[71] of R[64], where *P* values less than 0.05 were considered significant. We performed the boosted regression trees using the *gbm* package[72] in R[64], where 10-fold cross validation was used to determine the optimal number of iterations.

Effect of past climate change on spring phenology

Following Fu et al.[17], we assessed the effects of past climate warming on spring phenology. First, we calculated the mean temperature during growing season (May-September) across all the 2,322 sites in Europe from 1951 to 2015. Using a 15-year smoothing window, we identified the coldest and warmest periods: 1955-1969 and 2000-2014 over the past 60 years. Based on the mean temperature during previous growing season, we calculated the S_T of leaf unfolding during the two periods for each species at each site. One-way analysis of variance (ANOVA) was used to test the difference in the S_T of leaf unfolding during 1955-1969 and 2000-2014.

Structural equation modeling

We used a structural equation model (SEM) to analyze the relationships between climate, GPP_{max} and SOS from the 39 flux sites. The climate variables in the structural equation model included temperature, radiation, soil moisture, CO_2 and precipitation during previous growing season. Because the daily GPP started to increase from DOY 120, peaking at DOY 180, then decreased until DOY 300 (Fig. S2), the period between May and September was also selected as the growing season. This is also consistently with the period of growing season identified by the dates of leaf unfolding and leaf senescence in PEP725 network. The SEM was fitted using the *lavaan* package[73] in R 4.0.3[64].

All data analyses were conducted using R version 4.0.3[64].

Results

Temperature sensitivity (S_T, change in days per degree Celsius), is often used to describe the response of plant phenology to warmer temperatures. Here we calculated the S_T of spring phenology based on the mean temperatures during previous growing season between May and September and dates of spring leaf unfolding obtained from PEP725 network, and start of season (SOS) metrics extracted from Phenocam, and GIMMS NDVI$_{3g}$ images (see Methods). The calculated S_T of spring phenology based on the three datasets is shown in Fig. 2. Using the PEP725 network, the mean S_T of leaf unfolding across nine temperate tree species between 1951 and 2015 was −2.50 days·°C^{-1} (Fig. 2a). This suggested that a warmer previous growing season advanced leaf unfolding dates. The S_T were negative across all selected nine temperate tree species (Fig. 2b). The response of *Q. robur Linnaeus* to temperature of previous growing season were the strongest, with an average of −2.82 days·°C^{-1}, significantly stronger than *T. cordata* (−1.04 days·°C^{-1}) and *T. platyphyllos* (−1.16 days·°C^{-1}).

In addition to temperature, spring phenology has been reported to be influenced by other climate variables and autumn phenology. We used a linear mixed effects model to exclude these covariate effects and further examined the effects of temperature of previous growing season on spring leaf unfolding (see Methods). From our model, we consistently observed that leaf unfolding dates were advanced by increasing temperatures during previous growing season by an average of −2.67 days·°C^{-1} (Table S2). Using boosted regression tree, we found the temperature during previous growing season had the strongest effect on leaf unfolding dates (84.67%), followed by radiation (6.33%), soil moisture (3.93%), precipitation (3.32%), humidity (1.75%) (Fig. S1).

Our PEP725 results were corroborated by PhenoCam and remoting sensing data. Specifically, we observed a negative effect of temperature during previous growing season on SOS in deciduous broad-leaved forests, evergreen forests and grasslands using phenological metrics extracted from the PhenoCam network between 2000 and 2018 (Fig. 2c). According to the calculated S_T, the SOS in response to warming of the previous growing season was the strongest in deciduous broad-leaved forests, followed by evergreen forests and grasslands (Fig. 2c). Using the phenology metrics extracted from remote sensing dataset between 1982 and 2014, we also observed that increasing temperature during previous growing season advanced SOS across different vegetation types in the Norther Hemisphere (Fig. 2d). Among all the vegetation types, the S_T of the Tundra was the lowest, followed by Temperate Broadleaf & Mixed Forests and Savannas & Shrublands (Fig. 2d).

To test whether earlier spring phenology was driven by increased photosynthetic carbon assimilation, we further examined the relationship between daily maximum photosynthetic rate (GPP$_{max}$) of the previous growing season and SOS between 1992 and 2014 using FLUXNET data (see Methods). We found that the timing of SOS showed a significant negative correlation with the GPP$_{max}$ during the growing season between 1992 and 2014 (correlation coefficient = −0.36, $P<0.01$, Fig. 3a). This suggested that spring phenology tended to occur earlier with the increased photosynthetic carbon assimilation during previous growing season. To further test the carbon-driven hypothesis, we constructed a structural equation model (SEM) that included climate variables, GPP$_{max}$ and SOS (Fig. 3b). The SEM showed that spring phenology (SOS) was advanced by increased GPP$_{max}$ (slope = −2.331, $P<0.001$). In addition, the effect of temperature on GPP$_{max}$ was the strongest (slope = 0.319, $P<0.001$), followed by radiation (slope

= 1.28, $P<0.05$), while radiation (slope = 0.005, $P>0.05$), CO_2 (slope = 0.002, $P>0.05$) and

precipitation (slope = 0.001, $P>0.05$) almost had no effects on GPP_{max}. The detailed statistics

of the SEM are listed in Table S3.

To examine the potential effects of climate warming on leaf unfolding, we used the

PEP725 dataset to calculate and compare the S_T between the coldest and the warmest 15-year

periods: 1955-1969 and 2000-2014, respectively (Fig. 4). We found the S_T of leaf unfolding

decreased by 77.7% from -1.93 ± 0.08 days·$°C^{-1}$ during 1955-1969 to -0.43 ± 0.08 during

2000-2014 (Fig. 4a). Between 1955 and 1969, the S_T of early-successional species is -2.37

days·$°C^{-1}$ and -1.23 days·$°C^{-1}$ for late successional species. Between 2000 and 2014, S_T of

early-and late-successional species were -0.13 days·$°C^{-1}$ and -0.92 days·$°C^{-1}$, respectively. The

S_T of early successional species decreased more from the coldest to the warmest periods (-2.24

± 0.15 days·$°C^{-1}$) than that of late successional species (-0.31 ± 0.16 days·$°C^{-1}$) (Fig. 4b).

Discussion

Global warming advances bud break and leafing worldwide[21,74–77]. Using three long-term and

large-scale phenological datasets, we show that warmer temperatures of the previous growing

season drive earlier phenology in the following spring in the Northern Hemisphere. We also

find that warming increased photosynthetic carbon assimilation, suggesting a mechanism by

which global warming is triggering earlier spring phenology (Fig. 5).

In deciduous tree species, carbon gained through photosynthesis is often stored in the form

of non-structural carbohydrates (NSC-soluble carbohydrates and starch), which supports the

growth of buds and leaves in following spring before newly grown leaves can supply

photosynthates[78–80]. For instance, 95% of starch stored in the branches of *Fagus sylvatica* L.

and *Quercus petraea* (Matt.) Liebl was consumed when spring bud-break occurred[79]. Needle

growth of *Larix gmelinii* in spring drew nearly 50% of the carbohydrates fixed in previous

year[81,82]. Phloem girdling showed that deficient carbon storage can significantly delay the

timing of spring budbreak and reduce bud size[27].

During winter dormancy, temperate tree species also need to store sufficient carbohydrates

prior to leaf senescence for respiration to maintain baseline functions and protect cells from

frost damage and ensure survival[83,84]. Therefore, warmer temperatures in the previous growing

season may advance spring phenology by increasing carbon storage, supported by the negative

correlations between spring phenology and maximum photosynthetic rate in the previous year.

Recently, Zani et al.[32] has reported that increased carbon assimilation during the growing

season drives earlier autumn leaf senescence in temperate ecosystems. When leaf senescence

occurred earlier, trees advanced the endodormancy[5,85]. In this context, the requirement of

chilling units may be also fulfilled earlier. As a result, earlier autumn phenology facilitates an

earlier spring phenology[86]. Therefore, increased carbon assimilation may directly drive autumn

phenology, and, in turn, influence spring phenology. To test this hypothesis, we excluded the

covariate effect of autumn phenology and isolated the effect of temperature of the previous

growing season on leaf unfolding. The relationship was negative, confirming our hypothesis

and suggesting that increased carbon assimilation of previous season triggers an earlier spring

phenology.

We observed that early-successional species showed a stronger response to the warming

during growing season compared to late-successional species. In addition to temperature, spring

phenology is also under photoperiodic control[87]. Because photoperiod remains stable regardless of climate change, plants are expected to show relatively conservative climatic responses when they rely on photoperiod to determine spring phenology. However, photoperiod sensitivities often vary among species[87]. For example, late-successional species are reported to have a higher photoperiod sensitivity compared to early-successional species[87,88]. The higher photoperiod sensitivity of late-successional species may, therefore, explain their conservative climatic responses compared to early-successional species [87,89].

Recent studies have reported that the warming-induced earlier spring phenology has slowed down over the past decades[21,90,91]. Fu et al. (2015)[17] reported that S_T of leaf unfolding decreased by 40% from 4.0 ± 1.8 days·°C^{-1} during 1980-1994 to 2.3 ± 1.6 days·°C^{-1} during 1999-2013. The observed declining effect of warming on spring phenology is generally considered a result of chilling reduction in winter[92]. Here, we found the carbon-driven earlier spring phenology is also slowing down in recent decades, especially for early-successional species. Duffy et al.[12] showed that the mean temperatures in the warmest quarter passed the optimal for photosynthesis over the past decade, with a sharp declining photosynthesis. The increased heat and water stress of the last decades is the cause of a spreading growth decline of forests[93–95]. Therefore, the observed decline in the S_T may involve reductions in carbon assimilation by heat waves or drought events under global warming[96,97].

Conclusion

Despite the warming-induced spring phenology observed worldwide, the underlying causes and mechanisms of such events still remain under debate. To decipher this complex relationship, we used multiple long-term and large-scale datasets to provide evidence that spring phenology is advanced by warmer temperatures of the previous growing season. Correspondingly, we observed that leaf unfolding occurred earlier under enhanced maximum photosynthetic capability. These findings suggest that an increased carbon assimilation under global warming could be involved in the observed earlier leafing of trees. In addition, we observed a decline in the carry-over effect of growing-season warming on spring phenology likely due to the reduced photosynthetic carbon assimilation by heat and water stress under global warming. With an increase in projected drought frequency under warming scenarios[93,98], we expect that temperate trees will slow down the advancement of spring phenology. This may reduce the strength of forest carbon sinks under future climate conditions[17]. Our study provides new insights into the warming-induced change in spring phenology under global climate change to predict spring phenology and vegetation-atmosphere feedbacks under future climatic scenarios.

References

1. Gao, M. *et al.* Divergent changes in the elevational gradient of vegetation activities over the last 30 years. *Nat. Commun.* **10**, 2970 (2019).

2. Koen Hufkens *et al.* Ecological impacts of a widespread frost event following early spring leaf-out. *Glob. Chang. Biol.* **18**, 2365–2377 (2012).

3. Kari Saikkonen *et al.* Climate change-driven species' range shifts filtered by photoperiodism. *Nat. Clim. Chang.* **2**, 239–242 (2012).

4. Zeng, H., Jia, G., & Howard Epstein. Recent changes in phenology over the northern high latitudes detected from multi-satellite data. *Environ. Res. Lett.* **6**, 45508–45518 (2011).

5.   Marc Estiarte & Josep Peñuelas. Alteration of the phenology of leaf senescence and fall
in winter deciduous species by climate change: effects on nutrient proficiency. *Glob.*
Chang. Biol. **21**, 1005–1017 (2015).

6.   Richardson, A. *et al.* Climate change, phenology, and phenological control of vegetation
feedbacks to the climate system. *Agric. For. Meteorol.* **169**, 156–173 (2013).

7.   Josep Penuelas, This Rutishauser, & Iolanda Filella. Phenology Feedbacks on Climate
Change. *Science* **324**, 887–888 (2009).

8.   Fu, Y. H. *et al.* Three times greater weight of daytime than of night-time temperature on
leaf unfolding phenology in temperate trees. *New Phytol.* **212**, 590–597 (2016).

9.   Annette Menzel *et al.* European phenological response to climate change matches the
warming pattern. *Glob. Chang. Biol.* **12**, 1969–1976 (2006).

10.   Peuelas, J. & Iolanda Filella. Responses to a Warming World. *Science* **294**, 793–795
(2001).

11.   Piao, S. *et al.* Leaf onset in the northern hemisphere triggered by daytime temperature.
Nat. Commun. **6**, 6911 (2015).

12.   Duffy, K. A. *et al.* How close are we to the temperature tipping point of the terrestrial
biosphere? *Sci. Adv.* **7**, eaay1052 (2021).

13.   Keenan, T. F., Richardson, A. D. & Hufkens, K. On quantifying the apparent temperature
sensitivity of plant phenology. *New Phytol.* **225**, 1033–1040 (2020).

14.   Niu, S., Fu, Y., Gu, L. & Luo, Y. Temperature Sensitivity of Canopy Photosynthesis
Phenology in Northern Ecosystems. *Phenology: An Integrative Environmental Science*
503–519 doi:10.1007/978-94-007-6925-0_27 (2013).

15.   Meineke, E. K., Davis, C. C. & Davies, T. J. Phenological sensitivity to temperature
mediates herbivory. *Glob. Chang. Biol.* **27**, 2315–2327 (2021).

16.   Huanjiong Wang, Quansheng Ge, This Rutishauser, Yuxiao Dai, & Junhu Dai.
Parameterization of temperature sensitivity of spring phenology and its application in
explaining diverse phenological responses to temperature change. *Sci. Rep.* **5**, 8833
(2015).

17.   Fu, Y. H. *et al.* Declining global warming effects on the phenology of spring leaf unfolding.
Nature **526**, 104–107 (2015).

18.   Carol K Augspurger. Reconstructing patterns of temperature, phenology, and frost
damage over 124 years: Spring damage risk is increasing. *Ecology* **94**, 41–50 (2013).

19.   Jochner, S., Beck, I., Behrendt, H., Traidl-Hoffmann, C. & Menzel, A. Effects of extreme
spring temperatures on phenology: a case study from Munich and Ingolstadt. *Clim. Res.*
12, 101–112 (2010).

20.   Meng, F., Zhang, L. irong, Zhang, Z., Jiang, L. & Wang, Y. Enhanced spring temperature
sensitivity of carbon emission links to earlier phenology. *Sci. Total Environ.* **745**, 140999
(2020).

21.   Güsewell, R., Furrer, R., Gehrig, R. & Pietragalla, B. Changes in temperature sensitivity
of spring phenology with recent climate warming in Switzerland are related to shifts of
the preseason. *Glob. Chang. Biol.* **23**, 5189–5202 (2017).

22.   Wang, T. *et al.* The influence of local spring temperature variance on temperature
sensitivity of spring phenology. *Glob. Chang. Biol.* **20**, 1473–1480 (2014).

23. Huang, J.-G. *et al.* Intra-annual wood formation of subtropical Chinese red pine shows
better growth in dry season than wet season. *Tree Physiol.* **38**, 1225–1236 (2018).
24. Knowles, J. F., Scott, R. L., Biederman, J. A., Blanken, P. D. & Barron-Gafford, G. A.
Montane forest productivity across a semi-arid climatic gradient. *Glob. Chang. Biol.* **26**,
6945–6958 (2020).
25. Strimbeck, G. R., Trygve, D. K., Paul, G. S. & Paula, F. M. Dynamics of low-temperature
acclimation in temperate and boreal conifer foliage in a mild winter climate. *Tree Physiol.*
28, 1365–1374 (2008).
26. Smart, D. R. Exposure to elevated carbon dioxide concentration in the dark lowers the
respiration quotient of Vitis cane wood. *Tree Physiol.* **24**, 115–120 (2004).
27. Roxas, A. A., Orozco, J., Guzmán-Delgado, P., & Zwieniecki, M. A. Spring phenology is
affected by fall non-structural carbohydrates concentration and winter sugar redistribution
in three Mediterranean nut tree species. *Tree Physiol.* 1-14 (2021).
28. Palacio, S., Maestro, M., & G Montserrat-Martí. Seasonal dynamics of non-structural
carbohydrates in two species of mediterranean sub-shrubs with different leaf phenology.
Environ. Exp. Bot. **59**, 34–42 (2007).
29. Fierravanti, A., Rossi, S., Kneeshaw, D., De Grandpré, L. & Deslauriers, A. Low Non-
structural Carbon Accumulation in Spring Reduces Growth and Increases Mortality in
Conifers Defoliated by Spruce Budworm. *Front. For. Glob. Chang.* **2**, (2019).
30. Fajstavr, M., Giagli, K., Vavrcik, H., Gryc, V. & Urban, J. The effect of stem girdling on
xylem and phloem formation in Scots pine. *Silva Fennica* **51**, 1760 (2017).
31. Fajstavr, M. *et al.* The cambial response of Scots pine trees to girdling and water stress.
Iawa Journal **41**, 159–185 (2020).
32. Zani, D., Crowther, T. W., Mo, L., Renner, S. S. & Zohner, C. M. Increased growing-
season productivity drives earlier autumn leaf senescence in temperate trees. *Science* **370**,
1066–1071 (2020).
33. Lin, Y.-S., Medlyn, B. E. & Ellsworth, D. S. Temperature responses of leaf net
photosynthesis: the role of component processes. *Tree Physiol.* **32**, 219–231 (2012).
34. Smith, N. G. & Dukes, J. S. Plant respiration and photosynthesis in global-scale models:
incorporating acclimation to temperature and CO2. *Glob. Chang. Biol.* **19**, 45–63 (2013).
35. Dusenge, M. E., Duarte, A. G. & Way, D. A. Plant carbon metabolism and climate change:
elevated CO2 and temperature impacts on photosynthesis, photorespiration and
respiration. *New Phytol.* **221**, 32–49 (2019).
36. Huang, M. *et al.* Air temperature optima of vegetation productivity across global biomes.
Nat. Ecol. Evol. **3**, 772–779 (2019).
37. Smith, N. G., Lombardozzi, D., Tawfik, A., Bonan, G. & Dukes, J. S. Biophysical
consequences of photosynthetic temperature acclimation for climate. *J. Adv. Model.* **9**,
536–547 (2017).
38. Saxe, H., Cannell, M. G. R., Johnsen, Ø., Ryan, M. G. & Vourlitis, G. Tree and forest
functioning in response to global warming. *New Phytol.* **149**, 369–399 (2001).
39. Bigras, F. J. & Bertrand, A. Responses of Picea mariana to elevated CO2 concentration
during growth, cold hardening and dehardening: phenology, cold tolerance,
photosynthesis and growth. *Tree Physiol.* **26**, 875–888 (2006).

40. M.Ewa, J. & Reinhart, C. Effects of elevated atmospheric CO2 on phenology, growth and crown structure of Scots pine (Pinus sylvestris) seedlings after two years of exposure in the field. *Tree Physiol.* **19**, 289–300 (1999).

41. Penuelas & J. Phenology. Responses to a warming world. *Science* **294**, 793–795 (2001).

42. Rebecca, S.-D. *et al.* Divergent carbon cycle response of forest and grass-dominated northern temperate ecosystems to record winter warming. *Glob. Chang. Biol.* **26**, 1519–1531 (2019).

43. Templ, B. *et al.* Pan European Phenological database (PEP725): a single point of access for European data. *Int. J. Biometeorol.* **62**, 1109–1113 (2018).

44. Leys, C., Ley, C., Klein, O., Bernard, P. & Licata, L. Detecting outliers: Do not use standard deviation around the mean, use absolute deviation around the median. *J. Exp. Soc. Psychol.* **49**, 764–766 (2013).

45. Brown, M. T., Campbell, D. E., De Vilbiss, C. & Ulgiati, S. The geobiosphere emergy baseline: A synthesis. *Ecol. Modell.* **339**, 92–95 (2016).

46. Richardson, A. D. *et al.* Tracking vegetation phenology across diverse North American biomes using PhenoCam imagery. *Sci. Data.* **5**, 180028 (2018).

47. Klosterman, S. T. *et al.* Evaluating remote sensing of deciduous forest phenology at multiple spatial scales using PhenoCam imagery. *Biogeosciences* **11**, 4305–4320 (2014).

48. Zhang, Y. *et al.* Seasonal and interannual changes in vegetation activity of tropical forests in Southeast Asia. *Agric. For. Meteorol.* **224**, 1–10 (2016).

49. Pinzon, J. E. & Tucker, C. J. A Non-Stationary 1981–2012 AVHRR NDVI3g Time Series. *Remote Sens.* **6**, 6929–6960 (2014).

50. Wang, X. *et al.* No trends in spring and autumn phenology during the global warming hiatus. *Nat. Commun.* **10**, 2389 (2019).

51. Wang, X. *et al.* Validation of MODIS-GPP product at 10 flux sites in northern China. *Int. J. Remote Sens.* **34**, 587–599 (2013).

52. Julien, Y. & Sobrino, J. A. Global land surface phenology trends from GIMMS database. *Int. J. Remote Sens.* **30**, 3495–3513 (2009).

53. Zhang, X. *et al.* Monitoring vegetation phenology using MODIS. *Remote Sens. Environ.* **84**, 471–475 (2003).

54. Pastorello, G. *et al.* The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data. *Sci. Data.* **7**, 225 (2020).

55. Kalman & Dan. A singularly valuable decomposition: The SVD of a matrix. *The College Mathematics Journal* **39**, 2233–2241 (1996).

56. Huang, K. *et al.* Enhanced peak growth of global vegetation and its key mechanisms. *Nat. Ecol. Evol.* **2**, 1897–1905 (2018).

57. Tang, Y., Xu, X., Zhou, Z., Qu, Y. & Sun, Y. Estimating global maximum gross primary productivity of vegetation based on the combination of MODIS greenness and temperature data. *Ecol. Inform.* **63**, 101307 (2021).

58. Xia, J. *et al.* Joint control of terrestrial gross primary productivity by plant phenology and physiology. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 2788–2793 (2015).

59. Hu, Z. *et al.* Joint structural and physiological control on the interannual variation in productivity in a temperate grassland: A data-model comparison. *Glob. Chang. Biol.* **24**, 2965–2979 (2018).

60. Richardson, A. D. *et al.* Influence of spring and autumn phenological transitions on forest
ecosystem productivity. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **365**, 3227–3246 (2010).

61. Wu, C. *et al.* Interannual variability of net carbon exchange is related to the lag between
the end-dates of net carbon uptake and photosynthesis: Evidence from long records at two
contrasting forest stands. *Agric. For. Meteorol.* **164**, 29–38 (2012).

62. Cornes, R. C., Schrier, G. van der, Besselaar, E. J. M. van den & Jones, P. D. An Ensemble
Version of the E-OBS Temperature and Precipitation Data Sets. *J. Geophys. Res. Atmos.*
123, 9391–9409 (2018).

63. Hijmans, R. J. *raster: Geographic Data Analysis and Modeling*. (2020).

64. R Core Team. *R: A Language and Environment for Statistical Computing*. (2020).

65. Suonan, J., Classen, A. T., Sanders, N. J. & He, J.-S. Plant phenological sensitivity to
climate change on the Tibetan Plateau and relative to other areas of the world. *Ecosphere*
10, e02543 (2019).

66. Wang, C., Cao, R., Chen, J., Rao, Y. & Tang, Y. Temperature sensitivity of spring
vegetation phenology correlates to within-spring warming speed over the Northern
Hemisphere. *Ecol. Indic.* **50**, 62–68 (2015).

67. Elith, J., Leathwick, J. R. & Hastie, T. A working guide to boosted regression trees. *J*
Anim. Ecol. **77**, 802–813 (2008).

68. Davis, K. T. *et al.* Wildfires and climate change push low-elevation forests across a critical
climate threshold for tree regeneration. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 6193–6198
(2019).

69. Lemm, J. U. *et al.* Multiple stressors determine river ecological status at the European
scale: Towards an integrated understanding of river status deterioration. *Glob. Chang.*
Biol. **27**, 1962–1975 (2021).

70. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models
Using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

71. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in
Linear Mixed Effects Models. *J. Stat. Softw.* **82**, 1–26 (2017).

72. Greenwell, B., Boehmke, B., Cunningham, J. & Developers, G. B. M. *gbm: Generalized*
Boosted Regression Models. (2020).

73. Rosseel, Y. *lavaan: An R Package for Structural Equation Modeling*. 48 (2012).

74. Asch & Rebecca, G. Climate change and decadal shifts in the phenology of larval fishes
in the California Current ecosystem. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 4065–4074 (2015).

75. RaphaSaarn. False estimates of the advance of spring. *Nature* 600 (2001).

76. Wheeler, H. C., Hye, T. T., Schmidt, N. M., Svenning, J. C. & Forchhammer, M. C.
Phenological mismatch with abiotic conditions—implications for flowering in Arctic
plants. *Ecology* **96**, 775–787 (2015).

77. Yu, H., Luedeling, E. & Xu, J. Winter and spring warming result in delayed spring
phenology on the Tibetan Plateau. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 22151–22156 (2010).

78. Hartmann, H. & Trumbore, S. Understanding the roles of nonstructural carbohydrates in
forest trees – from what we can measure to what we want to know. *New Phytol.* **211**, 386–
403 (2016).

79. Tamir, K., Yann, V., & H Günter. Coordination between growth, phenology and carbon storage in three coexisting deciduous tree species in a temperate forest. *Tree Physiol.* **36**, 847–855 (2016).

80. Tixier, A., Gambetta, G. A., Godfrey, J., Orozco, J. & Zwieniecki, M. A. Non-structural Carbohydrates in Dormant Woody Perennials; The Tale of Winter Survival and Spring Arrival. *Front. For. Glob. Chang.* **2**, 18 (2019).

81. Kagawa, A. & Maximov, S. Seasonal course of translocation, storage and remobilization of 13C pulse-labeled photoassimilate in naturally growing Larix gmelinii saplings. *New Phytol.* **171**, 793–804 (2010).

82. Siegwolf *et al.* Examining the response of needle carbohydrates from Siberian larch trees to climate using compound-specific C-13 and concentration analyses. *Plant Cell Environ.* **38**, 2340–2352 (2015).

83. Erica, F., Eduardo, F., Helen, B. & Eike, L. A Conceptual Framework for Winter Dormancy in Deciduous Trees. *Agronomy* **10**, 241 (2020).

84. Majken, P., Brandt, A. U. & Lillie, A. Winter warming delays dormancy release, advances budburst, alters carbohydrate metabolism and reduces yield in a temperate shrub. *Aob Plants* **7**, plv024 (2015).

85. Malyshev, A. V. Warming Events Advance or Delay Spring Phenology by Affecting Bud Dormancy Depth in Trees. *Front. Plant Sci.* **11**, 856 (2020).

86. Liu, Q. *et al.* Modeling leaf senescence of deciduous tree species in Europe. *Glob. Chang. Biol.* **26**, (2020).

87. Basler, D. & Körner, C. Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agric. For. Meteorol.* **165**, 73–81 (2012).

88. Basler, D. & Körner, C. Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiol.* **34**, 377–388 (2014).

89. Körner, C. & Basler, D. Phenology Under Global Warming. *Science* **327**, 1461 (2010).

90. Daphné, A. *et al.* Warmer winters reduce the advance of tree spring phenology induced by warmer springs in the Alps. *Agric. For. Meteorol.* **252**, 220–230 (2018).

91. Vitasse, Signarbieux, & YSH. Global warming leads to more uniform spring phenology across elevations. *Proc. Natl. Acad. Sci. U.S.A.* **155**, (1004).

92. Clark, J. S., Salk, C., Melillo, J. & Mohan, J. Tree phenology responses to winter chilling, spring warming, at north and south range limits. *Funct. Ecol.* **28**, 1344–1355 (2014).

93. Brzostek, E. R. *et al.* Chronic water stress reduces tree growth and the carbon sink of deciduous hardwood forests. *Glob. Chang. Biol.* **20**, 2531–2539 (2014).

94. Julio Camarero, J. *et al.* Forest Growth Responses to Drought at Short- and Long-Term Scales in Spain: Squeezing the Stress Memory from Tree Rings. *Front. Ecol. Evol.* **6**, (2018).

95. Xu, P. *et al.* Impacts of Water Stress on Forest Recovery and Its Interaction with Canopy Height. *Int. J. Environ. Res. Public Health.* **15**, 1257 (2018).

96. Yashavanthakumar, K. J. *et al.* Impact of heat and drought stress on phenological development and yield in bread wheat. *Plant Physiol. Rep.* **26**, 357–367 (2021).

97. Choukri, H. *et al.* Heat and Drought Stress Impact on Phenology, Grain Yield, and Nutritional Quality of Lentil (Lens culinaris Medikus). *Front. Nutr.* **7**, (2020).

555   98.   Zhou, S., Zhang, Y., Williams, A. P. & Gentine, P. Projected increases in intensity,
556   frequency, and terrestrial carbon costs of compound drought and aridity events. *Sci. Adv.*
557   **5**, eaau5740 (2019).
558
559

562 **Fig. 1** Distributions of the phenological observation sites in this study. Blue dots represent the
563 2,300 sites selected from the PEP725 dataset across central Europe. Yellow and red dots
564 represent 78 sites in North America from the PhenoCam network and 39 FLUXNET sites,
565 respectively.

Fig. 2 Temperature sensitivities (S_T, change in days per degree Celsius) of spring phenology in response to increasing temperature during previous growing season. The calculated S_T was based on (**a**, **b**) records of spring leaf unfolding for nine temperate tree species at 2,300 sites in Europe, and phenological metrics extracted from (**c**) the PhenoCam network and (**d**) the GIMMS NDVI$_{3g}$ products for different biomes. DB, EN and GR in (**c**) represents deciduous broad-leaved forest, evergreen forest and grassland, respectively. In (**d**), the biomes included Boreal Forests/Taiga (BFT), Flooded Grasslands & Savannas (FGS), Temperate Broadleaf & Mixed Forests (TBM), Temperate Conifer Forests (TCF), Temperate Grasslands, Savannas & Shrublands (TGS), Tropical & Subtropical Coniferous Forests (TSC), Tropical &Subtropical Grasslands, Savannas & Shrublands (TSG), Tropical & Subtropical Moist Broadleaf Forest (TSM) and Tundra (TUN). The black dash lines indicate when the S_T is equal to zero.

582
583 **Fig. 3** (**a**) Relationship between spring phenology (SOS) and GPP_{max} and (**b**) the contructed
584 structural equation model using the data of 39 FLUXNET sites between 1992 and 2014. The
585 black dash line represents the fitted linear regression line ($SOS = 182.38 - 12.88 \times GPP_{max}$).　The
586 used variables in the structural equation model included climate variables (temperature,
587 radiation, soil moisture, CO_2 and precipitation), SOS and GPP_{max}.
588
589

590
Fig. 4 Temperature sensitivities (S_T, change in days per degree Celsius) of leaf unfolding in
early and late successional species during 1955-1969 and 2000-2014. The calculated S_T was
based on the temperature during previous growing season and leaf unfolding dates obtained
from the PEP725 database. The length of each box indicates the interquartile range, the
horizontal line inside each box the median, and the bottom and top of the box the first and third
quartiles respectively. The asterisks indicate a significant difference in the S_T 1955-1969 and
2000-2014 ($P<0.001$). The black dashed horizontal line indicates when the S_T is equal to zero.

Fig. 5 A schematic diagram of the earlier leaf-out in response to warming during previous
growing season. Warmer temperatures during the previous growing season drivers earlier spring
leaf-out by increasing the photosynthetic carbon assimilation.

Supporting Information

Table S1. List of the 9 temperate species selected from the PEP725 phenological network.

Number	Latin name	Abbreviated name	Successional type
1	*Aesculus hippocastanum* L.	*A. hippocastanum*	Early
2	*Betula pendula* Roth.	*B. pendula*	Early
3	*Alnus glutinosa* Medik.	*A. glutinosa*	Early
4	*Ribes grossularia*	*R. grossularia*	Early
5	*Fraxinus excelsior* L.	*F. excelsior*	Late
7	*Fagus sylvatica*	*F. sylvatica*	Late
7	*Quercus robur Linnaeus*	*Q. robur Linnaeus*	Late
8	*Tilia cordata* Mill.	*T. cordata*	Late
9	*Tilia platyphyllos* Scop.	*T. platyphyllos*	Late

608 **Table S2.** Results of linear mixed model that the effect of temperature during previous growing
609 season on spring phenology (SOS) after excluding the influence of other climatic factors
610 (radiation, precipitation, soil moisture, humidity) and autumn phenology (EOS).

Variables	Estimate	SE	*df*	*t* value	*P* value
Intercept	170.52	3.86	5.84	44.16	<0.001
Temperature	-2.67	5.59×10^{-2}	5.84×10^{4}	-47.85	<0.001
Radiation	-0.02	4.54×10^{-3}	5.48×10^{4}	-4.75	<0.001
Precipitation	0.00	2.37×10^{-3}	6.97×10^{4}	0.57	<0.001
Soil moisture	0.37	2.16×10^{-2}	1.76×10^{4}	17.14	<0.001
Humidity	-0.56	1.91×10^{-2}	7.01×10^{4}	-29.21	<0.001
EOS	0.07	2.77×10^{-3}	8.23×10^{4}	25.36	<0.001

613 **Table S3.** Statistics of the structural equation models (SEMs). To display model performance,
614 we calculated the Comparative Fit Index (CFI) and the root-mean square error (RMSE).

Statistics of the structural equation models (SEMs)						
Left-hand side	**Option**	**Right-hand side**	**Estimate**	**SE**	**Z value**	**_P_ value**
GPP_{max}	~	Radiation	0.005	0.004	1.179	>0.05
GPP_{max}	~	Soil moisture	0.167	0.026	6.347	<0.001
GPP_{max}	~	Temperature	0.319	0.033	9.569	<0.001
GPP_{max}	~	CO_2	0.002	0.007	0.325	>0.05
GPP_{max}	~	Precipitation	0.001	0.152	0.004	>0.05
SOS	~	GPP_{max}	-2.331	0.551	-4.229	<0.001
CFI = 0.89; RMSEA = 0.20						

Fig. S1 The relative influence of climatic factors during previous growing season on spring leaf
unfolding between 1984 and 2015 obtained from the PEP725 database.

617
618
619

Fig. S2 Daily gross primary productivity (GPP) changes in three vegetation types based on FLUXNET. The curve is obtained by average daily GPP approximately 400 site-year records of each vegetation using Generalized Additive Models (GAM) method.