Mathematical Modeling Reveals Quantitative Properties

of KEAP1-NRF2 Signaling

Shengnan Liu[1,2], Jingbo Pi[1], and Qiang Zhang[2]

[1] Program of Environmental Toxicology, School of Public Health, China Medical University, Shenyang, 110122, China

[2] Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, GA, 30322, USA

Corresponding author:

Qiang Zhang, M.D., Ph.D.

Gangarosa Department of Environment Health

Rollins School of Public Health

Emory University

1518 Clifton Road NE

Mailstop 1518-002-2BB

Atlanta, Georgia 30322

E-mail: qiang.zhang@emory.edu

ABSTRACT

In response to oxidative and electrophilic stresses, cells launch an NRF2-mediated transcriptional antioxidant program. The activation of NRF2 depends on a redox sensor, KEAP1, which acts as an E3-ligase adaptor to promote the ubiquitination and degradation of NRF2. While a great deal has been learned about this redox duo, its quantitative signaling properties are still largely unexplored. In the present study, we examined these properties including response time, half-life, maximal activation, and response steepness (ultrasensitivity) of NRF2, through mathematical modeling. The models describe, with increasing complexity, the reversible binding of KEAP1 dimer and NRF2 via the ETGE and DLG motifs, NRF2 production, KEAP1-dependent and independent NRF2 degradation, and perturbations by different classes of NRF2 activators. Our simulations revealed that at the basal condition, NRF2 molecules are largely sequestered by KEAP1, with the KEAP1-NRF2 complex distributed comparably in either an ETGE-bound only (open) state or an ETGE and DLG dual-bound (closed) state, corresponding to the unlatched and latched configurations of the conceptual hinge-latch model. With two-step ETGE binding, the open and closed states operate in cycle mode at the basal condition and transition to equilibrium mode at stressed conditions. Class I-V, electrophilic NRF2 activators, which modify redox-sensing cysteine residues of KEAP1, shift the balance to a closed state that is unable to degrade NRF2 effectively. When total NRF2 accumulates to a level that nearly saturates existing KEAP1, ultrasensitive NRF2 activation, i.e., a steep rise in the free NRF2 level, can occur. The ultrasensitivity results from two simultaneous mechanisms, zero-order degradation mediated by DLG binding and protein sequestration (molecular titration) mediated by ETGE binding. These response characteristics of class I-V activators do not require disruption of DLG binding to unlatch the KEAP1-NRF2 complex. In comparison, class VI NRF2 activators, which directly compete with NRF2 for KEAP1 binding, cause a shift to the unlatched, open state of KEAP1-NRF2 complex and ultimately its complete dissociation (unhinged), resulting in a fast release of free NRF2 followed by stabilization. Although class VI activators may

induce free NRF2 to higher levels, ultrasensitivity is lost due to lower free KEAP1 and thus its

NRF2-sequestering effect. Stress-induced nuclear NRF2 accumulation is enhanced when basal

nuclear NRF2 turnover constitutes a small load of NRF2 production. Our simulation further

demonstrated that optimal abundances of cytosolic and nuclear KEAP1 exist to maximize

ultrasensitivity. In summary, by simulating the dual role of KEAP1 in repressing NRF2, i.e.,

sequestration and promoting degradation, our mathematical modeling provides key novel

quantitative insights into the signaling properties of the KEAP1-NRF2 system, which may help

with the design of novel classes of NRF2 activators and inhibitors and understanding of the toxic

actions of environmental oxidative stressors.

Keywords: Oxidative Stress, KEAP1, NRF2, Ultrasensitivity, Protein Sequestration, Zero-order

degradation

INTRODUCTION

Under oxidative stress, the antioxidant capacity of cells is upregulated to meet the increasing demand for reactive species removal to maintain cellular redox homeostasis and limit cellular damage (Nguyen et al. 2003). Similar to many other cytoprotective responses, this adaptive antioxidant response is underpinned by a complex molecular circuitry of primarily negative feedback and incoherent feedforward nature, involving both posttranslational and transcriptional regulations (Zhang et al. 2010, Zhang et al. 2015). In mammalian cells, the main circuit mediating the transcriptional part of the antioxidant response is the KEAP1-NRF2-ARE pathway (Kobayashi et al. 2009). KEAP1 (Kelch ECH associating protein 1) is the molecular thiol-based sensor of ROS and other reactive species, which detects the redox status inside the cell and relays it to NRF2 (nuclear factor erythroid 2-related factor 2) (Dinkova-Kostova et al. 2002, Suzuki and Yamamoto 2017). As the master transcription factor, NRF2 partners with small Maf (sMaf) proteins to recognize promoter consensus sequences containing AREs (antioxidant response element) and induce a suite of target genes participating in antioxidant and detoxification reactions (Katsuoka et al. 2005, Kobayashi and Yamamoto 2005, Malhotra et al. 2010, Bellezza et al. 2018, Tonelli et al. 2018).

As the essential components for the transcriptional induction of antioxidant genes, the KEAP1 and NRF2 proteins and their interactions have been learned in great details in the past two decades (Itoh et al. 2010, Yamamoto et al. 2018, Paunkov et al. 2019, Baird and Yamamoto 2020). Tethered to the perinuclear actin cytoskeleton in the cytosol, KEAP1 functions as a homodimer (McMahon et al. 2006, Watai et al. 2007, Ogura et al. 2010). The KEAP1 peptide is composed of 624 amino acid residues forming five functional domains: NTR (N-terminal region), BTB (Broad complex, Tramtrack, and Bric-a-Brac), IVR (intervening region), DGR (double glycine repeat) or Kelch-repeat, and CTR (C-terminal region) (Canning et al. 2015, Dayalan Naidu and Dinkova-Kostova 2020). The BTB domain at the N-terminal is responsible for the

formation of KEAP1 homodimer (Zipper and Mulcahy 2002). The neighboring Kelch and CTR

domains (collectively termed as DC region) are responsible for the interaction of KEAP1 with

NRF2 (Li et al. 2004, Lo et al. 2006). As a redox sensor, KEAP1 contains 27 cysteine residues

distributed across the five domains, many of which can be modified or conjugated on the thiol

group by oxidants and electrophiles (Dinkova-Kostova et al. 2002, Yamamoto et al. 2008, Sekhar

et al. 2010).

The NRF2 protein is composed of 589 amino acids forming six functional domains, Neh1

through Neh6 (Moi et al. 1994, Tonelli et al. 2018). The Neh2 domain on the N-terminal is

responsible for the binding with the KEAP1 dimer (Itoh et al. 1999). Within Neh2, there exist two

conserved motifs in the N-to-C direction: DLG and ETGE, with an intervening sequence

containing 7 lysine residues that can be ubiquitinated (Tong et al. 2006, Tong et al. 2007). Both

motifs are involved in mediating the association between NRF2 and KEAP1 dimer. The ETGE

motif can bind to the DC region of one of the monomeric subunits of KEAP1 dimer, and the DLG

motif of the same NRF2 molecule binds to the DC region of the other subunit (McMahon et al.

2006). Therefore, the KEAP1-NRF2 complex exists at an internal molar ratio of 2:1 (Tong et al.

2006, Horie et al. 2021). The binding affinities between ETGE and KEAP1 and between DLG and

KEAP1 are substantially different, with ETGE nearly 100-fold higher than DLG (Lo et al. 2006,

Tong et al. 2006, Chen et al. 2011, Ichimura et al. 2013, Fukutomi et al. 2014). It is therefore

expected that the binding between KEAP1 and NRF2 occurs primarily in two sequential events:

an initial ETGE-mediated association forming an "open" KEAP1-NRF2 complex, and a

subsequent DLG-mediated intra-complex association forming a "closed" KEAP1-NRF2 complex

(Tong et al. 2006).

By interacting with CUL3 (Cullin 3) via its BTB and IVR domains, KEAP1 is an adaptor of

the KEAP1-CUL3-RBX1 E3 ubiquitin ligase complex (Kobayashi et al. 2004). When KEAP1 is

associated with NRF2 in the closed state, KEAP1 is able to enable the transfer of ubiquitin

molecules from the E2-ubiquitin conjugating enzyme bound to RBX1 (RING-box protein 1) to the

7 lysine residues in the intervening region between the DLG and ETGE motifs of NRF2 (Katoh et

al. 2005, Tong et al. 2006, Tong et al. 2007). Once ubiquitinated, NRF2 is rapidly degraded by

the proteasomal pathway (Kobayashi et al. 2006). Therefore, at basal conditions, NRF2 in the

cytosol has a very short half-life, mostly ranging between 6-20 min (Kwak et al. 2002, Alam et al.

2003, Itoh et al. 2003, Stewart et al. 2003, Kobayashi et al. 2004, He et al. 2006, Khalil et al.

2015, Crinelli et al. 2021). Under oxidative stress, certain sensor cysteine residues on KEAP1

are modified, which disables KEAP1's capability of mediating NRF2 ubiquitination (Yamamoto et

al. 2008, Sekhar et al. 2010, Suzuki and Yamamoto 2017). As a result, NRF2 is stabilized and

accumulates via *de novo* synthesis in the cytosol. Rising NRF2 then translocates into the nuclei

where it induces antioxidant and detoxification genes (Kobayashi and Yamamoto 2005, Itoh et al.

2010, Tonelli et al. 2018).

Despite the molecular details of KEAP1 and NRF2 interactions have been revealed to a

great extent, the quantitative signaling properties of the duo, culminating in NRF2 accumulation

and nuclear translocation, are still poorly understood. It has been demonstrated that the binding

between KEAP1 and NRF2 is not altered by oxidative stress, such that NRF2 does not dissociate

from KEAP1 (Eggler et al. 2005, He et al. 2006, Kobayashi et al. 2006). Since the discovery of

the two-site sequential binding scheme for KEAP1-NRF2 interaction, i.e., first through ETGE and

then through DLG, a hinge-latch model has been proposed (Tong et al. 2006, Tong et al. 2007,

Fukutomi et al. 2014). The model considers that the ETGE-mediated association (the hinge)

between KEAP1 and NRF2 is always engaged regardless of the presence of oxidative stressors.

However, oxidative stressors may disrupt the weaker DLG-mediated association (the latch),

rendering the closed KEAP1-NRF2 complex to revert to the open configuration (McMahon et al.

2006, Ogura et al. 2010). In the open state, KEAP1 can no longer mediate the ubiquitination of

NRF2, resulting in NRF2 stabilization. However, the validity of the hinge-latch model for KEAP1

cysteine-modifying, electrophilic oxidants (i.e., class I-V NRF2 inducers) becomes questionable

as emerging evidence suggests that these classes of compounds do not disrupt DLG binding

(Horie et al. 2021). Studies using Förster resonance energy transfer (FRET) revealed that the

association between KEAP1 and NRF2 may become even stronger when cells are exposed to

KEAP1 cysteine-modifying compounds (Baird et al. 2013).

The ETGE-mediated binding affinity between KEAP1 and NRF2 is high relative to their

cellular abundances, with the dissociation constant (K_d) ranging between 5-26 nM, as

summarized in Table S1 footnote (Lo et al. 2006, Tong et al. 2006, Chen et al. 2011, Ichimura et

al. 2013, Fukutomi et al. 2014), and the intracellular concentrations of KEAP1 dimer and NRF2 in

the order of hundreds of nM as observed in a variety of cell types (Iso et al. 2016). This suggests

that when KEAP1 is in excess relative to NRF2, as often the case at the basal condition, NRF2

molecules are largely sequestered by KEAP1, leaving free NRF2 only a very small fraction of its

total abundance. Such binding kinetics suggests that under oxidative stress, newly synthesized

NRF2 molecules will be still first sequestered by the remaining free KEAP1 reserve, and only

when it is nearly all filled by NRF2, will NRF2 becomes more available for nuclear translocation.

Therefore, the degree of NRF2 activation is in part regulated by the KEAP1 reserve capacity of

NRF2 sequestration. This mode of NRF2 activation is recently suggested in the floodgate

hypothesis (Iso et al. 2016, Suzuki and Yamamoto 2017, Yamamoto et al. 2018). If total NRF2

never accumulates to a level that can saturate existing KEAP1 molecules, nuclear NRF2

translocation and gene induction will remain muted. However, if total NRF2 can rise to a higher

level that nearly saturates KEAP1, from a quantitative signaling prospective, KEAP1-dependent

NRF2 degradation will operate near zero order and simultaneously NRF2 begins to escape

KEAP1 sequestration, both of which are robust ultrasensitive mechanisms that can produce a

steep rise in free NRF2 levels (Buchler and Louis 2008, Zhang et al. 2013, Ferrell and Ha 2014).

This amplified, nonlinear NRF2 activation can in turn induce antioxidant genes strongly. Therefore, the kinetic parameters governing the interactions between KEAP1 dimer and NRF2 seem to be critical to the quantitative behaviors of KEAP1-NRF2-ARE-mediated redox signal transduction.

From the perspective of effectively restoring redox homeostasis, the induction of these antioxidant genes needs to be launched timely and to levels that are sufficient to counteract the oxidative impacts exerted by the stressors (Zhang et al. 2010). Strong antioxidant induction would require signal amplification, i.e., ultrasensitivity, by which a small percentage change in the redox status can be transduced to induce a larger percentage change in the expression of antioxidant genes (Zhang et al. 2013, Ferrell and Ha 2014). a number of ultrasensitive mechanisms, including multistep signaling, homomultimerization, and autoregulation, have been revealed in the KEAP1-NRF2-ARE mediated transcriptional pathway (Zhang and Andersen 2007, Zhang et al. 2010). They operate collectively to ensure that the cellular antioxidant capacity can be adequately induced to levels matching the intensity of the oxidant insult.

Mathematical modeling plays a crucial role in understanding and predicting the quantitative behavior of redox pathways (Adimora et al. 2010, Selvaggio et al. 2018). Earlier modeling work including our own has included the KEAP1-NRF2 module in the larger context of the NRF2-mediated antioxidant response pathways (Zhang and Andersen 2007, Zhang et al. 2009, Hamon et al. 2014, Leclerc et al. 2014, Khalil et al. 2015, Xue et al. 2015, Kolodkin et al. 2020). However, in most of these studies the KEAP1-NRF2 module was treated as simplified degradation network motifs, yet the details of KEAP1-NRF2 interactions and especially the likely nonlinearity in signaling have not been explicitly and fully explored. In the present study, we developed a suite of mathematical models of detailed KEAP1-NRF2 interactions to explore the quantitative properties of NRF2 activation. With these models we examined the roles of open

192   and closed states of the KEAP1-NRF2 complex for the hinge-latch and floodgate hypotheses.

193   Our simulation predicts that ultrasensitive NRF2 activation may occur via zero-order protein

194   degradation and protein sequestration by KEAP1 under certain circumstances. Our

195   mathematical models provide key quantitative insights into the signaling properties of the

196   KEAP1-NRF2 module of the adaptive, cellular antioxidant response pathway.

 METHODS

Model structure

In keeping with the principle of parsimony and exploring the importance of molecular details, we

started with a minimal model capturing the basic interactions between KEAP1 and NRF2, and we

then progressively built more complexity into the model based on more recent quantitative

knowledge about the interactions. As a result of this evolution, a total of 6 models were explored

with increasing complexities, as summarized in Table 1. For all models, the following

assumptions were made.

(i) KEAP1 is treated as a single molecule of homodimer with two binding sites for NRF2 as the

dimer structure is required for NRF2 binding (Zipper and Mulcahy 2002).

(ii) Total KEAP1 abundance is a constant which is not altered by oxidative stress as extensively

demonstrated in experimental studies (Iso et al. 2016) and KEAP1 turnover (synthesis and

degradation) is not considered.

(iii) Since the binding affinity between KEAP1 and the ETGE motif of NRF2 is much higher than

the binding affinity between KEAP1 and the DLG motif of NRF2 (> 100-fold), as summarized

in Table S1 (Lo et al. 2006, Tong et al. 2006, Chen et al. 2011, Ichimura et al. 2013, Fukutomi

et al. 2014), for simplicity and following the concept of hinge-latch hypothesis (Yamamoto et

al. 2018), the initial interaction between KEAP1 and NRF2 is assumed to always start with

the binding between KEAP1 and ETGE while the binding between KEAP1 and DLG occurs

subsequently, as an intramolecular event.

(iv) Oxidation or conjugation of one monomeric subunit of the KEAP1 dimer by a class I-V NRF2

activator is sufficient to cause KEAP1 to lose its ability to mediate NRF2 degradation. The

oxidation or conjugation can occur to either free KEAP1 dimer or KEAP1 complexed with

NRF2 equally.

(v) For the Models (4a and 4b) with nuclear NRF2 translocation, cytosolic KEAP1 and nuclear

KEAP1 are kept as separate pools.

Model 1 is the most basic model, which captures the known essence of interactions between KEAP1 and NRF2 in the cytosol as shown in Fig. 1. In model 1, NRF2 is synthesized at a constant rate of k_0. Free NRF2 ($NRF2_{free}$) is degraded with a first-order rate constant of k_5, reflecting KEAP1-independent degradation such as the one mediated by the Neh6 domain involving the GSK-3, β-TrCP and Cul1 system (McMahon et al. 2004, Rada et al. 2011, Chowdhry et al. 2013, Hayes et al. 2015). $NRF2_{free}$ first binds to one of the monomeric subunits of the KEAP1 dimer through the ETGE domain with a second-order association rate constant k_1 and a first-order dissociation rate constant k_2, forming an intermediate complex $KEAP1\_NRF2_{open}$ (termed open state here). Since KEAP1 in the open state of the complex cannot execute its E3 ligase adaptor function (Katoh et al. 2005, Tong et al. 2006, Tong et al. 2007), NRF2 in $KEAP1\_NRF2_{open}$ is assumed to be degraded with a first-order rate constant of k_9 that is equal to k_5. As NRF2 is degraded, KEAP1 is recycled joining the free KEAP1 dimer pool. The NRF2 molecule in $KEAP1\_NRF2_{open}$ then further associates with the other unoccupied monomeric subunit of KEAP1 dimer through the DLG motif with a first-order association rate constant k_3 and a first-order dissociation rate constant k_4, forming the final complex $KEAP1\_NRF2_{closed}$ (termed closed state here). NRF2 in $KEAP1\_NRF2_{closed}$ is degraded with a first-order rate constant of k_6 which is much higher than k_5 and k_9, reflecting KEAP1-mediated ubiquitination and accelerated degradation of NRF2, and KEAP1 dimer is recycled. Class I-V oxidants and electrophiles can oxidize or conjugate KEAP1 (Yamamoto et al. 2008, Sekhar et al. 2010, Suzuki and Yamamoto 2017). In the model the oxidant converts $KEAP1$ to an oxidized form, $KEAP1_o$, with a second-order rate constant k_7. The same oxidation reaction is assumed to occur on the KEAP1 molecule in $KEAP1\_NRF2_{open}$ and $KEAP1\_NRF2_{closed}$ as well, forming $KEAP1_o\_NRF2_{open}$ and $KEAP1_o\_NRF2_{closed}$ respectively. $KEAP1_o$, $KEAP1_o\_NRF2_{open}$, and $KEAP1_o\_NRF2_{closed}$ can be reduced back to the respective original states with a first-order rate

constant k_8. Since there is no evidence that the association of KEAP1 with NRF2 alters the

kinetics of oxidation or conjugation of KEAP1 by oxidants, the same values of k_7 and k_8 are used

across all three oxidation/reduction reaction pairs. Since the alteration of NRF2 stability only

occurs in the closed state, NRF2 in $KEAP1_o\_NRF2_{open}$ is degraded with a first-order rate

constant of k'_9 that is equal to k_9. NRF2 in $KEAP1_o\_NRF2_{closed}$ is degraded with a first-order rate

constant of k'_6 that is much lower than k_6, reflecting the well-established fact that

oxidant-modified KEAP1 in the closed state loses its capability to mediate the ubiquitination and

degradation of NRF2 (Katoh et al. 2005, Tong et al. 2006, Tong et al. 2007). In both the k'_9 and k'_6

steps, $KEAP1_o$ is recycled joining the free $KEAP1_o$ pool. The binding between $NRF2_{free}$ and

$KEAP1_o$ through the ETGE domain is described by the second-order association rate constant

k'_1 and first-order dissociation rate constant k'_2, which are kept the same as k_1 and k_2 respectively

since class I-V oxidants do not alter the binding affinity between KEAP1 and NRF2 (Eggler et al.

2005, He et al. 2006, Kobayashi et al. 2006). The association and dissociation rate constants k'_3

and k'_4 for the intramolecular DLG binding between $KEAP1_o\_NRF2_{open}$ and $KEAP1_o\_NRF2_{closed}$

are also kept the same as k_3 and k_4 respectively, however, their values are varied to explore the

behavior of the hinge-latch hypothesis.

The detailed structure of Models 2, 3a, 3b, 4a, and 4b are presented in Figs. 1, 2A, 6A,

7A, and 8A, respectively. Briefly, in Model 2, the DLG-mediated internal binding kinetics (k_3 and

k_4) between $KEAP1\_NRF2_{open}$ and $KEAP1\_NRF2_{closed}$ is modified from Model 1 to simulate the

situation that the transitioning between the two states occurs in a cycle mode rather than an

equilibrium mode, as observed experimentally (Baird et al. 2013). In Models 3a and 3b, the

ETGE-mediated binding between KEAP1 and NRF2 is modified from the one-step mode as in

Models 1 and 2 to a two-step mode to simulate the situation that ETGE-mediated binding

involves an initial fast binding event followed by a subsequent slow binding event observed

experimentally (Fukutomi et al. 2014). This modification allows us to achieve the cycle mode of

operation without altering the DLG-mediated binding kinetics dramatically as done in Model 2.

Models 3a and 3b consider class I-V and VI NRF2 inducers as separate cases respectively.

Lastly, in Models 4a and 4b, translocation of NRF2 to the nucleus and its interaction with KEAP1

in the nucleus are considered, and the two models consider class I-V and VI NRF2 activators as

separate cases respectively.

Model parameters and ordinary differentiation equations (ODEs)

The values of most of the model parameters, including binding rate constants, degradation rate

constants, and abundances (concentrations) of KEAP1 and NRF2, were obtained or derived

from the literature. For those unknown parameter values, they were estimated based on other

constraints of the modeled system. References and details of the determination and calculation

of all parameter values are presented in Table S1 and its footnote. The unit of concentration of

the state variables is nM and time is second (S). The ODEs are presented in Tables S2-S6 and

algebraic equations calculating the concentrations of state variables in various combinations are

presented in Table S7. The steady-state concentrations of state variables at the basal and

maximally induced conditions are in Tables S8 and S9 respectively, and the steady-state

turnover fluxes of reactions at the basal and maximally induced conditions are in Tables S10 and

S11 respectively.

Modeling tools

The models were constructed and simulated in Berkeley Madonna (version 8.3.18, University of

California, Berkeley, CA) using the "Rosenbrock (stiff)" ODE solver. All model codes in Berkeley

Madonna format as well as in *R* format are available as additional Supplemental files.

Metrics of ultrasensitivity

In the present study, all oxidant-NRF2 dose-response (DR) curves were obtained once the

simulation has achieved steady state. The degree of ultrasensitivity of a steady-state DR curve

can be evaluated with two related metrics. First, the Hill coefficient, n_H, is approximated from the

equation

304
$$n_H = \frac{\ln 81}{\frac{\ln X_{0.9}}{\ln X_{0.1}}},$$
(1)

where $X_{0.9}$ and $X_{0.1}$ are the concentrations of an oxidant that produce 90% and 10% respectively

of the maximal NRF2 response (after subtracting the basal NRF2 levels) (Zhang et al. 2013). n_H

represents the overall steepness or global degree of ultrasensitivity of the DR curve. Second, we

evaluate the local response coefficient (*LRC*) of a DR curve by calculating all slopes of the curve

on dual-log scales, which are equivalent to the ratios of the fractional change in response (*R*) to

the fractional change in dose (*D*) (Goldbeter and Koshland 1982):

311
$$LRC = \frac{\mathrm{d}\ln R}{\mathrm{d}\ln D}.$$
(2)

The maximal |*LRC*| of a DR curve, LRC_{max}, represents the maximal amplification capacity of

KEAP1-NRF2-mediated signaling. Typical ultrasensitive responses have LRC_{max} values

substantially above 1. The comparison between n_H and *LRC* is important as these quantities are

not necessarily equivalent and depend on the basal response level and the shape of the DR

curve; thus, n_H alone can misrepresent the actual degree of signal amplification (Legewie et al.

2005, Zhang et al. 2013, Altszyler et al. 2017).

318
319

                                    Results

    Models 1 and 2

    Following the principle of parsimony, we started with the minimal model structure for Models 1

    and 2, which is depicted in Fig. 1 and detailed in Methods. In-depth analyses of the basic

    properties and response behaviors of the two models are provided in the Supplemental Material.

    Simulations of these two models provided the initial findings that the KEAP1-NRF2 interaction is

    capable of producing ultrasensitive activation of free NRF2 by class I-V compounds (Figs. S1C,

    S1D, and S3F), and the ultrasensitivity results from saturation of KEAP1 by NRF2 where both

    zero-order degradation and protein sequestration of NRF2 occur (Fig. S3G-S3H).

        In Model 1, however, the abundance ratio of the open and closed KEAP1-NRF2

    complexes (*KEAP1_NRF2*$_{open}$:*KEAP1_NRF2*$_{closed}$) remains constant at 1:1 at all times in all

    conditions (Figs. S1A-S1C and S2), indicating these two states are always at equilibrium to each

    other. Using FRET to track the open and closed states of KEAP1-NRF2 complex, Baird et al.

    observed that the two states diverge and do not follow an equilibrium mode of operation in a

    variety of chemically perturbed conditions (Baird et al. 2013). But rather, a "cyclic sequential

    attachment and regeneration" (abbreviated as "cycle") mode of operation was suggested. In this

    mode, because of the rapid degradation of NRF2 in the closed KEAP1-NRF2 complex, KEAP1 is

    quickly released (or regenerated) to join the free KEAP1 dimer pool and able to sequester newly

    synthesized NRF2 again, thus completing a global cycle for KEAP1. Under oxidative stress, this

    cycle is blocked as the NRF2 degradation-coupled release of KEAP1 from the closed

    KEAP1-NRF2 complex is inhibited, leading to accumulation of the closed state and depletion of

    free KEAP1 dimer. Flux analysis indicated that Model 1 operates in equilibrium mode because of

    the much higher association and dissociation fluxes through the DLG-binding step than the

    connected NRF2 turnover fluxes (Tables S10-S11). We thus evolved Model 1 into Model 2 by

    dramatically reducing the parameter values of k_3 and k_4 (the rate constants for DLG binding),

which indeed successfully made Model 2 behave in a cycle mode with divergent responses of

the open and closed KEAP1-NRF2 complexes (Figs. S3A-S3B, and S4). However, this "success"

was achieved by setting k_4, the dissociation rate constants for DLG binding, to a value that is

hundreds-fold lower than experimentally measured (Fukutomi et al. 2014). Importantly, in the

same study, it was also demonstrated that the first binding event, i.e., between KEAP1 and the

ETGE motif of NRF2, is a thermodynamically two-step process, involving an initial fast binding

step to form a transient, intermediate complex (termed *KEAP1_NRF2*$_{open1}$ here) first, followed by

a much slower second step that leads to a more stable configuration of the open complex

(termed *KEAP1_NRF2*$_{open2}$ here). As a minimal model, Models 1 and 2 only considered

single-step ETGE binding. We hypothesized that this second, slow ETGE-binding step may

account for the experimentally observed cycle mode of operation. We next set out to test this

hypothesis with Model 3a.

Model 3a (Two-Step ETGE-Binding Cycle Mode for Class I-V activator)

In Model 3a we added an extra, reversible step, $k_{1.1}$ and $k_{2.1}$, to account for the intramolecular

state transition between *KEAP1_NRF2*$_{open1}$ and *KEAP1_NRF2*$_{open2}$ (Fig. 2A), with k_4 restored to

the high value measured in (Fukutomi et al. 2014). As detailed in Table S1 footnote, we then

iteratively adjusted the values of k_0, k_3 and k_6 such that the basal *NRF2*$_{tot}$ level is still at 150 nM

and half-life at 10 min, and the basal open (*KEAP1_NRF2*$_{open}$ = *KEAP1_NRF2*$_{open1}$ +

KEAP1_NRF2$_{open2}$):closed (*KEAP1_NRF2*$_{closed}$) state ratio remains at 1:1 (Fig. 2B). NRF2 in

KEAP1_NRF2$_{open1}$ and *KEAP1_NRF2*$_{open2}$ were assumed to degrade with the same rate

constants (k_9=$k_{9.1}$=k_5).

At the basal condition, with the default parameter setting of Model 3a, the second step of

ETGE binding ($k_{1.2}$ and $k_{2.1}$) does not operate in equilibrium mode. This is because *flux*$_{k1.1}$ = 0.133

and *flux*$_{k2.1}$ = 1.73E-3 nM/S, thus only a tiny fraction of *KEAP1_NRF2*$_{open2}$ is returned to

KEAP1_NRF2 $KEAP1\_NRF2_{open1}$ (Table S10). Another small fraction is degraded through $flux_{k9.1}$ at 4.1E-3

nM/S. Over 95% of $KEAP1\_NRF2_{open2}$ is moved forward to become $KEAP1\_NRF2_{closed}$ at a net

flux ($flux_{k3}$ - $flux_{k4}$) of 0.127 nM/S. In contrast, both the first step of ETGE binding (k_1 and k_2) and

the step of DLG binding (k_3 and k_4) operate in equilibrium mode, with $flux_{k1}$ and $flux_{k2}$ at 16.48

and 16.33 nM/S respectively, and $flux_{k3}$ and $flux_{k4}$ at 14.19 and 14.07 nM/S respectively, all of

which are >100 fold higher than other connected turnover fluxes (Table S10). As a result, the

$NRF2_{free}$:$KEAP1\_NRF2_{open1}$ ratio, which is 1:9.4, is largely determined by the k_2:$(2*k_1*Keap1_{free})$

ratio, and the $KEAP1\_NRF2_{open2}$:$KEAP1\_NRF2_{closed}$ ratio, which is 1:5, is largely determined by

the k_4:k_3 ratio. At 58 and 14 nM respectively, $KEAP1\_NRF2_{open1}$ dominates $KEAP1\_NRF2_{open2}$,

accounting for 80% of the total open KEAP1-NRF2 complex (Fig. 2B and Table S8).

When setting k_0=0 to examine the decay of NRF2 species from their basal steady states,

$NRF2_{free}$:$KEAP1\_NRF2_{open1}$ and $KEAP1\_NRF2_{open2}$:$KEAP1\_NRF2_{closed}$ remain at the same

equilibrium ratios as above as all NRF2 species decrease (Fig. 2B). $NRF2_{free}$ and

$KEAP1\_NRF2_{open1}$ decrease quickly with a half-life of about 4-5 min, due primarily to the

depletion of $KEAP1\_NRF2_{open1}$ through $flux_{k1.1}$, which is about 8-fold greater than $flux_{k9}$. In

contrast, $KEAP1\_NRF2_{open2}$ and of $KEAP1\_NRF2_{closed}$ do not decrease as fast because of the

continued supply of KEAP1-NRF2 complex through $flux_{k1.1}$. Because of the differential decay

rates, the relative abundance of $KEAP1\_NRF2_{open1}$ and $KEAP1\_NRF2_{open2}$ switches positions

over time, with $KEAP1\_NRF2_{open2}$ becoming the dominant form of the open KEAP1-NRF2

complex eventually. Furthermore, the levels of $KEAP1\_NRF2_{open}$ and $KEAP1\_NRF2_{closed}$ diverge

quickly from the basal ratio of 1:1 to 1:2.5 by 15 min, and to about 1:4.5 eventually. Since toward

the end of the decay process $KEAP1\_NRF2_{open2}$ is the dominant form of the open KEAP1-NRF2

complex, this 1:4.5 ratio closely reflects the equilibrium ratio of

$KEAP1\_NRF2_{open2}$:$KEAP1\_NRF2_{closed}$, which is determined primarily by the k_4:k_3 ratio.

To examine the behavior of Model 3a when NRF2 in $KEAP1\_NRF2_{closed}$ is stabilized, we

first lowered k_6 to different values, while keeping $CLASS_{I-V}$=0. As k_6 decreases from the default

1.775E-3 S^{-1} (equivalent $t_{1/2}$=6.5 min) to 1.252E-4 (which is the default value of k'_6, equivalent

$t_{1/2}$=92 min), all NRF2 species (except $KEAP1\_NRF2_{open1}$) increase and reach steady states in

about 400 min (Fig. 2C). The open:closed ratio decreases and reaches about 1:2.8 at 1 h, and

settles to 1:3.5 at steady state, with $KEAP1\_NRF2_{open2}$ switching to the dominant form of the

open-state complex. When reaching steady states, $NRF2_{tot}$ increases by 5-fold, while $NRF2_{free}$

increases by a much greater fold, from 6.2 to 227 nM (36.6-fold). At this activated state, by

setting k_0=0, all NRF2 species decrease, with a half-life of 68 min for $NRF2_{tot}$, while $NRF2_{free}$

disappears much more quickly. By setting k_6 to even lower values, the maximal levels of both

$NRF2_{free}$ and $NRF2_{tot}$ increase but to a limited extent and the half-life of $NRF2_{tot}$ lengthens to 126

min in the extreme case when k_6=0. (Figs. S5A and S5C).

Therefore, at the basal condition, $KEAP1\_NRF2_{open1}$ is the dominant form of the open

KEAP1-NRF2 complex. However, upon perturbation, either by setting k_0=0 or setting k_6 to a

lower value than the default, the relative abundance between the two open states is switched,

such that $KEAP1\_NRF2_{open2}$ becomes dominant, and then the open:closed ratio will be following

the $KEAP1\_NRF2_{open2}$:$KEAP1\_NRF2_{closed}$ ratio, which is determined largely by k_3 and k_4 as an

equilibrium step. In this sense, although Model 3a behaves as shown above in a cycle mode

globally due to the slow $k_{1.1}$/$k_{2.1}$ steps, locally, some species of the KEAP1-NRF2 complexes still

maintain an equilibrium relationship, due to the high fluxes of the k_1/k_2 and k_3/k_4 binding steps.

With increasing $CLASS_{I-V}$ levels, the temporal behaviors of $NRF2_{tot}$ (Fig. 2D), $NRF2_{free}$

(Fig. 2E), and $KEAP1_{free}$ (Fig. 2F) are similar to Model 2. It takes a longer time for $NRF2_{tot}$ to

reach steady states, while the $NRF2_{free}$ response is initially delayed but its rising time is

shortened with increasing $CLASS_{I-V}$ levels. For steady-state dose-response relationships,

KEAP1_NRF2$_{\text{open2\_tot}}$ (*KEAP1_NRF2*$_{\text{open2}}$ + *KEAP1*$_\text{o}$*_NRF2*$_{\text{open2}}$) and *KEAP1_NRF2*$_{\text{closed\_tot}}$

(*KEAP1_NRF2*$_{\text{closed}}$ + *KEAP1*$_\text{o}$*_NRF2*$_{\text{closed}}$) both increase while remaining at a constant

equilibrium ratio with increasing *CLASS*$_{\text{I-V}}$ levels (Fig. 2G). In contrast, steady-state

KEAP1_NRF2$_{\text{open1\_tot}}$ (*KEAP1_NRF2*$_{\text{open1}}$ + *KEAP1*$_\text{o}$*_NRF2*$_{\text{open1}}$) first increases slightly then

decreases (Fig. 2G). Steady-state *NRF2*$_{\text{free}}$ exhibits an ultrasensitive, sigmoidal dose-response

with respect to *CLASS*$_{\text{I-V}}$ levels, with n_H of 1.78 and *LRC*$_{\text{max}}$ of 2.24 (Fig. 2G).

Flux analysis shows that the *total degradation rate* curve exhibits an S-shape as in Model

2 (Figs. 2H and 2I). However, because of the two-step ETGE binding, higher concentrations of

NRF2$_{\text{free}}$ are required to produce levels of turnover fluxes similar to Model 2, with the *flux*$_{\text{k6}}$ and

flux$_{\text{k9}}$ curves shifted to the right and closer to *flux*$_{\text{k5}}$. This shift leads to a shorter second phase of

the *total degradation rate* curve that is not as flat as in Model 2. As the k_6 value is varied

mimicking different stress levels, the intersection point between *synthesis rate* and *total*

degradation rate curves still swings quite dramatically, albeit not as dramatic as in Model 2 (Figs.

S3G and S3H). As shown in Figs. 2H and 2I, when k_6 is lowered from 5.325E-4 to 1.775E-4, a

3-fold decrease, the corresponding steady-state *NRF2*$_{\text{free}}$ concentration increases by 13-fold,

indicating signal amplification.

Effects of k_1 (k'_1) and k_2 (k'_2)

We next examined the effects of different parameters on the NRF2 response in Model 3a.

Enhancing the ETGE-mediated first-step binding affinity between free KEAP1 and free NRF2, by

increasing k_1 and k'_1 by 10-fold, only marginally decreases the basal *NRF2*$_{\text{tot}}$ level and half-life

(Fig. S6A) with nearly no effect on the steady-state dose-response curve (Fig. 3A). Neither the

basal levels of different open and closed KEAP1-NRF2 complexes nor their steady-state

dose-response curves are affected (Figs. S6E-S6H). In contrast, the basal *NRF2*$_{\text{free}}$ level

decreases dramatically and the ultrasensitivity of the dose-response curve is enhanced markedly

without much change in the maximal level (Fig. 3B). Decreasing k_1 and k'_1 by 10-fold appears to

have slightly larger albeit opposite effects on the various NRF2 species (Figs. 3A and S6), and

dramatically increases the $NRF2_{free}$ level and reduces its ultrasensitivity (Fig. 3B). The time delay

in the $NRF2_{free}$ response disappears with decreasing k_1 and k'_1 (Fig. S6C) and is further

increased with increasing k_1 and k'_1 (Fig. S6D). Varying k_2 and k'_2 has opposite effects as varying

k_1 and k'_1 (simulation results not shown).

Effects of $k_{1.1}$ ($k'_{1.1}$) and $k_{2.1}$ ($k'_{2.1}$)

We next examined the effects of the ETGE-mediated second-step binding, which is much slower

than the first step and is the key step for making Model 3a behave in a cycle mode. Increasing

$k_{1.1}$ and $k'_{1.1}$ shifts the balance between the two open states, causing a reduction in

$KEAP1\_NRF2_{open1\_tot}$ (Fig. S7E) but only a slight increase in $KEAP1\_NRF2_{open2\_tot}$ (Fig. S7F) and

$KEAP1\_NRF2_{closed\_tot}$ (Fig. S7H), resulting in a net decrease of the total open state

$KEAP1\_NRF2_{open\_tot}$ (Fig. S7G). As a result, the basal level of $NRF2_{tot}$ is reduced with a slight

decrease in its half-life (Fig. S7A) and the steady-state dose-response curve becomes steeper

(Fig. 3C). In comparison, the basal level of $NRF2_{free}$ is considerably reduced and the

ultrasensitivity of the dose-response curve is dramatically enhanced with little change in the

maximal response level (Fig. 3D). Decreasing $k_{1.1}$ and $k'_{1.1}$ has the opposite but generally larger

effects. It causes an increase in $KEAP1\_NRF2_{open1\_tot}$ (Fig. S7E) and a decrease in

$KEAP1\_NRF2_{open2}$ (Fig. S7F) and $KEAP1\_NRF2_{closed}$ (Fig. S7H), resulting in a net increase of

$KEAP1\_NRF2_{open\_tot}$ which seems to have a flat response to $CLASS_{I-V}$ (Fig. S7G). As $k_{1.1}$ and $k'_{1.1}$

are decreased by 10-fold, the basal level of $NRF2_{tot}$ is dramatically increased with its half-life

lengthened (Fig. S7A) and the steady-state dose-response curve becomes much shallower (Fig.

3C). The basal level of $NRF2_{free}$ is considerably elevated and the ultrasensitivity of its

dose-response curve is dramatically reduced (Fig. 3D). The time delay in the $NRF2_{free}$ response

disappears with decreasing $k_{1.1}$ and $k'_{1.1}$ (Fig. S7C) and is further increased with increasing $k_{1.1}$

and $k'_{1.1}$ (Fig. S7D). Varying $k_{2.1}$ and $k'_{2.1}$, especially when lowering the values, seems to affect

$KEAP1\_NRF2_{open1\_tot}$ the most, with a minimal effect on all other NRF2 species (Figs. 3E-3F and

S7I-S7P), which is consistent with the low backward flux nature of this second-step ETGE

binding, where the backward flux ($flux_{k2.1}$ + $flux_{k'2.1}$) is only a tiny fraction of the forward flux

($flux_{k1.1}$ + $flux_{k'1.1}$).

Effects of k_3 (k'_3) and k_4 (k'_4)

We next examined the effects of DLG-mediated binding. Increasing k_3 and k'_3 by 10-fold reduces

the $KEAP1\_NRF2_{open2\_tot}$ level dramatically across the range of $CLASS_{I-V}$ levels as expected (Fig.

S8F). However, it only marginally decreases the basal $KEAP1\_NRF2_{open1\_tot}$ (Fig. S8E) and

increases the basal $KEAP1\_NRF2_{closed\_tot}$ (Fig. S8H) levels. At high $CLASS_{I-V}$ levels,

$KEAP1\_NRF2_{open1\_tot}$ is suppressed considerably and $KEAP1\_NRF2_{closed\_tot}$ increases to higher

levels. These changes have slight effects on the basal $NRF2_{tot}$ level and its half-life (Fig. S8A),

and the steady-state dose-response curve (Fig. 3G). The basal $NRF2_{free}$ level decreases

marginally and the ultrasensitivity of the steady-state dose-response curve barely increases with

a slightly higher maximal level (Fig. 3H). Decreasing k_3 and k'_3 by 10-fold has opposite but larger

effects on the various species. With $KEAP1\_NRF2_{open2\_tot}$ at higher levels (Fig. S8F),

$KEAP1\_NRF2_{open1\_tot}$ (Fig. S8E) becomes higher and $KEAP1\_NRF2_{closed\_tot}$ (Fig. S8H) becomes

lower. Both basal $NRF2_{tot}$ and $NRF2_{free}$ levels increase and maximal response levels decrease,

reducing their ultrasensitivity (Figs. 3G and 3H). The time delay in the $NRF2_{free}$ response does

not appear to be affected by k_3 and k'_3 (Figs. S8B-S8D). Varying k_4 and k'_4 has opposite effects

as varying k_3 and k'_3, and reducing k_4 and k'_4 to zero thus making the DGL-mediated binding

irreversible has a similar effect as reducing k_4 and k'_4 by 10-fold (simulation results not shown).

Effects of hinge-latch mode of operation

The hinge-latch hypothesis states that under oxidative stress by class I-V oxidants, the

DLG-mediated binding is weakened, likely due to the cysteine modification on KEAP1 in multiple

domains, and the level of the closed KEAP1-NRF2 complex is reduced so that NRF2 is no longer

destabilized (Tong et al. 2006, Tong et al. 2007, Fukutomi et al. 2014). Here we used Model 3a to

explore the effects of the hinge-latch hypothesis. When setting k'_3 (which is the association rate

constant for the intramolecular binding between oxidized KEAP1 and DLG motif) to a lower value

(0.1 of default) to mimic a hinge-latch mode of operation, a high $CLASS_{I-V}$ level lead to increases

in both the open and closed states (Figs. 4A and 4B). However, the open state level is higher

than the closed state, which runs counter to the decreasing open:closed ratio under oxidative

stress as expected (Baird et al. 2013). The hinge-latch simulation also predicts more muted

maximal responses of $NRF2_{tot}$ (Fig. 4E) and $NRF2_{free}$ (Fig. 4F). Interestingly, increasing k'_3 to

simulate strengthened DLG binding under oxidative stress has the opposite effect: the

open:closed ratio further increases (Figs. 4C and 4D) and the $NRF2_{tot}$ (Fig. 4E) and $NRF2_{free}$ (Fig.

4F) dose-responses exhibit higher maximal levels and enhanced ultrasensitivity, although these

changes approach a limit as k'_3 is increased by > 10-fold. Changing the DLG binding affinity by

varying k'_4 has opposite effects as varying k'_3 (simulation results not shown). Therefore, with

current parameter settings, , the hinge-latch mode of operation is predicted to be less effective in

activating NRF2 by class I-V compounds.

Effects of KEAP1 abundance

The relative abundance of KEAP1 and NRF2 can have important effects on NRF2 activation.

The current default basal $NRF2_{tot}$:$KEAP1_{tot}$ ratio is about 1:4. Increasing $KEAP1_{tot}$ by up to

10-fold has little effect on the basal $NRF2_{tot}$ level and its half-life (Fig. S10A). This lack of effect is

because at the default $KEAP1_{tot}$ level, there is already sufficient KEAP1 to sequester the majority

of NRF2, so increasing $KEAP1_{tot}$ further does not alter the fraction of NRF2 in complex with

KEAP1 much, including the closed state which is most actively degraded. But the maximal level

of the dose-response curve of $NRF2_{tot}$ increases (Fig. 5A) and this occurs because NRF2 in

$KEAP1_o\_NRF2_{closed}$ is not degraded as readily as $NRF2_{free}$ or NRF2 in the open state. Increasing

total KEAP1 abundance reduces basal $NRF2_{free}$ and the maximal response levels dramatically

(Fig. 5B). The muted response is mostly due to the increased sequestering effect of higher

KEAP1 abundance. When $KEAP1_{tot}$ is reduced from its default value, basal $NRF2_{tot}$ levels and its

half-life increase (Fig. S10A), and the dose-response curve becomes shallower with lower

maximal response levels (Fig. 5A). Basal $NRF2_{free}$ increases dramatically with little further

increase in response to $CLASS_{I-V}$ at higher levels, indicating constitutive activation of NRF2 (Fig.

5B). These results suggest that there is an optimal NRF2:KEAP1 ratio that can maximize the

dynamic range of free NRF2 in response to oxidative stress. If the ratio is too low or too high, the

response of free NRF2 is muted.

Model 3b (Two-Step ETGE-Binding Cycle Mode for Class VI inducer)

Since Model 3a represents the most updated biology of KEAP1 and NRF2 interactions, the

remaining Models (3b, 4a and 4b) are based on this model. In Model 3b, we simulated class VI

NRF2 activators, which activate NRF2 by competing with NRF2 for binding to the DC domain of

KEAP1 (Hancock et al. 2012, Jiang et al. 2014, Lazzara et al. 2020). Model 3b keeps the

interactions between KEAP1 and NRF2 at the basal condition as in Model 3a, but differ in how

$CLASS_{VI}$ interacts with KEAP1 (Fig. 6A). We assume that a $CLASS_{VI}$ molecule can bind equally

to either of the two monomeric subunits in KEAP1 dimer that is not occupied by NRF2. It is thus

possible that a KEAP1 dimer can be occupied by 2 molecules of a $CLASS_{VI}$ compound such that

no NRF2 is able to bind to this KEAP1 dimer. This assumption is well justified as It has been

recently demonstrated that NRF2 can be progressively and ultimately completely liberated off

KEAP1 by increasing concentrations of p62 and other KEAP1-NRF2 interaction inhibitors (Horie

et al. 2021). Unlike the case with $CLASS_{I-v}$ activators, in response to $CLASS_{VI}$, $NRF2_{free}$

increases immediately without delay, followed by a slower rise over time to reach the steady

state in about 300 min (Fig. 6C). The initial rapid response of $NRF2_{free}$ is due to the titration of

KEAP1 by $CLASS_{VI}$, resulting in immediate liberation of NRF2 from the KEAP1-NRF2 complexes.

The subsequent slow $NRF2_{free}$ rise occurs because more KEAP1-NRF2 complex shifts away

from the rapidly-degrading closed state, resulting in NRF2 stabilization (Fig. 6E). Contrary to

Model 3a for $CLASS_{I-v}$, the higher the $CLASS_{VI}$ level, the long it takes for $NRF2_{free}$ to reach the

steady state (Fig. 6C). $NRF2_{tot}$ has a similar temporal profile to $NRF2_{free}$ except the initial

fast-rising phase (Fig. 6B). The steady-state dose responses of $NRF2_{free}$ and $NRF2_{tot}$ are shown

in Fig. 6D, with $NRF2_{free}$ exhibiting an n_H of 1.09 and LRC_{max} of 0.92. The $NRF2_{free}$ and $NRF2_{tot}$

responses to low $CLASS_{VI}$ levels are nearly flat, as $CLASS_{VI}$ molecules are first sequestered by

free KEAP1. Contrary to the decreasing open:closed ratio of KEAP1-NRF2 complexes under

$CLASS_{I-v}$, this ratio increases by $CLASS_{VI}$ (Fig. 6E). At high $CLASS_{VI}$ levels, the half-life of

$NRF2_{tot}$ approaches 40 min, which is also the half-lives of $NRF2_{free}$ and $KEAP1\_NRF2_{open\_tot}$ (Fig.

S11). We also explored the situation when only one KEAP1 monomeric subunit can be occupied

by class VI activators by setting both k'_7 and k'_8 to zero. As shown in Figs. 6F and 6G, this

configuration does not affect $NRF2_{tot}$, but weakens the $NRF2_{free}$ response as its maximal level

cannot reach as high as when both KEAP1 monomeric subunits can be occupied by class VI

activators. This more muted response is because without class VI activators blocking both

binding sites on KEAP1 dimer, NRF2 can still be sequestered by KEAP1 through the ETGE motif,

resulting in lower $NRF2_{free}$ levels.

Model 4a (With Nucleus for Class I-V activators)

Since NRF2 that translocates to the nucleus is what ultimately drives target gene expression, we

next explored the situation when a nuclear compartment is added. The following assumptions

were made regarding NRF2 translocation between the cytosol and nucleus (Fig. 7A). (i) The

binding kinetics between free nuclear NRF2 ($NRF2_{free\_nucleus}$) and free nuclear KEAP1 dimer

($KEAP1_{free\_nucleus}$) are the same as in the cytosol. (ii) KEAP1-mediated NNRF2 ubiquitination and

degradation does not occur in the nucleus. Therefore, the degradation rate constants of various

NRF2 species in the nucleus are the same as in the cytosol, except for the closed KEAP1-NRF2

complex, which is degraded with the same rate constant as other NRF2 species. (iii) NRF2

activators do not modify or bind to KEAP1 in the nucleus to regulate NRF2 stability.

At the basal condition, total nuclear NRF2 ($NRF2_{tot\_nucleus}$) is at 278 nM as observed in

RAW 264.7 cells (Iso et al. 2016) and a significant fraction of which is titrated by KEAP1 such

that $NRF2_{free\_nucleus}$ is at 180 nM (Table S8). When setting k_0=0, $NRF2_{tot\_cytosol}$, $NRF2_{tot\_nucleus}$, and

$NRF2_{tot\_cell}$ all decay but at different paces, with corresponding half-lives of about 11, 28, 18 min,

respectively (Fig. 7B). When viewed on log scale, it is apparent that $NRF2_{tot\_cell}$ decays in two

phases, a fast phase followed by a slow one (Fig. 7C). This two-phase decay profile of total

cellular NRF2 is caused by the fast cytosolic and slow nuclear decay and has been observed

experimentally in a variety of cell lines (Khalil et al. 2015). Under high stress when $CLASS_{I\text{-}v}$ = 1000

nM, the half-life of $NRF2_{tot\_cell}$ markedly lengthens to 55 min (Fig. 7D). In response to a range of

$CLASS_{I\text{-}v}$ levels, free and total NRF2 in both cytosol and nucleus rise and reach steady states in

about 300 min (Fig. S12). In contrast to Model 3a which does not have the nucleus compartment,

$NRF2_{free\_cytosol}$ rises to much lower levels (Fig. S12A) as most of it translocates into the nucleus

elevating $NRF2_{free\_nucleus}$ (Fig. S12C). The steady-state dose-response relationship for

$NRF2_{free\_nucleus}$ exhibits a shallow response, with n_H of 1.15 and of LRC_{max} of 0.31 (Fig. 7E). The

maximal response levels of $NRF2_{tot\_nucleus}$ and $NRF2_{free\_nucleus}$ increase by 2.6 and 3.5-fold

respectively, while those of $NRF2_{tot\_cytosol}$ and $NRF2_{free\_cytosol}$ both increase by about 3.2 and

3.1-fold, respectively (Tables S8 and S9). Thus, with a nuclear load, NRF2 activation is not as

robust as when the action is limited to the cytosol only. This lessor response contrasts with the

nearly 10-fold increase in nuclear NRF2 under exposure to DEM at 100 μM for 3 h observed in

RAW 264.7 cells which our model is partially based upon (Iso et al. 2016).

The overall muted response of Model 4a is due to the following reasons. At the basal

condition, the net influx of NRF2 from the cytosol to nucleus is $flux_{k10}$ - $flux_{k11}$ = 0.0434 nM/s,

which is about 22% of k_0 (0.1933), the NRF2 synthesis rate in the cytosol. Therefore, even if a

$CLASS_{I-V}$ activator can divert all synthesized NRF2 into the nucleus, the total nuclear NRF2 can

only increase by a maximal 4.45-fold (0.193/0.0434) with a constant nuclear NRF2 half-life. We

wondered if the relative abundance of nuclear KEAP1 and NRF2 plays a role in determining the

magnitude of the nuclear NRF2 response. When $KEAP1_{tot\_nucleus}$ abundance is increased (with

k_{10} adjusted simultaneously to maintain the same basal $NRF2_{tot\_cytosol}$ and $NRF2_{tot\_nucleus}$

concentrations), the simulations showed that both the basal and maximally-induced levels of

$NRF2_{free\_nucleus}$ decease because of the sequestering effect of KEAP1 (Fig. S13C). However, the

degree of ultrasensitivity of the $NRF2_{free\_nucleus}$ dose-response curve seems to be optimal when

$KEAP1_{tot\_nucleus}$ is at an intermediate abundance. Increasing $KEAP1_{tot\_nucleus}$ also leads to

changes in the maximal levels of $NRF2_{tot\_cytosol}$, $NRF2_{free\_cytosol}$, and $NRF2_{tot\_nucleus}$ (Fig. S13), but

the fold-increase of $NRF2_{tot\_nucleus}$ remains relatively low. These results suggest that other

mechanisms, as described in the Discussion, may operate *in vivo* to produce a more robust

nuclear NRF2 response. One possibility is a smaller nuclear NRF2 load at the basal condition.

As shown in Fig. S14, when reducing the basal $NRF2_{tot\_nucleus}$ level and thus the nuclear load

through simultaneously adjusting k_0 and k_{10} while maintaining the same basal $NRF2_{tot\_cytosol}$ level,

the magnitude of both $NRF2_{tot\_nucleus}$ and $NRF2_{free\_nucleus}$ responses improve considerably.

Model 4b (With Nucleus for Class VI activators)

We next considered the situation of a class VI activator which competes with NRF2 for binding to

KEAP1 in Model 4b (Fig. 8A). The model assumptions are similar to Model 4a and the class VI

activator only operates in the cytosol. Model 4b has an interesting dynamic. In response to a

range of $CLASS_{VI}$ levels, there is a quick spike in $NRF2_{free\_cytosol}$ within a couple of minutes

followed by a slow rise (Fig. 8B). Correspondingly, $NRF2_{tot\_cytosol}$ decreases immediately followed

by a slower increase before setting to steady states (Fig. 8C). The rapid increase in

$NRF2_{free\_cytosol}$ results from the immediate liberation of NRF2 from the KEAP1-NRF2 complex,

and the liberated NRF2 moves quickly into the nucleus, causing $NRF2_{free\_nucleus}$ (Fig. 8D) and

$NRF2_{tot\_nucleus}$ to rise quickly followed by a slower increase to steady states. Under high stress

when $CLASS_{VI}$ = 1000 nM, the half-life of $NRF2_{tot\_cell}$ lengthens to 40 min (Fig. 8E), shorter than

that in Model 4a. However, the steady-state $NRF2_{free\_nucleus}$ and $NRF2_{tot\_nucleus}$ levels can increase

to higher levels, maximally by 6 and 4.2-fold from their basal levels, respectively (Tables S8 and

S9). This is because by outcompeting NRF2 for KEAP1, $CLASS_{VI}$ can drive more NRF2 into the

nucleus (Fig. S12C vs. Fig. 8D). The steady-state dose-response curve $NRF2_{free\_nucleus}$ is shallow,

with n_H of 1.09 and of LRC_{max} of 0.46 (Fig. 8F). Interestingly, the steady-state dose-response

curve of $NRF2_{tot\_cytosol}$ monotonically decreases, from the basal 150 nM to 33 nM for higher levels

of $CLASS_{VI}$. This decrease occurs because KEAP1 dimer is gradually titrated away by $CLASS_{VI}$

activator, leaving fewer NRF2 in the KEAP1-bound form (Fig. 8G), and more NRF2 translocates

to the nucleus. As in Model 4a, varying nuclear KEAP1 and lowering basal nuclear load of NRF2

turnover can also improve the magnitude and ultrasensitivity of $NRF2_{free\_nucleus}$ (Figs. S15 and

S16).

DISCUSSION

NRF2 activation is an essential step toward the transcriptional induction of adaptive antioxidant

responses. It is mediated via a unique mechanism of protein stabilization where KEAP1 functions

as both a redox sensor and regulator. While the molecular interactions involved in this process

have been well characterized qualitatively and to some extent quantitatively, the quantitative

systems-level behaviors of this redox transducer module are still poorly understood. In the

present study, we explored the steady-state and dynamic behaviors of the KEAP1-NRF2

interactions through a series of mathematical models of increasing complexity. Our simulations

demonstrated that the kinetic details of the molecular interactions between KEAP1 and NRF2

play critical roles in determining the redox signaling properties.

Basal NRF2 half-life in relation to different NRF2 states

A prominent function of KEAP1 is to act as an E3 ligase adaptor to promote NRF2 ubiquitination

and degradation. This function is critically dependent on the configuration of the KEAP1-NRF2

complex. It is well-established that for the ubiquitination and degradation of NRF2 to occur, the

KEAP1-NRF2 complex has to be in the closed state, i.e., both of the two binding sites in the

cysteine-unmodified KEAP1 dimer have to be engaged by the ETGE and DLG motifs of the

same NRF2 molecule. Therefore, the fraction of this closed state and the rate at which NRF2

within this closed KEAP1-NRF2 complex is ubiquitinated and degraded are key determinants for

the half-life of NRF2 in the cytosol.

In the confine of the present model structure, NRF2 exists in three forms: free, open

KEAP1-NRF2 complex, and closed KEAP1-NRF2 complex. The relative abundances of these

forms at the basal steady state are determined by the binding parameters as well as the

degradation rate constant of each NRF2 form. Given the high binding affinity between KEAP1

and ETGE, it is expected in theory and shown by our simulation that the fraction of free cytosolic

NRF2 is very small, when KEAP1 is not limiting, and NRF2 exists predominantly in the complex

forms at the basal condition. Using FRET to track the open and closed states of the

KEAP1-NRF2 complex, Baird 2013 showed that at least in HEK293 cells, the open:closed ratio

of the KEAP1-NRF2 complex is near 1:1 under nonstressed conditions (Baird et al. 2013). The

half-life of total NRF2 in whole cells at basal conditions is short, mostly ranging between 6-20 min

depending on cell types (Kwak et al. 2002, Alam et al. 2003, Itoh et al. 2003, Stewart et al. 2003,

Kobayashi et al. 2004, He et al. 2006, Khalil et al. 2015, Crinelli et al. 2021). Since nuclear NRF2

is relatively more stable than cytosolic NRF2 (Itoh et al. 2003, Burroughs et al. 2018) and often

constitutes a considerable fraction of total NRF2 at the basal condition (Khalil et al. 2015, Iso et

al. 2016), it is expected that cytosolic NRF2 is actually degraded at even faster rates than

measured in whole cells. The comparable basal abundance of open and closed KEAP1-NRF2

complexes suggests that NRF2 in the closed form has to be degraded very fast with a half-life of

its own that is much shorter than the averaged half-life of total NRF2. In our models, parameter

k_6 governs the degradation of this NRF2 form. With an apparent half-life of cytosolic total NRF2

around 10 min at the basal condition, the default values of k_6 across the six models correspond

to an half-life of 5.7-6.6 min. In comparison, the half-lives of free NRF2 and NRF2 in the open

KEAP1-NRF2 complex, as determined by parameters k_5 and k_9 (and $k_{9.1}$ in the case of two-step

ETGE binding) respectively, are much longer, which is 40 min here, as reported for COS-1 and

HEK293T cells (McMahon et al. 2004, Rada et al. 2011). If k_5, k_9, and $k_{9.1}$ are set lower than the

current default value, k_6 needs to be even higher to maintain the same basal total NRF2 half-life.

Therefore, the turnover of basal NRF2 is predominantly routed through the closed KEAP1-NRF2

complex, and the apparent half-life of cytosolic total NRF2 is determined by the fraction of the

closed complex. In Model 1 which operates in an equilibrium mode, this fraction remains

constant at 50% at all times (Fig. S1A), therefore the instantaneous half-life of total NRF2 at any

given moment is fixed. In the remaining models which operate in a cycle mode, the fraction

increases dynamically and becomes dominant over other NRF2 forms during the decay process

(Figs. S3A and 2B), resulting in a nonlinear degradation of total NRF2 with shortening

instantaneous half-life. In Models 4a and 4b, which has the nucleus compartment, cellular total

NRF2 decays with a two-phase profile, which has been observed experimentally in a variety of

cell types (Khalil et al. 2015), reflecting the differential half-lives of cytosolic and nuclear NRF2.

Equilibrium vs cycle mode of operation

The comparable abundance of the open and closed states of the KEAP1-NRF2 complex at the

basal condition can be achieved in theory in two ways, depending on the transition fluxes ($flux_{k3}$

and $flux_{k4}$) between the two states relative to other turnover fluxes ($flux_{k5}$, $flux_{k6}$, $flux_{k9}$ and $flux_{k9.1}$).

If the transition fluxes are much higher than the turnover fluxes, then the open and closed states

of the KEAP1-NRF2 complex operate in an equilibrium mode, which means that the ratio of the

two states is predominantly determined by the k_3:k_4 ratio regardless of other parameter values.

Parameters k_3 and k_4 describe the DLG-mediated KEAP1 and NRF2 binding. In the literature, its

binding kinetics was determined *in vitro* by using mouse KEAP1-DC fragment and NRF2-Neh2

domain fragment (Tong et al. 2006) or extended DLG motif peptide (DLGex) (Fukutomi et al.

2014). However, *in vivo* the DLG binding is mostly an intra-molecular event within the open

KEAP1-NRF2 complex, since the ETGE motif has a much higher binding affinity for KEAP1 and

ETGE-mediated binding almost always occurs first to form the open-state complex (Stewart et al.

2003). In such *in vivo* scenario, k_3 is actually a first-order, as opposed to a second-order,

association rate constant while k_4 remains as a first-order dissociation rate constant. It is unclear

whether in the open state the DLG binding is enhanced since the DLG is in a closer vicinity to the

unoccupied KEAP1 binding site than the DLG in a free NRF2 molecule not yet attached to

KEAP1. Regardless, in our first trial, in Model 1 we used the k_4 value measured *in vitro* (Fukutomi

et al. 2014) and adjusted k_3, as detailed in Table S1 footnote, to achieve a 1:1 ratio for the basal

open:closed states. Examining the fluxes clearly revealed that with these parameter settings,

$flux_{k3}$ and $flux_{k4}$ are absolutely dominant over other turnover fluxes (Table S10). As shown in Fig.

S1, the open and closed KEAP1-NRF2 complexes remain at a 1:1 ratio in all perturbed

conditions including shutdown of NRF2 synthesis (Fig. S1A), stabilization of NRF2 in the closed

state (Fig. S1B), and under a wide range of $CLASS_{I\text{-}V}$ levels (Fig. S1C), demonstrating that Model

1 definitely operates in an equilibrium mode. However, the experimental study by Baird clearly

demonstrated that under various perturbations similar to above, the closed KEAP1-NRF2 state

will eventually dominate over the open state, thus negating an equilibrium mode of operation

(Baird et al. 2013). It was further suggested that the KEAP1-NRF2 interaction may operate

instead in a global cycle mode where the transition fluxes between the open and closed states

are not overwhelmingly higher than the turnover fluxes, such that KEAP1 in the complex is

almost always moved forward to the next state and eventually exists via the closed complex

along with NRF2 degradation, and recycled to join the free KEAP1 dimer pool.

The discrepancy between our Model 1's behavior and the cycle mode of operation led us

to Model 2 where the parameters k_3 and k_4 were lowered to alleviate the equilibrium mode of

operation. Model 2 indeed exhibits the behavior consistent with the cycle mode, where the

open:closed ratio decreases in all perturbed conditions (Figs. S3A, S3B, and S3F). The issue

with Model 2 is that parameter k_4, which describes the dissociation rate constant of DLG binding,

is 1.0E-4 S^{-1}, only about 1/2000 of the *in vitro* measured value (Table S1). This value translates

into an average lifetime of 167 min for the closed state before it can revert back into the open

state, which is considered too long for such weak binding (Fukutomi et al. 2014). In the same

study it was also demonstrated that ETGE-mediated binding is actually a two-step process,

involving an initial fast binding step followed by a subsequent slow binding step. We therefore

wondered whether the slow binding here may be responsible for the cycle mode behavior. When

this idea was implemented in Model 3, simulations indeed showed such effects on the

open:closed ratio under all perturbed conditions, including shutdown of NRF2 synthesis (Fig. 2B),

stabilization of NRF2 in the closed state (Fig. 2C), and under a wide range of $CLASS_{I\text{-}V}$ levels

(Fig. 2G). With the parameter setting for the second-step, slow ETGE binding ($k_{1.1}$ and $k_{2.1}$),

flux$_{k1.1}$ is much greater than *flux*$_{k2.1}$ (Table S10). *KEAP1_NRF2*$_{open1}$ is the dominant form of the

open state at the basal condition (Fig. 2B and Table S8), and is not in equilibrium with

KEAP1_NRF2$_{open2}$ (Fig. 2C and 2G). In contrast, *KEAP1_NRF2*$_{open2}$ is always in equilibrium with

KEAP1_NRF2$_{closed}$ at an approximate 1:5 ratio as determined by the k_4:k_3 ratio (Fig. 2G). During

CLASS$_{I-V}$ perturbation, *KEAP1_NRF2*$_{open1}$ decreases while *KEAP1_NRF2*$_{open2}$ increases and

becomes the dominant open form, resulting in an overall open:closed ratio that is close to 1:3.5.

Therefore, although Model 3 behaves globally in a cycle mode at the basal condition with

KEAP1_NRF2$_{open1}$ as the dominant open form, at high stress levels *KEAP1_NRF2*$_{open2}$ becomes

the dominant open form and the system switches to operate largely in equilibrium mode as far as

the overall open:closed ratio is concerned.

Hinge-latch hypothesis and class I-V vs. class VI NRF2 activators

An important theory of NRF2 activation is the hinge-latch hypothesis which postulates that the

ETGE-mediated association (i.e., the open-state complex) is always there functioning as a hinge

between KEAP1 dimer and NRF2, while the weaker DLG-mediated association can be latched

on (i.e., forming the closed-state complex) or off (i.e., reverting to the open state complex) by

oxidative stressors (Yamamoto et al. 2018). With Model 3a we tested the effects of hinge-latch

mode of operation on NRF2 activation by altering the k'_3:k'_4 ratio which governs the

intramolecular DLG binding affinity between oxidized/conjugated KEAP1 and NRF2. Our

simulations showed that when the DLG binding affinity is lowered to mimic a hinge-latch, the

maximally induced steady-state levels of total NRF2 and particularly free NRF2 is tangibly

reduced (Figs. 4E and 4F). In contrast, when the k'_3:k'_4 ratio is made higher, i.e., a strengthening

of the latched-on state under oxidative stress, there is an increase, albeit limited, in the maximal

NRF2 levels. These results suggest that a hinge-latch mode of operation may lead to a lessor

NRF2 response to class I-V compounds. The reason for the more muted response in our model

is because under oxidative stress, the closed KEAP1-NRF2 complex in which the KEAP1

molecule is modified on the sensor cysteine residues, i.e., $KEAP1_o\_NRF2_{closed}$, has a half-life

(determined by k'_6) even longer than those of NRF2 in the free or open complex forms. Therefore,

KEAP1 here reverses the normal role of promoting NRF2 degradation as at the basal condition,

and becomes instead protective of the NRF2 molecule. We also examined the situation when

$KEAP1_o\_NRF2_{closed}$ is not protective of NRF2 by setting k'_6 equal to k_5 such that it degrades with

the same half-life as free NRF2 and as open complex. In this case, the hinge-latch mode of

operation slightly improves the ultrasensitivity of free NRF2 and total NRF2 (Figs. S9E and S9F).

But regardless, when in the hinge-latch mode of operation, the open:closed ratio of

KEAP1-NRF2 complexes increases in response to high $CLASS_{I-V}$ levels (Figs. 4A-4B and

S9A-S9B), as opposed to the expected decrease, suggesting that the hinge-latch hypothesis

may not be valid for class I-V compounds-induced NRF2 activation. Indeed, using titration NMR

spectroscopy, the most recent study by Yamamoto group clearly demonstrated that

modifications of reactive cysteines of KEAP1 by class I-V oxidants and electrophiles, including

CDDO-Im and sulforaphane targeting Cys151 and 15d-PGJ2 targeting Cys288, do not break the

DLG-mediated binding (Horie et al. 2021).

Class VI compounds are those that can bind to the DC region of KEAP1 and thus disrupt

DLG-mediated, and also potentially, ETGE-mediated NRF2 binding. Therefore, the NRF2

stabilization effect of class VI compounds is indirect, by shifting KEAP1-NRF2 complex away

from the ubiquitinatible closed state. An endogenous ligand is p62, which has a

KEAP1-interacting region (KIR) containing a DPSTGE motif that is similar to the ETGE motif of

NRF2 (Lau et al. 2010, Jiang et al. 2015). The motif has similar or even higher binding affinities

for KEAP1 than the DLG motif of NRF2, depending on its phosphorylation status (Komatsu et al.

2010, Ichimura et al. 2013). Small-molecule compounds have also been identified recently as

disruptors of the protein-protein interaction between KEAP1 and NRF2, such as Cpd16 (Jiang et

al. 2014) and several others (Yasuda et al. 2017, Lazzara et al. 2020, Lee and Hu 2020). By

displacing DLG binding preferentially, these compounds make the KEAP1-NRF2 complex

function as a hinge-latch as recently demonstrated experimentally (Horie et al. 2021). Our Model

3b captures the hinge-latch behavior in response to class VI NRF2 activators (Fig. 6), and shows

that the open:closed ratio of KEAP1-NRF2 complex actually decreases with increasing

concentrations of class VI activator (Fig. 6E). Simulations of Model 3b also suggest that when the

two monomeric subunits of KEAP1 dimer can both be occupied by a class VI compound, NRF2

activation is more robust because of simultaneous sequestration of free KEAP1 by the

compound (Figs. 6D vs. 6F). Horie et al. indeed showed that with high enough concentrations,

p62 and small-molecule class I-V compounds can completely dissociate NRF2 from KEAP1

dimers, breaking the ETGE-mediated hinge (Horie et al. 2021). A recently identified endogenous

protein, FAM129B, has both DLG and ETGE motifs on the C terminal and can compete with

NRF2 for KEAP1 binding (Cheng 2019). FAM129B is found to be upregulated in many cancers

which have poor prognosis by promoting NRF2 activation and thus chemoresistance.

Maximal NRF2 activation, ultrasensitivity, and floodgate hypothesis

The maximal fold increase of total NRF2 is determined by the differential half-lives at the basal vs.

severely stressed conditions. With a basal half-life of about 10 min in the cytosol in our models,

and the lengthening of the half-life to over an hour under simulated oxidative stress such as in

Model 3a, total NRF2 increases by 5-fold (Fig. 2D and Tables S8-S9). As discussed above,

under oxidative stress, this model switches from KEAP1-mediated degradation to

KEAP1-mediated stabilization of NRF2, therefore the fold-increase can be even higher when

parameter k'_6 is of lower values (Figs. S5B and S5D). Conversely, the fold-increase becomes

smaller when k'_6 is higher (Fig. S5F). It is also evident that at the basal condition when there is a

higher fraction of the closed KEAP1-NRF2 complex that is rapidly degraded, the system is

poised to produce higher levels of NRF2 in response to stresses, as the closed complex

becomes stabilized by a class I-V compound or dissociated by a class VI compound. In Model 3b

which simulates class VI compounds, because there is no closed KEAP1-NRF2 complex with a

$CLASS_{VI}$ molecule attached, the maximal fold-increase of total NRF2 is limited by the half-lives of

free NRF2 and NRF2 in the open KEAP1-NRF2 complexes. In our study parameters k_5, k'_5, k_9,

k'_9, $k_{9.1}$ and $k'_{9.1}$ govern the degradation of these NRF2 species, which have an equal half-life of

40 min. As a result, the maximal fold-increase of total NRF2 in Model 3b cannot exceed 40/10 =

4-fold (Fig. 6B and Tables S8-S9).

As aforementioned in Introduction, for an adaptive stress response, it is ideal that some

degree of signal amplification, i.e., ultrasensitivity, can be embedded in the feedback circuit to

ensure robust resistance to perturbation. In Models 1, 2, and 3a, free NRF2 exhibits some decent

ultrasensitivity. Molecular mechanisms producing ultrasensitivity can arise from six common

ultrasensitive motifs (Zhang et al. 2013). In the KEAP1-NRF2 module here, it appears that both

zero-order degradation and protein sequestration (molecular titration) are at play simultaneously

to produce NRF2 ultrasensitivity, where the sequestration is mediated by ETGE binding and

zero-order degradation is mediated by saturation of DLG binding. As shown in Figs. S3G-S3H

and 2H-2I, KEAP1-mediated degradation of NRF2 in the closed KEAP1-NRF2 complex will

eventually saturate when all KEAP1 dimers are occupied by NRF2. Around this saturation point,

KEAP1-mediated NRF2 degradation becomes zero order such that any additional increase in

NRF2 will have to rely on KEAP1-independent mechanism to degrade. As a result of the

nonlinear, zero-order degradation, the steady-state total NRF2 abundance may experience

some steep changes around the point of KEAP1 saturation than when no saturation occurs.

Indeed, for Models 1-3b, the n_H of total NRF2 is between 1.17-1.35. But with LRC_{max} between

0.35-42, total NRF2 does not exhibit overt ultrasensitivity because of the high basal level. From

the perspective of free NRF2, this KEAP1 saturation point is also a moment when free NRF2 can

no longer be sequestered by KEAP1, and as a result any additional NRF2 synthesized *de novo*

will remain as free NRF2, leading to a steep increase in its abundance. Therefore, both

zero-order degradation (by providing more NRF2 overall) and protein sequestration are at play

simultaneously to produce the ultrasensitivity of free NRF2.

Conceivably, the abundance of total KEAP1 and whether NRF2 can accumulate to a level

that surpasses this abundance play a critical role in quantitative NRF2 activation. If total NRF2

can never increase to a level higher than KEAP1 dimer, then NRF2 cannot escape the

sequestration by KEAP1 and there will be no ultrasensitivity of free NRF2. This is first illustrated

by setting k'_6 to a higher value such that total NRF2 can only barely match the level of total

KEAP1 (Figs. S4F and S5F). The intracellular NRF2:KEAP1 ratio at basal conditions varies

among different cell types, which can be lower or higher 1:1 (Khalil et al. 2015, Iso et al. 2016).

Since nuclear NRF2 often constitutes a considerable fraction of total NRF2 at basal conditions

(Khalil et al. 2015, Iso et al. 2016), the cytosolic NRF2:KEAP1 ratio can be actually even lower

than the values reported for the whole cells. Varying the abundance of KEAP1 in the models has

some interesting results. By increasing KEAP1, its role in further destabilizing Nrf2 is limited

because it is already in excess, and as a result total NRF2 does not increase further (Fig. 5A).

But increasing KEAP1 will be more effective as a sequester to inhibit Nrf2. There seems to be an

optimal KEAP1 abundance that can produce the steepest $NRF2_{free}$ response (Fig. 5B). When

KEAP1 is too low, NRF2 is constitutively activated, but when KEAP1 is too high, free NRF2 is

constitutively suppressed.

This sequestering role of KEAP1 is consistent with the floodgate hypothesis which

postulates that stabilization of NRF2 due to loss of KEAP1 activity as an E3 ligase adaptor

protein is not sufficient to initialize NRF2 nuclear translocation; NRF2 has to accumulate to a

higher level to overflood the KEAP1 gate to move to the nucleus (Yamamoto et al. 2018). A

potential caveat of this mechanism is that it takes some time to produce enough NRF2 to

saturate KEAP1, therefore the free NRF2 response can be delayed as we demonstrated with our

models. However, it is likely that at the basal condition, the cytosolic NRF2:KEAP1 ratio is near

parity in some cells such that KEAP1 is near saturation. As a result, the system is at a tipping

point, poised to respond to a slight increase in oxidative stress to overwhelm KEAP1 and cause

an immediate and steep increase in free NRF2 (Baird et al. 2013).

While both Models 2 and 3a are cycle models, the ultrasensitivity of free NRF2 exhibited

by Model 3a, which has two-step ETGE binding, is somehow weaker than Model 2. As described

in Results, this is partly because a higher free NRF2 level is required to maintain the same

turnover fluxes through different NRF2 species (Figs. 2H and 2I) at the basal steady state in

Model 3a, resulting in a lesser zero-order degradation effect. The apparent dissociation constant

for the ETGE-mediated two-step binding is 7.54 nM, which is lower than the 20 nM used in Model

2. However, at the basal condition, free NRF2 is higher in Models 3a than Model 2 as a result of

the two-step binding and slow fluxes through the second-step binding. A higher basal level will

always reduce the degree of ultrasensitivity (Zhang et al. 2013), despite that Models 2 and 3a

have comparable maximally induced free NRF2 levels (Table S9). By increasing the binding

affinity of ETGE, e.g., through increasing k_1 and $k_{1.1}$, to reduce basal free NRF2, the

ultrasensitivity of Model 3a is improved dramatically (Figs. 3B and 3D).

With both n_H and LRC_{max} of free NRF2 close to unity, the ultrasensitivity for class VI

compounds as in Model 3b is basically absent. In contrast to class I-V compounds, a class VI

compound does not need to induce total NRF2 to a level that exceeds total KEAP1 to produce

tangible increase in free NRF2. This is because when a class VI compound can bind to both of

the monomeric subunits of KEAP1 dimer, it would titrate free KEAP1 away, essentially lowing the

amount of available KEAP1 that can sequester NRF2. This lowering of the "floodgate" can result

in a much higher level of free NRF2 that can be maximally induced by class VI compounds than

class I-V compounds (Figs. 6D vs. 2G and Table S9). However, because of the reduced

sequestration by KEAP1, the ultrasensitivity of free NRF2 is lost.

Response of nuclear NRF2

To transcriptionally regulate its target genes, NRF2 needs to translocate into the nucleus where it

dimerizes with sMaf to gain affinity for the AREs in promoters. The flux of nuclear translocation

constitutes a load to the cytosolic NRF2. At a constant NRF2 production rate in the cytosol, this

nuclear load is expected to alter the dynamics of NRF2 activation. With Models 4a and 4b we

made the assumptions that KEAP1 and NRF2 interactions in the nucleus follow the same kinetic

parameters as in the cytosol except that nuclear KEAP1 is not able to mediate the ubiquitination

and degradation of NRF2 and is not subject to redox modification by class I-V compounds or

binding by class VI compounds. With higher abundance of nuclear NRF2 than KEAP1, as

observed in RAW 264.7 cells and potentially many other cell types (Iso et al. 2016), nuclear

KEAP1 is nearly saturated by NRF2, resulting in low basal free nuclear KEAP1 dimer and high

free nuclear NRF2 levels. These configures result in a net nuclear influx of NRF2 that is 22% of

the NRF2 production rate in the cytosol (Table S10). Therefore, net nuclear importing of NRF2

constitutes a significant load of NRF2 production. It can thus be estimated that even under

oxidative stress that completely terminates cytosolic NRF2 degradation and all cytosolic NRF2

translocates into the nucleus, total nuclear NRF2 cannot increase by > 5-fold. Our simulations

confirmed this prediction (Figs. 7E and 8F, Tables S8-S9), and the fold increase of nuclear free

NRF2 is only slightly higher than the total. Class VI activators seem to has a larger effect on

maximal nuclear NRF2 (4.2-fold) than class I-V activators (2.6-fold). This is due to the

KEAP1-titrating effects of a class VI activator, which reduces free cytosolic KEAP1, pushing

more NRF2 into the nucleus. Nonetheless these lessor responses contrasts with the nearly

10-fold increase in nuclear NRF2 under exposure to DEM at 100 μM in RAW 264.7 cells which

our model is partially based upon (Iso et al. 2016). For nuclear Nr2 to increase to higher levels,

additional mechanisms have to be at play which are not included in our models. These include (i)

increased NRF2 production through transcriptional autoregulation, which has been confirmed in

many cell types including RAW 264.7 cells (Kwak et al. 2002, Pi et al. 2003, Pi et al. 2008); (ii)

reduced nuclear exporting of NRF2 due to redox-sensitive cysteine modification of the nuclear

export signal (NES) sequence in the Neh5 domain of NRF2 (Li et al. 2006); (iii) stabilization of

nuclear NRF2 under oxidative stress; (iv) lower nuclear NRF2 load at the basal condition such

that there is still more reserve capacity for nuclear NRF2 accumulation.

Free nuclear NRF2 does not exhibit overt ultrasensitivity in either of the two models. Part

of the reason is due to its high basal level and smaller fold increase of total nuclear NRF2

discussed above. However, a number of mechanisms that can potentially contribute to

ultrasensitivity have been confirmed in the KEAP1-NRF2 system. These mechanisms include (i)

positive transcriptional autoregulation of both NRF2 and sMaf (Kwak et al. 2002, Katsuoka et al.

2005), (ii) molecular titration of sMaf by inhibitor Bach1 (Igarashi et al. 1998), (iii) positive

feedback through NRF2 induction of p62 which can titrate KEAP1 away from NRF2 and also

promote KEAP1 autophagy (Katsuragi et al. 2016), and (iv) multi-step signaling through (a)

enhanced nuclear NRF2 accumulation due to redox modification of NES as mentioned above (Li

et al. 2006) and (b) redox-sensitive nuclear exporting of Bach1 (Suzuki et al. 2003,

Dhakshinamoorthy et al. 2005). It is highly likely that these mechanisms converge to produce

ultrasensitive nuclear free NRF2 accumulation.

Limitations

The KEAP1-NRF2 module has been modeled mathematically as part of larger networks. We

have constructed NRF2-mediated pathways of antioxidant induction and phase II enzyme

induction, containing negative feedback, incoherent feedforward, and a variety of ultrasensitive

motifs to understand the nonlinear dose-response relationship under oxidative stress (Zhang and

Andersen 2007, Zhang et al. 2009). Blis and his colleagues adapted these models to interpret

and predict antioxidant gene induction in human renal cells in response to cyclosporine (Hamon

et al. 2014), and glutathione depletion in liver microfluidic chips in response to flutamide (Leclerc

et al. 2014). Khalil et al. constructed a model of KEAP1-NRF2/sMaf-ARE activation and its

interaction with the peroxiredoxin and thioredoxin antioxidant enzymes in controlling intracellular

H_2O_2 levels and regulating the reduction of KEAP1, in which one-step ETGE binding was

considered (Khalil et al. 2015). Xue et al. observed a basal NRF2 cytosol-nucleus oscillation

behavior in cells with a period of about 2 hours for which they constructed a mathematical model

of negative feedback through NRF2 phosphorylation and dephosphorylation without involving

changes in the abundance (Xue et al. 2015). Kolodkin et al. has recently incorporated the

KEAP1-NRF2 component into an ROS dynamic network to explore the design principles relevant

to network-based therapies for Parkinson disease (Kolodkin et al. 2020). Compared to the

previous work, our present study provided a much more detailed analysis of the KEAP1-NRF2

module itself, which can be adapted and included in future systems-level models of antioxidant

and detoxification responses.

There are several limitations of the present study, however. In the models we have

limited the action of class I-V and VI compounds on the KEAP1 molecules in the cytosol only,

however it is possible that these compounds, especially class VI, may still compete for KEAP1 in

the nucleus to further drive NRF2 activation. We have assumed separate pools of cytosolic and

nuclear KEAP1 without exchange. However, it has been shown that KEAP1 may also control

postinduction repression of the NRF2-mediated antioxidant response by escorting NRF2 out of

the nucleus (Sun et al. 2007). The DLG-mediated binding kinetics has been measured *in vitro*

with peptide fragments of KEAP1 and NRF2 as inter-molecular event following the law of mass

action. Measuring the binding kinetics as an intra-molecular event as occurring with full-length

proteins will help reduce the uncertainty of model parameterization. It is also unclear whether the

modification of cysteine residues of both KEAP1 subunits of the dimer will have any differential

effects on NRF2 ubiquitination than only one subunit is modified. Lastly, the parameterization

and calibration of our models are based on experimental measurements from multiple cell types,

such as RAW 264.7 and HEK293 cells, and under various experimental conditions. Therefore,

the parameter values and model responses do not represent an ideal "average" cell. However,

we systemically varied parameters where applicable in our study to explore their effects on NRF2

response. All in all, future iterations of the KEAP1-NRF2 model should address these limitations

as more quantitative information, such as binding and degradation kinetics of all NRF2 forms in

complex with KEAP1, is obtained.

Conclusions

Robustly inducing antioxidant and detoxification genes to cope with cellular stress imposed by

oxidative and electrophilic chemicals requires timely and sufficient NRF2 accumulation and

translocation into the nucleus. KEAP1 plays a dual role in repressing NRF2 – promoting its

degradation to keep its total abundance low and sequestering to keep its free abundance low.

The floodgate hypothesis captures some of the dual actions of KEAP1 (Iso et al. 2016, Suzuki

and Yamamoto 2017). Our modeling revealed here that the quantitative aspect of protein

stabilization of NRF2 and nuclear translocation can be better understood as a water tank model

that overflows due to drain closure (Fig. 9A), which we believe is an improvement over the

floodgate analogy. Here, the water is poured into the large water tank at a constant rate, just as

NRF2 is produced in the cytosol. Since the drain is open, most of the water will leave the tank

with a small amount remaining and leaking to the small tank (nucleus). These events are like

NRF2 being actively degraded by KEAP1 and cytosolic and nuclear NRF2 levels are low. If a

stopper is partially put in place, the water will drain slowly, and the water level in the large tank

will rise, however it is still being held by the large tank without much going into the small tank.

This is like under mild stress, KEAP1-dependent NRF2 degradation is partially stopped, total NRF2 will increase but because of the sequestration by KEAP1, it still remains largely in the cytosol. Therefore, the height of the large water tank here is equivalent to the total cytosolic KEAP1 dimer. When the stopper is further pushed in to completely block the drain, the water level will rise and eventually overflow the large tank and flood the small tank. This is like under severe oxidative stress, KEAP1-mediated NRF2 degradation is totally shut down, NRF2 accumulates to a level exceeding cytosolic KEAP1 dimer, and free NRF2 rises sharply and translocates into the nucleus. Modification of KEAP1 cysteine by class I-V compounds is like slowing the drain without affecting the height of the large water tank, while binding of class VI compounds to both of the two subunits of KEAP1 dimer is like simultaneously slowing the drain and lowering the height of the large water tank. The differential action of this water-tank model can be captured by a reduced mathematical model of KEAP1-NRF2 interaction (Fig. 9B). The free nuclear NRF2 response to class I-V compounds is potentially more ultrasensitive than that to class VI compounds, while at lower concentrations class VI compounds may activate more nuclear NRF2 than class I-V compounds (Fig. 9C).

Quantitative understanding of NRF2 activation can have many implications. A detailed kinetic model like the one we presented here can help to explore the systems behavior of cellular oxidative stress responses. It may help to better understand cancer chemoresistance, where mutation in either NRF2 or KEAP1 can lead to constitutive NRF2 activation or a more prompt and robust activation in response to chemo-drugs. Our modeling suggests that threshold concentrations may exist for certain class I-V compounds and at suprathreshold concentrations they can readily activate NRF2, while to finely control NRF2 activation with precision, class VI compounds are preferred. The model may thus help with using a synthetic biology approach to improve current and design novel classes of NRF2 activators or inhibitors. The mechanistically-based KEAP1-NRF2 model can also help to understand the commonality of toxic

actions of environmental oxidative stressors.

ACKNOWLEDGEMENTS

This research was supported in part by the National Natural Science Foundation of China:

81830099 (J.P.), 82020108027 (J.P.) and 81602824 (S.L.); Liaoning Key Research and

Development Guidance Plan 2019JH8/10300012 (J.P.); NIEHS Superfund Research grant

P42ES04911, and NIEHS HERCULES grant P30ES019776.

1043

CONFLICT OF INTEREST

1045 The authors declare no conflict of interest.

1046

REFERENCES

Adimora, N. J., D. P. Jones and M. L. Kemp (2010). "A model of redox kinetics implicates the thiol
proteome in cellular hydrogen peroxide responses." Antioxid Redox Signal **13**(6): 731-743.

Alam, J., E. Killeen, P. Gong, R. Naquin, B. Hu, D. Stewart, J. R. Ingelfinger and K. A. Nath
(2003). "Heme activates the heme oxygenase-1 gene in renal epithelial cells by stabilizing Nrf2."
Am J Physiol Renal Physiol **284**(4): F743-752.

Altszyler, E., A. C. Ventura, A. Colman-Lerner and A. Chernomoretz (2017). "Ultrasensitivity in
signaling cascades revisited: Linking local and global ultrasensitivity estimations." PLoS One
12(6): e0180083.

Baird, L., D. Llères, S. Swift and A. T. Dinkova-Kostova (2013). "Regulatory flexibility in the
Nrf2-mediated stress response is conferred by conformational cycling of the Keap1-Nrf2 protein
complex." Proc Natl Acad Sci U S A **110**(38): 15259-15264.

Baird, L. and M. Yamamoto (2020). "The Molecular Mechanisms Regulating the KEAP1-NRF2
Pathway." Mol Cell Biol **40**(13).

Bellezza, I., I. Giambanco, A. Minelli and R. Donato (2018). "Nrf2-Keap1 signaling in oxidative
and reductive stress." Biochim Biophys Acta Mol Cell Res **1865**(5): 721-733.

Buchler, N. E. and M. Louis (2008). "Molecular titration and ultrasensitivity in regulatory
networks." J Mol Biol **384**(5): 1106-1119.

Burroughs, A. F., S. Eluhu, D. Whalen, J. S. Goodwin, A. M. Sakwe and I. J. Arinze (2018).
"PML-Nuclear Bodies Regulate the Stability of the Fusion Protein Dendra2-Nrf2 in the Nucleus."
Cellular Physiology and Biochemistry **47**(2): 800-816.

Canning, P., F. J. Sorrell and A. N. Bullock (2015). "Structural basis of Keap1 interactions with
Nrf2." Free Radic Biol Med **88**(Pt B): 101-107.

Chen, Y., D. Inoyama, A. N. Kong, L. J. Beamer and L. Hu (2011). "Kinetic analyses of
Keap1-Nrf2 interaction and determination of the minimal Nrf2 peptide sequence required for
Keap1 binding using surface plasmon resonance." Chem Biol Drug Des **78**(6): 1014-1021.

Chowdhry, S., Y. Zhang, M. McMahon, C. Sutherland, A. Cuadrado and J. D. Hayes (2013). "Nrf2
is controlled by two distinct β-TrCP recognition motifs in its Neh6 domain, one of which can be
modulated by GSK-3 activity." Oncogene **32**(32): 3765-3781.

Crinelli, R., C. Zara, L. Galluzzi, G. Buffi, C. Ceccarini, M. Smietana, M. Mari, M. Magnani and A.
Fraternale (2021). "Activation of NRF2 and ATF4 Signaling by the Pro-Glutathione Molecule
I-152, a Co-Drug of N-Acetyl-Cysteine and Cysteamine." Antioxidants (Basel) **10**(2).

Dayalan Naidu, S. and A. T. Dinkova-Kostova (2020). "KEAP1, a cysteine-based sensor and a
drug target for the prevention and treatment of chronic disease." Open Biol **10**(6): 200105.

Dhakshinamoorthy, S., A. K. Jain, D. A. Bloom and A. K. Jaiswal (2005). "Bach1 competes with
Nrf2 leading to negative regulation of the antioxidant response element (ARE)-mediated
NAD(P)H:quinone oxidoreductase 1 gene expression and induction in response to antioxidants."
J Biol Chem **280**(17): 16891-16900.

Dinkova-Kostova, A. T., W. D. Holtzclaw, R. N. Cole, K. Itoh, N. Wakabayashi, Y. Katoh, M.
Yamamoto and P. Talalay (2002). "Direct evidence that sulfhydryl groups of Keap1 are the
sensors regulating induction of phase 2 enzymes that protect against carcinogens and oxidants."
Proc Natl Acad Sci U S A **99**(18): 11908-11913.

Eggler, A. L., G. Liu, J. M. Pezzuto, R. B. van Breemen and A. D. Mesecar (2005). "Modifying
specific cysteines of the electrophile-sensing human Keap1 protein is insufficient to disrupt
binding to the Nrf2 domain Neh2." Proc Natl Acad Sci U S A **102**(29): 10070-10075.

Ferrell, J. E., Jr. and S. H. Ha (2014). "Ultrasensitivity part I: Michaelian responses and
zero-order ultrasensitivity." Trends Biochem Sci **39**(10): 496-503.

Ferrell, J. E., Jr. and S. H. Ha (2014). "Ultrasensitivity part II: multisite phosphorylation,
stoichiometric inhibitors, and positive feedback." Trends Biochem Sci **39**(11): 556-569.

Fukutomi, T., K. Takagi, T. Mizushima, N. Ohuchi and M. Yamamoto (2014). "Kinetic,
thermodynamic, and structural characterizations of the association between Nrf2-DLGex degron
and Keap1." Mol Cell Biol **34**(5): 832-846.

Goldbeter, A. and D. E. Koshland, Jr. (1982). "Sensitivity amplification in biochemical systems."
Q Rev Biophys **15**(3): 555-591.

Hamon, J., P. Jennings and F. Y. Bois (2014). "Systems biology modeling of omics data: effect of
cyclosporine a on the Nrf2 pathway in human renal cells." BMC Syst Biol **8**: 76.

Hancock, R., H. C. Bertrand, T. Tsujita, S. Naz, A. El-Bakry, J. Laoruchupong, J. D. Hayes and G.
Wells (2012). "Peptide inhibitors of the Keap1-Nrf2 protein-protein interaction." Free Radic Biol
Med **52**(2): 444-451.

Hayes, J. D., S. Chowdhry, A. T. Dinkova-Kostova and C. Sutherland (2015). "Dual regulation of
transcription factor Nrf2 by Keap1 and by the combined actions of β-TrCP and GSK-3." Biochem
Soc Trans **43**(4): 611-620.

He, X., M. G. Chen, G. X. Lin and Q. Ma (2006). "Arsenic induces NAD(P)H-quinone
oxidoreductase I by disrupting the Nrf2 x Keap1 x Cul3 complex and recruiting Nrf2 x Maf to the
antioxidant response element enhancer." J Biol Chem **281**(33): 23620-23631.

Horie, Y., T. Suzuki, J. Inoue, T. Iso, G. Wells, T. W. Moore, T. Mizushima, A. T. Dinkova-Kostova, T. Kasai, T. Kamei, S. Koshiba and M. Yamamoto (2021). "Molecular basis for the disruption of Keap1–Nrf2 interaction via Hinge & Latch mechanism." Communications Biology **4**(1): 576.

Ichimura, Y., S. Waguri, Y. S. Sou, S. Kageyama, J. Hasegawa, R. Ishimura, T. Saito, Y. Yang, T. Kouno, T. Fukutomi, T. Hoshii, A. Hirao, K. Takagi, T. Mizushima, H. Motohashi, M. S. Lee, T. Yoshimori, K. Tanaka, M. Yamamoto and M. Komatsu (2013). "Phosphorylation of p62 activates the Keap1-Nrf2 pathway during selective autophagy." Mol Cell **51**(5): 618-631.

Igarashi, K., H. Hoshino, A. Muto, N. Suwabe, S. Nishikawa, H. Nakauchi and M. Yamamoto (1998). "Multivalent DNA binding complex generated by small Maf and Bach1 as a possible biochemical basis for beta-globin locus control region complex." J Biol Chem **273**(19): 11783-11790.

Iso, T., T. Suzuki, L. Baird and M. Yamamoto (2016). "Absolute Amounts and Status of the Nrf2-Keap1-Cul3 Complex within Cells." Mol Cell Biol **36**(24): 3100-3112.

Itoh, K., J. Mimura and M. Yamamoto (2010). "Discovery of the negative regulator of Nrf2, Keap1: a historical overview." Antioxid Redox Signal **13**(11): 1665-1678.

Itoh, K., N. Wakabayashi, Y. Katoh, T. Ishii, K. Igarashi, J. D. Engel and M. Yamamoto (1999). "Keap1 represses nuclear activation of antioxidant responsive elements by Nrf2 through binding to the amino-terminal Neh2 domain." Genes Dev **13**(1): 76-86.

Itoh, K., N. Wakabayashi, Y. Katoh, T. Ishii, T. O'Connor and M. Yamamoto (2003). "Keap1 regulates both cytoplasmic-nuclear shuttling and degradation of Nrf2 in response to electrophiles." Genes Cells **8**(4): 379-391.

Jiang, T., B. Harder, M. Rojo de la Vega, P. K. Wong, E. Chapman and D. D. Zhang (2015). "p62 links autophagy and Nrf2 signaling." Free Radic Biol Med **88**(Pt B): 199-204.

Jiang, Z. Y., M. C. Lu, L. L. Xu, T. T. Yang, M. Y. Xi, X. L. Xu, X. K. Guo, X. J. Zhang, Q. D. You and H. P. Sun (2014). "Discovery of potent Keap1-Nrf2 protein-protein interaction inhibitor based on molecular binding determinants analysis." J Med Chem **57**(6): 2736-2745.

Katoh, Y., K. Iida, M. I. Kang, A. Kobayashi, M. Mizukami, K. I. Tong, M. McMahon, J. D. Hayes, K. Itoh and M. Yamamoto (2005). "Evolutionary conserved N-terminal domain of Nrf2 is essential for the Keap1-mediated degradation of the protein by proteasome." Arch Biochem Biophys **433**(2): 342-350.

Katsuoka, F., H. Motohashi, J. D. Engel and M. Yamamoto (2005). "Nrf2 transcriptionally activates the mafG gene through an antioxidant response element." J Biol Chem **280**(6): 4483-4490.

Katsuoka, F., H. Motohashi, T. Ishii, H. Aburatani, J. D. Engel and M. Yamamoto (2005). "Genetic evidence that small maf proteins are essential for the activation of antioxidant response element-dependent genes." Mol Cell Biol **25**(18): 8044-8051.

Katsuragi, Y., Y. Ichimura and M. Komatsu (2016). "Regulation of the Keap1–Nrf2 pathway by p62/SQSTM1." Current Opinion in Toxicology **1**: 54-61.

Khalil, H. S., A. Goltsov, S. P. Langdon, D. J. Harrison, J. Bown and Y. Deeni (2015). "Quantitative analysis of NRF2 pathway reveals key elements of the regulatory circuits underlying antioxidant response and proliferation of ovarian cancer cells." J Biotechnol **202**: 12-30.

Kobayashi, A., M. I. Kang, H. Okawa, M. Ohtsuji, Y. Zenke, T. Chiba, K. Igarashi and M. Yamamoto (2004). "Oxidative stress sensor Keap1 functions as an adaptor for Cul3-based E3 ligase to regulate proteasomal degradation of Nrf2." Mol Cell Biol **24**(16): 7130-7139.

Kobayashi, A., M. I. Kang, Y. Watai, K. I. Tong, T. Shibata, K. Uchida and M. Yamamoto (2006). "Oxidative and electrophilic stresses activate Nrf2 through inhibition of ubiquitination activity of Keap1." Mol Cell Biol **26**(1): 221-229.

Kobayashi, M., L. Li, N. Iwamoto, Y. Nakajima-Takagi, H. Kaneko, Y. Nakayama, M. Eguchi, Y. Wada, Y. Kumagai and M. Yamamoto (2009). "The antioxidant defense system Keap1-Nrf2 comprises a multiple sensing mechanism for responding to a wide range of chemical compounds." Mol Cell Biol **29**(2): 493-502.

Kobayashi, M. and M. Yamamoto (2005). "Molecular mechanisms activating the Nrf2-Keap1 pathway of antioxidant gene regulation." Antioxid Redox Signal **7**(3-4): 385-394.

Kolodkin, A. N., R. P. Sharma, A. M. Colangelo, A. Ignatenko, F. Martorana, D. Jennen, J. J. Briedé, N. Brady, M. Barberis, T. Mondeel, M. Papa, V. Kumar, B. Peters, A. Skupin, L. Alberghina, R. Balling and H. V. Westerhoff (2020). "ROS networks: designs, aging, Parkinson's disease and precision therapies." NPJ Syst Biol Appl **6**(1): 34.

Komatsu, M., H. Kurokawa, S. Waguri, K. Taguchi, A. Kobayashi, Y. Ichimura, Y. S. Sou, I. Ueno, A. Sakamoto, K. I. Tong, M. Kim, Y. Nishito, S. Iemura, T. Natsume, T. Ueno, E. Kominami, H. Motohashi, K. Tanaka and M. Yamamoto (2010). "The selective autophagy substrate p62 activates the stress responsive transcription factor Nrf2 through inactivation of Keap1." Nat Cell Biol **12**(3): 213-223.

Kwak, M. K., K. Itoh, M. Yamamoto and T. W. Kensler (2002). "Enhanced expression of the transcription factor Nrf2 by cancer chemopreventive agents: role of antioxidant response element-like sequences in the nrf2 promoter." Mol Cell Biol **22**(9): 2883-2892.

Lau, A., X. J. Wang, F. Zhao, N. F. Villeneuve, T. Wu, T. Jiang, Z. Sun, E. White and D. D. Zhang (2010). "A noncanonical mechanism of Nrf2 activation by autophagy deficiency: direct interaction between Keap1 and p62." Mol Cell Biol **30**(13): 3275-3285.

Lazzara, P. R., B. P. David, A. Ankireddy, B. G. Richardson, K. Dye, K. M. Ratia, S. P. Reddy and T. W. Moore (2020). "Isoquinoline Kelch-like ECH-Associated Protein 1-Nuclear Factor (Erythroid-Derived 2)-like 2 (KEAP1-NRF2) Inhibitors with High Metabolic Stability." J Med Chem **63**(12): 6547-6560.

Leclerc, E., J. Hamon, A. Legendre and F. Y. Bois (2014). "Integration of pharmacokinetic and NRF2 system biology models to describe reactive oxygen species production and subsequent glutathione depletion in liver microfluidic biochips after flutamide exposure." Toxicol In Vitro **28**(7): 1230-1241.

Lee, S. and L. Hu (2020). "Nrf2 activation through the inhibition of Keap1-Nrf2 protein-protein interaction." Med Chem Res **29**(5): 846-867.

Legewie, S., N. Bluthgen and H. Herzel (2005). "Quantitative analysis of ultrasensitive responses." FEBS J **272**(16): 4071-4079.

Li, W., S. W. Yu and A. N. Kong (2006). "Nrf2 possesses a redox-sensitive nuclear exporting signal in the Neh5 transactivation domain." J Biol Chem **281**(37): 27251-27263.

Li, X., D. Zhang, M. Hannink and L. J. Beamer (2004). "Crystal structure of the Kelch domain of human Keap1." J Biol Chem **279**(52): 54750-54758.

Lo, S. C., X. Li, M. T. Henzl, L. J. Beamer and M. Hannink (2006). "Structure of the Keap1:Nrf2 interface provides mechanistic insight into Nrf2 signaling." Embo j **25**(15): 3605-3617.

Malhotra, D., E. Portales-Casamar, A. Singh, S. Srivastava, D. Arenillas, C. Happel, C. Shyr, N. Wakabayashi, T. W. Kensler, W. W. Wasserman and S. Biswal (2010). "Global mapping of binding sites for Nrf2 identifies novel targets in cell survival response through ChIP-Seq profiling and network analysis." Nucleic Acids Res **38**(17): 5718-5734.

McMahon, M., N. Thomas, K. Itoh, M. Yamamoto and J. D. Hayes (2004). "Redox-regulated turnover of Nrf2 is determined by at least two separate protein domains, the redox-sensitive Neh2 degron and the redox-insensitive Neh6 degron." J Biol Chem **279**(30): 31556-31567.

McMahon, M., N. Thomas, K. Itoh, M. Yamamoto and J. D. Hayes (2006). "Dimerization of substrate adaptors can facilitate cullin-mediated ubiquitylation of proteins by a "tethering" mechanism: a two-site interaction model for the Nrf2-Keap1 complex." J Biol Chem **281**(34): 24756-24768.

Moi, P., K. Chan, I. Asunis, A. Cao and Y. W. Kan (1994). "Isolation of NF-E2-related factor 2

(Nrf2), a NF-E2-like basic leucine zipper transcriptional activator that binds to the tandem NF-E2/AP1 repeat of the beta-globin locus control region." Proc Natl Acad Sci U S A **91**(21): 9926-9930.

Nguyen, T., P. J. Sherratt and C. B. Pickett (2003). "Regulatory mechanisms controlling gene expression mediated by the antioxidant response element." Annu Rev Pharmacol Toxicol **43**: 233-260.

Ogura, T., K. I. Tong, K. Mio, Y. Maruyama, H. Kurokawa, C. Sato and M. Yamamoto (2010). "Keap1 is a forked-stem dimer structure with two large spheres enclosing the intervening, double glycine repeat, and C-terminal domains." Proc Natl Acad Sci U S A **107**(7): 2842-2847.

Paunkov, A., D. V. Chartoumpekis, P. G. Ziros and G. P. Sykiotis (2019). "A Bibliometric Review of the Keap1/Nrf2 Pathway and its Related Antioxidant Compounds." Antioxidants (Basel) **8**(9).

Pi, J., W. Qu, J. M. Reece, Y. Kumagai and M. P. Waalkes (2003). "Transcription factor Nrf2 activation by inorganic arsenic in cultured keratinocytes: involvement of hydrogen peroxide." Exp Cell Res **290**(2): 234-245.

Pi, J., Q. Zhang, C. G. Woods, V. Wong, S. Collins and M. E. Andersen (2008). "Activation of Nrf2-mediated oxidative stress response in macrophages by hypochlorous acid." Toxicol Appl Pharmacol **226**(3): 236-243.

Rada, P., A. I. Rojo, S. Chowdhry, M. McMahon, J. D. Hayes and A. Cuadrado (2011). "SCF/{beta}-TrCP promotes glycogen synthase kinase 3-dependent degradation of the Nrf2 transcription factor in a Keap1-independent manner." Mol Cell Biol **31**(6): 1121-1133.

Sekhar, K. R., G. Rachakonda and M. L. Freeman (2010). "Cysteine-based regulation of the CUL3 adaptor protein Keap1." Toxicol Appl Pharmacol **244**(1): 21-26.

Selvaggio, G., P. Coelho and A. Salvador (2018). "Mapping the phenotypic repertoire of the cytoplasmic 2-Cys peroxiredoxin - Thioredoxin system. 1. Understanding commonalities and differences among cell types." Redox Biol **15**: 297-315.

Stewart, D., E. Killeen, R. Naquin, S. Alam and J. Alam (2003). "Degradation of transcription factor Nrf2 via the ubiquitin-proteasome pathway and stabilization by cadmium." J Biol Chem **278**(4): 2396-2402.

Sun, Z., S. Zhang, J. Y. Chan and D. D. Zhang (2007). "Keap1 controls postinduction repression of the Nrf2-mediated antioxidant response by escorting nuclear export of Nrf2." Mol Cell Biol **27**(18): 6334-6349.

Suzuki, H., S. Tashiro, J. Sun, H. Doi, S. Satomi and K. Igarashi (2003). "Cadmium induces nuclear export of Bach1, a transcriptional repressor of heme oxygenase-1 gene." J Biol Chem

278(49): 49246-49253.

Suzuki, T. and M. Yamamoto (2017). "Stress-sensing mechanisms and the physiological roles of the Keap1-Nrf2 system during cellular stress." J Biol Chem **292**(41): 16817-16824.

Tonelli, C., I. I. C. Chio and D. A. Tuveson (2018). "Transcriptional Regulation by Nrf2." Antioxid Redox Signal **29**(17): 1727-1745.

Tong, K. I., Y. Katoh, H. Kusunoki, K. Itoh, T. Tanaka and M. Yamamoto (2006). "Keap1 recruits Neh2 through binding to ETGE and DLG motifs: characterization of the two-site molecular recognition model." Mol Cell Biol **26**(8): 2887-2900.

Tong, K. I., A. Kobayashi, F. Katsuoka and M. Yamamoto (2006). "Two-site substrate recognition model for the Keap1-Nrf2 system: a hinge and latch mechanism." Biol Chem **387**(10-11): 1311-1320.

Tong, K. I., B. Padmanabhan, A. Kobayashi, C. Shang, Y. Hirotsu, S. Yokoyama and M. Yamamoto (2007). "Different electrostatic potentials define ETGE and DLG motifs as hinge and latch in oxidative stress response." Mol Cell Biol **27**(21): 7511-7521.

Watai, Y., A. Kobayashi, H. Nagase, M. Mizukami, J. McEvoy, J. D. Singer, K. Itoh and M. Yamamoto (2007). "Subcellular localization and cytoplasmic complex status of endogenous Keap1." Genes Cells **12**(10): 1163-1178.

Xue, M., H. Momiji, N. Rabbani, G. Barker, T. Bretschneider, A. Shmygol, D. A. Rand and P. J. Thornalley (2015). "Frequency Modulated Translocational Oscillations of Nrf2 Mediate the Antioxidant Response Element Cytoprotective Transcriptional Response." Antioxid Redox Signal **23**(7): 613-629.

Yamamoto, M., T. W. Kensler and H. Motohashi (2018). "The KEAP1-NRF2 System: a Thiol-Based Sensor-Effector Apparatus for Maintaining Redox Homeostasis." Physiol Rev **98**(3): 1169-1203.

Yamamoto, T., T. Suzuki, A. Kobayashi, J. Wakabayashi, J. Maher, H. Motohashi and M. Yamamoto (2008). "Physiological significance of reactive cysteine residues of Keap1 in determining Nrf2 activity." Mol Cell Biol **28**(8): 2758-2770.

Yasuda, D., A. Yuasa, R. Obata, M. Nakajima, K. Takahashi, T. Ohe, Y. Ichimura, M. Komatsu, M. Yamamoto, R. Imamura, H. Kojima, T. Okabe, T. Nagano and T. Mashino (2017). "Discovery of benzo[g]indoles as a novel class of non-covalent Keap1-Nrf2 protein-protein interaction inhibitor." Bioorg Med Chem Lett **27**(22): 5006-5009.

Zhang, Q. and M. E. Andersen (2007). "Dose response relationship in anti-stress gene regulatory networks." PLoS Comput Biol **3**(3): e24.

Zhang, Q., S. Bhattacharya and M. E. Andersen (2013). "Ultrasensitive response motifs: basic
amplifiers in molecular signalling networks." Open Biol **3**(4): 130031.

Zhang, Q., S. Bhattacharya, J. Pi, R. A. Clewell, P. L. Carmichael and M. E. Andersen (2015).
"Adaptive Posttranslational Control in Cellular Stress Response Pathways and Its Relationship to
Toxicity Testing and Safety Assessment." Toxicol Sci **147**(2): 302-316.

Zhang, Q., J. Pi, C. G. Woods and M. E. Andersen (2009). "Phase I to II cross-induction of
xenobiotic metabolizing enzymes: a feedforward control mechanism for potential hormetic
responses." Toxicol Appl Pharmacol **237**(3): 345-356.

Zhang, Q., J. Pi, C. G. Woods and M. E. Andersen (2010). "A systems biology perspective on
Nrf2-mediated antioxidant response." Toxicol Appl Pharmacol **244**(1): 84-97.

Zipper, L. M. and R. T. Mulcahy (2002). "The Keap1 BTB/POZ dimerization function is required to
sequester Nrf2 in cytoplasm." J Biol Chem **277**(39): 36544-36552.
1289

1290
Tables

Table 1. KEAP1-NRF2 Model Features

Model #	Model structure	Cycle mode of operation	Two-step ETGE binding	NRF2 nuclear translocation	Class of NRF2 activator
1	Fig. 1				I-V
2	Fig. 1	X			I-V
3a	Fig. 2A	X	X		I-V
3b	Fig. 6A	X	X		VI
4a	Fig. 7A	X	X	X	I-V
4b	Fig. 8A	X	X	X	VI

Note: X denotes that a model has the corresponding feature. The structure of each model is illustrated in the Figures indicated.

1291

                                   Figure Legend

1293   **Figure 1. Structure of KEAP1-NRF2 Models 1 and 2.** The models feature one-step ETGE

1294   binding and interaction with class I-V activator. Short arrow bars next to a parameter symbol

1295   denote the direction of the reversible binding described by the parameter. Φ denotes degradation.

1296   These denotations apply to all other model structures.

1297

Figure 2. Structure, and dynamical and steady-state behaviors of Model 3a. (A) Structure of

Model 3a featuring two-step ETGE binding and interaction with class I-V activator. **(B)** Dynamical

changes of basal $NRF2_{free}$, $KEAP1\_NRF2_{open1}$, $KEAP1\_NRF2_{open2}$, $KEAP1\_NRF2_{closed}$, and

$NRF2_{tot}$ in response to termination of NRF2 synthesis (by setting k_0=0) starting at 0 min with k_6 at

default value. **(C)** Dynamical changes of various NRF2 species in response to stabilization of

NRF2 in $KEAP1\_NRF2_{closed}$ by setting k_6=1.252E-4 starting at 0 min and in response to

termination of NRF2 synthesis (by setting k_0=0) starting at 500 min. For simulations in (B) and

(C), $CLASS_{I-V}$ level is kept at zero. Dynamical changes of **(D)** $NRF2_{tot}$, **(E)** $NRF2_{free}$, and **(F)**

$KEAP1_{free\_tot}$ in response different levels of $CLASS_{I-V}$ with k'_6 at default value. **(G)** Steady-state

dose-response curves of various NRF2 species and $KEAP1_{free\_tot}$ on double-log scale with k'_6 at

default value. Shown are n_H and LRC_{max} for $NRF2_{free}$; n_H and LRC_{max} for $NRF2_{total}$ are 1.22 and

0.42 respectively (not shown). **(H-I)** Flux analyses for conditions when NRF2 in

$KEAP1\_NRF2_{closed}$ is stabilized by setting k_6 to 30% (H) and 10% (I) of default value.

1311

1312   **Figure 3. Effects of KEAP1-NRF2 binding parameters on NRF2 responses in Model 3a.**

1313   Effects of varying **(A-B)** k_1 (k'_1), **(C-D)** $k_{1.1}$ ($k'_{1.1}$), **(E-F)** $k_{2.1}$ ($k'_{2.1}$), and **(G-H)** k_3 (k'_3) on steady-state

1314   dose-response curves of $NRF2_{tot}$ (left panels) and $NRF2_{free}$ (right panels). Note k_i=k'_i for i = 1, 1.1,

1315   2.1, or 3; x0.1, x1, and x10 denote 0.1, 1, and 10 times default values.

1316

Figure 4. Effects of varying parameter k'_3 alone on NRF2 responses in Model 3a to test

hinge-latch hypothesis - with k'_6 at default value. Dynamical changes of $KEAP1\_NRF2_{open\_tot}$

and $KEAP1\_NRF2_{closed\_tot}$ in response to a high level of $CLASS_{I\text{-}V}$ at 1000 nM starting at 0 min

and in response to termination of NRF2 synthesis (by setting k_0=0) starting at 500 min, when k'_3

is **(A)** 0.1 and **(C)** 10 times of default value. Steady-state dose-response curves of

$KEAP1\_NRF2_{open\_tot}$ and $KEAP1\_NRF2_{closed\_tot}$ when k'_3 is **(B)** 0.1 and **(D)** 10 times of default

value. **(E-F)** Effects of varying k'_3 on steady-state dose-response curves of $NRF2_{tot}$ and $NRF2_{free}$

respectively.

1325

1326 **Figure 5. Effects of total KEAP1 abundance on NRF2 responses in Model 3a.** Steady-state

1327 dose-response curves of **(A)** $NRF2_{tot}$ and **(B)** $NRF2_{free}$ under different values of total KEAP1

1328 abundance relative to default value.

1329

Figure 6. Structure, and dynamical and steady-state behaviors of Model 3b. (A) Structure of

Model 3b featuring two-step ETGE binding and interaction with class VI activator. Dynamical

changes of **(B)** $NRF2_{tot}$ and **(C)** $NRF2_{free}$ in response different levels of $CLASS_{VI}$. **(D)**

Steady-state dose-response curves of $NRF2_{tot}$ and $NRF2_{free}$. Shown are n_H and LRC_{max} for

$NRF2_{free}$; n_H and LRC_{max} for $NRF2_{total}$ are 1.35 and 0.35 respectively (not shown). **(E)**

Steady-state dose-response curves of $KEAP1\_NRF2_{open\_tot}$, $KEAP1\_NRF2_{closed\_tot}$,

$ClassVI_1\_KEAP1$ (Class VI activator-KEAP1 complex containing one activator molecule) and

$ClassVI_2\_KEAP1$ (containing two activator molecules). **(F-G)** Steady-state oxidant-response

curves of $NRF2_{tot}$, $NRF2_{free}$, $KEAP1\_NRF2_{open\_tot}$, $KEAP1\_NRF2_{closed\_tot}$, and $ClassVI_1\_KEAP1$

under condition when only one class VI activator molecule is allowed to bind to KEAP1 by setting

$k'_7 = k'_8 = 0$.

1341

Figure 7. Structure, and dynamical and steady-state behaviors of Model 4a. (A) Structure of

Model 4a featuring two-step ETGE binding, nuclear NRF2 translocation, and interaction with

class I-V activator. **(B)** Dynamical changes of basal $NRF2_{tot\_cell}$, $NRF2_{tot\_nucleus}$, $NRF2_{tot\_cytosol}$,

$NRF2_{free\_nucleus}$, and $NRF2_{free\_cytosol}$ in response to termination of NRF2 synthesis (by setting $k_0=0$)

starting at 0 min. **(C)** $NRF2_{tot\_cell}$, $NRF2_{tot\_nucleus}$, and $NRF2_{tot\_cytosol}$ in (B) shown in log Y scale. **(D)**

Dynamical changes of various NRF2 species previously induced by a high level of $CLASS_{I-V}$ at

1000 nM in response to termination of NRF2 synthesis (by setting $k_0=0$) starting at 0 min. **E)**

Steady-state dose-response curves of various NRF2 species. n_H and LRC_{max} of $NRF2_{free\_nucleus}$

curve are indicated.

1351

Figure 8. Structure, and dynamical and steady-state behaviors of Model 4b. (A) Structure of

Model 4b featuring two-step ETGE binding, nuclear NRF2 translocation, and interaction with

class VI activator. Dynamical changes of **(B)** $NRF2_{free\_cytosol}$, **(C)** $NRF2_{tot\_cytosol}$, and **(D)**

$NRF2_{free\_nucleus}$ in response to different levels of $CLASS_{VI}$. **(E)** Dynamical changes of various

NRF2 species previously induced by a high level of $CLASS_{VI}$ at 1000 nM in response to

termination of NRF2 synthesis (by setting $k_0=0$) starting at 0 min. **(F-G)** Steady-state

dose-response curves of various NRF2 species and KEAP1 species respectively.

1359

Figure 9. Water-tank analogy and reduced KEAP1-NRF2 mathematical model. (A)

Schematic illustration of the water-tank analogy for KEAP1-dependent NRF2 degradation,

sequestration, and nuclear translocation. Large tank: cytosol, small tank: nucleus, height of large

tank: total cytosolic KEAP1 abundance, water: NRF2, tap: NRF2 production, drain:

KEAP1-mediated NRF2 degradation, stopper: oxidant or NRF2 inducer. To reduce clutter for

clarity, KEAP1-independent NRF2 degradation and nuclear NRF2 degradation are not shown. **(B)**

Reduced KEAP1-NRF2 mathematical model for NRF2 activation by class I-V and class VI

activators. **(C)** Predicted differential free nuclear NRF2 dose-response for class I-V and class VI

activators.

1369

Figure 1

Figure 2

H

k_6 = 5.325E-4 (0.3 × default)

I

k_6 = 1.775E-4 (0.1 × default)

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

A

Oxidative stressor level

B

C