Extreme Heterogeneity in Genomic Differentiation between Phenotypically Divergent Songbirds: A Test of Mitonuclear Co-introgression

Ellen Nikelski[1, *], Alexander S. Rubtsov[2], and Darren Irwin[1]

[1] Department of Zoology, and Biodiversity Research Centre, 6270 University Blvd., University of British Columbia, Vancouver, BC, Canada

[2] State Darwin Museum, Moscow, Russia

[*] Present address: Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

Corresponding Author:

Ellen Nikelski, Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada. Email: ellen.nikelski@mail.utoronto.ca

Keywords:

mitonuclear co-introgression, mtDNA, mitonuclear gene, genetic differentiation, chromosome Z, Aves

Abstract:

Comparisons of genomic variation among closely related species often show more differentiation

in mitochondrial DNA (mtDNA) and sex chromosomes than in autosomes, a pattern expected

due to the differing effective population sizes and evolutionary dynamics of these genomic

components. Yet, introgression can cause species pairs to deviate dramatically from general

differentiation trends. The yellowhammer (*Emberiza citrinella*) and the pine bunting

(*E. leucocephalos*) are hybridizing avian sister species that differ greatly in appearance and

moderately in nuclear DNA, but that show no mtDNA differentiation. This mitonuclear

discordance is best explained by adaptive mtDNA introgression—a process that can select for

co-introgression at nuclear genes with mitochondrial functions (mitonuclear genes). To better

understand the extent of mitonuclear discordance and characterize nuclear differentiation

patterns in this system, we investigated genome-wide differentiation between allopatric

yellowhammers and pine buntings and compared it to what was seen previously in mtDNA. We

found significant nuclear differentiation that was highly heterogeneous across the genome, with a

particularly wide differentiation peak on the sex chromosome Z. We further tested for

preferential introgression of mitonuclear genes and found statistical support for this process in

yellowhammers. A role for mitonuclear coevolution in this system is supported by a stronger

signal of co-introgression in genes coding for subunits of the mitoribosome and electron

transport chain complexes. Altogether, our study emphasizes the extreme variation seen in

differentiation across genomic components and study systems as well as highlights the

ramifications of this variation in species evolution.

Introduction

Evolution in eukaryotes is shaped by changes in multiple genomic components that differ in their modes of inheritance: mitochondrial DNA (mtDNA) is usually inherited through the matrilineal line, autosomes are inherited through both parental lines and sex chromosomes are inherited differentially depending on the sex of both parent and offspring (Avise, 2000). During speciation, populations of a single species diverge genetically, often in isolation, with the strength and pattern of genetic differentiation varying across the different genomic components (reviewed in Coyne & Orr, 2004; reviewed in Price, 2008). This variation arises as a result of differences in each component's rate of evolution as well as the degree to which each component contributes towards reproductive isolation and is resistant to gene flow between populations in secondary contact. Most commonly, speciating taxa will show clear differentiation in mtDNA (eg. Hebert et al. 2004; Kerr et al. 2007), moderate differentiation in sex chromosomes (eg. Thornton & Long, 2002; Borge et al. 2005; Lu & Wu, 2005; Harr, 2006; Ruegg et al. 2014; Sackton et al. 2014), and comparatively modest differentiation across autosomes (Harr, 2006; Nadeau et al. 2012; Irwin et al. 2018).

In most cases, mtDNA shows strong differentiation between speciating populations (e.g. Hebert et al. 2004; Kerr et al. 2007). These patterns are partially driven by the mitochondrial genome's uniparental inheritance and haploid nature which decrease its effective population size to ¼ that of autosomal DNA (Moore, 1995). Combined with a relatively high mutation rate (Lynch et al. 2006), this low effective population size leads to the rapid fixation of mitochondrial mutations via genetic drift and strong mtDNA differentiation between taxa. Additionally, a lack of recombination across the mitochondrial genome can further contribute to mtDNA divergence through rampant genetic hitchhiking (Hill, 2020). Here, positive selection for an adaptive

mtDNA mutation causes the fixation of genetic variants across the mitochondrial genome

resulting in strong differentiation between speciating taxa.

Sex chromosomes—specifically the Z (Borge et al. 2005; Ruegg et al. 2014; Sackton et

al. 2014) and X chromosomes (Thorton & Long, 2002; Lu & Wu, 2005; Harr, 2006)—show

moderate levels of genetic differentiation between speciating populations, but often much greater

genetic differentiation than autosomes. This trend is driven by two important characteristics of

Z/X chromosomes. First, on Z/X chromosomes, beneficial recessive mutations are immediately

exposed to selection in the heterogametic sex allowing these advantageous variants to fix more

rapidly than they would on autosomes—a phenomenon known as the "faster Z/X effect" (Meisel

& Connallon, 2013; Irwin, 2018). Second, because Z/X chromosomes are inherited as either one

or two copies depending on the sex, these chromosomes have a lower effective population size

than autosomes (Mank et al. 2010; Irwin, 2018). This lower effective population size allows for

the fixation of a greater number of neutral and slightly deleterious mutations due to less effective

purifying selection and a larger role of genetic drift. Working in tandem, these two

characteristics drive genetic divergence of Z/X chromosomes between speciating populations.

Across autosomes, genetic differentiation between speciating taxa is relatively modest

compared to mtDNA and sex chromosomes and, interestingly, it is often characterized by

"islands of differentiation"—peaks of high relative differentiation that appear within a

background of low relative differentiation (Harr, 2006; Nadeau et al. 2012; Hejase et al. 2020).

Explanations for these "islands" often invoke reduced gene flow between speciating taxa during

secondary contact (Wu, 2001) and/or repeated bouts of selection prior to and following

secondary contact (Cruickshank and Hahn, 2014; Irwin et al. 2018). In the former scenario,

differentiation peaks are hypothesized to house the loci responsible for reproductive barriers

between taxa making them resistant to the homogenizing influence of gene flow (Wu, 2001). In

the latter scenario, differentiation islands are described as genomic areas that have experienced

recurrent selection or selective sweeps (Cruickshank and Hahn, 2014; Irwin et al. 2018). These

events reduce genetic diversity first in the ancestral population and then in both daughter

populations producing relative peaks in genetic differentiation between taxa.

Despite the general differentiation trends discussed above, an increasing number of

studies have reported differentiation patterns that greatly diverge from what is normally seen

between speciating populations (Irwin et al. 2009; Yannic et al. 2010; Bryson et al. 2012). In

these studies, genomic components that are normally highly divergent between taxa show

remarkably low differentiation when compared to other genomic components and observable

phenotypes. In particular, cases of "mitonuclear discordance", where mtDNA shows

dramatically low differentiation compared to nuclear DNA, have been identified across lineages

(Toews & Brelsford, 2012). One hypothesis that may explain mitonuclear discordance is

introgression of mitochondrial haplotypes from one population into another following secondary

contact and hybridization. Introgression of mtDNA can occur neutrally due to various processes

such as sex-biased dispersal or hybrid zone movement (Toews & Brelsford, 2012); however, the

extreme degree of discordance observed in certain systems (e.g. Alves et al. 2008; Irwin et al.

2009), suggests that this process may also be driven by selection and occur adaptively by

providing a fitness advantage to the receiving population.

Introgression of foreign mtDNA is thought to provide fitness advantages through two

major avenues. First, because variation in mtDNA has been associated with variation in

mitochondrial efficiency under different abiotic conditions (e.g. Ballard et el. 2007), mtDNA

introgression may allow the receiving population to better adapt and survive within a changing or

novel environment (Hulsey et al. 2016; Sloan et al. 2017; Hill, 2019). For example, mtDNA

introgression has been tentatively associated with thermal adaptation in populations of rabbits

(Alves et al. 2008) and cichlids (Hulsey et al. 2016). The second way that mtDNA introgression

can provide a fitness advantage to a receiving population is by replacing a mitochondrial genome

with a high mutational load (Sloan et al. 2017; Hill, 2019). As described earlier, the low effective

population size (Moore, 1995) and high mutation rate (Lynch et al. 2006) of mtDNA can result

in rapid fixation of mitochondrial mutations, including deleterious mutations. This tendency

combined with genetic hitchhiking of deleterious mutations (Hill, 2020), can lead to a high

mtDNA mutational load which may decrease mitochondrial efficiency. Through introgression of

a foreign mitochondrial haplotype with a lower mutational load, the receiving population may be

able to regain greater mitochondrial function which translates into higher fitness (Llopart et al.

2014; Hulsey et al. 2016; Sloan et al. 2017; Hill, 2019).

Adaptive mtDNA introgression presents a compelling hypothesis for the appearance of

extreme "mitonuclear discordance" between speciating taxa. However, this idea becomes

complicated when we consider recent work that suggests strong coevolution between the

mitochondrial and nuclear genomes (Hill, 2019). In most bilaterian animals, the mitochondrial

genome has been reduced to 37 genes. As a result of this low gene content, all mitochondrial

processes—including oxidative phosphorylation and transcription, translation and replication of

mtDNA—are reliant on more than 1000 proteins encoded by specific "mitonuclear genes" across

the nuclear genome (Calvo & Mootha, 2010; Lotz et al. 2014). Tight associations between

mitochondrial and mitonuclear proteins imply tight coevolution between the mitochondrial and

nuclear genomes due to selection for mitochondrial efficiency: changes in one create selective

pressure for compatible changes in the other (Gershoni et al. 2009; Burton & Barreto, 2012; Hill,

2019). This association also suggests that hybridization and gene flow between speciating

populations will be selected against due to recombination of coevolved mitochondrial and

mitonuclear genes exposing genetic incompatibilities (mitonuclear incompatibilities) within

hybrid individuals. In this way, mitonuclear coevolution has the potential to contribute towards

reproductive isolation between taxa and to strongly select against mtDNA introgression.

Nevertheless, if mtDNA introgression is highly adaptive such that its fitness advantages

outweigh the fitness disadvantages of mitonuclear incompatibilities, this process may still be

able to occur. In this scenario, researchers hypothesize that mtDNA introgression will select for

co-introgression of mitonuclear genes to maintain coevolved genotypes and optimum

mitochondrial function (Sloan et al. 2017; Hill, 2019). Evidence for mitonuclear co-introgression

has been found in some systems (Beck et al. 2015; Morales et al. 2018; Wang et al. 2021), but

the significance of this process in genome-wide nuclear differentiation and general speciation

trends is still up for debate.

The yellowhammer (Passeriformes: Emberizidae: *Emberiza citrinella*) and pine bunting

(*E. leucocephalos*) system (Figure 1) is one that may capture the complex interplay between

genetic differentiation and mitonuclear coevolution. Thought to have diverged in isolation during

the Pleistocene glaciations (Irwin et al. 2009), this Eurasian avian sister pair is highly divergent

in plumage and moderately divergent in song and ecology (Panov et al. 2003; Rubtsov &

Tarasov, 2017). Yet, despite their differences, yellowhammers and pine buntings hybridize

extensively in a large and seemingly expanding secondary contact zone in central and western

Siberia (Panov et al. 2003; 2007; Rubtsov, 2007; Irwin et al. 2009; Rubtsov & Tarasov, 2017).

Previous genomic work has identified mitonuclear discordance between allopatric

yellowhammers and pine buntings (Irwin et al. 2009) as they are nearly identical in mtDNA but

show moderate differentiation in nuclear AFLP (Amplified Fragment Length Polymorphism)

markers. To explain these results, Irwin et al. (2009) suggested that mtDNA introgressed

adaptively from one species into the other during a previous selective sweep, and this hypothesis

was supported by several statistical tests performed on the mtDNA haplotype network. Based on

mitonuclear theory, such mtDNA introgression could select for similar introgression at

mitonuclear genes to maintain optimum mitochondrial function (Sloan et al. 2017; Hill, 2019).

The resulting lack of mitonuclear incompatibilities between taxa due to this co-introgression

could facilitate their continued hybridization and hinder the build-up of reproductive barriers.

With this opposition between the observed strong nuclear differentiation and the potential for

mitonuclear co-introgression, the fate of the yellowhammer and pine bunting system remains

uncertain. Depending on which way the scales tip, yellowhammers and pine buntings may

continue to diverge and speciate or they may continue to hybridize and eventually collapse into

one interbreeding population.

Here, we present a largescale comparison of DNA sequence variation across the nuclear

genomes of allopatric yellowhammers and pine buntings. With this data, we address key

questions regarding genetic differentiation and mitonuclear coevolution in this system. First,

what is the degree and structure of genetic differentiation between yellowhammers and pine

buntings across the nuclear genome? Earlier AFLP analyses identified clear differentiation of

nuclear markers between yellowhammers and pine buntings (Irwin et al. 2009), but those results

were not based on actual DNA sequences and only captured a small portion of the nuclear

genome. Comparing patterns of differentiation across the nuclear genome enables better

understanding of the extent of mitonuclear discordance between taxa and of which genomic

regions or components (e.g. sex chromosomes) show particularly high differentiation. Second, is

there an over-representation of known mitonuclear genes within genomic regions that show the

strongest signals of being introgressed–a pattern consistent with mitonuclear co-introgression?

Evidence of mitonuclear co-introgression and the resulting lack of mitonuclear incompatibilities

could explain the continued, extensive hybridization seen between yellowhammers and pine

buntings and implicate this process as a force that counters divergence and the evolution of

strong reproductive barriers between groups. By answering these questions, we hope to provide

insight on the evolutionary trajectory of yellowhammers and pine buntings (i.e. whether it is one

of continued population divergence or of population merging) and also to explore how the

interplay between genetic differentiation and mitonuclear coevolution influences the speciation

process more generally.

Materials and Methods

Sampling

We included 109 blood and tissue samples in this study: 53 phenotypic yellowhammers,

42 phenotypic pine buntings, and 14 other members of Emberizidae (one *Emberiza aureola*

[yellow-breasted bunting], one *Emberiza calandra* [corn bunting], one *Emberiza cioides*

[meadow bunting], one *Emberiza hortulana* [ortolan bunting], four *Emberiza stewarti* [white-

capped bunting], and six *Emberiza cirlus* [cirl bunting]) to put variation between yellowhammers

and pine buntings into a deeper phylogenetic context (Figure 1A; Table 1; Supplementary Table

1). A total of 91 of our samples were included in the AFLP analysis of Irwin et al. (2009) while

18 samples were examined for the first time as part of the present research.

When possible, body measurements and photographs were taken of live birds or museum

skins. Yellowhammer and pine bunting males were scored phenotypically and sorted into

phenotypic classes based on the protocols presented in Panov et al. (2003) and Rubtsov &

Tarasov (2017). Briefly, each male received a score from 0-7 for three plumage traits, where 0

represents a phenotypically pure yellowhammer and 7 represents a phenotypically pure pine

bunting. These three traits were (1) the amount of chestnut plumage (vs. yellow or white) at the

brow, (2) the amount of chestnut plumage (vs. yellow or white) at the throat, and (3) background

plumage colour, meaning the strength of yellow—ranging from bright yellow to pure white—in

head and body plumage that did not show brown or black streaking. Phenotypic classes included:

pure *citrinella* (PC), almost *citrinella* (SC, for "semi-*citrinella*"), *citrinella* hybrid (CH), yellow

hybrid (YH), white hybrid (WH), *leucocephalos* hybrid (LH), almost *leucocephalos* (SL, for

"semi-*leucocephalos*") and pure *leucocephalos* (PL) (Rubtsov & Tarasov, 2017). Any SC and SL

individuals that appeared in the allopatric zones were grouped together with PC and PL

individuals respectively and treated as phenotypic yellowhammers and phenotypic pine buntings

in subsequent analyses (Figure 1B).

DNA extraction and genotyping-by-sequencing

DNA was extracted from samples using a standard phenol-chloroform method. We then

divided the DNA samples into four genotyping-by-sequencing (GBS) libraries (Elshire et al.

2011). The 109 samples included in this study were sequenced together with 226 yellowhammer,

pine bunting and hybrid DNA samples collected near and within the sympatric zone as part of a

larger project (Nikelski et al. in prep). The libraries were prepared as per the protocol described

by Alcaide et al. (2014) with modifications specified by Geraldes et al. (2019) except that we

maintained a 300-400 bp fragment size during size selection. Paired-end sequencing was

completed by Genome Québec using an Illumina HiSeq 4000 system, producing more than 1.2

billion reads, each 150 bp in length, across the four GBS libraries.

Genotyping-by-sequencing data filtering

We processed the reads following Irwin et al. (2016; 2018), as summarized here. Reads

were demultiplexed using a custom perl script designed by Baute et al. (2016). Next, reads were

trimmed for quality using Trimmomatic version 0.36 (Bolger et al. 2014) with the parameters:

TRAILING:3, SLIDINGWINDOW:4:10, MINLEN:30. Trimmed reads were aligned to the zebra

finch reference genome (*Taeniopygia guttata* version 3.2.4; Warren et al. 2010) using the

program BWA-MEM (Li & Durbin, 2009) and a BAM file of this information was created for

each individual using the programs Picard (http://broadinstitute.github.io/picard/) and SAMtools

(Li et al. 2009). BAM files were converted into GVCF files using the HaplotypeCaller command

as part of GATK version 3.8 (McKenna et al. 2010). We then combined information from the

individuals in two ways to create 1) a genome-wide "variant site" VCF file containing only

variant site information, and 2) a series of chromosome-specific "info site" VCF files containing

information on both variant and invariant sites with sufficient coverage.

To create the genome-wide "variant site" VCF file, we used the GenotypeGVCFs

command in GATK to identify and isolate single nucleotide polymorphisms (SNPs) among

individuals. This command also converted the variant site information into a single VCF file

encompassing the entire nuclear genome. Using a combination of VCFtools (Danecek et al.

2011) and GATK, we filtered the VCF file to remove indels and non-biallelic SNPs. We also

discarded loci with QD < 2.0, MQ < 40.0, FS > 60.0, SOR > 3.0, or ReadPosRankSum < -8.0.

Finally, loci with more than 60% missing genotypes were removed. The average coverage of

variable sites in the resulting VCF file was 16.59.

To convert GVCF files into "info site" VCF files, we similarly employed the

GenotypeGVCFs command in GATK with the addition of the -allSites and -L flags to retain

invariant sites and split the information into chromosome-specific files. The resulting VCF files

were filtered using VCFtools and GATK to remove indels, sites with more than two alleles, sites

with more than 60% missing data, sites with MQ values lower than 20 and sites with

heterozygosities greater than 60% (to avoid potential paralogs).

Variant site analyses

The genome-wide "variant site" VCF file was analyzed using modified versions of the R

scripts described in Irwin et al. (2018), and all of our analyses used R version 3.6.2 (R Core

Team, 2014). A total of 374,780 SNPs were identified among allopatric yellowhammers and pine

buntings. For each of these SNPs, we calculated sample size, allele frequency, and Weir and

Cockerham's F_{ST} (Weir & Cockerham, 1984). Genetic differentiation between yellowhammers

and pine buntings was then visualized using a principal components analysis (PCA) generated

with the pca command and the svdImpute method to account for any missing genomic data using

the pcaMethods package (Stacklies et al. 2007). A Manhattan plot of F_{ST} values for 349,807

SNPs with known genomic locations was created using the package qqman (Turner, 2018).

Differentiation across the genome

To thoroughly investigate nuclear differentiation between allopatric yellowhammers and

pine buntings, we performed further analysis on both variant and invariant loci within "info site"

VCF files using R scripts described in Irwin et al. (2018).

We calculated Weir and Cockerham's F_{ST}, between-group nucleotide differentiation (π_B)

and within-group nucleotide variation (π_W) within nonoverlapping windows of available

sequence data across each chromosome. The first window was positioned at the "start" of each

chromosome as described in the zebra finch reference genome (Warren et al. 2010). We used a

window size of 2,000 bp of sequenced data rather than 10,000 bp (as in Irwin et al. 2018), to

visualize narrow peaks in relative and absolute differentiation within our dataset. We hereafter

refer to these windows as "genomic windows."

We developed a new R script to calculate a Tajima's D value (Tajima, 1989) for each of

the genomic windows. Values of Tajima's D were used to identify areas of the genome where

patterns of variation in yellowhammer and pine bunting populations deviated from a neutral

model. Significantly negative Tajima's D implies that the ratio of common vs. rare alleles is

lower than expected under neutrality, likely because of a selective sweep or population

expansion following a bottleneck. Significantly positive Tajima's D suggests that the ratio of

common vs. rare alleles is higher than expected under neutrality, potentially stemming from

balancing selection or a rapid population contraction.

Phylogenetic comparison with other Emberizidae species

We employed whole-genome averages of π_B between allopatric yellowhammers and pine

buntings as well as among these focal species and six other Emberizidae species (*Emberiza*

aureola, Emberiza calandra, Emberiza cioides, Emberiza cirlus, Emberiza hortulana and

Emberiza stewarti) to estimate a phylogeny. A list of average π_B values for each species pair was

converted into a distance matrix and used to create an unrooted neighbour-joining tree. This tree

was constructed using the ape package (Paradis & Schliep, 2019) and the BioNJ algorithm

(Gascuel, 1997) with *Emberiza aureola* set as the outgroup (Alström et al. 2008). The phylogeny

was created to provide further support for the sister relationship between yellowhammers and

pine buntings that had previously been hypothesized using mitochondrial markers (Alström et al.

2008; Irwin et al. 2009) but that was questioned in some studies (Rubtsov & Opeav, 2012). In

creating this phylogeny we were also able to better investigate the degree of mitonuclear

discordance within the system using a greater amount of nuclear data.

304 Signals of mitonuclear co-introgression

To test for preferential mitonuclear gene introgression (compared to other genomic

regions) between allopatric yellowhammers and pine buntings, we asked whether there was an

association between a list of known mitonuclear genes and a list of genomic windows that have

statistical characteristics most consistent with introgression. We call the latter "putative

introgression windows" (PIWs), and we explain their identification in detail below.

The mitonuclear genes that we analyzed for signals of introgression were all protein

coding with products that interacted directly with mtDNA or an immediate product of the

mitochondrial genome (i.e. protein or RNA). For these nuclear-encoded genes, any change in

mtDNA including those caused by introgression would likely cause selection for co-

introgression of compatible alleles (Gershoni et al. 2009; Burton & Barreto, 2012; Hill, 2019).

Mitonuclear genes that met these criteria included those that encode protein subunits of ATP

synthase or the first, third and fourth complex of the electron transport chain (ETC), assembly

and ancillary proteins involved in the formation of the ETC, and proteins that are part of the

transcription, translation or DNA replication machinery within the mitochondria (Diodato et al.

2013; Greber and Ban 2016; Hill, 2019). After removing any genes that were not annotated in

the zebra finch reference genome or that lacked a specific location on the reference genome, a

total of 162 mitonuclear genes remained for analysis (Supplementary Table 2).

Given the ongoing hybridization observed between yellowhammers and pine buntings

and the evidence for mtDNA introgression and replacement, it is reasonable to assume that

regions of the nuclear genome have also moved between the species. For each taxon, we

identified a subset of genomic windows that showed statistical properties most consistent with

substantial introgression, although we cannot be certain of introgression for any particular PIW.

We identified PIWs as those that possessed a low π_B average and a low Tajima's D average.

Low π_B indicates high similarity between the nucleotide sequences of the two groups as would

be expected if alleles had introgressed from one taxon into the other. Low Tajima's D suggests a

past selective sweep within a population which would also be expected if an adaptive allele had

introgressed from a separate taxon and swept throughout the receiving population. For this

analysis, our quantitative criteria for a PIW were a Tajima's D value within the lowest 5% of the

available windowed values and a π_B value within the lowest 30% of the available windowed

values. The Tajima's D threshold was kept relatively low to capture windows with particularly

strong signals of selection that could be associated with adaptive introgression while the π_B

threshold was left higher to identify an appreciable number of PIWs for analysis.

Using a custom R script, each mitonuclear gene was assigned to the genomic window

that minimized the absolute difference between the location of the mitonuclear gene centre and

the location of the window centre. We then determined the number of mitonuclear genes that

occurred within the PIWs of each taxon.

We conducted a Fisher's Exact test for both yellowhammers and pine buntings to

determine whether the proportion of mitonuclear genes within PIWs was significantly different

from what would be expected based on the total proportion of protein coding genes appearing

within these windows. To complete this analysis, a list of the 14,008 protein-coding genes

annotated in the zebra finch reference genome (Warren et al. 2010) was compiled. We removed

our 162 mitonuclear genes from this list and assigned the remaining 13,846 non-mitonuclear

genes to genomic windows using the methodology described above. The proportion of

mitonuclear and non-mitonuclear genes appearing in PIWs were then compared separately in

yellowhammers and pine buntings.

Results

When comparing allopatric yellowhammers and pine buntings, we identified 374,780

variable SNPs within our "variant site" VCF file and 13,703,455 invariant and 699,122 variant

sites across thirty autosomes and the Z chromosome within our "info site" VCF files following

filtering. In the latter "info site" files, we designated a total of 7,187 genomic windows (of 2000

sequenced bp each) across the genome, with each window covering an average distance of about

139 kilobases.

Phylogenetic comparison with other Emberizidae species

An unrooted neighbour-joining tree of average π_B values between yellowhammers, pine

buntings and six other Emberizidae species (Figure 2) depicted similar species relationships as

were identified previously using mitochondrial markers (Alström et al. 2008; Irwin et al. 2009).

Relative branch lengths were also similar, with the major exception being the branch length

between yellowhammers and pine buntings which was much longer in our analysis using nuclear

DNA. To put this into context, we determined the relative genetic distance between

yellowhammers and pine buntings versus the genetic distance between *E. stewarti* and either

member of the yellowhammer/pine bunting clade for our nuclear phylogeny and for the

previously calculated mitochondrial phylogeny (Irwin et al. 2009). The nuclear ratio was 11.4

times greater than the mitochondrial ratio which corroborates the presence of strong mitonuclear

discordance between yellowhammers and pine buntings. These results also support the

hypothesis of an extended period of divergence between yellowhammers and pine buntings

followed by adaptive mtDNA introgression.

Overall genetic differentiation

Based on 374,780 SNPs, the genome-wide F_{ST} estimate was 0.0232 between allopatric

yellowhammers and pine buntings. Despite this low average, a PCA based on the same SNP

genotypes separated yellowhammers and pine buntings into tight genetic clusters (Figure 3).

Two pine buntings were outliers along PC1, while the remaining yellowhammers and pine

buntings separated into distinct groups mainly along PC2. Further investigation into these

outliers revealed that they were males from the same location, but a kinship analysis completed

as part of a separate study did not find close kinship between the two pine buntings that could

explain their position (Nikelski et al. in prep). We also examined the PC1 loadings and found

that the signal for PC1 position was broadly distributed across the nuclear genome, rather than

being concentrated in a few highly influential regions (Supplementary Figure 1). Finally, we

temporarily removed one of the outliers and re-ran the PCA. This caused the other outlier to fall

into the pine bunting cluster, but revealed a further yellowhammer outlier (Supplementary Figure

2). Removal of this yellowhammer outlier in addition to one member of the pine bunting outlier

pair in turn revealed another yellowhammer outlier (Supplementary Figure 3). It is unclear what

is responsible for these outliers, but the distinct yellowhammer and pine bunting genetic clusters

remained intact in all the PCAs considered.

Differentiation across the genome

Relative differentiation between allopatric yellowhammers and pine buntings was highly heterogeneous across the nuclear genome with peaks in F_{ST} seen on most of the larger chromosomes (Figures 4, 5, 6A-B; Supplementary Figure 4). Chromosome Z in particular showed a large peak in F_{ST} with several SNPs possessing values close to one. In fact, F_{ST} for the Z chromosome was 0.1246—more than five times larger than the genome-wide F_{ST}.

Patterns of between-group nucleotide differentiation (π_B) and within-group nucleotide variation (π_W) were also heterogenous across the genome and comparable to each other in magnitude: genome-wide π_B = 0.0041; genome-wide π_W for both taxa= 0.0040 (Figure 5; Supplementary Figure 4). Because between-group and within-group nucleotide differentiation are so intimately related in their evolution and calculation, it is expected that windowed averages of these two statistics will show a highly positive relationship. In fact, most windowed π_B and π_W averages fell near a 1:1 association line (Figure 6C) which is equivalent to no or little differentiation. However, some genomic windows showed much reduced π_W compared to π_B; these were the windows with high F_{ST}. Additionally, we detected a weak but highly significant negative correlation between the windowed averages of F_{ST} and π_B (Spearman's Rank Correlation: -0.1196, p < 2.2 × 10^{-16}; Figure 6D) as is hypothesized if peaks in relative differentiation are products of repeated selective events (Cruickshank & Hahn, 2014; Irwin et al. 2018).

Finally, we found that Tajima's D varied across the genome but was mostly negative (Figure 5; Supplementary Figure 4), consistent with a history of population growth and/or

selective sweeps. The average genome-wide Tajima's D was similar between populations: -1.377

for yellowhammers and -1.335 for pine buntings.

Signals of mitonuclear co-introgression

Of the 7187 genomic windows identified across the nuclear genome, we classified 244

(3.4%) as PIWs within yellowhammers and 222 (3.1%) as PIWs within pine buntings. Average

values of π_B and Tajima's D in yellowhammer PIWs were 0.0016 and -2.3751 respectively, and

0.0019 and -2.3369 in pine bunting PIWs respectively. In non-PIWs, the average values of π_B

and Tajima's D were 0.0042 and -1.3416 in yellowhammers and 0.0042 and -1.3031 in pine

buntings. Of the PIWs identified in yellowhammer and pine bunting populations, 71 were shared

between the taxa. It should be noted that sharing of some PIWs is expected given that the

contribution of π_B to window selection was identical for both taxa (in contrast, Tajima's D was

calculated separately for yellowhammers and pine buntings).

Our examination of the gene content within yellowhammer PIWs revealed that they

contained a higher percentage (7.4%) of mitonuclear genes (12 of the 162 genes considered) than

of non-mitonuclear genes (4.1%; 574 of the 13,846 genes considered). This difference was

statistically significant (Fisher's Exact test: $p = 0.04714$), providing evidence for mitonuclear

genes preferentially appearing within yellowhammer PIWs. Pine bunting PIWs contained 4.3%

of the mitonuclear genes (7 of the 162 genes considered) and 3.3% of the non-mitonuclear genes

(455 of the 13,846 genes considered), a difference that was not statistically significant (Fisher's

Exact test: $p = 0.3806$).

The twelve mitonuclear genes that appeared within yellowhammer PIWs were: APOPT1,

COX5A, COX17, LARS2, MRPL1, MRPL27, MRPL32, MRPS7, MRPS25, NDUFC1, SSBP1

and UQCR11 (Table 2). Five of these genes encode protein subunits of the mitoribosome, three

encode structural subunits of the ETC, two encode assembly factors of the ETC, one encodes a

mitochondrial aminoacyl-tRNA synthetase and one encodes a single-stranded DNA-binding

protein involved in mtDNA replication. Two genes each appear on chromosomes 2, 4 and 18

while the rest appear on separate chromosomes. Interestingly, three of the five putatively

introgressed genes associated with the ETC are specifically associated with complex IV.

The seven mitonuclear genes that appeared within pine bunting PIWs were: ATP5H,

COX5A, MRPL2, MRPL58, MRPS7, MRPS14 and NDUFB4 (Table 2). Four of these genes

encode protein subunits of the mitoribosome and three encode structural subunits of the ETC.

Three genes appear on chromosome 18—with two genes sharing the same genomic window—

while the rest of the genes appear on separate chromosomes. The COX5A and MRPS7 genes

was found in both yellowhammer and pine bunting PIWs.

Discussion

Yellowhammers and pine buntings show negligible mtDNA differentiation (Irwin et al.

2009) but are well differentiated phenotypically (Panov et al. 2003, Rubtsov & Tarasov, 2017)

and moderately differentiated in AFLP nuclear markers (Irwin et al. 2009). In the wake of this

mitonuclear discordance, Irwin et al. (2009) proposed that mtDNA adaptively introgressed

between taxa following a period of allopatric isolation; however, due to the limited information

provided by AFLP analyses, the extent of mitonuclear discordance and of nuclear differentiation

between yellowhammers and pine buntings remained unknown lending some uncertainly to this

hypothesis. In the present study, analysis of genetic variation across the nuclear genome

identified heterogeneous nuclear differentiation between allopatric populations with strong

differentiation peaks that separated taxa into distinct genetic clusters. This result supports

yellowhammers and pine buntings experiencing a period of separate evolution followed by the

hybridization within their current contact zone in western and central Siberia. Our phylogenetic

analysis showing a longer branch length between yellowhammers and pine buntings based on

nuclear markers—when compared to a phylogeny based on mtDNA—also corroborates a

hypothesis of recent mtDNA introgression and mitochondrial haplotype replacement in this

system likely driven by selection (Irwin et al. 2009). Additionally, our finding that mitonuclear

genes are over-represented in putative introgression windows in yellowhammers is consistent

with mitonuclear co-introgression from pine buntings into yellowhammers.

Though genetically distinct, the genome-wide F_{ST} between allopatric yellowhammers and

pine buntings (0.0232) was comparable to or sometimes lower than the averages seen between

avian subspecies (e.g., subspecies of barn swallow: 0.017-0.026 [Scordato et al. 2017]; myrtle

warbler and Audubon's warbler: 0.077-0.106 [Irwin et al. 2018]; yellow- and red-shafted

northern flickers: 0.098 [Manthey et al. 2016]). This low genome-wide F_{ST} contrasts with the

moderate F_{ST} averages reported from an analysis of AFLP markers performed on the same

populations: 0.078 based on allele frequencies and 0.140 based on band frequencies (Irwin et al.

2009). However, the present study also revealed that relative differentiation was highly

heterogeneous across the nuclear genome with F_{ST} peaks on various chromosomes. It is possible

that the previous AFLP analysis captured a disproportionate number of loci within these

differentiation peaks, thereby inflating F_{ST} estimates. This comparison highlights the caution that

should be taken when interpreting genome-wide averages as highly variable genetic

differentiation landscapes can cause large variability in F_{ST} estimates when they are based on a

limited and non-random sample of loci.

The F_{ST} peaks seen between yellowhammers and pine buntings on larger autosomes and

most significantly on the Z chromosome are consistent with the "islands of differentiation" often

noted in comparisons of closely related taxa (Harr, 2006; Nadeau et al. 2012; Irwin et al. 2018).

In contrast to these islands, large regions of close similarity in π_B and π_W suggests high gene

flow between taxa across much of the nuclear genome. This scenario is consistent with the

observed extensive hybridization between these taxa (Panov et al. 2003; 2007; Rubtsov, 2007;

Rubtsov & Tarasov, 2017). Nevertheless, the high F_{ST} islands, those with much reduced π_W

compared to π_B, can be explained by divergent selection causing low gene flow in these regions.

It is unlikely that this pattern is the result of genetic drift over an extended period of geographic

separation, as this would result in most genomic regions deviating slightly from $\pi_B = \pi_W$

congruence rather than the observed pattern of extreme heterogeneity. Instead, this trend

suggests that selection acted in a way that lowered π_W relative to π_B within "islands of

differentiation". Considering that high F_{ST} regions were associated with relatively low values of

π_B, we propose that differentiation islands in this system are most consistent with a model

invoking repeated bouts of selection that lower nucleotide diversity (Cruickshank & Hahn, 2014;

Irwin et al. 2018). A sweep-before-differentiation model (Irwin et al. 2018), where F_{ST} peaks are

produced by adaptive selective sweeps between populations followed by adaptive selection at the

same regions in local populations, is particularly in line with the extensive hybridization

presently observed between yellowhammers and pine buntings.

Of the "islands of differentiation" identified between taxa, the tallest and widest was

found on the Z chromosome. Greater differentiation on the Z chromosome compared to

autosomes is a common observation when comparing closely related species (Borge et al. 2005;

Ruegg et al. 2014; Sackton et al. 2014) and is consistent with stronger positive selection (the

"faster Z/X effect") and with less efficient purifying selection on this chromosome (Mank et al.

2010; reviewed in Meisel & Connallon, 2013; reviewed in Irwin, 2018). However, the large

region of the Z chromosome that have F_{ST} values near zero suggests that additional factors are

involved in producing this island of differentiation.

One possible explanation for the large differentiation island on chromosome Z could be

that it corresponds with an area of low recombination—a region of connected loci that tend to be

inherited together, leading to linked selection of nearby loci. Strong divergent selection acting on

one SNP within this region would act similarly on all the loci that are linked to it such that a

wide, highly divergent genomic block would become fixed and appear as an "island" between

taxa (reviewed in Cutter & Payseur, 2013). Areas of low recombination and linkage are often

associated with inversion polymorphisms (reviewed in Smukowski & Noor, 2011) as different

orientations of an inversion experience little successful recombination (reviewed in Kirkpatrick,

2010). Further research is warranted to characterize the nature of this differentiated region as

well as whether it houses an inversion polymorphism.

While numerous "islands of differentiation" were observed between yellowhammers and

pine buntings implying moderate genetic divergence, mtDNA introgression has the potential to

homogenize the nuclear genomes of these taxa at mitonuclear genes by selecting for co-

introgression of compatible alleles (Beck et al. 2015; Sloan et al. 2017; Morales et al. 2018). In

the yellowhammer and pine bunting system, we found statistical support for the preferential

introgression of mitonuclear genes into yellowhammers, but no statistical support for the

preferential introgression into pine buntings. More specifically, our analysis showed that the

proportion of introgressing mitonuclear genes was 1.7X higher in yellowhammers versus pine

buntings. While these results suggest mitonuclear introgression primarily in the direction of pine

buntings into yellowhammers, we also noted some consistency in the functions of putatively

introgressing mitonuclear genes in both taxa.

Three of the mitonuclear genes within yellowhammer PIWs and three within pine bunting

PIWs encode structural subunits of the ETC. The ETC is broken into five protein complexes

which, through a series of enzymatic reactions, perform oxidative phosphorylation to produce

ATP necessary for organism survival (reviewed in Ernster & Schatz, 1981). Four of the five ETC

complexes are made up of subunits encoded by both the nuclear and mitochondrial genome (Hill,

2019) and correct fit between differentially encoded subunits is essential for the flow of electrons

and protons across the ETC. To put this in perspective, changing even a single amino acid in one

subunit can significantly disrupt its ability to interact with other subunits within an ETC complex

(e.g. Gershoni et al. 2014). Because of the tight interactions within complexes and the

consequences of subunit incompatibility, introgression of mtDNA is expected to select for co-

introgression of mitonuclear genes encoding ETC structural subunits. Such co-introgression has

been detected between differentially adapted populations of eastern yellow robin where mtDNA

introgression between populations was followed by similar introgression of mitonuclear genes

encoding subunits of complex I (Morales et al. 2018) and between different species of

Drosophila where introgression and replacement of the mtDNA of one species during

hybridization selected for co-introgression of genes that encode subunits of complex IV (Beck et

al. 2015).

Of the ETC complexes, complex IV showed the strongest signal of co-introgression in

the yellowhammer and pine bunting system. Three of the genes within yellowhammer PIWs and

one gene within pine bunting PIWs were associated with this complex. As well, the gene

COX5A—a structural subunit of complex IV—appeared in both sets of PIWs. It is unlikely that

this gene introgressed in both directions, but it is possible that COX5A adaptively swept in both

populations. In this situation, a particularly adaptive allele may have appeared in one species and

swept to high frequency before co-introgressing into the other species following mtDNA

introgression. Interestingly, the COX5A gene was one of the subunits that co-introgressed in the

Drosophila example discussed above (Beck et al. 2015) lending some support to its particular

importance to mitonuclear coevolution. More generally, complex IV is often used as a model for

studying mitonuclear interactions due to its distinctive structure where a core of mitochondrial-

encoded subunits is surrounded by nuclear-encoded subunits (Saraste, 1999). With such an

excess of mitonuclear interactions, incompatibility involving complex IV has been investigated

and detected in several systems including within primate xenomitochondrial cybrids (Barrientos

et al. 2000) and between different species of *Drosophila* (Sackton et al. 2003). Furthermore,

work by Osada & Akashi (2012) has provided strong evidence for compensatory coevolution

between mitonuclear genes related to complex IV—including COX5A—and mtDNA among

primates particularly at interacting amino acids of differentially encoded subunits. Altogether,

these results suggest a crucial role for complex IV in mitonuclear coevolution as it may relate to

divergence and speciation between yellowhammers and pine buntings.

Another group of mitonuclear genes that appeared consistently within the yellowhammer

and pine bunting PIWs were those encoding subunits of the mitoribosome (five in

yellowhammer PIWs and four in pine bunting PIWs). MRPS7, like COX5A, appeared in both

yellowhammer and pine bunting PIWs suggesting that this gene may have adaptively swept

through both taxa. Unlike the protein-protein interactions occurring within ETC complexes,

mitonuclear interactions in the mitoribosome are between nuclear-encoded proteins and

mitochondrial-encoded RNA (Hill, 2019). Protein subunits associate closely with rRNA during

the formation of a mitoribosome, but also interact with mRNA and tRNA during the synthesis of

mitochondrial proteins (Greber & Ban, 2016). Currently, research is limited on the extent and

importance of interactions between mitoribosomal subunits and mitochondrial RNA. However,

the fact that interactions between components are extensive and necessary for the synthesis of

mitochondrial proteins suggests close coevolution between mtDNA and genes encoding

mitoribosomal subunits that could strongly select for mitonuclear co-introgression following

mtDNA introgression.

To summarize, yellowhammers and pine buntings are sister taxa that are divergent in

appearance, song, and ecology (Panov et al. 2003; Rubtsov & Tarasov, 2017) yet vary greatly in

their genomic differentiation from virtually none (at the mitochondrial genome) to nearly fixed

(the differentiation peak on the Z chromosome). These patterns are best explained by a period of

differentiation while geographically separated, followed by hybridization and mitochondrial

DNA introgression. We observed evidence for preferential mitonuclear gene introgression

(compared to introgression of other genes) from pine buntings into yellowhammers, as well as a

tendency for mitonuclear genes encoding structural components of the ETC and the

mitoribosome to appear within the PIWs of both taxa. One intriguing possibility is that

mitonuclear co-introgression could have resulted in reduced mitonuclear incompatibilities

between yellowhammers and pine buntings (Gershoni et al. 2009; Burton & Barreto, 2012; Hill,

2019), thereby contributing to their current extensive hybridization within central Siberia (Panov

et al. 2003; 2007; Rubtsov, 2007; Rubtsov & Tarasov, 2017). This idea leads into the question—

which can be addressed through a close examination of genomic variation within the hybrid

zone—of whether the observed islands of differentiation on the Z and autosomes are sufficient in

stabilizing yellowhammers and pine buntings as separate entities where they hybridize, or

whether the two taxa are gradually merging into a single species.

Author contributions:

E. N., D.I., and A.S.R. conceived of this study. A.S.R. collected samples. E. N. and A.S.R.

completed molecular techniques. E.G.M.N. conducted data analysis and wrote this manuscript

with input from D.I. and A.S.R.

Acknowledgements:

For providing valuable feedback, we thank Dolph Schluter, Eric Taylor, Judith Mank, Elizabeth

Natola, Rashika Ranasinghe, Kenneth Askelson, Finola Fogarty, Quinn McCallum, Ana

Barreira, Jamie Clarke, Armando Geraldes and Jessica Irwin. For their kindness and support

during field work, we thank the Tazeev family and Madelyn Ore. For providing additional

samples, we thank The Bell Museum, The Burke Museum of Natural History and Culture, The

Field Museum, The State Darwin Museum, The Swedish Museum of Natural History, The

Zoological Museum of the Zoological Institute of the Russian Academy of Sciences, the

Zoological Museum of the University of Copenhagen and their accompanying personnel. Major

research funding was provided by the Natural Sciences and Engineering Research Council of

Canada (NSERC CGSM award to E.N, Discovery Grants RGPIN-2017-03919 and RGPAS-

2017-507830 awarded to D.I.) and by the Werner and Hildegard Hesse research awards

(Research award in Ornithology and Fellowship in Ornithology awarded to E.G.M.N. by the

University of British Columbia).

Data Accessibility Statement:

Raw DNA sequencing reads are available on the NBCI Sequence Read Archive (BioProject

PRJNA768601). Read processing codes, barcodes, genotype data and R codes associated with

statistical analyses will be made available on Dryad upon publication acceptance.

Conflict of Interest Statement:

None declared

Bibliography

Alcaide, M., Scordato, E. S. C., Price, T. D., & Irwin, D. E. (2014). Genomic divergence in a
ring species complex. *Nature, 511*(7507), 83-85. doi:10.1038/nature13285

Alström, P., Olsson, U., Lei, F., Wang, H., Gao, W., & Sundberg, P. (2008). Phylogeny and
classification of the old world *Emberizini* (*Aves, Passeriformes*). *Molecular*
Phylogenetics and Evolution, 47(3), 960-973. doi:10.1016/j.ympev.2007.12.007

Alves, P. C., Melo-Ferreira, J., Freitas, H., & Boursot, P. (2008). The ubiquitous mountain hare
mitochondria: Multiple introgressive hybridization in hares, genus *Lepus*. *Philosophical*
Transactions. Biological Sciences, 363(1505), 2831-2839. doi:10.1098/rstb.2008.0053

Avise, J. C. (2000). Phylogeography: The history and formation of species. Harvard University
Press.

Ballard, J. W. O., Melvin, R. G., Katewa, S. D., & Maas, K. (2007). Mitochondrial DNA
variation is associated with measurable differences in life-history traits and mitochondrial
metabolism in *Drosophila simulans. Evolution, 61*(7), 1735-1747. doi:10.1111/j.1558-
5646.2007.00133.x

Barrientos, A., Müller, S., Dey, R., Wienberg, J., & Moraes, C. T. (2000). Cytochrome c oxidase
assembly in primates is sensitive to small evolutionary variations in amino acid sequence.
Molecular Biology and Evolution, 17(10), 1508-1519.
doi:10.1093/oxfordjournals.molbev.a026250

Baute, G. J., Owens, G. L., Bock, D. G., & Rieseberg, L. H. (2016). Genome-wide genotyping-
by-sequencing data provide a high-resolution view of wild *Helianthus* diversity, genetic
structure, and interspecies gene flow. *American Journal of Botany, 103*(12), 2170-2177.
doi:10.3732/ajb.1600295

Beck, E. A., Thompson, A. C., Sharbrough, J., Brud, E., & Llopart, A. (2015). Gene flow
between *Drosophila yakuba* and *Drosophila santomea* in subunit V of cytochrome c
oxidase: A potential case of cytonuclear cointrogression. *Evolution, 69*(8), 1973-1986.
doi:10.1111/evo.12718

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics, 30*(15), 2114-2120. doi:10.1093/bioinformatics/btu170

Borge, T., Webster, M. T., Andersson, G., & Saetre, G. (2005). Contrasting patterns of
polymorphism and divergence on the Z chromosome and autosomes in two *Ficedula*
flycatcher species. *Genetics, 171*(4), 1861-1873. doi:10.1534/genetics.105.045120

Burton, R. S., & Barreto, F. S. (2012). A disproportionate role for mtDNA in Dobzhansky–
Muller incompatibilities? *Molecular Ecology, 21*(20), 4942-4957.
doi:10.1111/mec.12006

Bryson, J., Robert W, De Oca, A. N., Jaeger, J. R., & Riddle, B. R. (2010). Elucidation of cryptic
diversity in a widespread Nearctic treefrog reveals episodes of mitochondrial gene
capture as frogs diversified across a dynamic landscape. *Evolution, 64*(8), 2315-
2330. doi:10.1111/j.1558-5646.2010.01014.x

Calvo, S. E., & Mootha, V. K. (2010). The mitochondrial proteome and human disease. *Annual*
Review of Genomics and Human Genetics, 11(1), 25-44.
doi:10.1146/annurev-genom-082509-141720

Coyne, J. A., & Orr, H. A. (2004). *Speciation*. Sinauer Associates.

Cruickshank, T. E., & Hahn, M. W. (2014). Reanalysis suggests that genomic islands of
speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology,*
23(13), 3133-3157. doi:10.1111/mec.12796

Cutter, A. D., & Payseur, B. A. (2013). Genomic signatures of selection at linked sites: Unifying
the disparity among species. *Nature Reviews. Genetics, 14*(4), 262-274.
doi:10.1038/nrg3425

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R.
E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R. & 1000 Genomes
Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics,*
27(15), 2156-2158. doi:10.1093/bioinformatics/btr330

Diodato, D., Ghezzi, D., & Tiranti, V. (2014). The mitochondrial aminoacyl tRNA synthetases:
Genes and syndromes. *International Journal of Cell Biology, 2014*(2014), 1-11.
doi:10.1155/2014/787956

Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler, E. S., & Mitchell,
S. E. (2011). A robust, simple genotyping-by-sequencing (GBS) approach for high
diversity species. *PloS One, 6*(5), e19379. doi:10.1371/journal.pone.0019379

Ernster, L., & Schatz, G. (1981). Mitochondria: A historical review. *The Journal of Cell*
Biology, 91(3), 227s-255s. doi:10.1083/jcb.91.3.227s

Gascuel, O. (1997). BIONJ: An improved version of the NJ algorithm based on a simple model
of sequence data. *Molecular Biology and Evolution, 14*(7), 685-695.
doi:10.1093/oxfordjournals.molbev.a025808

Geraldes, A., Askelson, K. K., Nikelski, E., Doyle, F. I., Harrower, W. L., Winker, K., & Irwin,
D. E. (2019). Population genomic analyses reveal a highly differentiated and endangered

genetic cluster of northern goshawks (*Accipiter gentilis laingi*) in Haida Gwaii.
Evolutionary Applications, 12(4), 757-772. doi:10.1111/eva.12754

Gershoni, M., Levin, L., Ovadia, O., Toiw, Y., Shani, N., Dadon, S., Barzilai, N., Bergman, A.,
Atzmon, G., Wainstein, J., Tsur, A., Nijtmans, L. G. J., Glaser, B., & Mishmar, D.
(2014). Disrupting mitochondrial-nuclear coevolution affects OXPHOS complex I
integrity and impacts human health. *Genome Biology and Evolution, 6*(10), 2665-2680.
doi:10.1093/gbe/evu208

Gershoni, M., Templeton, A. R., & Mishmar, D. (2009). Mitochondrial bioenergetics as a major
motive force of speciation. *Bioessays, 31*(6), 642-650. doi:10.1002/bies.200800139

Greber, B. J., & Ban, N. (2016). Structure and function of the mitochondrial ribosome. *Annual*
Review of Biochemistry, 85(1), 103-132. doi:10.1146/annurev-biochem-060815-014343

Harr, B. (2006). Genomic islands of differentiation between house mouse subspecies. *Genome*
Research, 16(6), 730-737. doi:10.1101/gr.5045006

Hebert, P. D., Penton, E. H., Burns, J. M., Janzen, D. H., & Hallwachs, W. (2004). Ten species
in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly
Astraptes fulgerator. *Proceedings of the National Academy of Sciences*, *101*(41), 14812-
14817. doi:10.1073/pnas.0406166101

Hejase, H. A., Salman-Minkov, A., Campagna, L., Hubisz, M. J., Lovette, I. J., Gronau, I., &
Siepel, A. (2020). Genomic islands of differentiation in a rapid avian radiation have been
driven by recent selective sweeps. *Proceedings of the National Academy of*
Sciences, *117*(48), 30554–30565. doi:10.1073/pnas.2015987117
736

Hill, G. E. (2019). *Mitonuclear Ecology* (First ed.). Oxford University Press.

Hill, G. E. (2020). Genetic hitchhiking, mitonuclear coadaptation, and the origins of mt DNA
barcode gaps. *Ecology and Evolution, 10*(17), 9048-9059. doi:10.1002/ece3.6640

Hulsey, C. D., Bell, K. L., García-de-León, F. J., Nice, C. C., & Meyer, A. (2016). Do relaxed
selection and habitat temperature facilitate biased mitogenomic introgression in a
narrowly endemic fish? *Ecology and Evolution, 6*(11), 3684- 3698.
doi:10.1002/ece3.2121

Irwin, D. E. (2018). Sex chromosomes and speciation in birds and other ZW systems. *Molecular*
Ecology, 27(19), 3831-3851. doi:10.1111/mec.14537

Irwin, D. E., Alcaide, M., Delmore, K. E., Irwin, J. H., & Owens, G. L. (2016). Recurrent
selection explains parallel evolution of genomic regions of high relative but low absolute
differentiation in a ring species. *Molecular Ecology, 25*(18), 4488-4507.
doi:10.1111/mec.13792

Irwin, D. E., Milá, B., Toews, D. P. L., Brelsford, A., Kenyon, H. L., Porter, A. N., Grossen, C.,
Delmore, K. E., Alcaide, M., & Irwin, J. H. (2018). A comparison of genomic islands of
differentiation across three young avian species pairs. *Molecular Ecology, 27*(23), 4839-
4855. doi:10.1111/mec.14858

Irwin, D. E., Rubtsov, A. S., & Panov, E. N. (2009). Mitochondrial introgression and
replacement between yellowhammers (*Emberiza citrinella*) and pine buntings (*Emberiza*
leucocephalos) (Aves: Passeriformes). *Biological Journal of the Linnean Society, 98*(2),
422-438. doi:10.1111/j.1095-8312.2009.01282.x

Kerr, K. C. R., Stoeckle, M. Y., Dove, C. J., Weigt, L. A., Francis, C. M., & Hebert, P. D. N.
(2007). Comprehensive DNA barcode coverage of North American birds. *Molecular*
*Ecology Notes, 7(*4), 535-543. doi:10.1111/j.1471-8286.2007.01670.x

Kirkpatrick, M. (2010). How and why chromosome inversions evolve. *PLoS Biology, 8*(9),
e1000501. doi:10.1371/journal.pbio.1000501

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics, 25*(14), 1754-1760. doi:10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., &
Durbin, R. (2009). The sequence alignment map format and SAMtools. *Bioinformatics,*
25(16), 2078-2079. doi:10.1093/bioinformatics/btp352

Llopart A., Herrig D., Brud E., & Stecklein Z. (2014) Sequential adaptive introgression of the
mitochondrial genome in *Drosophila yakuba* and *Drosophila santomea*. *Molecular*
Ecology, 23 (5), 1124–1136. doi:10.1111/mec.12678
782

783 Lotz, C., Lin, A. J., Black, C. M., Zhang, J., Lau, E., Deng, N., Wang, Y., Zong, N. C., Choi, J.
784    H., Xu, T., Liem, D. A., Korge, P., Weiss, J. N., Hermjakob, H., Yates, J. R., III.,
785    Apweiler, R., & Ping, P. (2014). Characterization, design, and function of the
786    mitochondrial proteome: From organs to organisms. *Journal of Proteome Research,*
787    *13*(2), 433-446. doi:10.1021/pr400539j

789 Lu, J., & Wu, C. (2005). Weak selection revealed by the whole-genome comparison of the X
790    chromosome and autosomes of human and chimpanzee. *Proceedings of the National*
791    *Academy of Sciences, 102*(11), 4063-4067. doi:10.1073/pnas.0500436102

793 Lynch, M., Koskella, B., & Schaack, S. (2006). Mutation pressure and the evolution of organelle
794    genomic architecture. *Science, 311*(5768), 1727-1730. doi:10.1126/science.1118884

796 Mank, J. E., Nam, K., & Ellegren, H. (2010). Faster-Z evolution is predominantly due to genetic
797    drift. *Molecular Biology and Evolution, 27*(3), 661-670. doi:10.1093/molbev/msp282

799 Manthey, J. D., Geiger, M., & Moyle, R. G. (2017). Relationships of morphological groups in
800    the northern flicker superspecies complex (*Colaptes auratus* & *C. chrysoides*).
801    *Systematics and Biodiversity, 15*(3), 183-191. doi:10.1080/14772000.2016.1238020

803 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella,
804    K., Altshuler, D., Gabriel, S., Daley, M., & DePristo, M. A. (2010). The genome analysis
805    toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data.
806    *Genome Research, 20*(9), 1297-1303. doi:10.1101/gr.107524.110

808 Meisel, R. P., & Connallon, T. (2013). The faster-X effect: Integrating theory and data. *Trends in*
809    *Genetics, 29*(9), 537-544. doi:10.1016/j.tig.2013.05.009

811 Moore, W. S. (1995). Inferring phylogenies from mtDNA variation: Mitochondrial-gene trees
812    versus nuclear-gene trees. *Evolution, 49*(4), 718-726. doi:10.2307/2410325

814 Morales, H. E., Pavlova, A., Amos, N., Major, R., Kilian, A., Greening, C., & Sunnucks, P.
815    (2018). Concordant divergence of mitogenomes and a mitonuclear gene cluster in bird
816    lineages inhabiting different climates. *Nature Ecology & Evolution, 2*(8), 1258-1267.
817    doi:10.1038/s41559-018-0606-3

Nadeau, N. J., Whibley, A., Jones, R. T., Davey, J. W., Dasmahapatra, K. K., Baxter, S. W.,
Quail, M. A., Joron, M., Ffrench-Constant, R. H., Blaxter, M. L., Mallet, J., & Jiggins, C.
D. (2012). Genomic islands of divergence in hybridizing *Heliconius* butterflies identified
by large-scale targeted sequencing. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, 367(1587), 343-353. doi:10.1098/rstb.2011.0198

Osada, N., & Akashi, H. (2012). Mitochondrial-nuclear interactions and accelerated
compensatory evolution: Evidence from the primate cytochrome c oxidase complex.
Molecular Biology and Evolution, 29(1), 337-346. doi:10.1093/molbev/msr211

Panov, E. N., Rubtsov, A. S., & Monzikov, D. G. (2003). Hybridization between yellowhammer
and pine bunting in Russia. *Dutch Birding, 25,* 17-31.

Panov, E. N., Rubtsov, A. S., & Mordkovich, M. V. (2007). New data on the relationships
between two species of buntings (*Emberiza citrinella* and *E. leucocephalos*) hybridizing
in the areas of overlap of their ranges, *Zoologicheskii Zhurnal, 86*(11), 1362–1378.

Paradis, E., & Schliep, K. (2019). ape 5.0: An environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics, 35*(3), 526-528.
doi:10.1093/bioinformatics/bty633

Price, T. (2008). *Speciation in birds*. Roberts and Co.

R Core Team (2014). R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing. Retrieved from http://www.r-project.org/

Rubtsov, A.S. 2007. Variability of songs of the yellowhammer (*Emberiza citrinella*) and pine
bunting (*Emberiza leucocephala*) as an evidence of population structure and evolutionary
history of the species, *Zoologicheskii Zhurnal, 86*(7), 863–876.

Rubtsov, A. S., & Opaev, A. S. (2012). Phylogeny reconstruction of the yellowhammer
(*Emberiza citrinella*) and pine bunting (*Emberiza leucocephala*) based on song and
morphological characters. *Biology Bulletin of the Russian Academy of Sciences, 39*(9),
715-728. doi:10.1134/S1062359012090087

Rubtsov, A. S., & Tarasov, V. V. (2017). Relations between the yellowhammer (*Emberiza*
citrinella) and the pine bunting (*Emberiza leucocephalos*) in the forested steppe of the
Trans-Urals. *Biology Bulletin, 44*(9), 1059-1072. doi:10.1134/S1062359017090114

Ruegg, K., Anderson, E. C., Boone, J., Pouls, J., & Smith, T. B. (2014). A role for migration-
linked genes and genomic islands in divergence of a songbird. *Molecular Ecology,*
23(19), 4757-4769. doi:10.1111/mec.12842

Sackton, T. B., Corbett-Detig, R. B., Nagaraju, J., Vaishna, L., Arunkumar, K. P., & Hartl, D. L.
(2014). Positive selection drives faster-z evolution in silkmoths. *Evolution, 68*(8), 2331-
2342. doi:10.1111/evo.12449

Sackton, T. B., Haney, R. A., & Rand, D. M. (2003). Cytonuclear coadaptation in *Drosophila*:
Disruption of cytochrome c oxidase activity in backcross genotypes. *Evolution, 57*(10),
2315-2325. doi:10.1111/j.0014-3820.2003.tb00243.x

Saraste, M. (1999). Oxidative phosphorylation at the fin de siècle. *Science, 283*(5407), 1488-
1493. doi:10.1126/science.283.5407.1488

Scordato, E. S. C., Wilkins, M. R., Semenov, G., Rubtsov, A. S., Kane, N. C., & Safran, R. J.
(2017). Genomic variation across two barn swallow hybrid zones reveals traits associated
with divergence in sympatry and allopatry. *Molecular Ecology, 26*(20), 5676-
5691. doi:10.1111/mec.14276

Sloan, D. B., Havird, J. C., & Sharbrough, J. (2017). The on-again, off-again relationship
between mitochondrial genomes and species boundaries. *Molecular Ecology, 26*(8),
2212-2236. doi:10.1111/mec.13959

Smukowski, C. S., & Noor, M. A. F. (2011). Recombination rate variation in closely related
species. *Heredity, 107*(6), 496-508. doi:10.1038/hdy.2011.44

Stacklies, W., Redestig, H., Scholz, M., Walther, D., & Selbig, J. (2007). pcaMethods-a
bioconductor package providing PCA methods for incomplete data. *Bioinformatics,*
23(9), 1164-1167. doi:10.1093/bioinformatics/btm069

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics, 123*(3), 585-595.

Thornton, K., & Long, M. (2002). Rapid divergence of gene duplicates on the *Drosophila*
melanogaster X chromosome. *Molecular Biology and Evolution, 19*(6), 918-925.
doi:10.1093/oxfordjournals.molbev.a004149

Toews, D. P. L., & Brelsford, A. (2012). The biogeography of mitochondrial and nuclear
discordance in animals. *Molecular Ecology, 21*(16), 3907-3930. doi:10.1111/j.1365-
294X.2012.05664.x

Turner, S. (2018). qqman: An R package for visualizing GWAS results using Q-Q and
Manhattan plots. *Journal of Open Source Software, 3*(25), 731. doi:10.21105/joss.00731

Wang, S., Ore, M. J., Mikkelsen, E. K., Lee-Yaw, J., Toews, D. P. L., Rohwer, S., & Irwin, D.
(2021). Signatures of mitonuclear coevolution in a warbler species complex. *Nature*
Communications, in press. doi:10.1038/s41467-021-24586-8

Warren, W. C., Clayton, D. F., Ellegren, H., Arnold, A. P., Hillier, L. W.,Künstner, A., Searle,
S., White, S., Vilella, A. J., Fairley, S., Heger, A., Kong, L., Ponting, C. P., Jarvis, E. D.,
Mello, C. V., Minx, P., Lovell, P., Velho, T. A. F., Ferris, M., … Wilson, R. K. (2010).
The genome of a songbird. *Nature, 464*(7289), 757–762. doi:10.1038/nature08819

Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population
structure. *Evolution, 38*(6), 1358-1370. doi:10.1111/j.1558-5646.1984.tb05657.x

Wu, C. (2001). The genic view of the process of speciation. *Journal of Evolutionary*
Biology, 14(6), 851-865. doi:10.1046/j.1420-9101.2001.00335.x

Yannic, G., Dubey, S., Hausser, J., & Basset, P. (2010). Additional data for nuclear DNA give
new insights into the phylogenetic position of *Sorex granarius* within the *Sorex araneus*
group. *Molecular Phylogenetics and Evolution, 57*(3), 1062-1071.
doi:10.1016/j.ympev.2010.09.015

Tables

Table 1. Geographic locations and sample sizes of the sites included in this study. Sampling locations may include multiple sites that appeared too close together to be shown in detail in Figure 1A. Full details for the sites included in each sampling locations can be found in Supplementary Table 1. The sampling location numbers that appear in the "Sampling Location" column correspond to those that appear in red in Figure 1A. The "Sample Size" columns describes the total number of samples collected from a particular location.

Sampling Location	Latitude (°N)	Longitude (°E)	*E. citrinella* sample size	*E. leucocephalos* sample size
1	57.99	12.49	1	0
2	59.81	17.05	1	0
3	51.71	18.61	1	0
4	55.28	20.97	5	0
5	65.86	21.48	2	0
6	51.38	35.84	3	0
7	55.97	38.50	18	0
8	61.45	38.67	12	0
9	43.54	40.47	1	0
10	65.85	44.24	1	0
11	58.33	44.76	1	0
12	51.20	57.27	7	0
13	49.64	110.17	0	2
14	50.66	115.09	0	17
15	51.12	118.56	0	15
16	50.56	143.08	0	8
		Total	**53**	**42**

Table 2. Identities, chromosomal locations, windowed Tajima's D values and functions of mitonuclear genes that appeared within 244 yellowhammer PIWs or within 222 pine bunting PIWs. In the "Mitonuclear Gene Function" column, ETC stands for "Electron Transport Chain". The "*" indicates a mitonuclear gene that appeared in both yellowhammer and pine bunting PIWs. The "[†]" indicates two genes that appeared within the same PIW.

Mitonuclear Gene	Chromosome where mitonuclear gene is found	Windowed Tajima's D Value	Mitonuclear Gene Function
Yellowhammer (E. citrinella)			
APOPT1	5	-2.207	Assembly factor/ancillary protein for ETC complex IV
COX5A*	10	-2.420	Structural subunit of ETC complex IV
COX17	1	-2.509	Assembly factor/ancillary protein for ETC complex IV
LARS2	2	-2.238	Mitochondrial aminoacyl-tRNA synthetase
MRPL1	4	-2.207	Mitochondrial large ribosomal subunit protein
MRPL27	18	-2.214	Mitochondrial large ribosomal subunit protein
MRPL32	2	-2.306	Mitochondrial large ribosomal subunit protein
MRPS7*	18	-2.628	Mitochondrial small ribosomal subunit protein
MRPS25	12	-2.323	Mitochondrial small ribosomal subunit protein
NDUFC1	4	-2.399	Structural subunit of ETC complex I
SSBP1	1A	-2.362	Single stranded DNA-binding protein
UQCR11	28	-2.499	Structural subunit of ETC complex III
Pine bunting (E. leucocephalos)			
ATP5H[†]	18	-2.601	Structural subunit of ETC complex V
COX5A*	10	-2.545	Structural subunit of ETC complex IV

MRPL2	3	-2.247	Mitochondrial large ribosomal subunit protein
MRPL58[†]	18	-2.601	Mitochondrial large ribosomal subunit protein
MRPS7*	18	-2.252	Mitochondrial small ribosomal subunit protein
MRPS14	8	-2.173	Mitochondrial small ribosomal subunit protein
NDUFB4	1	-2.304	Structural subunit of ETC complex I

Figures

A.

E. citrinella ——— 🟡   **E. leucocephalos** - - - - 🔴

B.

Emberiza citrinella

Pure Citrinella (PC)   Almost Citrinella (SC)

Emberiza leucocephalos

Pure Leucocephalos (PL)   Almost Leucocephalos (SL)

Figure 1. A) Map of sampling locations included in this study. Red numbers accompanying each location correspond to the sampling location numbers appearing in Table 1 which also describes sample sizes. Sampling locations may include multiple sites that appeared too close together to be shown in detail in this figure. Full details for the sites included in each sampling location can be found in Supplementary Table 1. Sampling location points are coloured based on the taxon caught in each area: yellowhammer (*Emberiza citrinella*; yellow) and pine bunting (*Emberiza leucocephalos*; brown). The solid black line indicates the geographic range of the yellowhammer and the dashed black line indicates the geographic range of the pine bunting as described in Irwin et al. (2009). **B)** Photos of plumage variation between yellowhammers and pine buntings. Each photo represents one of four phenotypic classes: PC, SC, PL and SL. Individuals with a PC and SC phenotypic class were grouped together as *Emberiza citrinella* and individuals with a PL and SL phenotypic class were grouped together as *Emberiza leucocephalos*. All photos are credited to Dr. Alexander Rubtsov.

Figure 2. Unrooted neighbour-joining tree of Emberizidae species constructed based on average absolute between-population nucleotide diversity (π_B). Sample sizes for each species are as follows: *E. aureola* = 1, *E. calandra* = 1, *E. cioides* = 1, *E. hortulana* = 1, *E. cirlus* = 6, *E. stewarti* = 4, *E. citrinella* = 53 and *E. leucocephalos* = 42.

Figure 3. PCA of genetic variation between allopatric yellowhammers (yellow; n = 53) and allopatric pine buntings (brown; n = 42), based on 374,780 genome-wide SNPs. PC1 and PC2 explain 3.6% and 2.9%, respectively, of the variation among individuals.

Figure 4. Relative differentiation (F_{ST}) of 349,807 genome-wide SNPs identified between allopatric yellowhammers (n = 53) and allopatric pine buntings (n = 42), with chromosomes represented with alternating black and grey. Narrow regions of elevated differentiation can be seen on many autosomes, and there are broad regions of high differentiation on the Z chromosome.

Figure 5. Patterns of genetic variation comparing allopatric yellowhammers (n = 53) and allopatric pine buntings (n = 42) across three chromosomes (2, 5 and Z). Relative nucleotide differentiation (F_{ST}), absolute between-population nucleotide differentiation (π_B), absolute within-population nucleotide variation (π_W) and Tajima's D (Taj$_D$) are shown as 2000 bp windowed averages across each chromosome. F_{ST} and π_B are shown as purple lines to indicate that values were calculated as a comparison between allopatric yellowhammers and pine buntings. π_W and Taj$_D$ are shown as two separate lines (yellow = yellowhammers, brown = pine buntings) to indicate that values were calculated separately for each population.

Figure 6. A) A histogram of average relative differentiation (F_{ST}) values calculated for 2000 bp windows across the nuclear genome when comparing allopatric yellowhammers (n=53) with allopatric pine buntings (n=42). **B)** A truncated version of Figure 6A that depicts the high F_{ST} tail of the histogram. The limits of the y-axis were set as (0,50) with the double dotted line indicating bars of the histogram that have been truncated. **C)** Mean absolute within-group nucleotide variation (π_W) of allopatric yellowhammers and allopatric pine buntings plotted against absolute between-group nucleotide differentiation (π_B). Each dot represents the average value taken from a 2000 bp window of sequenced data across the nuclear genome. The black line indicates where mean within-group nucleotide variation equals between-group nucleotide differentiation. Increasing values of relative differentiation (F_{ST}) calculated for each window are shown in darker shades of blue. **D)** Association between relative differentiation and absolute between-group nucleotide differenation of allopatric yellowhammers and allopatric pine buntings. Each black dot represents average values calculated from a 2000 bp window of sequenced data. A cubic spline fit between the variables is shown as a purple line.