# Mitonuclear co-introgression opposes genetic differentiation between phenotypically divergent songbirds

Ellen Nikelski[1, *], Alexander S. Rubtsov[2], and Darren Irwin[1]

[1] Department of Zoology, and Biodiversity Research Centre, 6270 University Blvd., University of British Columbia, Vancouver, BC, Canada

[2] State Darwin Museum, Moscow, Russia

[*] Present address: Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

Running Title:

Co-introgression opposes differentiation

Corresponding Author:

Ellen Nikelski, Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada. Email: ellen.nikelski@mail.utoronto.ca

Keywords:

mitonuclear co-introgression, mtDNA, mitonuclear gene, genetic differentiation, chromosome Z, Aves

*Abstract*

Comparisons of genomic variation among closely related species often show more differentiation

in mitochondrial DNA (mtDNA) and sex chromosomes than in autosomes, a pattern expected

due to the relative effective population sizes of these genomic components.  Differential

introgression can cause some species pairs to deviate dramatically from this pattern. The

yellowhammer (*Emberiza citrinella*) and the pine bunting (*E. leucocephalos*) are hybridizing

avian sister species that differ greatly in appearance but show no mtDNA differentiation. This

discordance might be explained by mtDNA introgression—a process that can select for co-

introgression at nuclear genes with mitochondrial functions (mitonuclear genes). We investigated

genome-wide nuclear differentiation between yellowhammers and pine buntings and compared it

to what was seen previously in the mitochondrial genome. We found clear nuclear differentiation

that was highly heterogeneous across the genome, with a particularly wide differentiation peak

on the sex chromosome Z. We further tested for preferential introgression of mitonuclear genes

and detected evidence for such biased introgression in yellowhammers. Mitonuclear co-

introgression can remove post-zygotic incompatibilities between species and may contribute to

the continued hybridization between yellowhammers and pine buntings despite their clear

morphological and genetic differences. As such, our results highlight the potential ramifications

of co-introgression in species evolution.

*Introduction*

Evolution in eukaryotes is shaped by changes in multiple genomic components that differ

in their modes of inheritance: mitochondrial DNA (mtDNA) is usually inherited through the

matrilineal line, autosomes are inherited through both parental lines and sex chromosomes are

inherited differentially depending on the sex of both parent and offspring (Avise, 2000). There is

often much variation among these genomic components in the degree of genetic differentiation

between related populations or species (reviewed in Coyne & Orr, 2004; reviewed in Price,

2008), suggesting that their dynamics differ during the process of speciation of a single species

into two or more. This variation can arise through differences in both the rate at which specific

DNA sequences evolve and the degree to which different components contribute towards genetic

incompatibilities that reduce gene flow between populations. A common pattern observed

between speciating taxa is clear differentiation in mtDNA (eg. Hebert et al. 2004; Kerr et al.

2007), moderate differentiation in sex chromosomes (eg. Thornton & Long, 2002; Borge et al.

2005; Lu & Wu, 2005; Harr, 2006; Ruegg et al. 2014; Sackton et al. 2014), and comparatively

modest differentiation across autosomes (Harr, 2006; Nadeau et al. 2012; Irwin et al. 2018).

Measures of mtDNA differentiation are often used to identify and classify genetically

distinct populations (eg. Hebert et al. 2004; Kerr et al. 2007) and to infer their histories (Moore,

1995; Zink & Barrowclough, 2008). Due to its uniparental inheritance, mtDNA has one quarter

the effective population size and coalescence time of autosomal nuclear DNA (Moore, 1995).

This characteristic combined with mtDNA's relatively high mutation rate (Lynch et al. 2006)

mean that genetic differences arise and fix relatively quickly, creating patterns of clear mtDNA

differentiation between recently diverged populations.

Sex chromosomes are another genomic region that often shows higher between-

population genetic differentiation compared to autosomes between speciating taxa, in both Z/W

(Borge et al. 2005; Ruegg et al. 2014; Sackton et al. 2014) and X/Y systems (Thorton & Long,

2002; Lu & Wu, 2005; Harr, 2006). To explain this "faster Z/X effect," researchers have noted

that, because beneficial recessive mutations on the Z or X chromosome are immediately exposed

to selective forces in the heterogametic sex, fixation of these mutations should proceed faster

than if the mutations appeared on autosomes (reviewed in Meisel & Connallon, 2013; Irwin,

2018). Also contributing to genetic differentiation on the Z and X chromosomes are the lower

effective population sizes of these chromosomes compared to autosomes (Mank et al. 2010;

reviewed in Irwin, 2018). A lower effective population size allows for the fixation of a greater

number of slightly deleterious mutations due to less effective purifying selection and a larger role

of genetic drift. It is likely that both forces—the faster Z/X effect and less effective purifying

selection—contribute to the moderate amount of genetic differentiation seen between the sex

chromosomes of diverging taxa (Thorton & Long, 2002; Borge et al. 2005; Lu & Wu, 2005;

Harr, 2006; Ruegg et al. 2014; Sackton et al. 2014).

Differentiation across autosomes, which tends to be lower than on mtDNA and sex

chromosomes, can be highly heterogeneous. In fact, many researchers report "islands of

differentiation" on autosomes where peaks of high relative differentiation are found against a

background of low relative differentiation (e.g., Harr, 2006; Nadeau et al. 2012; Hejase et al.

2020). Explanations for these "islands" usually invoke reduced gene flow (reviewed in Wu,

2001) and/or repeated bouts of selection (Cruickshank and Hahn, 2014; Irwin et al. 2018). In the

former scenario, differentiation peaks are hypothesized to house the loci responsible for

reproductive barriers between interacting taxa and, as a result, they are resistant to the gene flow

that homogenizes the rest of the nuclear genome. In contrast, explanations invoking repeated

selection hypothesize that differentiation islands are areas of the genome that experienced

repeated reductions in genetic diversity as a result of selection or selective sweeps in both

ancestral and daughter populations.

Despite the general patterns of differentiation discussed above, an increasing number of

studies report remarkably low differentiation between populations at what are normally highly

divergent genetic components when compared to other genetic regions or observable phenotypes

(e.g., Irwin et al. 2009; Yannic et al. 2010; Bryson et al. 2012). In a number of cases, mtDNA

shows dramatically low differentiation when compared to differentiation of the nuclear genome,

a pattern referred to as "mitonuclear discordance" (reviewed in Toews & Brelsford, 2012).

Discordance between marker types may be explained by hybridization and introgression between

populations, perhaps due to a selective advantage of the introgressing genetic region. For

example, Hulsey et al. (2016) documented low mtDNA differentiation—likely due to

introgression—and clear differentiation in nuclear DNA (nucDNA) between two hybridizing

cichlid species (Hulsey & García de León, 2013). The researchers further reported high mtDNA

differentiation between isolated populations of cichlids at genetic sites associated with thermal

tolerance and a significant correlation between mtDNA divergence and water temperature

(Hulsey et al. 2016). Altogether, these results suggest that mtDNA introgression produced the

discordance seen between marker types and that this outcome was potentially driven by adaptive

selection for tolerance of extreme water temperatures.

The hypothesis of adaptive introgression increases in complexity if we consider the

potential for coevolution between genomic components. Research investigating coevolution

between mitochondrial and nuclear genomes is relatively novel as mtDNA was often treated as a

neutral marker in past evolutionary research (Avise, 2000). Nevertheless, recent empirical and

theoretical work has provided greater context regarding how mitonuclear coevolution may

influence the progression of differentiation and speciation between taxa (Hill, 2019).

Mitonuclear ecology is the study of how forces acting on the mitochondrial and nuclear

genomes interact to influence ecological and evolutionary processes (Hill, 2019). Best known for

aerobic respiration, the mitochondrion is the site of the electron transport chain (ETC)

responsible for oxidative phosphorylation in eukaryotic organisms (reviewed in Ernster &

Schatz, 1981). Due to the mitochondrial genome's reduced size of about 37 genes, proper

functioning of the ETC as well as transcription, translation and replication of mtDNA is reliant

on about 1500 proteins encoded by mitonuclear genes within the nuclear genome (Calvo &

Mootha, 2010; Lotz et al. 2014). This interplay between mtDNA and nucDNA implies

coevolution between the two genomes such that changes in one places selection on the other for

compensatory changes that reduce genetic incompatibility and maintain mitochondrial function

(Gershoni et al. 2009; Burton & Barreto, 2012; Hill, 2019). Secondary contact and hybridization

between differentiated populations can result in hybrid breakdown due to mismatches between

coevolved combinations of mtDNA and mitonuclear genes, contributing to reproductive isolation

and eventual speciation of the groups. Yet, introgression of an adaptive mitochondrial haplotype

from one population into another could select for similar introgression of compatible alleles at

mitonuclear genes (eg. Beck et al. 2015; Morales et al. 2018; Wang et al. 2021). Such co-

introgression would act as a homogenizing force and decrease genetic differentiation between

populations in both the mitochondrial and nuclear genomes.

The potential for introgression between speciating populations depends on the degree to

which they are in geographic contact during their period of differentiation. In many cases,

closely related species occur in allopatry, but in others, taxa meet in areas of contact where they

may interbreed (reviewed in Coyne & Orr, 2004; reviewed in Price, 2008). Successful

hybridization and backcrossing allow for the introduction of genetic variation from one

population into the other and for adaptive introgression or co-introgression between taxa at

particular genomic regions which can create the discordant differentiation patterns noted in many

systems (reviewed in Toews & Brelsford, 2012).  To look for evidence of adaptive co-

introgression, we must therefore examine systems where two species are or have hybridized

previously and where different genomic components show discordant patterns of genetic

differentiation.

An example of one such system includes the yellowhammer (Passeriformes:

Emberizidae: *Emberiza citrinella*) and the pine bunting (*Emberiza leucocephalos*)—an avian

species pair thought to have diverged during the Pleistocene glaciations when they were

separated on either side of Eurasia by an area of unsuitable habitat (Irwin et al. 2009). These taxa

are highly divergent in plumage and moderately divergent in song and ecology (Panov et al.

2003; Rubtsov & Tarasov, 2017). Despite their differences, yellowhammers and pine buntings

hybridize extensively in a large contact zone in central and western Siberia (Panov et al. 2003;

2007; Rubtsov, 2007; Irwin et al. 2009; Rubtsov & Tarasov, 2017). Genomic work has identified

mitonuclear discordance between allopatric yellowhammers and pine buntings (Irwin et al. 2009)

as they possess almost no mtDNA divergence but show moderate differentiation in nuclear

AFLP (Amplified Fragment Length Polymorphism) markers. To explain these results, Irwin et

al. (2009) suggested that mtDNA may have introgressed from one species into the other during a

previous selective sweep, and this hypothesis was supported by several statistical tests performed

on the mtDNA haplotype network. Such mtDNA introgression could select for co-introgression

at mitonuclear genes if sizeable genetic differentiation had developed between yellowhammers

and pine buntings (Sloan et al. 2017; Hill, 2019; Wang et al. 2021) which is implied by their

divergent behaviour, ecology and appearance. Mitonuclear co-introgression and the resulting

lack of mitonuclear incompatibility could facilitate the continued hybridization seen between

yellowhammers and pine buntings and prevent the build-up of reproductive isolation and full

speciation between these taxa (Hill, 2019).

Here we present the first largescale comparison of DNA sequence variation across the

nuclear genomes of allopatric yellowhammers and pine buntings. We address several key

questions regarding genetic differentiation in this system. First, does nucDNA differentiation

between yellowhammers and pine buntings resemble that of mtDNA (virtually none), that of

plumage phenotype (very strong differentiation), or something in between? The earlier AFLP

results suggested clear differentiation of nuclear markers between groups (Irwin et al. 2009), but

those results were not based on actual DNA sequences. Substantial nuclear differentiation would

provide stronger support for the hypothesis that there has been mitochondrial introgression and

replacement. Second, what is the structure of differentiation across the nuclear genome? The

degree of heterogeneity in differentiation across the genome can be used to test whether adaptive

introgression may have occurred in this system, and whether certain regions of the genome (e.g.,

the sex chromosomes) may be especially important during divergence. Third, is there an over-

representation of known mitonuclear genes within putatively introgressing nucDNA regions–a

pattern consistent with mitonuclear co-introgression? Evidence of mitonuclear co-introgression

and the resulting loss of mitonuclear incompatibility could offer a possible explanation for the

extensive hybridization seen between yellowhammers and pine buntings as well as implicate this

process as a homogenizing force that counters divergence between these groups.

***Materials and Methods***

**Sampling**

We included 109 blood and tissue samples in this study: 53 phenotypic yellowhammers,

42 phenotypic pine buntings, and 14 other members of Emberizidae (one *Emberiza aureola*

[yellow-breasted bunting], one *Emberiza calandra* [corn bunting], one *Emberiza cioides*

[meadow bunting], one *Emberiza hortulana* [ortolan bunting], four *Emberiza stewarti* [white-

capped bunting], and six *Emberiza cirlus* [cirl bunting]) to put variation between yellowhammers

and pine buntings into a deeper phylogenetic context (Figure 1A; Table 1; Supplementary Table

1). A total of 91 samples were included in the AFLP analysis of Irwin et al. (2009) while 18

samples were examined for the first time as part of the present research.

When possible, body measurements and photographs were taken of live birds or museum

skins. Yellowhammer and pine bunting males were also scored phenotypically and sorted into

phenotypic classes based on the protocols presented in Panov et al. (2003) and Rubtsov &

Tarasov (2017). Briefly, each male received a score from 0-7 for background plumage colour,

the amount of chestnut plumage (vs. yellow or white) at the brow and the amount of chestnut

plumage (vs. yellow or white) at the throat. For background colour, birds were assessed on the

strength of yellow—ranging from bright yellow to pure white—in head and body plumage that

did not show brown or black streaking. Phenotypic scores of 0 are consistent with a

phenotypically pure yellowhammer and scores of 7 are consistent with a phenotypically pure

pine bunting. Phenotypic classes included: pure citrinella (PC), almost citrinella (SC), citrinella

hybrid (CH), yellow hybrid (YH), white hybrid (WH), leucocephalos hybrid (LH), almost

leucocephalos (SL) and pure leucocephalos (PL) (Rubtsov & Tarasov, 2017). Unless stated

otherwise, any SC and SL individuals that appeared in the allopatric zones were grouped

together with PC and PL individuals respectively and treated as phenotypic yellowhammers and

phenotypic pine buntings in subsequent analyses (Figure 1B).

**DNA extraction and genotyping-by-sequencing**

DNA was extracted from samples using a standard phenol-chloroform method. We then

divided the DNA samples into four genotyping-by-sequencing (GBS) libraries (Elshire et al.

2011). The 109 samples included in this study were sequenced in libraries together with 226

yellowhammer, pine bunting and hybrid DNA samples collected near and within the sympatric

zone as part of a larger project (Nikelski et al. in prep). The libraries were prepared as per the

protocol described by Alcaide et al. (2014) with the modifications specified by Geraldes et al.

(2019) except that we maintained a 300-400 bp fragment size during size selection. Paired-end

sequencing was completed by Genome Québec using the Illumina HiSeq 4000 system,

producing more than 1.2 billion reads, 150 bp in length, across the four GBS libraries.

**Genotyping-by-sequencing data filtering**

Processing of GBS reads for the samples analyzed in this study was done in conjunction

with reads from the samples included in the larger project mentioned above. We processed the

reads following Irwin et al. (2016; 2018), as summarized here. Reads were demultiplexed using a

custom perl script designed by Baute et al. (2016). Next, reads were trimmed for quality using

Trimmomatic version 0.36 (Bolger et al. 2014) with the parameters: TRAILING:3,

SLIDINGWINDOW:4:10, MINLEN:30. Trimmed reads were aligned to the zebra finch

reference genome (*Taeniopygia guttata* version 3.2.4; Warren et al. 2010) using the program

BWA-MEM (Li & Durbin, 2009) and a BAM file of this information was created for each

individual using the programs Picard (http://broadinstitute.github.io/picard/) and SAMtools (Li et

al. 2009). Then, BAM files were converted into GVCF files using the HaplotypeCaller command

as part of GATK version 3.8 (McKenna et al. 2010). We processed the resulting GVCF files in

two ways to create 1) a genome-wide "variant site" VCF file containing information only on

variant sites, and 2) a series of chromosome-specific "info site" VCF files containing information

on both variant and invariant sites with sufficient coverage.

To create the genome-wide "variant site" VCF file, we used the GenotypeGVCFs

command in GATK version 3.8 to identify and isolate single nucleotide polymorphisms (SNPs)

among individuals. This command also converted the variant site information into a single VCF

file encompassing the entire nuclear genome. Using a combination of VCFtools (Danecek et al.

2011) and GATK, we filtered the VCF file to remove indels and non-biallelic SNPs. We also

discarded loci with QD < 2.0, MQ < 40.0, FS > 60.0, SOR > 3.0, or ReadPosRankSum < -8.0.

Finally, loci with more than 60% missing genotypes were removed. The average coverage of

variable sites in the resulting VCF file was 16.59.

To convert GVCF files into "info site" VCF files, we similarly employed the

GenotypeGVCFs command in GATK with the addition of the -allSites and -L flags to retain

invariant sites and split the information into chromosome-specific files. The resulting VCF files

were filtered using VCFtools and GATK to remove indels, sites with more than two alleles, sites

with more than 60% missing genomic data, sites with MQ values lower than 20 and sites with

heterozygosities greater than 60% (to avoid potential paralogs). Use of these filters simplified

calculations in downstream analyses and ensured that these analyses were restricted to sites with

sufficient data.

All following statistical analyses were completed using R version 3.6.2 (R Core Team,

2014).

**Variant site analyses**

The genome-wide "variant site" VCF file was analyzed using modified versions of the R scripts described in Irwin et al. (2018). A total of 374,780 SNPs were identified among allopatric yellowhammers and pine buntings. For each of these SNPs we calculated sample size, allele frequency, and Weir and Cockerham's $F_{ST}$ (Weir & Cockerham, 1984). Genetic differentiation between yellowhammers and pine buntings was then visualized using a principal components analysis (PCA) generated with the pca command and the svdImpute method to account for any missing genomic data using the pcaMethods package (Stacklies et al. 2007). The PC1 loadings were also graphed as a Manhattan plot using the package qqman (Turner, 2018). Finally, to examine the spread of variant sites across the genome and identify areas of high differentiation, the $F_{ST}$ values of 349,807 SNPs were graphed as a Manhattan plot. The remaining SNPs did not possess known genomic locations and, therefore, could not be included in the plot.

**Differentiation across the genome**

To thoroughly investigate genomic differentiation between allopatric yellowhammers and pine buntings, we performed further analysis on both variant and invariant loci within "info site" VCF files using R scripts described in Irwin et al. (2018).

We calculated Weir and Cockerham's $F_{ST}$, between-group nucleotide distance ($\pi_B$) and within-group nucleotide diversity ($\pi_W$) for nonoverlapping windows of available sequence data across each chromosome. The first window was positioned at the "start" of each chromosome as described in the zebra finch reference genome (Warren et al. 2010). We used a window size of 2000 bp of sequenced data rather than 10,000 bp (as in Irwin et al. 2018), to visualize narrow peaks in relative and absolute differentiation within our dataset.

We developed a new R script to calculate a Tajima's D value (Tajima, 1989) for each of

the 2000 bp windows. Values of Tajima's D were used to identify areas of the genome where

patterns of variation in yellowhammer and pine bunting populations deviated from models of

neutrality. Significantly negative Tajima's D implies that there are more rare alleles in a

population than expected under neutrality, likely because of a selective sweep or population

expansion following a bottleneck. Significantly positive Tajima's D suggests that there are fewer

rare alleles in a population than expected under neutrality, potentially stemming from balancing

selection or rapid population contraction.

**Phylogenetic comparison with other Emberizidae species**

We employed whole-genome averages of $\pi_B$ between allopatric yellowhammers and

allopatric pine buntings as well as among these focal species and six other Emberizidae species

(*Emberiza aureola, Emberiza calandra, Emberiza cioides, Emberiza cirlus, Emberiza hortulana*

*and Emberiza stewarti*) to estimate a phylogeny. A list of average $\pi_B$ values for each species pair

was converted into a distance matrix and used to create an unrooted neighbour-joining tree. This

tree was constructed using the ape package (Paradis & Schliep, 2019) and the BioNJ algorithm

(Gascuel, 1997) with *Emberiza aureola* set as the outgroup (Alström et al. 2008).

**Signals of mitonuclear co-introgression**

To test for signals of mitonuclear gene introgression between allopatric yellowhammers

and pine buntings, we compiled a list of mitonuclear genes and a list of 2000 bp putative

introgression windows (hereafter referred to as "introgression windows") and then tested for an

association between them. We explain these steps in detail below. If mitonuclear genes were

found within introgression windows statistically more often than was expected due to chance, it

would provide support for mitonuclear gene introgression potentially in response to the adaptive

mtDNA introgression (Hill, 2019) hypothesized to have occurred in this system (Irwin et al.

2009).

To start, we created a list of mitonuclear genes to analyze for signals of introgression. We

chose mitonuclear genes that were protein-coding and whose protein products interacted directly

with mtDNA or an immediate product of the mitochondrial genome (i.e. protein or RNA). For

these nuclear-encoded genes, any change in mtDNA including those caused by introgression

would likely cause selection for co-introgression of compatible alleles (Gershoni et al. 2009;

Burton & Barreto, 2012; Hill, 2019). Mitonuclear genes that met these criteria included those

that encode protein subunits of ATP synthase or the first, third and fourth complex of the ETC,

assembly and ancillary proteins involved in the formation of the ETC, or proteins that are part of

the transcription, translation or DNA replication machinery within the mitochondria. This list of

genes was created using information from Figure 2.3 and Table 2.1 in Hill (2019). After

removing any genes that were not annotated in the zebra finch reference genome or that lacked a

specific location on the reference genome, a total of 134 mitonuclear genes remained for analysis

(Supplementary Table 2).

Next, we identified introgression windows across the genome as those possessing both a

low Tajima's D value and a low $\pi_B$ value. Low $\pi_B$ indicates high similarity between the

nucleotide sequences of two groups as would be expected if mitonuclear gene alleles had

introgressed from one taxon into the other. Low Tajima's D suggests a past selective sweep

within a population which would also be expected if an adaptive mitonuclear allele had

introgressed from a separate taxon and swept throughout the receiving population. For this

analysis, our quantitative criteria for an introgression window were a Tajima's D value within

the lowest 5% of the available windowed values and a $\pi_B$ value within the lowest 30% of the

available windowed values. Out of the 7187 windows described across the genome, 244

introgression windows were identified for yellowhammers and 222 introgression windows were

identified for pine buntings. As well, of the introgression windows identified in yellowhammer

and pine bunting populations, 71 were shared between the taxa. It should be noted that sharing of

some introgression windows is expected given that the contribution of $\pi_B$ to window selection

was identical for both taxa (in contrast, Tajima's D was calculated separately for yellowhammers

and pine buntings).

Following window classification, we employed a custom R script to determine how many

mitonuclear genes occurred within introgression windows. To do this, we identified the genomic

centre position of each mitonuclear gene as well as the average genomic position of each of the

genomic windows. We then calculated the absolute difference between mitonuclear gene centres

and average window positions. Mitonuclear genes were assumed to occur within whatever

window minimized this difference. With this information, we were able to determine the number

of mitonuclear genes that occurred within the introgression windows of each taxon.

The number of mitonuclear genes within introgression windows were compared to what

would be expected if these genes were distributed randomly across the genome using separate

two-tailed binomial tests for yellowhammers and pine buntings. Because genes are often not

distributed randomly and may appear more densely packed in certain genomic regions

(International Chicken Genome Sequencing Consortium, 2004), we also conducted a Fisher's

Exact test for both yellowhammers and pine buntings to determine whether the proportion of

mitonuclear genes within introgression windows was significantly different from what would be

expected based on the total proportion of protein coding genes appearing within these windows.

*Results*

348   When comparing allopatric yellowhammers and pine buntings, following filtering we

identified 374,780 variable SNPs within our "variant site" VCF file and 13,703,455 invariant and

699,122 variant sites across thirty autosomes and the Z chromosome within our "info site" VCF

files. In the latter "info site" files, we designated a total of 7187 genomic windows (of 2000

sequenced bp each) across the genome, with each window covering an average distance of about

139 kilobases.

**Phylogenetic comparison with other Emberizidae species**

355   An unrooted neighbour-joining tree of average $\pi_B$ values between yellowhammers, pine

buntings and six other Emberizidae species (Figure 2) depicted similar species relationships as

were identified previously using mitochondrial markers (Alström et al. 2008; Irwin et al. 2009).

As well, similar relative branch lengths between taxa were recovered with the exception of that

between yellowhammers and pine buntings. In terms of the relative genetic distance ($\pi_B$)

between yellowhammers and pine buntings compared to the distance between each of those and

*E. stewarti*, nuclear genetic distance was 11.4 times greater than mitochondrial genetic distance.

This corroborates the presence of mitonuclear discordance between the taxa where nucDNA is

much more deeply diverged than mtDNA, supporting the hypothesis of an extended period of

divergence between yellowhammers and pine buntings followed by mtDNA introgression.

**Overall genetic differentiation**

366   Based on 374,780 SNPs considered all together, our genome-wide $F_{ST}$ estimate was

0.0232 between allopatric yellowhammers and pine buntings. Despite this low average, a PCA

based on the same SNP genotypes separated yellowhammers and pine buntings into tight genetic

clusters (Figure 3). PC1 explained 3.6% of the variation among individuals while PC2 explained

2.9% of the variation. Two pine buntings were outliers along PC1, while the remaining

yellowhammers and pine buntings separated into distinct clusters mainly along PC2. Further

investigation into these outliers revealed that they were males from the same location. A kinship

analysis completed as part of a separate study did not find close kinship between the two pine

buntings that could explain their position as outliers (Nikelski et al. in prep). An examination of

the PC1 loadings for each of the SNPs revealed that the signal for the PC1 positioning was

broadly distributed across the genome, rather than being concentrated in a few specific regions

(Supplementary Figure 1). To explore the causes of these outliers, we temporarily removed one

of them and re-ran the PCA. This caused the other outlier to fall into the pine bunting cluster, but

did reveal a further yellowhammer outlier (Supplementary Figure 2). Removal of this

yellowhammer outlier in addition to one member of the pine bunting outlier pair in turn revealed

another yellowhammer outlier (Supplementary Figure 3). It is unclear what is responsible for

these outliers, but the distinct yellowhammer and pine bunting genetic clusters remained intact in

all the PCAs considered.

**Differentiation across the genome**

Relative differentiation between allopatric yellowhammers and pine buntings was highly

heterogeneous across the nuclear genome with peaks in $F_{ST}$ seen on most of the larger

chromosomes (Figures 4, 5; Supplementary Figure 4). Chromosome Z in particular showed a

large peak in $F_{ST}$ with several SNPs possessing values close to one. In fact, $F_{ST}$ for the Z

chromosome was 0.1246—more than five times larger than the genome-wide $F_{ST}$.

Patterns of between-group nucleotide diversity ($\pi_B$) and within-group nucleotide

diversity ($\pi_W$) were also heterogenous across the genome and comparable to each other in

magnitude: genome-wide $\pi_B$ = 0.0041; genome-wide $\pi_W$ for both taxa= 0.0040 (Figure 5;

Supplementary Figure 4). Because between-group and within-group nucleotide diversity are so

intimately related in their evolution and calculation, it is expected that windowed averages of

these two statistics will show a highly positive relationship. In fact, most windowed $\pi_B$ and $\pi_W$

averages fell near a 1:1 association line (Figure 6) which is equivalent to no or little

differentiation. However, some genomic windows showed much reduced $\pi_W$ compared to $\pi_B$;

these were the windows with high $F_{ST}$. Additionally, we detected a weak negative correlation

between the windowed averages of $F_{ST}$ and $\pi_B$ (Spearman's Rank Correlation: -0.1196, p < 2.2 ×

$10^{-16}$; Figure 7) as is hypothesized if peaks in relative differentiation are products of repeated

selective events (Cruickshank & Hahn, 2014; Irwin et al. 2018).

Finally, we found that Tajima's D varied across the genome but was mostly negative

(Figure 5; Supplementary Figure 4), consistent with a history of population growth and/or

selective sweeps. The average genome-wide Tajima's D was similar between populations: -1.377

for yellowhammers and -1.335 for pine buntings.

**Signals of mitonuclear co-introgression**

Of the 7187 genomic windows identified across the nuclear genome, we classified 244 as

introgression windows within yellowhammers and 222 as introgression windows within pine

buntings (Table 2). Average values of $\pi_B$ and Tajima's D in yellowhammer introgression

windows were 0.0016 and -2.3751 respectively, and 0.0019 and -2.3369 in pine buntings

respectively.

Nine mitonuclear genes—6.7% of the 134 mitonuclear genes considered—appeared

within yellowhammer introgression windows (Table 2). This finding was significant in a two-

tailed binomial test (p = 0.04952) indicating that mitonuclear genes appeared in yellowhammer

introgression windows more often than would be expected if they were assigned to windows

randomly. However, this finding was not significant in a Fisher's Exact test (p = 0.1311) which

takes into account the differing densities of genes across the nuclear genome. Four mitonuclear

genes appeared within pine bunting introgression windows—3.0% of the genes considered. This

result was statistically insignificant in both a two-tailed binomial test (p = 1) and a Fisher's Exact

test (p = 1) indicating that mitonuclear genes did not appear in pine bunting introgression

windows more often than would be expected due to chance. Overall, the significant signal of

introgression in yellowhammers and insignificant signal of introgression in pine buntings could

indicate that mitonuclear gene introgression—if it occurred—was biased in the direction of pine

buntings into yellowhammers.

The nine mitonuclear genes that appeared within yellowhammer introgression windows

are: APOPT1, COX5A, COX17, MRPL1, MRPL27, MRPL32, NDUFC1, mtSSB, UQCR11

(Table 3). Three of these genes encode protein subunits of the mitochondrial ribosome, three

encode structural subunits of the ETC, two encode assembly factors of the ETC and one encodes

a single-stranded DNA-binding protein involved in mtDNA replication. All putatively

introgressed genes appear on separate autosomes except for two genes that appear on

chromosome 4. Interestingly, three of the five putatively introgressed genes associated with the

ETC are specifically associated with complex IV.

The four mitonuclear genes that appeared within pine bunting introgression windows are:

ATP5H5I, COX5A, MRPL2 and NDUFB4 (Table 4). All four genes appeared on separate

autosomes with three of these genes encoding structural subunits of the ETC and one encoding a

protein subunit of the mitochondrial ribosome. The COX5A gene, which encodes a structural

subunit of ETC complex IV, was found in both yellowhammer and pine bunting introgression

windows.

***Discussion***

Yellowhammers and pine buntings show negligible mtDNA differentiation (Irwin et al.

2009) but are well differentiated phenotypically (Panov et al. 2003, Rubtsov & Tarasov, 2017).

Prior to this study, a possible explanation for this pattern was simply rapid phenotypic evolution

between two genetically similar sister taxa. Our analysis of nucDNA variation has shown clear

separation of allopatric yellowhammers and pine buntings, with strong differentiation peaks in

specific parts of the nuclear genome. This result points to these taxa experiencing a long period

of separate evolution followed by the hybridization now observed within a large contact zone in

western and central Siberia. These results combined with our phylogenetic analysis showing a

longer branch length between yellowhammers and pine buntings based on nuclear markers—

when compared to a phylogeny based on mtDNA (Irwin et al. 2009)—support recent mtDNA

introgression and mitochondrial haplotype replacement in this system likely driven by selection.

Our analyses also provided some evidence for the disproportionate introgression of mitonuclear

genes between taxa which is consistent with co-introgression discussed in current mitonuclear

theory (Hill, 2019).

Though genetically distinct, the genome-wide $F_{ST}$ between allopatric yellowhammers and

pine buntings (0.0232) was comparable to or sometimes lower than the averages seen between

avian subspecies (e.g., subspecies of barn swallow: 0.017-0.026 [Scordato et al. 2017]; myrtle

warbler and Audubon's warbler: 0.077-0.106 [Irwin et al. 2018]; yellow- and red-shafted

northern flickers: 0.098 [Manthey et al. 2016]). This low genome-wide $F_{ST}$ contrasts with the

moderate $F_{ST}$ averages reported from an analysis of 63 AFLP markers performed on the same

populations: 0.078 based on allele frequencies and 0.140 based on band frequencies (Irwin et al.

2009). The present study reveals that relative differentiation was highly heterogeneous across the

nuclear genome with $F_{ST}$ peaks on various chromosomes. It is possible that the previous AFLP

analysis captured a disproportionate number of loci within these differentiation peaks, thereby

inflating $F_{ST}$ estimates. This comparison highlights the caution that should be taken when

interpreting genome-wide averages as highly variable genetic differentiation landscapes can

cause large variability in $F_{ST}$ estimates when they are based on a limited number of loci.

The $F_{ST}$ peaks seen between yellowhammers and pine buntings on larger autosomes and

most significantly on the Z chromosome are consistent with the "islands of differentiation" often

noted in comparisons of other closely related taxa (Harr, 2006; Nadeau et al. 2012; Irwin et al.

2018). In contrast to these islands, the large regions of close similarity in $\pi_B$ and $\pi_W$ suggests

high gene flow between taxa at those regions. This scenario is consistent with the observed

extensive hybridization between these taxa (Panov et al. 2003; 2007; Rubtsov, 2007; Rubtsov &

Tarasov, 2017). Nevertheless, the high $F_{ST}$ regions, those with much reduced $\pi_W$ compared to

$\pi_B$, indicate that they have had low gene flow presumably as a result of divergent selection. It is

unlikely that this pattern can be explained by genetic drift over an extended period of geographic

separation, as this would result in most genomic regions deviating slightly from $\pi_B = \pi_W$

congruence rather than the observed pattern of extreme heterogeneity. Instead, the pattern

suggests that selection acted in a way that lowered $\pi_W$ relative to $\pi_B$ within "islands of

differentiation". Considering that high $F_{ST}$ regions were associated with relatively low values of

$\pi_B$, we propose that differentiation islands in this system are most consistent with a model

invoking repeated bouts of selection that lower nucleotide diversity (Cruickshank & Hahn, 2014;

Irwin et al. 2018). A sweep-before-differentiation model (Irwin et al. 2018) where $F_{ST}$ peaks are

produced by adaptive selective sweeps between populations followed by adaptive selection at the

same regions in local populations is particularly in line with the extensive hybridization presently

observed between yellowhammers and pine buntings.

Of the "islands of differentiation" identified between taxa, the tallest and widest was

found on the Z chromosome. Greater differentiation on the Z chromosome compared to

autosomes is a common observation when comparing closely related species (Borge et al. 2005;

Ruegg et al. 2014; Sackton et al. 2014) and is consistent with the "faster Z/X effect" that has

been explained by less efficient purifying selection and/or more positive selection on this

chromosome (Mank et al. 2010; reviewed in Meisel & Connallon, 2013; reviewed in Irwin,

2018). However, the large regions of the Z chromosome that have $F_{ST}$ values near zero suggests

that additional factors are involved in producing the large and wide island of differentiation on

the Z.

One possible explanation for a large differentiation island could be that it corresponds

with an area of low recombination—a region of connected loci that tend to be inherited together,

leading to linked selection of nearby loci. Strong divergent selection acting on one SNP would

act similarly on all the loci that are linked to it such that a wide, highly divergent genomic block

would become fixed and appear as an "island" between taxa (reviewed in Cutter & Payseur,

2013). Areas of low recombination and linkage are often associated with inversion

polymorphisms (reviewed in Smukowski & Noor, 2011) as different orientations of an inversion

experience little successful recombination (reviewed in Kirkpatrick, 2010). An exploration of a

potential chromosomal inversion within the yellowhammer and pine bunting system is being

investigated as part of a separate study and is supported by preliminary evidence (Nikelski et al.

in prep).

While numerous "islands of differentiation" were observed between yellowhammers and

pine buntings implying moderate genetic divergence between them, mtDNA introgression has

the potential to homogenize their nuclear genomes at mitonuclear genes by selecting for co-

introgression of compatible alleles (Beck et al. 2015; Sloan et al. 2017; Morales et al. 2018).

Consistent with this idea, a two-tailed binomial test supported preferential introgression of

mitonuclear genes in allopatric yellowhammers. Because a comparable signal of introgression

was not found in allopatric pine buntings, we suggest that mitonuclear co-introgression could

have occurred in the direction of pine buntings into yellowhammers. Yet, these results must be

interpreted with caution due to limitations in introgression window identification. Because we

employed reduced-representation sequencing that only captures a small portion of the nuclear

genome, we have limited resolution in detecting signals of introgression over narrow genomic

regions. This may have contributed to the variation we saw in the statistical significance of

mitonuclear gene introgression depending on whether gene densities were considered.

Nevertheless, the fact that a significant signal of introgression was detected despite limitations is

intriguing especially when considered in conjunction with the identities of the mitonuclear genes

found in introgression windows.

Three of the mitonuclear genes within yellowhammer introgression windows and three

within pine bunting introgression windows encoded structural subunits of the ETC. The ETC is

broken into five protein complexes which, through a series of enzymatic reactions, perform

oxidative phosphorylation to produce ATP necessary for organism survival (reviewed in Ernster

& Schatz, 1981). Four of the five ETC complexes are made up of subunits encoded by both the

nuclear and mitochondrial genome (Hill, 2019) and correct fit between differentially encoded

subunits is essential for the flow of electrons and protons across the ETC. To put this in

perspective, changing even a single amino acid in one subunit can significantly disrupt its ability

to interact with other subunits within a ETC complex (eg. Gershoni et al. 2014). Because of the

tight interactions within complexes and the consequences of subunit incompatibility,

introgression of mtDNA is expected to select for co-introgression of mitonuclear genes encoding

ETC structural subunits. Such co-introgression has been detected between differentially adapted

populations of eastern yellow robin where mtDNA introgression between populations was

followed by similar introgression of mitonuclear genes encoding subunits of complex I (Morales

et al. 2018) and between different species of *Drosophila* where introgression and replacement of

the mtDNA of one species during hybridization selected for co-introgression of genes that

encode subunits of complex IV (Beck et al. 2015).

Of the ETC complexes, complex IV showed the strongest signal of co-introgression in

the yellowhammer and pine bunting system. Three of the genes within yellowhammer

introgression windows and one gene within pine bunting introgression windows were associated

with this complex. Interestingly, gene COX5A—a structural subunit of complex IV—appeared

in both sets of introgression windows. It is unlikely that this gene introgressed in both directions,

but it is possible that COX5A adaptively swept in both populations. In this situation, a

particularly adaptive allele may have appeared in one species and swept to high frequency before

co-introgressing into the other species following mtDNA introgression. The COX5A gene was

also one of the subunits that co-introgressed in the *Drosophila* example discussed above (Beck et

al. 2015) lending some support to its particular importance to mitonuclear compatibility. More

generally, complex IV is often used as a model for studying mitonuclear interactions due to its

distinctive structure of a core of mtDNA-encoded subunits surrounded by nucDNA-encoded

subunits (Saraste, 1999). With such an excess of mitonuclear interactions, incompatibility

involving complex IV has been investigated and detected in several systems including within

primate xenomitochondrial cybrids (Barrientos et al. 2000) and between different species of

*Drosophila* (Sackton et al. 2003). Furthermore, work by Osada & Akashi (2012) has provided

strong evidence for compensatory co-evolution between mitonuclear genes related to complex

IV and mtDNA among primates particularly at interacting amino acids of differentially encoded

subunits. Altogether, these results suggest a crucial role for complex IV in mitonuclear co-

evolution as it may relate to divergence and speciation between taxa.

Another group of mitonuclear genes that appeared to preferentially introgress within the

yellowhammer and pine bunting system were those encoding subunits of the mitoribosome.

Unlike the protein-protein interactions occurring within ETC complexes, mitonuclear

interactions in the mitoribosome are between nuclear-encoded proteins and mitochondrial-

encoded RNA (Hill,. 2019). Protein subunits associate closely with rRNA during the formation

of a mitoribosome, but also interact with mRNA and tRNA during the synthesis of mitochondrial

proteins (Greber & Ban, 2016). Currently, research is limited on the extent and importance of

interactions between mitoribosomal subunits and mitochondrial RNA. However, the fact that

interactions between components are extensive and necessary for the synthesis of the

mitochondrial proteins suggests close co-evolution between mtDNA and genes encoding

mitoribosomal subunits that could strongly select for mitonuclear co-introgression following

mtDNA introgression.

In summary, yellowhammers and pine buntings are sister taxa that are divergent in

appearance, song, and ecology (Panov et al. 2003; Rubtsov & Tarasov, 2017) yet vary in their

genomic differentiation from virtually none (at the mitochondrial genome) to nearly fixed (the

differentiation peak on the Z chromosome). These patterns are best explained by a period of

differentiation while geographically separated, followed by hybridization and introgression. We

found some evidence of mitonuclear gene introgression in the direction of pine buntings into

yellowhammers that is consistent with mitonuclear co-introgression. This occurred preferentially

in mitonuclear genes encoding structural components of both the ETC and the mitoribosome,

potentially due to mitonuclear incompatability. Mitonuclear incompatibilities are thought to

represent an important post-zygotic reproductive barrier between taxa (Gershoni et al. 2009;

Burton & Barreto, 2012; Hill, 2019), meaning mitonuclear co-introgression has the potential to

weaken species boundaries. Support for such breakdown may be seen in the extensive and

dynamic hybrid zone between yellowhammers and pine buntings (Panov et al. 2003; 2007;

Rubtsov, 2007; Rubtsov & Tarasov, 2017). Further, careful examination of genetic

differentiation and reproductive barriers within the yellowhammer and pine bunting hybrid zone

would shed light on the possibility of their merging in the future. As well, the inclusion of

analyses that compare mtDNA and mitonuclear gene differentiation in a wider range of systems

would help to clarify the potentially important role that mitonuclear interactions play in the

merging or diverging of species.

***Author contributions:***

E. N., D.I., and A.S.R. conceived of this study. A.S.R. collected samples. E. N. and A.S.R.

completed molecular techniques. E.G.M.N. conducted data analysis and wrote this manuscript

with input from D.I. and A.S.R.

*Acknowledgements:*

For providing valuable feedback, we thank Dolph Schluter, Eric Taylor, Judith Mank, Elizabeth

Natola, Rashika Ranasinghe, Kenneth Askelson, Finola Fogarty, Quinn McCallum, Ana

Barreira, Jamie Clarke, Armando Geraldes and Jessica Irwin. For their kindness and support

during field work, we thank the Tazeev family and Madelyn Ore. For providing additional

samples, we thank The Bell Museum, The Burke Museum of Natural History and Culture, The

Field Museum, The State Darwin Museum, The Swedish Museum of Natural History, The

Zoological Museum of the Zoological Institute of the Russian Academy of Sciences, the

Zoological Museum of the University of Copenhagen and their accompanying personnel. Major

research funding was provided by the Natural Sciences and Engineering Research Council of

Canada (NSERC CGSM award to E.N, Discovery Grants RGPIN-2017-03919 and RGPAS-

2017-507830 awarded to D.I.) and by the Werner and Hildegard Hesse research awards

(Research award in Ornithology and Fellowship in Ornithology awarded to E.G.M.N. by the

University of British Columbia).

*Data Accessibility Statement:*

Raw DNA sequencing reads will be made available on the NBCI Sequence Read Archive upon

publication acceptance. Read processing codes, barcodes, genotype data and R codes associated

with statistical analyses will be made available on Dryad upon publication acceptance.

*Conflict of Interest Statement:*

None declared

***Bibliography***

Alcaide, M., Scordato, E. S. C., Price, T. D., & Irwin, D. E. (2014). Genomic divergence in a
621           ring species complex. *Nature, 511*(7507), 83-85. doi:10.1038/nature13285

623      Alström, P., Olsson, U., Lei, F., Wang, H., Gao, W., & Sundberg, P. (2008). Phylogeny and
624           classification of the old world *Emberizini* (*Aves, Passeriformes*). *Molecular*
625           *Phylogenetics and Evolution, 47*(3), 960-973. doi:10.1016/j.ympev.2007.12.007

627      Avise, J. C. (2000). Phylogeography: The history and formation of species. Harvard University
628           Press.

630      Barrientos, A., Müller, S., Dey, R., Wienberg, J., & Moraes, C. T. (2000). Cytochrome c oxidase
631           assembly in primates is sensitive to small evolutionary variations in amino acid sequence.
632           *Molecular Biology and Evolution, 17*(10), 1508-1519.
633           https://doi.org/10.1093/oxfordjournals.molbev.a026250

635      Baute, G. J., Owens, G. L., Bock, D. G., & Rieseberg, L. H. (2016). Genome-wide genotyping-
636           by-sequencing data provide a high-resolution view of wild *Helianthus* diversity, genetic
637           structure, and interspecies gene flow. *American Journal of Botany, 103*(12), 2170-2177.
638           doi:10.3732/ajb.1600295

640      Beck, E. A., Thompson, A. C., Sharbrough, J., Brud, E., & Llopart, A. (2015). Gene flow
641           between *Drosophila yakuba* and *Drosophila santomea* in subunit V of cytochrome c
642           oxidase: A potential case of cytonuclear cointrogression. *Evolution, 69*(8), 1973-1986.
643           doi:10.1111/evo.12718

645      Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina
646           sequence data. *Bioinformatics, 30*(15), 2114-2120. doi:10.1093/bioinformatics/btu170

648      Borge, T., Webster, M. T., Andersson, G., & Saetre, G. (2005). Contrasting patterns of
649           polymorphism and divergence on the Z chromosome and autosomes in two *Ficedula*
650           flycatcher species. *Genetics, 171*(4), 1861-1873. doi:10.1534/genetics.105.045120

Burton, R. S., & Barreto, F. S. (2012). A disproportionate role for mtDNA in Dobzhansky–
        Muller incompatibilities? *Molecular Ecology, 21*(20), 4942-4957.
        doi:10.1111/mec.12006

Bryson, J., Robert W, De Oca, A. N., Jaeger, J. R., & Riddle, B. R. (2010). Elucidation of cryptic
        diversity in a widespread Nearctic treefrog reveals episodes of mitochondrial gene
        capture as frogs diversified across a dynamic landscape. *Evolution, 64*(8), 2315-
        2330. https://doi.org/10.1111/j.1558-5646.2010.01014.x

Calvo, S. E., & Mootha, V. K. (2010). The mitochondrial proteome and human disease. *Annual
        Review of Genomics and Human Genetics, 11*(1), 25-44.
        doi:10.1146/annurev-genom-082509-141720

Coyne, J. A., & Orr, H. A. (2004). *Speciation*. Sinauer Associates.

Cruickshank, T. E., & Hahn, M. W. (2014). Reanalysis suggests that genomic islands of
        speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology,
        23*(13), 3133-3157. doi:10.1111/mec.12796

Cutter, A. D., & Payseur, B. A. (2013). Genomic signatures of selection at linked sites: Unifying
        the disparity among species. *Nature Reviews. Genetics, 14*(4), 262-274.
        doi:10.1038/nrg3425

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R.
        E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R. & 1000 Genomes
        Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics,
        27*(15), 2156-2158. doi:10.1093/bioinformatics/btr330

Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler, E. S., & Mitchell,
        S. E. (2011). A robust, simple genotyping-by-sequencing (GBS) approach for high
        diversity species. *PloS One, 6*(5), e19379. doi:10.1371/journal.pone.0019379

Ernster, L., & Schatz, G. (1981). Mitochondria: A historical review. *The Journal of Cell
        Biology, 91*(3), 227s-255s. doi:10.1083/jcb.91.3.227s

Gascuel, O. (1997). BIONJ: An improved version of the NJ algorithm based on a simple model
of sequence data. *Molecular Biology and Evolution, 14*(7), 685-695.
doi:10.1093/oxfordjournals.molbev.a025808

Geraldes, A., Askelson, K. K., Nikelski, E., Doyle, F. I., Harrower, W. L., Winker, K., & Irwin,
D. E. (2019). Population genomic analyses reveal a highly differentiated and endangered
genetic cluster of northern goshawks (*Accipiter gentilis laingi*) in Haida Gwaii.
*Evolutionary Applications, 12*(4), 757-772. doi:10.1111/eva.12754

Gershoni, M., Levin, L., Ovadia, O., Toiw, Y., Shani, N., Dadon, S., Barzilai, N., Bergman, A.,
Atzmon, G., Wainstein, J., Tsur, A., Nijtmans, L. G. J., Glaser, B., & Mishmar, D.
(2014). Disrupting mitochondrial-nuclear coevolution affects OXPHOS complex I
integrity and impacts human health. *Genome Biology and Evolution, 6*(10), 2665-2680.
https://doi.org/10.1093/gbe/evu208

Gershoni, M., Templeton, A. R., & Mishmar, D. (2009). Mitochondrial bioenergetics as a major
motive force of speciation. *Bioessays, 31*(6), 642-650. doi:10.1002/bies.200800139

Greber, B. J., & Ban, N. (2016). Structure and function of the mitochondrial ribosome. *Annual
Review of Biochemistry, 85*(1), 103-132. doi:10.1146/annurev-biochem-060815-014343

Harr, B. (2006). Genomic islands of differentiation between house mouse subspecies. *Genome
Research, 16*(6), 730-737. doi:10.1101/gr.5045006

Hebert, P. D., Penton, E. H., Burns, J. M., Janzen, D. H., & Hallwachs, W. (2004). Ten species
in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly
*Astraptes fulgerator*. *Proceedings of the National Academy of Sciences*, *101*(41), 14812-
14817. doi:10.1073/pnas.0406166101

Hejase, H. A., Salman-Minkov, A., Campagna, L., Hubisz, M. J., Lovette, I. J., Gronau, I., &
Siepel, A. (2020). Genomic islands of differentiation in a rapid avian radiation have been
driven by recent selective sweeps. *Proceedings of the National Academy of
Sciences*, *117*(48), 30554–30565. doi:10.1073/pnas.2015987117
720

Hill, G. E. (2019). *Mitonuclear Ecology* (First ed.). Oxford University Press.

Hulsey, C. D., Bell, K. L., García-de-León, F. J., Nice, C. C., & Meyer, A. (2016). Do relaxed
selection and habitat temperature facilitate biased mitogenomic introgression in a
narrowly endemic fish? *Ecology and Evolution, 6*(11), 3684- 3698.
https://doi.org/10.1002/ece3.2121

Hulsey, C. D., & García-de-León, F. J. (2013). Introgressive hybridization in a trophically
polymorphic cichlid. *Ecology and Evolution, 3*(13), 4536-4547.
https://doi.org/10.1002/ece3.841

International Chicken Genome Sequencing Consortium. (2004). Sequence and comparative
analysis of the chicken genome provide unique perspectives on vertebrate evolution.
*Nature, 432*(7018), 695-716. doi:10.1038/nature03154

Irwin, D. E. (2018). Sex chromosomes and speciation in birds and other ZW systems. *Molecular*
*Ecology, 27*(19), 3831-3851. doi:10.1111/mec.14537

Irwin, D. E., Alcaide, M., Delmore, K. E., Irwin, J. H., & Owens, G. L. (2016). Recurrent
selection explains parallel evolution of genomic regions of high relative but low absolute
differentiation in a ring species. *Molecular Ecology, 25*(18), 4488-4507.
doi:10.1111/mec.13792

Irwin, D. E., Milá, B., Toews, D. P. L., Brelsford, A., Kenyon, H. L., Porter, A. N., Grossen, C.,
Delmore, K. E., Alcaide, M., & Irwin, J. H. (2018). A comparison of genomic islands of
differentiation across three young avian species pairs. *Molecular Ecology, 27*(23), 4839-
4855. doi:10.1111/mec.14858

Irwin, D. E., Rubtsov, A. S., & Panov, E. N. (2009). Mitochondrial introgression and
replacement between yellowhammers (*Emberiza citrinella*) and pine buntings (*Emberiza*
*leucocephalos*) (Aves: Passeriformes). *Biological Journal of the Linnean Society, 98*(2),
422-438. doi:10.1111/j.1095-8312.2009.01282.x

Kerr, K. C. R., Stoeckle, M. Y., Dove, C. J., Weigt, L. A., Francis, C. M., & Hebert, P. D. N.
(2007). Comprehensive DNA barcode coverage of North American birds. *Molecular*
*Ecology Notes, 7(*4), 535-543. doi:10.1111/j.1471-8286.2007.01670.x

Kirkpatrick, M. (2010). How and why chromosome inversions evolve. *PLoS Biology, 8*(9),
e1000501. doi:10.1371/journal.pbio.1000501

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics, 25*(14), 1754-1760. doi:10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., &
Durbin, R. (2009). The sequence alignment map format and SAMtools. *Bioinformatics,*
*25*(16), 2078-2079. doi:10.1093/bioinformatics/btp352

Lotz, C., Lin, A. J., Black, C. M., Zhang, J., Lau, E., Deng, N., Wang, Y., Zong, N. C., Choi, J.
H., Xu, T., Liem, D. A., Korge, P., Weiss, J. N., Hermjakob, H., Yates, J. R., III.,
Apweiler, R., & Ping, P. (2014). Characterization, design, and function of the
mitochondrial proteome: From organs to organisms. *Journal of Proteome Research,*
*13*(2), 433-446. doi:10.1021/pr400539j

Lu, J., & Wu, C. (2005). Weak selection revealed by the whole-genome comparison of the X
chromosome and autosomes of human and chimpanzee. *Proceedings of the National*
*Academy of Sciences, 102*(11), 4063-4067. doi:10.1073/pnas.0500436102

Lynch, M., Koskella, B., & Schaack, S. (2006). Mutation pressure and the evolution of organelle
genomic architecture. *Science, 311*(5768), 1727-1730. doi:10.1126/science.1118884

Mank, J. E., Nam, K., & Ellegren, H. (2010). Faster-Z evolution is predominantly due to genetic
drift. *Molecular Biology and Evolution, 27*(3), 661-670. doi:10.1093/molbev/msp282

Manthey, J. D., Geiger, M., & Moyle, R. G. (2017). Relationships of morphological groups in
the northern flicker superspecies complex (*Colaptes auratus & C. chrysoides*).
*Systematics and Biodiversity, 15*(3), 183-191.
https://doi.org/10.1080/14772000.2016.1238020

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella,
K., Altshuler, D., Gabriel, S., Daley, M., & DePristo, M. A. (2010). The genome analysis
toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data.
*Genome Research, 20*(9), 1297-1303. doi:10.1101/gr.107524.110

Meisel, R. P., & Connallon, T. (2013). The faster-X effect: Integrating theory and data. *Trends in*
*Genetics, 29*(9), 537-544. doi:10.1016/j.tig.2013.05.009

Moore, W. S. (1995). Inferring phylogenies from mtDNA variation: Mitochondrial-gene trees
        versus nuclear-gene trees. *Evolution, 49*(4), 718-726. doi:10.2307/2410325

Morales, H. E., Pavlova, A., Amos, N., Major, R., Kilian, A., Greening, C., & Sunnucks, P.
        (2018). Concordant divergence of mitogenomes and a mitonuclear gene cluster in bird
        lineages inhabiting different climates. *Nature Ecology & Evolution, 2*(8), 1258-1267.
        doi:10.1038/s41559-018-0606-3

Nadeau, N. J., Whibley, A., Jones, R. T., Davey, J. W., Dasmahapatra, K. K., Baxter, S. W.,
        Quail, M. A., Joron, M., Ffrench-Constant, R. H., Blaxter, M. L., Mallet, J., & Jiggins, C.
        D. (2012). Genomic islands of divergence in hybridizing *Heliconius* butterflies identified
        by large-scale targeted sequencing. *Philosophical Transactions of the Royal Society B:
        Biological Sciences, 367*(1587), 343-353. doi:10.1098/rstb.2011.0198

Osada, N., & Akashi, H. (2012). Mitochondrial-nuclear interactions and accelerated
        compensatory evolution: Evidence from the primate cytochrome c oxidase complex.
        *Molecular Biology and Evolution, 29*(1), 337-346.
        https://doi.org/10.1093/molbev/msr211

Panov, E. N., Rubtsov, A. A., & Monzikov, D. G. (2003). Hybridization between yellowhammer
        and pine bunting in Russia. *Dutch Birding, 25,* 17-31.

Panov, E. N., Rubtsov, A. S., & Mordkovich, M. V. (2007). New data on the relationships
        between two species of buntings (*Emberiza citrinella* and *E. leucocephalos*) hybridizing
        in the areas of overlap of their ranges, *Zoologicheskii Zhurnal, 86*(11), 1362–1378.

Paradis, E., & Schliep, K. (2019). ape 5.0: An environment for modern phylogenetics and
        evolutionary analyses in R. *Bioinformatics, 35*(3), 526-528.
        doi:10.1093/bioinformatics/bty633

Price, T. (2008). *Speciation in birds*. Roberts and Co.

R Core Team (2014). R: A language and environment for statistical computing. Vienna, Austria:
        R Foundation for Statistical Computing. Retrieved from http://www.r-project.org/

Rubtsov, A.S. 2007. Variability of songs of the yellowhammer (*Emberiza citrinella*) and pine
bunting (*Emberiza leucocephala*) as an evidence of population structure and evolutionary
history of the species, *Zoologicheskii Zhurnal, 86*(7), 863–876.

Rubtsov, A. S., & Tarasov, V. V. (2017). Relations between the yellowhammer (*Emberiza
citrinella*) and the pine bunting (*Emberiza leucocephalos*) in the forested steppe of the
Trans-Urals. *Biology Bulletin, 44*(9), 1059-1072. doi:10.1134/S1062359017090114

Ruegg, K., Anderson, E. C., Boone, J., Pouls, J., & Smith, T. B. (2014). A role for migration-
linked genes and genomic islands in divergence of a songbird. *Molecular Ecology,
23*(19), 4757-4769. doi:10.1111/mec.12842

Sackton, T. B., Corbett-Detig, R. B., Nagaraju, J., Vaishna, L., Arunkumar, K. P., & Hartl, D. L.
(2014). Positive selection drives faster-z evolution in silkmoths. *Evolution, 68*(8), 2331-
2342. doi:10.1111/evo.12449

Sackton, T. B., Haney, R. A., & Rand, D. M. (2003). Cytonuclear coadaptation in *Drosophila*:
Disruption of cytochrome c oxidase activity in backcross genotypes. *Evolution, 57*(10),
2315-2325. https://doi.org/10.1111/j.0014-3820.2003.tb00243.x

Saraste, M. (1999). Oxidative phosphorylation at the fin de siècle. *Science, 283*(5407), 1488-
1493. https://doi.org/10.1126/science.283.5407.1488

Scordato, E. S. C., Wilkins, M. R., Semenov, G., Rubtsov, A. S., Kane, N. C., & Safran, R. J.
(2017). Genomic variation across two barn swallow hybrid zones reveals traits associated
with divergence in sympatry and allopatry. *Molecular Ecology, 26*(20), 5676-
5691. https://doi.org/10.1111/mec.14276

Sloan, D. B., Havird, J. C., & Sharbrough, J. (2017). The on-again, off-again relationship
between mitochondrial genomes and species boundaries. *Molecular Ecology, 26*(8),
2212-2236. doi:10.1111/mec.13959

Smukowski, C. S., & Noor, M. A. F. (2011). Recombination rate variation in closely related
species. *Heredity, 107*(6), 496-508. doi:10.1038/hdy.2011.44

Stacklies, W., Redestig, H., Scholz, M., Walther, D., & Selbig, J. (2007). pcaMethods-a
bioconductor package providing PCA methods for incomplete data. *Bioinformatics,*
*23*(9), 1164-1167. doi:10.1093/bioinformatics/btm069

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics, 123*(3), 585-595.

Thornton, K., & Long, M. (2002). Rapid divergence of gene duplicates on the *Drosophila*
*melanogaster* X chromosome. *Molecular Biology and Evolution, 19*(6), 918-925.
doi:10.1093/oxfordjournals.molbev.a004149

Toews, D. P. L., & Brelsford, A. (2012). The biogeography of mitochondrial and nuclear
discordance in animals. *Molecular Ecology, 21*(16), 3907-3930. doi:10.1111/j.1365-
294X.2012.05664.x

Turner, S. (2018). qqman: An R package for visualizing GWAS results using Q-Q and
Manhattan plots. *Journal of Open Source Software, 3*(25), 731. doi:10.21105/joss.00731

Wang, S., Ore, M. J., Mikkelsen, E. K., Lee-Yaw, J., Toews, D. P. L., Rohwer, S., & Irwin, D.
(2021). Signatures of mitonuclear coevolution in a warbler species complex. *Nature*
*Communications*, in press.  doi:10.1038/s41467-021-24586-8

Warren, W. C., Clayton, D. F., Ellegren, H., Arnold, A. P., Hillier, L. W.,Künstner, A., Searle,
S., White, S., Vilella, A. J., Fairley, S., Heger, A., Kong, L., Ponting, C. P., Jarvis, E. D.,
Mello, C. V., Minx, P., Lovell, P., Velho, T. A. F., Ferris, M., … Wilson, R. K. (2010).
The genome of a songbird. *Nature, 464*(7289), 757–762. doi:10.1038/nature08819

Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population
structure. *Evolution, 38*(6), 1358-1370. doi:10.1111/j.1558-5646.1984.tb05657.x

Wu, C. (2001). The genic view of the process of speciation. *Journal of Evolutionary*
*Biology, 14*(6), 851-865. doi:10.1046/j.1420-9101.2001.00335.x

Yannic, G., Dubey, S., Hausser, J., & Basset, P. (2010). Additional data for nuclear DNA give
new insights into the phylogenetic position of *Sorex granarius* within the *Sorex araneus*

group. *Molecular Phylogenetics and Evolution, 57*(3), 1062-
1071. https://doi.org/10.1016/j.ympev.2010.09.015

Zink, R. M., & Barrowclough, G. F. (2008). Mitochondrial DNA under siege in avian
phylogeography. *Molecular Ecology, 17*(9), 2107-2121. doi:10.1111/j.1365-
294X.2008.03

# *Tables*

**Table 1.** Geographical locations and sample sizes of the sites included in this study. Sampling locations may include multiple sites that appeared too close together to be shown in detail in Figure 1A. Full details for the sites included in each sampling locations can be found in Supplementary Table 1. The sampling location numbers that appear in the "Sampling Location" column correspond to those that appear in red in Figure 1A. The "Sample Size" columns describes the total number of samples collected from a particular location.

| Sampling Location | Latitude (°N) | Longitude (°E) | *E. citrinella* sample size | *E. leucocephalos* sample size |
|---|---|---|---|---|
| 1 | 57.99 | 12.49 | 1 | 0 |
| 2 | 59.81 | 17.05 | 1 | 0 |
| 3 | 51.71 | 18.61 | 1 | 0 |
| 4 | 55.28 | 20.97 | 5 | 0 |
| 5 | 65.86 | 21.48 | 2 | 0 |
| 6 | 51.38 | 35.84 | 3 | 0 |
| 7 | 55.97 | 38.50 | 18 | 0 |
| 8 | 61.45 | 38.67 | 12 | 0 |
| 9 | 43.54 | 40.47 | 1 | 0 |
| 10 | 65.85 | 44.24 | 1 | 0 |
| 11 | 58.33 | 44.76 | 1 | 0 |
| 12 | 51.20 | 57.27 | 7 | 0 |
| 13 | 49.64 | 110.17 | 0 | 2 |
| 14 | 50.66 | 115.09 | 0 | 17 |
| 15 | 51.12 | 118.56 | 0 | 15 |
| 16 | 50.56 | 143.08 | 0 | 8 |
| | | **Total** | **53** | **42** |

**Table 2.** Summary statistics calculated while conducting mitonuclear co-introgression analysis. A total of 7187 windows, each of 2000 bp of obtained sequence, were considered when determining introgression windows. A total of 134 mitonuclear genes were investigated for signals of co-introgression. "*" indicates a significant p-value.

| Species | # of introgression windows identified | % of mitonuclear genes appearing in introgression windows | Binomial test p-value | Fisher's Exact test p-value |
|---|---|---|---|---|
| Yellowhammer | 244 | 6.7 | 0.04952* | 0.1311 |
| Pine bunting | 222 | 3.0 | 1 | 1 |

**Table 3.** Identities, chromosomal locations, windowed Tajima's D values and functions of mitonuclear genes that appeared within 244 yellowhammer introgression windows. In the "Mitonuclear Gene Function" column, ETC stands for "Electron Transport Chain". Mitonuclear gene names are written as they appear in Hill (2019).

| Mitonuclear Gene | Chromosome where mitonuclear gene is found | Windowed Tajima's D Value | Mitonuclear Gene Function |
|---|---|---|---|
| APOPT1 | 5 | -2.207 | Assembly factor/ancillary protein for ETC complex IV |
| COX5A | 10 | -2.420 | Structural subunit of ETC complex IV |
| COX17 | 1 | -2.509 | Assembly factor/ancillary protein for ETC complex IV |
| MRPL1 | 4 | -2.207 | Mitochondrial large ribosomal subunit protein |
| MRPL27 | 18 | -2.214 | Mitochondrial large ribosomal subunit protein |
| MRPL32 | 2 | -2.306 | Mitochondrial large ribosomal subunit protein |
| NDUFC1 | 4 | -2.399 | Structural subunit of ETC complex I |
| mtSSB | 1A | -2.362 | Single stranded DNA-binding protein |
| UQCR11 | 28 | -2.499 | Structural subunit of ETC complex III |

**Table 4.** Identities, chromosomal locations, windowed Tajima's D values and functions of mitonuclear genes that appeared within 222 pine bunting introgression windows. In the "Mitonuclear Gene Function" column, ETC stands for "Electron Transport Chain". Mitonuclear gene names are written as they appear in Hill (2019).

| Mitonuclear Gene | Chromosome where mitonuclear gene is found | Windowed Tajima's D Value | Mitonuclear Gene Function |
|---|---|---|---|
| ATP5H5I | 18 | -2.601 | Structural subunit of ETC complex V |
| COX5A | 10 | -2.545 | Structural subunit of ETC complex IV |
| MRPL2 | 3 | -2.247 | Mitochondrial large ribosomal subunit protein |
| NDUFB4 | 1 | -2.304 | Structural subunit of ETC complex I |

*Figures*

**A.**

**B.**

### Emberiza citrinella

Pure Citrinella (PC)

Almost Citrinella (SC)

### Emberiza leucocephalos

Pure Leucocephalos (PL)

Almost Leucocephalos (SL)

**Figure 1. A)** Map of sampling locations included in this study. Red numbers accompanying each location correspond to the sampling location numbers appearing in Table 1 which also describes sample sizes. Sampling locations may include multiple sites that appeared too close together to be shown in detail in this figure. Full details for the sites included in each sampling location can be found in Supplementary Table 1. Sampling location points are coloured based on the taxon caught in each area: yellowhammer (*Emberiza citrinella*; yellow) and pine bunting (*Emberiza leucocephalos*; brown). The solid black line indicates the geographic range of the yellowhammer and the dashed black line indicates the geographic range of the pine bunting as described in Irwin et al. (2009). **B)** Photos of plumage variation between yellowhammers and pine buntings. Each photo represents one of four phenotypic classes: PC, SC, PL and SL. Individuals with a PC and SC phenotypic class were grouped together as *Emberiza citrinella* and individuals with a PL and SL phenotypic class were grouped together as *Emberiza leucocephalos*. All photos are credited to Dr. Alexander Rubtsov.

**Figure 2.** Unrooted neighbour-joining tree of Emberizidae species constructed based on average absolute between-population nucleotide diversity ($\pi_B$). Sample sizes for each species are as follows: *E. aureola* = 1, *E. calandra* = 1, *E. cioides* = 1, *E. hortulana* = 1, *E. cirlus* = 6, *E. stewarti* = 4, *E. citrinella* = 53 and *E. leucocephalos* = 42.

**Figure 3.** PCA of genetic variation between allopatric yellowhammers (yellow; n = 53) and allopatric pine buntings (brown; n = 42), based on 374,780 genome-wide SNPs. PC1 and PC2 explain 3.6% and 2.9%, respectively, of the variation among individuals.

**Figure 4.** Relative differentiation ($F_{ST}$) of 349,807 genome-wide SNPs identified between allopatric yellowhammers (n = 53) and allopatric pine buntings (n = 42), with chromosomes represented with alternating black and grey. Narrow regions of elevated differentiation can be seen on many autosomes, and there are broad regions of high differentiation on the Z chromosome.

**Figure 5.** Patterns of genetic variation comparing allopatric yellowhammers (n = 53) and allopatric pine buntings (n = 42) across three chromosomes (2, 5 and Z). Relative nucleotide differentiation ($F_{ST}$), absolute between-population nucleotide diversity ($\pi_B$), absolute within-population nucleotide diversity ($\pi_W$) and Tajima's D (Taj$_D$) are shown as 2000 bp windowed averages across each chromosome. $F_{ST}$ and $\pi_B$ are shown as purple lines to indicate that values were calculated as a comparison between allopatric yellowhammers and pine buntings. $\pi_W$ and Taj$_D$ are shown as two separate lines (yellow = yellowhammers, brown = pine buntings) to indicate that values were calculated separately for each population.

**Figure 6.** Mean absolute within-group nucleotide diversity ($\pi_W$) of allopatric yellowhammers (n = 53) and allopatric pine buntings (n =42) plotted against absolute between-group nucleotide diversity ($\pi_B$). Each dot represents the average value taken from a 2000 bp window of sequenced data across the nuclear genome. The black line indicates where mean within-group nucleotide diversity equals between-group nucleotide diversity. Increasing values of relative differentiation ($F_{ST}$) calculated for each window are shown in darker shades of blue.

**Figure 7.** Association between relative differentiation ($F_{ST}$) and absolute between-group nucleotide diversity ($\pi_B$) of allopatric yellowhammers (n=53) and allopatric pine buntings (n = 42). Each black dot represents average values calculated from a 2000 bp window of sequenced data. A cubic spline fit between the variables is shown as a purple line.