Quantitative Restoration of Immune Defense in Old Animals Determined by Naïve Antigen-Specific CD8 T cell Numbers

Jennifer L. Uhrlaub[1,2*], Mladen Jergović[1,2], Christine M. Bradshaw[1,2], Sandip Sonar[1,2], Christopher P. Coplen[1,2], Jarrod Dudakov[3,4], Kristy O. Murray[5,6], Marion C. Lanteri[7], Michael P. Busch[7], Marcel R. M. van den Brink[8,9], and Janko Nikolich-Žugich[1,2, 3*]

1- Department of Immunobiology, University of Arizona College of Medicine, Tucson, AZ, USA

2- University of Arizona Center on Aging, University of Arizona, College of Medicine, Tucson, Tucson, AZ, USA

3- Program in Immunology, Clinical Research Division, and Immunotherapy Integrated Research Center, Fred Hutchinson Cancer Research Center, Seattle, WA 98109, USA;

4-Department of Immunology, University of Washington, Seattle, WA 98109, USA;

5- Department of Pediatrics, Section of Pediatric Tropical Medicine and National School of Tropical Medicine, Baylor College of Medicine, Houston, TX 77030, USA

6-William T. Shearer Center for Human Immunobiology, Texas Children's Hospital, Houston, TX 77030, USA

7- Blood Systems Research Institute, Vitalant Research Institute, San Francisco, CA, 94118, USA

8- Department of Medicine and Immunology Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

9-Immunology Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

Supported by the USPHS awards AG052359, AG020719, and AI091816-01 and the Bowman Endowed Chair in Medical Sciences

*Address correspondence to Janko Nikolich-Žugich at nikolich@arizona.edu or Jennifer Uhrlaub at juhrlaub@email.arizona.edu or by mail at P.O. Box 245221, Department of

Immunobiology, University of Arizona College of Medicine-Tucson, 1501 N. Campbell

Ave., Tucson, AZ 85724.

<u>ABSTRACT</u>

Older humans and animals often exhibit reduced immune responses to infection

and vaccination, and this often directly correlates to the numbers and frequency of

naïve T (Tn) cells. We found such a correlation between reduced numbers of blood

CD8+ Tn cells and severe clinical outcomes of West Nile virus (WNV) in both humans

naturally exposed to, and mice experimentally infected with, WNV.

To examine possible causality, we sought to increase the number of CD8 Tn

cells by treating C57BL/6 mice with IL-7 complexes (IL-7C, anti-IL-7 mAb bound to IL-

7), shown previously to efficiently increase peripheral T cell numbers by homeostatic

proliferation. T cells underwent robust expansion following IL-7C administration to old

mice increasing the number of total T cells (>four-fold) and NS4b:H-2D^b-restricted

antigen-specific CD8 T cells (two-fold). This improved the numbers of NS4b-specific

CD8 T cells detected at the peak of the response against WNV, but not survival of WNV

challenge. IL-7C treated old animals also showed no improvement in WNV-specific

effector immunity (neutralizing antibody and *in vivo* T cell cytotoxicity). To test

quantitative limits to which CD8 Tn cell restoration could improve protective immunity,

we transferred graded doses of Ag-specific precursors into old mice and showed that

injection of 5,400 (but not of 1,800 or 600) adult naïve WNV-specific CD8 T cells

significantly increased survival after WNV. These results set quantitative limits to the

level of Tn reconstitution necessary to improve immune defense in older organisms and

are discussed in light of targets of immune reconstitution.

Keywords: Immune aging/ immune rejuvenation/ IL-7/ CD8 T cells

Introduction

Prolonged T cell suppression after myeloablative conditioning is a major clinical problem with considerable clinical implications for therapies such as hematopoietic cell transplantation. Consequently, much work has gone into developing therapies that can boost T cell reconstitution, with varying degrees of success, most of which focus on stimulating thymic function (reviewed in (Granadier, Iovino, Kinsella, & Dudakov, 2021).

Aging is also accompanied by a variable degree of lymphopenia (Groarke & Young, 2019) due to combined effects of early-life thymic involution and late-life defects in lymphocyte maintenance in secondary lymphoid organs (SLO) (rev. in (Nikolich-Žugich, 2018). This lymphopenia in the last third of life correlates to decreased ability of the immune system to respond to infection and vaccination (rev, in (Nikolich-Žugich, 2018). Age-related lymphopenia affects all lymphocytes over time, but is drastically evident for naïve T (Tn) cells that reside in SLO such as the spleen and lymph nodes (LN) (Thompson, Smithey, Surh, & Nikolich-Žugich, 2017). Tn cells are the most diverse T cell subset, in charge of defending the organism against new infection, which transforms Tn cells (CD62LhiCD44lo in mice) specific for the invading infection into effector (Te, CD62LloCD44hi) and subsequently into memory (Tem – effector memory, also CD62LloCD44hi; and Tcm- central memory T cells, CD62LloCD44hiCD49hi). With aging, disturbances in SLO maintenance of Tn cells leads to the accumulation of antigen-independent memory cells (virtual memory; Tvm, CD62LloCD44hiCD49dlo) impacting their phenotype and function and diminishing their ability to protect the organism against infection (Rev. in (Nikolich-Žugich, 2018). It is well established that both CD4 and CD8 Tn cells diminish in numbers by the last third of life in mice (Thompson et al., 2017), non-human primates (Cicin-Sain et al., 2007; Cicin-Sain et al., 2010), and humans ((Groarke & Young, 2019; Wertheimer et al., 2014), albeit with slightly different kinetics depending on the species and compartment sampled. Associations between Tn lymphopenia and decreased immunity in old age were reported. Studies have shown that the CD8 response to live single-cycle poxvirus vaccination directly correlates to the CD8 Tn numbers in non-human primates (Cicin-Sain et al., 2010). Correlations were also found in a study of long-term repeated

apheresis subjects, where 15-20% of PBMC were retained by the apheresis machine,

and there was a correlation between the number of donations, lymphopenia, and

reactivation of persistent herpesviruses (e.g. Varicella-Zoster virus) (Zhao et al., 2021).

Moreover, severity of COVID-19 in older adults has been shown to correlate to reduced

numbers of CD4 and CD8 Tn cells (Rydyznski Moderbacher et al., 2020). A limitation of

these studies is that they did not independently control for aging, and, in most cases,

baseline samples were not available to answer whether this was a preexisting condition,

a virus-induced lymphodepletion, or a combination of both. All of these intriguing

correlative studies raise the possibility of causal association between the (age-related)

Tn lymphopenia and functional immunity decline with aging.

Reconstitution of T lymphopoiesis and maintenance in older organisms, also

known as immune rejuvenation, has been a highly sought goal in regenerative

immunotherapy. Success in that regard has been partial, with no clear translation into

an accepted medical protocol. Several treatments in humans and mice transiently

(usually up to 45-60 days in mice) reawaken thymic production of Tn cells, including

surgical (T. S. P. Heng et al., 2005) and various forms of chemical (Sutherland et al.,

2005) sex steroid ablation (SSA), and treatment with various growth factors, including

the human growth hormone (Taub, Murphy, & Longo, 2010) and the keratinocyte growth

factor (KGF) (Min et al., 2007). Functional evidence that immunity was improved

following such treatments in *bona fide* old mice was less abundant (T. S. Heng et al.,

2012; Min et al., 2007) and did not involve testing of immune protection using lethal

infectious challenge. We have recently discovered that KGF- and pharmacological SSA-

induced thymic rebound readily led to restoration, followed by a decline over time, of

both thymic cellularity and of the influx of recent thymic emigrants (RTE) into the blood

(Thompson et al., 2019) in old mice. Such RTE, however, sparsely populated old

spleens and lymph nodes (LN), and treated mice showed no improvement in survival

following challenge with West Nile virus (Thompson et al., 2019). That led to

investigations of LN involution with age, and its role in immune senescence and

reduced immune defense with aging.

As part of these studies, we sought to conclusively address the role of age-linked

lymphopenia in reduced immune defense with aging. Here, we report using IL-7

complexes (IL-7C, IL-7 bound to an anti-IL-7 Ab, (Boyman, Ramsey, Kim, Sprent, &

Surh, 2008) to increase lymph node cellularity by means of homeostatic proliferation

and found a doubling of CD8 Tn precursors specific for the immunodominant NS4b

epitope of WNV. However, most of these cells were no longer true naïve CD8 cells, but

rather antigen-independent Tvm. Upon challenge with WNV, we found no improvement

of immune protection, and no increase in either CD8 effector T cells or their cytotoxic

function, consistent with our prior findings that Tvm cells do not proliferate well upon

stimulation. These mice also did not produce robust anti-WNV antibodies, suggesting

that CD4 and/or B cell defects were not remedied either. To evaluate whether an

adequate reconstitution of true naïve cells could protect old mice against lethal

challenge, we transferred graded amounts of naïve CD8+ NS4b/Db-specific TCR

transgenic cells. We found that numbers of Ag-specific Tn cells needed to confer

protection to old mice were slightly above the numbers of Ag-specific Tn precursors

present in young adult mice, suggesting that both the numbers and the quality of

precursors are important in reconstitution. These results establish the numerical limits

of CD8 Tn reconstitution and are discussed in light of strategies for immune

reconstitution with aging.

RESULTS

Older age and reduced naïve T cell numbers are associated with WNV disease severity.

Previously unencountered and emerging viral infections often cause increased disease severity and mortality in older populations. Indeed, patients who have experienced the most severe symptoms caused by WNV were significantly older than those who had milder disease ((Campbell, Marfin, Lanciotti, & Gubler, 2002) and Figure 1A), and similar findings have been made in other emerging infectious diseases including SARS-CoV, Chikungunya virus, and SARS-CoV-2. Consistent with our prior data in general geriatric populations (Wertheimer et al., 2014) our cohort of WNV-exposed participants exhibited significantly lower levels of circulating CD8, but not CD4, Tn cells when compared individuals who experienced asymptomatic infection (Figure 1B, C). We determined T cell phenotype frequencies using a combination of CCR7, CD45RA, CD28 and CD95, to define naïve T cells as CCR7hi,CD45RAhi,CD28int/hi and CD95low. Participants who experienced the most severe symptoms, meningitis and encephalitis, exhibited a measurable loss of both Tn CD4 and CD8 T cells as compared to at least one other group. In a retrospective study, where a limited number of patient samples were collected long after asymptomatic infection or disease recovery, it is impossible to determine whether this was a cause, an epiphenomenon, or an effect, of their severe infection. To examine the nature of associations between age, Tn cell deficit, and severe WNV disease, we examined this issue in an established mouse model of age-related susceptibility to WNV (Brien, Uhrlaub, Hirsch, Wiley, & Nikolich-Zugich, 2009).

The murine model recapitulates the descriptive findings from human subjects.

It is well established that the naïve T cell compartment contracts with age and is proportionally less well represented than the T cells carrying memory phenotypes (Zhang, Weyand, & Goronzy, 2021). A direct correlation with disease severity, however, has not been conclusively established. To determine if our WNV mouse model is valid to interrogate this association, we measured the number of circulating CD8 Tn cells in WNV-infected adult (3-5 month old) and old (19-22 month old) mice on day 7 p.i.; a time

point prior to onset of WNV disease (Figure 2A). Mice that will eventually succumb to

WNV exhibited lower numbers of circulating naïve CD8 and CD4 T cells at this early

time point when compared to those that will survive (Figure 2B, C). This disparity held

regardless of age, despite the fact that old mice exhibit reduced numbers of Tn cells

overall (Figure 2B, C). These data support the possibility that a numeric advantage in

naïve T cells is directly linked to WNV resistance regardless of age. These results are

also consistent with the human data from Figure 1 with the exception that there is also a

robust loss of naïve CD4 T cells in susceptible mice. Of note, there was overlap in CD8

Tn numbers between humans with severe and asymptomatic WNV disease, as well as

between mice that succumbed and survived WNV infection. This suggests that the

numbers of CD8 Tn cells are not the only determinant of increased WNV resistance,

consistent with the extensive literature on the multipronged nature of immune defense

against WNV (Brien et al., 2009; Purtha et al., 2007; Richner et al., 2015). The overlap

in the human and mouse data provides confidence that this mouse model can be used

to examine whether, and to what extent, reversing age-induced lymphopenia and,

specifically, a deficit in Tn cells, may afford improved immune protection against WNV.

IL-7 complex treatment increases the number of circulating Tn cells

IL-7 is a pleiotropic cytokine that provides an indispensable homeostatic survival

signal to Tn cells, and that drives Tn and other T cell subsets into homeostatic

proliferation under the conditions of lymphopenia. Delivered *in vivo* as a cytokine-

antibody complex, IL-7C has been shown to drive naïve T cell proliferation and increase

thymopoeisis in unmanipulated as well as lymphopenic mice (Boyman, Kovar,

Rubinstein, Surh, & Sprent, 2006). Despite its therapeutic impact in adult mice, this

model has not been rigorously interrogated for reversal of age-related lymphopenia and

reduced susceptibility to infection. We treated old mice with IL-7C (3 i.p. injections,

every other day), and sacrificed them on days 7, 10, and 14 post-first IL-7C treatment to

assess total spleen and LN content (schema, Figure 3A). Total cellularity of secondary

lymphoid tissues (spleen and LNs) remained elevated through day 14 post-first

treatment (Figure 3B), an expansion mirrored by the numbers of CD4 and CD8 T cells

(Figure 3C, D). Additionally, numbers of CD4 and CD8 Tn cells in treated old mice were

no longer statistically significantly reduced relative to those in adult mice (Figure 3E, F).

The numerical increase in CD4 T cells was driven by split increases in Tcm and Tvm

phenotype cells, neither of which were independently statistically significant (Figure 3G,

H). In the CD8 pool, T cells with a CM or Tvm phenotype dominated the increase in

numbers (Figure 3I, J). To determine whether the expansion of T cells by IL-7C also

included CD8 T cells specific for the immunodominant epitope of WNV, $NS4b_{2488}$, we

enumerated the number of these CD8 T cell precursors on day 13 post-treatment in

unimmunized mice (Figure 3K). Similar to our prior work with other immunodominant

CD8 epitopes (Rudd, Venturi, Davenport, & Nikolich-Zugich, 2011; Smithey, Li, Venturi,

Davenport, & Nikolich-Žugich, 2012), we found a marked ~62% loss of NS4b-specific

CD8 precursors in old animals compared to adult counterparts (Figure 3F).   We found

that IL-7C treatment led to a numerical increase in CD8 precursors specific for the WNV

NS4b immunodominant epitope that, on average, no longer statistically differed from

that in adult animals (Figure 3K). The response between individual animals, however,

was heterogenous, with many animals exhibiting no increase, and others exhibiting up

to a two-fold increase in precursor numbers. Further, and as would be expected, NS4b-

specific precursors were of a similar phenotype as compared to the rest of the CD8 T

cell pool (Table 1).

Table 1		Adult	Old	Old+IL7-C
Tn	%CD44low CD62Lhigh of Db-NS4b CTLp (+/-SEM)	65.1 (2.6)	56.2 (4.2)	47.6 (7.1)
Tvm	%CD49dlow of CM Db-NS4b+ CTLp (+/-SEM)	93.7 (2.0)	75.7 (6.8)	83.8 (4.5)

218 The relative improvement in total T cell numbers, from days 10 to 14, including an

219 overall increase in WNV-specific CD8 precursors on day 13 following IL-7C treatment

220 suggested that functional immunity could be improved within this temporal window,

221 because lymphopenia in old mice was measurably reduced and residual IL-7C (if any)

222 did not cause further T cell proliferation in secondary lymphoid tissues.

IL-7c treated old mice gain numbers, but not function, of WNV-specific CD8 T cell
responses

To conclusively test whether this increase in T cell numbers is sufficient to overcome
infectious disease susceptibility, we challenged IL-7C treated mice with WNV. Based
on the above kinetics of IL-7C action, we chose day 13 as an optimal time point for
challenge, reasoning that the mice would benefit from an expanded T cell pool at the
time when IL-7C is no longer driving further proliferation of T cells, and should not
influence the development of a functional immune response to WNV. Adult (3-4
months); old (19-22 months); and IL-7C treated old (19-22 months) mice were
challenged with 1000pfu of WNV and followed for WNV-specific CD8 T cell responses
and survival (Figure 4A). On day 7 post-infection, old mice treated with IL-7C showed a
3-fold improvement in the number of WNV NS4b-specific CD8 T cells measured in the
blood (Figure 4B). However, this improvement failed to translate to better survival
(Figure 4C), despite the fact that we observed roughly equivalent numbers of WNV-
specific CD8 T cells in old IL-7C treated mice compared to adult unmanipulated mice.
When we segregated each experimental group into animals that died or survived WNV
challenge, we found that those that survived again exhibited a numeric advantage in
WNV-specific cells over those that succumbed (Figure 4D). The effect, however, was
not robust enough to ensure that the group, as a whole, survives better than their
untreated counterparts.

Multiple effector mechanisms of adaptive immunity must synergize to clear WNV
infection. These include potent neutralizing antibodies as well as CD4 and CD8
cytotoxic T cell effector function. To mechanistically determine whether any functional
improvement in immunity was restored by IL-7C treatment we interrogated the immune
response in IL7-C treated old mice in a step-wise manner. IL-7C have the potential to
enhance B cell lymphopoiesis, and to thereby increase B cell numbers, although they
are not expected to directly affect the mature B cell compartment (Boyman et al., 2008).
Previous studies have reported that IL-7C co-delivered with immunization can boost
humoral immune responses through enhanced generation of germinal center B cells
and Tfh cells (Seo et al., 2014). Therefore, we first analyzed neutralizing antibody titers

at day 7 post-WNV infection as a proxy for generalized improvement in B cell

responses. We detected no improvement in WNV neutralization by sera of old mice

treated with IL7-C relative to untreated controls (Figure 4E). We next examined

whether WNV-specific CD8 or CD4 T cells exhibited improved killing function following

IL-7C treatment by measuring cytotoxicity in vivo (Figure 4F, G). We included targets

specific for one WNV-specific CD4 immunodominant epitope, ENV_{641}, in this assay

since CD4 T cells are known to mount cytotoxic responses against WNV in adult mice

and data about their specific contribution to viral clearance with age is limited (Brien,

Uhrlaub, & Nikolich-Zugich, 2008). We found that, despite numeric increases in

circulating NS4b-specific CD8 T cells, IL-7C treated old mice did not have improved

cytotoxic capacity to clear $NS4b_{2488}$ peptide-loaded targets as compared to untreated

old mice (Figure 4F). There was also no improvement in killing of the ENV_{641}-loaded

targets (Figure 4G), consistent with the fact that we saw no increase in CD4 Tn cell

numbers (Fig. 3E). These data indicate an overall lack of improvement in cytotoxic

function in old mice treated with IL-7C. As these assessments were done prior to the

onset of disease, we measured memory antibody responses in WNV survivors as an

indication of whether IL7-C treatment impacted overall humoral immunity. We

determined that neither total envelope nor DIII lateral ridge specific antibodies were

improved by IL7-C treatment prior to WNV infection (Figure H, I). In fact, when

comparing the survivors of WNV infection, it appears that old mice treated with IL-7C

established an overall lower level of circulating WNV-specific antibodies at this point in

memory (Figure H, I). Our interpretation of this data is that the expansion of CD4 T cells

by IL-7C treatment does not afford a short- or lasting- impact on antibody responses to

WNV infection. At face value, these results suggest that numerical improvement in CD8

and CD4 T cells afforded by IL-7C treatment is not sufficient to improve functional

immunity.

To investigate why a numeric increase in antigen-specific precursors was not

able to restore T cell responses we considered the possibility that the expansion

induced via IL7-C treatment may push the cells into an exhausted state, impairing their

ability to participate in the WNV response. T cells with an 'exhausted' phenotype are

typically seen in the course of chronic infections and cancer (Collier et al., 2021; Reiner,

2021). The number of divisions and other factors required to induce this state are not

fully understood. To investigate whether homeostatic proliferation driven by IL-7C

treatment increases the proportion of responding WNV-specific T cells with an

exhausted phenotype we measured exhaustion markers in both adult and old mice

treated with IL7-C on day 7 of WNV. We included an adult cohort in these experiments

to dissociate aging from treatment-associated phenotypes. Our data confirm that a

higher percentage of WNV-specific CD8 T cells in aged mice express PD-1, LAG-3, or

2B4 regardless of whether they have been treated with IL7-C (Figure 5A, B, C). The

expression of these molecules could contribute to their poor cytotoxic function, and

overall lack of improvement in survival, experienced by the old animals. These data

suggest that the homeostatic proliferation induced by IL7-C treatment is not *per se*

pushing T cells into an exhausted state. Rather, consistent with some reports (Decman

et al., 2012) we hypothesize that the age of the animal may have already determined

the phenotype of the available T cell pool and this state is maintained through the

expansion initiated by IL7-C treatment to deleterious effect.

However, the fact that a similar increase in precursor numbers induced by IL-7C

treatment did not lead to either improved immunity or improved cytotoxicity, suggested

that some other property of reconstituted cells may be of critical importance. To directly

address whether an increase in CD8 Tn precursors specific for an invading pathogen

can overcome the age-related defects in immune defense and afford improved survival

to old mice, we performed adoptive transfer experiments. We transferred graded levels

of WNV-specific CD8 T cells (NS4b$_{2488}$ specific, referred to as WNV-I (Aguilar-

Valenzuela et al., 2018)) the day before challenge into old mice to boost the number of

precursors available to respond. These transgenic T cells have a naïve phenotype at

time of transfer and are not yet primed or cytotoxic. After challenge, the transferred

cells quantitatively restored the WNV-specific CD8 T cell numbers measured in the

blood on day 7 post-infection (Figure 6A). The differences between old mice treated

with IL-7C and those receiving adoptively transferred cells was that the former mounted

a response from endogenous, old precursors, and these precursors were driven into

homeostatic proliferation by IL-7C; the latter received naïve, unperturbed NS4b-specific

precursors from adult donors. Both the age and the immediate prior proliferation

induced by IL-7C could have impacted their response. More importantly, in the transfer

experiments, we were able to significantly improve survival in old mice following WNV

challenge with 5,400 transferred WNV-specific CD8+ Tn cells; but not with the two lower

transfer boluses of 1,800 and 600 cells (Figure 6B).

To put these numbers in the context of physiological precursor numbers available

to old and adult animals, we determined the efficacy of engraftment of transferred cells.

Some estimates of transfer efficiency posit that only about 10% of the transferred T cells

engraft in i.v. injected mice (Blattman et al., 2002; Moon et al., 2009). We enumerated

the total number of NS4b-specific CD8 T cells that were either endogenous (CD45.2) or

transgenic (CD45.1). Indeed, our experiments confirmed that 24h after transfer, we

could recover between 12-15% of transgenic cells (Supp. Fig 1A). Note here that the

numbers of engrafted cells were quantified in direct comparison to the number of

endogenous naïve precursors recovered from adult or old mice as they were

enumerated alongside each other (Supp. Fig 2). This means that a dose of 5,400

transferred cells reconstituted about 650-810 NS4b precursors, that were effectively

capable to significantly protect the old animal against WNV with survival not significantly

different from untreated adult mice. Of interest, an adult mouse contains 500-700 NS4b-

specific CD8+ precursors. Importantly, old mice that received 5400 WNV-I Tg CD8 T

cells efficiently recruited them to the WNV response (Figure 6C) and exhibited improved

cytotoxic function measured by increased killing (Figure 6D). This offers a plausible

mechanism for improved survival of this group (Figure 6B). We conclude that restoring

the precursor numbers of an immunodominant CD8 Tn specificity to adult levels, and

with cells that have a Tn phenotype, may be enough to confer a significant degree of

protection to an old mouse against WNV, despite concurrent defects in many other

arms of the immune response.

DISCUSSION

Immune therapies that seek to overcome age-induced lymphopenia are an

attractive strategy to reduce susceptibility to infectious diseases in older adults.

Conceptually, increasing the total cellular resources of the immune system, regardless

of antigen specificity, would provide the best solution for broad-based protection against

diverse challenges. IL-7C treatment has been shown to increase peripheral T cell

numbers even in the absence of thymic output (Boyman et al., 2008). In this work, we

formally demonstrated that IL-7C treatment can transiently increase T cell numbers in

old mice that have suffered substantial thymic involution, predominantly due to an

expansion of virtual central memory Tvm phenotype cells. We found significant, but

variable, expansion of the numbers of CD8 precursors specific for WNV, and an overall

increase in both CD8 and CD4 T cell numbers. However, these expanded cells did not

protect mice from WNV infection, and we could find no evidence that the cytotoxic

function of either CD8 or CD4 T cells was improved in IL-7C treated old mice. While we

did not enumerate B cells, their protective responses were similarly flat, with no

improvement in neutralization capacity, despite an increase in total numbers of CD4 T

cells. Therefore, under our experimental conditions, IL-7C treatment showed no

appreciable immune protection benefit to the old mice, similar to our prior studies with

unconjugated IL-7 in non-human primates (Okoye et al., 2015).

Given that protection against WNV normally requires a coordinated and multi-

faceted response, it may not be surprising that the increase in T cells alone cannot

augment protection. It was more puzzling, however, that we could find no evidence of

improved responses from the numerically increased T cell pool. To further explore this

question and test whether we could improve protection against WNV by boosting a

single arm of protective immunity, we performed adoptive transfer of graded numbers of

transgenic CD8 T cells. We demonstrated that sufficient numbers of transferred cells

(approximately equivalent to the precursor content of a single adult mouse) were able to

significantly improve survival, albeit not to restore it to the levels of adult mice. These

results begin to provide quantitative targets of CD8 T cell reconstitution, and also

provide a proof-of-principle that the aged immune system can coordinate an effective

immune response if the proper T cell resources are provided.

One key unanswered question remained regarding the relationship between

precursor T cell numbers and their quality. Specifically, adoptively transferred old mice

contained similar numbers of overall precursors relative to the IL-7C treated old mice,

suggesting that somehow the quality of the two precursor populations was different. To

address the potential differences, we considered prior work from ours and other groups

on age-related changes in Tn cell maintenance. Specifically, with aging, an increasing

number of Tn cells change their maintenance pattern and become Tvm (Chiu, Martin,

Stolberg, & Chensue, 2013; Decman et al., 2012; Rudd et al., 2011) to the point that

most of the central memory phenotype cells in SPF mice are of Tvm origin. While work

from other groups (Lee, Hamilton, Akue, Hogquist, & Jameson, 2013) and us

(Renkema, Li, Wu, Smithey, & Nikolich-Žugich, 2014) demonstrated that Tvm cells in

young adult mice exhibit strong proliferation and enhanced effector function relative to

Tn cells, we have also shown that with age Tvm cells exhibit reduced proliferative

response to cognate antigen (Renkema et al., 2014). By contrast, the response to

homeostatic cytokines remains preserved in Tvm cells. Additional studies will be

necessary to directly compare in vivo antiviral function and protective capacity of Tvm

and Tn cells of different ages with and without prior expansion by IL-7C and similar

treatments. Moreover, as these studies were focused on improving immune protection

against acute lethal challenge, it remains to be investigated whether and to what extent

the interventions used here, including IL-7C, could potentially improve responses to

vaccination, including memory responses. That notwithstanding, the work presented

here places quantitative limits on desirable levels of immune reconstitution and opens

provocative questions about their quality, that remain to be answered in the future.

EXPERIMENTAL PROCEDURES

Study Subjects and Blood Samples

This study was approved by the Institutional Review Boards at the University of Arizona (Tucson, AZ; # 080000673; currently #2102460536), the Oregon Health and Science University (Portland, OR; # IRB00003007); the University of Texas Health Science Center at Houston, TX; and the Blood Systems Research Institute, San Francisco, CA. – presently Vitalant Research Institute. Exclusion criteria included known immunosuppressive pathology, stroke, cancer, or use of steroids within the last 5 years. WNV-infected donors were enrolled by the University of Texas at Houston, 2006–2009 or Blood Systems Research Institute (BSRI), 2009 – 2011. Samples, demographics (Supp. Table I), and symptoms data were collected and analyzed as previously described (Lanteri et al., 2014) after the subjects provided an informed consent approved by the UCSF Committee on Human Research (protocol #10-01255) or the Committee for the Protection of Human Subjects at the University of Texas Health Science Center at Houston (HSC-SPH-03-039), respectively. Blood was drawn into heparinized Vacutainer CPT tubes (BD Bioscience) and processed fresh at respective sites per manufacturer's recommendations to isolate peripheral blood mononuclear cells (PBMC) and plasma; K2-EDTA tubes were used to determine complete blood counts. PBMCs were frozen in 90% fetal bovine serum (FBS) and 10% DMSO.

Flow Cytometry – human samples

Frozen PBMCs were thawed in RPMI medium supplemented with 10% FBS, penicillin and streptomycin in the presence of DNAse (Millipore Sigma) then rested overnight in X-Vivo medium (Lonza) supplemented with 5% human male AB serum. $1–3 \times 10^6$ PBMC were stained with a saturating concentration of the following antibodies: CD3 AF700 (clone SP34-2, BD Biosciences), CD4 eFluor450 (clone OKT4, eBioscience), CD8β ECD (2ST8.5H7, Beckman Coulter), CD95 APC (clone DX2, BD Biosciences), and CD28 PercpCy5.5 (clone CD28.2, BioLegend) in 1XPBS, 2% FBS and next incubated with LIVE/DEAD® Fixable Dead Cell Stain (ThermoFisher Sci), both for 30

min at 4°C. Flow cytometry acquisition was performed on a BD Bioscience Fortessa and

analyzed using FlowJo software (Tree Star).

Mice

Mouse experiments were carried out in strict accordance with the recommendations in

the Guide for the Care and Use of Laboratory Animals of the National Institutes of

Health. Protocols were approved by the Institutional Animal Care and Use Committee at

the University of Arizona (IACUC 19-580, PHS Assurance No. A-3248-01. Old (19-22

months) and adult (3-5 months) were obtained from the National Institute on Aging

Rodent Resource or Jax laboratories. Mice in each experiment were age matched

within 4 weeks. WNV-I mice were obtained from Drs Jason Netland and. Michael J.

Bevan (Aguilar-Valenzuela et al., 2018). Footpad (f.p.) injections and retro-orbital

intravenous cell transfers were performed under isoflurane anesthesia. IL-7 cytokine

was obtained from the Biological Resources Branch (BRB), Developmental

Therapeutics Program (DTP), Division of Cancer Treatment and Diagnosis (DCTD),

National Cancer Institute (NCI). IL-7R antibody (clone M25) was purchased from

BioXcell. 0.59ug of IL-7 was complexed with 2.9ug of M25 antibody for 30 min at RT

then administered via intraperitoneal injection in 200uL USP saline on days 1, 3, and 5

of the experiment. Euthanasia was performed by isoflurane overdose or cervical

dislocation.

Virus

West Nile virus NY 385–99, isolated from the liver of a snowy owl, and in its second

passage, was a kind gift from Dr. Robert Tesh (University of Texas Medical Branch at

Galveston, Galveston, TX). Virus was prepared as previously described (Brien et al.,

2009). Mice were infected by footpad (f.p.) route with 1,000 pfu WNV/50 µl/mouse and

monitored for survival until 45 days p.i.

Tissue harvest, cell counts, flow cytometry, neutralization titers, antibody ELISA

Blood was collected via retro-orbital route and subjected to hypotonic lysis of red blood

cells. Spleen and peripheral lymph nodes (inguinal, brachial, axillary, and cervical) were

collected and disassociated over 40uM mesh. Cells were resuspended and stained

overnight at 4°C in a saturating concentration of antibodies against CD49d (9C10,

MFR4.B), CD44 (IM7), CD62L (MEL-14), CD3 (17A2), CD4 (RM4–5), CD8a (53–6.7),

CD45.1 (A20), CD244.2 (m2B4 (B6) 458.1), CD223 (LAG-3, C9B7W), PD-1 (29F.1A12)

(eBioscience and BioLegend) and H2-Db/NS4b tetramers (NIH Tetramer Facility,

Atlanta, GA) followed by Live/Dead viability dye (ThermoFisher Sci). Cytotoxic

precursor enumeration was accomplished as described in (Uhrlaub, Smithey, &

Nikolich-Žugich, 2017). NS4b-Tg CD45.1 CD8 T cells were purified with Miltenyi Biotec

reagents and columns using negative magnetic selection. Cell counts and purity (>95%)

were confirmed by FCM analysis and count beads before adoptive transfer.

Optimization procedures for flow cytometry are based on current standards and

included full-minus-one (FMO) gating controls. Samples were acquired on a BD

LSRFortessa cytometer (BD Biosciences) or Cytek Aurora (Cytek Biosciences) and

analyzed with FlowJo software (Tree Star). WNV neutralization titers were performed as

described in (Wong et al., 2020) and scored as a 90% reduction in viral inoculum. WNV

ELISA's for both total ENV and DIII were also performed as described in (Wong et al.,

2020) and directly reported as O.D.

In Vivo Killing Assay

Splenocytes from C57BL/6 mice were disassociated into a single cell suspension and

labeled with 1µM CellTrace Far Red (ThermoFisher Sci). Targets were then divided

into two populations: $NS4b_{2488}$ targets were labeled with Cell Trace Violet

(ThermoFisher Sci) 5mM for 10^{-6}M peptide loaded targets and 0.5mM for non-loaded

negative control) and ENV_{641} targets were labeled with CFSE (ThermoFisher Sci) 5mM

for 10^{-6}M peptide loaded targets and 0.5mM for non-loaded negative control. All

labeling was done as described in manufacturer instructions. Peptide loading was 1

hour at 37°C. After thorough washing, targets were pooled and injected intravenously

(i.v.) into infected and naïve mice. After 4 hours mice were sacrificed spleen

disassociated to a single cell suspension. Splenocytes were gated on total targets

(CellTrace Far Red positive) then CellTrace Violet high and low or CFSE high and low.

Calculation of killing: 100% - [(% peptide bearing targets IMMUNE recipients / % no

peptide targets IMMUNE recipients) / (% peptide bearing targets NAIVE recipients / %
no peptide targets NAIVE recipients) x 100].

STATISTICS:

GraphPad Prism® version 9.2.0 (GraphPad Software) was used for data analysis. All
experiments were performed at least twice. Significance was determined as described
in figure legends.

ACKNOWLEDGEMENTS:

Authors wish to thank Drs Jason Netland and Michael J. Bevan for sharing the WNV-I
TCR transgenic mice; the NIH tetramer facility for providing Class I MHC tetramers; Dr.
Deepta Bhattacharya for providing reagents; Dr. Anne Wertheimer for assistance with
human sample processing; and Drs. Nancy R. Manley, Lauren I. R. Ehrlich, Bonnie J.
LaFleur, Gregory Sempowski, and Laura Hale for their critical reading of the manuscript
and suggestions. Graphics were created in Biorender.com.

CONFLICT OF INTEREST:

J.N.Ž. is co-chair of the scientific advisory board of and receives research funding from
Young Blood Institute, Inc.

AUTHOR CONTRIBUTIONS:

J.L.U., J.N-Ž., J.D. and M.R.M.vdB. designed the research; J.L.U., M.J., C.M.B., S.S.,
and C.P.C. performed experiments and analyzed the data; K.O.M., M.C.L., M.P.B.
contributed human samples; J.L.U. prepared the figures; J.L.U. and J.N-Ž. wrote the
manuscript; all authors edited the manuscript.

DATA AVAILABILITY:

All data available upon request.

REFERENCES

Aguilar-Valenzuela, R., Netland, J., Seo, Y. J., Bevan, M. J., Grakoui, A., & Suthar, M. S. (2018). Dynamics of Tissue-Specific CD8(+) T Cell Responses during West Nile Virus Infection. *J Virol, 92*(10). doi:10.1128/jvi.00014-18

Blattman, J. N., Antia, R., Sourdive, D. J., Wang, X., Kaech, S. M., Murali-Krishna, K., . . . Ahmed, R. (2002). Estimating the precursor frequency of naive antigen-specific CD8 T cells. *J Exp Med, 195*(5), 657-664. doi:10.1084/jem.20001021

Boyman, O., Kovar, M., Rubinstein, M. P., Surh, C. D., & Sprent, J. (2006). Selective stimulation of T cell subsets with antibody-cytokine immune complexes. *Science, 311*(5769), 1924-1927. doi:10.1126/science.1122927

Boyman, O., Ramsey, C., Kim, D. M., Sprent, J., & Surh, C. D. (2008). IL-7/anti-IL-7 mAb complexes restore T cell development and induce homeostatic T Cell expansion without lymphopenia. *J Immunol, 180*(11), 7265-7275. doi:10.4049/jimmunol.180.11.7265

Brien, J. D., Uhrlaub, J. L., Hirsch, A., Wiley, C. A., & Nikolich-Zugich, J. (2009). Key role of T cell defects in age-related vulnerability to West Nile virus. *J Exp Med, 206*(12), 2735-2745. doi:10.1084/jem.20090222

Brien, J. D., Uhrlaub, J. L., & Nikolich-Zugich, J. (2008). West Nile virus-specific CD4 T cells exhibit direct antiviral cytokine secretion and cytotoxicity and are sufficient for antiviral protection. *J Immunol, 181*(12), 8568-8575. doi:10.4049/jimmunol.181.12.8568

Campbell, G. L., Marfin, A. A., Lanciotti, R. S., & Gubler, D. J. (2002). West Nile virus. *The Lancet Infectious Diseases, 2*(9), 519-529. doi:https://doi.org/10.1016/S1473-3099(02)00368-7

Chiu, B. C., Martin, B. E., Stolberg, V. R., & Chensue, S. W. (2013). Cutting edge: Central memory CD8 T cells in aged mice are virtual memory cells. *J Immunol, 191*(12), 5793-5796. doi:10.4049/jimmunol.1302509

Cicin-Sain, L., Messaoudi, I., Park, B., Currier, N., Planer, S., Fischer, M., . . . Nikolich-Zugich, J. (2007). Dramatic increase in naive T cell turnover is linked to loss of naive T cells from old primates. *Proc Natl Acad Sci U S A, 104*(50), 19960-19965. doi:10.1073/pnas.0705905104

Cicin-Sain, L., Smyk-Pearson, S., Currier, N., Byrd, L., Koudelka, C., Robinson, T., . . . Nikolich-Zugich, J. (2010). Loss of naive T cells and repertoire constriction predict poor response to vaccination in old primates. *J Immunol, 184*(12), 6739-6745. doi:10.4049/jimmunol.0904193

Collier, D. A., Ferreira, I., Kotagiri, P., Datir, R. P., Lim, E. Y., Touizer, E., . . . Gupta, R. K. (2021). Age-related immune response heterogeneity to SARS-CoV-2 vaccine BNT162b2. *Nature*. doi:10.1038/s41586-021-03739-1

Decman, V., Laidlaw, B. J., Doering, T. A., Leng, J., Ertl, H. C., Goldstein, D. R., & Wherry, E. J. (2012). Defective CD8 T cell responses in aged mice are due to quantitative and qualitative changes in virus-specific precursors. *J Immunol, 188*(4), 1933-1941. doi:10.4049/jimmunol.1101098

Granadier, D., Iovino, L., Kinsella, S., & Dudakov, J. A. (2021). Dynamics of thymus function and T cell receptor repertoire breadth in health and disease. *Semin Immunopathol, 43*(1), 119-134. doi:10.1007/s00281-021-00840-5

Groarke, E. M., & Young, N. S. (2019). Aging and Hematopoiesis. *Clin Geriatr Med, 35*(3), 285-293. doi:10.1016/j.cger.2019.03.001

Heng, T. S., Reiseger, J. J., Fletcher, A. L., Leggatt, G. R., White, O. J., Vlahos, K., . . . Boyd, R. L. (2012). Impact of sex steroid ablation on viral, tumour and vaccine responses in aged mice. *PLoS One, 7*(8), e42677. doi:10.1371/journal.pone.0042677

Heng, T. S. P., Goldberg, G. L., Gray, D. H. D., Sutherland, J. S., Chidgey, A. P., & Boyd, R. L. (2005). Effects of Castration on Thymocyte Development in Two Different Models of Thymic Involution. *The Journal of Immunology, 175*(5), 2982-2993. doi:10.4049/jimmunol.175.5.2982

Lanteri, M. C., Lee, T. H., Wen, L., Kaidarova, Z., Bravo, M. D., Kiely, N. E., . . . Busch, M. P. (2014). West
Nile virus nucleic acid persistence in whole blood months after clearance in plasma: implication
for transfusion and transplantation safety. *Transfusion, 54*(12), 3232-3241.
doi:10.1111/trf.12764
Lee, J. Y., Hamilton, S. E., Akue, A. D., Hogquist, K. A., & Jameson, S. C. (2013). Virtual memory CD8 T
cells display unique functional properties. *Proc Natl Acad Sci U S A, 110*(33), 13498-13503.
doi:10.1073/pnas.1307572110
Min, D., Panoskaltsis-Mortari, A., Kuro, O. M., Holländer, G. A., Blazar, B. R., & Weinberg, K. I. (2007).
Sustained thymopoiesis and improvement in functional immunity induced by exogenous KGF
administration in murine models of aging. *Blood, 109*(6), 2529-2537. doi:10.1182/blood-2006-
08-043794
Moon, J. J., Chu, H. H., Hataye, J., Pagán, A. J., Pepper, M., McLachlan, J. B., . . . Jenkins, M. K. (2009).
Tracking epitope-specific T cells. *Nat Protoc, 4*(4), 565-581. doi:10.1038/nprot.2009.9
Nikolich-Žugich, J. (2018). The twilight of immunity: emerging concepts in aging of the immune system.
Nat Immunol, 19(1), 10-19. doi:10.1038/s41590-017-0006-x
Okoye, A. A., Rohankhedkar, M., Konfe, A. L., Abana, C. O., Reyes, M. D., Clock, J. A., . . . Picker, L. J.
(2015). Effect of IL-7 Therapy on Naive and Memory T Cell Homeostasis in Aged Rhesus
Macaques. *J Immunol, 195*(9), 4292-4305. doi:10.4049/jimmunol.1500609
Purtha, W. E., Myers, N., Mitaksov, V., Sitati, E., Connolly, J., Fremont, D. H., . . . Diamond, M. S. (2007).
Antigen-specific cytotoxic T lymphocytes protect against lethal West Nile virus encephalitis. *Eur J
Immunol, 37*(7), 1845-1854. doi:10.1002/eji.200737192
Reiner, S. L. (2021). A Regenerative Perspective on Successful and Failed T-Cell Immunity. *Cold Spring
Harb Perspect Biol, 13*(7). doi:10.1101/cshperspect.a037937
Renkema, K. R., Li, G., Wu, A., Smithey, M. J., & Nikolich-Žugich, J. (2014). Two separate defects affecting
true naive or virtual memory T cell precursors combine to reduce naive T cell responses with
aging. *J Immunol, 192*(1), 151-159. doi:10.4049/jimmunol.1301453
Richner, J. M., Gmyrek, G. B., Govero, J., Tu, Y., van der Windt, G. J., Metcalf, T. U., . . . Diamond, M. S.
(2015). Age-Dependent Cell Trafficking Defects in Draining Lymph Nodes Impair Adaptive
Immunity and Control of West Nile Virus Infection. *PLoS pathogens, 11*(7), e1005027.
doi:10.1371/journal.ppat.1005027
Rudd, B. D., Venturi, V., Davenport, M. P., & Nikolich-Zugich, J. (2011). Evolution of the antigen-specific
CD8+ TCR repertoire across the life span: evidence for clonal homogenization of the old TCR
repertoire. *J Immunol, 186*(4), 2056-2064. doi:10.4049/jimmunol.1003013
Rydyznski Moderbacher, C., Ramirez, S. I., Dan, J. M., Grifoni, A., Hastie, K. M., Weiskopf, D., . . . Crotty,
S. (2020). Antigen-Specific Adaptive Immunity to SARS-CoV-2 in Acute COVID-19 and
Associations with Age and Disease Severity. *Cell, 183*(4), 996-1012.e1019.
doi:10.1016/j.cell.2020.09.038
Seo, Y. B., Im, S. J., Namkoong, H., Kim, S. W., Choi, Y. W., Kang, M. C., . . . Sung, Y. C. (2014). Crucial roles
of interleukin-7 in the development of T follicular helper cells and in the induction of humoral
immunity. *J Virol, 88*(16), 8998-9009. doi:10.1128/jvi.00534-14
Smithey, M. J., Li, G., Venturi, V., Davenport, M. P., & Nikolich-Žugich, J. (2012). Lifelong persistent viral
infection alters the naive T cell pool, impairing CD8 T cell immunity in late life. *J Immunol,
189*(11), 5356-5366. doi:10.4049/jimmunol.1201867
Sutherland, J. S., Goldberg, G. L., Hammett, M. V., Uldrich, A. P., Berzins, S. P., Heng, T. S., . . . Boyd, R. L.
(2005). Activation of Thymic Regeneration in Mice and Humans following Androgen Blockade.
The Journal of Immunology, 175(4), 2741-2753. doi:10.4049/jimmunol.175.4.2741

Taub, D. D., Murphy, W. J., & Longo, D. L. (2010). Rejuvenation of the aging thymus: growth hormone-
mediated and ghrelin-mediated signaling pathways. *Curr Opin Pharmacol, 10*(4), 408-424.
doi:10.1016/j.coph.2010.04.015
Thompson, H. L., Smithey, M. J., Surh, C. D., & Nikolich-Žugich, J. (2017). Functional and Homeostatic
Impact of Age-Related Changes in Lymph Node Stroma. *Front Immunol, 8*, 706.
doi:10.3389/fimmu.2017.00706
Thompson, H. L., Smithey, M. J., Uhrlaub, J. L., Jeftić, I., Jergović, M., White, S. E., . . . Nikolich-Žugich, J.
(2019). Lymph nodes as barriers to T-cell rejuvenation in aging mice and nonhuman primates.
Aging Cell, 18(1), e12865. doi:10.1111/acel.12865
Uhrlaub, J. L., Smithey, M. J., & Nikolich-Žugich, J. (2017). Cutting Edge: The Aging Immune System
Reveals the Biological Impact of Direct Antigen Presentation on CD8 T Cell Responses. *The*
Journal of Immunology, 199(2), 403. doi:10.4049/jimmunol.1700625
Wertheimer, A. M., Bennett, M. S., Park, B., Uhrlaub, J. L., Martinez, C., Pulko, V., . . . Nikolich-Žugich, J.
(2014). Aging and cytomegalovirus infection differentially and jointly affect distinct circulating T
cell subsets in humans. *J Immunol, 192*(5), 2143-2155. doi:10.4049/jimmunol.1301721
Wong, R., Belk, J. A., Govero, J., Uhrlaub, J. L., Reinartz, D., Zhao, H., . . . Bhattacharya, D. (2020). Affinity-
Restricted Memory B Cells Dominate Recall Responses to Heterologous Flaviviruses. *Immunity,*
53(5), 1078-1094.e1077. doi:10.1016/j.immuni.2020.09.001
Zhang, H., Weyand, C. M., & Goronzy, J. J. (2021). Hallmarks of the aging T-cell system. *Febs j*.
doi:10.1111/febs.15770
Zhao, J., Gabriel, E., Norda, R., Höglund, P., Baden, L., Diedrich, B. A., . . . Edgren, G. (2021). Frequent
platelet donation is associated with lymphopenia and risk of infections: A nationwide cohort
study. *Transfusion, 61*(2), 464-473. doi:10.1111/trf.16175

FIGURE LEGENDS:

Figure 1

Older adults experience more severe WNV disease and have a dearth of naïve

CD8 T cells.

A cohort of 52 patients with WNV disease severity clinically classified as Asymptomatic,

WN Fever (WNF), WN Meningitis (WNM), and WN meningitis and encephalitis (WNME)

were retrospectively recruited to participate in our study. The age of the patients

experiencing the most severe disease is significantly older than asymptomatic (A).

Naïve (CD95^{low} and $^{CD28int/high}$, also confirmed additionally to be CCR7hi and CD45RA+)

CD4 and CD8 T cells were phenotyped by FCM and counts were extrapolated from

clinical cell blood count (CBC) data collected on the fresh blood sample. Statistical

significance evaluated by Kruskal-Wallis with Dunn's multiple comparisons test between

all groups, +SEM. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$.

Figure 2

Decreased numbers of naïve CD8 T cells in both young and old mice correlate with increased susceptibility to WNV.

Adult (3-5 months) and old (19-22 months) C57BL/6 mice were infected with WNV, 1000pfu, f.p.; bled on day 7 for FCM; and followed for susceptibility (A, schema). Absolute numbers of naïve CD8 or CD4 T cells were enumerated per mL of blood on day 7 post-infection by FCM (B, C). Data compiled from two independent experiments. Statistical significance evaluated by t-test, median. $*P \leq 0.05$; $**P \leq 0.01$; $****P \leq 0.0001$.

Figure 3

IL-7C treatment expands the total number of T cells available in secondary lymphoid tissues.

Old (19-22 months) C57BL/6 mice were treated with IL-7C on days 1, 3, and 5 of the experiment and sacrificed on days 7, 10, and 14 post-first treatment alongside untreated adult (3-5 months) and old (19-22 months) (schema, A). The total number of lymphocytes (B), CD4 T cells (C), and CD8 T cells (D) are increased at indicated day(s) post-first treatment. The total number of naïve, CD44 low and CD62L high, CD4 and CD8 T cells are not significantly increased at any day post-treatment (E, F). The expansion of CD4 T cells is not dominated by either CM (CD44 high, CD62L high, and CD49d high) or virtual memory (CD44 high, CD62L high, and CD49d low) phenotyped cells (G,H). Both CM and VM phenotype CD8 T cells are expanded at day 10 post-first treatment (I,J). WNV-specific CD8 CTLp (Db-NS4b$_{2488}$) are not reliably increased by IL-7C treatment (K). (B-J) Data shown is representative of two experiments with similar results and statistical significance evaluated by one-way Anova with Dunn's multiple comparisons test. All comparisons are to untreated old mice. $*P \leq 0.05$; $**P \leq 0.01$. (K) IL-7C treated group is day 13 post-first treatment. Data compiled from three independent experiments and statistical significance evaluated by t-test comparison to O, median. $*P \leq 0.05$.

Figure 4

IL-7C treatment of old mice improves WNV-specific CD8 T cell responses without

impacting survival.

Adult (3-5 months), Old (19-22 months), and Old (19-22 months) mice treated with IL-

7C 13 days before challenge are infected with WNV (1000pfu, f.p.; n=45-52 per group).

On day 7 post-WNV infection mice are bled and monitored for survival (schema, A).

Number of WNV-specific CD8 T cells per mL of blood on day 7 post-WNV (B). WNV

survival at day 45 post-infection (C). Number of WNV-specific CD8 T cells from (B)

broken by those that survived and those that died from WNV in each group (C).

Neutralizing antibody titers on day 7 post-WNV broken by those that survived and those

that from died from WNV in each group (E). *In vivo* killing assay on day 7 post-WNV for

NS4b$_{2488}$ (F) and ENV$_{641}$ (G). Total WNV-ENV and DIII-specific memory antibody levels

>45 days post-infection (Figure H, I). Data compiled from two experiments. Statistical

significance evaluated by anova in (B, F, G) and t-test in (D) and (E) as represented on

graphs. Significance of survival evaluated by Log-rank (Mantel-Cox) test. *$P \leq 0.05$; **P

≤ 0.01; ***$P \leq 0.001$; ****$P \leq 0.0001$.

Figure 5

IL7-C treatment does not induce T cell exhaustion.

Adult (3-5 months) and Old (19-22 months) mice were treated with IL-7C 13 days before

infection with WNV (1000pfu, f.p.). On day 7 post-WNV infection mice were bled and

circulating PBMC's were analyzed by FCM for expression of PD1, CD244 (2B4), and

CD223 (LAG-3) on NS4b$_{2488}$+ CD8 T cells (A-C). Data compiled from two experiments.

Statistical significance evaluated by anova in (all groups) and t-test in (by age) as

represented on graphs. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$.

Figure 6

Adoptive transfer of WNV-I Tg CD8 T cells rescues old mice from WNV

susceptibility.

WNV-I CD8 T cells (5400, 1600, or 600) were transferred to groups of old mice (19-22

months) alongside adult (3-5 months) and Old (19-22months) mice the day before WNV

challenge (1000pfu, f.p.). On day 7 post-infection mice were bled and total numbers of

circulating WNV-specific CD8 T cells were enumerated per mL of blood (A). Mice were

monitored for survival to day 45 post-infection (B). The frequency of donor WNV-I CD8

T cells (CD45.1) contributing to the response in (A) is shown in (C). *In vivo* killing assay

on day 7 post-WNV for $NS4b_{2488}$ (D). Data compiled (A-C) or representative (D) of two

experiments. Statistical significance evaluated by anova (A, C, D) with multiple

comparisons test. Significance of survival evaluated by Log-rank (Mantel-Cox) test. *P

≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤ 0.0001.