Increasing temporal variance leads to stable species range limits

John W. Benning
University of Wyoming
Department of Botany
Laramie, WY, USA

Ruth A. Hufbauer
Colorado State University
Department of Agricultural Biology and Graduate Degree Program in Ecology
Fort Collins, CO, USA

Christopher Weiss-Lehman
University of Wyoming
Department of Botany
Laramie, WY, USA

Abstract

What prevents populations of a species from adapting to the novel environments outside the species' geographic distribution? Previous models highlighted how gene flow across spatial environmental gradients determines species expansion vs. extinction and the location of species range limits. However, space is only one of two axes of environmental variation — environments also vary in time, and we know temporal environmental variation has important consequences for population demography and evolution. We used analytical and individual-based evolutionary models to explore how temporal variation in environmental conditions influences the spread of populations across a spatial environmental gradient. We find that temporal variation greatly alters our predictions for range dynamics compared to temporally static environments. When temporal variance is equal across the landscape, the fate of species (expansion vs. extinction) is determined by the interaction between the degree of temporal autocorrelation in environmental fluctuations and the steepness of the spatial environmental gradient. When the magnitude of temporal variance changes across the landscape, stable range limits form where this variance increases maladaptation sufficiently to prevent local persistence. These results illustrate the pivotal influence of temporal variation on the likelihood of populations colonizing novel habitats and the location of species range limits.

Key words: species range limits; temporal stochasticity; adaptation; geographic distributions; invasion; fluctuating selection

Introduction

The limits of species' geographic distributions have long puzzled biologists [1,2]. The causes of distribution, or range, limits fit into two broad categories — either populations have not colonized suitable areas outside their current range margin (a distribution limited by dispersal) or the environment outside that margin is unsuitable, keeping population growth below replacement (a distribution limited by adaptation). The latter case begs the question of why populations do not simply adapt to the novel environmental conditions beyond their range edge. This question is especially perplexing as this exact process — adaptation to novel environments — presumably gave rise to the species' *current* distribution.

Spatial gradients in the environment clearly play a major role in determining range limits — as the environment changes across space, so do the observed flora and fauna. Climbing a mountain slope, one can readily observe how community composition changes with elevation. But *within* species, there is ample evidence for populations successfully adapting to all sorts of abiotic and biotic environmental gradients [3–8]. What causes this adaptive process to cease and a range limit to form? Theory has shed light on the mechanisms underlying the relationship between spatial environmental gradients, local adaptation, and species distributions. For a species occupying a landscape with a spatial gradient (e.g., in soil pH, precipitation, etc.), theory tells us that several mechanisms may constrain adaptation, and subsequently, expansion at the edge of a range. Steep environmental gradients and high gene flow can swamp adaptation at the range edge to create stable range limits [9,but see 10]. Metapopulation dynamics [11] and biotic interactions [12,13] also can enforce range limits. Recent simulation approaches have highlighted how demography, genetic drift, expansion load, and not only the slope, but the shape (e.g., linear vs. non-linear) of the environmental gradient influence adaptation, range expansion,

and the formation of stable range limits [14–17]. In all of these theoretical treatments, the spatial

environmental gradient, which is modeled as a spatial gradient in phenotypic optima, is key to

understanding when populations can expand and when stable range limits form.

However, habitats vary not only in space, but also in time. Indeed, in nature, temporal

variation in the environment is the rule rather than the exception [18–20]. Variation in weather

within and between years provides the most obvious illustration of abiotic temporal variation.

Biotic environments also fluctuate (stochastically or regularly) through time as populations of

predators, mutualists, pathogens, and competitors wax and wane. Both theoretical and empirical

work demonstrate that temporal variation has important consequences for population

demography [21–24,reviewed in 25] and evolution [20,26–31]. Demographic models have

illustrated that a population's extinction risk generally increases as temporal variation increases

[23,32], and experimental work has supported these theoretical predictions [24,33]. Viewed

through an evolutionary lens, temporal variation can generate fluctuating selection due to

phenotypic optima changing through time [34]. These changes in phenotypic optima can lower

expected individual fitness via increases in genetic load, i.e., the reduction in fitness due to an

individual's deviation from an optimal phenotype [35]. With phenotypic optima changing

through time, even a phenotype perfectly adapted to the long-term mean optimum will

experience fitness costs due to genetic load [36,37]. In microcosm experiments, temporally

fluctuating environments have also been shown to impede adaptation to directional

environmental changes, presumably due to relaxed selection during benign periods [30].

However, Holt et al. [38] showed how temporal environmental fluctuations can have positive

effects and facilitate adaptation in sink habitats if the sink environment becomes more benign

long enough to increase population growth and adaptation. Similarly positive effects of temporal

variation were observed in an experiment exposing diatoms to increasing temperatures [31],

where increased population sizes during benign periods resulted in more effective selection.

Though our understanding of the prevalence of fluctuating selection in natural populations is still

incomplete and plagued by sampling error [39,40], several recent rigorous studies do show

strong temporal fluctuations in selection [e.g., 34,41,42], implying fluctuating phenotypic optima

through time.

Because temporal variation can influence key aspects of population demography and

evolution, there is reason to expect it influences species range dynamics, as well [43]. However,

most range limit models to date assume temporally constant environments. Yet in nature, the

environment will always vary in both space and time, and it is easy to imagine myriad ways that

temporal variation might affect the fate of populations spreading across a spatial gradient. If the

environment is changing directionally through time (e.g., glacial advance, warming

temperatures), theory has shown how the rate of environmental change, the amount of genetic

variance, and the steepness of an underlying spatial gradient all influence population persistence

[44,45]. Recent work by Holt et al. [43] provided some of the first insights into the influence of

non-directional temporal variation on range dynamics, showing how temporal variation in

competition can modulate the size of an established species' range, and that temporal variation in

immigration increases the probability of establishment in sink habitats. However, we lack range

limit models exploring the spread of a species across continuous spatial gradients with non-

directional temporal variation in environmental conditions. Such temporal variation in the

environment could slow adaptation along a spatial gradient via fluctuating selection.

Alternatively, positive demographic effects of temporal variation, such as high fecundity in a

relatively favorable year, could boost population sizes and increase range expansion and the

efficacy of natural selection [38]. Temporal variation will directly affect extinction probabilities

via demography [23], and could further influence adaptation by influencing levels of genetic

variation [46]. Importantly, just as environmental variation through space can manifest with

different patterns (e.g., linear / non-linear gradients), environments can vary through time in

different ways. First, environmental variation can exhibit temporal autocorrelation patterns that

are negative (e.g. dry years tend to be followed by wet years and vice versa), positive (e.g. dry

years tend to be followed by another dry year), or reflect uncorrelated random noise [47].

Second, temporal variance may not be equal across space. For instance, positive relationships

between demographic variability and distance from the center of a species' range suggests that

habitats on the edge of the range may sometimes be more temporally variable than habitats in the

core [48,49].

In this paper, we use analytical and simulation models to ask how temporal variation in

the environment influences population expansion and the formation of range limits across spatial

gradients. In our models, temporal variation manifests as temporal stochasticity in phenotypic

optima, with optima fluctuating generation to generation around the patch's long-term mean

optimum. We explore two geographic modes of temporal variation (Fig. 1):

1. **Varying intercept**: all points in space experience environmental change equally through

124   time. This would be the case if, say, temperatures or precipitation changed across a

125   landscape by a similar magnitude and direction year to year (e.g., a landscape-wide

126   drought). It manifests as stochasticity in the intercept of the spatial gradient in phenotypic

127   optima, fluctuating around the long-term mean spatial gradient (Fig. 1b)

128   2. **Varying slope**: the magnitude of environmental change through time increases away

129   from the landscape center. This would be the case if, say, there were increased temporal

variation in inundation at the edge of a wetland compared to the center, or increased

temporal variation in rainfall as one goes from mesic to arid habitat. It manifests as

stochasticity in the slope of the spatial gradient in phenotypic optima, with a constant

intercept at the range center. Thus, the phenotypic optimum displays larger fluctuations

around the long-term average with increasing distance from the range center (Fig. 1c).

We first derive a simple analytical model to describe how, under these two different

geographic modes, temporal variation imposes genetic load on populations across a landscape.

We then explore the influence of temporal environmental variation on range expansion and

dynamics using an individual-based, forward-in-time genetic simulation model. In the model, a

single population colonizes the center of a spatial environmental gradient, and we track the

demography, evolution, and spread of populations across the landscape with the environment

varying in space and time. We vary both the steepness of the underlying spatial environmental

gradient and the geographic mode of temporal variation as defined above. We also vary the

temporal correlation pattern of the environment from negative to positive autocorrelation.

Figure 1. Conceptual diagram illustrating how temporal variation is incorporated in the model. (a) illustrates how spatial environmental variation is modeled in existing range limit models, with a temporally constant spatial gradient in phenotypic optima. In our model, there is intergenerational change in phenotypic optima around the mean optimum at each point in space, according to two "geographic modes" of temporal variation. In the "varying intercept" scenario (b), the intercept of the spatial phenotypic optimum gradient fluctuates such that all patches experience the same magnitude and direction of deviation from the mean optimum each generation. In the "varying slope" scenario (c), the slope of the spatial gradient fluctuates stochastically around the mean slope each generation. The gradient is "anchored" at the center of the landscape, thus, the magnitude of temporal variance increases with distance from the landscape center. The blue dots (b, c) represent the initial founding population in the simulation, which then spreads (or does not spread) across the landscape. Black arrows indicate the magnitude of temporal variance at that point on the landscape [equal at all points in "varying intercept" and increasing with distance from the landscape center in "varying slope"].

Materials and Methods

Data Availability Statement

166   All SLiM and R code, simulation parameters, and the simulation results needed to

167 reproduce the figures in this manuscript, are uploaded for review and will be made available on

168 Figshare upon acceptance.

Expected genetic load due to temporal variation

We first present a simple analytical model to describe how our two geographic modes of

temporal environmental variation (varying intercept and varying slope) affect expected

individual fitness. Consider a species distributed across a landscape, with populations occupying

discrete patches. The environment varies across space (x), producing a spatial gradient in

phenotypic optima with slope b (Fig. 1). Each population comprises individuals perfectly

adapted to the long-term mean phenotypic optimum at that point (patch x) in space. As the

environment varies stochastically through time, mismatches arise between individuals'

phenotypes and the current phenotypic optimum, even though these individuals are perfectly

adapted to the long-term mean phenotypic optimum within their patch. We model this mismatch

as genetic load, which is the fitness cost incurred by an individual due to deviation from the

phenotypic optimum [35]. As in Lande and Shannon [36], we define genetic load for an

individual i as

182
$$L(\theta_{x,t}) = \frac{s}{2}(z_i - \theta_{x,t})^2 \qquad (1)$$

where $\theta_{x,t}$ is the phenotypic optimum in patch x at time t, s is the strength of stabilizing

selection, and z_i is the individual's phenotype. In our varying intercept scenario (Fig. 1b),

$\theta_{x,t} = \varepsilon_t + bx$, while in our varying slope scenario (Fig. 1c), $\theta_{x,t} = (\varepsilon_t + b)x$. In both cases, b

is the slope of the gradient in phenotypic optima, x is the patch position along the gradient, and

ε_t is a normally distributed random variable defining the stochastic deviation in either the

intercept or slope of the spatial environmental gradient.. For now, we assume uncorrelated

temporal fluctuations with a fixed variance of τ (i.e., $\varepsilon_t \sim N(0, \tau)$). For an individual perfectly

adapted to the expected phenotypic optimum (i.e., $z_i = bx$), genetic load in the varying intercept

scenario becomes

$$L(\theta_{x,t}) = \frac{s}{2}\varepsilon_t{}^2 \tag{2}$$

Following mathematical rules for transformations of normally distributed random variables

yields

$$L(\theta_{x,t}) \sim N\left(\frac{s\tau}{2}, \frac{s^2\tau^2}{2}\right) \tag{3}$$

Similarly, genetic load for an individual with $z_i = bx$ in the varying slope scenario is

$$L(\theta_{x,t}) = \frac{s}{2}(\varepsilon_t x)^2 \sim N\left(\frac{s x^2 \tau}{2}, \frac{s^2 x^4 \tau^2}{2}\right) \tag{4}$$

Therefore, the mean and variance in genetic load increase with spatial position (x) in the varying

slope scenario, but are constant in space in the varying intercept scenario.

Effects of temporal variation on range dynamics

The analytical model above describes individuals' expected genetic load due to temporal

stochasticity in phenotypic optima in our two landscape scenarios. However, the model does not

account for gene flow, density dependence, genetic drift, demographic stochasticity, varying

patterns of temporal autocorrelation, or range colonization dynamics. To explore these factors,

we used SLiM [50], a forward-in-time population genetic modeling software, to build a complex,

individual-based simulation of species range dynamics in a spatially and temporally varying

environment. SLiM has recently come to the fore as a flexible, fast, and powerful tool to model

individual genomes under a wide variety of spatially and temporally explicit evolutionary

scenarios. Our simulation model builds most directly on the modeling frameworks of Polechová

and Barton [14] and Bridle et al. [16], which explored the role of spatial environmental gradients

on range dynamics. We build on the insights from those models by asking how temporal

variation in the environment influences population spread and the formation of range limits .

Genetics and mutation

In our simulations, individuals were diploid and either male or female, with obligate

sexual reproduction, and a single chromosome 100,000 bp long. There were two mutation types:

1) neutral mutations, and 2) mutations that contributed additively to a quantitative trait [i.e.,

biallelic quantitative trait loci (QTL) with no dominance]. The overall mutation rate was set to 1

$\times$ 10^{-7} mutations per base position on a gamete per generation (Table S1), and mutations were 10

times more likely to be neutral than QTLs. QTL effect size was drawn from a standard normal

distribution (i.e. $N(0, 1)$). Recombination rate was set to 1 $\times$ 10^{-5} (probability of a cross-over

event between any two adjacent bases per genome per generation).

Mating and population dynamics

223   The simulated landscape comprised a one dimensional array of 201 patches. This is most

224 analogous to a natural landscape that approximates one dimension, like a river, river corridor,

225 mountain ridge, or valley. Each patch hosted a local population subject to density-dependent

226   regulation, with carrying capacity (for a perfectly adapted population) constant across the

227   landscape (here, K = 50). Following Bridle et al. [16], individual fitness (W_i) was calculated as

$$W_i = 2 + 2r(1 - N_x/K) - L(\theta)$$ (5)

Where r is the population maximum rate of increase (set to $r = 0.8$; Table S1), N_x is the number

of individuals in population x, K is the carrying capacity of each patch (set to $K = 50$), and $L(\theta)$

defines the genetic load as defined in the analytical model above (Eq. 1). Thus, the first part of

equation (5) describes standard logistic growth and the second part introduces the fitness cost

scaled by the deviation of an individual's phenotypic value from the local optimum (i.e., genetic

load). For females, W_i defined the mean of a Poisson distribution for the number of offspring

that female produced (female fecundity); for males, W_i defined the likelihood of a male mating

(W_i corresponded to a male's weight in a weighted draw from the pool of available males in a

population). Males could mate multiple times. Any $W_i < 0$ was set to $W_i = 0.00001$ (as the mean

for a Poisson distribution in SLiM must be greater than zero). After mating occurred within a

population, offspring dispersed according to a Poisson dispersal kernel with mean (m) = 0.8. The

direction of dispersal (left or right along the gradient) was unbiased and random. Individuals set

to disperse beyond the edges of the landscape were instead routed to the most distal patch in that

direction. To avoid any potential edge effects, one end condition for our simulations (see below)

was if the population of either of the most distal patches reached half the carrying capacity or

more. It is important to note that dispersal (m) and carrying capacity (K) are both hard to estimate

and highly variable in natural populations; for our main simulations we use values for these

parameters near the mode of their distributions as estimated by Polechová and Barton [14]. We

explored the sensitivity of our results to varying m and K in Supplementary Material S1.

Environmental variation

Environmental variation manifested as changes in the optimum phenotype of the

quantitative trait across space and time. The slope of the spatial environmental gradient (b)

represents the change in phenotypic optima across the landscape (Fig. 1a). To illustrate how b

manifests as a fitness cost of dispersal (i.e., genetic load due to spatial variation in optima), let us

assume $s = 0.125$ (used throughout; Table S1). Then the genetic load incurred by dispersal to

patch $x+1$ for an individual perfectly adapted to patch x is,

255
$$L(\theta) = -0.125b^2/2 \tag{6}$$

Thus if $b = 1$, a female perfectly adapted to patch x will experience a fitness cost of $L(\theta) = $

0.0625 upon migration to patch $x+1$, or a fitness decrease of ~3.125%.

Temporal environmental variation was implemented as intergenerational stochasticity in

the intercept or slope of the spatial gradient in patch phenotypic optima using the same equations

presented for the analytical model above. To incorporate temporal autocorrelation, the random

deviations in intercept or slope were defined as:

262
$$\varepsilon_t = a\varepsilon_{t-1} + d\Phi_t \tag{7}$$

where $\varepsilon_0 = 0$, d scales the magnitude of noise $\Phi_t \sim Normal(0, \tau)$, and a represents the degree of

temporal autocorrelation. When $a = 0$ and $d = 1$, Equation 7 is equivalent to the distribution of

stochastic deviations in phenotypic optima used in the analytical model above. Temporal

autocorrelation could be positive ($0 < a \leq 1$), negative ($-0.99 \leq a < 0$), or uncorrelated

($a = 0$) temporal stochasticity, with autocorrelation lag equal to one generation (Fig. S1). (We

prevented a from going all the way to -1 because, due to $\varepsilon_0 = 0$, this would be identical to $a = $

1.) We set $d = (1 - a^2)^{0.5}$ so that $Var(\varepsilon)$ over the full time series used in our simulations was

approximately equal for all values of a.

τ differed between the two geographic modes of temporal variation. For the "varying

intercept" scenario we set $\tau_{intercept} = 4$. This resulted in temporal variation in selection

gradients, quantified as $\sigma_\beta \approx 0.14$, being within the range of estimates from natural populations

(median $\sigma_\beta = 0.099$ in de Villemeruil et al. [42], averaging across birds and mammals; full

derivation of σ_β presented in Supp mat. S2). For the "varying slope" scenario, we set $\tau_{slope} =$

0.0025 which resulted in patch conditions mimicking the "varying intercept" scenario at 40

patches away from the landscape center (Fig. 2). We also ran simulations with the intercept and

slope varying simultaneously; in this "varying intercept and slope" scenario, respective τ values

remained the same ($\tau_{intercept} = 4$ and $\tau_{slope} = 0.0025$).

Simulation process

Burn-in

Each simulation began with a burn-in period of 20,000 generations. At the start of the

burn-in, 100 genetically homogenous, perfectly adapted, founding individuals were initiated in

the central patch of the landscape. Mating and dispersal happened as described above, but the

landscape was limited to 21 patches wide (10 patches on either side of the founding patch).

Carrying capacity in each patch was 100 individuals (thus, landscape-wide K = 2,100). There

was a modest spatial environmental gradient in optima across these 21 patches (b = 0.5), and

modest, uncorrelated temporal fluctuations in the intercept of the spatial gradient in optima

[deviations drawn from Normal(0,1)]. The goal of this burn-in period was to generate

independent replicates of genetically variable source populations that had evolved in landscapes

of similar spatial and temporal heterogeneity for each simulation run, while allowing enough

time for the different simulation burnins to converge on similar levels of genetic diversity.

During the burn-in, mean heterozygosity of neutral mutations (π) in the central population

usually reached an equilibrium by 10,000 generations (Fig. S2). At generation 20,000, a random

subset of 50 individuals was selected from the 21-patch landscape and migrated to the central

patch for the "founding event". All other individuals were then removed from the simulation.

Main simulation

After the burn-in period, the main simulation began with the prescribed parameters and

the available landscape expanded to 201 patches, with 50 individuals in the founding central

patch. Simulations ended after 20,000 generations, or if all populations went extinct, or if at least

one of the most peripheral landscape patches reached a population size at least half the carrying

capacity (i.e., the species had filled the entire landscape). We ran 1000 simulations for each of

the three temporal variation scenarios (varying intercept, varying slope, varying intercept and

slope) with parameter values pulled randomly from uniform distributions: b[0-3], $a_{intercept}$[-

0.99-1], a_{slope}[-0.99-1]. To compare our model with the model of Polechová & Barton [14], we

also ran 200 simulations, varying b but with $a_{intercept} = 1$ and $a_{slope} = 1$ (i.e., a temporally

constant environment), and compared our results to results predicted from that model (Supp. mat.

S3).

Results

Analytical model

312　　　　As previous work has shown [29,36,38], temporal variation in phenotypic optima results

313　in a genetic load on individuals (Fig. 2). In our varying intercept scenario, this load is constant

314　across space, while in the varying slope scenario, load increases non-linearly with distance from

315　the landscape center. Genetic load increases with the magnitude of temporal variance ( ) and the

316　strength of stabilizing selection (*s*).

Figure 2. Results of the analytical model estimating expected genetic load across space due to
temporal stochasticity in phenotypic optima, for the varying intercept (black; Fig. 1b) and
varying slope (gray; Fig. 1c) scenarios. Solid lines represent lower magnitude of temporal
variance (　　　　and　　　　　　); dashed lines represent higher magnitude of
temporal variance (　　　　and　　　　)

Simulation model

With a temporally constant environment (i.e., $a = 1$) expansion was prevented where the spatial gradient slope (b) was ⬚ 2.5, somewhat steeper than the threshold predicted from the model of Polechová & Barton [14], b 1.96 (Supp. mat. S3).

Varying intercept

When temporal variation in the environment was equal in all patches on the landscape ("varying intercept" scenario; Fig. 1b), a clear relationship emerged between the degree of temporal autocorrelation (a) and the slope of the underlying spatial gradient (b) in determining whether populations spread across the landscape or went extinct (Fig. 3). Range expansion was favored when temporal variation was more positively autocorrelated and spatial gradients were shallow. The ultimate fate of the species in each simulation was either eventual extinction or continual expansion; stable range limits did not form, though the rate of range expansion slowed as spatial gradients steepened (light blue points in Fig. 3). (Here and below we use "species" to describe the group of populations on the landscape.)

As in previous models, steep spatial gradients introduced a strong fitness cost of dispersal, allowing drift to overpower selection and stymie adaptation. Temporal variation exacerbated the negative effects of dispersal by introducing a fitness cost even for individuals well-adapted to their patch's long-term mean optimum; i.e., temporal fluctuations in optima meant that no genotype could completely escape maladaptation across generations. In environments with temporal variation, populations were often able to adapt to the underlying spatial gradient if it was not too steep. However, a large deviation from the mean optimum in a generation had large negative demographic consequences due to individuals being strongly

346   maladapted in most patches. If then the following generation experienced a strongly different

347   optimum, which is more likely as temporal autocorrelation becomes more negative, extinction

348   risk was high (Fig. 3b,c). Thus, temporal variation introduced an extinction risk due to

349   fluctuating phenotypic optima that had strong effects on mean fitness [e.g., 29,36,38].

Figure 3. Range dynamics under the "varying intercept" scenario. (a) When temporal variation
manifests identically across the landscape, range expansion is favored when temporal variation
(*a*, X-axis) is more positively autocorrelated and spatial gradients (*b*, Y-axis) are shallow. (A
temporal autocorrelation of 1 is a temporally static environment, as modeled in previous range
limit models.) Each point represents a simulation (N = 1000). No stable range limits formed in
this scenario; species either continually expanded or went extinct. All blue points are simulations
where species were able to spread, and red/orange points are simulations where species went
extinct; points are colored by how quickly they expanded or went extinct. (b-d) Details of one
simulation (marked with crosshairs in panel a) prior to extinction. Panels track (b) temporal
changes in patch phenotypic optima, (c) landscape-wide population abundance, and (d) median
population genetic variance (for the trait conferring adaptation to the gradient) across generations
prior to extinction (grey dashed line).

Varying slope

When the magnitude of temporal variance increased away from the landscape center

("varying slope" scenario), stable range limits formed (Fig. 4). The exact location of the range

limit fluctuated over time as extinction/colonization dynamics played out at the range edge; thus,

we define stable range limits as the most distal patches on either side of the founding patch that

did not go extinct for at least 950 of the 1000 generations before the end of the simulation (which

lasted 20,000 generations). Range limits formed where increasing temporal variance resulted in

increasing genetic load (Fig. 2), causing populations to fail to adapt to the optimum of edge

patches because temporal fluctuations in optima caused populations in those patches to often go

extinct (or close enough to extinction to strongly undermine genetic variance and adaptive

potential).

Range width (distance between range limits) remained relatively constant across a wide

range of spatial gradient slopes, but then began to decrease with steeper spatial gradients (Fig.

4a). The high fitness costs of dispersal across steep gradients (due to maladaptation) combined

with the negative fitness effects of temporally fluctuating optima to increase extinction risk more

quickly as populations moved away from the center. This led to narrower range widths than for

species spreading across shallower gradients. When temporal variation was positively

autocorrelated, range width generally increased. If that positive autocorrelation was particularly

strong ($a \approx 1$), sometimes the landscape filled completely, especially when the spatial gradient

was shallow (open points in Fig. 4). Because environmental conditions at the landscape core

were stable, species usually only went extinct with very steep spatial gradients.

The results presented above for the varying intercept and varying slope scenarios were

qualitatively robust to changes in carrying capacity (K) and dispersal (m) (Supp. mat. S1). In

general, larger carrying capacities decrease the parameter space where populations go extinct,

likely due to lower demographic stochasticity [e.g., 51] and more efficient selection [e.g., 52]. In

contrast, higher dispersal lowered the maximum spatial gradient slope where expansion was

possible, likely because this introduced a higher mean fitness cost of dispersal [e.g., 14].

409   When both the slope and the intercept of the spatial gradient varied through time, there

410   was a contraction in the parameter space where range expansion was possible (Supp mat. S4;

411   Figs. S8, S9, S10). In general, species persistence and range expansion required positive

412   temporal autocorrelation in the gradient intercept (i.e., $a_{intercept} > 0$). Range width was overall

413   smaller and more variable when the spatial gradient intercept varied along with the slope

414   (compare Figures 4a and S8b).

Figure 4. Range dynamics under the "varying slope" scenario. When temporal variance increases away from the landscape center, stable range limits form where temporal variance is too high to allow further adaptation and persistence. (a) Effects of the spatial gradient slope (X-axis) and degree of temporal autocorrelation (point color) on a species' ultimate range width (Y-axis). (Range width is the number of patches between the species' two stable range limits.) Each point represents a simulation (N = 1000) at the end of 20,000 generations. Open points near the top of the Y-axis are simulations where the species reached at least one edge of the landscape. (b) abundance, (c) temporal variation in optima, (d) genetic variance, and (e) mean phenotype of populations in a single simulation ($b = 0.17$, $a = 0.48$; marked with crosshairs in panel a) after formation of stable range limits (dashed red lines), averaging over the final 1000 generations of the simulation. We define stable range limits as the most distal patches that did not go extinct for at least 950 of the 1000 generations prior to the end of the simulation. (c) shows the standard deviation of the temporal changes in phenotypic optima, to illustrate how temporal variance increases away from the landscape center. In (e), black points show the mean phenotype and the solid blue line indicates the underlying mean spatial gradient in phenotypic optima.

Discussion

Our model results show that, for the parameter space explored here, temporal variation in

the environment substantially alters our predictions for range dynamics across spatially variable

landscapes. When environmental conditions change from generation to generation similarly

across the landscape ("varying intercept" scenario), the ultimate fate of species is determined by

the interaction between the degree of temporal autocorrelation in environmental fluctuations and

the slope of the spatial environmental gradient. As found in previous results [14], species

extinction becomes more likely as the slope of the environmental gradient increases. However,

positive autocorrelation of temporal variation can allow populations to expand along even very

steep gradients. Negative autocorrelation, though, increases demographic fluctuations and

increases extinction risk due to fluctuating phenotypic optima resulting in frequent maladaptation

and high genetic load. When temporal variation in the environment increases toward the

periphery of a landscape ("varying slope" scenario), stable range limits form. Populations persist

where temporal variance is modest, but where temporal variance becomes too large, the

associated genetic load reduces fitness such that population adaptation and persistence is not

possible. The ultimate width of the species' range is primarily a function of the underlying

spatial environmental gradient. When spatial gradients vary in both intercept and slope through

time, range expansion overall becomes less likely. Together these results illustrate how temporal

variation in the environment can have a pivotal influence on the likelihood of a species

colonizing a new landscape and the location of a taxon's range limit.

When temporal variance has no spatial structure ("varying intercept"), we can clearly

delineate the parameter space where colonizing populations go extinct, versus where they expand

across the landscape. When the underlying spatial gradient is shallow, expansion can occur

across a wide range of temporal correlation patterns, but increasingly positive autocorrelation is

required for expansion as the spatial gradient steepens. Environmental noise in nature seems to

largely range from random to positively autocorrelated [47]. Indeed, these simulations indicate

that if environmental conditions were strongly negatively autocorrelated through time,

colonization and expansion would be very rare. Increasing temporal variability is one predicted

(and observed) consequence of contemporary climate change [53], and our simulations suggest

this increased variability could reduce the probability that populations will be able to

successfully track climatic changes via shifting spatial distributions. For example, the upslope

colonization process of an alpine plant due to warming could be stymied if temporal variation is

augmented by climate change. Our model suggests increased temporal environmental variation

due to climate change could further reduce fitness of edge populations, potentially hampering

their ability to track or adapt to changing mean conditions. Furthermore, if a species' current

range limit is due in part to increased temporal environmental variation, then models forecasting

future distributions built solely using the mean, and not variance, of predicted climate will likely

be inadequate.

The biogeographic fact that all species have limited distributions is often at odds with the

ability of evolutionary range limit models to produce *stable* range limits. For instance, when

Barton [10] adjusted Kirkpatrick and Barton's foundational work [9] to allow genetic variance to

evolve, range expansion was continuous and limits failed to form. Polechová & Barton [14]

provided a solution to this conundrum by incorporating genetic drift and demographic

stochasticity in their models, but still found that stable limits only formed with nonlinear spatial

gradients [see also [15,16]]. Similarly, in our model there are no stable range limits when there is

no spatial trend in temporal variance (i.e., varying intercept scenario). However, we do see stable

range limits form when there is a spatial gradient in the magnitude of temporal variance — i.e.,

temporal variance, and the resulting genetic load, increases nonlinearly with distance from the

center of the landscape (varying slope scenario) — even across linear spatial gradients. When

viewed collectively, our results and those of Polechová & Barton [14], Polechová [15], and

Bridle et al. [16] highlight how nonlinear spatiotemporal trends may be key in generating stable

range limits. This work indicates we should perhaps expect qualitatively different patterns in

how environments change through time and space at the edge versus the core of a species' range.

Our results suggest that there is a critical threshold at which temporal variation in the

environment can stop range expansion and enforce a stable range limit. Do we find evidence for

such a pattern in nature? The idea that environments at the edge of species' ranges tend to be

more temporally variable has been assumed more often than empirically shown, but some

demographic studies do suggest range edge habitats to be more temporally variable than range

core habitats [48,49,54–57]. Beyond indirect inference of temporal environmental variation via

demographic data, there are surprisingly few direct measurements of temporal environmental

variability across species ranges. One exception is Eckhart et al. [58], who showed that for the

California annual plant *Clarkia xantiana* ssp. *xantiana*, interannual variability in precipitation

increased toward the subspecies' eastern range margin. Our model supports the notion that

increased variability in precipitation could contribute to the stable range limit observed in this

species.

These results lead to several testable predictions. First, does temporal variation in the

environment increase toward species range edges? For climatic variables this would be fairly

straightforward to test, as we have excellent data for both species distributions and long-term

weather. With such data we could also ask whether, looking across species distributions, we see

a positive relationship between the steepness of putatively important spatial gradients across a

species range and the temporal autocorrelation in that environmental variable. Figure 3 would

suggest this relationship — colonization across steep spatial gradients should only be possible in

fairly stable environments (i.e., environments with positive temporal correlation). For example,

we might expect to observe that species with populations distributed across steep altitudinal

gradients experience more positively correlated temporal variation than species spread across

more shallow spatial gradients (e.g., a primarily latitudinal rather than altitudinal gradient).

Here we have focused on spatio-temporal environmental variation and its influence on

adaptation in populations of a single species. Our understanding of species range dynamics could

be further improved by extending this model to include other genetic, life history, and ecological

factors that can potentially have large influences on population dynamics and spread in nature.

As Antonovics [59] pointed out, evolution of *multiple* traits may often be required for

populations to expand into novel habitats. Genetic correlations between traits whose evolution is

required to colonize novel habitat may greatly influence the probability of colonization [60,61].

Incorporating phenotypic plasticity and its evolution would also be a valuable extension of the

current model [62]. In terms of life history, the effects of overlapping generations may be very

important in temporally variable environments [46], especially if an organism's life cycle

includes dormant stages (e.g., seed banks). The addition of species interactions such as

competition would further illuminate how ecological phenomena interact with evolutionary

processes to modulate range dynamics [12,63,64]. Simulating spread across a patchy landscape,

as opposed to the smooth gradients in trait optima modeled here, could also substantially change

our predictions for population colonization and spread. Lastly, temporal variation as modeled

here is strongly spatially autocorrelated — *i.e.*, adjacent patches always experience temporal

variation in optima in the same direction, positive or negative. It would be fruitful to relax this

assumption to allow less spatial autocorrelation in temporal variation, as the effects of gene flow

on adaptation would likely differ.

Species range limits are as ubiquitous as they are puzzling. What prevents adaptation

from allowing species distributions to continually grow by "accretion like the rings of a tree"

[65]? Recent theory highlights the importance of genetic drift and nonlinear spatial gradients in

controlling range dynamics and the location of stable range limits [14–16]. Here we showed how

temporal variation in the environment, an ever-present feature of natural systems, strongly

contributes to determining whether a population will expand into novel habitat or go extinct, and

can readily enforce stable range limits. By expanding range dynamics models to the temporal

dimension, we gain more realistic, comprehensive insight into the mechanisms and processes

underlying biogeographic patterns, insights of great relevance to invasion biology, the limits to

adaptation, and the fate of populations with environmental change.

Acknowledgements

538 We thank three anonymous reviewers for their insightful comments that greatly improved the
539 manuscript.

Funding

541 This work was supported by grants from the National Science Foundation (DBI-2010892 to

542 JWB, EPS-2019528 to CWL, and DEB-2121980 to RAH). Any opinions, findings, and

543 conclusions expressed in this material are those of the authors and do not necessarily reflect the

544 views of the National Science Foundation.

References

1. Angert AL, Bontrager MG, Ågren J. 2020 What Do We Really Know About Adaptation at Range Edges? *Annu. Rev. Ecol. Evol. Syst.* **51**, 341–361.

2. Sexton JP, McIntyre PJ, Angert AL, Rice KJ. 2009 Evolution and Ecology of Species Range Limits. *Annu. Rev. Ecol. Evol. Syst.* **40**, 415–436.

3. Antonovics J, Bradshaw AD. 1970 Evolution in closely adjacent plant populations. VIII. Clinal patterns at a mine boundary. *Heredity* **25**, 349–362.

4. Etterson JR. 2004 Evolutionary potential of Chamaecrista fasciculata in relation to climate change. I. Clinal patterns of selection along an environmental gradient in the great plains. *Evolution* **58**, 1446–1458.

5. Dice LR. 1940 Ecologic and Genetic Variability within Species of Peromyscus. *Am. Nat.* **74**, 212–221.

6. Johnson NC, Wilson GWT, Bowker MA, Wilson JA, Miller RM. 2010 Resource limitation is a driver of local adaptation in mycorrhizal symbioses. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 2093–2098.

7. Briscoe Runquist RD *et al.* 2020 Context Dependence of Local Adaptation to Abiotic and Biotic Environments: A Quantitative and Qualitative Synthesis. *Am. Nat.* **195**, 412–431.

8. Torres Dowdall J, Handelsman CA, Ruell EW, Auer SK, Reznick DN, Ghalambor CK. 2012 Fine-scale local adaptation in life histories along a continuous environmental gradient in Trinidadian guppies. *Funct. Ecol.* **26**, 616–627.

9. Kirkpatrick M, Barton NH. 1997 Evolution of a species' range. *Am. Nat.* **150**, 1–23.

10. Barton N. 2001 Adaptation at the edge of a species' range. In *Integrating ecology and evolution in a spatial context*, books.google.com.

11. Holt RD, Keitt TH. 2000 Alternative causes for range limits: a metapopulation perspective. *Ecol. Lett.*

12. Case TJ, Taper ML. 2000 Interspecific Competition, Environmental Gradients, Gene Flow, and the Coevolution of Species' Borders. *Am. Nat.* **155**, 583–605.

13. Case TJ, Holt RD, McPeek MA, Keitt TH. 2005 The community context of species' borders: ecological and evolutionary perspectives. *Oikos*

14. Polechová J, Barton NH. 2015 Limits to adaptation along environmental gradients. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 6401–6406.

15. Polechová J. 2018 Is the sky the limit? On the expansion threshold of a species' range. *PLoS Biol.* **16**, e2005372.

16. Bridle JR, Kawata M, Butlin RK. 2019 Local adaptation stops where ecological gradients steepen or are interrupted. *Evol. Appl.* **12**, 1449–1462.

17. Gilbert KJ, Sharp NP, Angert AL, Conte GL, Draghi JA, Guillaume F, Hargreaves AL, Matthey-

Doret R, Whitlock MC. 2017 Local Adaptation Interacts with Expansion Load during Range
Expansion: Maladaptation Reduces Expansion Load. *Am. Nat.* **189**, 368–380.

18. Ruokolainen L, Lindén A, Kaitala V, Fowler MS. 2009 Ecological and evolutionary dynamics under
coloured environmental variation. *Trends Ecol. Evol.* **24**, 555–563.

19. Toszogyova A, Storch D. 2019 Global diversity patterns are modulated by temporal fluctuations in
primary productivity. *Glob. Ecol. Biogeogr.* **28**, 1827–1838.

20. Bell G. 2010 Fluctuating selection: the perpetual renewal of adaptation in variable environments.
Philos. Trans. R. Soc. Lond. B Biol. Sci. **365**, 87–97.

21. Boyce MS, Haridas CV, Lee CT, The Nceas Stochastic Demography Working Group. 2006
Demography in an increasingly variable world. *Trends Ecol. Evol.* **21**, 141–148.

22. Fullerton MR, Walters JR, Will RE, Loss SR. 2021 Interannual climate variation influences nest
initiation date and nest productivity of the Red-cockaded Woodpecker at the northwestern edge of its
range. *Ornithological Applications* **123**, duab013.

23. Ripa J, Lundberg P. 1996 Noise colour and the risk of population extinctions. *Proceedings of the
Royal Society of London. Series B: Biological Sciences* **263**, 1751–1753.

24. Rescan M, Grulois D, Ortega-Aboud E, Chevin L-M. 2020 Phenotypic memory drives population
growth and extinction risk in a noisy environment. *Nat Ecol Evol* **4**, 193–201.

25. Lande R, Engen of BS, Engen S, Sæther B-E, Saether of PE. 2003 *Stochastic Population Dynamics
in Ecology and Conservation*. Oxford University Press.

26. Bürger R, Lynch M. 1995 Evolution and extinction in a changing environment: A quantitative-
genetic analysis. *Evolution* **49**, 151–163.

27. Hairston NG, Dillon TA. 1990 Fluctuating selection and response in a population of freshwater
copepods. *Evolution* **44**, 1796–1805.

28. Grant PR, Grant BR. 1995 Predicting microevolutionary responses to directional selection on
heritable variation. *Evolution* **49**, 241–251.

29. Chevin L-M, Cotto O, Ashander J. 2017 Stochastic Evolutionary Demography under a Fluctuating
Optimum Phenotype. *Am. Nat.* **190**, 786–802.

30. Hao Y-Q, Brockhurst MA, Petchey OL, Zhang Q-G. 2015 Evolutionary rescue can be impeded by
temporary environmental amelioration. *Ecol. Lett.* **18**, 892–898.

31. Schaum C-E, Buckling A, Smirnoff N, Studholme DJ, Yvon-Durocher G. 2018 Environmental
fluctuations accelerate molecular evolution of thermal tolerance in a marine diatom. *Nat. Commun.*
9, 1719.

32. Wissel C, Stöcker S. 1991 Extinction of populations by random influences. *Theor. Popul. Biol.* **39**,
315–328.

33. Pimm SL, Jones HL, Diamond J. 1988 On the Risk of Extinction. *Am. Nat.* **132**, 757–785.

34. Chevin L-M, Visser ME, Tufto J. 2015 Estimating the variation, autocorrelation, and environmental

sensitivity of phenotypic selection. *Evolution* **69**, 2319–2332.

35. Whitlock MC, Davis B. 2011 Genetic Load. *eLS*. (doi:10.1002/9780470015902.a0001787.pub2)

36. Lande R, Shannon S. 1996 THE ROLE OF GENETIC VARIATION IN ADAPTATION AND
POPULATION PERSISTENCE IN A CHANGING ENVIRONMENT. *Evolution* **50**, 434–437.

37. Chevin L-M. 2013 Genetic constraints on adaptation to a changing environment. *Evolution* **67**, 708–
721.

38. Holt RD. 2004 Temporal variation can facilitate niche evolution in harsh sink environments. *Am.*
Nat. **164**, 187–200.

39. Siepielski AM, DiBattista JD, Carlson SM. 2009 It's about time: the temporal dynamics of
phenotypic selection in the wild. *Ecol. Lett.* **12**, 1261–1276.

40. Morrissey MB, Hadfield JD. 2012 Directional selection in temporally replicated studies is
remarkably consistent. *Evolution* **66**, 435–442.

41. Bonnet T, Postma E. 2018 Fluctuating selection and its (elusive) evolutionary consequences in a
wild rodent population. *J. Evol. Biol.* **31**, 572–586.

42. de Villemereuil P *et al.* 2020 Fluctuating optimum and temporally variable selection on breeding
date in birds and mammals. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 31969–31978.

43. Holt RD, Barfield M, Peniston JH. 2022 Temporal variation may have diverse impacts on range
limits. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **in press**.

44. Pease CM, Lande R, Bull JJ. 1989 A Model of Population Growth, Dispersal and Evolution in a
Changing Environment. *Ecology* **70**, 1657–1664.

45. Polechová J, Barton N, Marion G. 2009 Species' range: adaptation in space and time. *Am. Nat.* **174**,
E186–204.

46. Ellner S, Hairston NG. 1994 Role of Overlapping Generations in Maintaining Genetic Variation in a
Fluctuating Environment. *Am. Nat.* **143**, 403–417.

47. Vasseur DA, Yodzis P. 2004 The color of environmental noise. *Ecology* **85**, 1146–1152.

48. Gerst KL, Angert AL, Venable DL. 2011 The effect of geographic range position on demographic
variability in annual plants. *J. Ecol.* , no–no.

49. Angert AL. 2009 The niche, limits to species' distributions, and spatiotemporal variation in
demography across the elevation ranges of two monkeyflowers. *Proc. Natl. Acad. Sci. U. S. A.* **106**,
19693–19698.

50. Haller BC, Messer PW. 2019 SLiM 3: Forward Genetic Simulations Beyond the Wright-Fisher
Model. *Mol. Biol. Evol.* **36**, 632–637.

51. Lande R. 1993 Risks of Population Extinction from Demographic and Environmental Stochasticity
and Random Catastrophes. *Am. Nat.* **142**, 911–927.

52. Samani P, Bell G. 2010 Adaptation of experimental yeast populations to stressful conditions in

relation to population size. *J. Evol. Biol.* **23**, 791–796.

53. IPCC. 2021 *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to*
the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge
University Press.

54. Curnutt JL, Pimm SL, Maurer BA. 1996 Population Variability of Sparrows in Space and Time.
Oikos **76**, 131–144.

55. Doak DF, Morris WF. 2010 Demographic compensation and tipping points in climate-induced range
shifts. *Nature* **467**, 959–962.

56. Villellas J, Morris WF, García MB. 2013 Variation in stochastic demography between and within
central and peripheral regions in a widespread short-lived herb. *Ecology* **94**, 1378–1388.

57. Williams CK, Ives AR, Applegate RD. 2003 Population dynamics across geographical ranges: Time-
series analyses of three small game species. *Ecology* **84**, 2654–2667.

58. Eckhart VM, Geber MA, Morris WF, Fabio ES, Tiffin P, Moeller DA. 2011 The geography of
demography: long-term demographic studies and species distribution models reveal a species border
limited by adaptation. *Am. Nat.* **178**, S26–43.

59. Antonovics J. 1976 The Nature of Limits to Natural Selection. *Ann. Mo. Bot. Gard.* **63**, 224–247.

60. Etterson JR, Shaw RG. 2001 Constraint to adaptive evolution in response to global warming. *Science*
294, 151–154.

61. Duputié A, Massol F, Chuine I, Kirkpatrick M, Ronce O. 2012 How do genetic correlations affect
species range shifts in a changing environment? *Ecol. Lett.* **15**, 251–259.

62. King JG, Hadfield JD. 2019 The evolution of phenotypic plasticity when environments fluctuate in
time and space. *Evol Lett* **3**, 15–27.

63. Benning JW, Moeller DA. 2019 Maladaptation beyond a geographic range limit driven by
antagonistic and mutualistic biotic interactions across an abiotic gradient. *Evolution* **73**, 2044–2059.

64. Legault G, Bitters ME, Hastings A, Melbourne BA. 2020 Interspecific competition slows range
expansion and shapes range boundaries. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 26854–26860.

65. Mayr E. 1963 *Animal species and evolution*. Belknap Press of Harvard University Press.

678