Kv4.3 channel downregulation mediates chronic post-lesional pacemaker acceleration in surviving dopamine substantia nigra neurons

Lora Kovacheva[1], Josef Shin, Navid Farassat[*] & Jochen Roeper[1]

Institute of Neurophysiology, Neuroscience Center, Goethe University, Frankfurt, Germany.

*Present address: Eye Center, Medical Center, Medical Faculty, Albert-Ludwig University Freiburg, Germany

[1] Corresponding authors:

Prof. Dr. Jochen Roeper & Lora Kovacheva

Institute for Neurophysiology

Neuroscience Center, Goethe University

Theodor-Stern-Kai 7, 60590 Frankfurt, Germany

e-mail: roeper@em.uni-frankfurt.de & email: lorask@gmail.com

Tel: +49-69-6301-84091/92

Abstract

Substantia nigra dopamine (SN DA) neurons are progressively lost in Parkinson disease (PD). While the molecular and cellular mechanisms of their differential vulnerability and degeneration have been extensively studied, we still know very little about potential functional adaptations of those SN DA neurons that – at least for some time – manage to survive during earlier stages of PD. We utilized a partial lesion 6-OHDA mouse model to characterize initial electrophysiological impairments and chronic adaptations of surviving identified SN DA neurons, both *in vivo* and *in vitro*. Early after lesion (3 weeks), we detected a selective loss of *in vivo* burst firing in surviving SN DA neurons, which was accompanied by *in vitro* pacemaker instability. In contrast, late after lesion (>2 months), *in vivo* firing properties of surviving SN DA neurons had recovered in the presence of 2-fold accelerated pacemaking *in vitro*. Finally, we show that this chronic cell-autonomous adaptation in surviving SN DA neurons was mediated by Kv4.3 channel downregulation. Our study demonstrates substantial homeostatic plasticity of surviving SN DA neurons after a single-hit non-progressive lesion, which might contribute to the phenotype of initially surviving SN DA neurons in PD.

Keywords

dopamine, 6-OHDA, Kv4.3 channel, substantia nigra, accelerated pacemaker

Introduction

One clinically relevant hallmark of Parkinson disease (PD) is the progressive degeneration of vulnerable neuronal populations, including substantia nigra dopamine (SN DA) neurons (Giguere et al., 2018; Kalia and Lang, 2015; Poewe et al., 2017). At the time of clinical diagnosis, more than 50% of SN DA neurons, in particular those in ventrolateral regions, are already lost (Gibb and Lees, 1991; Kordower et al., 2013). Moreover, Kordower and colleagues (Kordower *et al.*, 2013) demonstrated that most remaining SN DA neurons were rapidly lost (or lose their DA phenotype) within the first few years of clinically manifest PD. While neuropathological studies map the time course of degeneration with cellular resolution and also allow for a molecular characterization of the remaining neurons, they cannot investigate the functional properties of surviving neurons. *In vivo* recordings of putative human SN DA neurons in the context of DBS surgery provided a first glimpse of electrically active DA survivors (Dragicevic et al., 2015; Schiemann et al., 2012; Zaghloul et al., 2009), but a systematic investigation of these DA neurons is not feasible in a therapeutic setting. Thus, studying functional adaptations of surviving DA neurons in the context of PD is the domain of animal models, which themselves often capture only partial aspects of the disease. Among the various rodent PD models, 6-hydroxydopamine (6-OHDA) lesions have been extensively used to study the DA-depleted basal ganglia (Chen et al., 2021; Fieblinger et al., 2018; Kravitz et al., 2010; Parker et al., 2018; Parker et al., 2016; Rubi and Fritschy, 2020; Sharott et al., 2005; Sitzia et al., 2020; Ungerstedt, 1968; West et al., 2018). By comparison, a smaller number of studies applying only partial lesions focused on the properties of the remaining DA midbrain system, including its long-term adaptations and drug responses (Bez et al., 2016). These studies demonstrated that – in contrast to PD – 6-OHDA DA lesions are non-progressive (i.e. single-hit) and follow a stereotypical time course: an initial phase dominated by cell loss, inflammation and behavioral impairment for about 3 weeks, which is followed by a chronic phase (studied for up to 20 month in mice), where partial behavioral recovery, neurochemical and molecular adaptations and – to a certain degree – axonal sprouting of DA neurons occur (Bez *et al.*, 2016; Cenci and Bjorklund, 2020; Kirik et al., 1998; Schwarting and Huston, 1996; Winkler et al., 2002). Single cell

electrophysiological investigations of viable post-lesional DA neurons, however, have been

exceedingly rare with only Hollerman & Grace providing a first data set (Hollerman and Grace, 1990).

They found 6-0HDA-lesion-size-dependent changes in *in vivo* firing properties of putative DA neurons

in rats, among them a decrease of *in vivo* burst firing after 50-60% (sized) lesions (Hollerman and

Grace, 1990).

In contrast, cell-autonomous characteristics of surviving SN DA neurons after 6-OHDA lesion have not

yet been studied. Here, we aim to close this gap by exploring the electrophysiological properties *in*

vivo and *in vitro* of surviving SN DA neurons in a well-characterized, intra-striatal unilateral 6-OHDA

lesion mouse model, where about 50% of SN DA neurons survive. We selected two time points of

electrophysiological investigation based on our long-term behavioral characterization of motor

phenotypes. In particular, we studied the dynamics of spontaneous locomotion and turning behavior

in a drug-free setting. This identified a 3-week post-lesion time point (i.e. early phase), where motor

impairment was prominent, and a stable late phase (>68 days post-lesion), where turning asymmetry

had completely recovered. In contrast to the significant behavioral differences between early and late

post-lesional time points, the extent of the striatal lesion and the degree of stereologically-determined

TH-positive cell numbers were stable. This pointed to a relevant contribution of early functional

impairment and late adaptation in a stable pool of surviving SN DA neurons as detailed below.

Results

Early behavioral impairments and long-term adaptations after partial lesion of substantia nigra dopamine neurons

In order to identify potential compensatory adaptations in surviving SN DA neurons, we utilized a C57Bl6N mouse model with a partial 6-OHDA lesion of the nigrostriatal pathway. The neurotoxin 6-OHDA was unilaterally infused into the right dorsal striatum, causing a partial degeneration of DA neurons in the ipsilateral substantia nigra, which was quantified by unbiased stereology (Fig. 1A). We continuously characterized motor behavior of the model (i.e. open field, cylinder test) until two different time points, either 21-days post-lesion (early phase) or more than 60 days post-lesion (late phase). Initially, animals were sacrificed for TH immunohistochemistry in midbrain and striatum. As shown in Fig. 1B, at the early phase, the number of TH-positive (i.e. dopaminergic) SN neurons was reduced post-6-OHDA (lower left panels) compared to control mice, where only vehicle (ACSF with ascorbic acid) was infused into the dorsal striatum (upper left panels). The right panel in Fig. 1B displays the stereological results of TH-positive SN neurons for N=9 vehicle-treated and N=9 6-OHDA-infused mice. The data were normalized to the stereological results of respective contralateral SN DA neurons. In contrast to vehicle, where no ipsilateral loss of SN DA neurons was detected, the number of ipsilateral surviving SN DA neurons in the early post-6-OHDA phase was on average reduced to about 40% (Fig. 1B; vehicle: 100 ± 2.644 %; 6-OHDA: 36.85 ± 3.463%; p < 0.0001, Mann-Whitney test). We also carried out the unbiased stereology for TH-positive SN neurons in the late phase and detected a similar degree of loss (Suppl. Fig. 1C, vehicle: 102.6 ± 7.9 %; 6-OHDA: 39.96 ± 3.7 %; p = 0.0262, Mann-Whitney test). These results indicated that SN DA cell loss was stable throughout our observation period of more than 2 months. In addition, we analyzed the axonal compartment of DA midbrain neurons by determining striatal TH-optical densities, both in early and late phase (Suppl. Fig. 1). Similar to cell counts, we found an about 50% reduction of TH-immunosignal in the ipsilateral dorsal

striatum, both in early and late phase, whereas the ventral striatum was only mildly affected (ca. 20%

reduction). In summary, our model induces a stable loss of SN DA neurons, associated with a stable

reduction of striatal TH-immunoreactivity throughout the entire observation period. In other words,

in accordance with previous chronic 6-OHDA studies (Bez *et al.*, 2016; Cenci and Bjorklund, 2020;

Schwarting and Huston, 1996), this model showed no progression and is therefore suitable to study

functional adaptations in a stable pool of surviving SN DA neurons.

Fig 1 C shows our continuous behavioral monitoring of unilateral 6-OHDA and vehicle-treated mice

before and up to 68 days post-infusion. Based on previous studies (Cenci and Bjorklund, 2020;

Schwarting and Huston, 1996), we focused on the dynamics of drug-free spontaneous turning

behavior during open field locomotion (Fig. 1C, lefts panels). In contrast to vehicle-infused mice, which

displayed a stable symmetric ratio (ca. 50%) of ipsi- to contral-lateral turning throughout the entire

experiment, 6-OHDA-infused mice showed a dramatic shift to ipsilateral turning immediately after

treatment (Figure 1C, right panel). Interestingly, ipsilateral turns occurred in long sequences of up to

40 individuals turns, a pattern not observed in controls (Suppl. Fig. 2). Importantly for this study, these

turning sequences as well as the overall ipsilateral bias gradually but eventually completely recovered

over 2-months post treatment (two-way-ANOVA, p-value across time $p<0.0001$, p-value across groups

$p<0.0001$, significant difference between vehicle and 6-OHDA group till day 40, Šídák's multiple

comparisons test). Analysis of turning features, such as diameter or velocity, revealed that the

recovered contralateral turns were similar to turns performed by vehicle-treated mice (Suppl. Fig. 3).

This could be indicative for a degree of functional recovery and/or functional compensation within the

stable pool of surviving DA neurons. General locomotion, measured as performed track length per

session, recovered in parallel to the turning bias (Suppl. Fig. 1). However, other motor behaviors, such

as contralateral paw use during rearing in the cylinder test, did not show recovery throughout the

experimental period (Fig. 1D) (21st post-infusion day: vehicle $38.5 \pm 5.24\%$, 6-OHDA $5.73 \pm 1.35\%$, $p <$

0.0001, Mann-Whitney test; 68th post-infusion day: vehicle $63.67\pm 6.6\%$, 6-OHDA $3.167 \pm 1.8\%$, $p <$

0.0001, Mann-Whitney test), which is consistent with the partial but stable loss of SN DA neurons and

reduction of dorsal striatal TH.

To investigate this model's homeostatic mechanisms from the perspective of the surviving SN DA

neurons, we performed *in vivo* extracellular recordings and *in vitro* whole-cell patch-clamp recordings

once within a period with strong turning bias, at the 21st day after lesion (early phase), and at a later

time point – after reaching stable baseline levels (late phase).

Figure 1. Unilateral striatal 6-OHDA mouse model with stable loss of SN DA neurons resulted in

delayed partial behavioral recovery.

(A) Experimental design, illustrating timeline of behavioral assays with termination points – groups

ended either on the 21st, or later than the 64th post-infusion day. (B) Left panel: TH-DAB staining of

the SN, 10x magnification and 60x magnification for vehicle and 6-OHDA-infused mouse. Scalebar left

200µm, scalebar right 25µm. Right panel: ratio of ipsilateral (infusion side) to contralateral side of

surviving TH-positive neurons in SN at 21st post-infusion day. (C) Left: spontaneous locomotion of mice

in open field arena for a 10 min session. Left, examples of an ACSF-infused mouse (vehicle), right

example of a 6-OHDA-infused mouse at baseline (upper panels), 4th post-infusion day (middle panels)

and 68th post-infusion day (lower panels). Right: ratio of ipsilateral to contralateral turning behavior

for all experimental groups plotted against session days. Note recovery in the 6-OHDA-treated mice

from day 4 to day 68 after initial strong turning bias (>90%). (D) Cylinder test quantified by ratio of

contra- to ipsilateral forepaw use. Note significant loss of contralateral forepaw involvement, both at

21 and 64 days. All data are presented as mean ± standard error of mean (SEM).

Surviving SN DA neurons in the early post-6-OHDA phase exhibit impaired firing properties both

in vivo and *in vitro*

In the early post-6-OHDA phase, we explored the *in vivo* spontaneous activity of surviving identified

SN DA neurons in isoflurane-anesthetized mice, using single-unit extracellular recordings combined

with juxtacellular labelling and post-hoc TH immunohistochemistry (see schema Fig. 2A). Figure 2B

shows the *in vivo* extracellular electrical activity (top panel) and identification of a representative SN

DA neuron (lower right panel) from a control mouse, 3 weeks after vehicle infusion. Note the typical

combination of irregular single spike firing with occurrence of transient fast burst discharges (bursts

were defined by the 80/160ms Grace criteria; see methods for details). The interspike interval (ISI)

histogram, which captures ISIs from the entire recording time of 10 min for this cell, displays the typical

features of an *in vivo* bursty DA neuron, i.e. a bimodal ISI distribution representing intra- and interburst

intervals over the dynamic range between ~30-1000 ms (lower left panel). This recorded and labelled

cell was localized in the medial SN and characterized as TH-immunopositive. In comparison, Fig. 2C

displays representative *in vivo* activity of a surviving identified SN DA neuron 3 weeks after 6-OHDA-infusion. Here, while irregular single-spike pacemaking was present similar to controls, fast burst events were almost completely absent (top panel). In accordance, the ISI histogram of this neuron, for the entire 10 min of recording, showed a unimodal, Gaussian-like distribution without fast intraburst and slow interburst intervals (Fig. 2C, lower left panel). This recorded and labelled cell was also localized in the medial SN and identified as TH-immunopositive.

Similar to these two example cells, all *in vivo* recorded DA neurons from the early phase were successfully juxtacellularly labeled with neurobiotin (NB), post-hoc immunohistochemically reconstructed and localized within the ventral midbrain (Fig 2D, vehicle: n =28, N =15; 6-OHDA: n =18, N =16). For best comparison, we restricted our analysis to those DA midbrain neurons localized in the medial half of the SN (mSN), where a sufficient number of surviving and electrically active DA neurons could be sampled after 6-OHDA lesion. In contrast, we only managed to detect a very small number of active DA neurons in the lateral SN (lSN), the most vulnerable part of the DA midbrain (Figure 2D). Consequently, Figure 2 E-H compares the group data of surviving mSN DA neurons at early phase between vehicle and 6-OHDA treatment. We detected only a small reduction (< 40 %) of overall firing rate for post-6-OHDA mSN DA neurons compared to vehicle-treated controls (Fig.2E; vehicle: mean frequency = 7.5 ± 0.92 Hz, 6-OHDA: mean frequency = 4.71 ± 0.6 Hz, p = 0.0184, Mann-Whitney test) and no differences in overall firing regularity expressed as CV (Fig. 2F; vehicle: CV = 53.96 ± 7.78 %, 6-OHDA: CV = 36.06 ± 3.7 %, p = 0.1694, Mann-Whitney test). In contrast, the reduction of bursting - expressed as the percentage of spikes fired in bursts (SFB) and burst rate - was dramatic (ca. 10-fold) and highly significant for mSN DA neurons in the 6-OHDA group compared to vehicle controls (Fig. 2G, vehicle: SFB = 31.22 ± 7.71; 6-OHDA: SFB = 2.24 ± 1.09 %; p = 0.0032, Welch's t test; Fig. 2H, vehicle: burst rate = 0.34 ± 0.09 Hz, 6-OHDA: burst rate = 0.04 ± 0.02 Hz; p = 0.0066, Welch's t test). All other analyzed firing parameters identified no additional differences between the two early groups (Suppl. Fig. 4). In summary, we detected a dramatic compression of the dynamic *in vivo* firing range in mSN DA neurons surviving the 6-OHDA lesion. In addition to the degree of DA cell loss, this functional

impairment of surviving DA neurons might be a novel contributing factor to the extensive motor

impairment, observed during the early post-6-OHDA phase.

Burst discharge in DA neurons is orchestrated by the interplay of patterned synaptic inputs with their

intrinsic excitability, giving rise to different types of bursting (Otomo et al., 2020). To identify a possible

contribution of cell-autonomous changes in intrinsic excitability of surviving mSN DA neurons to their

impaired *in vivo* bursting, we also studied these cells *in vitro* in synaptic isolation (see Schema 2I).

Figure 2J shows representative stable and regular pacemaker activity of identified mSN DA neuron

from a vehicle-infused control (lower right panel), as well as a subthreshold response to

hyperpolarizing current injection (lower left panel). In contrast, pacemaker activity of mSN DA neurons

post-6-OHDA infusion was unstable and characterized by intermittent periods of action potential

failure (see Fig. 2K, upper panel). However, subthreshold responses were similar to controls (lower

right panel). Analogous to the *in vivo* experiments, all *in vitro* recorded DA neurons were neurobiotin

(NB) filled, identified, and mapped, resulting in an equivalent, medial clustering of surviving SN DA

neurons (Fig. 2L). When *in vitro* pacemaker properties were compared between identified mSN DA

neurons from the vehicle and the 6-OHDA groups (Fig. 2L), we detected no differences in mean firing

rates (Fig. 2M, vehicle: firing rate = 1.699 ± 0.17 Hz, 6-OHDA: firing rate = 2.425± 0.25 Hz, p = 0.1448,

Mann-Whitney test), rebound delays and sag-amplitudes (Fig. 2O, vehicle: rebound delay = 584.9 ±

95.8ms, 6-OHDA: rebound delay = 429.6 ± 56.1ms, p = 0.1738, Mann-Whitney test; Fig. 2P, vehicle:

sag-amplitude = 15.3 ± 1.1 mV, 6-OHDA: sag-amplitude = 13.5 ± 0.96 mV, p = 0.1395, Mann-Whitney

test), but a highly significant 5-fold increase of pacemaker irregularity expressed as CV (Fig.2N, vehicle:

CV = 6.531 ± 0.7147 %; 6-OHDA: CV = 33.99 ± 5.788; p = 0.0007, Mann-Whitney test). More detailed

analysis revealed that the increased CV was mainly mediated by intermittent phases of firing failure,

similar to the one shown in Fig. 2K (Suppl. Fig. 5). Further analysis, including a comparison of on-cell

and whole-cell recordings, did not reveal additional differences. In short, we provided the first data

set in the literature that surviving SN DA neurons 3-weeks after a 6-OHDA infusion are functionally

impaired displaying a 10-fold reduced *in vivo* burst activity. This was accompanied by a 5-fold decrease

of *in vitro* cell-autonomous pacemaker stability. In order to address the question whether these

altered properties were stable adaptions or reflected only transient impairment, we also studied *in*

vivo and *in vitro* electrophysiology of surviving mSN DA neurons in the late phase, characterized by

partial behavioral recovery.

Figure 2. Surviving SN DA neurons at early post-6-OHDA phase exhibited a 10-fold decrease in *in*

***vivo* bursting and a 5-fold decree in *in vitro* regularity**

(A), (I) Experimental design, illustrating timeline of behavioral assays, followed by terminal *in vivo* juxtacellular recordings (A) or by terminal *in vitro* whole-cell recordings (I) at the 21st post-infusion day. (B), (C) Top: 10s original recording trace of spontaneous *in vivo* single-unit activity from SN DA neurons in vehicle (B) and 6-OHDA-infused mouse (C). Scalebars: 1s, 0.2mV. Below, left: corresponding ISI histograms. Below, right: corresponding confocal images of juxtacellularly labelled and immunohistochemically identified DA neuron. Note the sparse bursting of the surviving SN DA neuron from the 6-OHDA-infused mouse at early phase. (D), (L) Anatomical mapping of all extracellularly recorded and juxtacellularly labelled DA neurons (D), and of all in vitro recorded and filled DA neurons (L), projected to bregma -3.08 mm. Location of example SN DA neurons in (B), (C), (J), (K) are highlighted. Smaller symbols represent DA neurons that have been recorded and identified, but not included in the group data analysis in (E-H). (E-H) Scatter dot-plots, showing significant decrease of *in vivo* mean firing rate (E), percentage of SFB (G) and burst rate (H) and no significant differences in CV (F) between the vehicle and 6-OHDA-infused mice. Note the 10-fold decrease in SFB for 6-OHDA-infused mice. (J), (K) Top: 10s original recording trace of *in vitro* whole-cell recording of spontaneous SN DA neuron activity in a vehicle (J) and 6-OHDA-infused (K) mouse. Scalebars: 1s, 20mV. Note that the 6-OHDA DA neuron has a highly irregular pacemaking. Below, left: corresponding to hyperpolarizing current injection. Below, right: confocal images of NB filled and immunohistochemically identified DA neuron. (M-P) Scatter dot-plots, showing no difference in *in vitro* mean firing rate (M), rebound delay (O) or sag-component (P) and a 5-fold increase in CV (N). Immunohistochemical imaging for all four DA neurons are displayed in 10x and 60x magnifications (green, TH; red, NB), scalebars: 200µm, 20µm. All data are presented as mean ± standard error of mean (SEM).

Surviving mSN DA neurons in the late 6-OHDA phase display full recovery of *in vivo* burst firing associated with accelerated *in vitro* pacemaking

We studied and analyzed surviving mSN DA neurons in the late 6-OHDA phase in an identical fashion to the early phase (compare Fig. 3A/I and Fig. 2A/I). In contrast to the 10-fold difference in burst rate in early phase, we found no significant electrophysiological *in vivo* differences between the two groups in the late phase (compare Fig 3B with Fig. 3C). Among others (Suppl. Fig. 6), this implies that mean firing frequencies, CV, SFB and burst rates were similar in chronically surviving mSN DA neurons compared to controls (Fig. 3E, vehicle: mean firing rate = 6.95 ± 0.82 Hz; 6-OHDA: mean firing rate = 7.53 ± 1.12 Hz; p = 0.6038, Mann-Whitney test; Fig. 3F, vehicle: CV = 54.88 ± 12.23% ; 6-OHDA: CV = 53.33 ± 12.65%, p > 0.9999, Mann-Whitney test; Fig. 3G, vehicle: SFB = 20.2 ± 9.11%, 6-OHDA: SFB = 22.17 ± 8.87%, p = 0.8784, Mann-Whitney test; Fig. 3H, vehicle: burst rate = 0.1728 ± 0.08 Hz; 6-OHDA: burst rate = 0.3071 ± 0.15Hz, p = 0.4308, Mann-Whitney test). This suggests that surviving mSN DA neurons might have fully recovered more than 2-months after lesion. However, the *in vivo* results cannot distinguish between simple recovery (e.g. functional restoration) or adaptive homeostatic plasticity. Therefore, we also studied *in vitro* pacemaker properties of the DA neurons in the late post-6-OHDA phase. Here, we found clear evidence for chronic functional remodeling of surviving mSN DA neurons. When comparing intrinsic pacemaker frequencies, we noticed that late 6-OHDA survivors discharged almost 2-fold faster compared to vehicle-controls (compare Fig. 3J with 3K, Fig. 3M vehicle: firing rate = 1.474 ± 0.1653 Hz; 6-OHDA: firing rate = 2.713 ± 0.2057 Hz, p = 0.0016, Mann-Whitney test), while maintaining a stable and regular pattern (Fig. 3N, vehicle: CV = 9.515 ± 1.7 %, 6-OHDA: CV = 16.10 ± 6.661 %, p = 0.175, Mann-Whitney test). In line with this pacemaker acceleration, the sag component was increased, and the rebound delay shortened (Fig 3O, vehicle: rebound delay = 629.6 ± 96.38 ms, 6-OHDA: rebound delay = 364.2 ± 57.6 ms, p = 0.0216, Mann-Whitney test; Fig. 3P, vehicle: sag-amplitude = 15.06 ± 0.9487 mV, 6-OHDA: sag-amplitude = 19.11 ± 0.9118 mV, p = 0.0072, Mann-Whitney test). Other *in vitro* spike features were not significantly different between the two groups (Suppl. Fig. 7). Interestingly, breaking into whole-cell recording mode further accelerated firing

frequency, while only marginally increasing firing irregularity (Suppl. Fig. 7F). In summary, complete

functional recovery of surviving mSN DA neurons was – at least in part – achieved by homeostatic

plasticity of intrinsic excitability. Precisely, surviving mSN DA neurons not only recovered from instable

pacemaking at the early phase, but managed to double their discharge rates in late phase. As this

phenotype is reminiscent to the chronic adaptation of mSN DA neurons to mutant α-synuclein

overexpression, (Subramaniam et al., 2014a), we aimed to identify the underlying biophysical and

molecular changes.

Figure 3. Surviving SN DA neurons at late 6-OHDA phase recovered *in vivo* burst firing and doubled

their intrinsic pacemaker frequency *in vitro*.

(A), (I) Experimental design, illustrating timeline of behavioral assays, followed by terminal *in vivo* juxtacellular recordings (A) or by terminal *in vitro* whole-cell recordings (I) after >64 post-infusion days. (B), (C) Top: 10s original recording trace of spontaneous *in vivo* single-unit activity from SN DA neurons in vehicle (B) and 6-OHDA-infused mouse (C). Scalebars: 1s, 0.2mV. Below, left: corresponding ISI histograms. Below, right: corresponding confocal images of juxtacellularly labelled and immunohistochemically identified DA neuron. (D), (L) Anatomical mapping of all extracellularly recorded and juxtacellularly labelled DA neurons (D), and of all *in vitro* recorded and filled DA neurons (L), projected to bregma -3.40 mm (D) and -3.08mm (L). Location of example SN DA neurons in (B), (C), (J), (K) are highlighted. Smaller symbols represent DA neurons that have been recorded and identified, but not included in the group data analysis in (E-H). (E-H) Scatter dot-plots, showing no significant difference in mean firing rate (E), CV (F), SFB (G) and burst rate (H). (J), (K) Top: 10s original recording trace of *in vitro* whole-cell recording of spontaneous activity from SN DA neurons in a vehicle (J) and 6-OHDA-infused (K) mouse. Scalebars: 1s, 20mV. Note that the 6-OHDA DA neuron has enhanced, but regular pacemaking. Below, left, corresponding to hyperpolarizing current injection. Below, right, confocal images of NB filled and immunohistochemically identified DA neuron. (M-P) Scatter dot-plots, showing doubling of the firing rate (M), no difference in CV (N), decrease in rebound delay (O) and increase of sag-component (P) for the 6-OHDA-treated mice in comparison to vehicle group. Immunohistochemical imaging for all four DA neurons are displayed in 10x and 60x magnifications (green, TH; red, NB), scalebars: 200µm, 20µm. All data are presented as mean ± standard error of mean (SEM).

Accelerated *in vitro* pacemaking of surviving mSN DA neurons is caused by downregulation of Kv4.3 channels

To test whether Kv4.3-mediated A-type channels are responsible for differences in late phase

pacemaker rate between 6-OHDA and controls, we carried out a pharmacological occlusion

experiment using 1µM AmmTx3, a selective Kv4.3 channel blocker (see scheme Fig. 4A). The presence

of 1µM AmmTx3 not only accelerated discharge rates in both identified mSN DA neurons from vehicle-

and 6-OHDA-infused mice (see Fig. 4BC for representative cells; Fig. 4D for mapping) but, importantly,

also eliminated significant rate differences between the two treatment groups (Fig. 4E, vehicle: mean

firing rate = 4.54 ± 0.18Hz, 6-OHDA: mean firing rate = 4.84 ± 0.28 Hz, p = 0.285; Fig. 4 F, vehicle: CV =

7.84 ± 1.26 %, 6-OHDA: CV = 5.53 ± 0.44%, p = 0.2517; both Mann-Whitney test). This result

demonstrated that differences in Kv4.3 channel function were a main driver of the late post-6OHDA

fast pacemaker phenotype. In addition, 1µM AmmTx3 also abolished more subtle differences in

rebound delays and sag amplitudes (Fig. 4G, vehicle: rebound delay = 31.71 ± 2.4ms, 6-OHDA: rebound

delay = 28.4 ± 4.0ms, p = 0.0952; Fig. 4H. vehicle: sag-amplitude = 15.61 ± 1.1mV, 6-OHDA: sag-

amplitude = 16.43 ± 0.95mV, p = 0.7434; both Mann-Whitney test, Suppl. Fig. 8). In summary, these

experiments demonstrated that reduced Kv4.3 function and/or expression are mediators of the post-

6-OHDA late phase phenotype of accelerated *in vitro* pacing.

Next, we aimed to reveal whether this adaptive phenotype was associated with altered Kv4.3 protein

expression. We performed Kv4.3 immunohistochemistry and confocal imaging in vehicle and 6-OHDA

treated mice both on the ipsilateral and contralateral infusion side (see scheme in Fig. 4I). We defined

DA-selective ROIs by using TH-immunopositive areas to measure the intensity of Kv4.3-immunosignals

exclusively within DA neurons (i.e. TH-signal based mask) (Subramaniam *et al.*, 2014a). Figure 4J and

4K compare Kv4.3-immunoreactivities in the midbrain between/from contralateral control side and

ipsilateral affected side in a 6-OHDA-treated mouse in the late post-6-OHDA phase. Note the reduction

of TH-immunoreactivity on the ipsilateral side (upper panels). Higher resolution of Kv4.3-

immunoreactivity in surviving TH-positive neurons showed lower Kv4.3-immunosignal on the treated

side. Quantitative comparison of TH-ROIs of the entire SN between contralateral control side and 6-

OHDA-treated side revealed a significant reduction of average Kv4.3-immunosignal on the 6-OHDA

treated side (Figure 4L, D = 0.2014, p < 0.0001, two-sample Kolmogorov–Smirnov test). This reduction

was also detected in a selective analysis of Kv4.3-immunosignals in the medial SN (Figure 4M,

contralateral side: 73.03 ± 0.389, ipsilateral side: 54.8 ± 0.486, p < 0.0001, unpaired t-test), the region

where we carried out our functional characterization. In addition, Kv4.3 signal intensities showed

weak correlation to the size of TH-positive ROIs ($R2 = 0.0041$, Sy.x = 1227, N = 4874, p <0.0001, simple

linear regression, Suppl. Fig. 9). As Kv4.3 channel subunits are expected to be mostly located on the

cell membrane, we also investigated whether Kv4.3-immunosignal was different across cell

compartments. A decrease of Kv4.3-immunosignal was observed both in cell membrane and

cytoplasm ROIs (Suppl. Fig. 9). In contrast, analogous quantification in vehicle-treated mice revealed

no contra-ipsilateral differences in Kv4.3-immunoreactivity (Suppl. Fig. 9). In summary, these

immunohistochemical experiments provided evidence for a downregulation of Kv4.3 channel subunits

in mSN DA neurons, surviving a partial 6-OHDA lesion. This phenotype was therefore distinct from the

overexpression of Kv4.3 channel protein, observed in mutant α-synuclein mice (Subramaniam *et al.*,

2014a). Thus, we provided first evidence for a selective functional adaptation of SN DA neurons in

response to a single-hit non-progressive lesion.

Figure 4. Enhanced pacemaker frequency of SN DA neurons at the late post-6-OHDA phase was mediated by K$_v$4.3 channel downregulation

(A), (I) Experimental design, illustrating timeline of behavioral assays, followed by terminal *in vitro* whole-cell recordings under K$_v$4.3 channel blocker AmmTx3 (A) or by multi-labeling immunohistochemistry for exploring K$_v$4.3-channel expression differences (I) after >64 post-infusion days. (B), (C) Top: 10s original recording traces of *in vitro* whole-cell recording of spontaneous activity from SN DA neurons in AmmTx3-preincubated slices for a vehicle (B) and 6-OHDA-infused (C) mouse.

Below, left: corresponding hyperpolarizing current injection. Scalebars: 1s, 20mV. Below, right:

confocal images of NB filled and immunohistochemically identified DA neuron (green, TH; red, NB).

10x , 60x magnification, scalebars: 200µm, 20µm. (D) Anatomical mapping of all *in vitro* recorded and

filled DA neurons, projected to bregma -2.92 mm. (E-H) Scatter dot-plots, showing loss of differences

in *in vitro* electrophysiological parameters. (J), (K) Top: 4x magnification of midbrain of a 6-OHDA-

infused mouse, >64 days post-lesion – contralateral side (J), and corresponding ipsilateral side (K).

Middle: 60x magnification in the highlighted area from 4x image (green, TH; red, Kv4.3). Bottom, left:

zoom-in on an example ROI (highlighted in 60x image). Bottom, right: color-coded K_v4.3-channel

immunohistochemical signal intensity in the example ROI. Note Kv4.3-channel signal decrease in

surviving DA neurons on the ipsilateral to the injection side. (L) Histogram showing intensity of Kv4.3

immunosignals for all TH-positive ROIs, from ipsilateral, lesioned, side (in red) and from contralateral

side (in grey). Inset, same data shown as a cumulative distribution. Note a clear right-shift to lower

intensities for the ipsilateral side. (M) Comparison of mean TH-colocalized K_v4.3 immunosignals from

medial SN from ipsilateral, lesioned, side, and contralateral, as control side. All data are presented as

mean ± standard error of mean (SEM).

Discussion

Our study provides the first combined *in vitro* & *in vivo* electrophysiological characterization of identified DA neurons in the substantia nigra, surviving a partial 6-OHDA lesion. Studying these surviving SN DA neurons at two time points, we discovered time-dependent post-6-OHDA selective differences of firing properties both *in vivo* and *in vitro*. Early after lesion and coinciding with prominent behavioral impairments in our model, we detected a selective and dramatic (about 10-fold) reduction of *in vivo* burst firing in identified surviving SN DA neurons. This is reminiscent of an early study by Hollermann & Grace (Hollerman and Grace, 1990), who reported reduced bursting in putative SN DA neurons one week after partial lesion. At the early time point, we also found unstable *in vitro* pacemaker activity in these cells, while the mean firing rates were not different compared to those from vehicle-infused controls. In contrast, late (>2 months) after lesion, where substantial behavioral recovery had occurred, we detected no differences in *in vivo* firing rates and patterns – including burst discharges – of surviving SN DA neurons between 6-OHDA- and vehicle-infused mice. Surprisingly, the *in vitro* pacemaker rate was almost twice as fast in post-6-OHDA DA survivors compared to controls. Finally, we identified Kv4.3 channel downregulation as the cause for this chronic post-lesional pacemaker acceleration. In essence, we revealed for the first time the homeostatic adaptations of electrophysiological properties in surviving SN DA neurons.

Regarding the functional phenotype of surviving SN DA neurons, we identified a sequence of early but transient impairment, followed by chronic functional adaptation. While we did not investigate the mechanisms underlying the early impairment of *in vivo* burst firing, we focused our in-depth analysis on the chronic phase. Here, we identified a slowly-developing chronic Kv4.3 channel-mediated mechanism of pacemaker adaptation. Previously, we had already identified Kv4.3 channels in SN DA neurons as a pathophysiological target in a transgenic mutant (A53T-SNCA) α-synuclein mouse model (without cell loss) (Subramaniam *et al.*, 2014a). In this model, we found a pacemaker acceleration caused by oxidative impairment of Kv4.3 channels. In contrast to the post-lesional SN DA neurons in

this study, the activity of SN DA neurons in the α-synuclein model was accelerated also *in vivo.* Thus, α-synuclein pathology causes an allostatic shift of the *in vivo* firing rate setpoint of SN DA neurons, while post-lesional SN DA neurons recover their *in vivo* firing setpoint by homeostatic pacemaker plasticity. Interestingly, both models converge on the pacemaker potassium channel Kv4.3 but in a different fashion. While Kv4.3 subunits were downregulated on protein level in post-lesional SN DA neurons, mutant α-synuclein induced oxidative Kv4.3 dysfunction as well as protein upregulation (Subramaniam *et al.*, 2014a). We are aware that neither of the mouse models fully capture the pathophysiology of PD, where α-synuclein pathology, cell loss, inflammation and compensatory mechanisms are expected to occur side by side (Surmeier et al., 2017b). At least at the transcriptional level, we know that Kv4.3 mRNA is upregulated in human SN DA neurons from PD patients (Dragicevic *et al.*, 2015; Schiemann *et al.*, 2012). This is similar to α-synuclein models and opposite to post-lesional Kv4.3 downregulation, shown here, suggesting that Kv4.3 channel mediated compensatory mechanisms are finally overwhelmed during PD progression. Moreover, the results of this study imply that α-synuclein pathology interferes with the homeostatic capacity of surviving SN DA neurons, in particular the regulation of Kv4.3 channels. Indeed, we recently showed a Kv4.3 gain-of-function phenotype for vagal motoneurons in response to mutant SNCA expression (Chiu et al., 2021). Thus, Kv4.3 channels – a major "brake" of the pacemaker rate (Khaliq and Bean, 2008; Liss et al., 2001; Tarfa et al., 2017) – is an emerging downstream target across several PD models. It would be interesting to further investigate functional and molecular regulation of this potassium channel in chronic and progressive PD models, where pathology and cell loss go hand in hand.

In addition to cell-autonomous adaptions in surviving SN DA neurons, we assume that their chronic *in vivo* electrophysiological phenotype is also shaped by network level plasticity in response to dopamine-depletion. Our finding that the mean *in vivo* discharge rates were not different to controls in the presence of an accelerated intrinsic pacemaker strongly suggests a shift of the synaptic excitation-inhibition (E-I) balance towards more inhibition. Numerous studies have found altered synaptic inhibition in the dopamine-depleted basal ganglia (recently reviewed in (Zhang et al., 2021)). In particular, Heo and colleagues recently demonstrated a chronic E-I balance shift toward more

inhibition across several PD models including 6-OHDA (Heo et al., 2020). They identified a substantial

contribution of additional GABA synthesis and release from reactive astrocytes in the midbrain. It has

also been well-established that basal ganglia GABA neurons fire in more synchronized hence more

effective fashion after dopamine depletion (Cagnan et al., 2015; Evans et al., 2020; Milosevic et al.,

2018; Phillips et al., 2020; Tinkhauser et al., 2020; Wichmann et al., 2018). This has recently been

confirmed by elegant *in vivo* single cell resolved studies using either gCAMP-based calcium monitoring

or *in vivo* patch-clamp approaches (Ketzef et al., 2017; Kravitz *et al.*, 2010; Parker *et al.*, 2018; Parker

et al., 2016; Sitzia *et al.*, 2020). The chronic acceleration of pacemaker rate in response to enhanced

inhibition would also be expected within the framework of homeostatic plasticity (Turrigiano, 2012).

However, the mechanisms of firing rate set-point control, recently elucidated for hippocampal

neurons (Styr et al., 2019), is still unknown for SN DA neurons.

Our study has several limitations. As already discussed, we explored a single-hit partial lesion model,

while PD is a chronic progressive neurodegenerative process. While we demonstrate that

electrophysiological phenotypes per se are flexible and change in response to lesion, we also have to

consider distinct DA subpopulations within the SN. We recently showed that lateral SN DA neurons

projecting to DLS, the most vulnerable DA subpopulation (corresponding to the ventro-lateral SN DA

neurons in humans (Duzel et al., 2009; McCutcheon et al., 2019)) possess a distinct *in vivo* firing

phenotype in control mice (Farassat et al., 2019). Therefore, recording from this population in PD

models would be highly relevant. However, in our current partial 6-OHDA lesion model, these lateral

SN DA neurons were too severely lost to allow for more than anecdotal post-lesion

electrophysiological analysis (see Fig. 3). Here, a milder version of the current model would be needed

to identify their homeostatic plasticity. Regarding the surviving DA neuron in the medial SN, we

previously showed that they have distinct projection targets including DLS, DMS and lateral shell of

nucleus accumbens (Farassat *et al.*, 2019). In the current study, we have not aimed for identification

of axonal projections to avoid potentially confounding effects of additional brain surgery and to not

further complicate already challenging experiments. Nevertheless, this important issue should be

addressed in follow-up studies by e.g. molecular subtyping approaches (Heymann et al., 2020; Poulin

et al., 2018; Poulin et al., 2020; Saunders et al., 2018).

Finally, we would like to speculate about the possible implications of our findings for PD. In principle,

homeostatic responses, as identified here, could have a dual nature. On one hand, they might render

the neuron more robust, thereby slowing the rate of cell loss. On the other hand, the homeostatic

adaptations may amplify innate vulnerability and accelerate disease progression. The latter would be

a variant of the "stressful pacemaker" hypothesis of DA vulnerability (Chan et al., 2007; Surmeier,

2007; Surmeier et al., 2017a) and its potential clinical implications (Guzman et al., 2018; Liss and

Striessnig, 2019; Ortner, 2021; Ortner et al., 2017; Parkinson Study Group, 2020). In summary, we

provide the first steps towards defining the homeostatic repertoire of surviving DA neurons, but

deeper insights are necessary to rationally design neuroprotective interventions in PD.

Methods

Animals

Male C57Bl6/N mice (Charles River Laboratories) were used for the study. Mice were 8 weeks old, group housed and maintained on a 12-hour light-dark cycle. All experiments and procedures involving mice were approved by the German Regierungspräsidium Darmstadt (V54-19c20/15-F40/30). In total, 115 mice were used for this study (see table below).

	Early phase		Late phase	
	Mice (#)	Cells (#)	Mice (#)	Cells (#)
In vivo juxta & behavior – vehicle	15	12 – medial/ 28 – all	12	10 – medial/ 26 – all
In vivo juxta & behavior – 6-OHDA	16	9 – medial/ 16 – all	10	9 – medial/ 11 – all
In vitro – vehicle	3	16	2	9
In vitro – 6-OHDA	6	37	5	25
In vitro & AmmTx3 – vehicle	-	-	3	15
In vitro & AmmTx3 – 6-OHDA	-	-	3	14
	Mice (#)		Mice (#)	
Kv4.3 Immunochemistry	8		8	
SN quantification – vehicle	9		3	
SN quantification – 6-OHDA	9		3	
Sum = 115				

Stereotactic 6-OHDA infusion

All surgeries were performed under general anesthesia in areflexic state. Prior to the induction of anesthesia, a premedication of 0.2 mg/kg atropine (atropine-sulfate, Braun Melsungen AG, Melsungen) was given as an intraperitoneal (i.p.) injection to stabilize circulation. Anesthesia was induced in a plastic chamber, which was flooded with 5% Isoflurane (Florene®, AbbVie Deutschland GmbH & Co. KG, Ludwigshafen, Germany) in pure oxygen (0.4 l/min). For maintenance of anesthesia, isoflurane was delivered through a breathing mask with a flow rate of 0.35 l/min and its concentration was regulated to 1.5-2.2% using an adjustable vaporizer (Uno, Zevenaar, Netherlands). The depth of anesthesia was controlled by testing the toe pinch reflex and the breathing rate (1-2Hz). Body temperature (36°C) and respiration were constantly monitored. Lidocaine/prilocaine ointment (25

mg/g, Emla® creme, AstraZeneca GmbH, 22876 Wedel) was applied prior to surgery and after suturing

of the wound as local anesthetics. Additional analgesia was provided by subcutaneous injection of

carprofen (4 mg/kg in NaCl, 1:100, Rimadyl®, Pfilzer GmbH, Berlin, Germany) after infusion. Eye

lubricant (Visidic, Bausch and Lomb, Berlin, Germany) was used to protect eyes from desiccation.

Desipramine hydrochloride (20 mg/kg, Sigma Aldrich) was injected i.p. 20-40 min before intracranial

infusions to prevent 6-OHDA uptake by noradrenergic neurons. The desipramine solution was

prepared in sterile, isotonic NaCl solution (B. Braun Melsungen AG, Germany) at the day of surgery.

The infusion solutions are based on sterile artificial cerebrospinal fluid (ACSF, Harvard Apparatus,

Holliston, MA, USA) with 0.2% L-ascorbic acid (used also as a vehicle solution). The 6-OHDA solution

(0.02% 6-hydroxydopamine hydrochloride in ACSF with 0.2% L-ascorbic acid) was prepared at the day

of infusion, stored on ice, and shielded from light.

During surgery, the animals were placed on a heating pad and were fixed in a stereotactic frame

(Model 1900, Kopf Instruments, Tujunga, USA) with a stereotactic arm and a connected three-way

digital positioning display. The scalp was opened by a longitudinal cut to expose the skull with bregma

and lambda on display. With a centering scope (Model 1915, Kopf Instruments, Tujunga, USA), the

bregma-lambda distance was measured and examined for suitable anatomy (4.4±0.2 mm distance).

Afterwards, the skull was aligned to a reference frame with a stereotaxic alignment indicator (Model

1905, Kopf Instruments, Tujunga, USA) and the manipulator system was referenced to bregma.

Using a stereotaxic drill (Model 1911, Kopf Instruments, Tujunga, USA) with a 500 µm diameter drill

bit, a hole above the right striatum was drilled (coordinates: ML: +1.9 mm, AP: +0.5 mm to bregma).

ACSF or 6-OHDA solution were loaded to a 10 µl NanoFil syringe (World Precision Instruments Inc.,

Sarasota, FL, USA) with a 35G blunt needle, which was mounted on a MicroSyringe Pump (UMP3-1,

World Precision Instruments) and controlled by a SYSMicro4 Controller (World Precision Instruments).

Using the stereotactic arm, the needle was slowly lowered (about 750 µm/min) to a position of -2.2

mm below the brain surface (infusion site coordinates: ML: +1.9 mm, AP: +0.5 mm, DV: -2.2 mm to

bregma). Anatomical references are based on Franklin and Paxinos (2008). A volume of 6 µl was

infused with a flow rate of 250 nl/min. Once the volume was infused, the needle rested for 5 minutes

in that position before it was slowly moved out of the brain. Directly before and after infusion, proper

functioning of the syringe system and the needle was checked. Finally, after suture the animal was

placed on a heating pad for full recovery. Oats, wet food pallets and water were placed inside the cage

to ease consumption.

Behavioral testing

Open field

Spontaneous locomotion (track length, wall distance, time in center and number of rearings) and

rotations of all mice were monitored in open field (50 × 50 cm, center 30 × 30 cm; red illumination, 3

lx) for 10 min in 3 baseline sessions and every 4th or 7th day post-infusion of ACSF/6-OHDA till the day

of *in vivo* or *in vitro* experiment (e.g. 21st or >68th post-operative day). The open field was cleaned

before and after each mouse with 0,1% acetic acid in distilled water. Using a video tracking system

(Viewer II/III, Biobserve) spontaneous behavior was recorded and analyzed both online and offline.

Data was extracted from Viewer as Excel-tables and the final analysis was made with custom made

Matlab-scripts.

Cylinder test

Forelimb use during explorative activity was explored with cylinder test. The test was performed at

corresponding termination time point (20-21st and 64th post-infusion day). Mice were placed

individually in a glass beaker (9 cm diameter, 19 cm height) at room light and were video recorded

with a camera (Logitech HD Webcam C615) for about 5 min. No habituation was allowed before video

recording. The glass cylinder was cleaned before and after every mouse with 0,1% acetic acid in

distilled water. Only weight-bearing wall contacts made by each and both forelimb on the cylinder

wall were scored. Wall exploration was expressed in terms of the percentage of contralateral to the

infusion side (in the 6-OHDA-infused mice also impaired forepaw) to all forelimb wall contacts.

In vivo electrophysiology

Extracellular recording

In vivo extracellular single-unit activities of SN and VTA neurons were recorded in ACSF-infused

(vehicle) and 6-OHDA-infused mice, similar procedures were used in other studies from our lab

(Farassat *et al.*, 2019; Schiemann *et al.*, 2012; Subramaniam *et al.*, 2014a). Briefly, mice were

anesthetized (isoflurane; induction 4.5-5%, maintenance 1–2% in 0.4 l/min O2) and placed into a

stereotactic frame. The craniotomies were performed as described above to target the lateral SN (AP:

-3.08 mm, ML: 1.4 mm) and medial SN (AP: -3.08 mm, ML: 0.9 mm). Borosilicate glass electrodes (10–

25 MΩ; Harvard Apparatus, Holliston, MA, USA) were made using a horizontal puller (DMZ-Universal

Puller, Zeitz, Germany) and filled with 0.5 M NaCl, 10 mM HEPES (pH 7.4) and 1.5% neurobiotin (Vector

Laboratories, Burlingame, CA, USA). A micromanipulator (SM-6; Luigs & Neumann, Ratingen,

Germany) was used to lower the electrodes to the recording site. The single-unit activity of each

neuron was recorded for at least 10 minutes at a sampling rate of 12.5 kHz (for firing pattern analyses),

and then for another one minute at a sampling rate of 20 kHz (for the fine analysis of AP waveforms).

Signals were amplified 1000x (ELC-03M; NPI electronics, Tamm, Germany), notch- and bandpass-

filtered 0.3–5000 Hz (single-pole, 6 dB/octave, DPA-2FS, NPI electronics) and recorded on a computer

with an EPC-10 A/D converter (Heka, Lambrecht, Germany). Simultaneously, the signals were

displayed on an analog oscilloscope and an audio monitor (HAMEG Instruments CombiScope HM1508;

AUDIS-03/12M NPI electronics). Midbrain DA neurons were initially identified by their broad biphasic

AP (> 1.2 ms duration) and slow frequency (1–8 Hz) (Grace and Bunney, 1984a; b; Ungless and Grace,

2012). AP duration was determined as the interval between the start of initial upward component and

the minimum of following downward component.

Juxtacellular labeling of single neurons

In order to identify the anatomical location and neurochemical identity of the recorded neurons, they were labeled post-recording with neurobiotin using the juxtacellular *in vivo* labeling technique (Pinault, 1996). Microiontophoretic currents were applied (1–10 nA positive current, 200ms on/off pulse, ELC-03M, NPI Electronics) via the recording electrode in parallel to the monitoring of single-unit activity. Labeling was considered successful, when: the firing pattern of the neuron was modulated during current injection (i.e., increased activity during on-pulse and absence of activity in the off-pulse), the process was stable for at least 20s, and was followed by the recovery of spontaneous activity. This procedure allowed for the exact mapping of the recorded DA neuron within the SN and VTA subnuclei (Franklin and Paxinos, 2008) using custom written scripts in Matlab (MathWorks, Natick, MA, USA), combined with neurochemical identification using TH-immunostaining.

In vitro electrophysiology

Slice preparation

Animals were anesthetized by intraperitoneal injection of ketamine (250 mg/kg, Ketaset, Zoetis) and medetomidine-hydrochloride (2.5 mg/kg, Domitor, OrionPharma) prior to intracardial perfusion using ice-cold ACSF consisting of the following (in mM): 50 sucrose, 125 NaCl, 2.5 KCl, 25 NaHCO3, 1.25 NaH2PO4, 2.5 glucose, 6 MgCl2, 0.1 CaCl2 and 2.96 kynurenic acid (Sigma-Aldrich), oxygenated with 95% O2 and 5% CO2. Rostral coronal midbrain slices (bregma: -2.92 mm to -3.16 mm) were sectioned at 250 μm using a vibrating blade microtome (VT1200s, Leica). Slices were incubated for 1 h before recordings in a 37°C bath with oxygenated extracellular solution with extra 1μM AmmTx3, containing the following (in mM): 22.5 sucrose, 125 NaCl, 3.5 KCl, 25 NaHCO3, 1.25 NaH2PO4, 2.5 glucose, 1.2 MgCl2 and 1.2 CaCl2.

 In vitro patch-clamp recordings

Slices were placed in a heated recording chamber (37°C) that was perfused with oxygenated

extracellular solution at 2-4 ml min^{-1}. CNQX (20 μM; Biotrend), gabazine (SR95531, 4 μM; Biotrend)

and DL-AP5 (10 μM; Cayman Chemical) were added to inhibit excitatory and inhibitory synaptic

transmission. Neurons were visualized using infrared differential interference contrast

videomicroscopy with a digital camera (VX55, Till Photonics) connected to an upright microscope

(Axioskop 2, FSplus, Zeiss). Patch pipettes were pulled from borosilicate glass (GC150TF-10; Harvard

Apparatus LTD) using a temperature-controlled, horizontal pipette puller (DMZ-Universal Puller,

Zeitz). Patch pipettes (4-6 MΩ) were filled with a solution containing the following (in mM): 135 KGlu,

5 KCl, 10 HEPES, 0.1 EGTA, 5 MgCl2, 0.075 CaCl2, 5 NaATP, 1 LiGTP, 0.1% neurobiotin, adjusted to a

pH of 7.35 with KOH. Recordings were performed using an EPC-10 patch-clamp amplifier (Heka

electronics) with a sampling rate of 20 kHz and a low-pass filter (Bessel, 5 kHz). For analysis, action

potential thresholds (mV) were determined at $dV_m/dt > 10$ mV/ms.

Immunohistochemistry

Following *in vivo* recordings, animals were transcardially perfused, as described previously (Farassat

et al., 2019; Schiemann *et al.*, 2012; Subramaniam *et al.*, 2014a). Fixed brains were sectioned into

60μm (midbrain) or 100μm (forebrain) coronal sections using a vibrating microtome (VT1000S, Leica).

In vitro slices were fixed in paraformaldehyde after finishing the experiment. Sections were rinsed in

PBS and then incubated (in blocking solution (0.2 M PBS with 10% horse serum, 0.5% Triton X-100,

0.2% BSA). Afterwards, sections were incubated in carrier solution (room temperature, overnight)

with the following primary antibodies: polyclonal guinea pig anti-tyrosine hydroxylase (TH; 1:1000;

Synaptic Systems), monoclonal mouse anti-TH (1:1000; Synaptic Systems) or polyclonal rabbit anti-TH

(1:1000; Synaptic Systems); mouse anti-Kv4.3 (1:1000, Alomone Labs). In sequence, sections were

again washed in PBS and incubated (room temperature, overnight) with the following secondary

antibodies: goat anti-guinea pig 488, goat anti-rabbit 488, goat anti-mouse 488, goat anti-mouse 568

(all 1:750; Thermofisher). Streptavidin AlexaFluor-568 and Streptavidin AlexaFluor-488 (both 1:1000;

Invitrogen) were used for identifying neurobiotin-filled cells. Sections were then rinsed in PBS and

mounted on slides with fluorescent mounting medium (Vectashield, Vector Laboratories).

DAB immunocytochemistry

For DAB (3,3'-diaminobenzidine) staining procedures, a Vectastain ABC Staining Kit (Vector

Laboratories) was used. Coronal sections of midbrain (30 µm) areas were cut and rinsed in PBS (3x10

min). Similar to previous immunolabeling procedures, unspecific antigen binding sites were blocked

by incubation of the sections with blocking solution (60 min, room temperature). Subsequently,

sections were incubated with primary antibody against TH (rabbit anti TH) overnight, rinsed in PBS

(3x10 min), and were incubated with biotinylated secondary antibodies (biotinylated anti-rabbit) for

two hours at RT. In parallel, an avidin-biotin complex (ABC) was formed by pre-incubation of avidin

(1:1000) with biotinylated HRP (1:1000) in PBS for two hours at room temperature. Sections were

rinsed in PBS (3x10 min) prior to incubation with ABC solution (60 min, room temperature). Next,

sections were rinsed in PBS (2x10 min) and Tris-buffer (1x10 min). Finally, DAB oxidation was

developed by application of 2 % H_2O_2, 2 % $NiCl_2$ and 4 % DAB in Tris-buffer using a DAB Substrate Kit

(Vector Laboratories, Burlingame, USA). $NiCl_2$ enhances sensitivity and intensity of DAB precipitation

product. DAB oxidation was developed for 2 to 5 minutes and was stopped with Tris-buffer once a

specific high-contrast signal was detectable. Sections were rinsed in Tris-buffer (3x10 min) and

transferred onto gelatin-covered slides, air-dried overnight, and dehydrated in consecutive ascending

alcohol concentrations (50 %, 70 %, 90 % and 2x 100 %; 10 min each) followed by dehydration in xylol

(2x 100 %; 10 min each). Finally, sections were mounted under glass coverslips with hardening

mounting medium (Vectamount, Vector Laboratories, Burlingame, USA).

Unbiased stereology measurements

For quantification of total cell loss, TH-DAB labeled SN DA neurons were counted using unbiased stereology based on optical dissection (Gundersen, 1986). In coronal sections (30 μm), the region of interest was selected based on anatomical landmarks including the medial lemniscus, which separates SN and adjacent VTA. Stereological counting provides unbiased data based on random, systematic sampling using an optical fractionator. This method involves counting of neurons with an optical dissector, a three-dimensional probe placed through a reference space (Gundersen, 1986). The optical dissector forms a focal plane with a thin depth of field through the selected sections. Objects in focus of this focal plane are located within the reference section and are counted, while objects outside of the focal plane are not counted. On top of the optical dissector, a counting frame is applied. Counting frames ensure that all neurons have equal probabilities of being selected, regardless of shape, size, orientation, and distribution. To avoid counting of capped neurons at the border of a section, an additional guard zone was deployed at the upper and lower borders of each section. DA neurons within the counting frame as well as those crossing the green line (acceptance line) were counted, while DA neurons crossing the red line (rejection line) excluded. Moreover, only neurons with a detectable nucleus in focus within the optical dissector were counted. For quantification of total cell loss, StereoInvestigator software (V5, MicroBrightField, Colchester, USA) was used in combination with BX61 microscope (Olympus, Hamburg, Germany). The region of interest was selected and marked using a low magnification objective lens (2x, NA 0.25, Olympus) and 12-30 serial sections of 30 μm thickness were counted, covering the entire caudo-rostral extent of the SN. To count the number of DA neurons in the area of the SN pars compacta, a high magnification oil-immersion objective lens (100x, NA 1.30, Olympus) was used.

Optical density measurements

Optical density measurements of TH-DAB labeled striatal sections were performed using ImageJ software (http://rsbweb.nih.gov/ij/). Following TH-DAB labeling and TH-immunohistochemistry,

images of five coronal sections (100 μm) covering the rostrocaudal axis of the striatum were captured

using laser-scanning microscope (Nikon Eclipse90i, Nikon GmbH). Images were gray scale converted

and mean gray values of desired striatal areas were encircled and measured. Unspecific mean gray

values were measured in a defined cortical area (100x100 pixels) that displayed no specific TH signal

due to the absence of DA innervation and were subtracted. The ventral edge of lateral ventricles

served as an anatomical landmark to separate dorsal and ventral areas. For all animals, the

measurement from the ipsilateral to the infusion side were divided by the contralateral side to

calculate the relative optical density of the striatum.

Immunohistochemical Kv4.3 channel signal quantification

A Nikon Eclipse 90i microscope was used for fluorescent signal detection, excitation wavelength of

488 nm for TH-signal and 568 nm for Kv4.3-channel signal. From each animal, 4 midbrain slices

covering the caudal, intermedial and rostral regions, were selected and imaged for overview with 4x

magnification. Then 60x magnification was used to acquire data from 4 areas within each SN (4 images

on the ipsilateral and 4 images on the contralateral to infusion side). All images were acquired using

the same laser and camera settings. Images were exported from Nikon NIS-Elements Advanced

Research (Version 4.20.03) software as 8-bit TIFF files for quantification. Data was analyzed using

custom made Python 3.0 scripts with matplotlib, numpy, scimage and scipy modules. First, TH

immunosignals were converted to a binary image via Otsu-thresholding algorithm. Then, the resulting

binary image was used as a mask for Kv4.3 channel immunosignal detection. For all ROIs surface areas

and mean Kv.4.3 channel signal intensity was measured. By applying erosion and dilatation algorithms

on the ROIs, membrane and cytoplasm areas were segregated, allowing isolation of Kv4.3 channel

immunosignal intensity for these cell compartments. Background Kv4.3 channel immunosignal was

quantified in TH-immunosignal areas below the Otsu-threshold. All data were then grouped according

to medio-lateral and ipsi-/contralateral position for both vehicle and 6-OHDA group. Graphs and

statistical analysis for this data was performed using Python custom made scripts.

Statistical Analysis

Spike train analyses

Spike time-stamps were extracted by thresholding above noise levels with IgorPro 6.02 (WaveMetrics,

Lake Oswego, OR, USA). Firing pattern properties such as mean frequency, coefficient of variation (CV)

and bursting measures were analyzed using custom scripts in Matlab. In order to estimate burstiness

and intra-burst properties, we used the burst detection methods described in Grace & Bunney (Grace

and Bunney, 1984a; b). All non-burst related ISIs (excluding all ISIs that followed the Grace and Bunney

criteria, as well as all pre- and post-burst ISIs) were used to calculate the single spike firing frequency

and single spike coefficient of variation.

For analysis of general firing patterns, autocorrelation histograms (ACH) were plotted using custom

Matlab scripts. We used established criteria for classification of *in vivo* firing patterns based on visual

inspection of autocorrelograms (Farassat *et al.*, 2019; Schiemann *et al.*, 2012; Subramaniam *et al.*,

2014a): single spike-oscillatory (≥3 equidistant peaks with decreasing amplitudes), single spike-

irregular (<3 peaks, increasing from zero approximating a steady state), bursty-irregular (narrow peak

with steep increase at short ISIs) and bursty-oscillatory (narrow peak reflecting fast intraburst ISIs

followed by clear trough and repetitive broader peaks).

Statistics

Categorical data is represented as stacked bar graphs. To investigate the assumption of normal

distribution, we performed the single-sample Kolmogorov-Smirnov test. The Mann-Whitney-Test,

one-/two-way ANOVA were performed in non-parametric data to determine statistical significance.

Categorical parameters, such as ACH-based firing pattern, were analyzed with the Chi-squared test.

Statistical significance level was set to $p < 0.05$. All data values are presented as means ± SEM.

Statistical tests were performed using GraphPad Prism 9 (GraphPad Software, San Diego, CA, USA),

Matlab and Python. The scatter plots are represented with median or mean ± SEM. The resulting *p*

values were compared with Bonferroni-corrected α-level or Tukey *post hoc* comparison. A value of p

≤ 0.05 was considered to be statistically significant; $p \leq 0.05 =$ *$p \leq 0.005 =$ **$p \leq 0.0005 =$ ***. Graphs

were plotted using GraphPad Prism software (9.0c), Matlab and Python.

Acknowledgements

724 This study was supported by research grants to JR (DFG CRC 1080 and CRC 1451). LK is a MD/PhD

725 candidate at TransMed, Gutenberg University Mainz. We thank Beatrice Fischer and Jasmine Sonntag

726 for technical assistance, Alexander Prinz for preliminary data on post-6-OHDA stereology and Kauê

727 Machado Costa for support in Matlab based analysis and mentoring to LK.

Additional information

Author contributions

730 LK & JR designed the study. LK performed the lesions, behavioral & *in vivo* electrophysiology, JS

731 performed the *in vitro* experiments. Analysis was carried out jointly by LK, JS, NF and JR. NF taught LK

732 *in vivo* electrophysiology and juxtacellular labelling. LK & JR wrote the manuscript.

Declaration of Interests

734 The authors declare no conflict of interests.

Author ORCIDs

736 Jochen Roeper https://orcid.org/0000-0003-2145-8742

737 Lora Kovacheva https://orcid.org/0000-0001-6999-1533

738 Josef Shin https://orcid.org/0000-0002-3556-125X

Ethics

740 Animal experimentation: All experiments and procedures involving mice were approved by the

741 German Regierungspräsidium Darmstadt (V54-19c20/15-F40/30).

References

Bez, F., Francardo, V., and Cenci, M.A. (2016). Dramatic differences in susceptibility to l-DOPA-induced dyskinesia between mice that are aged before or after a nigrostriatal dopamine lesion. Neurobiol Dis *94*, 213-225. 10.1016/j.nbd.2016.06.005.

Cagnan, H., Duff, E.P., and Brown, P. (2015). The relative phases of basal ganglia activities dynamically shape effective connectivity in Parkinson's disease. Brain *138*, 1667-1678. 10.1093/brain/awv093.

Cenci, M.A., and Bjorklund, A. (2020). Animal models for preclinical Parkinson's research: An update and critical appraisal. Prog Brain Res *252*, 27-59. 10.1016/bs.pbr.2020.02.003.

Chan, C.S., Guzman, J.N., Ilijic, E., Mercer, J.N., Rick, C., Tkatch, T., Meredith, G.E., and Surmeier, D.J. (2007). 'Rejuvenation' protects neurons in mouse models of Parkinson's disease. Nature *447*, 1081-1086. 10.1038/nature05865.

Chen, L., Daniels, S., Kim, Y., and Chu, H.Y. (2021). Cell Type-Specific Decrease of the Intrinsic Excitability of Motor Cortical Pyramidal Neurons in Parkinsonism. J Neurosci *41*, 5553-5565. 10.1523/JNEUROSCI.2694-20.2021.

Chiu, W.H., Kovacheva, L., Musgrove, R.E., Arien-Zakay, H., Koprich, J.B., Brotchie, J.M., Yaka, R., Ben-Zvi, D., Hanani, M., Roeper, J., and Goldberg, J.A. (2021). α-Synuclein-induced Kv4 channelopathy in mouse vagal motoneurons drives nonmotor parkinsonian symptoms. Sci Adv *7*. 10.1126/sciadv.abd3994.

Dragicevic, E., Schiemann, J., and Liss, B. (2015). Dopamine midbrain neurons in health and Parkinson's disease: emerging roles of voltage-gated calcium channels and ATP-sensitive potassium channels. Neuroscience *284*, 798-814. 10.1016/j.neuroscience.2014.10.037.

Duzel, E., Bunzeck, N., Guitart-Masip, M., Wittmann, B., Schott, B.H., and Tobler, P.N. (2009). Functional imaging of the human dopaminergic midbrain. Trends Neurosci *32*, 321-328. 10.1016/j.tins.2009.02.005.

Evans, R.C., Twedell, E.L., Zhu, M., Ascencio, J., Zhang, R., and Khaliq, Z.M. (2020). Functional Dissection of Basal Ganglia Inhibitory Inputs onto Substantia Nigra Dopaminergic Neurons. Cell Rep *32*, 108156. 10.1016/j.celrep.2020.108156.

Farassat, N., Costa, K.M., Stojanovic, S., Albert, S., Kovacheva, L., Shin, J., Egger, R., Somayaji, M., Duvarci, S., Schneider, G., and Roeper, J. (2019). In vivo functional diversity of midbrain dopamine neurons within identified axonal projections. Elife *8*. 10.7554/eLife.48408.

Fieblinger, T., Zanetti, L., Sebastianutto, I., Breger, L.S., Quintino, L., Lockowandt, M., Lundberg, C., and Cenci, M.A. (2018). Striatonigral neurons divide into two distinct morphological-physiological phenotypes after chronic L-DOPA treatment in parkinsonian rats. Sci Rep *8*, 10068. 10.1038/s41598-018-28273-5.

Gibb, W.R., and Lees, A.J. (1991). Anatomy, pigmentation, ventral and dorsal subpopulations of the substantia nigra, and differential cell death in Parkinson's disease. J Neurol Neurosurg Psychiatry *54*, 388-396. 10.1136/jnnp.54.5.388.

Giguere, N., Burke Nanni, S., and Trudeau, L.E. (2018). On Cell Loss and Selective Vulnerability of Neuronal Populations in Parkinson's Disease. Front Neurol *9*, 455. 10.3389/fneur.2018.00455.

Grace, A.A., and Bunney, B.S. (1984a). The control of firing pattern in nigral dopamine neurons: burst firing. J Neurosci *4*, 2877-2890.

Grace, A.A., and Bunney, B.S. (1984b). The control of firing pattern in nigral dopamine neurons: single spike firing. J Neurosci *4*, 2866-2876.

Gundersen, H.J. (1986). Stereology of arbitrary particles. A review of unbiased number and size estimators and the presentation of some new ones, in memory of William R. Thompson. J Microsc *143*, 3-45.

Guzman, J.N., Ilijic, E., Yang, B., Sanchez-Padilla, J., Wokosin, D., Galtieri, D., Kondapalli, J., Schumacker, P.T., and Surmeier, D.J. (2018). Systemic isradipine treatment diminishes calcium-dependent mitochondrial oxidant stress. J Clin Invest *128*, 2266-2280. 10.1172/JCI95898.

Heo, J.Y., Nam, M.H., Yoon, H.H., Kim, J., Hwang, Y.J., Won, W., Woo, D.H., Lee, J.A., Park, H.J., Jo, S., et al. (2020). Aberrant Tonic Inhibition of Dopaminergic Neuronal Activity Causes Motor Symptoms in Animal Models of Parkinson's Disease. Curr Biol *30*, 276-291.e279. 10.1016/j.cub.2019.11.079.

Heymann, G., Jo, Y.S., Reichard, K.L., McFarland, N., Chavkin, C., Palmiter, R.D., Soden, M.E., and Zweifel, L.S. (2020). Synergy of Distinct Dopamine Projection Populations in Behavioral Reinforcement. Neuron *105*, 909-920 e905. 10.1016/j.neuron.2019.11.024.

Hollerman, J.R., and Grace, A.A. (1990). The effects of dopamine-depleting brain lesions on the electrophysiological activity of rat substantia nigra dopamine neurons. Brain Res *533*, 203-212. 10.1016/0006-8993(90)91341-d.

Kalia, L.V., and Lang, A.E. (2015). Parkinson's disease. Lancet *386*, 896-912. 10.1016/S0140-6736(14)61393-3.

Ketzef, M., Spigolon, G., Johansson, Y., Bonito-Oliva, A., Fisone, G., and Silberberg, G. (2017). Dopamine Depletion Impairs Bilateral Sensory Processing in the Striatum in a Pathway-Dependent Manner. Neuron *94*, 855-865 e855. 10.1016/j.neuron.2017.05.004.

Khaliq, Z.M., and Bean, B.P. (2008). Dynamic, nonlinear feedback regulation of slow pacemaking by A-type potassium current in ventral tegmental area neurons. J Neurosci *28*, 10905-10917. 10.1523/JNEUROSCI.2237-08.2008.

Kirik, D., Rosenblad, C., and Bjorklund, A. (1998). Characterization of behavioral and neurodegenerative changes following partial lesions of the nigrostriatal dopamine system induced by intrastriatal 6-hydroxydopamine in the rat. Exp Neurol *152*, 259-277. 10.1006/exnr.1998.6848.

Kordower, J.H., Olanow, C.W., Dodiya, H.B., Chu, Y., Beach, T.G., Adler, C.H., Halliday, G.M., and Bartus, R.T. (2013). Disease duration and the integrity of the nigrostriatal system in Parkinson's disease. Brain *136*, 2419-2431. 10.1093/brain/awt192.

Kravitz, A.V., Freeze, B.S., Parker, P.R., Kay, K., Thwin, M.T., Deisseroth, K., and Kreitzer, A.C. (2010). Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. Nature *466*, 622-626. 10.1038/nature09159.

Liss, B., Franz, O., Sewing, S., Bruns, R., Neuhoff, H., and Roeper, J. (2001). Tuning pacemaker frequency of individual dopaminergic neurons by Kv4.3L and KChip3.1 transcription. EMBO J *20*, 5715-5724. 10.1093/emboj/20.20.5715.

Liss, B., and Striessnig, J. (2019). The Potential of L-Type Calcium Channels as a Drug Target for Neuroprotective Therapy in Parkinson's Disease. Annu Rev Pharmacol Toxicol *59*, 263-289. 10.1146/annurev-pharmtox-010818-021214.

McCutcheon, R.A., Abi-Dargham, A., and Howes, O.D. (2019). Schizophrenia, Dopamine and the Striatum: From Biology to Symptoms. Trends Neurosci *42*, 205-220. 10.1016/j.tins.2018.12.004.

Milosevic, L., Kalia, S.K., Hodaie, M., Lozano, A.M., Fasano, A., Popovic, M.R., and Hutchison, W.D. (2018). Neuronal inhibition and synaptic plasticity of basal ganglia neurons in Parkinson's disease. Brain *141*, 177-190. 10.1093/brain/awx296.

Ortner, N.J. (2021). Voltage-Gated Ca(2+) Channels in Dopaminergic Substantia Nigra Neurons: Therapeutic Targets for Neuroprotection in Parkinson's Disease? Front Synaptic Neurosci *13*, 636103. 10.3389/fnsyn.2021.636103.

Ortner, N.J., Bock, G., Dougalis, A., Kharitonova, M., Duda, J., Hess, S., Tuluc, P., Pomberger, T., Stefanova, N., Pitterl, F., et al. (2017). Lower Affinity of Isradipine for L-Type Ca(2+) Channels during Substantia Nigra Dopamine Neuron-Like Activity: Implications for Neuroprotection in Parkinson's Disease. J Neurosci *37*, 6761-6777. 10.1523/JNEUROSCI.2946-16.2017.

Otomo, K., Perkins, J., Kulkarni, A., Stojanovic, S., Roeper, J., and Paladini, C.A. (2020). In vivo patch-clamp recordings reveal distinct subthreshold signatures and threshold dynamics of midbrain dopamine neurons. Nat Commun *11*, 6286. 10.1038/s41467-020-20041-2.

Parker, J.G., Marshall, J.D., Ahanonu, B., Wu, Y.W., Kim, T.H., Grewe, B.F., Zhang, Y., Li, J.Z., Ding, J.B., Ehlers, M.D., and Schnitzer, M.J. (2018). Diametric neural ensemble dynamics in parkinsonian and dyskinetic states. Nature *557*, 177-182. 10.1038/s41586-018-0090-6.

Parker, P.R., Lalive, A.L., and Kreitzer, A.C. (2016). Pathway-Specific Remodeling of Thalamostriatal Synapses in Parkinsonian Mice. Neuron *89*, 734-740. 10.1016/j.neuron.2015.12.038.

Parkinson Study Group, S.-P.D.I.I.I.I. (2020). Isradipine Versus Placebo in Early Parkinson Disease: A Randomized Trial. Ann Intern Med *172*, 591-598. 10.7326/M19-2534.

Phillips, R.S., Rosner, I., Gittis, A.H., and Rubin, J.E. (2020). The effects of chloride dynamics on substantia nigra pars reticulata responses to pallidal and striatal inputs. Elife *9*. 10.7554/eLife.55592.

Pinault, D. (1996). A novel single-cell staining procedure performed in vivo under electrophysiological
control: morpho-functional features of juxtacellularly labeled thalamic cells and other central neurons
with biocytin or Neurobiotin. J Neurosci Methods *65*, 113-136. 10.1016/0165-0270(95)00144-1.
Poewe, W., Seppi, K., Tanner, C.M., Halliday, G.M., Brundin, P., Volkmann, J., Schrag, A.E., and Lang,
A.E. (2017). Parkinson disease. Nat Rev Dis Primers *3*, 17013. 10.1038/nrdp.2017.13.
Poulin, J.F., Caronia, G., Hofer, C., Cui, Q., Helm, B., Ramakrishnan, C., Chan, C.S., Dombeck, D.A.,
Deisseroth, K., and Awatramani, R. (2018). Mapping projections of molecularly defined dopamine
neuron subtypes using intersectional genetic approaches. Nat Neurosci *21*, 1260-1271.
10.1038/s41593-018-0203-4.
Poulin, J.F., Gaertner, Z., Moreno-Ramos, O.A., and Awatramani, R. (2020). Classification of Midbrain
Dopamine Neurons Using Single-Cell Gene Expression Profiling Approaches. Trends Neurosci *43*, 155-
169. 10.1016/j.tins.2020.01.004.
Rubi, L., and Fritschy, J.M. (2020). Increased GABAergic transmission in neuropeptide Y-expressing
neurons in the dopamine-depleted murine striatum. J Neurophysiol *123*, 1496-1503.
10.1152/jn.00059.2020.
Saunders, A., Macosko, E.Z., Wysoker, A., Goldman, M., Krienen, F.M., de Rivera, H., Bien, E., Baum,
M., Bortolin, L., Wang, S., et al. (2018). Molecular Diversity and Specializations among the Cells of the
Adult Mouse Brain. Cell *174*, 1015-1030 e1016. 10.1016/j.cell.2018.07.028.
Schiemann, J., Schlaudraff, F., Klose, V., Bingmer, M., Seino, S., Magill, P.J., Zaghloul, K.A., Schneider,
G., Liss, B., and Roeper, J. (2012). K-ATP channels in dopamine substantia nigra neurons control
bursting and novelty-induced exploration. Nat Neurosci *15*, 1272-1280. 10.1038/nn.3185.
Schwarting, R.K., and Huston, J.P. (1996). The unilateral 6-hydroxydopamine lesion model in
behavioral brain research. Analysis of functional deficits, recovery and treatments. Prog Neurobiol *50*,
275-331. 10.1016/s0301-0082(96)00040-8.
Sharott, A., Magill, P.J., Harnack, D., Kupsch, A., Meissner, W., and Brown, P. (2005). Dopamine
depletion increases the power and coherence of beta-oscillations in the cerebral cortex and
subthalamic nucleus of the awake rat. Eur J Neurosci *21*, 1413-1422. 10.1111/j.1460-
9568.2005.03973.x.
Sitzia, G., Mantas, I., Zhang, X., Svenningsson, P., and Chergui, K. (2020). NMDA receptors are altered
in the substantia nigra pars reticulata and their blockade ameliorates motor deficits in experimental
parkinsonism. Neuropharmacology *174*, 108136. 10.1016/j.neuropharm.2020.108136.
Styr, B., Gonen, N., Zarhin, D., Ruggiero, A., Atsmon, R., Gazit, N., Braun, G., Frere, S., Vertkin, I.,
Shapira, I., et al. (2019). Mitochondrial Regulation of the Hippocampal Firing Rate Set Point and Seizure
Susceptibility. Neuron *102*, 1009-1024 e1008. 10.1016/j.neuron.2019.03.045.
Subramaniam, M., Althof, D., Gispert, S., Schwenk, J., Auburger, G., Kulik, A., Fakler, B., and Roeper, J.
(2014a). Mutant alpha-synuclein enhances firing frequencies in dopamine substantia nigra neurons by
oxidative impairment of A-type potassium channels. J Neurosci *34*, 13586-13599.
10.1523/JNEUROSCI.5069-13.2014.
Surmeier, D.J. (2007). Calcium, ageing, and neuronal vulnerability in Parkinson's disease. Lancet
Neurol *6*, 933-938. 10.1016/S1474-4422(07)70246-6.
Surmeier, D.J., Halliday, G.M., and Simuni, T. (2017a). Calcium, mitochondrial dysfunction and slowing
the progression of Parkinson's disease. Exp Neurol *298*, 202-209. 10.1016/j.expneurol.2017.08.001.
Surmeier, D.J., Obeso, J.A., and Halliday, G.M. (2017b). Selective neuronal vulnerability in Parkinson
disease. Nat Rev Neurosci *18*, 101-113. 10.1038/nrn.2016.178.
Tarfa, R.A., Evans, R.C., and Khaliq, Z.M. (2017). Enhanced Sensitivity to Hyperpolarizing Inhibition in
Mesoaccumbal Relative to Nigrostriatal Dopamine Neuron Subpopulations. J Neurosci *37*, 3311-3330.
10.1523/JNEUROSCI.2969-16.2017.
Tinkhauser, G., Torrecillos, F., Pogosyan, A., Mostofi, A., Bange, M., Fischer, P., Tan, H., Hasegawa, H.,
Glaser, M., Muthuraman, M., et al. (2020). The Cumulative Effect of Transient Synchrony States on
Motor Performance in Parkinson's Disease. J Neurosci *40*, 1571-1580. 10.1523/JNEUROSCI.1975-
19.2019.
Turrigiano, G. (2012). Homeostatic synaptic plasticity: local and global mechanisms for stabilizing
neuronal function. Cold Spring Harb Perspect Biol *4*, a005736. 10.1101/cshperspect.a005736.

Ungerstedt, U. (1968). 6-Hydroxy-dopamine induced degeneration of central monoamine neurons. Eur J Pharmacol *5*, 107-110. 10.1016/0014-2999(68)90164-7.

Ungless, M.A., and Grace, A.A. (2012). Are you or aren't you? Challenges associated with physiologically identifying dopamine neurons. Trends Neurosci *35*, 422-430. 10.1016/j.tins.2012.02.003.

West, T.O., Berthouze, L., Halliday, D.M., Litvak, V., Sharott, A., Magill, P.J., and Farmer, S.F. (2018). Propagation of beta/gamma rhythms in the cortico-basal ganglia circuits of the parkinsonian rat. J Neurophysiol *119*, 1608-1628. 10.1152/jn.00629.2017.

Wichmann, T., Bergman, H., and DeLong, M.R. (2018). Basal ganglia, movement disorders and deep brain stimulation: advances made through non-human primate research. J Neural Transm (Vienna) *125*, 419-430. 10.1007/s00702-017-1736-5.

Winkler, C., Kirik, D., Bjorklund, A., and Cenci, M.A. (2002). L-DOPA-induced dyskinesia in the intrastriatal 6-hydroxydopamine model of parkinson's disease: relation to motor and cellular parameters of nigrostriatal function. Neurobiol Dis *10*, 165-186. 10.1006/nbdi.2002.0499.

Zaghloul, K.A., Blanco, J.A., Weidemann, C.T., McGill, K., Jaggi, J.L., Baltuch, G.H., and Kahana, M.J. (2009). Human substantia nigra neurons encode unexpected financial rewards. Science *323*, 1496-1499. 10.1126/science.1167342.

Zhang, W., Xiong, B.R., Zhang, L.Q., Huang, X., Yuan, X., Tian, Y.K., and Tian, X.B. (2021). The Role of the GABAergic System in Diseases of the Central Nervous System. Neuroscience. 10.1016/j.neuroscience.2021.06.037.