Recovery of the full *in vivo* firing range in post-lesion surviving DA SN neurons associated with Kv4.3-mediated pacemaker plasticity

Lora Kovacheva[1*], Josef Shin[1,2], Josefa Zaldivar-Diez[1,3], Johanna Mankel[1], Navid Farassat[1,4], Kauê Machado Costa[1,5], Poonam Thakur[1,6], José Obeso[3] & Jochen Roeper[1*]

1 Institute of Neurophysiology, Neuroscience Center, Goethe University, Frankfurt, Germany

2 Department of Neurology, University Medical Centre of the Johannes Gutenberg University Mainz, Mainz, Germany

3 HM CINAC (Centro Integral de Neurociencias Abarca Campal), Hospital Universitario HM Puerta del Sur, HM Hospitales. Madrid, Spain.
CIBERNED, Instituto de Salud Carlos III, Madrid, Spain.
Medical School, CEU-San Pablo University, Madrid, Spain

4 Eye Center, Medical Center-University of Freiburg, Freiburg im Breisgau, Germany

5 Department of Psychology, University of Alabama at Birmingham, AL, USA

6 School of Biology, Indian Institute of Science Education and Research (IISER)-Thiruvananthapuram, Kerala, India

[*]Corresponding authors:
Dr. Lora Kovacheva & Prof. Dr. Jochen Roeper
Institute for Neurophysiology
Neuroscience Center, Goethe University
Theodor-Stern-Kai 7, 60590 Frankfurt, Germany
e-mail: roeper@em.uni-frankfurt.de & lorask@gmail.com
webpage: https://www.neurophys-frankfurt.de
Tel: +49-69-6301-84091/92

Abstract

Dopamine (DA) neurons in the substantia nigra (SN) control several essential functions, including the voluntary movement, learning and motivated behavior. Healthy DA SN neurons show diverse firing patterns *in vivo*, ranging from slow pacemaker-like activity (1-10 Hz) to transient high frequency bursts (<100 Hz), interspersed with pauses that can last hundreds of milliseconds. Recent *in vivo* patch experiments have started to reveal the subthreshold mechanisms underlying this physiological diversity, but the impact of challenges like cell loss on the *in vivo* activity of adult DA SN neurons, and how these may relate to behavioral disturbances, are still largely unknown. We investigated the *in vivo* electrophysiological properties of surviving SN DA neurons after partial unilateral 6-OHDA lesions, a single-hit, non-progressive model of neuronal cell loss. We show that mice subjected to this model have an initial motor impairment, measured by asymmetrical rotations in the open field test, which recovered over time. At 3 weeks post-lesion, when open field locomotion was strongly impaired, surviving DA SN neurons showed a compressed *in vivo* dynamic firing range, characterized by a 10-fold reduction of *in vivo* burst firing compared to controls. This *in vivo* phenotype was accompanied by pronounced *in vitro* pacemaker instability. In contrast, in the chronic post-lesion phase (>2 months), where turning symmetry in open field locomotion had recovered, surviving SN DA neurons displayed the full dynamic range of *in vivo* firing, including *in vivo* bursting, similar to controls. The normalized *in vivo* firing pattern was associated with a 2-fold acceleration of stable *in vitro* pacemaking, mediated by Kv4.3 potassium channel downregulation. Our findings demonstrate the existence of a homeostatic pacemaker plasticity mechanism in surviving DA SN neurons after pronounced cell loss.

Keywords

dopamine, 6-OHDA, Kv4.3 potassium channel, substantia nigra, pacemaker, homeostasis, intrinsic plasticity

Introduction

Midbrain dopamine (DA) neurons in the substantia nigra (SN) project mainly to the dorsal striatum (DS), where they release dopamine in an activity-dependent manner, further shaped by additional mechanisms at the distal axon (Liu et al., 2022, Kramer et al., 2023). The resulting temporal and spatial dynamics of DA release in DS are relevant for multiple functions, including the control of voluntary movements, as evidenced by the cardinal features of Parkinson disease (PD), which is characterized by the death of SN DA neurons. The *in vivo* activity pattern of healthy SN DA neurons is complex, ranging from slow pacemaker-like activity (1-10 Hz) to transient high frequency bursts (<100 Hz), and interspersed with pauses that can last hundreds of milliseconds. However, our understanding if and how SN DA neuron electrical activity adapts to cell loss, relevant during aging or PD, is limited.

We have previously shown that overexpression of mutant A53T-α-synuclein induced an age-dependent acceleration of *in vivo* and *in vitro* firing frequencies. This was associated with redox-mediated dysfunction of Kv4.3 channels and an upregulation of Kv4.3 expression, potentially representing a homeostatic attempt to regulate pacemaker frequency (Subramaniam, Althof, et al. 2014). This study, however, left a more fundamental question unanswered: do adult DA SN neurons have the ability for homeostatic regulation of firing pattern in response to challenge? For this purpose, we employed the well-established 6-Hydroxy-Dopamine (6-OHDA) chemical lesion model in order to induce an unilateral loss of about half of the DA SN neurons. 6-OHDA has been widely used for generating non-progressive loss of DA SN neurons (Rubi and Fritschy 2020; P. R. L. Parker, Lalive, and Kreitzer 2016; Ungerstedt 1968). We opted for an unilateral, partial, intrastriatal 6-OHDA model, previously well characterized by Bez et al. (Bez, Francardo, and Cenci 2016). Bez and colleagues, as well as others, demonstrated that these partial 6-OHDA DA lesions are non-progressive and follow a stereotypical time course: an initial damage phase dominated by cell loss, inflammation and behavioral impairment for about 3 weeks, which is followed by a chronic phase (studied for up to 20 month in mice), characterized by partial behavioral recovery, neurochemical and molecular adaptations, and – to a certain degree – axonal sprouting of remaining DA neurons (Bez, Francardo,

and Cenci 2016; Cenci and Björklund 2020; Kirik, Rosenblad, and Björklund 1998; Schwarting and

Huston 1996; Winkler et al. 2002). This framework, with its distinct phases of impairment and

subsequent partial functional recovery, provides a well-suited platform to investigate if and how the

in vivo firing properties of surviving DA SN neurons change over time. Previous electrophysiological

investigations of viable post-6-OHDA DA neurons have been limited, with only Hollerman & Grace

providing a pioneering dataset (Hollerman and Grace 1990). They found lesion-size-dependent

changes in *in vivo* firing properties of putative DA neurons in rats, including a decrease of *in vivo* burst

firing occurring with larger lesions (Hollerman and Grace 1990). Therefore, conducting

electrophysiological investigations at two distinct time-points within the context of a non-progressive

partial 6-OHDA lesion is expected to provide novel insights into the homeostatic properties of

identified DA SN neurons.

Here, we explore the electrophysiological properties, both *in vivo* and *in vitro,* of surviving SN DA

neurons in an intra-striatal unilateral 6-OHDA lesion mouse model, titrated to about 40% surviving SN

DA neurons. We selected two time points for recordings based on the continuous behavioral

characterization of the motor phenotypes.

Early behavioral impairments and long-term adaptations after partial lesion of substantia

nigra dopamine neurons

To characterize electrophysiological properties in surviving and identified SN DA neurons, we

employed a C57Bl6N mouse model with a partial 6-OHDA lesion of the nigrostriatal pathway. We

explored different 6-OHDA concentrations (0.2µg/µl - 2µg/µl) and volumes (2µL and 6µL) resulting in

a range of 0.4µg to 12µg 6-OHDA) to titrate the ipsilateral loss of DA SN neurons (Suppl. Fig. 1E for

details). To target 40-50% cell loss, we slowly (250 nl/min) infused 6 µl of 2µg/µl 6-OHDA (i.e. 24 min

total infusion time, total of 12 µg 6-OHDA) into the right DS. For controls, we infused vehicle (ACSF)

with the same rate and same volume into the right DS without observing any damage (Fig. 1B; Suppl.

Fig. 1-C). The loss of TH-positive SN neurons was quantified by unbiased stereology at 21 days and 3

months post-lesion (Fig. 1A).

We continuously characterized the motor phenotype of the model (i.e. open field locomotion, cylinder

test) until the two different end points, either 21-days post-lesion (early phase) or more than 60 days

post-lesion (late phase). In the first cohorts, the mice were sacrificed for TH immunohistochemistry in

midbrain and striatum. As shown in Fig. 1B, at the early phase, the number of TH-positive (i.e.

dopaminergic) SN neurons was reduced post-6-OHDA (lower left panels) compared to vehicle-control

mice (upper left panels). The right panel in Fig. 1B displays the stereological results of TH-positive SN

neurons for N=9 vehicle-treated and N=9 6-OHDA-infused male mice 21 days after their respective

infusions. The data were normalized to the stereological results of respective contralateral SN DA

neurons. In contrast to vehicle-infusions, where no ipsilateral loss of SN DA neurons was detected, the

number of ipsilateral surviving SN DA neurons in the early post-6-OHDA phase was on average reduced

to about 40% (Fig. 1B; vehicle: 100 ± 2.64 %; 6-OHDA: 36.85 ± 3.46%; p < 0.0001, Mann-Whitney test).

We also carried out the unbiased stereology for TH-positive SN neurons in the late phase and detected

a very similar degree of loss (Suppl. Fig. 1C, vehicle: 102.6 ± 7.9 %; 6-OHDA: 39.96 ± 3.7 %; p = 0.0262,

Mann-Whitney test; please add N numbers). Importantly, these results indicated that SN DA cell loss

was non-progressive throughout our observation period of more than 2 months. In addition, we

analyzed the axonal compartment of DA midbrain neurons by determining striatal TH-optical

densities, both for early and late phase (Suppl. Fig. 1). Similar to the midbrain DA cell body counts, we

found a stable reduction of about 50% of TH-immunosignal in the ipsilateral DS, again normalized to

the contralateral side (Suppl. Fig. 1A 21st day - vehicle: 98.4 ± 2.2 %; 6-OHDA: 42.5 ± 3.0 %; Suppl. Fig.

1B > 64 days - vehicle: 94.4 ± 2.7 %; 6-OHDA: 48.3 ± 3 %), both in early and late phase. By comparison,

the ventral striatum was only mildly affected (ca. 20 % reduction, Suppl. Fig. 1A 21st day - vehicle: 94.6

± 2.0 %; 6-OHDA: 81.4 ± 3.0 %; Suppl. Fig. 1B > 64 days - vehicle: 97.8 ± 2.0 %; 6-OHDA: 75.2 ± 5.0 %).

In summary and in accordance with previous studies, our model induced stable 60% loss of SN DA

neurons, associated with a stable 50% reduction of DS TH-immunoreactivity throughout the entire

observation period (Bez, Francardo, and Cenci 2016; Cenci and Björklund 2020; Schwarting and Huston

1996). This stability provided a suitable framework to identify time-dependent homeostatic changes

in surviving SN DA neurons.

Fig 1 C shows the data from continuous behavioral monitoring of unilateral 6-OHDA and vehicle-

treated mice before and up to 68 days post-infusion. Based on previous studies (Cenci and Björklund

2020; Schwarting and Huston 1996), we focused on the dynamics of drug-free spontaneous turning

behavior during open field locomotion (Fig. 1C, lefts panels). In contrast to vehicle-infused mice, which

displayed a stable symmetric ratio (ca. 50%) of ipsi- to contral-lateral turning throughout the entire

experiment, 6-OHDA-infused mice showed a dramatic shift to ipsilateral turning immediately after

treatment (Figure 1C, right panel). Interestingly, ipsilateral turns occurred in long sequences of up to

40 individual turns, a pattern not observed in controls (Suppl. Fig. 3). Importantly, these turning

sequences as well as the overall ipsilateral bias gradually recovered completely over a 2-months post-

lesion period (two-way-ANOVA, p-value across time p<0.0001, p-value across groups p<0.0001,

significant difference between vehicle and 6-OHDA group till day 40, Šídák's multiple comparisons

test). Analysis of turning features, such as diameter or velocity, revealed that the recovered

contralateral turns were similar to turns performed by vehicle-treated mice (Suppl. Fig. 4). Also, open

field locomotion, quantified as total track length per session, recovered almost completely (Suppl. Fig.

1D). However, other motor behavior, the contralateral paw use during spontaneous rearing in the

cylinder test, did not show significant recovery throughout the experimental period (Fig. 1D) (21st

post-infusion day: vehicle 38.5 ± 5.2%, 6-OHDA 5.7 ± 1.4%, p < 0.0001, Mann-Whitney test; 68th post-

infusion day: vehicle 63.7± 6.6%, 6-OHDA 3.2 ± 1.8%, p < 0.0001, Mann-Whitney test). Finally,

repeating the continuous behavioral monitoring and the quantification of TH immunohistochemistry

in midbrain and striatum with female C57Bl6N mice, infused with either 6-OHDA or vehicle, showed

very similar results and gave no indications for sex differences (Suppl. Fig. 2).

To investigate the functional properties in immunohistochemically-identified surviving SN DA neurons,

we performed *in vivo* extracellular recordings combined with juxtacellular neurobiotin (NB) labeling

and *in vitro* whole-cell patch-clamp recordings combined with neurobiotin filling for two timepoints:

at an early timepoint (21st day after lesion, early phase), where a strong turning asymmetry was

present, and at a later time period (> 64 days post-lesion, late phase) when turning symmetry had

recovered.

Surviving SN DA neurons in the early post-6-OHDA phase exhibit impaired firing properties

both *in vivo* and *in vitro*

At the early post-6-OHDA phase, we explored the *in vivo* spontaneous activity of surviving identified

SN DA neurons in isoflurane-anesthetized mice, using single-unit extracellular recordings combined

with juxtacellular neurobiotin labelling and post-hoc TH immunohistochemistry (see schema Fig. 2A).

Figure 2B shows the *in vivo* extracellular electrical activity (top panel) and identification of a

representative SN DA neuron (lower right panel) from a control mouse, 3 weeks after vehicle infusion. Note the typical combination of irregular single spike firing with the occurrence of transient fast burst discharges (bursts were defined by the 80/160ms Grace criteria; see methods for details) and longer pauses. The interspike interval (ISI) histogram, which captures ISIs from the entire recording time of 10 min for this cell, displays the typical features of an *in vivo* bursty DA neuron, i.e. a bimodal ISI distribution representing intra- and interburst intervals over a broad dynamic range between ~30-1000 ms (lower left panel). This recorded and labelled cell was localized in the medial SN and characterized as TH-immunopositive. In comparison, Fig. 2C displays the representative *in vivo* activity of a surviving and identified SN DA neuron 3 weeks after 6-OHDA-infusion. Here, while irregular single-spike pacemaking was present similar to controls, fast burst events and slow pauses were almost completely absent (top panel). Indeed, the ISI histogram of this neuron, for the entire 10 min of recording, showed a compressed dynamic firing range and was well-described by a unimodal, Gaussian-like distribution without fast intraburst and slow interburst intervals (Fig. 2C, lower left panel). This recorded and labelled cell was also localized in the medial SN and identified as TH-immunopositive.

All *in vivo* recorded DA neurons, like the two representative cells shown above, from the early phase were successfully juxtacellularly labeled with NB, post-hoc immunohistochemically identified and localized within the ventral midbrain (Fig 2D, vehicle: n =28, N =15; 6-OHDA: n =18, N =16). For best comparison, we restricted our analysis to those DA midbrain neurons localized in the medial half of the SN (mSN), where it was possible to sample a sufficient number of surviving and spontaneously active DA neurons after 6-OHDA lesion. In contrast, we only managed to detect a small number of active DA neurons in the lateral SN (lSN), the most vulnerable part of the DA midbrain (Figure 2D). Consequently, Figure 2 E-H compares the group data of surviving mSN DA neurons at early phase between vehicle and 6-OHDA treatment. We detected an about 60% reduction of the mean firing rates for post-6-OHDA mSN DA neurons compared to vehicle-treated controls (Fig.2E; vehicle: mean frequency = 7.5 ± 0.9 Hz, 6-OHDA: mean frequency = 4.7 ± 0.6 Hz, p = 0.0184, Mann-Whitney test) and no significant differences in overall firing regularity (Fig. 2F; vehicle: CV = 53.9 ± 7.8 %, 6-OHDA: CV =

36.1 ± 3.7 %, p = 0.1694, Mann-Whitney test). Moreover, we observed a significant 10-fold reduction

of bursting - expressed as the percentage of spikes fired in bursts (SFB) and burst rate - in mSN DA

neurons in the 6-OHDA group compared to vehicle controls (Fig. 2G, vehicle: SFB = 31.2 ± 7.7; 6-OHDA:

SFB = 2.2 ± 1.1 %; p = 0.0032, Welch's t test; Fig. 2H, vehicle: burst rate = 0.34 ± 0.09 Hz, 6-OHDA: burst

rate = 0.04 ± 0.02 Hz; p = 0.0066, Welch's t test). .

All other analyzed firing parameters, such as intra-burst firing frequencies, identified no additional

differences between the two groups (Suppl. Fig. 5A-G). In summary, we detected a rightward shift to

lower firing frequencies with a corresponding compression of the dynamic *in vivo* firing range in mSN

DA neurons surviving the 6-OHDA lesion (Suppl. Fig. 5I). Thus, in addition to DA cell loss, the

dysfunction of surviving DA neurons might be a novel contributing factor to the extensive motor

impairment , observed during the early post-6-OHDA phase. In particular, transient bursts of DA SN

neurons have been shown to provide start and stop signals for motor control (see discussion for

details).

Burst discharges in DA neurons are orchestrated by the interplay of patterned synaptic inputs with

their intrinsic excitability, giving rise to different types of *in vivo* bursting (Otomo et al. 2020). To

identify a potential contribution of cell-autonomous changes in intrinsic excitability of surviving mSN

DA neurons to their impaired *in vivo* dynamics, we also studied these cells *in vitro* in synaptic isolation

(see Schema 2I). Figure 2J shows a representative, stable and regular pacemaker activity of an

identified mSN DA neuron from a vehicle-infused control (lower right panel), as well as a subthreshold

response to hyperpolarizing current injection (lower left panel). In contrast, pacemaker activity of mSN

DA neurons post-6-OHDA infusion was unstable and characterized by intermittent periods of action

potential failure (see Fig. 2K, upper panel). However, subthreshold responses were similar to controls

(lower right panel). Analogous to the *in vivo* experiments, all *in vitro* recorded DA neurons were NB

filled, identified, and mapped, resulting in a similar medial localization of surviving SN DA neurons (Fig.

2L). When *in vitro* pacemaker properties were compared between identified mSN DA neurons from

the vehicle and the 6-OHDA groups (Fig. 2L), we detected no differences in mean firing rates (Fig. 2M,

vehicle: firing rate = 1.7 ± 0.2 Hz, 6-OHDA: firing rate = 2.4 ± 0.3 Hz, p = 0.1448, Mann-Whitney test),

rebound delays or sag-amplitudes (Fig. 2O, vehicle: rebound delay = 584.9 ± 95.8 ms, 6-OHDA:

rebound delay = 429.6 ± 56.1 ms, p = 0.1738, Mann-Whitney test; Fig. 2P, vehicle: sag-amplitude =

15.3 ± 1.1 mV, 6-OHDA: sag-amplitude = 13.5 ± 0.96 mV, p = 0.1395, Mann-Whitney test). In contrast,

a significant, about 5-fold increase of pacemaker irregularity expressed as CV was detected (Fig.2N,

vehicle: CV = 6.5 ± 0.7 %; 6-OHDA: CV = 33.9 ± 5.8; p = 0.0007, Mann-Whitney test). Detailed

observation of the raw traces revealed that the increased CV was mainly mediated by a combination

of intermittent phases of firing failure, similar to the one shown in Fig. 2K, and episodes of higher

frequency firing (as quantified in the ISI distribution histogram, Suppl. Fig. 6H). We also showed that

these differences were preserved both in on-cell and whole-cell recordings indicating the latter did

not induce a differential bias in any of the two groups (e.g. different calcium buffering) (Suppl. Fig. 6

F-G).

In short, this data demonstrates functional alterations in DA SN neurons surviving three weeks after a

6-OHDA infusion. These neurons exhibited a compressed dynamic range, including a 10-fold reduction

of *in vivo* bursting. Furthermore, we observed a 5-fold decrease of *in vitro* cell-autonomous pacemaker

stability with intermittent failure of action potential firing. This compromised intrinsic *in vitro*

excitability likely contributes to the 10-fold reduction of high frequency *in vivo* firing.

In order to address the question of whether these altered properties represent stable adaptations or

reflect a transient state of damage, we also studied *in vivo* and *in vitro* electrophysiology of surviving

mSN DA neurons in the late phase after lesion, characterized by partial behavioral recovery of motor

functions.

Surviving mSN DA neurons in the late post-6-OHDA phase display full recovery of the *in vivo*

dynamic range associated with an accelerated *in vitro* pacemaker

We recorded and analyzed surviving mSN DA neurons in the late 6-OHDA phase analogous to the early

phase (compare Fig. 3A/I and Fig. 2A/I). In contrast to the early phase, we found no significant

differences in the electrophysiological *in vivo* properties between the post-lesional and vehicle-

infused groups in the late phase (compare Fig 3B with Fig. 3C). Among others (Suppl. Fig. 7), this

implies that mean firing frequencies, dynamic ranges, CV, SFB and burst rates did not differ in

chronically surviving mSN DA neurons compared to those in controls (Fig. 3E, vehicle: mean firing rate

= 6.9 ± 0.8 Hz; 6-OHDA: mean firing rate = 7.5 ± 1.1 Hz; p = 0.6038, Mann-Whitney test; Fig. 3F, vehicle:

CV = 54.9 ± 12.2% ; 6-OHDA: CV = 53.3 ± 12.6%, p > 0.9999, Mann-Whitney test; Fig. 3G, vehicle: SFB

= 20.2 ± 9.1%, 6-OHDA: SFB = 22.2 ± 8.9%, p = 0.8784, Mann-Whitney test; Fig. 3H, vehicle: burst rate

= 0.17 ± 0.08 Hz; 6-OHDA: burst rate = 0.30 ± 0.15 Hz, p = 0.4308, Mann-Whitney test). This suggests

that surviving mSN DA neurons show full recovery of their electrophysiological properties within 2-

months after the lesion. However, our *in vivo* results alone cannot distinguish between two scenarios:

first, a slow functional recovery process that eventually leads to return of the physiological activity

range or second, the return to the full dynamic firing range *in vivo* is more than simple repair to the

status quo ante but involves a homeostatic adaptation of the functional properties of the DA neuron

itself. To resolve this question, we also recorded the *in vitro* pacemaker properties of the surviving DA

mSN neurons in the late post-6-OHDA phase. Here, we found clear evidence for allostatic adaptation

of these DA neurons (i.e the presence of new homeostatic setpoint). When comparing intrinsic

pacemaker frequencies, we noticed that late 6-OHDA survivors discharged almost 2-fold faster

compared to vehicle-controls (compare Fig. 3J with 3K, Fig. 3M vehicle: firing rate = 1.7 ± 0.2 Hz; 6-

OHDA: firing rate = 2.7 ± 0.2 Hz, p = 0.0035, Mann-Whitney test). In contrast to speed, no significant

difference in pacemaker regularity was observed (Fig. 3N, vehicle: CV = 9.19 ± 1.30 %, 6-OHDA: CV =

16.10 ± 6.66 %, p = 0.1245, Mann-Whitney test). Also, the sag components and the rebound delay

were not significantly different (Fig 3O, vehicle: rebound delay = 512.8 ± 93.8 ms, 6-OHDA: rebound

delay = 375.0 ± 58.9 ms, p = 0.1994, Mann-Whitney test; Fig. 3P, vehicle: sag-amplitude = 16.4 ± 1.2

mV, 6-OHDA: sag-amplitude = 19.1 ± 0.9 mV, p = 0.0528, Mann-Whitney test). Other *in vitro* spike

features were also not significantly different between the two groups (Suppl. Fig. 8). This accelerated

pacemaker phenotype was also observed in metabolically intact DA neurons recorded in the on-cell

configuration but further accelerated in the whole-cell mode (Suppl. Fig. 8 F-G).

In summary, functional recovery of the *in vivo* dynamic firing range of surviving mSN DA neurons was

associated with homeostatic plasticity of pacemaking (i.e. an allostatic shift of pacemaker setpoint).

Surviving mSN DA neurons not only recovered from instable pacemaking at the early phase but

doubled their discharge rates in late phase.

Accelerated pacemaker caused by downregulation of Kv4.3 channels in late surviving mSN

DA neurons

As Kv4.3 A-type channels have been shown to be powerful regulators of DA pacemaker frequency

(Subramaniam, Althof, et al. 2014; Subramaniam, Kern, et al. 2014; Liss et al. 2001; Khaliq and Bean

2008; Tarfa, Evans, and Khaliq 2017), we tested their potential role for the observed pacemaker

acceleration in late phase DA mSN neurons by pharmacological occlusion experiments using 1µM

AmmTx3, a selective Kv4.3 channel blocker (see scheme Fig. 4A). The presence of 1µM AmmTx3 not

only accelerated discharge rates in both identified mSN DA neurons from vehicle- and 6-OHDA-infused

mice (see Fig. 4BC for representative cells; Fig. 4D for mapping) but, importantly, also eliminated

significant rate differences between the two treatment groups (Fig. 4E, vehicle: mean firing rate = 4.5

± 0.2 Hz, 6-OHDA: mean firing rate = 4.8 ± 0.3 Hz, p = 0.285; Fig. 4 F, vehicle: CV = 7.8 ± 1.3 %, 6-OHDA:

CV = 5.5 ± 0.4 %, p = 0.2517; both Mann-Whitney test). This result demonstrated that differences in

Kv4.3 channel function were the main driver of the late post-6OHDA accelerated pacemaker

phenotype. In addition, 1µM AmmTx3 also abolished the more subtle differences in rebound delays

and sag amplitudes (Fig. 4G, vehicle: rebound delay = 31.7 ± 2.4 ms, 6-OHDA: rebound delay = 28.4 ±

4.0 ms, p = 0.0952; Fig. 4H. vehicle: sag-amplitude = 15.6 ± 1.1 mV, 6-OHDA: sag-amplitude = 16.4 ±

0.9 mV, p = 0.7434; both Mann-Whitney test, Suppl. Fig. 9). In summary, these experiments

demonstrated that reduced Kv4.3 channel function in post-6-OHDA late phase DA mSN neurons had

a causal role in the phenotype of accelerated *in vitro* pacing.

To directly quantify the gating properties of transient outward (A-type) potassium currents mediated

by Kv4.3 channels, we carried out whole-cell voltage-clamp recordings (see scheme Fig. 4I-O). As

reported before (Liss et al. 2001), voltage-steps from negative holding potentials to the relevant

subthreshold range (e.g. -60 mV to -30 mV) activated large, fast inactivating outward currents in the

range of 1-10 nA in vehicle injected controls. In contrast, the same voltage steps activated smaller fast-

inactivating outward currents in post-6-OHDA late phase DA mSN neurons. These experiments

demonstrated about 50% smaller maximal A-type Kv4.3 potassium currents in late surviving DA mSN

neurons from 6-OHDA-treated mice compared to vehicle-treated controls (Fig. 4 M. vehicle: Kv4.3

channel current at -30mV = 3.8 ± 0.4 nA , 6-OHDA: Kv4.3 channel current at -30mV = 1.7 ± 0.3 nA, p <

0.0001, Mann-Whitney test). When comparing voltage-dependent gating parameters, we found no

changes in the biophysical channel properties apart from differences in the slope of the steady-state

activation curves (Fig. 4M, Suppl. Fig. 10 A-C).

By comparison, recordings of HCN currents in whole-cell voltage-clamp recordings showed no

differences in current amplitudes between the groups (Fig. 4N, vehicle: ΔHCN = 90.8 ± 10.7 pA, 6-

OHDA: ΔHCN = 94.9 ± 19.1 pA, p = 0.7375, Mann-Whitney test). These results substantiated the

selective role of reduced Kv4.3 function in causing the accelerated pacemaker in the post-6-OHDA late

phase surviving mSN DA neurons. As the reduction of Kv4.3 function might be caused by a number of

different mechanisms, including e.g. reduced protein expression and/or membrane delivery of Kv4.3

subunits as well as post-translational modifications of existing Kv4.3 channel complexes (e.g.

phosphorylation, redox-modification), we decided to assess Kv4.3 protein expression via

immunohistochemistry.

We performed Kv4.3 immunohistochemistry and confocal imaging in vehicle and 6-OHDA treated mice

both on the ipsilateral and contralateral infusion side (see scheme in Fig. 5A). We defined DA-selective

ROIs by using TH-immunopositive areas to determine the intensity of Kv4.3-immunosignals exclusively

within DA neurons (i.e. TH-signal based mask) (Subramaniam, Althof, et al. 2014). Figure 5B and 5C

compare Kv4.3-immunoreactivities in the midbrain between the contralateral control side and the

affected ipsilateral side during the late post-6-OHDA phase. Ipsilateral post-lesional TH-positive

neurons showed reduced Kv4.3-immunosignals compared to the control side. Semi-quantitative

comparison of TH-ROIs of the entire SN between contralateral control side and 6-OHDA-treated side

revealed a significant reduction of average Kv4.3-immunosignal on the 6-OHDA treated side of about

25% (Figure 5D, contralateral side n = 4004, ipsilateral side, n = 2645, D = 0.2014, p < 0.0001, two-

sample Kolmogorov–Smirnov test). A more detailed analysis of regional differences within the SN

revealed that the reduction in Kv4.3 immunoreactivity was significant for the medial SN (Figure 5E,

contralateral side, n = 4004: 73.03 ± 0.39, ipsilateral side, n = 2645: 54.8 ± 0.49, p < 0.0001, unpaired

t-test), where we recorded Kv4-currents, but even more pronounced in the lateral parts of the SN

(Suppl. Fig. 11C). As Kv4.3 channel subunits are expected to be mainly located on the cell membrane,

we also investigated whether Kv4.3-immunosignals were differentially distributed across cell

compartments. A robust, about 17% reduction of Kv4.3-immunosignals was observed in ipsilateral

post-lesional TH-positive neurons both for cell membrane and cytoplasm ROIs (Suppl. Fig. 11D). In

contrast, analogous quantification in vehicle-treated mice revealed no contra-ipsilateral differences

in Kv4.3-immunoreactivity (Suppl. Fig. 11 E-F). In summary, these immunohistochemical experiments

provided evidence for a reduced expression of Kv4.3 channel proteins in late post-6OHDA mSN DA

neurons. Reduced Kv4.3 protein expression is consistent with a reduction of functional Kv4.3 channels

in surviving mSN DA neurons as described above and suggests that pacemaker acceleration is

mediated by a reduced number of Kv4.3 channels.

Discussion

Our study provides the first combined *in vitro* & *in vivo* electrophysiological characterization of identified DA neurons in the substantia nigra surviving a partial 6-OHDA lesion. Studying these surviving SN DA neurons at two time points, we discovered time-dependent post-6-OHDA-selective differences of firing properties both *in vivo* and *in vitro*. Early after lesion, and coinciding with prominent behavioral impairments, we detected a selective and dramatic (about 10-fold) reduction of *in vivo* burst firing in identified surviving SN DA neurons. This is reminiscent of an early study by Hollermann & Grace (Hollerman and Grace 1990), who reported reduced bursting in putative SN DA neurons one week after a partial lesion. Considering the Berretta et al 2005 study (Berretta et al. 2005), these early electrophysiological effects may reside from 6-OHDA triggering a cascade of events, e.g. hyperpolarization via activation of K-ATP channels. At this early time point, we also found unstable and more irregular *in vitro* pacemaker activity in these cells, with no differences in mean firing rates, compared to those from vehicle-infused controls.

Conversely, at a later post-lesion time point (>2 months), when partial motor recovery had occurred, we detected no differences in the *in vivo* firing range – including high frequency bursts– between surviving SN DA neurons post-6-OHDA and those from vehicle-infused mice. Surprisingly, the *in vitro* pacemaker rates in post-6-OHDA DA survivors were not only stable but almost twice as fast compared to those from vehicle-controls. Finally, we pinpointed functional and protein Kv4.3 channel downregulation as the leading cause for this chronic post-lesional pacemaker acceleration, while the HCN channel function remained relatively stable. In essence, we identified a slow recovery of the dynamic *in vivo* firing range of surviving post-lesional DA SN neurons, linked to an allostatic acceleration of the intrinsic pacemaker by Kv4.3 channel downregulation.

Our results suggest a simple homeostatic scheme, where reducing the population size of DA SN neurons by half is at least in part compensated by a two-fold increase of pacemaker speed in the

remaining half of the DA SN cell population. Given the 10-fold reduction of *in vivo* bursting in the initial

period after the lesion, it further suggests that the pacemaker setpoint rate is coupled to the *in vivo*

high frequency burst rate in DA mSN neurons. It is possible that a burst-related calcium signal might

be detected and integrated over time to control pacemaker rate via e.g. tuning Kv4.3 expression.

Recent studies have implicated the mitochondrial calcium uniporter (MCU) in coupling bursts to

mitochondrial Ca^{2+}, acting as a calcium sensor for neuronal firing rate homeostasis (Katsenelson et al.

2022). Future mechanistic studies will be necessary to investigate how *in vivo* bursting and intrinsic

pacemaking are coupled in distinct DA subpopulations. It is however interesting to note that the very

first descriptive study of in vivo SN DA firing patterns had already found a positive correlation between

the degree of in vivo burst firing and single spike firing rate (A. A. Grace and Bunney 1984). In

agreement with this proposal, it was also previously found that an about 4-fold-reduction of *in vivo*

burst rate via DA-selective NR1 knockout resulted in a scaled reduction of *in vivo* firing rate (Zweifel

et al. 2009).

In contrast to the homeostatic role of flexible Kv4.3 expression identified in this study, we previously

identified Kv4.3 channels in SN DA neurons as a pathophysiological target in a transgenic mutant

(A53T-SNCA) α-synuclein mouse model (without cell loss) (Subramaniam, Althof, et al. 2014). In this

model, we found a pacemaker acceleration caused by oxidative impairment of Kv4.3 channels.

However, the activity of SN DA neurons in the α-synuclein model was accelerated also *in vivo.* Thus,

mutant α-synuclein expression also caused a shift of the *in vivo* firing range of SN DA neurons towards

higher frequency. While Kv4.3 subunits were downregulated at the protein level in post-lesional SN

DA neurons, mutant α-synuclein induced oxidative Kv4.3 dysfunction as well as protein upregulation

(Subramaniam, Althof, et al. 2014). It will be interesting to test in follow up studies if and how α-

synuclein pathology affects firing and burst rate homeostasis in DA SN neurons.

Beyond pacemaker adaptations in surviving SN DA neurons, we assume that their chronic *in vivo*

electrophysiological phenotype might also be reshaped by network level plasticity in response to

dopamine-depletion.

Our finding that the mean *in vivo* discharge rates were not different to controls in the presence of an

accelerated intrinsic pacemaker strongly suggests a shift of the synaptic excitation-inhibition (E-I)

balance towards more inhibition. Numerous studies have found altered synaptic inhibition in the

dopamine-depleted basal ganglia (recently reviewed in (Zhang et al. 2021)). In particular, Heo and

colleagues recently demonstrated a chronic E-I balance shift toward more inhibition across several PD

models including 6-OHDA (Heo et al. 2020). They identified a substantial contribution of additional

GABA synthesis and release from reactive astrocytes in the midbrain. It has also been well-established

that basal ganglia GABA neurons fire in more synchronized hence more effective fashion after DA

depletion (Cagnan, Duff, and Brown 2015; Milosevic et al. 2018; Phillips et al. 2020; Tinkhauser et al.

2020; Wichmann, Bergman, and DeLong 2018; Evans et al. 2020). This has recently been confirmed by

elegant *in vivo* single cell resolved studies using either gCAMP-based calcium monitoring or *in vivo*

patch-clamp approaches (Ketzef et al. 2017; Kravitz et al. 2010; J. G. Parker et al. 2018; P. R. L. Parker,

Lalive, and Kreitzer 2016; Sitzia et al. 2020). The accelerated pacemaker in response to enhanced net

inhibition would also be expected within the framework of homeostatic plasticity (Turrigiano 2012).

Our study has several limitations. First, we explored a single-hit, non-progressive partial lesion model,

which needs to be differentiated from *state-of-the-art* PD rodent models, which develop disease-

relevant α-synuclein pathology and show progressive loss of DA SN neurons (Thakur et al. 2017).

Second, we do not differentiate between DA SN subtypes. We recently showed that lateral SN DA

neurons projecting to DLS possess a distinct *in vivo* firing phenotype compared to medial SN DA

neurons projecting to either DLS or DMS in mice (Farassat et al. 2019). Therefore, recording from

lateral DA SN population in lesion models would be important. However, in our current partial 6-OHDA

lesion model, these lateral SN DA neurons were too severely lost to allow for more than anecdotal

post-lesion electrophysiological analysis (see Fig. 3). A milder version of the current model would be

needed to identify their properties and responses to partial lesion. Regarding the surviving DA neuron

in the medial SN, we previously showed that they have distinct projection targets including DLS, DMS

and lateral shell of nucleus accumbens (Farassat et al. 2019). In the current study, we have not aimed

for identification of axonal projections to avoid potentially confounding effects of additional brain

surgery. Nevertheless, this important issue should be addressed in follow-up studies by e.g. molecular

subtyping approaches (Heymann et al. 2020; Poulin et al. 2018; Saunders et al. 2018). Third, we are

aware that *in vivo* recordings carried out in isoflurane do not display the full dynamic spectrum of DA

SN firing compared to awake, freely moving animals (we have discussed the aspect extensively in

Farassat et al. 2019).

Finally, we would like to speculate about the possible implications of our findings for PD. In principle,

cell-loss induced pacemaker plasticity, as identified here, might have a dual nature. On one hand, they

might render the surviving DA SN neurons more robust in defending their phenotype and function. On

the other hand, the allostatic pacemaker acceleration may amplify the innate vulnerability of an

already at baseline metabolically and oxidatively challenged neuron type. The latter would be an

extension of the "stressful pacemaker" hypothesis of DA vulnerability (Chan et al. 2007; Surmeier

2007; Surmeier, Obeso, and Halliday 2017; Surmeier, Halliday, and Simuni 2017) with its potential

clinical implications(Guzman et al. 2018; Liss and Striessnig 2019; Ortner et al. 2017; Ortner 2021; The

Parkinson Study Group STEADY-PD 2020) . In summary, we identified the homeostatic capacity and

pacemaker allostasis of surviving DA SN neurons in response to cell loss.

Methods

Animals

Male and female C57Bl6/N mice (Charles River Laboratories) were used for the study. The mice were 8 weeks old, group housed and maintained on a 12-hour light-dark cycle. All experiments and procedures involving mice were approved by the German Regierungspräsidium Darmstadt (V54-19c20/15-F40/30). In total, 152 mice were used for this study (see table below).

	Early phase		Late phase	
	Mice (#)	Cells (#)	Mice (#)	Cells (#)
In vivo juxta & behavior – vehicle	15	12 – medial/ 28 – all	12	10 – medial/ 26 – all
In vivo juxta & behavior – 6-OHDA	16	9 – medial/ 16 – all	10	9 – medial/ 11 – all
In vitro – vehicle	3	16	3	12
In vitro – 6-OHDA	6	37	5	25
In vitro & AmmTx3 – vehicle	-	-	3	15
In vitro & AmmTx3 – 6-OHDA	-	-	3	14
In vitro voltage clamp experiments – vehicle	-	-	3	17
In vitro voltage clamp experiments – 6-OHDA	-		3	15
Female behavior experiments – vehicle, 6-OHDA	-		10	
Female histology – vehicle, 6-OHDA	-		8	
6-OHDA dose response	12			
	Mice (#)		Mice (#)	
Kv4.3 Immunochemistry	8		8	
SN quantification – vehicle	9		3	
SN quantification – 6-OHDA	9		3	
Sum = 152				

Stereotactic 6-OHDA infusion

All surgeries were performed under general anesthesia in areflexic state. Prior to the induction of anesthesia, a premedication of 0.2 mg/kg atropine (atropine-sulfate, Braun Melsungen AG,

Melsungen) was given as an intraperitoneal (i.p.) injection to stabilize circulation. Anesthesia was induced in a plastic chamber, which was flooded with 5% Isoflurane (Florene®, AbbVie Deutschland GmbH & Co. KG, Ludwigshafen, Germany) in pure oxygen (0.4 l/min). For maintenance of anesthesia, isoflurane was delivered through a breathing mask with a flow rate of 0.35 l/min and its concentration was regulated to 1.5-2.2% using an adjustable vaporizer (Uno, Zevenaar, Netherlands). The depth of anesthesia was controlled by testing the toe pinch reflex and the breathing rate (1-2Hz). Body temperature (36°C) and respiration were constantly monitored. Lidocaine/prilocaine ointment (25 mg/g, Emla® creme, AstraZeneca GmbH, 22876 Wedel) was applied prior to surgery and after suturing of the wound as local anesthetics. Additional analgesia was provided by subcutaneous injection of carprofen (4 mg/kg in NaCl, Rimadyl®, Pfilzer GmbH, Berlin, Germany) after infusion. Eye lubricant (Visidic, Bausch and Lomb, Berlin, Germany) was used to protect eyes from desiccation.

Desipramine hydrochloride (20 mg/kg, Sigma Aldrich) was injected i.p. 20-40 min before intracranial infusions to prevent 6-OHDA uptake by noradrenergic neurons. The desipramine solution was prepared in sterile, isotonic NaCl solution (B. Braun Melsungen AG, Germany) at the day of surgery. The infusion solutions are based on sterile artificial cerebrospinal fluid (ACSF, Harvard Apparatus, Holliston, MA, USA) with 0.02% L-ascorbic acid (used also as a vehicle solution). The 6-OHDA solution (0.2% 6-hydroxydopamine hydrochloride in ACSF with 0.02% L-ascorbic acid) was prepared at the day of infusion, stored on ice, and shielded from light.

During surgery, the animals were placed on a heating pad and were fixed in a stereotactic frame (Model 1900, Kopf Instruments, Tujunga, USA) with a stereotactic arm and a connected three-way digital positioning display. The scalp was opened by a longitudinal cut to expose the skull with bregma and lambda on display. With a centering scope (Model 1915, Kopf Instruments, Tujunga, USA), the bregma-lambda distance was measured and examined for suitable anatomy (4.4 ± 0.2 mm distance). Afterwards, the skull was aligned to a reference frame with a stereotaxic alignment indicator (Model 1905, Kopf Instruments, Tujunga, USA) and the manipulator system was referenced to bregma.

Using a stereotaxic drill (Model 1911, Kopf Instruments, Tujunga, USA) with a 500 µm diameter drill

bit, a hole above the right striatum was drilled (coordinates: ML: +1.9 mm, AP: +0.5 mm to bregma).

ACSF or 6-OHDA solution were loaded to a 10 µl NanoFil syringe (World Precision Instruments Inc.,

Sarasota, FL, USA) with a 35G blunt needle, which was mounted on a MicroSyringe Pump (UMP3-1,

World Precision Instruments) and controlled by a SYSMicro4 Controller (World Precision Instruments).

Using the stereotactic arm, the needle was slowly lowered (about 750 µm/min) to a position of -2.2

mm below the brain surface (infusion site coordinates: ML: +1.9 mm, AP: +0.5 mm, DV: -2.2 mm to

bregma). Anatomical references are based on Franklin and Paxinos (2008). A volume of 6 µl was

infused with a flow rate of 250 nl/min. Once the volume was infused, the needle rested for 5 minutes

in that position before it was slowly moved out of the brain. Directly before and after infusion, proper

functioning of the syringe system and the needle was checked. Finally, after suture the animal was

placed on a heating pad for full recovery. Oats, wet food pallets and water were placed inside the cage

to ease consumption.

Behavioral testing

Open field

Spontaneous locomotion (track length, wall distance, time in center and number of rearings) and

rotations of all mice were monitored in open field (50 × 50 cm, center 30 × 30 cm; red illumination, 3

lx) for 10 min in 3 baseline sessions and every 4th or 7th day post-infusion of ACSF/6-OHDA till the day

of *in vivo* or *in vitro* experiment (e.g. 21st or >68th post-operative day). The open field was cleaned

before and after each mouse with 0,1% acetic acid in distilled water. Using a video tracking system

(Viewer II/III, Biobserve) spontaneous behavior was recorded and analyzed both online and offline.

Data was extracted from Viewer as Excel-tables and the final analysis was made with custom made

Matlab-scripts.

 Cylinder test

Forelimb use during explorative activity was explored with cylinder test. The test was performed at

corresponding termination time point (20-21st and 64th post-infusion day). Mice were placed

individually in a glass beaker (9 cm diameter, 19 cm height) at room light and were video recorded

with a camera (Logitech HD Webcam C615) for about 5 min. No habituation was allowed before video

recording. The glass cylinder was cleaned before and after every mouse with 0,1% acetic acid in

distilled water. Only weight-bearing wall contacts made by each and both forelimb on the cylinder

wall were scored. Wall exploration was expressed in terms of the percentage of contralateral to the

infusion side (in the 6-OHDA-infused mice also impaired forepaw) to all forelimb wall contacts.

In vivo electrophysiology

Extracellular recording

In vivo extracellular single-unit activities of SN and VTA neurons were recorded in ACSF-infused

(vehicle) and 6-OHDA-infused mice, similar procedures were used in other studies from our lab

(Farassat et al. 2019; Schiemann et al. 2012; Subramaniam, Althof, et al. 2014). Briefly, mice were

anesthetized (isoflurane; induction 4.5-5%, maintenance 1–2% in 0.4 l/min O2) and placed into a

stereotactic frame. The craniotomies were performed as described above to target the lateral SN (AP:

-3.08 mm, ML: 1.4 mm) and medial SN (AP: -3.08 mm, ML: 0.9 mm). Borosilicate glass electrodes (10–

25 MΩ; Harvard Apparatus, Holliston, MA, USA) were made using a horizontal puller (DMZ-Universal

Puller, Zeitz, Germany) and filled with 0.5 M NaCl, 10 mM HEPES (pH 7.4) and 1.5% neurobiotin (Vector

Laboratories, Burlingame, CA, USA). A micromanipulator (SM-6; Luigs & Neumann, Ratingen,

Germany) was used to lower the electrodes to the recording site. The single-unit activity of each

neuron was recorded for at least 10 minutes at a sampling rate of 12.5 kHz (for firing pattern analyses),

and then for another one minute at a sampling rate of 20 kHz (for the fine analysis of AP waveforms).

Signals were amplified 1000x (ELC-03M; NPI electronics, Tamm, Germany), notch- and bandpass-

filtered 0.3–5000 Hz (single-pole, 6 dB/octave, DPA-2FS, NPI electronics) and recorded on a computer

with an EPC-10 A/D converter (Heka, Lambrecht, Germany). Simultaneously, the signals were

displayed on an analog oscilloscope and an audio monitor (HAMEG Instruments CombiScope HM1508;

AUDIS-03/12M NPI electronics). Midbrain DA neurons were initially identified by their broad biphasic

AP (> 1.2 ms duration) and slow frequency (1–8 Hz) (Anthony A. Grace and Bunney 1984; Ungless and

Grace 2012). AP duration was determined as the interval between the start of initial upward

component and the minimum of following downward component.

Juxtacellular labeling of single neurons

In order to identify the anatomical location and neurochemical identity of the recorded neurons, they

were labeled post-recording with neurobiotin using the juxtacellular *in vivo* labeling technique (Pinault

1996). Microiontophoretic currents were applied (1–10 nA positive current, 200ms on/off pulse, ELC-

03M, NPI Electronics) via the recording electrode in parallel to the monitoring of single-unit activity.

Labeling was considered successful, when: the firing pattern of the neuron was modulated during

current injection (i.e., increased activity during on-pulse and absence of activity in the off-pulse), the

process was stable for at least 20s, and was followed by the recovery of spontaneous activity. This

procedure allowed for the exact mapping of the recorded DA neuron within the SN and VTA subnuclei

(Franklin and Paxinos 2012) using custom written scripts in Matlab (MathWorks, Natick, MA, USA),

combined with neurochemical identification using TH-immunostaining.

In vitro electrophysiology

Slice preparation

Animals were anesthetized by intraperitoneal injection of ketamine (250 mg/kg, Ketaset, Zoetis) and

medetomidine-hydrochloride (2.5 mg/kg, Domitor, OrionPharma) prior to intracardial perfusion using

ice-cold ACSF consisting of the following (in mM): 50 sucrose, 125 NaCl, 2.5 KCl, 25 NaHCO3, 1.25

NaH2PO4, 2.5 glucose, 6 MgCl2, 0.1 CaCl2 and 2.96 kynurenic acid (Sigma-Aldrich), oxygenated with

95% O2 and 5% CO2. Rostral coronal midbrain slices (bregma: -2.92 mm to -3.16 mm) were sectioned

at 250 μm using a vibrating blade microtome (VT1200s, Leica). Slices were incubated for 1 h before

recordings in a 37°C bath with oxygenated extracellular solution with extra 1μM AmmTx3, containing

the following (in mM): 22.5 sucrose, 125 NaCl, 3.5 KCl, 25 NaHCO3, 1.25 NaH2PO4, 2.5 glucose, 1.2

MgCl2 and 1.2 CaCl2.

In vitro patch-clamp recordings

Slices were placed in a heated recording chamber (37°C) that was perfused with oxygenated

extracellular solution at 2-4 ml min$_{-1}$. CNQX (20 μM; Biotrend), gabazine (SR95531, 4 μM; Biotrend)

DL-AP5 (10 μM; Cayman Chemical) were added to inhibit excitatory and inhibitory synaptic

transmission. For voltage clamp recordings, TTX (500nM; Tocris) was added to the extracellular

solution. Neurons were visualized using infrared differential interference contrast videomicroscopy

with a digital camera (VX55, Till Photonics) connected to an upright microscope (Axioskop 2, FSplus,

Zeiss). Patch pipettes were pulled from borosilicate glass (GC150TF-10; Harvard Apparatus LTD) using

a temperature-controlled, horizontal pipette puller (DMZ-Universal Puller, Zeitz). Patch pipettes (4-6

MΩ) were filled with a solution containing the following (in mM): 135 KGlu, KCl, 10 HEPES, 0.1 EGTA,

5 MgCl2, 0.075 CaCl2, 5 NaATP, 1 LiGTP, 0.1% neurobiotin, adjusted to a pH of 7.35 with KOH.

Recordings were performed using an EPC-10 patch-clamp amplifier (Heka electronics) with a sampling

rate of 20 kHz and a low-pass filter (Bessel, 5 kHz). For voltage clamp recordings only experiments with

uncompensated series resistance <10MO were included in this study and series resistance was

electronically compensated 75%. Neurons were held at a holding potential of -40mV to minimize HCN

activation. To determine Kv4.3 activation kinetics neurons were hyperpolarized to -80mV for 500ms

followed by varying voltage steps from -60mV to -20mV in increments of 5mV for 1s. For inactivation

kinetics neurons were hyperpolarized from -120mV to -20mV in increments of 10 mV for 1s followed

by a fixed voltage step to -20mV for 1s. For analysis of HCN currents neurons were hyperpolarized

from -80mV to -120mV in increments of 20mV for 1s. For analysis, action potential thresholds (mV)

were determined at $dV_m/dt > 10$ mV/ms.

Immunohistochemistry

Following *in vivo* recordings, animals were transcardially perfused, as described previously (Farassat

et al. 2019; Schiemann et al. 2012; Subramaniam, Althof, et al. 2014). Fixed brains were sectioned into

60μm (midbrain) or 100μm (forebrain) coronal sections using a vibrating microtome (VT1000S, Leica).

In vitro slices were fixed in paraformaldehyde after finishing the experiment. Sections were rinsed in

PBS and then incubated (in blocking solution (0.2 M PBS with 10% horse serum, 0.5% Triton X-100,

0.2% BSA). Afterwards, sections were incubated in carrier solution (room temperature, overnight)

with the following primary antibodies: polyclonal guinea pig anti-tyrosine hydroxylase (TH; 1:1000;

Synaptic Systems), monoclonal mouse anti-TH (1:1000; Synaptic Systems) or polyclonal rabbit anti-TH

(1:1000; Synaptic Systems); mouse anti-Kv4.3 (1:1000, Alomone Labs). In sequence, sections were

again washed in PBS and incubated (room temperature, overnight) with the following secondary

antibodies: goat anti-guinea pig 488, goat anti-rabbit 488, goat anti-mouse 488, goat anti-mouse 568

(all 1:750; Thermofisher). Streptavidin AlexaFluor-568 and Streptavidin AlexaFluor-488 (both 1:1000;

Invitrogen) were used for identifying neurobiotin-filled cells. Sections were then rinsed in PBS and

mounted on slides with fluorescent mounting medium (Vectashield, Vector Laboratories).

DAB immunocytochemistry

For DAB (3,3'-diaminobenzidine) staining procedures, a Vectastain ABC Staining Kit (Vector

Laboratories) was used. Coronal sections of midbrain (30 μm) areas were cut and rinsed in PBS (3x10

min). Similar to previous immunolabeling procedures, unspecific antigen binding sites were blocked

by incubation of the sections with blocking solution (60 min, room temperature). Subsequently,

sections were incubated with primary antibody against TH (rabbit anti TH) overnight, rinsed in PBS

(3x10 min), and were incubated with biotinylated secondary antibodies (biotinylated anti-rabbit) for

two hours at RT. In parallel, an avidin-biotin complex (ABC) was formed by pre-incubation of avidin

(1:1000) with biotinylated HRP (1:1000) in PBS for two hours at room temperature. Sections were

rinsed in PBS (3x10 min) prior to incubation with ABC solution (60 min, room temperature). Next,

sections were rinsed in PBS (2x10 min) and Tris-buffer (1x10 min). Finally, DAB oxidation was

developed by application of 2 % H_2O_2, 2 % $NiCl_2$ and 4 % DAB in Tris-buffer using a DAB Substrate Kit

(Vector Laboratories, Burlingame, USA). $NiCl_2$ enhances sensitivity and intensity of DAB precipitation

product. DAB oxidation was developed for 2 to 5 minutes and was stopped with Tris-buffer once a

specific high-contrast signal was detectable. Sections were rinsed in Tris-buffer (3x10 min) and

transferred onto gelatin-covered slides, air-dried overnight, and dehydrated in consecutive ascending

alcohol concentrations (50 %, 70 %, 90 % and 2x 100 %; 10 min each) followed by dehydration in xylol

(2x 100 %; 10 min each). Finally, sections were mounted under glass coverslips with hardening

mounting medium (Vectamount, Vector Laboratories, Burlingame, USA).

Unbiased stereology measurements

For quantification of total cell loss, TH-DAB labeled SN DA neurons were counted using unbiased

stereology based on optical dissection (Gundersen 1986). In coronal sections (30 μm), the region of

interest was selected based on anatomical landmarks including the medial lemniscus, which separates

SN and adjacent VTA. Stereological counting provides unbiased data based on random, systematic

sampling using an optical fractionator. This method involves counting of neurons with an optical

dissector, a three-dimensional probe placed through a reference space (Gundersen 1986). The optical

dissector forms a focal plane with a thin depth of field through the selected sections. Objects in focus

of this focal plane are located within the reference section and are counted, while objects outside of

the focal plane are not counted. On top of the optical dissector, a counting frame is applied. Counting

frames ensure that all neurons have equal probabilities of being selected, regardless of shape, size,

orientation, and distribution. To avoid counting capped neurons at the border of a section, an

additional guard zone was deployed at the upper and lower borders of each section. DA neurons

within the counting frame as well as those crossing the green line (acceptance line) were counted,

while DA neurons crossing the red line (rejection line) excluded. Moreover, only neurons with a

detectable nucleus in focus within the optical dissector were counted. For quantification of total cell

loss, StereoInvestigator software (V5, MicroBrightField, Colchester, USA) was used in combination

with BX61 microscope (Olympus, Hamburg, Germany). The region of interest was selected and marked

using a low magnification objective lens (2x, NA 0.25, Olympus) and 12-30 serial sections of 30 μm

thickness were counted, covering the entire caudo-rostral extent of the SN. To count the number of

DA neurons in the area of the SN pars compacta, a high magnification oil-immersion objective lens

(100x, NA 1.30, Olympus) (counting frame, 50 × 50 μm; sampling grid, 125 × 100 μm) was used. After

counting was finished, the total number of neurons was calculated by the software using the formula

described by West et al (West 1993).

Optical density measurements

Optical density measurements of TH-DAB labeled striatal sections were performed using ImageJ

software (http://rsbweb.nih.gov/ij/). Following TH-DAB labeling and TH-immunohistochemistry,

images of five coronal sections (100 μm) covering the rostrocaudal axis of the striatum were captured

using laser-scanning microscope (Nikon Eclipse90i, Nikon GmbH). Images were gray scale converted

and mean gray values of desired striatal areas were encircled and measured. Unspecific mean gray

values were measured in a defined cortical area (100x100 pixels) that displayed no specific TH signal

due to the absence of DA innervation and were subtracted. The ventral edge of lateral ventricles

served as an anatomical landmark to separate dorsal and ventral areas. For all animals, the

measurement from the ipsilateral to the infusion side were divided by the contralateral side to

calculate the relative optical density of the striatum.

Immunohistochemical Kv4.3 channel signal quantification

A Nikon Eclipse 90i microscope was used for fluorescent signal detection, excitation wavelength of 488 nm for TH-signal and 568 nm for Kv4.3-channel signal. From each animal, 4 midbrain slices covering the caudal, intermedial and rostral regions, were selected and imaged for overview with 4x magnification. Then 60x magnification was used to acquire data from 4 areas within each SN (4 images on the ipsilateral and 4 images on the contralateral to infusion side). All images were acquired using the same laser and camera settings. Images were exported from Nikon NIS-Elements Advanced Research (Version 4.20.03) software as 8-bit TIFF files for quantification. Data was analyzed using custom made Python 3.0 scripts with matplotlib, numpy, scimage and scipy modules. First, TH immunosignals were converted to a binary image via Otsu-thresholding algorithm to detect TH areas bigger than 400 pixels. Then, the resulting binary image was used as a mask for Kv4.3 channel immunosignal detection. For all ROIs surface areas and mean Kv.4.3 channel signal intensity was measured. By applying erosion and dilatation algorithms on the ROIs, membrane and cytoplasm areas were segregated, allowing isolation of Kv4.3 channel immunosignal intensity for these cell compartments. Background Kv4.3 channel immunosignal was quantified in TH-immunosignal areas below the Otsu-threshold. All data were then grouped according to medio-lateral and ipsi-/contralateral position for both vehicle and 6-OHDA group. Graphs and statistical analysis for this data was performed using Python custom made scripts.

Statistical Analysis

Spike train analyses

Spike time-stamps were extracted by thresholding above noise levels with IgorPro 6.02 (WaveMetrics, Lake Oswego, OR, USA). Firing pattern properties such as mean frequency, coefficient of variation (CV) and bursting measures were analyzed using custom scripts in Matlab. In order to estimate burstiness and intra-burst properties, we used the burst detection methods described in Grace & Bunney (A. A. Grace and Bunney 1984; Anthony A. Grace and Bunney 1984; Ungless and Grace 2012). All non-burst

related ISIs (excluding all ISIs that followed the Grace and Bunney criteria, as well as all pre- and post-

burst ISIs) were used to calculate the single spike firing frequency and single spike coefficient of

variation.

For analysis of general firing patterns, autocorrelation histograms (ACH) were plotted using custom

Matlab scripts. We used established criteria for classification of *in vivo* firing patterns based on visual

inspection of autocorrelograms (Farassat et al. 2019; Schiemann et al. 2012; Subramaniam, Althof, et

al. 2014): single spike-oscillatory (≥3 equidistant peaks with decreasing amplitudes), single spike-

irregular (<3 peaks, increasing from zero approximating a steady state), bursty-irregular (narrow peak

with steep increase at short ISIs) and bursty-oscillatory (narrow peak reflecting fast intraburst ISIs

followed by clear trough and repetitive broader peaks).

Statistics

Categorical data is represented as stacked bar graphs. To investigate the assumption of normal

distribution, we performed the single-sample Kolmogorov-Smirnov test. The Mann-Whitney-Test,

one-/two-way ANOVA were performed in non-parametric data to determine statistical significance.

Categorical parameters, such as ACH-based firing pattern, were analyzed with the Chi-squared test.

Statistical significance level was set to $p < 0.05$. All data values are presented as means ± SEM.

Statistical tests were performed using GraphPad Prism 9 (GraphPad Software, San Diego, CA, USA),

Matlab and Python. The scatter plots are represented with median or mean ± SEM. The resulting p

values were compared with Bonferroni-corrected α-level or Tukey *post hoc* comparison. A value of p

≤ 0.05 was considered to be statistically significant; $p \leq 0.05$ = * $p \leq 0.005$ = ** $p \leq 0.0005$ = ***. Graphs

were plotted using GraphPad Prism software (9.0c), Matlab and Python.

Acknowledgements

706 This study was supported by research grants to JR (DFG, CRC 1451). LK is a MD/PhD candidate at

707 TransMed, Gutenberg University Mainz. We thank Beatrice Fischer and Jasmine Sonntag for technical

708 assistance, Alexander Prinz for preliminary data on post-6-OHDA stereology.

Additional information

Author contributions

LK & JR designed the study. LK performed the lesions, behavioral & *in vivo* electrophysiology, JS & JM performed the *in vitro* experiments. JZ performed dose response lesion and female experiments. Analysis was carried out jointly by LK, JZ, KMC, JO and JR. NF taught LK *in vivo* electrophysiology and juxtacellular labelling. LK & JR wrote the manuscript.

Declaration of Interests

The authors declare no conflict of interests.

Author ORCIDs

Jochen Roeper   https://orcid.org/0000-0003-2145-8742

Lora Kovacheva   https://orcid.org/0000-0001-6999-1533

Josef Shin   https://orcid.org/0000-0002-3556-125X

Kauê Machado Costa   https://orcid.org/0000-0002-5562-6495

Poonam Thakur   https://orcid.org/0000-0002-4462-5550

Josefa Zaldivar-Diez   https://orcid.org/0000-0003-2317-3898

Ethics

Animal experimentation: All experiments and procedures involving mice were approved by the German Regierungspräsidium Darmstadt (V54-19c20/15-F40/30).

References

Berretta, Nicola, Peter S. Freestone, Ezia Guatteo, Denis De Castro, Raffaella Geracitano, Giorgio Bernardi, Nicola B. Mercuri, and Janusz Lipski. 2005. "Acute Effects of 6-Hydroxydopamine on Dopaminergic Neurons of the Rat Substantia Nigra Pars Compacta in Vitro." *NeuroToxicology* 26

(5): 869–81. https://doi.org/10.1016/j.neuro.2005.01.014.

Bez, Francesco, Veronica Francardo, and M. Angela Cenci. 2016. "Dramatic Differences in
Susceptibility to L-DOPA-Induced Dyskinesia between Mice That Are Aged before or after a
Nigrostriatal Dopamine Lesion." *Neurobiology of Disease* 94 (October): 213–25.
https://doi.org/10.1016/j.nbd.2016.06.005.

Cagnan, Hayriye, Eugene Paul Duff, and Peter Brown. 2015. "The Relative Phases of Basal Ganglia
Activities Dynamically Shape Effective Connectivity in Parkinson's Disease." *Brain* 138 (6): 1667–
78. https://doi.org/10.1093/brain/awv093.

Cenci, M. Angela, and Anders Björklund. 2020. "Animal Models for Preclinical Parkinson's Research:
An Update and Critical Appraisal." *Progress in Brain Research* 252: 27–59.
https://doi.org/10.1016/bs.pbr.2020.02.003.

Chan, C. Savio, Jaime N. Guzman, Ema Ilijic, Jeff N. Mercer, Caroline Rick, Tatiana Tkatch, Gloria E.
Meredith, and D. James Surmeier. 2007. "'Rejuvenation' Protects Neurons in Mouse Models of
Parkinson's Disease." *Nature* 447 (7148): 1081–86. https://doi.org/10.1038/nature05865.

Evans, Rebekah C., Emily L. Twedell, Manhua Zhu, Jefferson Ascencio, Renshu Zhang, and Zayd M.
Khaliq. 2020. "Functional Dissection of Basal Ganglia Inhibitory Inputs onto Substantia Nigra
Dopaminergic Neurons." *Cell Reports* 32 (11): 108156.
https://doi.org/10.1016/j.celrep.2020.108156.

Farassat, Navid, Kauê Machado Costa, Strahinja Stojanovic, Stefan Albert, Lora Kovacheva, Josef Shin,
Richard Egger, et al. 2019. "In Vivo Functional Diversity of Midbrain Dopamine Neurons within
Identified Axonal Projections." *ELife* 8 (October). https://doi.org/10.7554/eLife.48408.

Franklin, KBJ, and George Paxinos. 2012. *Paxinos and Franklin's The Mouse Brain in Stereotaxic
Coordinates. Elsevier*.

Grace, A. A., and B. S. Bunney. 1984. "The Control of Firing Pattern in Nigral Dopamine Neurons: Burst
Firing." *Journal of Neuroscience* 4 (11): 2877–90. https://doi.org/10.1523/jneurosci.04-11-
02877.1984.

Grace, Anthony A., and Benjamin S. Bunney. 1984. "The Control of Firing Pattern in Nigral Dopamine
Neurons: Single Spike Firing." *Journal of Neurosc* 4 (11): 2866–76.

Gundersen, H. J.G. 1986. "Stereology of Arbitrary Particles." *Journal of Microscopy* 143 (1): 3–45.
https://doi.org/10.1111/j.1365-2818.1986.tb02764.x.

Guzman, Jaime N., Ema Ilijic, Ben Yang, Javier Sanchez-Padilla, David Wokosin, Dan Galtieri, Jyothisri
Kondapalli, Paul T. Schumacker, and D. James Surmeier. 2018. "Systemic Isradipine Treatment
Diminishes Calcium-Dependent Mitochondrial Oxidant Stress." *Journal of Clinical Investigation*
128 (6): 2266–80. https://doi.org/10.1172/JCI95898.

Heo, Jun Young, Min Ho Nam, Hyung Ho Yoon, Jeongyeon Kim, Yu Jin Hwang, Woojin Won, Dong Ho
Woo, et al. 2020. "Aberrant Tonic Inhibition of Dopaminergic Neuronal Activity Causes Motor
Symptoms in Animal Models of Parkinson's Disease." *Current Biology* 30 (2): 276-291.e9.
https://doi.org/10.1016/j.cub.2019.11.079.

Heymann, Gabriel, Yong S. Jo, Kathryn L. Reichard, Naomi McFarland, Charles Chavkin, Richard D.
Palmiter, Marta E. Soden, and Larry S. Zweifel. 2020. "Synergy of Distinct Dopamine Projection
Populations in Behavioral Reinforcement." *Neuron* 105 (5): 909-920.e5.
https://doi.org/10.1016/j.neuron.2019.11.024.

Hollerman, Jeffrey R., and Anthony A. Grace. 1990. "The Effects of Dopamine-Depleting Brain Lesions on the Electrophysiological Activity of Rat Substantia Nigra Dopamine Neurons." *Brain Research* 533 (2): 203–12. https://doi.org/10.1016/0006-8993(90)91341-D.

Katsenelson, Maxim, Ilana Shapira, Eman Abbas, Kristina Jevdokimenko, Boaz Styr, Antonella Ruggiero, Saba Aïd, et al. 2022. "IGF-1 Receptor Regulates Upward Firing Rate Homeostasis via the Mitochondrial Calcium Uniporter." *Proceedings of the National Academy of Sciences of the United States of America* 119 (33): 1–10. https://doi.org/10.1073/pnas.2121040119.

Ketzef, M., G. Spigolon, Y. Johansson, A. Bonito-Oliva, G. Fisone, and G. Silberberg. 2017. "Dopamine Depletion Impairs Bilateral Sensory Processing in the Striatum in a Pathway-Dependent Manner." *Neuron* 94 (4): 855-865.e5. https://doi.org/10.1016/j.neuron.2017.05.004.

Khaliq, Zayd M., and Bruce P. Bean. 2008. "Dynamic, Nonlinear Feedback Regulation of Slow Pacemaking by A-Type Potassium Current in Ventral Tegmental Area Neurons." *Journal of Neuroscience* 28 (43): 10905–17. https://doi.org/10.1523/JNEUROSCI.2237-08.2008.

Kirik, Deniz, Carl Rosenblad, and Anders Björklund. 1998. "Characterization of Behavioral and Neurodegenerative Changes Following Partial Lesions of the Nigrostriatal Dopamine System Induced by Intrastriatal 6-Hydroxydopamine in the Rat." *Experimental Neurology* 152 (2): 259–77. https://doi.org/10.1006/exnr.1998.6848.

Kravitz, Alexxai V., Benjamin S. Freeze, Philip R.L. Parker, Kenneth Kay, Myo T. Thwin, Karl Deisseroth, and Anatol C. Kreitzer. 2010. "Regulation of Parkinsonian Motor Behaviours by Optogenetic Control of Basal Ganglia Circuitry." *Nature* 466 (7306): 622–26. https://doi.org/10.1038/nature09159.

Liss, Birgit, Oliver Franz, Sabine Sewing, Ralf Bruns, Henrike Neuhoff, and Jochen Roeper. 2001. "Tuning Pacemaker Frequency of Individual Dopaminergic Neurons by Kv4.3L and KChip3.1 Transcription." *EMBO Journal* 20 (20): 5715–24. https://doi.org/10.1093/emboj/20.20.5715.

Liss, Birgit, and Jorg Striessnig. 2019. "The Potential of L-Type Calcium Channels as a Drug Target for Neuroprotective Therapy in Parkinson's Disease." *Annual Review of Pharmacology and Toxicology* 59: 263–89. https://doi.org/10.1146/annurev-pharmtox-010818-021214.

Milosevic, Luka, Suneil K. Kalia, Mojgan Hodaie, Andres M. Lozano, Alfonso Fasano, Milos R. Popovic, and William D. Hutchison. 2018. "Neuronal Inhibition and Synaptic Plasticity of Basal Ganglia Neurons in Parkinson's Disease." *Brain* 141 (1): 177–90. https://doi.org/10.1093/brain/awx296.

Ortner, Nadine J. 2021. "Voltage-Gated Ca2+ Channels in Dopaminergic Substantia Nigra Neurons: Therapeutic Targets for Neuroprotection in Parkinson's Disease?" *Frontiers in Synaptic Neuroscience* 13 (February): 1–11. https://doi.org/10.3389/fnsyn.2021.636103.

Ortner, Nadine J., Gabriella Bock, Antonios Dougalis, Maria Kharitonova, Johanna Duda, Simon Hess, Petronel Tuluc, et al. 2017. "Lower Affinity of Isradipine for L-Type Ca2+ Channels during Substantia Nigra Dopamine Neuron-like Activity: Implications for Neuroprotection in Parkinson's Disease." *Journal of Neuroscience* 37 (28): 6761–77. https://doi.org/10.1523/JNEUROSCI.2946-16.2017.

Otomo, Kanako, Jessica Perkins, Anand Kulkarni, Strahinja Stojanovic, Jochen Roeper, and Carlos A. Paladini. 2020. "In Vivo Patch-Clamp Recordings Reveal Distinct Subthreshold Signatures and Threshold Dynamics of Midbrain Dopamine Neurons." *Nature Communications* 11 (1): 1–15. https://doi.org/10.1038/s41467-020-20041-2.

Parker, Jones G., Jesse D. Marshall, Biafra Ahanonu, Yu Wei Wu, Tony Hyun Kim, Benjamin F. Grewe, Yanping Zhang, et al. 2018. *Diametric Neural Ensemble Dynamics in Parkinsonian and Dyskinetic*

States. *Nature*. Vol. 557. https://doi.org/10.1038/s41586-018-0090-6.

Parker, Philip R.L., Arnaud L. Lalive, and Anatol C. Kreitzer. 2016. "Pathway-Specific Remodeling of
Thalamostriatal Synapses in Parkinsonian Mice." *Neuron* 89 (4): 734–40.
https://doi.org/10.1016/j.neuron.2015.12.038.

Phillips, Ryan S., Ian Rosner, Aryn H. Gittis, and Jonathan E. Rubin. 2020. "The Effects of Chloride
Dynamics on Substantia Nigra Pars Reticulata Responses to Pallidal and Striatal Inputs." *ELife* 9:
1–35. https://doi.org/10.7554/ELIFE.55592.

Pinault, Didier. 1996. "A Novel Single-Cell Staining Procedure Performed in Vivo under
Electrophysiological Control: Morpho-Functional Features of Juxtacellularly Labeled Thalamic
Cells and Other Central Neurons with Biocytin or Neurobiotin." *Journal of Neuroscience Methods*
65 (2): 113–36. https://doi.org/10.1016/0165-0270(95)00144-1.

Poulin, Jean Francois, Giuliana Caronia, Caitlyn Hofer, Qiaoling Cui, Brandon Helm, Charu
Ramakrishnan, C. Savio Chan, Daniel A. Dombeck, Karl Deisseroth, and Rajeshwar Awatramani.
2018. "Mapping Projections of Molecularly Defined Dopamine Neuron Subtypes Using
Intersectional Genetic Approaches." *Nature Neuroscience* 21 (9): 1260–71.
https://doi.org/10.1038/s41593-018-0203-4.

Rubi, Lena, and Jean Marc Fritschy. 2020. "Increased GABAergic Transmission in Neuropeptide Y-
Expressing Neurons in the Dopamine-Depleted Murine Striatum." *Journal of Neurophysiology*
123 (4): 1496–1503. https://doi.org/10.1152/jn.00059.2020.

Saunders, Arpiar, Evan Z. Macosko, Alec Wysoker, Melissa Goldman, Fenna M. Krienen, Heather de
Rivera, Elizabeth Bien, et al. 2018. "Molecular Diversity and Specializations among the Cells of
the Adult Mouse Brain." *Cell* 174 (4): 1015-1030.e16. https://doi.org/10.1016/j.cell.2018.07.028.

Schiemann, Julia, Falk Schlaudraff, Verena Klose, Markus Bingmer, Susumu Seino, Peter J. Magill,
Kareem A. Zaghloul, Gaby Schneider, Birgit Liss, and Jochen Roeper. 2012. "K-ATP Channels in
Dopamine Substantia Nigra Neurons Control Bursting and Novelty-Induced Exploration." *Nature*
Neuroscience 15 (9): 1272–80. https://doi.org/10.1038/nn.3185.

Schwarting, R. K.W., and J. P. Huston. 1996. "Unilateral 6-Hydroxydopamine Lesions of Meso-Striatal
Dopamine Neurons and Their Physiological Sequelae." *Progress in Neurobiology* 49 (3): 215–66.
https://doi.org/10.1016/S0301-0082(96)00015-9.

Sitzia, Giacomo, Ioannis Mantas, Xiaoqun Zhang, Per Svenningsson, and Karima Chergui. 2020. "NMDA
Receptors Are Altered in the Substantia Nigra Pars Reticulata and Their Blockade Ameliorates
Motor Deficits in Experimental Parkinsonism." *Neuropharmacology* 174 (May): 108136.
https://doi.org/10.1016/j.neuropharm.2020.108136.

Subramaniam, Mahalakshmi, Daniel Althof, Suzana Gispert, Jochen Schwenk, Georg Auburger, Akos
Kulik, Bernd Fakler, and Jochen Roeper. 2014. "Mutant α-Synuclein Enhances Firing Frequencies
in Dopamine Substantia Nigra Neurons by Oxidative Impairment of A-Type Potassium Channels."
Journal of Neuroscience 34 (41): 13586–99. https://doi.org/10.1523/JNEUROSCI.5069-13.2014.

Subramaniam, Mahalakshmi, Beatrice Kern, Simone Vogel, Verena Klose, Gaby Schneider, and Jochen
Roeper. 2014. "Selective Increase of in Vivo Firing Frequencies in DA SN Neurons after
Proteasome Inhibition in the Ventral Midbrain." *European Journal of Neuroscience* 40 (6): 2898–
2909. https://doi.org/10.1111/ejn.12660.

Surmeier, D. James. 2007. "Calcium, Ageing, and Neuronal Vulnerability in Parkinson's Disease."
Lancet Neurology 6 (10): 933–38. https://doi.org/10.1016/S1474-4422(07)70246-6.

Surmeier, D. James, Glenda M. Halliday, and Tanya Simuni. 2017. "Calcium, Mitochondrial Dysfunction
and Slowing the Progression of Parkinson's Disease." *Experimental Neurology* 298: 202–9.
https://doi.org/10.1016/j.expneurol.2017.08.001.

Surmeier, D. James, José A. Obeso, and Glenda M. Halliday. 2017. "Selective Neuronal Vulnerability in
Parkinson Disease." *Nature Reviews Neuroscience* 18 (2): 101–13.
https://doi.org/10.1038/nrn.2016.178.

Tarfa, Rahilla A., Rebekah C. Evans, and Zayd M. Khaliq. 2017. "Enhanced Sensitivity to Hyperpolarizing
Inhibition in Mesoaccumbal Relative to Nigrostriatal Dopamine Neuron Subpopulations." *Journal
of Neuroscience* 37 (12): 3311–30. https://doi.org/10.1523/JNEUROSCI.2969-16.2017.

Thakur, Poonam, Ludivine S. Breger, Martin Lundblad, Oi Wan Wan, Bengt Mattsson, Kelvin C. Luk,
Virginia M.Y. Lee, John Q. Trojanowski, and Anders Björklund. 2017. "Modeling Parkinson's
Disease Pathology by Combination of Fibril Seeds and α-Synuclein Overexpression in the Rat
Brain." *Proceedings of the National Academy of Sciences of the United States of America* 114
(39): E8284–93. https://doi.org/10.1073/pnas.1710442114.

The Parkinson Study Group STEADY-PD. 2020. "Isradipine Versus Placebo in Early Parkinson Disease A
Randomized Trial The." *Ann Intern Med* 63 (8): 1–18. https://doi.org/10.7326/M19-
2534.Isradipine.

Tinkhauser, Gerd, Flavie Torrecillos, Alek Pogosyan, Abteen Mostofi, Manuel Bange, Petra Fischer,
Huiling Tan, et al. 2020. "The Cumulative Effect of Transient Synchrony States on Motor
Performance in Parkinson⇔s Disease." *Journal of Neuroscience* 40 (7): 1571–80.
https://doi.org/10.1523/JNEUROSCI.1975-19.2019.

Turrigiano, Gina. 2012. "Homeostatic Synaptic Plasticity: Local and Global Mechanisms for Stabilizing
Neuronal Function." *Cold Spring Harbor Perspectives in Biology* 4 (1): 1–18.
https://doi.org/10.1101/cshperspect.a005736.

Ungerstedt, Urban. 1968. "6-Hydroxy-Dopamine Induced Degeneration of Central Monoamine
Neurons." *European Journal of Pharmacology* 5 (1): 107–10. https://doi.org/10.1016/0014-
2999(68)90164-7.

Ungless, Mark A., and Anthony A. Grace. 2012. "Are You or Aren't You? Challenges Associated with
Physiologically Identifying Dopamine Neurons." *Trends in Neurosciences* 35 (7): 422–30.
https://doi.org/10.1016/j.tins.2012.02.003.

West, Mark J. 1993. "New Stereological Methods for Counting Neurons." *Neurobiology of Aging* 14
(4): 275–85. https://doi.org/10.1016/0197-4580(93)90112-O.

Wichmann, Thomas, Hagai Bergman, and Mahlon R. DeLong. 2018. "Basal Ganglia, Movement
Disorders and Deep Brain Stimulation: Advances Made through Non-Human Primate Research."
Journal of Neural Transmission 125 (3): 419–30. https://doi.org/10.1007/s00702-017-1736-5.

Winkler, Christian, Deniz Kirik, Anders Björklund, and M. Angela Cenci. 2002. "L-DOPA-Induced
Dyskinesia in the Intrastriatal 6-Hydroxydopamine Model of Parkinson's Disease: Relation to
Motor and Cellular Parameters of Nigrostriatal Function." *Neurobiology of Disease* 10 (2): 165–
86. https://doi.org/10.1006/nbdi.2002.0499.

Zhang, W., B.R. Xiong, L.Q. Zhang, X. Huang, X. Yuan, Y.K. Tian, and X.B. Tian. 2021. "The Role of the
GABAergic System in Diseases of the Central Nervous System." *Neuroscience*.

Zweifel, Larry S., Jones G. Parker, Collin J. Lobb, Aundrea Rainwater, Valeriez Wall, Jonathan P. Fadok,
Martin Darvas, et al. 2009. "Disruption of NMDAR-Dependent Burst Firing by Dopamine Neurons

Provides Selective Assessment of Phasic Dopamine-Dependent Behavior." *Proceedings of the*
National Academy of Sciences of the United States of America 106 (18): 7281–88.
https://doi.org/10.1073/pnas.0813415106.

Figures

Figure 1. Unilateral striatal 6-OHDA mouse model with stable loss of SN DA neurons resulted in delayed partial behavioral recovery.

(A) Experimental design, illustrating timeline of behavioral assays with termination points – groups ended either on the 21st, or later than the 64th post-infusion day. (B) Left panel: TH-DAB staining of the SN, 10x magnification and 60x magnification for vehicle and 6-OHDA-infused mouse. Scalebar left 200µm, scalebar right 25µm. Right panel: ratio of ipsilateral (infusion side) to contralateral side of surviving TH-positive neurons in SN at 21st post-infusion day. (C) Left: spontaneous locomotion of mice in open field arena for a 10 min session. Left, examples of an ACSF-infused mouse (vehicle), right example of a 6-OHDA-infused mouse at baseline (upper panels), 4th post-infusion day (middle panels) and 68th post-infusion day (lower panels). Right: ratio of ipsilateral to contralateral turning behavior for all experimental groups plotted against session days. Note recovery in the 6-OHDA-treated mice from day 4 to day 68 after initial strong turning bias (>90%). (D) Cylinder test quantified by ratio of contra- to ipsilateral forepaw use. Note significant loss of contralateral forepaw involvement, both at 21 and 64 days. All data are presented as mean ± standard error of mean (SEM).

Figure 2. Surviving SN DA neurons at early post-6-OHDA phase exhibited a compressed dynamic range with a 10-fold decrease in *in vivo* bursting and a 5-fold decree in *in vitro* pacemaker regularity

(A), (I) Experimental design, illustrating timeline of behavioral assays, followed by terminal *in vivo* juxtacellular recordings (A) or by terminal *in vitro* whole-cell recordings (I) at the 21st post-infusion day. (B), (C) Top: 10s original recording trace of spontaneous *in vivo* single-unit activity from SN DA neurons in vehicle (B) and 6-OHDA-infused mouse (C). Scalebars: 1s, 0.2mV. Below, left: corresponding ISI histograms. Below, right: corresponding confocal images of juxtacellularly labelled and immunohistochemically identified DA neuron. Note the sparse bursting of the surviving SN DA neuron from the 6-OHDA-infused mouse at early phase. (D), (L) Anatomical mapping of all extracellularly recorded and juxtacellularly labelled DA neurons (D), and of all in vitro recorded and filled DA neurons (L), projected to bregma -3.08 mm. Location of example SN DA neurons in (B), (C), (J), (K) are highlighted. Smaller symbols represent DA neurons that have been recorded and identified, but not included in the group data analysis in (E-H). (E-H) Scatter dot-plots, showing significant decrease of *in vivo* mean firing rate (E), percentage of SFB (G) and burst rate (H) and no significant differences in CV (F) between the vehicle and 6-OHDA-infused mice. Note the 10-fold decrease in SFB for 6-OHDA-infused mice. (J), (K) Top: 10s original recording trace of *in vitro* whole-cell recording of spontaneous SN DA neuron activity in a vehicle (J) and 6-OHDA-infused (K) mouse. Scalebars: 1s, 20mV. Note that the 6-OHDA DA neuron has a highly irregular pacemaking. Below, left: corresponding to hyperpolarizing current injection. Below, right: confocal images of NB filled and immunohistochemically identified DA neuron. (M-P) Scatter dot-plots, showing no difference in *in vitro* mean firing rate (M), rebound delay (O) or sag-component (P) and a 5-fold increase in CV (N). Immunohistochemical imaging for all four DA neurons are displayed in 10x and 60x magnifications (green, TH; red, NB), scalebars: 200μm, 20μm. All data are presented as mean ± standard error of mean (SEM).

Figure 3. Surviving SN DA neurons at late 6-OHDA phase recovered *in vivo* burst firing and doubled their intrinsic pacemaker frequency *in vitro*.

(A), (I) Experimental design, illustrating timeline of behavioral assays, followed by terminal *in vivo* juxtacellular recordings (A) or by terminal *in vitro* whole-cell recordings (I) after >64 post-infusion days. (B), (C) Top: 10s original recording trace of spontaneous *in vivo* single-unit activity from SN DA neurons in vehicle (B) and 6-OHDA-infused mouse (C). Scalebars: 1s, 0.2mV. Below, left: corresponding ISI histograms. Below, right: corresponding confocal images of juxtacellularly labelled and immunohistochemically identified DA neuron. (D), (L) Anatomical mapping of all extracellularly recorded and juxtacellularly labelled DA neurons (D), and of all *in vitro* recorded and filled DA neurons (L), projected to bregma -3.40 mm (D) and -3.08mm (L). Location of example SN DA neurons in (B), (C), (J), (K) are highlighted. Smaller symbols represent DA neurons that have been recorded and identified, but not included in the group data analysis in (E-H). (E-H) Scatter dot-plots, showing no significant difference in mean firing rate (E), CV (F), SFB (G) and burst rate (H). (J), (K) Top: 10s original recording trace of *in vitro* whole-cell recording of spontaneous activity from SN DA neurons in a vehicle (J) and 6-OHDA-infused (K) mouse. Scalebars: 1s, 20mV. Note that the 6-OHDA DA neuron has enhanced, but regular pacemaking. Below, left, corresponding to hyperpolarizing current injection. Below, right, confocal images of NB filled and immunohistochemically identified DA neuron. (M-P) Scatter dot-plots, showing doubling of the firing rate (M), no difference in CV (N), decrease in rebound delay (O) and increase of sag-component (P) for the 6-OHDA-treated mice in comparison to vehicle group. Immunohistochemical imaging for all four DA neurons are displayed in 10x and 60x magnifications (green, TH; red, NB), scalebars: 200µm, 20µm. All data are presented as mean ± standard error of mean (SEM).

Figure 4. Enhanced pacemaker frequency of SN DA neurons at the late post-6-OHDA phase was mediated by K$_v$4.3 channel downregulation

(A), (I) Experimental design, illustrating timeline of behavioral assays, followed by terminal *in vitro* whole-cell recordings under K$_v$4.3 channel blocker AmmTx3 (A) or *in vitro* voltage-clamp whole-cell recordings (I) after >64 post-infusion days. (B), (C) Top: 10s original recording traces of *in vitro* whole-cell recording of spontaneous activity from DA SN neurons in AmmTx3-preincubated slices for a vehicle (B) and 6-OHDA-infused (C) mouse. Below, left: corresponding hyperpolarizing current injection. Scalebars: 1s, 20mV. Below, right: confocal images of NB filled and immunohistochemically identified DA neuron. (D), (L) Anatomical mapping of all in vitro recorded and filled DA neurons, projected to bregma -2.92 mm (D) and -3.16mm (L). (E-H) Scatter dot-plots, showing no differences in mean firing rate (E), coefficient of variation (F), rebound delay duration (G) and sag amplitude (H). (J), (K) Original recording trace of in vitro voltage-clamp recordings from DA SN neurons in a vehicle (J) and 6-OHDA-infused (K) mouse. Top, right: Zoom in from a K$_v$4.3 channel activation. Top, left: Zoom in from a K$_v$4.3 channel inactivation. Below, right: Zoom in from HCN-channel activation. Below, right, confocal images of NB filled and immunohistochemically identified DA neuron. Scalebars first row: 3nA, 100ms. Scalebars below, left: 5nA, 500ms. Note the small K$_v$4.3 channel in/activation peak in a surviving DA neuron from a 6-OHDA mouse in comparison the one from an ACSF treated mouse. Immunohistochemical imaging for all four DA neurons are displayed in 10x and 60x magnifications (green, TH; red, NB), scalebars: 200μm, 20μm. All data are presented as mean ± standard error of mean (SEM). (M) Scatter dot-plots, showing a significant, half of maximum K$_v$4.3 channel conductance in surviving DA neurons. (N) Scatter dot-plots, showing no difference in HCN channel delta current (I$_{peak}$ - I$_{steady}$) between groups. (O) Normalized K$_v$4.3 current (I$_{peak}$ - I$_{steady}$) at different voltage steps,

resulting in inactivation and activation curve for both groups. All data are presented as mean ± standard error
of mean (SEM).

Figure 5. Surviving DA SN neurons at the late post-6-OHDA phase showed lower K$_v$4.3 channel

immunohistochemical signal

(A) Experimental design, illustrating timeline of behavioral assays, followed by multi-labeling

immunohistochemistry for exploring K$_v$4.3-channel expression differences after >64 post-infusion days. (B), (C)

Top: 4x magnification of midbrain of a 6-OHDA-infused mouse, >64 days post-lesion – contralateral side (B), and

corresponding ipsilateral side (C). Middle: 60x magnification in the highlighted area from 4x image (green, TH;

red, K$_v$4.3). Bottom, left: zoom-in on an example ROI (highlighted in 60x image). Bottom, right: color-coded

Kv4.3-channel immunohistochemical signal intensity in the example ROI. Note K$_v$4.3-channel signal decrease in

surviving DA neurons on the ipsilateral to the injection side. (D) Histogram showing intensity of K$_v$4.3

immunosignals for all TH-positive ROIs, from ipsilateral, lesioned, side (in red) and from contralateral side (in

grey). Inset, same data shown as a cumulative distribution. Note a clear right-shift to lower intensities for the

ipsilateral side. (E) Comparison of mean TH-colocalized K$_v$4.3 immunosignals from medial SN from ipsilateral,

lesioned, side, and contralateral, as control side. All data are presented as mean ± standard error of mean (SEM).

1006

1007

1008

1009   Supplementary Figures
1010

1011   **Supplementary Figure 1.**

1012

(A), (B) Comparison of ipsilateral, infusion side, to contralateral side as percentage of relative TH
immunohistochemistry signal in the dorsal striatum (dSTR) and ventral striatum (vSTR), at the corresponding
early (A) and late phase (B). Note the stability of TH density loss in the dorsal striatum through time. (C) Ratio of
ipsilateral (infusion side) to contralateral side of surviving TH-positive neurons in SN in the late phase. (D) Mean
track length from all mice for each open field session. Note the post-infusion drop in performed track in the 6-
OHDA groups, which gradually recovers. (Infusion day marked as a thin gray line.) (E) Ratio of TH+ cell loss in SNc
(ipsilateral (infusion side) to contralateral side) >21st post-infusion day from different infused 6-OHDA doses.

Supplementary **Figure** **2.**

1021

(A) Ratio of ipsilateral to contralateral turning behavior for female ACSF and 6-OHDA infused groups plotted

against session days and with underlayed the corresponding ratio from the male ACSF and 6-OHDA groups as a

mean-line. (B) Ratio of ipsilateral (infusion side) to contralateral side of surviving TH-positive neurons in SN at

>64st post-infusion day from female ACSF and 6-OHDA infused mice. (C) Comparison of ipsilateral, infusion side,

to contralateral side as percentage of relative TH immunohistochemistry signal in the dorsal striatum (dSTR) at

>64st post-infusion day from female ACSF and 6-OHDA infused mice.

Supplementary   Figure   3.

1030   For two representative vehicle- and 6-OHDA-treated mice schematic representation of turning sequences. Left

1031   turn sequences are plotted as negative values, whereas right turn sequences are plotted as positive values. The

1032   count represents the number of same-direction turns within a sequence. On the left panels are turning

1033   sequences for the control animal and on the right panels for the 6-OHDA-infused animal, correspondingly in a

1034   baseline session (A), 4th post-operative day (B), 20th post-operative day (C) and 64th post-operative day (D).

1035

1038   Correlation pair plots comparing turning features, such as velocity (cm/s), diameter (cm) and duration for all

1039   contralateral turns (left panels) and all ipsilateral turns performed from all vehicle-infused mice (in orange) and

1040   all 6-OHDA-treated mice (in blue) across different time points (baseline, 4th, 20th and 64th post-infusion day).

Supplementary **Figure** **5.**

Scatter dot-plots, showing no significant difference of in vivo mean intra-burst (IB) frequency (A), mean

maximum (max) intra-burst (IB) frequency (B), burst duration (C), number of spikes per burst (D), single spike

frequency (SSF) (E), single spike coefficient of variance (CV) (F) and action potential (AP) width (G) (H) between

the vehicle and 6-OHDA-infused mice in the early phase. (H) Normalized stacked bar plots of different in vivo

firing patterns based on ACH. (I) ISI distributions from all in vivo recorded and labeled mSN DA neurons from

vehicle- and 6-OHDA-treated mice in the early phase. Inset, cumulative representation of the same distributions.

1049

Supplementary **Figure** **6.**

1051

Scatter dot-plots, showing no significant difference of in vitro afterhyperpolarization (AHP) minimum (A),

threshold (B), spike amplitude (C), action potential (AP) width (D) and input resistance (E) between the vehicle

and 6-OHDA-infused mice in the early phase. (F), (G) Paired scatter dot-plots of firing frequency (F) and

coefficient of variance (G) during on-cell recording and whole-cell recording in the early phase. (H) ISI

distributions from all in vitro whole-cell recorded and labeled mSN DA neurons from vehicle- and 6-OHDA-

treated mice in the early phase. Inset, cumulative representation of the same distributions. (I) Paired scatter

dot-plot indicating no significant difference between the mean voltage during spike-pauses and the rest of the

firing activity.

1060

1061

 Supplementary Figure 7.

1063

Scatter dot-plots, showing no significant difference of in vivo mean intra-burst (IB) frequency (A), mean

1065 maximum (max) intra-burst (IB) frequency (B), burst duration (C), number of spikes per burst (D), single spike

1066 frequency (SSF) (E), single spike coefficient of variance (CV) (F) and action potential (AP) width (G) (H) between

1067 the vehicle and 6-OHDA-infused mice in the late phase. (H) Normalized stacked bar plots of different in vivo

1068 firing patterns based on ACH. (I) ISI distributions from all in vivo recorded and labeled mSN DA neurons from

1069 vehicle- and 6-OHDA-treated mice in the late phase. Inset, cumulative representation of the same distributions.

Supplementary Figure **8.**

1071

Scatter dot-plots, showing no significant difference of in vitro afterhyperpolarization (AHP) minimum (A),

threshold (B), spike amplitude (C), action potential (AP) width (D) and input resistance (E) between the vehicle

and 6-OHDA-infused mice in the late phase. (F), (G) Paired scatter dot-plots of firing frequency (F) and coefficient

of variance (G) during on-cell recording and whole-cell recording in the late phase. (H) ISI distributions from all

in vitro whole-cell recorded and labeled mSN DA neurons from vehicle- and 6-OHDA-treated mice in the late

phase. Inset, cumulative representation of the same distributions.

Supplementary Figure 9.

1079

Scatter dot-plots, showing minor significant difference of in vitro afterhyperpolarization (AHP) minimum (A), and

no difference in threshold (B), spike amplitude (C), action potential (AP) width (D) and input resistance (E)

between the vehicle and 6-OHDA-infused mice in the late phase under 1µM AmmTx3. (F), (G) Paired scatter dot-

plots of firing frequency (F) and coefficient of variance (G) during on-cell recording and whole-cell recording in

the late phase under 1µM AmmTx3. (H) ISI distributions from all in vitro whole-cell recorded and labeled mSN

DA neurons from vehicle- and 6-OHDA-treated mice in the late phase under 1µM AmmTx3. Inset, cumulative

representation of the same distributions.

1087

1088   **Supplementary Figure 10.**

1089

1090   (A)-(D) Measurements of transient outward (A-type) potassium biophysical parameters in whole-cell voltage-

1091   clamp recordings of Kv4.3 channel activation and inactivation in vehicle relative to 6-OHDA infused mice. (E), (F)

1092   No difference in series resistance (Rs) or in slow capacitance between groups.

1093

Supplementary **Figure** **11.**

1095

(A) Example illustration of an image from neuronal TH-immunosignal, transformed as a TH mask and overlayed

on top of Kv4.3 immunosignal. Further segregation of the membrane and cytoplasm compartment with the

corresponding Kv4.3 immunosignal intensity. (B) Distributions of all ROIs area sizes based on TH-masks from

vehicle- (gray) and 6-OHDA-treated mice (red) in the late phase. Inset, cumulative representation of the same

distributions. (C) Scatter dot-plots, illustrating the medio-lateral gradient of Kv4.3 immunosignal on the

contralateral and ipsilateral side from 6-OHDA-treated mice in the late phase. (D) Scatter dot-plots, showing

significant decrease in Kv4.3 immunosignal in both membrane and cytoplasm compartments in the ipsilateral

(6-OHDA-infused) side compared to the contralateral (control) side. Membrane: p = 0.0003, contralateral side

67.2 ± 0.9, ipsilateral side 56.7 ± 1.1 (about 16% reduction); Cytoplasm: p < 0.0001, contralateral side 55.8 ± 0.9,

ipsilateral side 45.9 ± 1.1 (about 18% reduction) (E) Top: 4x magnification of midbrain of a vehicle-infused mouse,

>64 days post-lesion – contralateral side (left panel), and corresponding ipsilateral side (right panel). Middle: 60x

1107 magnification in the highlighted area from 4x image (green, TH; red, Kv4.3). Bottom, left: zoom-in on an example

1108 ROI (highlighted in 60x image). Bottom, right: color-coded Kv4.3-channel immunohistochemical signal intensity

1109 in the example ROI. (F) Histogram showing intensity of Kv4.3 immunosignals for all TH-positive ROIs, from

1110 ipsilateral, vehicle-infused, side (in orange) and from contralateral side (in green). Inset, same data shown as a

1111 cumulative distribution.

1112

1113

Figure 1. Unilateral striatal 6-OHDA mouse model with stable loss of SN DA neurons resulted in delayed partial behavioral recovery.

(A) Experimental design, illustrating timeline of behavioral assays with termination points – groups ended either on the 21st, or later than the 64th post-infusion day. (B) Left panel: TH-DAB staining of the SN, 10x magnification and 60x magnification for vehicle and 6-OHDA-infused mouse. Scalebar left 200μm, scalebar right 25μm. Right panel: ratio of ipsilateral (infusion side) to contralateral side of surviving TH-positive neurons in SN at 21st post-infusion day. (C) Left: spontaneous locomotion of mice in open field arena for a 10 min session. Left, examples of an ACSF-infused mouse (vehicle), right example of a 6-OHDA-infused mouse at baseline (upper panels), 4th post-infusion day (middle panels) and 68th post-infusion day (lower panels). Right: ratio of ipsilateral to contralateral turning behavior for all experimental groups plotted against session days. Note recovery in the 6-OHDA-treated mice from day 4 to day 68 after initial strong turning bias (>90%). (D) Cylinder test quantified by ratio of contra- to ipsilateral forepaw use. Note significant loss of contralateral forepaw involvement, both at 21 and 64 days. All data are presented as mean ± standard error of mean (SEM).

Figure 2. Surviving SN DA neurons at early post-6-OHDA phase exhibited a compressed dynamic range with a 10-fold decrease in *in vivo* bursting and a 5-fold decree in *in vitro* pacemaker regularity

(A), (I) Experimental design, illustrating timeline of behavioral assays, followed by terminal *in vivo* juxtacellular recordings (A) or by terminal *in vitro* whole-cell recordings (I) at the 21st post-infusion day. (B), (C) Top: 10s original recording trace of spontaneous *in vivo* single-unit activity from SN DA neurons in vehicle (B) and 6-OHDA-infused mouse (C). Scalebars: 1s, 0.2mV. Below, left: corresponding ISI histograms. Below, right: corresponding confocal images of juxtacellularly labelled and immunohistochemically identified DA neuron. Note the sparse bursting of the surviving SN DA neuron from the 6-OHDA-infused mouse at early phase. (D), (L) Anatomical mapping of all extracellularly recorded and juxtacellularly labelled DA neurons (D), and of all in vitro recorded and filled DA neurons (L), projected to bregma -3.08 mm. Location of example SN DA neurons in (B), (C), (J), (K) are highlighted. Smaller symbols represent DA neurons that have been recorded and identified, but not included in the group data analysis in (E-H). (E-H) Scatter dot-plots, showing significant decrease of *in vivo* mean firing rate (E), percentage of SFB (G) and burst rate (H) and no significant differences in CV (F) between the vehicle and 6-OHDA-infused mice. Note the 10-fold decrease in SFB for 6-OHDA-infused mice. (J), (K) Top: 10s original recording trace of *in vitro* whole-cell recording of spontaneous SN DA neuron activity in a vehicle (J) and 6-OHDA-infused (K) mouse. Scalebars: 1s, 20mV. Note that the 6-OHDA DA neuron has a highly irregular pacemaking. Below, left: corresponding to hyperpolarizing current injection. Below, right: confocal images of NB filled and immunohistochemically identified DA neuron. (M-P) Scatter dot-plots, showing no difference in *in vitro* mean firing rate (M), rebound delay (O) or sag-component (P) and a 5-fold increase in CV (N). Immunohistochemical imaging for all four DA neurons are displayed in 10x and 60x magnifications (green, TH; red, NB), scalebars: 200μm, 20μm. All data are presented as mean ± standard error of mean (SEM).

Figure 3. Surviving SN DA neurons at late 6-OHDA phase recovered *in vivo* burst firing and doubled their intrinsic pacemaker frequency *in vitro*.

(A), (I) Experimental design, illustrating timeline of behavioral assays, followed by terminal *in vivo* juxtacellular recordings (A) or by terminal *in vitro* whole-cell recordings (I) after >64 post-infusion days. (B), (C) Top: 10s original recording trace of spontaneous *in vivo* single-unit activity from SN DA neurons in vehicle (B) and 6-OHDA-infused mouse (C). Scalebars: 1s, 0.2mV. Below, left: corresponding ISI histograms. Below, right: corresponding confocal images of juxtacellularly labelled and immunohistochemically identified DA neuron. (D), (L) Anatomical mapping of all extracellularly recorded and juxtacellularly labelled DA neurons (D), and of all *in vitro* recorded and filled DA neurons (L), projected to bregma -3.40 mm (D) and -3.08mm (L). Location of example SN DA neurons in (B), (C), (J), (K) are highlighted. Smaller symbols represent DA neurons that have been recorded and identified, but not included in the group data analysis in (E-H). (E-H) Scatter dot-plots, showing no significant difference in mean firing rate (E), CV (F), SFB (G) and burst rate (H). (J), (K) Top: 10s original recording trace of *in vitro* whole-cell recording of spontaneous activity from SN DA neurons in a vehicle (J) and 6-OHDA-infused (K) mouse. Scalebars: 1s, 20mV. Note that the 6-OHDA DA neuron has enhanced, but regular pacemaking. Below, left, corresponding to hyperpolarizing current injection. Below, right, confocal images of NB filled and immunohistochemically identified DA neuron. (M-P) Scatter dot-plots, showing doubling of the firing rate (M), no difference in CV (N), decrease in rebound delay (O) and increase of sag-component (P) for the 6-OHDA-treated mice in comparison to vehicle group. Immunohistochemical imaging for all four DA neurons are displayed in 10x and 60x magnifications (green, TH; red, NB), scalebars: 200μm, 20μm. All data are presented as mean ± standard error of mean (SEM).

Figure 4. Enhanced pacemaker frequency of SN DA neurons at the late post-6-OHDA phase was mediated by K$_v$4.3 channel downregulation

(A), (I) Experimental design, illustrating timeline of behavioral assays, followed by terminal *in vitro* whole-cell recordings under K$_v$4.3 channel blocker AmmTx3 (A) or *in vitro* voltage-clamp whole-cell recordings (I) after >64 post-infusion days. (B), (C) Top: 10s original recording traces of *in vitro* whole-cell recording of spontaneous activity from DA SN neurons in AmmTx3-preincubated slices for a vehicle (B) and 6-OHDA-infused (C) mouse. Below, left: corresponding hyperpolarizing current injection. Scalebars: 1s, 20mV. Below, right: confocal images of NB filled and immunohistochemically identified DA neuron. (D), (L) Anatomical mapping of all in vitro recorded and filled DA neurons, projected to bregma -2.92 mm (D) and -3.16mm (L). (E-H) Scatter dot-plots, showing no differences in mean firing rate (E), coefficient of variation (F), rebound delay duration (G) and sag amplitude (H). (J), (K) Original recording trace of in vitro voltage-clamp recordings from DA SN neurons in a vehicle (J) and 6-OHDA-infused (K) mouse. Top, right: Zoom in from a K$_v$4.3 channel activation. Top, left: Zoom in from a K$_v$4.3 channel inactivation. Below, right: Zoom in from HCN-channel activation. Below, right, confocal images of NB filled and immunohistochemically identified DA neuron. Scalebars first row: 3nA, 100ms. Scalebars below, left: 5nA, 500ms. Note the small K$_v$4.3 channel in/activation peak in a surviving DA neuron from a 6-OHDA mouse in comparison the one from an ACSF treated mouse. Immunohistochemical imaging for all four DA neurons are displayed in 10x and 60x magnifications (green, TH; red, NB), scalebars: 200μm, 20μm. All data are presented as mean ± standard error of mean (SEM). (M) Scatter dot-plots, showing a significant, half of maximum K$_v$4.3 channel conductance in surviving DA neurons. (N) Scatter dot-plots, showing no difference in HCN channel delta current (I$_{peak}$ - I$_{steady}$) between groups. (O) Normalized K$_v$4.3 current (I$_{peak}$ - I$_{steady}$) at different voltage steps, resulting in inactivation and activation curve for both groups. All data are presented as mean ± standard error of mean (SEM).

Figure 5. Surviving DA SN neurons at the late post-6-OHDA phase showed lower K$_v$4.3 channel immunohistochemical signal

(A) Experimental design, illustrating timeline of behavioral assays, followed by multi-labeling immunohistochemistry for exploring K$_v$4.3-channel expression differences after >64 post-infusion days. (B), (C) Top: 4x magnification of midbrain of a 6-OHDA-infused mouse, >64 days post-lesion – contralateral side (B), and corresponding ipsilateral side (C). Middle: 60x magnification in the highlighted area from 4x image (green, TH; red, K$_v$4.3). Bottom, left: zoom-in on an example ROI (highlighted in 60x image). Bottom, right: color-coded Kv4.3-channel immunohistochemical signal intensity in the example ROI. Note K$_v$4.3-channel signal decrease in surviving DA neurons on the ipsilateral to the injection side. (D) Histogram showing intensity of K$_v$4.3 immunosignals for all TH-positive ROIs, from ipsilateral, lesioned, side (in red) and from contralateral side (in grey). Inset, same data shown as a cumulative distribution. Note a clear right-shift to lower intensities for the ipsilateral side. (E) Comparison of mean TH-colocalized K$_v$4.3 immunosignals from medial SN from ipsilateral, lesioned, side, and contralateral, as control side. All data are presented as mean ± standard error of mean (SEM).

Supplementary Figure 1.

(A), (B) Comparison of ipsilateral, infusion side, to contralateral side as percentage of relative TH immunohistochemistry signal in the dorsal striatum (dSTR) and ventral striatum (vSTR), at the corresponding early (A) and late phase (B). Note the stability of TH density loss in the dorsal striatum through time. (C) Ratio of ipsilateral (infusion side) to contralateral side of surviving TH-positive neurons in SN in the late phase. (D) Mean track length from all mice for each open field session. Note the post-infusion drop in performed track in the 6-OHDA groups, which gradually recovers. (Infusion day marked as a thin gray line.) (E) Ratio of TH+ cell loss in SNc (ipsilateral (infusion side) to contralateral side) >21st post-infusion day from different infused 6-OHDA doses.

Supplementary Figure 2.

(A) Ratio of ipsilateral to contralateral turning behavior for female ACSF and 6-OHDA infused groups plotted against session days and with underlayed the corresponding ratio from the male ACSF and 6-OHDA groups as a mean-line. (B) Ratio of ipsilateral (infusion side) to contralateral side of surviving TH-positive neurons in SN at >64st post-infusion day from female ACSF and 6-OHDA infused mice. (C) Comparison of ipsilateral, infusion side, to contralateral side as percentage of relative TH immunohistochemistry signal in the dorsal striatum (dSTR) at >64st post-infusion day from female ACSF and 6-OHDA infused mice.

Supplementary Figure 3.

For two representative vehicle- and 6-OHDA-treated mice schematic representation of turning sequences. Left turn sequences are plotted as negative values, whereas right turn sequences are plotted as positive values. The count represents the number of same-direction turns within a sequence. On the left panels are turning sequences for the control animal and on the right panels for the 6-OHDA-infused animal, correspondingly in a baseline session (A), 4th post-operative day (B), 20th post-operative day (C) and 64th post-operative day (D).

Supplementary Figure 4.

Correlation pair plots comparing turning features, such as velocity (cm/s), diameter (cm) and duration for all contralateral turns (left panels) and all ipsilateral turns performed from all vehicle-infused mice (in orange) and all 6-OHDA-treated mice (in blue) across different time points (baseline, 4th, 20th and 64th post-infusion day).

Supplementary Figure 5.

Scatter dot-plots, showing no significant difference of in vivo mean intra-burst (IB) frequency (A), mean maximum (max) intra-burst (IB) frequency (B), burst duration (C), number of spikes per burst (D), single spike frequency (SSF) (E), single spike coefficient of variance (CV) (F) and action potential (AP) width (G) (H) between the vehicle and 6-OHDA-infused mice in the early phase. (H) Normalized stacked bar plots of different in vivo firing patterns based on ACH. (I) ISI distributions from all in vivo recorded and labeled mSN DA neurons from vehicle- and 6-OHDA-treated mice in the early phase. Inset, cumulative representation of the same distributions.

Supplementary Figure 6.

Scatter dot-plots, showing no significant difference of in vitro afterhyperpolarization (AHP) minimum (A), threshold (B), spike amplitude (C), action potential (AP) width (D) and input resistance (E) between the vehicle and 6-OHDA-infused mice in the early phase. (F), (G) Paired scatter dot-plots of firing frequency (F) and coefficient of variance (G) during on-cell recording and whole-cell recording in the early phase. (H) ISI distributions from all in vitro whole-cell recorded and labeled mSN DA neurons from vehicle- and 6-OHDA-treated mice in the early phase. Inset, cumulative representation of the same distributions. (I) Paired scatter dot-plot indicating no significant difference between the mean voltage during spike-pauses and the rest of the firing activity.

Supplementary Figure 7.

Scatter dot-plots, showing no significant difference of in vivo mean intra-burst (IB) frequency (A), mean maximum (max) intra-burst (IB) frequency (B), burst duration (C), number of spikes per burst (D), single spike frequency (SSF) (E), single spike coefficient of variance (CV) (F) and action potential (AP) width (G) (H) between the vehicle and 6-OHDA-infused mice in the late phase. (H) Normalized stacked bar plots of different in vivo firing patterns based on ACH. (I) ISI distributions from all in vivo recorded and labeled mSN DA neurons from vehicle- and 6-OHDA-treated mice in the late phase. Inset, cumulative representation of the same distributions.

Supplementary Figure 8.

Scatter dot-plots, showing no significant difference of in vitro afterhyperpolarization (AHP) minimum (A), threshold (B), spike amplitude (C), action potential (AP) width (D) and input resistance (E) between the vehicle and 6-OHDA-infused mice in the late phase. (F), (G) Paired scatter dot-plots of firing frequency (F) and coefficient of variance (G) during on-cell recording and whole-cell recording in the late phase. (H) ISI distributions from all in vitro whole-cell recorded and labeled mSN DA neurons from vehicle- and 6-OHDA-treated mice in the late phase. Inset, cumulative representation of the same distributions.

Supplementary Figure 9.

Scatter dot-plots, showing minor significant difference of in vitro afterhyperpolarization (AHP) minimum (A), and no difference in threshold (B), spike amplitude (C), action potential (AP) width (D) and input resistance (E) between the vehicle and 6-OHDA-infused mice in the late phase under 1μM AmmTx3. (F), (G) Paired scatter dot-plots of firing frequency (F) and coefficient of variance (G) during on-cell recording and whole-cell recording in the late phase under 1μM AmmTx3. (H) ISI distributions from all in vitro whole-cell recorded and labeled mSN DA neurons from vehicle- and 6-OHDA-treated mice in the late phase under 1μM AmmTx3. Inset, cumulative representation of the same distributions.

Supplementary Figure 10.

(A)-(D) Measurements of transient outward (A-type) potassium biophysical parameters in whole-cell voltage-clamp recordings of Kv4.3 channel activation and inactivation in vehicle relative to 6-OHDA infused mice. (E), (F) No difference in series resistance (Rs) or in slow capacitance between groups.

Supplementary Figure 11.

(A) Example illustration of an image from neuronal TH-immunosignal, transformed as a TH mask and overlayed on top of Kv4.3 immunosignal. Further segregation of the membrane and cytoplasm compartment with the corresponding Kv4.3 immunosignal intensity. (B) Distributions of all ROIs area sizes based on TH-masks from vehicle- (gray) and 6-OHDA-treated mice (red) in the late phase. Inset, cumulative representation of the same distributions. (C) Scatter dot-plots, illustrating the medio-lateral gradient of Kv4.3 immunosignal on the contralateral and ipsilateral side from 6-OHDA-treated mice in the late phase. (D) Scatter dot-plots, showing significant decrease in Kv4.3 immunosignal in both membrane and cytoplasm compartments in the ipsilateral (6-OHDA-infused) side compared to the contralateral (control) side. Membrane: p = 0.0003, contralateral side 67.2 ± 0.9, ipsilateral side 56.7 ± 1.1 (about 16% reduction); Cytoplasm: p < 0.0001, contralateral side 55.8 ± 0.9, ipsilateral side 45.9 ± 1.1 (about 18% reduction) (E) Top: 4x magnification of midbrain of a vehicle-infused mouse, >64 days post-lesion – contralateral side (left panel), and corresponding ipsilateral side (right panel). Middle: 60x magnification in the highlighted area from 4x image (green, TH; red, Kv4.3). Bottom, left: zoom-in on an example ROI (highlighted in 60x image). Bottom, right: color-coded Kv4.3-channel immunohistochemical signal intensity in the example ROI. (F) Histogram showing intensity of Kv4.3 immunosignals for all TH-positive ROIs, from ipsilateral, vehicle-infused, side (in orange) and from contralateral side (in green). Inset, same data shown as a cumulative distribution.