Patient-derived triple negative breast cancer organoids provide robust model systems that recapitulate tumor intrinsic characteristics

Sonam Bhatia[1], Melissa Kramer[1#], Suzanne Russo[1#], Payal Naik[1#], Gayatri Arun[1$], Kyle Brophy[1], Peter Andrews[1], Cheng Fan[2], Charles M. Perou[2], Jonathan Preall[1], Taehoon Ha[1], Dennis Plenker[1$$], David A. Tuveson[1], Arvind Rishi[3], J Erby Wilkinson[4], W. Richard McCombie[1], Karen Kostroff[5] and David L. Spector[1*]

[1]Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 11724
[2]University of North Carolina, Lineberger Comprehensive Cancer Center, Department of Genetics, Chapel Hill, N.C. 27599
[3]Northwell Health, Department of Pathology, Lake Success, NY 11042
[4]University of Michigan, Department of Pathology, Ann Arbor, MI 48109, USA
[5]Northwell Health, Department of Surgical Oncology, Lake Success, NY 11042

[#]Contributed equally
[$]Currently at: Envisagenics Inc.
[$$]Currently at: Loxo Oncology at Eli Lilly
[*]Corresponding author: spector@cshl.edu

Conflict of Interest

C.M.P is an equity stock holder and consultant of BioClassifier LLC; C.M.P is also listed as an inventor on patent applications for the Breast PAM50 Subtyping assay.

Abstract

Triple negative breast cancer (TNBC) is an aggressive form of breast cancer with poor patient outcomes, and an unmet clinical need for targeted therapies and better model systems. Here, we developed and comprehensively characterized a diverse biobank of normal and breast cancer patient-derived organoids (PDOs) with a focus on TNBCs. PDOs recapitulated patient tumor intrinsic properties and a subset of PDOs can be propagated for long-term culture (LT-TNBCs). Single cell profiling of PDOs identified cell types and gene candidates affiliated with different aspects of cancer progression. The LT-TNBC organoids exhibit signatures of aggressive MYC-driven basal-like breast cancers and are largely comprised of luminal progenitor (LP)-like cells. The TNBC LP-like cells are distinct from normal LPs and exhibit hyperactivation of NOTCH and MYC signaling. Overall, our study validates TNBC PDOs as robust models for understanding breast cancer biology and progression, paving the way for personalized medicine and tailored treatment options.

Statement of Significance

A comprehensive analysis of TNBC patient-derived organoids is presented by genomic, transcriptomic, and *in-vivo* analyses, providing insights into cellular heterogeneity and mechanisms of tumorigenesis at the single cell level.

Introduction

1   Breast cancer is characterized into several histopathological subtypes based on the expression of various receptors: estrogen (ER), progesterone (PR) and human epidermal growth factor receptor 2 (HER2/ERBB2). Molecular subtyping revealed multiple subgroups based on gene expression patterns that have been refined overtime and constitute luminal A, luminal B, HER2-enriched, basal-like, and normal-like breast cancers (1–3). Luminal A (ER+/PR+/Her2-) is the most prevalent and constitutes ~70% of all breast cancer cases, luminal B (ER+/PR+/HER2+) make up ~12% of cases, HER2-amplified (ER-/PR-/HER2+) represents ~5% of total cases and the basal or triple negative breast cancer (TNBC, ER-/PR-/HER2-) makes up ~12% of total breast cancer cases (4). The majority of luminal breast cancers exhibit low proliferation rates while the TNBCs are highly aggressive and proliferative in nature. Since TNBCs lack ER/PR and HER2 receptors, patients are exempt from any targeted endocrine or HER2 therapies resulting in non-specific cytotoxic chemotherapy as the standard of care. The majority, but not all, TNBCs fall under the basal-like subgroup of breast cancer by gene expression based PAM50 profiling and a small subset belong to the highly invasive claudin-low subgroup (1,4–6). Further molecular characterization of TNBCs revealed heterogeneity within this subset and at least 6 molecular subtypes have been identified that can be used to design more targeted therapies (7,8). TNBCs are typically presented as high-grade carcinomas (9) and show a significantly poor 5-year prognosis compared to other subtypes (10–12). Therefore, there is an unmet clinical need to better understand triple-negative disease progression and identify more precise druggable targets, and specific therapies for this cohort of breast cancer patients.

3-D organoid models have gained traction over recent years as the next generation

of models for studying disease and development (reviewed in (13,14)). While cell lines,

spheroids and mouse model-derived organoids have been the primary *in vitro* model

systems for studying cancer biology; patient-derived organoid (PDO) models have now

been developed for a variety of cancers including those originating in the colon (15),

pancreas (16), ovary (17,18), prostate (19) and breast (20). Sachs et al. developed a

methodology for growing human breast tumors *ex vivo* as organoid models that has now

been expanded to the growth of organoids from normal mammary tissue as well (21,22).

While cancer cell lines have been a valuable resource for providing insights into cancer

biology and drug development (23–25), PDO models are not only patient-specific but also

provide a three-dimensional context which is closer to that of the actual tumor

microenvironment. PDOs, therefore, represent unique model systems to study disease

progression and for the identification and validation of better treatment options. However,

we do not yet fully understand the extent to which these models recapitulate the cellular

heterogeneity and complexities of triple negative disease.

Here, we developed a diverse breast cancer PDO biobank and performed

comprehensive genomic, transcriptomic and cellular characterization of organoids with

an emphasis on TNBCs. Using genomic assays we show that our organoid models

recapitulate pathogenic single nucleotide variants (SNVs) and copy number alterations

(CNAs) of breast cancers as portrayed in large scale breast cancer genomic datasets

(10,26–28) and reveal lesser studied cancer driver genes. Transcriptomically, our biobank

recapitulates the various subtypes and signatures of breast cancers, with a subset of

organoids exhibiting signature profiles associated with poor patient outcomes. *In-vivo*

transplants of these organoids highly recapitulated the patient-tumor morphology,

providing strong evidence of retention of individual tumor intrinsic properties in long-term

organoid cultures, even for models derived from rare BCs. In addition, we find that while

normal PDOs retain the major cell types found within the mammary epithelium, the TNBC

PDOs have lost this lineage specificity and are predominantly enriched for luminal

progenitor (LP)-like cells. Single cell RNA-sequencing (scRNA-seq) of TNBC and normal

PDOs identified differential signatures between the tumor and normal LP cells, providing

insights into putative mechanisms of tumorigenesis. Lastly, we identified cells with various

gene expression signatures in TNBC organoids that can be used to model and perturb

various aspects of cancer biology, including tumorigenesis, hypoxia response, and EMT.

Overall, our comprehensive characterization of TNBC organoids identified them as valid

cancer models for studying cancer biology and for applications in precision medicine.

Materials and Methods:

Patient Material

Tumor resections from breast cancer patients along with the distal and adjacent normal

tissue were obtained from Northwell Health in accordance with Institutional Review Board

protocol IRB-03-012 and IRB 20-0150. Specific information for all samples is available in

Table S1. The collection of genomic and phenotypic data was consistent with 45 CFR

Part 46 (Protection of Human Subjects) and the NIH Genomic Data Sharing (GDS)

Policy. Informed consent ensured that the de-identified materials collected, the models

created, and data generated from them can be shared without exceptions with

researchers in the scientific community.

Patient-derived organoid culture: Patient-derived organoids were established and

propagated using a previously published protocol (20). In summary, the tissues were

manually cut into smaller pieces and treated with 2mg/ml collagenase IV in base media

(ADF+++: Advanced DMEM-F12 (Invitrogen 12634-034) with 1x Glutamax (Invitrogen

12634-034), 10mM Hepes (Invitrogen 15630-056), 100U/ml Pen-Strep (Invitrogen 15140-

122)) at 37°C for 45-90mins with gentle agitation to break the tissue into small clusters of

cells. The suspension was intermittently resuspended by pipetting multiple times to

ensure proper digestion of the tissue. The cell suspension was centrifuged at 300g for

5mins and the pellet was treated with red blood cell lysis buffer (Cat # 11814389001,

Sigma) for 5mins at room temperature if it appeared bloody. The suspension was washed

2x with ADF+++ and plated in a matrigel (lot test for concentration of 8-10mg/ml, Cat #

356231, Corning) dome on pre-warmed tissue culture plates. The dome was incubated

at 37°C for 15mins and supplemented with completed medium: 10% R-Spondin1

conditioned medium, 5nM Neuregulin 1 (Peprotech 100-03), 5ng/ml FGF7 (Peprotech

100-19), 20ng/ml FGF10 (Peprotech 100-26), 5ng/ml EGF (Peprotech AF-100-15),

100ng/ml Noggin (Peprotech 120-10C), 500nM A83-01 (Tocris 2939), 5uM Y-27632

(Abmole Y-27632), 1.2uM SB202190 (Sigma S7067), 1x B27 supplement (Gibco 17504-

44), 1.25mM N-Acetylcysteine (Sigma A9165), 5mM Nicotinamide (Sigma N0636),

50ug/ml Primocin (Invitrogen ant-pm-1) in ADF+++. The organoids were passaged every

15-30 days using TrypLE™ (Thermo Fischer 12605028) to break down the organoids into

smaller clusters of cells and re-plating them in matrigel domes as described above. For

tumor scrapings, the tumor surface was shaved on multiple sides to collect material which

was subsequently manually broken down, treated with red blood cell lysis buffer and

seeded in matrigel followed by regular PDO culture.

Organoid models labelled with the prefix HCM-CSHL were acquired as part of the

Human Cancer Model Initiative (HCMI) https://ocg.cancer.gov/programs/HCMI and a

subset of those models are or will be available for access from ATCC. The data for these

models can be accessed here: dbGaP accession number phs001486. Organoid

nomenclature: prefixes LNS, NH, DS, HCM-CSHL are de-identified patient IDs and are

not distinct in any features other than protocols used for sample acquisition; prefix NM

designates true normal samples collected from patients undergoing reductive

mammoplasty (NM: Normal Mammoplasty); suffixes: T=tumor, N=normal, ND=normal

distal, NAdj=normal adjacent, and TSc= tumor scraping. Organoid freeze thaws are

indicated in parenthesis: (passage frozen down)passage after thaw eg. NH85TSc (p4)p4.

Organoid DNA and RNA extraction: Organoid RNA was extracted using TRIzol®

(Thermo Fischer 15596018) RNA extraction protocol. DNA was extracted by removing

matrigel from organoids using ice cold PBS or TrypLE following by DNA extraction using

Qiagen DNeasy Blood and Tissue kit (Qiagen 69504) with elution in nuclease free water

(Thermo Fischer/Ambion 4387936).

Targeted gene panel sequencing: We performed capture based targeted gene panel

sequencing (65) for a panel of potential cancer driver genes. Briefly, we used a panel of

143 cancer genes with a total of ~4000 probes for capture. The captured DNA was paired-

end sequenced with 150bp reads and a coverage of about 300-500x. Library preparation

and sequencing of the targeted gene panel was performed by the CSHL Next Generation

Sequencing Core Facility. We developed an analysis pipeline to prioritize identification of

driver mutations. The sequencing reads are aligned to the hg19 reference genome using

BWA (66), followed by conversion to BAM format and sorting with Samtools (67), removal

of PCR duplicates with Picard (https://broadinstitute.github.io/picard/), and filtering with

Bamtools (68) for mapping quality and proper read pairing. Coverage of the target regions

is assessed for breadth and depth using Picard HSmetrics to ensure adequate coverage

for confident variant detection. Variants are then called using VarScan2 (69) in somatic

mode to stratify germline versus somatic variants, and are annotated with Annovar (70)

to cover a broad range of variant assessment tools. We then select rare loss of function

variants (nonsense, frameshift, splice site) with frequency less than 1% in the Gnomad,

ExAC, EVS and 1000 Genomes databases. Missense and in-frame indel variants are

selected if they are noted as pathogenic by ClinVar (71), or if they are both rare (<1% in

all genome databases) and annotated as pathogenic by COSMIC (72), or if they are both

rare and found to be present in the TCGA cohorts. Finally, missense variants are selected

if they are annotated as potentially deleterious by the ensemble tools REVEL (73) and

MCAP (74). Variants that were deleterious by REVEL and MCAP but did not have

population level data were discarded from the final oncoplot. Oncoplots are generated

from these candidate variants using Maftools (75).

RNA-seq: All RNA samples were quality controlled using a nanodrop followed by a

bioanalyzer (RNA nano-kit Cat # 5067-1511) and only samples with RIN >7 were used

for RNA sequencing. 750ng of RNA was used to prepared un-stranded RNA-seq libraries

using Illumina TruSeq RNA Library prep kit v2 (RS-122-2001) and sequenced as 75bp

paired-end reads.

RNA-seq analysis: The sequencing fastq files were quality checked using fastQC to

make sure the reads were of consistent quality between different runs. The reads were

aligned using STAR-aligner STAR-2.5.3a (76) using the following parameters:

```
147  --outFilterMismatchNmax 8 --alignIntronMax 1000000 --alignMatesGapMax
148  1000000      --outSAMstrandField   intronMotif   --outFilterIntronMotifs
149  RemoveNoncanonicalUnannotated
```
150 against the gencode v27 gtf reference file. Any PCR duplicates were marked in the

151 aligned files using STAR with the following parameters:

```
152  --runMode        inputAlignmentsFromBAM        --bamRemoveDuplicatesType
153  UniqueIdenticalNotMulti --runThreadN $thread --inputBAMfile $input_bam
154  --outSAMtype BAM SortedByCoordinate
155
```
followed by removing duplicated reads using `samtools view -bF 0x400`. HTseq-count

was used to count the reads per gene using the gencode v27 gtf file. The counts files

were exported into R 4.1.0 and analyzed for differential expression using DeSeq2 1.32.0

(77). Concordance between technical replicates was ensured using PCA and sample

distance matrix before summing them together for downstream analysis. Sample distance

matrix was generated using euclidean distances between the samples and hierarchical

clustering was performed using "ward.D2" linkage method followed by cutree with

kmeans=6 with the R package "stats" (v4.1.0). For signature correlation, all the samples

were used and an unsupervised clustering was performed for 838 previously curated

gene expression signatures (37–39) and visualized using Java TreeView v1.2.0. For

individual signature comparisons signature scores in experimental groups were

compared using Kruskal-Wallis test followed by pairwise comparisons using Wilcoxon

rank-sum test. Family-wise error rate was adjusted using Bonferroni-Holm method.

SMASH copy number: SMASH was performed as per the published protocol (32)

starting with 750ng of genomic DNA. SMASH was performed in batches of 10 samples

and sequenced on a MiSeq PE150bp run. The SMASH analysis is based on identification

of Maximal Unique Matches (MUMs) to the human genome in all read pairs (32). These

MUMs were filtered to remove matches <20 bp, matches with <4 bp of excess unique

sequence, and matches on read 2 that are within 1000 bases of the matches from read

1. Raw copy number profiles are then generated from the remaining 3–4 matches per

read pair which are then added to empirically sized bins spanning the genome. Regions

with identical copy are expected to yield similar bin counts using these empirical bins. We

next perform GC correction by normalizing counts based on LOWESS smoothing of count

vs. GC content data in each bin. Final copy number profiles are normalized so that the

autosome has an average copy number of 2. Plots were generated with G-Graph

MUMdex software (https://mumdex.com/) and IGV browser v2.9.2 (78).

Organoid Histology: Organoid domes in complete medium were scraped from the tissue

culture plate and collected in falcon tube precoated with BSA. The organoids were

collected and washed 1x with PBS by spinning at 300g for 5mins. Organoid harvesting

solution (Cat # 3700-100-01, Trevigen) was added to the organoids (3x the volume of

matrigel) and incubated at 4°C on ice for 30minutes to ensure that matrigel was removed

and the organoids were concentrated at the bottom. The organoids were washed 1x with

ample PBS and fixed with fresh 4% PFA at room temperature for 10minutes. 1:1 (v/v)

BSA was added to the tube and spun at 300g for 5mins to remove the PFA. The organoids

were washed 2X with ample amounts of PBS and embedded in 2% agarose in dH2O).

The agarose organoid molds were then paraffin embedded and cut into 5um sections.

Organoid Hematoxylin & Eosin (H&E) staining and Immunohistochemistry (IHC):

H&E and IHC staining were performed at the CSHL Histology Core Facility. PFA fixed

organoids in agarose were processed in Thermo Excelsior ES processor and embedded

with Thermo HistoStar embedding system following manufacturer's protocols. Paraffin

blocks were cut into 5um sections and mounted onto positively charged slides (VWR

superfrost plus micro slide).

For H&E staining, slides were stained in a Leica Multistainer (ST5020). Briefly,

slides were deparaffinized and rehydrated and then stained in hematoxylin (Hematoxylin

560 MX, Leica) for 1 min, followed by destaining in Define MX-aq (Leica) for 30 sec, bluing

in Blue Buffer 8 (Leica) for 1min and subsequently stained in eosin (EOSIN 515 LT, Leica)

for 30sec. After dehydration, coverslips were placed onto glass slides using a robotic

coversliper (Leica CV5030).

IHC slides were stained in Discovery Ultra automatic IHC stainer (Roche) following

standard protocols. Briefly, slides were subjected to antigen retrieval (Benchmark Ultra

CC1, Roche) at 96˚C for 1hr; primary antibodies were incubated at 37˚C for 1hr and

Discovery multimer detection system (Discovery OmniMap HRP, Discovery DAB and

Purple, Roche) was used to detect and amplify immuno-signals. Antibodies used: Ki67

(Spring Bioscience, #M3062, 1:500).

Organoid formation assay: Organoids were processed using TrypLE and 1500 single

cells per well of a 96 well plate were seeded in 10% matrigel + complete growth medium.

Cell viability was assessed using Cell Titre Glo 3D luminescence assay (Promega G9683

CellTiter-Glo 3D Cell Viability Assay). Baseline cells were measured using Cell Titre Glo

3D assay at 24 hrs (d1) post seeding and growth was measured at 6 days after seeding

(d6). Each organoid line was evaluated for multiple passages n=2 or n=3 per PDO.

For MYCi and DAPT experiments: Organoids were processed to single cells using

TrypLE and seeded in 50ul matrigel domes as 10,000 cells per well of a 24 well-plate.

Complete medium or medium supplemented with DMSO, DAPT (Selleckchem S2215

DAPT) and MYC-inhibitor (Selleckchem S8906 MYCi975) were added to the respective

wells. For normal PDOs, organoids were dissociated with trypLE and live cells were

sorted using fluorescence activated cell sorting (FACS) based on staining of CD46f,

EPCAM and 7AAD. Organoids were allowed to form for 12 days, images were acquired

using microscope and organoids were manually counted for each condition. Statistical

analysis was performed using one-way ANOVA test followed by pairwise comparisons

using two-sample t-test. Family-wise error rate was adjusted using Bonferroni-Hold

method.

Organoid proliferation index analysis: Paraffin embedded organoid sections were IHC

stained for Ki67. Slides were scanned and viewed using Aperio ImageScope 12.3.3.

Images were analyzed in FIJI (79). Briefly, the images were deconvolved using Colour

Deconvolution for hematoxylin and DAB, converted to 8-bit binary images and analyzed

using the BioVoxxel Toolbox plugin (https://www.biovoxxel.de/#/) to evalute %Ki67

positive cells per organoid. Multiple passages for each organoid line were evaluated, n=2

or n=3 per PDO.

Drug dose response assays: Organoids were digested into single cells and seeded as

1500 cells/well of a 384 wp (USA-Scientific Cat # 5678-1976) as suspension cultures in

complete medium with 10% matrigel using a liquid handler. Organoids were incubated at

37C for 24hrs and drugs were added using a drug dispenser Beckman Echo 650. The

organoids were incubated for 5 days at 37C and assayed using the CellTiter-Glo® 3D

Cell Viability Assay (Promega Cat # G9681) and read for luminescence using the

EnVision 2105 plate reader. The data was analyzed using GraphPad Prism v9.0.0. The

screens were performed in 3 technical replicates with 2-4 experimental replicates per

organoid line. The raw reads were first normalized against untreated controls and a non-

linear model was fit for the mean, after removing any detected outliers, using the

log(inhibitor) vs normalized response with a variable slope.

Animals: Six-week-old female NOD scid mice (NOD.Cg-*Prkdc*scid/J) were obtained from

the Jackson laboratory (JAX stock #001303) and acclimated at the Cold Spring Harbor

Laboratory Animal Shared Resource for a minimum of 1week. All animal experiments

were performed in accordance with the Institutional Animal Care and Use Committee.

***In-vivo* transplant experiments**: TNBC organoids were harvested using the organoid

harvesting solution (Trevigen Cat# 3700-100-01) and manually counted. Organoids were

resuspended in a 1:1 mixture of PBS:matrigel and 50K organoids were injected into the

bilateral mammary fat pads by the fourth nipple of female NOD scid mice (NOD.Cg-

Prkdc[scid]/J). Mice were anesthetized with 1.5-2% isoflurane and weighed before the

injections. The animals were regularly monitored for their weight, tumor size and any other

discrepancies. Mice were sacrificed when the tumors reached any of the following end-

points: 2cm tumors, ulceration, visible necrosis, blistering of tumors or deteriorating health

of the mice. At end-point, dissections were performed and the tumors along with lungs,

liver, lymph nodes and the femur were fixed in 4% PFA overnight at 4°C. If the tumors

were not observed the mammary fat pads were collected instead. The transplant

experiments were done with 2 independent passages of PDOs, with 4-6 injections per

PDO per passage.

Fixed tumors and tissues were processed for histology as above. Metastases and

micro-metastases were assessed using IHC with a human mitochondria antibody

(Millipore MAB1273 Anti-Mitochondria clone 113-1).

Flow Cytometry: Organoids were scraped in the culture medium and washed 1X with

PBS. TrypLE™ was used to fully digest the organoids into single cells. The cells were

counted, diluted to 200,000 cells/100ul and stained in 100ul of ADF+++ using anti-Epcam

(1:50), anti-CD49f (1:50), 7AAD (1:50). The following antibodies were used: PE Mouse

IgG2a, κ Isotype Control (BD 555574), APC Mouse IgG2b κ Isotype Control RUO (BD

555745), Alexa Fluor® 647 anti-human CD326 (EpCAM) Antibody (Biolegend 324212),

PE anti-human/mouse CD49f Antibody (Biolegend 313612). The cells were read using a

BD Fortessa and analyzed using the FACS DIVA and FlowJo v10 software.

Single cell RNA-seq: Organoids were digested into single cells using TrypLE™,

resuspended in 0.04% BSA in PBS as 1 million cells/ml. ~12,000 cells were loaded into

one well of a 10x Chromium microfluidics chip. Single cell barcoding and libraries were

prepared using the 10x Chromium v3 chemistry (Cat # V3 reagents 1000075, V3 chips -

1000153 or NextGEM reagents- 1000121, chips 1000120). Libraries were quality

checked using a Bioanalyzer HS kit for cDNA yield and final library size and qubit to

quantify.

Single cell analysis was performed in three different batches (Table S7). Batch 3

was a multiplexed pool of 4 samples, which were demultipexed using a custom genotype-

aware pipeline.　At the time of 10X Genomics library preparation, ~20,000 cells from each

of the four organoids were set aside to prepare low-input bulk RNA-seq libraries tagged

with unique i7 barcodes. These bulk libraries share the same adapter structure as 10X

Genomics libraries, and were spiked into the Illumina NextSeq500 flow cell at a 5% molar

ratio to obtain ~5M reads per organoid.　These barcoded bulk libraries were then used to

create reference VCF files using cellSNP v0.3.2 by searching a list of 7.4M common

human　　　　SNPs　　　　from　　　　the　　　　1000　　　　Genomes　　　　Project

(http://ufpr.dl.sourceforge.net/project/cellsnp/SNPlist/genome1K.phase3.SNP_AF5e2.ch

r1toX.hg38.vcf.gz).　Genotype profiles were filtered to include only positions with < 10%

minor allele frequency and >20 UMI counts.　In parallel, per-cell VCF files were generated

from the multiplexed single cell library using the cellSNP with the same parameters.　Cells

from the single-cell pool were assigned to their respective donors using Vireo v0.4.2 (80).

Genotyping low-input bulk RNAseq library prep:

RT Primer Design:

CTACACGACGCTCTTCCGATCTSSSSSSSSNNNNNNNNNNVVVVVTTTTTTTTT

TTTTTTTTTTTTTTTTTTTTTTVN

where:

SSSSSSSS = 8bp sample barcode, with following multi-plexing key:

GACAGTGC=HCM-CSHL-0366-C50

GAGTTAGC=NH85TSc

GATGAATC=NH95T

GCCAAGAC=NH93T

NNNNNNNNNNVVVVV = 15bp UMI with 5 non-T residues at 3' end

Template Switch Oligo: AAGCAGTGGTATCAACGCAGAGTGAATrGrGrG

cDNA_amplification_Forward:

AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCG

cDNA_amplification_Reverse: AAGCAGTGGTATCAACGCAGAGT

RT was performed using the SuperScript IV (Life Technologies #18091050) according to

the manufacturer's instructions except for the addition of 1uM Template Switch Oligo

during first strand synthesis. All custom oligos were synthesized by IDT. After cDNA

amplification, molar concentration was estimated using the Agilent Bioanalyzer 2100,

libraries were pooled at an equimolar ratio, and prepared for Illumina sequencing using

the Nextera XT DNA Library Prep Kit (Illumina) according to the manufacturer's

instructions. Final fragmented libraries were again checked and quantified by Bioanalyzer

prior to mixing at a 1:20 molar ratio with 10X Genomics libraries for sequencing.

Sequencing and Mapping: The libraries were sequenced on Illumina NextSeq 500 High

Output 75 cycle kits using the read format: 8bp (i7 index) x 28bp (Read 1) x 56bp

(Read2). 10X Genomics libraries were mapped using Cell Ranger version 4.0.0 (10X

Genomics) with default settings and a custom genome reference based on the

comprehensive gene annotation set from Gencode Release 32 (GRCh38.p13)

(http://ftp.ebi.ac.uk/pub/databases/gencode/Gencode_human/release_32/gencode.v32.

annotation.gtf.gz). For the multiplexed pool, sample identities for each cell were assigned

using Vireo as described above, such that each sample could be subset from the pooled

matrix during analysis.

Filtering, feature selection, clustering, and other secondary analyses were carried

out in R using Seurat v4.0.3 (https://satijalab.org/seurat/) (47,81). Gene set enrichment

analyses were performed using GSEA v4.1.0 (82). Following gene sets were used for the

various signature scores: adult human breast epithelium markers from (49), Mammary

epithelial   lineage   scores   from   (40),   NOTCH   signaling:

REACTOME_SIGNALING_BY_NOTCH,   BMP2   targets: LEE_BMP2_TARGETS_UP,

MYC signature from (83), Hypoxia signature: HALLMARK_HYPOXIA, Basal mammary

stem cell signature from (40).

Quantification and statistical analysis: Statistical analyses were performed using R

(version 4.1.0) on RStudio and GraphPad Prism software (v9.1.2, GraphPad Software,

San Diego, California USA, www.graphpad.com). Specific tests are indicated in the figure

legends along with the statistical significance.

Materials, data and code availability: Organoid lines generated under the HCMI project (starting with HCM) will be available for purchase from the American Type Culture Collection (ATCC). All raw data will be available to download from dbGaP (phs002722.v1.p1 and phs001486). Copy number segment file is available in Table S2, transcriptome signature scores are in Table S4. Metadata for the different PDOs is available in Table S1. All original code will be deposited at https://github.com/bhatia-sonam/manuscript-v1

Results:

Establishment and somatic variant profiling of a diverse patient-derived breast cancer organoid biobank

Breast cancer tissues along with paired normal breast tissues were acquired from female patients and developed into 3-D organoids. In addition to breast cancer tissues, we also collected normal reductive mammoplasty samples from 10 cancer-free patients to generate normal PDOs for downstream analyses. The majority of tumor samples were from patients with invasive ductal carcinomas (56/87, Fig. 1A), 11/87 were from invasive lobular carcinomas, 4/87 were from metastatic lymph nodes and a small percentage from other categories (Fig 1A, Table S1). The majority of the tumor samples (43/87) were luminal BC as defined by immunohistochemistry (ER/PR+/HER2-), 37/87 were from TNBCs (ER-PR-HER2-), 2/87 were from ER/PR+/HER2+ BCs, 2/87 were from ER-/PR-/HER2+ subtype, and 3/87 samples belonged to post-treatment residual tissue with no visible carcinoma (NA) (Fig 1B). Samples were collected from patients of various age

groups (Fig 1C) and diverse ethnic and racial backgrounds (Fig 1D). An increased

proportion of TNBC tumor samples was observed from black patients with African or west-

Indian heritage (Fig 1D, Table S1) which recapitulates the higher incidence of TNBCs in

the African-American community (29,30). Using a DNA-seq panel of 143 cancer driver

genes, we identified pathogenic single nucleotide variants (SNVs) in 49 tumor organoid

lines (Fig 1E); *PIK3CA* was mutated in 33% of the tumor organoid lines, the majority of

which were from patients with luminal breast cancer. 24% of the organoid lines (45% of

TNBC PDOs 10/22) had a pathogenic *TP53* mutation and primarily represented the high

grade TNBC cohort (Fig 1E). Two of the organoid lines are derived from a patient with

rare breast cancer showing adenoid cystic carcinoma (AdCC) like morphology, NH87T

(primary tumor) and HCM-CSHL-0655-C50 (lymph node metastasis from the same

patient) (Table S1). The AdCC organoids show a non-traditional TNBC mutation profile

with pathogenic mutations in *APC*, *KDM6A* and *NOTCH1* (Fig 1E), which were previously

observed in AdCCs of the breast (31). Mutations in *KMT2C* and *GATA3* were also

observed in a variety of luminal breast cancer organoids, *CDH1* mutations were present

in 10% of PDOs all of which belonged to invasive lobular carcinomas, and *ARID1B*

mutations were found in a few TNBC-derived organoids (Fig 1E). Overall, the mutation

profiles of patient-derived BC organoids are in concordance with the mutational

landscape of BCs (10,26). A relatively large subset of tissues (24/87) that resulted in

cultured organoids did not have pathogenic SNVs (Fig 1F, Table S1), while 10/87

samples dropped off in culture at early passages (Fig 1F). The PDOs with pathogenic

SNVs showed a range of growth properties *in vitro* with subsets exhibiting long-term

continued expansion while others seem to have more limited cultures either in terms of

maximum passages reached or limited expansion abilities (Table S1).

A comparison of the SNV profile between several pairs of patient tumor and PDOs

showed high concordance of the pathogenic SNVs between the tumor and tumor derived

organoids (Fig S1A), except in some cases where the SNVs are present at a much lower

variant allele frequency (VAF) in the tumor, for instance, NH84TT-P0 showed <25% TP53

VAF compared to 100% VAF in NH84TT-p10 PDO (Fig S1A) suggesting normal

contamination in the tumor tissues and a successful outgrowth of cancer cells in organoid

cultures. Longitudinal analyses for some patient-derived organoids showed the stability

of primary driver mutations overtime in culture (Fig S1B). To determine whether the

amount of tumor material was a limiting step for generation of organoids, we collected

tumor scrapings (see methods) from a subset of tumor samples. A comparison of tumor

tissue (labeled TT) and the scrapings (labeled Sc) from three different patients (Fig S1C)

showed high concordance and indicates that, if necessary, small amounts of tumor

material can be used to generate organoids. We also examined the pathogenic SNVs in

TNBC tissue samples (p0) that did not result in a successful generation of organoids

(establishment or long-term cultures) and 6/10 had TP53 mutations and some samples

also had BRCA1/2 mutations (Fig S1D), the latter were largely absent in successful

organoid cultures (Fig 1E).

Breast cancer organoids are enriched for copy number alterations

Having identified pathogenic driver mutations for the tumor-derived organoids, we

performed copy number analysis using SMASH (short multiply aggregated sequence

homologies) (32) for various PDOs. TP53 mutated TNBC PDOs make up the majority of

the ER- subset and are highly genomically altered compared to the luminal ER+

organoids (Fig 2A, B, Table S2). Some organoids that were deemed to be derived from

tumor based on pathogenic SNVs (eg. LNS12T, LNS18T, NH06T etc.) showed no

prominent CNAs. Some TNBC PDOs that showed trace SNVs such as NH58T, NH72T

and NH66T also showed minimal CNAs (Fig 1E and 2A). In keeping with large scale

genomic datasets (10,26), a consistent gain of chromosome 1q (chr1q) and loss of chr16q

was observed in the luminal organoids where CNAs were detected (Fig 2A). The TP53

mutated TNBC organoids were highly genomically aberrant where multiple lines showed

a gain of chromosomes (chrs) 1q, 8q, 19q and chrs 7, 20 and 21; loss of chrs 3p, 4, 5,

17p and Xp was also observed in multiple organoid lines (Fig 2A). These TNBC PDOs

also showed copy number loss of chr4q and chr5q (Fig 2A) that have been previously

reported to be over-represented in basal-like breast cancers (33). Of note, the AdCC-like

organoids, NH87T (primary tumor) and HCM-CSHL-0655-C50 (lymph node metastasis

from the same patient), were considerably less genomically aberrant as compared to the

other TNBC organoids and showed focal amplification of chr8q and deletion of chr6p (Fig

2A, C-D). Interestingly, the chrX deletion was only observed in the primary tumor (NH87T)

sample (Fig 2A, C) and a focal deletion was observed in chr2 of the lymph met HCM-

CSHL-0655-C50 (Fig 2C). The AdCC organoid CNA profiles are consistent with the low

CNA profiles that are typically observed in AdCC-like breast cancers (31).

For a subset of these samples, we compared the copy number profiles of the

primary tumor with the paired tumor and normal organoids and observed more

pronounced CNAs in the organoids compared to the tumors suggesting that the organoid

culture enriches for tumor cells (Fig 2C-D). As observed in the SNV data (Fig S1) the

NH84T tumor (p0 with <25% VAF of pathogenic TP53) was likely very heterogenous and

contained a significant population of normal cells, however, the tumor cells successfully

outgrew the normal and resulted in a highly pure tumor organoid culture over multiple

passages (Fig 2C-D). For HCM-CSHL-0366-C50 and NH87T the early passage p1 or

tumor tissue (p0) respectively, had relatively less normal contamination and maintained

their copy number profiles over time in culture (Fig 2C-D). While copy number profiles

were enriched for in successful tumor PDOs, for some cultures we observed loss of copy

number alterations in the organoids over-time, suggesting a normal outgrowth (Fig S1E).

Since tumor organoids are more enriched for tumor cells compared to the p0 tumor

tissue, we profiled the putative cancer driver gene panel in individual organoid samples.

TNBC PDOs typically had a higher copy number alteration frequency of these cancer

driver genes (top bar-graph in Fig 2B). *TGFB2*, *MDM3*, *AKT3*, *DDR2* and *INSRR* showed

the highest frequency of alteration (Fig 2B) and are all genes present on chr1q that is

amplified in both luminal and TNBC PDOs (Fig 2A-B). Interestingly, we also found a

higher frequency of deletion of *KDM6A*, *ARAF* and *RPS6KA3* tumor suppressors, located

on chrXp, specifically in the TNBC-derived organoids (arrows in Fig 2B). Loss of

PPP2R2A (chr8p), which was previously reported to be a tumor suppressor in breast

cancer (10), was also identified in 38% of the samples, the majority of which are of TNBC

subtype (Fig 2B). Furthermore, there is an over-representation of loss of chr3p in the

TNBC samples, that results in the deletion of potential tumor suppressors: *ACVR2B*,

BAP1, *CTNNB1*, *MLH1*, *MYF88* and *PBRM1* (Fig 2B). Copy number loss of chr6q is also

common in TNBC organoids and results in loss of *ARID1B*, *MAP3K4*, and *PARK2*, a

master regulator of G1/S cyclins (34). *ESR1*, the gene encoding estrogen receptor (ER),

present on chr6q is also frequently lost in these TNBC organoids (Fig 2A-B). We also

identified gains of *CALR*, *JAK3*, *KEAP1*, *PIN1* and *SMARCA4* that are associated with

the amplification of chr19p and amplification of mismatch repair (MMR) genes *MSH2* and

MSH6 located on chr2p (Fig 2A-B) in subsets of TNBC PDOs. While MMR genes are

commonly mutated in various cancers, their overexpression was recently associated with

aggressive prostate cancers (35) and was shown to promote genomic instability in yeast

(36) but their role is yet to be determined in TNBC disease progression.

Taken together, the SNV and copy number profiling of PDOs shows robust

retention of genomic features of various types of breast cancers, including luminal, TNBC,

and rare AdCC-like carcinomas. While there might be small alterations overtime in PDO

cultures, the key pathogenic mutations and overall copy number profiles are conserved

throughout organoid culture (Fig S1, 2C-D) supporting their utilization as valid cancer

models. Additionally, we find overrepresentation of some lesser studied copy number

variants in our data such as loss of tumor suppressors *RPS6KA3*, *PPP2R2A* and *PARK2*

and copy number gains of MMR genes *MSH2* and *MSH6* that might have important

unexplored consequences in breast cancer progression.

A subset of TNBC organoids recapitulate signatures of aggressive basal-like breast

cancers

Next, we performed RNA-seq on various tumor and normal-derived organoids to

profile their transcriptomes (Table S3). Hierarchical clustering of samples by euclidean

distance divided them into six groups (Fig S2A). We used supervised clustering with 838

previously curated gene expression signatures (37–39) (Table S4) and found similar

patterns of clustering among the various organoid lines (Fig 3A). Group 4 is largely

comprised of the true normal organoids (labeled with prefix NM in Fig S2A) derived from

breast tissues obtained from individuals undergoing reductive mammoplasty (Normal in

Fig 3A-B, S2A, C), paired normal PDOs derived from normal tissue adjacent or distal to

the tumor and very few tumor-derived organoids; this cluster showed an enrichment of

normal mammary stem cell (MaSC) signatures and low proliferation related expression

profiles (Fig 3A-B). Group 1 is comprised of a mixture of either paired normal organoids

or tumor organoids that did not have a strong driver mutation (eg. *PIK3CA* or *TP53*) (Fig

3A, S2A, 1E). This group had a signature profile that was distinct from the true normal

group (Group 4) and is most similar to luminal cell signatures (Fig 3A). The samples in

this group also have a higher proliferation signature compared to the true normal Group

4 (Fig 3A). Group 5 mostly contained PDOs derived from Luminal BC and showed luminal

like gene-expression signatures (Fig 3A-B). Signature profiling also uncovered immune

and stromal signatures that highly corresponded to samples in Group 3 (Fig 3A, S2C).

Interestingly, two of the samples in this group NH63T and NH54T exhibited limited

propagation in culture due to stromal outgrowth leading us to hypothesize that these

samples had some fibroblast-like cells. We also found a strong interferon (IFN) signature

in this group along with some additional PDOs from Group 6 (Fig 3A, S2C).

Groups 2 and 6 are comprised mostly of TNBC organoids and all organoid lines in

these groups could be propagated to long-term cultures (>passage10, labeled LT-

TNBCs) and showed continued expansion (Fig 3A, S2A). The only exception is the

NH48N normal sample, which was confirmed to be mostly tumor by copy number analysis

and identical to its counterpart NH48T (Fig S2B). The most prominent signatures of

Groups 2 and 6 correspond to basal-like breast cancer gene-sets that are also defined

by luminal progenitor (LP) like signatures (Fig 3A-B, S2C-D). LP-like gene expression has

previously been shown to be associated with basal-like breast cancers (40). The

organoids in this group also showed enrichment of proliferation signatures (Fig 3A-B) and

the majority had a *TP53* mutation which was in conjunction with the *PIK3CA* mutations

for three samples (HCM-CSHL-0773-C50, HUB4T and HCM-CSHL-0155-C50) (Fig 3A,

S2A). We also observed a MYC amplification signature in this subgroup (Fig 3B) that was

accompanied by the copy number amplification of *cMYC* in many of these PDOs (Fig

S2E). Additionally, gene sets associated with hypoxia, glycolysis, angiogenesis and fetal

MaSC (fMaSC) metabolism signatures were also enriched in LT-TNBCs (Fig 3A-B, S2C).

The *VEGF* 13-gene signature showed high correlation with basal-like breast cancers (29),

has prognostic significance and is associated with poor outcome in breast and other

cancers (41). Similarly, fMaSC metabolism signature is a refined 8-gene signature, which

primarily comprises genes associated with glycolysis and fatty acid metabolism, and was

previously shown to be associated with TNBCs and metastatic TNBC lesions (42). This

suggests that the PDOs with these signatures are likely associated with tumors linked to

poor outcomes.

In order to understand the complexity of organoid cultures we profiled the growth

properties of various TNBC and normal organoids. TNBC organoids appear as densely

filled balls of cells which is in stark contrast to the normal organoids that have a

predominant acinar structure with a central lumen and some organoids that were filled

(NM04N in Fig 3C). The TNBC-organoids showed varying degrees of propagation in

culture where some lines could be propagated for over 10 passages (LT-TNBCs) with

continued expansion while many short-term culture lines dropped out of culture at various

points (Fig 3D, S3A) and had limited material to assay. Low starting material, limited

proliferation, normal and stromal outgrowths were among the primary reasons for short

term culture growth (Fig S3A). We focused on the LT-TNBCs for all downstream

analyses. IHC for the proliferation marker, Ki-67, in confluent PDO cultures showed large

variability of expression amongst the various organoid lines and also within each culture

(Fig 3C- lower panels). In concordance with expression signatures (Fig 3B), LT-TNBC

organoids were highly proliferative compared to normal and luminal organoids (Fig 3C,

S3B-C). There was a lot of heterogeneity within organoid cultures with some organoids

being mostly comprised of proliferating cells, for instance NH85TSc and NH95T, while

other cultures were more mixed (Fig S3C). Normal-derived organoids typically had <10%

of proliferating cells per organoid and showed luminal-basal organization with a hollow

lumen, while TNBC-derived organoids tend to be highly proliferative and undifferentiated

with little to no observed cellular organization (Fig S3C). When seeded as single cells, we

observed a range of organoid formation amongst the various lines (Fig S3D) which did

not always correlate with the proliferative index of the PDOs, and might be due to the

requirement of cell-cell contact in some but not all PDOs that is apparent when seeding

at low density as single cells.

Of note, none of the TNBCs propagated in culture showed expression profiles of

the more mesenchymal-like claudin-low subgroup (6), possibly because the culture

conditions are more favorable towards the propagation of epithelial cells. Overall, the

gene expression signatures of the normal and tumor-derived organoids depict the

expression profiles of luminal and basal-like breast cancers. The TNBC organoids that

can be propagated to long-term cultures (LT-TNBCs), Groups 2 and 6, had very classic

basal-like breast cancer signatures and were associated with proliferation (Fig 3A-D),

hypoxia (Fig 3A-B) and *c-MYC* amplification gene expression signatures (Fig 3B, S2E).

Furthermore, about 40% of basal-breast cancers exhibit *c-MYC* amplification (26) and the

majority of our LT-TNBC organoids show an enrichment of *c-MYC* (Fig S2E) signature

suggesting that our organoid system results in long-term expansion of TNBCs that are *c-*

MYC driven and have a LP-like basal breast cancer signature.

As a proof-of-concept, we tested whether these PDOs were amenable to drug

response assays, and performed dose response curves using 6 different PDO lines with

three different drugs (Fig S3E). TNBC PDOs (Fig S3D) showed high sensitivity to the

cytotoxic chemotherapeutic agent paclitaxel compared to the luminal line NH53T and the

TNBC-met HCM-CSHL-0773-C50 (Fig S3E). Conversely, capivasertib (AZD5363), a pan-

AKT inhibitor used in PIK3C mutated cancers (43), showed greater sensitivity against the

PIK3CA mutated luminal NH53T PDO (Fig S3E). Similarly, afatinib a targeted inhibitor for

EGFR, showed the most activity in the EGFR amplified NH84T PDO line (Fig S3E).

TNBC organoids can recapitulate tumor morphology *in-vivo*

Next, we examined the *in-vivo* tumor forming abilities of these patient-derived

cultures. We transplanted organoids from eight different TNBC (LT-TNBCs) lines into the

mammary-fat pads of NOD-SCID mice and assayed their tumor formation and metastatic

potential over time (Fig 4A). We found striking differences in the tumor forming ability

between the different organoid lines (Fig 4B) despite having highly aberrant genomic

profiles (Fig 2, 4C). TNBC organoids HCM-CSHL-0366-C50, NH85TSc and NH87T

resulted in palpable masses around 30 days and steady tumor formation in 100% of

injection sites (Fig 4B, Table S6). NH85TSc-derived tumors grew rapidly and resulted in

visibly necrotic masses that needed to be resected at ~100 days. HCM-CSHL-0366-C50

resulted in primary tumors that also developed fluid filled cysts on top of the tumors that

drained on their own. NH87T and HCM-CSHL-0655-C50 are the AdCC-like primary tumor

and lymph-node metastasis from the same patient. Interestingly, while NH87T formed

tumors rapidly, HCM-CSHL-0655-C50 did not and the tumors remained small (Fig 4B,

Table S6), despite both of these lines having equivalent growth properties *in vitro* (Fig 3,

S3). This observation was consistent for the two independent transplant experiments

done with different passage organoids (Table S6). NH95T resulted in small tumor masses

at multiple injection sites that were slow to grow while NH84T and NH93T had small

tumors observable only in fat-pad histology sections of some sites and NH64T did not

result in any primary tumors (Figs 4B, S4A, Table S6) despite being highly genomically

aberrant (Figs 4C, 1-2) and having high proliferation and organoid formation rates *in-vitro*

(Figs S3C-D).

Remarkably, the PDO-derived xenograft (PDO-X) tumors resulting from HCM-

CSHL-0366-C50, NH85TSc, NH87T and NH95T organoids recapitulated the morphology

of the patient tumor despite previously being in culture for up to 18 passages. The tumor

and the organoid-xenograft from patient HCM-CSHL-0366-C50 were the most distinct

and showed squamous differentiation with pleomorphic cells that were variable in size

and had cells with either oval or pale nuclei with prominent nucleoli (Fig 4D). We also

observed cells that had thin spinous connections with adjacent tumor cells which were

also present in the organoids (black arrows Fig 4D). NH85TSc tumor and organoid-

xenografts both showed highly undifferentiated morphology with large necrotic areas and

stromal compartments infiltrating in between the tumors (Fig 4D). Lastly, NH87T tumor

was a TNBC type with AdCC like features that are characterized by cribriform

architectural patterns and pseudo-lumens (31,44). While the organoid-derived xenograft

had a lower stromal composition compared to the patient tumor it still recapitulated the

key features of the original tumor, including the presence of cuboidal cells, cribriform

architecture and pseudo-lumens (yellow arrows Fig 4D). This particular subtype is also

characterized by the expression of CD117 and CK5/6 (45). We performed IHC for CD117

and CK5/6 on PDO-X tumor sections from NH87T and observed positive membrane

labeling for CD117 along with areas of high and low CK5/6 labeling as observed in the

clinical IHC labeling of the patient slide (Fig S4C).

While NH95T formed tumors at a much lower efficiency (Fig 4A, B), the tumors

that did form had morphological similarities to the original patient tumor (Fig 4D). The

patient tumor cells showed an organization pattern which was recapitulated in the

organoid-derived xenografts and to some extent in the NH95T organoids (Fig 4D). Of

note, patients from whom the fast-growing organoids were derived, i.e., HCM-CSHL-

0366-C50, NH85 and NH87, presented with poor diagnoses and outcomes (TMN staging

in Fig 4B). Patient HCM-CSHL-0366-C50 had rapid metastasis to the brain and

succumbed from the disease, patient NH85 had local recurrence in the lymph node and

patient NH87 presented with a positive lymph node at initial diagnosis (Table S1). Using

IHC with a human mitochondrial antibody we only observed micro-metastasis and single-

cell metastasis in our experiments (Fig 4D, S4B, Table S6), however, altering the

experimental conditions such as: prolonging the end-point, using a more

immunocompromised NOD/SCID gamma (NSG) mouse model or tail-vein injections

might result in more metastatic lesions and will be examined in future studies. Thus far,

our data shows that TNBC PDOs recapitulate the tumor intrinsic properties of the original

tumors at genomic, transcriptomic and morphological levels.

TNBC derived organoids are enriched for luminal-progenitor-like cells

To fully assess the utility of PDOs as cancer models, we next asked what were the

cell types represented within these organoids and how did they differ from the normal

derived PDOs. In order to profile the cell-types present within the normal and TNBC

organoids, we used a combination of flow cytometry and single cell RNA-seq (scRNA-

seq). We assayed for mammary epithelial lineages using EPCAM and CD49f as luminal

and basal cell markers respectively (21,22,40,46). As previously shown (22), the normal

derived organoids recapitulate the EPCAM+ luminal lineage, EPCAM+/CD49f+ luminal

progenitor cells and EPCAM-/CD49f+ basal cell lineages (NM07NL in Fig 5A). The

populations slightly fluctuated between the different normal lines and passages but

overall, the different lineages were observed in all normal organoids (Fig 5A-C, S5A,

Table S7). TNBC organoids were less heterogeneous in the expression of these markers,

and as observed by RNA-seq signatures (Fig 3B, S2D), showed an enrichment of luminal

progenitor-like cells (Fig 5A-B) which had co-expression of EPCAM and CD49f but

showed slightly different and patient-specific expression patterns of these markers (Fig

5A, S5A). TNBC organoids also showed a higher gene expression of EPCAM and CD49f

(gene *ITGA6*) compared to the normal organoids (Fig S5B). Some TNBC PDOs, for

example NH84T, had a more luminal cell flow-cytometry profile, despite clustering with

the other LT-TNBCs at the transcriptome level. Interestingly, metastatic TNBC lines HCM-

CSHL-0773-C50 and HUB4T also had an enrichment of more luminal-like cells (Fig 5B,

S5D). NH85TSc which was derived from a patient that later relapsed and HCM-CSHL-

0366-C50, which showed rapid progression in the patient, also had a more luminal cell

profile with higher percentage of EPCAM only cells (Fig 5A-B) despite showing gene

signatures of the basal-like BCs. The EPCAM/CD49f profile of the individual lines was

stable overtime at different passages (Fig S5C). Additionally, while always observed in

the normal-organoids, we rarely observed CD49f only cells in the TNBC-organoids (Fig

5A-B, S5A, D). We performed longitudinal analysis for one of the patient-derived organoid

lines and observed an enrichment of the tumor LPs in culture over time in early passages

(Fig 5C) which coincided with a more pronounced copy number profile of the TNBC

organoids (Fig 5D) suggesting LP-like cells being the predominant cancer cell population

in TNBCs and potentially the cell of origin of these cancers.

In order to better understand the cellular composition of these organoids, we

performed scRNA-seq analysis on three normal and seven TNBC organoid lines. The

study was done in three experimental batches (Table S7) and processed where each of

the samples was quality controlled and filtered to remove cells with high mitochondrial

gene content and low gene identification (i.e. dead cells). The filtered matrix was SC-

transformed using Seurat (47) and the samples were integrated to account for the

different batches (Fig 5E). Clustering of data identified 19 clusters (Fig 5E) with some

clusters being highly representative of the normal organoids (clusters 0, 3, 5, 6, 9, 11 in

Fig 5F) while other clusters were largely comprised of tumor cells. Tumor organoids were

distinct from normal, with some overlap in clusters 0, 3 and 9 (Fig 5F-H). Expression

profiles of *EPCAM* and *ITGA6* (gene encoding CD49f) (Fig 5H) recapitulated the data

from flow cytometry (Fig 5A-B) where tumor organoids are predominantly of the luminal

progenitor nature as measured by co-expression of *EPCAM* and *ITGA6* (CD49f) while the

normal organoids recapitulate the three broad mammary epithelial lineages: mature

luminal (EPCAM+), luminal progenitors (LPs, EPCAM+/CD49f+) and myoepithelial/basal-

like cells (Epcam-low/CD49f+). Unlike the normal organoids, all tumor cells were

EPCAM+ and a subset of those had low *ITGA6* (Fig 5H). In concordance with the flow

cytometry data (Fig 5A), the *ITGA6*-low cells were largely present in HCM-CSHL-0366-

C50 and NH85TSc. NH93T and the NH95T organoid line had similar profiles and

occupied similar space as the normal LP cells (Fig 5H). Lines NH87T and HCM-CSHL-

0655-C50 are paired primary and lymph node metastasis samples from the same patient

and occupied very similar spaces with some overlap with normal LP cells (Fig 5H), while

NH64T, NH85TSc and HCM-CSHL-0366-C50 are very distinct from the normal lines.

Interestingly, there was a significant overlap between the clusters identified between

NH85TSc and HCM-CSHL-0366-C50 samples (Fig 5H) in keeping with their

transcriptome similarity (Fig 3A, S2A) and rapid tumor progression *in vivo* (Fig 4A).

This data builds on previous studies that have correlated a LP-like expression

signature with basal-like breast cancers (40) and have shown that LP cells are the cell of

origin of BRCA-mutated basal breast cancers (48). We suggest that LP-like cells are the

possible cell of origin for a broader subset of TNBCs. Interestingly, a higher percent of

EPCAM+ only cells seems to be associated with a greater degree of disease progression,

however, this needs to be further investigated.

Tumor LP-like cells exhibit altered expression and have an upregulation of NOTCH

and MYC downstream pathways

Since TNBC organoids had a large proportion of LP-like cells that seemed distinct

from the normal LP cells (Figs 5, S5) we performed integrated single cell analysis by SC-

transforming individual samples and performing an anchor-dependent integration for all

individual samples. We identified thirteen clusters between the tumor and normal cells

(Fig 6A) out of which clusters 6, 8, 10 and 9 represented cell cycle clusters (Fig S6A) with

cluster 6 representing a population of G1-S phase cells, cluster 8 representing S-phase

cells, cluster 9 S-G2M transition cells and cluster 10 marking G2M cells while the

remaining clusters represented G1 cells (Fig S6A-C). The cell clusters identified

represented biologically meaningful cell-types recently annotated by single cell

sequencing of adult human breast epithelium (49). Cell types identified included: XBP1,

AGR2 expressing mature luminal cells (Mat lum, cluster 1), APOE, KRT6A expressing

basal stem-like cells (basal SC, cluster 4), TAGLN, TIMP1 expressing myoepithelial cells

(Myo, cluster 3), SLPI expressing luminal progenitor cells (LP, cluster 7) and LTF

expressing secretory cells (LP sec, cluster 2) (Fig 6B, S6C) (49). To further validate the

lineage specification in organoids we overlaid the published mammary epithelial gene-

sets (40) and computed a score for each of the three mammary epithelial cells. The

normal cell clusters showed a much stronger cell type enrichment score while the tumor

cells despite being predominantly EPCAM+/CD49f+ showed a more diffused enrichment

of these signatures, suggesting some cell type heterogeneity within these tumor

organoids (Fig 6C). In normal organoids, Clusters 2,7 and 11 had a higher enrichment for

luminal progenitor (LP) cell score, while cluster 1 showed an enrichment for mature

luminal (mature Lum) cells and cluster 3 was predominantly of the basal mammary stem

cell (Basal SC) compartment (Fig 6C, S6C). The tumor organoids showed varied

expression profiles of various luminal/basal markers (Fig S6D) and a diffused cluster

specific enrichment of lineage scores (Fig 6C, S6E), suggesting that while they have lost

proper cell type specification, the clusters identified using this integration approach have

some similarity to the normal cell lineages.

      With the aim to identify mechanisms that underlie the tumor luminal progenitor-like

(LP-like) cells in TNBCs, we performed a differential expression analysis between tumor

and normal cells of clusters 2, 7 and 11. We identified 1103 significantly differentially

expressed genes (p_val_adj<0.05) between the tumor LPs vs normal LPs (Table S8).

GSEA on the differentially expressed genes showed an enrichment of NOTCH signaling

related genes in the tumor LPs versus an enrichment of BMP2 targets in the normal LPs

(Fig 6D). Leading edge genes from these gene sets showed a consistent downregulation

of NOTCH signaling related genes, *RPS27A*, *YBX1*, *JAG1*, *MDK*, *MYC* and *SEM1*, and

an upregulation of BMP2 target genes, *LTF*, *MGP*, *KRT16*, *KRT7*, *PLAAT3* and *NTRK2*

in normal LPs from all three patients (Fig 6E). MDK, JAG1, YBX1 are ligands of NOTCH1,

while *MYC* and *RSP27A* are downstream targets. While NOTCH activity was present in

normal LP cells, it was more pronounced in the TNBC organoids (Fig S6F).

      A further investigation into differentially regulated pathways showed upregulation

of genes involved in ribosomal biogenesis and translation in the tumor LPs. We performed

motif enrichment analysis on the differentially expressed LP genes and found a significant

enrichment of MYC binding sites in the tumor LP expressed genes (Fig S6G). We used

published MYC signatures to assess MYC activity in tumor and normal organoids and

found a higher ubiquitous enrichment of MYC activation in tumor organoid cells, while in

normal organoids it seemed to be higher in the proliferating cell clusters (Fig S6F).

Similarly, while the NOTCH pathway was hyper-active in tumor LPs it was ubiquitously

active across all tumor cells (Fig S6F). We tested whether NOTCH and MYC activation

was required for organoid formation and seeded NH95T, NH85TSc and HCM-CSHL-

0366-C50 as single cells in regular organoid growth medium in the presence of DAPT

(NOTCH inhibitor) or MYCi975 (MYC inhibitor) (50) and allowed for organoids to form for

12 days. We saw a significant reduction in organoid formation in the presence of both

MYC and NOTCH inhibitors for NH85TSc and NH95T but not for HCM-CSHL-0366-C50

(Fig 6F). While HCM-CSHL-0366-C50 did not show reduction in the number of organoids

formed, we did observe a significant difference in organoid size with DAPT and MYCi975

suggesting growth defects on inhibition of these pathways in all lines (Fig S6H). We

further tested whether inhibition of NOTCH and MYC is necessary for the formation of

normal PDOs. Since, normal PDOs have multiple populations we flow-sorted the basal

stem-like CD49f+ve population and EPCAM+/CD49f+ LP cells and seeded them as single

cells in the presence of DAPT or MYCi975 (Fig S6I). While DAPT had no effect on

generation of normal PDOs from both LP and basal stem-like cells, MYC inhibition

resulted in no organoid formation.

Thus, our data shows that TNBC organoids have an underlying LP expression

signature which is driven by LP-like cells that exhibit altered expression from normal LPs

by hyperactivation of NOTCH and MYC signaling. These pathways are necessary for the

formation of some, but not all, TNBC organoids and when perturbed result in proliferation

defects in all tumor lines examined.

TNBC organoids are comprised of heterogenous cancer cell populations

Since normal organoids have clearly defined lineages, the clustering of the

integrated dataset (Fig 6) was largely driven by the normal organoids. To assess the

heterogeneity that exists within TNBC organoid cultures we performed a similar integrated

analysis on the TNBC organoids only and identified 13 different cell clusters (Fig 7A,

S7A). Clusters 3,9,6,8 and 5 correspond to the different cell cycle phases (Fig 7A) and

the remaining clusters are G1 cells (Fig 7A-B). We identified the markers that uniquely

define each of the clusters (Fig 7C, Table S9) and performed GSEA on the marker genes

to identify the phenotypes associated with each cell cluster (Fig 7D). Cluster 0

represented a mixed cell cluster with some enrichment of mature luminal-like cells.

Cluster 1 was defined as Basal-like 1 as it showed a positive enrichment of a basal breast

cancer gene-set including specific markers *SAT1*, *GABRP*, *TM4SF1*, *TTYH1*, *KRT16* and

KRT6A genes (Table S9). Cluster 2 is mesenchymal-like due to the selective expression

of mesenchyme genes including *CCN2*, *TPM1* and *LAMB1*. Cluster 7, MGP-high cluster,

is a relatively distinct cluster with a very specific high expression of *MGP* (Matrix Gla

Protein) in all TNBC organoid lines (Fig 7C, S7E). MGP is normally expressed in smooth

muscle cells perhaps suggesting that these cells might have contractile abilities. Cluster

12 had a very high expression of ribosomal and translation related genes (Fig 7C-D).

Clusters 4,10 and 11 are the most distinct of the tumor cell clusters and are identified by

the unique expression of certain genes. Cluster 10 represents an NFKB-active cluster

that was also found in normal organoids (Fig S6C), albeit with some differences in gene

expression. As in normal organoids (Fig 6A, S6B, 8C). this cluster shows the expression

of *CXCL1*, *CXCL3*, *NFKBIA* etc. Cluster 11 is also defined by a basal breast cancer

signature (Fig 7C-D), is labeled Basal-like 2 and shows a selective expression of *KRT17*.

Cluster 4 is a hypoxia cell cluster that shows selective high expression of *EGLN3*,

NDRG1, *VEGFA* and likely represents cells at the center of these organoids (Fig 7C-D).

This cluster is represented by a higher hypoxia score (Fig 7C-D), which interestingly

correlates with basal stem cell (Basal SC) signatures and shows comparatively low

activity of MYC and NOTCH pathways (Fig 7E and S7D-E). We also noted that the cluster

specific genes for all clusters were fairly well conserved between the different PDO lines

(Fig S7F). Overall, this data suggests that while the TNBC organoids retain patient

intrinsic properties (Fig 1-4) there are common cell signatures that define the cell types

within these organoids.

The distinct gene expression patterns within these cell clusters also suggests that

the TNBC organoids are comprised of multiple cell types that are known to be involved in

tumorigenesis, tumor progression and metastasis, and can therefore be used as models

to study the various aspects of breast cancer progression.

Discussion:

We developed a patient-derived biobank of normal and breast cancer organoids

from a diverse group of patients with a focus on the highly aggressive TNBC subtype.

Our biobank is heterogenous in terms of ethnic/racial backgrounds, patient age and

breast cancer subtypes. A comprehensive genomic, transcriptomic, and cellular

characterization of these PDO models demonstrate their faithful recapitulation of the

patient tumor intrinsic properties and hence validates them as cancer models to study

various aspects of breast cancer progression and treatment.

Patient-derived organoids are an exciting step towards personalized medicine with

a promise to be used for real-time drug screens for guiding patient treatment (14). When

successfully established, breast cancer PDOs retain the genomic and transcriptomic

features of breast cancers and their parent tissues. We found that, the TNBC organoids

that showed long-term robust growth had activated MYC signaling, an LP-like gene

expression signature and were overwhelmingly composed of LP-like cells. As these are

among the most aggressive forms of breast cancer, organoid models appear to be an

exciting avenue for their study. In addition to replicating the tumor specific features *in-*

vitro, TNBC PDOs when transplanted into NOD/SCID mice generated tumors with

remarkable morphological similarity to that of the original patient tumors, despite being in

long-term organoid culture for up-to 18 passages. Long-term cultured organoids, thus,

maintain the intrinsic ability to represent the tumor from which they were derived when

placed in an *in vivo* environment. Interestingly, in our study not all PDOs readily generated

primary tumors *in vivo* despite being highly proliferative and genomically aberrant

organoids. The association of PDOs that generated tumors rapidly, with worse outcomes

and diagnosis, suggests that there might be a biologically relevant explanation for this

observed difference and must be investigated further along with long-term patient follow

up information. Patient-derived xenograft (PDX) models also retain the histopathology of

the original patient tumors (51,52) and a recent pre-print study showed successful

derivation of breast cancer organoids from PDX models (53). Our models are

complementary to this system and represent an opportunity to derive PDOs first, perform

in-vitro assays and drug screens, followed by PDOX derivation for *in-vivo* validation

studies.

While successfully established PDOs faithfully recapitulate the patient tumor

properties, the efficiency of establishment of these cultures is currently less than ideal.

The common challenges we faced during establishing organoid cultures included normal

outgrowth in some tumor organoid cultures, stromal outgrowth, non-proliferating or

dormant tumor cells, or limited growth potential *ex-vivo*. Since the medium composition

for tumor organoid growth also supports the culture of normal mammary organoids (22)

the purity of the starting tumor tissue might govern the time to generation of tumor-only

organoid cultures and early passages should be meticulously tested at a genomic level

before any pharmacological studies are carried out with these models. Furthermore,

some subtypes of breast cancer might be challenging to culture including the less

proliferative Luminal A subtype or the more mesenchymal claudin-low group of TNBCs.

Efforts to culture tumor-organoids in reduced complexity medium (53), might alleviate

some of these challenges and needs to be examined in future studies. The PDO culture

conditions are comprised of complex growth factors including, EGF, FGF10, Neuregulin,

R-spondin etc. (20), and it remains to be tested whether the removal or alteration of these

components would be beneficial in selecting out cancer cells over normal epithelial cells,

resulting in higher efficiency in the establishment of TNBC PDOs. However, the

simultaneous growth of normal and TNBC organoids in the same culture conditions allows

for comparison of similar cell types within these organoids.

We used an integrated scRNA-seq approach to compare normal LP cells with the

tumor LP-like cells and identified hyperactivation of NOTCH and MYC signaling in the

tumor compared to normal LPs. LP cells were previously shown to be the cell of origin of

BRCA1 mutated basal-like BCs (40,48) and are speculated to be involved in

tumorigenesis of all basal-like BCs due to similarity of gene expression profiles (40). Our

data provides strong support for this hypothesis by showing that TNBC PDOs are largely

comprised of LP-like cells and suggests that hyperactivation of NOTCH and MYC

signaling might be relevant in tumorigenesis from LP-cells (Fig 7F). Mouse studies have

shown that Notch signaling is important in the maintenance of luminal lineage of the

normal mammary gland (54–57) and overactivation of Notch in luminal progenitors

resulted in hyperplasia and acquisition of self-renewal properties (54). Notch activation

drives the luminal fate specificity in normal mammary gland and can reprogram committed

basal mammary cells to an ER- luminal cell fate via multipotent embryonic cell states

(55,58), suggesting a role for Notch in fate specification and cellular plasticity. While our

data provides strong support for hyperactivation of NOTCH signaling in LP-like cells in

TNBCs it remains to be tested whether these cells arose from activation of NOTCH in

normal LP cells or from reprograming of normal basal cells. Our data suggest that

activation of MYC along with NOTCH might also have a role in this transformation. Given

the complementarity of the NOTCH and MYC pathways, further studies are needed to

gain better resolution of this mechanism. The PDO system, thus, can be exploited for

identification of dysregulated pathways in cancers which can then be perturbed in the

normal PDOs to understand aspects of the origins of cancers.

A detailed analysis of the cell type heterogeneity of TNBC organoids also revealed

the presence of MYC/NOTCH-activity low and hypoxia high cells amongst various other

cancer relevant cell types. Recent studies have shown that cancer cells can enter a MYC-

low diapause-like state of dormancy upon chemotherapy treatment and result in therapy

escape (59,60). Furthermore, single cell analysis from TNBC tumors with residual disease

after chemotherapy treatment showed an enrichment of hypoxia, angiogenesis, EMT and

ECM degradation related genes in the persistent tumors (61). We hypothesize that in

PDOs with these expression signatures, the cells that exhibit a MYC-low/hypoxia-high

profile are the likely candidates for chemotherapy escape (Fig 7F). Additionally, fate

mapping and other studies have shown that tumor cells exposed to hypoxia have a higher

tendency to result in metastasis (62,63). It remains to be tested to what extent these MYC-

low/hypoxia-high cells contribute towards metastasis and/or chemotherapy resistance,

and these patient-derived organoids will be an invaluable tool to answer these questions.

In summary, we developed a diverse biobank of BC organoids with a focus on

TNBC-derived organoids. We have thoroughly characterized these models as valid

systems that mimic the various aspects of patients' tumors, including genomic

alternations, transcriptomic signatures, cell type specificity and morphological

characteristics. Comparison of TNBC and normal-derived organoids provides important

insights into the mechanisms regulating tumorigenesis that can then be validated by

perturbation in normal PDOs (21). This comparison can also be used to identify novel

tumor specific targets that may play an important role in tumor growth and progression

(64). These next generation cancer models and the data derived from them offer vast

utilities and can be used for drug-screens, co-culture experiments, metabolomics, and

fate mapping studies to better understand the mechanisms driving cancers and for

identifying more specific treatment options.

Acknowledgements

We would like to thank members of the Spector lab for critical discussions and

advice throughout the course of this study. We thank Bodu Liu for critical review of this

manuscript. We also thank the various members of NCI-Human Cancer Model Initiative

(HCMI) for general advice and inputs on organoid cultures. We acknowledge the CSHL

Cancer Center Shared Resources (Animal, Flow Cytometry, Histology, Microscopy,

Organoid, Next-Gen Sequencing, and Single Cell Biology) for services and technical

expertise (NCI 2P3OCA45508). The sequencing analysis was performed using

equipment purchased through NIH grant S10OD028632-01. We thank Dr. Zihua Wang

for advice and a protocol for SMASH library preparation, Dr. Qing Gao for assistance with

histological sample preparation and optimization, Jill Habel for assistance with mouse

experiments. We are grateful for the support of Dr. Tawfiqui Bhuiya (Northwell Health) for

tumor tissue sections and regret his untimely passing. We deeply appreciate the efforts

of the Northwell Health Biobanking team for sample acquisition and the patients and their

families who donated tissues for research. This research was supported by

CSHL/Northwell Health support (D.L.S.), NCI 5P01CA013106-Project 3 (D.L.S.), Leidos

Biomedical HHSN26100008 (D.A.T, D.L.S), and the Manhasset Women's Coalition

Against Breast Cancer (S.B.). This research was also supported by NCI Breast SPORE

program P50-CA58223 (C.M.P), and U01CA238475-01 (C.M.P).

References

1. Parker JS, Mullins M, Cheang MCU, Leung S, Voduc D, Vickery T, et al. Supervised risk predictor of breast cancer based on intrinsic subtypes. Journal of clinical oncology : official journal of the American Society of Clinical Oncology. 2009 Mar;27(8):1160–7.

2. Perou C, Sørlie T, Eisen M, Rijn MV, Jeffrey S, Rees C, et al. Molecular portraits of human breast tumours. Nature. 2000;406:747–52.

3. Sørlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, et al. Repeated observation of breast tumor subtypes in independent gene expression data sets. Proceedings of the National Academy of Sciences of the United States of America. 2003;100:8418–23.

4. Prat A, Perou CM. Deconstructing the molecular portraits of breast cancer. Molecular Oncology. 2011 Feb;5(1):5–23.

5. Herschkowitz J, Simin K, Weigman VJ, Mikaelian I, Usary J, Hu Z, et al. Identification of conserved gene expression features between murine mammary carcinoma models and human breast tumors. Genome Biology. 2006;8:R76–R76.

6. Prat A, Parker JS, Karginova O, Fan C, Livasy C, Herschkowitz JI, et al. Phenotypic and molecular characterization of the claudin-low intrinsic subtype of breast cancer. Breast Cancer Research. 2010 Sep 2;12(5):R68.

7. Lehmann BD, Pietenpol JA, Tan AR. Triple-Negative Breast Cancer: Molecular Subtypes and New Targets for Therapy. American Society of Clinical Oncology Educational Book. 2015;(35):e31–9.

8. Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, et al. Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. The Journal of clinical investigation. 2011 Jul;121(7):2750–67.

9. Howlader N, Altekruse SF, Li CI, Chen VW, Clarke CA, Ries LAG, et al. US incidence of breast cancer subtypes defined by joint hormone receptor and HER2 status. Journal of the National Cancer Institute. 2014 Apr;106(5).

10. Curtis C, Shah SP, Chin S-F, Turashvili G, Rueda OM, Dunning MJ, et al. The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. Nature. 2012 Jun;486(7403):346–52.

11. Liedtke C, Mazouni C, Hess KR, Andre F, Tordai A, Mejia JA, et al. Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. Journal of clinical oncology : official journal of the American Society of Clinical Oncology. 2008 Mar;26(8):1275–81.

12. Rueda OM, Sammut S-J, Seoane JA, Chin S-F, Caswell-Jin JL, Callari M, et al. Dynamics of breast-cancer relapse reveal late-recurring ER-positive genomic subgroups. Nature. 2019 Mar;567(7748):399–404.

13. Fujii M, Sato T. Somatic cell-derived organoids as prototypes of human epithelial tissues and diseases. Nature Materials. 2021 Feb;20(2):156–69.

14. Tuveson D, Clevers H. Cancer modeling meets human organoid technology. Science. 2019 Jun 7;364(6444):952–5.

15. Sato T, Stange DE, Ferrante M, Vries RGJ, van Es JH, van den Brink S, et al. Long-term Expansion of Epithelial Organoids From Human Colon, Adenoma, Adenocarcinoma, and Barrett's Epithelium. Gastroenterology. 2011 Nov 1;141(5):1762–72.

16. Boj SF, Hwang C-I, Baker LA, Chio IIC, Engle DD, Corbo V, et al. Organoid Models of Human and Mouse Ductal Pancreatic Cancer. Cell. 2015 Jan 15;160(1):324–38.

17. Hoffmann K, Berger H, Kulbe H, Thillainadarasan S, Mollenkopf H-J, Zemojtel T, et al. Stable expansion of high-grade serous ovarian cancer organoids requires a low-Wnt environment. The EMBO journal. 2020 Mar;39(6):e104013–e104013.

18. Kopper O, de Witte CJ, Lohmussaar K, Valle-Inclan JE, Hami N, Kester L, et al. An organoid platform for ovarian cancer captures intra- and interpatient heterogeneity. Nature medicine. 2019 May;25(5):838–49.

19. Beshiri ML, Tice CM, Tran C, Nguyen HM, Sowalsky AG, Agarwal S, et al. A PDX/Organoid Biobank of Advanced Prostate Cancers Captures Genomic and Phenotypic Heterogeneity for Disease Modeling and Therapeutic Screening. Clinical cancer research : an official journal of the American Association for Cancer Research. 2018 Sep;24(17):4332–45.

20. Sachs N, de Ligt J, Kopper O, Gogola E, Bounova G, Weeber F, et al. A Living Biobank of Breast Cancer Organoids Captures Disease Heterogeneity. Cell. 2018 Jan;172(1–2):373-386.e10.

21. Dekkers JF, Whittle JR, Vaillant F, Chen H-R, Dawson C, Liu K, et al. Modeling Breast Cancer Using CRISPR-Cas9-Mediated Engineering of Human Breast Organoids. Journal of the National Cancer Institute. 2020 May;112(5):540–4.

22. Rosenbluth JM, Schackmann RCJ, Gray GK, Selfors LM, Li CM-C, Boedicker M, et al. Organoid cultures from normal and cancer-prone human breast tissues preserve complex epithelial lineages. Nature Communications. 2020 Dec;11(1):1711–1711.

23. Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature. 2012 Mar;483(7391):603–7.

24. Garnett MJ, Edelman EJ, Heidorn SJ, Greenman CD, Dastur A, Lau KW, et al. Systematic identification of genomic markers of drug sensitivity in cancer cells. Nature. 2012 Mar;483(7391):570–5.

25. Yu C, Mannan AM, Yvone GM, Ross KN, Zhang Y-L, Marton MA, et al. High-throughput identification of genotype-specific cancer vulnerabilities in mixtures of barcoded tumor cell lines. Nature Biotechnology. 2016 Apr;34(4):419–23.

26. Koboldt DC, Fulton RS, McLellan MD, Schmidt H, Kalicki-Veizer J, McMichael JF, et al. Comprehensive molecular portraits of human breast tumours. Nature. 2012;490(7418):61–70.

27. Nik-Zainal S, Davies H, Staaf J, Ramakrishna M, Glodzik D, Zou X, et al. Landscape of somatic mutations in 560 breast cancer whole-genome sequences. Nature. 2016 Jun;534(7605):47–54.

28. Shah SP, Roth A, Goya R, Oloumi A, Ha G, Zhao Y, et al. The clonal and mutational evolution spectrum of primary triple-negative breast cancers. Nature. 2012;486(7403):395–9.

29. Perou CM. Molecular Stratification of Triple-Negative Breast Cancers. The Oncologist. 2011 Jan;16(S1):61–70.

30. Prakash O, Hossain F, Danos D, Lassak A, Scribner R, Miele L. Racial Disparities in Triple Negative Breast Cancer: A Review of the Role of Biologic and Non-biologic Factors. Frontiers in Public Health. 2020;8:762–762.

31. Fusco N, Geyer FC, De Filippo MR, Martelotto LG, Ng CKY, Piscuoglio S, et al. Genetic events in the progression of adenoid cystic carcinoma of the breast to high-grade triple-negative breast cancer. Modern pathology : an official journal of the United States and Canadian Academy of Pathology, Inc. 2016 Nov;29(11):1292–305.

32. Wang Z, Andrews P, Kendall J, Ma B, Hakker I, Rodgers L, et al. SMASH, a fragmentation and sequencing method for genomic copy number analysis. Genome research. 2016 Jun;26(6):844–51.

33. Weigman VJ, Chao H-H, Shabalin AA, He X, Parker JS, Nordgard SH, et al. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. Breast Cancer Res Treat. 2012 Jun;133(3):865–80.

34. Gong Y, Zack TI, Morris LGT, Lin K, Hukkelhoven E, Raheja R, et al. Pan-cancer genetic analysis identifies PARK2 as a master regulator of G1/S cyclins. Nature genetics. 2014 Jun;46(6):588–94.

35. Wilczak W, Rashed S, Hube-Magg C, Kluth M, Simon R, Büscheck F, et al. Up-regulation of mismatch repair genes MSH6, PMS2 and MLH1 parallels development of genetic instability

and is linked to tumor aggressiveness and early PSA recurrence in prostate cancer. Carcinogenesis. 2017 Jan 1;38(1):19–27.

36. Chakraborty U, Dinh TA, Alani E. Genomic Instability Promoted by Overexpression of Mismatch Repair Factors in Yeast: A Model for Understanding Cancer Progression. Genetics. 2018 Jun 1;209(2):439–56.

37. Fan C, Prat A, Parker JS, Liu Y, Carey LA, Troester MA, et al. Building prognostic models for breast cancer patients using clinical variables and hundreds of gene expression signatures. BMC Medical Genomics. 2011 Dec;4(1):3–3.

38. Garcia-Recio S, Thennavan A, East MP, Parker JS, Cejalvo JM, Garay JP, et al. FGFR4 regulates tumor subtype differentiation in luminal breast cancer and metastatic disease. Journal of Clinical Investigation. 2020 Aug;130(9):4871–87.

39. Gatza ML, Silva GO, Parker JS, Fan C, Perou CM. An integrated genomics approach identifies drivers of proliferation in luminal-subtype human breast cancer. Nature Genetics. 2014 Oct;46(10):1051–9.

40. Lim E, Vaillant F, Wu D, Forrest NC, Pal B, Hart AH, et al. Aberrant luminal progenitors as the candidate target population for basal tumor development in BRCA1 mutation carriers. Nature Medicine. 2009 Aug;15(8):907–13.

41. Hu Z, Fan C, Livasy C, He X, Oh DS, Ewend MG, et al. A compact VEGF signature associated with distant metastases and poor outcomes. BMC Medicine. 2009 Mar;7(1):9–9.

42. Giraddi RR, Chung C-Y, Heinz RE, Balcioglu O, Novotny M, Trejo CL, et al. Single-Cell Transcriptomes Distinguish Stem Cell State Changes and Lineage Specification Programs in Early Mammary Gland Development. Cell reports. 2018 Aug;24(6):1653-1666.e7.

43. Davies BR, Greenwood H, Dudley P, Crafter C, Yu D-H, Zhang J, et al. Preclinical Pharmacology of AZD5363, an Inhibitor of AKT: Pharmacodynamics, Antitumor Activity, and Correlation of Monotherapy Activity with Genetic Background. Mol Cancer Ther. 2012 Apr 1;11(4):873–87.

44. Marchiò C, Weigelt B, Reis-Filho JS. Adenoid cystic carcinomas of the breast and salivary glands (or "The strange case of Dr Jekyll and Mr Hyde" of exocrine gland carcinomas). Journal of clinical pathology. 2010 Mar;63(3):220–8.

45. Weigelt B, Horlings H, Kreike B, Hayes M, Hauptmann M, Wessels L, et al. Refinement of breast cancer classification by molecular characterization of histological special types. The Journal of Pathology. 2008;216(2):141–50.

46. Raouf A, Zhao Y, To K, Stingl J, Delaney A, Barbara M, et al. Transcriptome Analysis of the Normal Human Mammary Cell Commitment and Differentiation Process. Cell Stem Cell. 2008 Jul 3;3(1):109–18.

47. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al. Comprehensive Integration of Single-Cell Data. Cell. 2019 Jun 13;177(7):1888-1902.e21.

48. Molyneux G, Geyer FC, Magnay F-A, McCarthy A, Kendrick H, Natrajan R, et al. BRCA1 Basal-like Breast Cancers Originate from Luminal Epithelial Progenitors and Not from Basal Stem Cells. Cell Stem Cell. 2010 Sep 3;7(3):403–17.

49. Nguyen QH, Pervolarakis N, Blake K, Ma D, Davis RT, James N, et al. Profiling human breast epithelial cells using single cell RNA sequencing identifies cell diversity. Nature Communications. 2018 May 23;9(1):2028.

50. Han H, Jain AD, Truica MI, Izquierdo-Ferrer J, Anker JF, Lysy B, et al. Small-Molecule MYC Inhibitors Suppress Tumor Growth and Enhance Immunotherapy. Cancer cell. 2019 Nov;36(5):483-497.e15.

51. Bruna A, Rueda OM, Greenwood W, Batra AS, Callari M, Batra RN, et al. A Biobank of Breast Cancer Explants with Preserved Intra-tumor Heterogeneity to Screen Anticancer Compounds. Cell. 2016 Sep;167(1):260-274.e22.

52. DeRose YS, Wang G, Lin Y-C, Bernard PS, Buys SS, Ebbert MTW, et al. Tumor grafts derived from women with breast cancer authentically reflect tumor pathology, growth, metastasis and disease outcomes. Nature Medicine. 2011 Nov;17(11):1514–20.

53. Guillen KP, Fujita M, Butterfield AJ, Scherer SD, Bailey MH, Chu Z, et al. A breast cancer patient-derived xenograft and organoid platform for drug discovery and precision oncology. bioRxiv. 2021 Mar 11;2021.02.28.433268.

54. Bouras T, Pal B, Vaillant F, Harburg G, Asselin-Labat M-L, Oakes SR, et al. Notch Signaling Regulates Mammary Stem Cell Function and Luminal Cell-Fate Commitment. Cell Stem Cell. 2008 Oct;3(4):429–41.

55. Lilja AM, Rodilla V, Huyghe M, Hannezo E, Landragin C, Renaud O, et al. Clonal analysis of Notch1-expressing cells reveals the existence of unipotent stem cells that retain long-term plasticity in the embryonic mammary gland. Nature Cell Biology. 2018 Jun;20(6):677–87.

56. Rodilla V, Dasti A, Huyghe M, Lafkas D, Laurent C, Reyal F, et al. Luminal Progenitors Restrict Their Lineage Potential during Mammary Gland Development. PLOS Biology. 2015 Feb 17;13(2):e1002069.

57. Yalcin-Ozuysal Ö, Fiche M, Guitierrez M, Wagner K-U, Raffoul W, Brisken C. Antagonistic roles of Notch and p63 in controlling mammary epithelial cell fates. Cell Death & Differentiation. 2010 Oct;17(10):1600–12.

58. Bland P, Howard BA. Mammary lineage restriction in development. Nature Cell Biology. 2018 Jun;20(6):637–9.

59. Dhimolea E, de Matos Simoes R, Kansara D, Al'Khafaji A, Bouyssou J, Weng X, et al. An Embryonic Diapause-like Adaptation with Suppressed Myc Activity Enables Tumor Treatment Persistence. Cancer Cell. 2021 Feb;39(2):240-256.e11.

60. Rehman SK, Haynes J, Collignon E, Brown KR, Wang Y, Nixon AML, et al. Colorectal Cancer Cells Enter a Diapause-like DTP State to Survive Chemotherapy. Cell. 2021 Jan;184(1):226-242.e21.

61. Kim C, Gao R, Sei E, Brandt R, Hartman J, Hatschek T, et al. Chemoresistance Evolution in Triple-Negative Breast Cancer Delineated by Single-Cell Sequencing. Cell. 2018;1–15.

62. Godet I, Shin YJ, Ju JA, Ye IC, Wang G, Gilkes DM. Fate-mapping post-hypoxic tumor cells reveals a ROS-resistant phenotype that promotes metastasis. Nature Communications. 2019;10(1):4862–4862.

63. Zhao Y, Fu X, Lopez JI, Rowan A, Au L, Fendler A, et al. Selection of metastasis competent subclones in the tumour interior. Nature Ecology & Evolution. 2021 May 17;1–13.

64. Chang K-C, Diermeier SD, Yu AT, Brine LD, Russo S, Bhatia S, et al. MaTAR25 lncRNA regulates the Tensin1 gene to impact breast cancer progression. Nature Communications. 2020 Dec;11(1):6438–6438.

65. Chen R, Im H, Snyder M. Whole-Exome Enrichment with the Roche NimbleGen SeqCap EZ Exome Library SR Platform. Cold Spring Harb Protoc. 2015 Jul 1;2015(7):pdb.prot084855.

66. Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform. Bioinformatics. 2009 Jul 15;25(14):1754–60.

67. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078–9.

68. Barnett DW, Garrison EK, Quinlan AR, Strömberg MP, Marth GT. BamTools: a C++ API and toolkit for analyzing and managing BAM files. Bioinformatics. 2011 Jun 15;27(12):1691–2.

69. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2: Somatic mutation and copy number alteration discovery in cancer by exome sequencing. Genome Res. 2012 Mar 1;22(3):568–76.

70. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. Nucleic Acids Research. 2010 Sep 1;38(16):e164–e164.

71. Landrum MJ, Lee JM, Benson M, Brown GR, Chao C, Chitipiralla S, et al. ClinVar: improving access to variant interpretations and supporting evidence. Nucleic Acids Research. 2018 Jan 4;46(D1):D1062–7.

72. Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, et al. COSMIC: the Catalogue Of Somatic Mutations In Cancer. Nucleic Acids Research. 2019 Jan 8;47(D1):D941–7.

73. Ioannidis NM, Rothstein JH, Pejaver V, Middha S, McDonnell SK, Baheti S, et al. REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. The American Journal of Human Genetics. 2016 Oct 6;99(4):877–85.

74. Jagadeesh KA, Wenger AM, Berger MJ, Guturu H, Stenson PD, Cooper DN, et al. M-CAP eliminates a majority of variants of uncertain significance in clinical exomes at high sensitivity. Nature Genetics. 2016 Dec;48(12):1581–6.

75. Mayakonda A, Lin D-C, Assenov Y, Plass C, Koeffler HP. Maftools: efficient and comprehensive analysis of somatic variants in cancer. Genome Res. 2018 Nov;28(11):1747–56.

76. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics (Oxford, England). 2013 Jan;29(1):15–21.

77. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology. 2014 Dec 5;15(12):550.

78. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative genomics viewer. Nat Biotechnol. 2011 Jan;29(1):24–6.

79. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012 Jul;9(7):676–82.

80. Huang Y, McCarthy DJ, Stegle O. Vireo: Bayesian demultiplexing of pooled single-cell RNA-seq data without genotype reference. Genome Biology. 2019 Dec 13;20(1):273.

81. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. Nat Biotechnol. 2018 May;36(5):411–20.

82. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. PNAS. 2005 Oct 25;102(43):15545–50.

83. Chandriani S, Frengen E, Cowling VH, Pendergrass SA, Perou CM, Whitfield ML, et al. A core MYC gene expression signature is prominent in basal-like breast cancer but only partially overlaps the core serum response. PloS one. 2009 Aug;4(8):e6693–e6693.

Figure Legends:

Figure 1. Establishment and somatic variant profiling of the breast cancer organoid biobank. A.) Summary of cancer type of the various tumor tissues that were used to generate organoids. IDC: Invasive ductal carcinoma, ILC: Invasive lobular carcinoma, Met-Lym: lymph node metastasis, IMC: Invasive mucinous carcinoma, NR: no residual tumor seen, DCIS: Ductal carcinoma in-situ, other: see Table S1 **B.)** Histopathological subtypes of the tumor tissues, ER/PR: Estrogen receptor (ER) and/or Progesterone receptor (PR), NA: not assessed **C.)** Age at diagnosis of the various subgroups of patient tumor tissues **D.)** Subtype specific, self-identified racial and ethnic breakdown of the patients represented in this biobank **E.)** Pathogenic single nucleotide variants (SNVs) identified in putative cancer driver genes in patient-derived organoids. **F.)** Proportion of organoids with pathogenic SNVs identified. Pathogenic SNVs: SNVs called pathogenic by ClinVar, COSMIC or REVEL and MCAP scores from targeted gene-panel sequence (49 samples) or whole exome sequencing (1 sample) (see Table S1), limited cultures: cultures where organoids were established at early passages (p0-p1) but could not be propagated, NA: not assessed

Figure 2. Copy number alterations (CNAs) enriched in the organoid models. A.) Copy number profiles, from IGV, of the various ER+ and ER- tumor derived organoids, along with the summary of overall copy number alterations across all samples. Side panel shows the pathogenic SNVs identified in that sample from Fig 1E. **B.)** Copy number amplifications or deletions identified in putative cancer driver genes (from Fig 1E). **C.)**

Copy number across different passages of three different sets of patient-derived organoids. **D.)** Magnified view of the chromosome regions in the red boxes in C.

Figure 3. A subset of TNBC organoids recapitulate signatures of aggressive basal-like breast cancers. A.) Molecular signatures associated with the different organoid lines. The sample legends are type: Normal= reductive mammoplasty derived normal organoids, Paired Normal= Adjacent or Distal to the tumor paired normal, Normal outgrowth= no pathogenic mutations were found, Luminal= ER/PR+ organoids; driver mutation: Other= trace mutations (see Fig 1E), None= no pathogenic mutations were found, NA= not assessed **B.)** Box-plots showing the breast cancer related and TNBC-specific scores for various gene signatures associated with poor outcomes. Each dot represents a different PDO; Luminal N=12, Normal N= 7, TNBC N= 19, TNBC met= 4. Differences in experimental groups were compared using Kruskal-Wallis test followed by pairwise comparisons using Wilcoxon rank-sum test. Bonferroni-Holm method was used to adjust the family-wise error (** adjusted p-value < 0.005, * adjusted p-value < 0.05) **C.)** Light microscopy images of the various TNBC- and normal (NM04N) derived organoid lines, along with Ki67-IHC, scale bars=100μm **D.)** Distribution of maximum passage numbers tested for the various TNBC PDOs. Long-term cultures (long) are defined by p>10 with continued expansion

Figure 4. TNBC Organoids can recapitulate tumor morphology and progression *in-vivo*. A.) Overview of PDO xenotransplant experiment using TNBC PDOs. **B.)** Box plots showing the end point tumor volume for the various organoid lines transplanted into the

fat-pads of NOD-SCID mice. Each dot represents tumor volume from one injection, N1= experiment 1, N2= experiment 2. ## For NH84T microscopic primary tumors observed in histology sections from 1/10 sites. ** For NH93T microscopic primary tumors observed in histology sections from 6/10 sites. TMN staging: pathologic TMN staging from patient pathology report (Table S1) **C.)** CNV profiles of the PDOs lines selected for *in vivo* transplant experiments **D.)** H&E images of the paired patient tumor tissue, TNBC patient-derived organoids (PDO) and xenografts generated form patient-derived organoids (PDO-X). S= Stroma, T= tumor, N=necrosis. Black arrows point to the cells with spinous connections with adjacent tumor cells. Yellow arrows point to the pseudo-lumen observed within AdCC-like breast cancers. Last column shows human mitochondria IHC in the lung and the liver from a representative mouse injected with the respective PDO (scale bar=100μm).

Figure 5. TNBC organoids are enriched in luminal progenitor-like cells. A.) Representative flow-cytometry plots for normal derived organoid (NM07NL) and various TNBC organoids stained for CD49f-PE on the x-axis and EPCAM-AF647 on the y-axis. The gates are subsets of live single cells within each organoid line and represent various cell types of the mammary epithelium. L=EPCAM-high mature luminal cells, LP= EPCAM+CD49f+ luminal progenitors, B= CD49f+ basal cells, S= stromal compartment. **B.)** Quantitation of the L, LP and B gates in panel B for multiple TNBC and normal organoid lines over multiple passages (see Table S7). Data-points are plotted as mean ±SEM using GraphPad Prism. NM0s: comprises multiple normal mammoplasty derived organoids from different patients **C.)** Flux of the epithelial cells during the early passages

of organoid derivation for normal distal (DS97ND) and TNBC tumor (DS97T) samples from the same patient **D.)** Copy number plots of the TNBC organoids DS97T over multiple passages **E.)** UMAP plot of batch corrected scRNA-seq data from 3 normal and 7 TNBC organoids. Numbers on the plot represent cluster IDs. **F. & G.)** UMAP plot in A but **F.)** separated by the normal and tumor samples and **G.**) colored by cell cycle. **H.)** UMAP plot in E split by individual tumor and normal samples and showing normalized expression of *EPCAM* and *ITGA6* (gene encoding CD49f) expression patterns in individual cells.

Figure 6. Tumor LP-like cells exhibit altered gene expression and have an upregulation of NOTCH and MYC downstream pathways. A.) UMAP plots for Integrated scRNA-seq data for all samples. The numbers indicate cluster IDs. **B.)** Marker expression of various cell type specific genes in the adult human breast epithelium (49). **C.)** Plots showing combined scores for the three mammary epithelial lineages: LP score: Luminal Progenitor score, Mature Lum score: Mature luminal score, MaSC score: Mammary stem cell score (40). Dashed region indicates LP clusters 2,7 and 11 that were used to perform differential expression analysis between normal and tumor LPs. **D.)** GSEA plots showing enrichment of the differentially expressed genes between normal and tumor LPs. **E.)** Violin plots showing combined expression in clusters 2,7 and 11 of the leading-edge NOTCH signaling genes and BMP2 target genes as identified in E. **F.)** Organoid formation from single cells. Significance was assessed by two-way ANOVA ns= not significant, ** pvalue<0.005, * pvalue<0.05

Figure 7. TNBC Organoids are comprised of heterogenous cell populations. A.) UMAP plot of TNBC only integrated scRNA-seq data showing clusters identified and cell cycle phases. **B.)** Distribution of cells in each of the G1 clusters identified per organoid line. **C.)** Dot-plot showing the marker genes for each of the G1 clusters and the associated phenotypic identity of that cell cluster. **D.)** Enrichment scores from GSEA of each of the G1 clusters that showed strong enrichment of some specific pathways or phenotypes. Enrichment score is represented by -10*NES*padj.value. **E**.) Combined gene set scores for the various phenotypes. Top panel: green= MYC signature, pink= Hypoxia signature. Bottom panel: green= Basal mammary stem cell (SC) signature, pink= Hypoxia signature. White represents positive correlation of the two signatures. **F.)** Schematic (created using BioRender.com) showing the cellular composition and heterogeneity observed in normal vs TNBC PDOs when cultured in matrigel. TNBC PDOs retain the tumor SNV/CNA profiles, are largely comprised of LP-like cells that might have originated from normal LP cells by the hyperactivation of NOTCH/MYC pathways. TNBC PDOs also have cells with signatures of hypoxia which is anti-correlated with NOTCH/MYC and positively associated with basal mammary stem cell signatures.

Figure 1

Figure 2

Figure 3

A. Normal PDOs | LT-TNBC PDOs — group, type, driver mutation

Breast Cancer Gene Expression Signatures:
- Immune + Stromal signatures
- IFN
- Basal BC (LP high)
- Normal MaSC
- Luminal
- LP low
- Angiogenesis/Hypoxia
- Proliferation

Color scale: 3 / -3

group
- Group1
- Group2
- Group3
- Group4
- Group5
- Group6

driver mutation
- TP53, PIK3CA
- PIK3CA
- TP53
- APC
- MAP3K13
- Other
- None
- NA
- Normal

type
- Normal
- Paired Normal
- Normal outgrowth
- TumorTissue
- Luminal
- Her2
- TNBC
- TNBC-met

B. Breast Cancer Signatures
- Proliferation
- UNC Basal
- TCGA Luminal

Signatures Associated with Poor Outcomes
- GSEA MYC signature
- VEGF signature 13genes
- GSEA Hypoxia & Glycolysis
- fMaSC Metabolism

(x-axis categories: Luminal, Normal, TNBC, TNBC Met)

C.
NH84T TNBC | NH95T TNBC | NH64T TNBC | NH87T TNBC | HCM-CSHL-0655-C50 TNBC-met | NM04N Normal
Bright Field / Ki67-IHC

NH93T TNBC | NH85TSc TNBC | HCM-CSHL-0366-C50 TNBC | NH67T TNBC
Bright Field / Ki67-IHC

D. TNBC PDO passage
max passage # (y-axis 0–25); x-axis: short, long

Figure 4

Figure 5

A. NM07NL NH93T NH95T NH87T HCM-CSHL-0655-C50 NH85TSc HCM-CSHL-0366-C50 NH84T

EPCAM-AF647

CD49f-PE

B. Epcam+ Epcam+CD49f+ CD49f_high

% of live cells

C. d69 d83 d110

DS97ND DS97T

CD49f-PE

D. DS97T Copy Number Profiles

E. Combined UMAP plot

F. Normal TNBC

G. Cell Cycle Phase

H. TNBC PDOs Normal PDOs

HCM-CSHL-0366-C50 NH85TSc NH64T HCM-CSHL-0655-C50 lymph-met NH87T 1° tumor NH95T NH93T NH87ND NM04N1 NM05N

EPCAM

ITGA6

Paired PDOs

Figure 6

Figure 7