Acetylation regulates the oligomerization state and activity of RNase J, the major ribonuclease of *Helicobacter pylori*

Alejandro TEJADA-ARRANZ[1,2,+], Maxime BOUILLOUX-LAFONT[1], Xue-Yuan PEI[3], Thibaut DOUCHÉ[4], Mariette MATONDO[4], Allison WILLIAMS[5], Bertrand RAYNAL[6], Ben F LUISI[3] and Hilde DE REUSE[1*]

[1] Unité Pathogenèse de *Helicobacter*, CNRS UMR2001, Département de Microbiologie, Institut Pasteur, Paris, France.

[2] Université de Paris, Sorbonne Paris Cité, Paris, France.

[3] Department of Biochemistry, University of Cambridge, Tennis Court Road, Cambridge CB2 1GA, UK.

[4] Plateforme Protéomique, Unité de Spectrométrie de Masse pour la Biologie, C2RT, USR CNRS 2000, Institut Pasteur, Paris, France.

[5] University of California San Francisco, Cellular Molecular Pharmacology School, School of Medicine, San Francisco CA, USA

[6] Plateforme de biophysique moléculaire, Département de Biologie structurale et chimie, Institut Pasteur, Paris, France.

+ Present address: Biozentrum, University of Basel, Switzerland

*Corresponding author: hdereuse@pasteur.fr

Institut Pasteur, Unité Pathogenèse de *Helicobacter*

28, rue du Docteur Roux

75724 PARIS Cedex 15 FRANCE

Tel: +33 1 40 61 36 41

Abstract

In *Helicobacter pylori*, post-transcriptional regulation strongly relies on the activity of an RNA degradosome, composed of the essential ribonuclease RNase J and the DEAD-box RNA helicase RhpA. Here, we describe post-translational modifications of this protein complex that affect its activity. Cell-extracted RNase J is acetylated on multiple residues, one of which, K649, strongly impacts RNase J oligomerization, which in turn influences ribonuclease activity. Corroborating the link between oligomerization and activity, mutations targeting K649 and other residues affect the dimerization and *in vitro* activity of RNase J. Our crystal structure of RNase J reveals three loops that gate access to the active site and rationalizes how oligomerization state influences activity. The acetylated residues of RNase J are important for *H. pylori* morphology, highlighting that the modifications affect the RNase J cellular function. We propose acetylation as a regulatory level controlling the activity of RNase J and the *H. pylori* RNA degradosome.

Key words: post-transcriptional regulation, RNA degradosome, RNase J, *Helicobacter pylori*, post-translational modifications, protein acetylation.

Introduction

Post-transcriptional regulation is one of the most important levels of control of gene expression in every kingdom of life. Ribonucleases (RNases) are key enzymes in this regulation, that are involved in RNA maturation and degradation. In several bacteria, major RNases act in concert with DEAD-box RNA helicases, that help to unfold RNA secondary structures, within protein machines called RNA degradosomes[1].

We previously demonstrated the existence of a minimal RNA degradosome in *Helicobacter pylori*[2]. This Gram-negative bacterial pathogen chronically colonizes the stomach of half of the human population worldwide. Infection leads to chronic gastritis and, in some cases, to further gastric pathologies such as peptic ulcers, MALT lymphoma, or gastric adenocarcinoma that causes 800,000 deaths each year worldwide[3]. *H. pylori* possesses a small genome (1.6 Mb) with few transcriptional regulators, and several lines of evidence indicate that post-transcriptional regulation plays a major role in gene expression control in this organism[4–6]. The *H. pylori* RNA degradosome is composed of RNase J, a 77.6 kDa enzyme that possesses both 5′-3′ exoribonuclease and endoribonuclease activities, and RhpA, a DEAD-box RNA helicase, with a molecular weight of 55.8 kDa[2,7]. RNase J is essential in *H. pylori* and plays a major role in the degradation of both mRNAs and antisense RNAs [8]. RhpA is the sole DEAD-box RNA helicase encoded by the *H. pylori* genome and is essential for colonization in the mouse infection model[7]. Furthermore, we recently demonstrated that, in *H. pylori*, the RNA degradosome is compartmentalized into clusters at the inner membrane whose formation is regulated and likely represents a major level of control of its activity[9].

RNase J-encoding genes are present in 57 % of sequenced bacteria[1],

including the Gram-positive organisms *Bacillus subtilis*, *Staphylococcus aureus*

and *Deinococcus radiodurans*. *B. subtilis* and *S. aureus* encode two paralogues

of RNase J, called J1 and J2 while *H. pylori, Streptomyces pyogenes* and *D.*

radiodurans possess only one gene for this protein. Compared to the RNase J

proteins of these three organisms, the *H. pylori* enzyme possesses a unique N-

terminal extension of 132 residues that is predicted to be predominantly

disordered[2,9]. In *B. subtilis*, while both RNase J paralogues (*Bsu*RNases J1 and

J2) have equivalent endoribonuclease activities, *Bsu*RNase J2 seems to lack

the 5'-3' exoribonuclease activity that predominates in *Bsu*RNase J1[10].

Furthermore, these two proteins interact and form a heterotetramer *in vitro* that

has different cleavage specificity *in vitro* than each protein alone[10]. *In vitro*

analysis of the RNase J1 enzyme from *S. aureus* showed that it predominantly

assembles into dimers, but also forms tetramers, independently of divalent

cations[11]. The single RNase J from *S. pyogenes* forms dimers and tetramers *in*

vitro without divalent cations being observed in the dimerization surface[12]. In *D.*

radiodurans, RNase J forms a dimer in a Mn-dependent fashion *in vitro*, and the

interaction between the two monomers is mediated by the C-terminal domain of

*Dra*RNase J[13]. Several crystal structures of RNase J proteins have been

published[12–14], where RNase J is generally found to be a tetramer (dimer of

dimers). However, it is unknown whether the different oligomeric states of

RNase J detected *in vitro* have differences in activity.

Almost nothing is known about post-translational modifications of these

protein complexes and whether this might influence their activity. Acetylation

has emerged as a major post-translational regulatory mechanism in

bacteria[15,16]. Although the acetylation of lysine, cysteine, serine and threonine

residues has been reported, lysine acetylation is the most prominent

modification, particularly on its ε-amino group[15]. This modification has been

shown to regulate multiple properties of the target proteins, including enzymatic

activity, DNA-binding capability or stability, among others[16]. Interestingly, some

proteins involved in RNA metabolism have been reported to be lysine

acetylated, including (i) *Escherichia coli* RNase R, whose acetylation regulates

its stability[17]; (ii) *E. coli* RNase II, regulated at the activity level[18]; and (iii) the

CshA DEAD-box RNA helicase from *B. subtilis*, where acetylation has been

proposed to influence its interaction with RNA polymerase independently from

its association with RNase Y, the major ribonuclease of the *B. subtilis* RNA

degradosome[19].

The central importance of RNase J in post-transcriptional regulation in *H.*

pylori and its participation in an RNase J-based degradosome prompted us to

investigate the structure, oligomeric state and post-translational regulation of

the *H. pylori* RNase J and to assess its stoichiometry in the RNA degradosome.

Here, we report the crystal structure of *H. pylori* RNase J without its N-terminal

extension. We also show that RNase J is in equilibrium between monomeric

and dimeric states *in vitro* and that, together with its monomeric RNA helicase

partner RhpA, assembles into a minimal degradosome complex with a 1:1

stoichiometry. Our data suggest that the dimeric form constitutes the active form

of RNase J *in vitro*. Furthermore, we found for the first time that RNase J is

acetylated at several residues by multiple mechanisms. We concluded that the

acetylation state regulates the oligomerization state and activity of RNase J *in*

vitro and its activity in *H. pylori*, and thereby constitutes an important level of

control of the RNA degradosome.

Results

Structure of the *H. pylori* RNase J

To better understand the structure-function relationship of the *H. pylori*

RNase J protein, we first decided to determine its crystal structure. Several

attempts to purify sufficient amounts of full-length RNase J from *H. pylori* to

obtain crystals were unsuccessful. As we previously reported[9], the N-terminal

part of RNase J (first 120 amino acids) that is specific of the *Helicobacter*

species is predicted to be an intrinsically disordered region (IDR). After

truncating this region, well diffracting crystals were obtained from this RNase J

form comprising residues 137-691 (from *H. pylori* strain 26695, uniprot P56185).

The crystal structure of *H. pylori* RNase J was solved at 2.75 Å by molecular

replacement using the *Streptomyces coelicolor* homologue as a search model

(Table S1). One molecule occupies the asymmetric unit of the crystal, and

through crystal symmetry a homotetramer can be generated that is organized

as a dimer-of-dimers (Figure 1A). The core region of the enzyme, residues 137-

586, encompasses the beta-CASP and beta-lactamase domains that define this

family[20], but there are no detectable metal cofactors in the map at the catalytic

site. An X-ray Zn-peak scan also confirmed that there are no detectable Zn ions

in the crystal.

The structure predicted by the Robetta server, based on machine learning[21],

is in excellent agreement with the X-ray model for the core, with the notable

exception of the three loop regions 174-183, 431-441 and 575-586 at the

entrance to the active site, for which the predicted loop positions are closer to

those seen in the crystal structure of the *S. coelicolor* RNase J (Figure 1B-C).

The electron density over these regions is well defined in the *H. pylori* map and
modelled with high confidence. Comparing the structures, it is apparent that the
position of the loops will enforce a different path for the RNA substrate in *H.*
pylori RNase J compared with the *S. coelicolor* enzyme (Figure 1C).

The C-terminal domain (CDT), from residues 587 to 691, is poorly ordered in
the crystal lattice and the density was not sufficiently well resolved to model the
polypeptide. The Robetta model for the C-terminal domain fits well and fitted
into the map, showing good agreement with the predicted helical segments.
The predicted model is also in agreement with a high confidence homology
model prepared by PHYRE2 using the *Thermus thermophilus* homolog as
template (PDB 3BK2). This domain makes a self-complementary interaction
through crystal lattice contact. The Robetta model predicted with low confidence
an extended helical region of the N-terminal residues 1-136 that were not
included in the crystallized protein, and this region seems unlikely to have a
compact fold. The final model to fit the X-ray data is a fusion of the fitted crystal
structure of the core and the Robetta model for the C-terminal domain with side
chains removed due to poor resolution of this part of the electron density map.

In the dimeric unit, the protomer interfaces are formed by self-complementary
interactions made mostly by the CTD. The surface area buried at the CTD-CTD
interface is estimated to be 1165 $Å^2$. The beta-CASP and beta-lactamase
domains of RNase J also make a self-complementary interface in the principal
dimer, with an estimated area of 672 $Å^2$ that is a little more than half of the
CTD-CTD contribution. Stronger interactions typically have a surface area of
more than 2000 $Å^2$ [22]. The interface between the beta-CASP and beta-
lactamase domains is very similar to that seen in other dimers of RNase J, for

instance in the homologue of *S. coelicolor*[12]. The interfaces that are formed

between the dimer units to make the tetramer are also made by the beta-CASP

and beta-lactamase interfaces, and these have a total buried area of 2070 Å^2.

An overlay of RNase J of *H. pylori* and of *S. coelicolor*, where an RNA

substrate is bound, suggests that the path of the RNA to the active site cannot

be the same for the two enzymes, and that the nucleic acid would need to be

rerouted in the *H. pylori* enzyme. The obstructions are caused by exposed

peptide loop regions that are positioned to accommodate the CTD. It is possible

that changes in the oligomerization state will impact on the loops and access to

the active site.

It is notable that the CTD has extensive charged side chains on the surface

of proteins that likely form complementary charge interactions. Such an

organization may stabilize the fold of the small domain[23–25]. As the self-

complementary interactions of the C-terminal domain are a key component of

the assembly, any changes in the stability of the CTD may impact on the

oligomeric state. This prompted us to investigate the *in vitro* oligomeric state of

RNase J and its RhpA partner.

In vitro* analysis of the oligomeric state of the two partners of the *H. pylori

RNA degradosome

The crystal structure of *H. pylori* RNase J and of several other RNase J

proteins correspond to tetramers. However, for some of these proteins, other

oligomeric states (dimers and monomers) were found in solution and the activity

of the different forms was unclear. To evaluate the oligomeric state of *H. pylori*

RNase J and RhpA *in vitro*, we performed analytical ultracentrifugation (AUC)

with His-tagged versions of RNase J and RhpA purified from *E. coli* (Supp

Figure S1). The sedimentation coefficient distribution of RNase J indicates that

this protein forms two oligomeric species *in vitro* with concentration-independent

sedimentation coefficients (S_0) of 5.8 and 8.8 (Fig. 2A). Based on the structural

data of *H. pylori* RNase J presented above, the calculation of the sedimentation

coefficients with the HYDROPRO software was 5.4 and 8.9 for a monomer and

dimer, respectively. Moreover, the calculated sedimentation for the monomer

suggests that the C-terminal end of the monomer is probably flexible and less

extended in solution, explaining the small difference in sedimentation

coefficient. On the other hand, the calculated sedimentation for the dimer is fully

compatible with the extended arrangement of the C-terminal end, showing that

dimerization locks the structure in an extended conformation. Therefore, we

concluded that the two forms of RNase J observed by AUC are indeed a

monomer and a dimer (Figure 2A) that have a structural arrangement in solution

which is compatible with the observed crystal structure. The monomeric

population predominates (~60%) over the dimer (~40%) independently of the

protein concentration, showing that these species were not in equilibrium but

were dependent on other modifications.

AUC of RhpA shows behavior as a monomer independent of concentration

with an S_0 of 3.85 (Figure 2B).

RNase J and RhpA form a 1:1 minimal degradosome complex

Next, we wanted to evaluate the stoichiometry of the protein complex formed

by RNase J and RhpA. We mixed both proteins at 5:1, 1:1 and 1:5 ratios and

subjected them to AUC (Figure 2C). When using a 5:1 RNase J:RhpA ratio, we

identified three peaks corresponding to both proteins alone (RNase J monomer

and dimer and RhpA monomer). With a 1:1 RNase J:RhpA ratio, three peaks

could be detected. Based on the previous calculations, these peaks are

proposed to correspond to (i) RhpA monomer alone for the 3.85S peak, (ii) a

dimer of RNase J for the 8.8S peak, and (iii) a minor peak with a sedimentation

coefficient of 4.8S that would correspond to the RNase J-RhpA complex. This

latter peak was much more prominent when using a 1:5 ratio of RNase J:RhpA,

where only two peaks could be detected, corresponding to RhpA alone and

most probably the complex with a coefficient of 4.8S. From this peak and the

available frictional coefficients, we calculated a mass for the complex of

approximately 124 kDa that is only compatible with a complex with a 1:1

stoichiometry. Interestingly, neither the monomer nor the dimer of RNase J was

detected under this condition, and all RNase J was involved in complex

formation. We thus conclude that RNase J and RhpA form a 1:1 stoichiometry

complex *in vitro*.

Bacterial two hybrid (BACTH) to identify residues important for the

oligomerization of RNase J

As AUC results show that the formation of oligomer is not an equilibrium and

is independent of concentration, we then searched for the potential residues

involved in the self-interaction of RNase J using bacterial two hybrid (BACTH)

experiments that test the interaction of protein pairs in *E. coli*. First, a consistent

interaction of wild type RNase J with itself was detected (Figure 3A). Then,

different RNase J mutants were constructed and tested. As previously reported

for other RNase J proteins[13], we found that the C-terminal region of RNase J is

essential for this interaction. Next, in order to pinpoint residues that are

important for RNase J multimerization, we constructed RNase J mutants in

which nine charged residues of this C-terminal domain were individually

replaced by alanine residues. As shown in Figure 3A, several mutants were

significantly affected in their capacity to homo-oligomerize, including RNase J

E618A, E654A, K660A, E663A and K649A. These data show that charged

residues at the C-terminal domain of RNase J are important for its self-

interaction.

These residues are not at the predicted dimerization interface of the C-

terminal domain, but they are likely to be involved in stabilizing the domain

through interactions with side chains of complementary charge.

RNase J is acetylated at several lysine residues

RNase R and RNase II from *E. coli* have been reported to be post-

translationally modified by acetylation but no such modification was reported on

RNase J. A mass spectrometry (MS) analysis was performed on an RNase J-

FLAG fusion protein expressed from the *H. pylori* chromosome (Supp Figure

S1) under the control of its native promoter. The RNase J-FLAG protein was

pulled down from *H. pylori* crude lysates using anti-FLAG antibodies and

extracted from a polyacrylamide gel. The experiment was performed on three

biological replicates from cultures in exponential phase. We found that nine

lysine residues of RNase J, spread across all protein domains, are consistently

acetylated: K134, K140, K323, K337, K397, K511, K547, K634 and K649 (Supp

Figure S2 and Supp Table S2).

Lysine 649 is important for the oligomerization state of RNase J

Lysine 649 (K649), localized at the RNase J C-terminal domain, was particularly interesting since this residue was both acetylated and required for RNase J homo-oligomerization in two hybrid experiments. Therefore, AUC analyses were performed with purified RNase J variants (Supp Figure S1) to test the importance of K649 in RNase J homo-oligomerization. As shown in Figure 3B, the purified RNase J K649A mutant assembles exclusively as a dimer *in vitro*, with a complete loss of the monomeric population seen with the WT protein. Circular dichroism was performed to verify that this effect was not due to a defect in the folding of RNase J K649A. The circular dichroism spectra of wild-type and K649A variants of RNase J did not show significant differences (Supp Figure S3) and the estimated proportions of the different secondary structure elements were maintained (Supp Table S3). This shows that mutation K649A does not introduce a major folding defect in RNase J secondary structure. Thus, K649 is important for RNase J dimerization both *in vitro* and in *E. coli*.

Acetylation mimicry of Lysine 649 affects the oligomeric state of RNase J

Given that K649 is acetylated, we next investigated whether RNase J acetylation at this residue could represent a control level of its oligomeric state. Therefore, we constructed RNase J mutants in which K649 was either replaced by an arginine residue (K649R), that mimics a non-acetylated state but maintains a positive charge at this position, or replaced by a glutamine residue (K649Q) which mimics an acetylated state while neutralizing the positive charge [a previously validated strategy[26]]. As shown in Figure 3C, the AUC

sedimentation profile of RNase J K649R is similar to that of the wild type

enzyme. In contrast, RNase J K649Q shows a significantly increased proportion

of dimers over the monomers, similarly to the K649A mutant (Figure 3B-C).

These data show that the positive charge of K649 is important for the stability of

the RNase J monomeric state and that oligomerization is controlled by the

charge state of K649.

Mutations in K649 alter the exoribonuclease activity of RNase J

Since RNase J WT forms both monomers and dimers *in vitro*, we next

assessed whether the oligomeric state and the presence of a charge at position

649 were correlated with differences in the *in vitro* activity of this protein. The

exoribonuclease activity of RNase J was measured with a 24-nt long single

stranded RNA substrate from the 5'-untranslated region (UTR) of *rnj*, upon

which we previously found RNase J to be active[8]. This substrate was labeled

with 6-carboxyfluorescein (6FAM) at either the 3' or 5'-ends. As a control, a

mutant inactivated at the RNase J active site, D299A, was constructed. We

found, as expected, that the D299A inactive mutant did not degrade the

substrate while the wild type RNase J was active in its 5'-3' exoribonucleolytic

mode (Figure 4A-B and Table 1). This activity is evident from the rapid depletion

in the intact substrate bands labelled in 5' (Figure 4A) or 3' (Figure 4B) and,

from the additional accumulation of the 1nt product upon digestion of the 5'-

labeled substrate (Figure 4A).

The RNase J mutant with K649 replaced by R, that mimics a non-acetylated

state, and which behaves as monomers and dimers like the WT protein, was

strongly affected in its exoribonuclease activity (Figure 4 and Table 1). For the

K649R variant, the half-lives of the 3'- and 5'-labeled substrates increased by 5 and 10-fold, respectively (Table 1).

In contrast, the protein variants with K649 replaced either by A (that abolishes both the positive charge at this position and the acetylation) or by Q (acetylation-mimicking mutation) and that both mainly assembled into dimers, presented an activity that was significantly higher than that of the RNase J K649R variant. The half-lives of 3' and 5'-labeled substrates were 13 and 49-fold lower for the K649A variant and 2.5 and 4-fold lower for the K649Q variant, respectively. The half-lives of the substrates were calculated from three independent experiments for the wild type and for every RNase J variant and are shown in Table 1.

From these data, we conclude that, for these mutants, the dimeric form of RNase J is most probably the active form *in vitro*. Altogether, we identified a residue, K649, whose charge is important for RNase J activity and oligomerization and that is acetylated in *H. pylori* cells. This indicates a mechanism of control of RNase J activity that is based on the allosteric control of its dimerization by acetylation.

The activity of the RNase J-RhpA complex is not affected by K649 mutations

It is not known whether the oligomerization state of RNase J can affect the formation of the RNA degradosome complex. Therefore, we wanted to determine whether mutations of K649 affect the activity of the RNase J-RhpA complex. For this, we used a 45-nt long RNA substrate with hairpin structures at both ends to prevent the exoribonuclease activity from the 5'-extremity and

labeled it with 6FAM at the 3' end [as in [13], Supp Figure S4A]. First, the activity

of RNase J WT and mutant proteins alone was tested on this substrate. As

compared to the linear 3'-labeled substrate, the degradation kinetics of this

substrate by RNase J was significantly reduced but not abolished (half-life of

the substrate is >4 fold lower, Table 1), suggesting that RNase J might attack

this substrate by its endoribonuclease activity and then proceeds by

exoribonuclease activity. As we previously reported, in a functional

degradosome complex, RhpA will unwind the secondary structure elements and

allow RNase J to act in its exoribonucleolytic mode. Most interestingly, the

activity tests with a mixture of RNase J and RhpA in a 1:1 ratio resulted in a

significant decrease in the half-life of the substrate to 6.3±2.9 min as compared

to the activity of RNase J alone (approx. 31.4±17.6 min), compatible with a

functional complex (Table 1). The half-life of the structured substrate incubated

with the complex between RhpA and the RNase J K649A variant (that has

increased exoribonuclease activity) was not affected (half-life 6.7±0.5 min)

(Supp Figure S4B and Table 1). We conclude that the activity of the RNase J-

RhpA complex is not significantly affected by the K649A mutation.

374   Residue D185 is important for oligomeric state and the activity of RNase J

The RNase J structure suggests that residue D185 may indirectly support the

stability of the monomer, as this residue is in an interaction network that

supports a helical extension involving residues 584-594 that supports

interactions with the C-terminal domain. In agreement with our predictions, AUC

analysis of the RNase J D185A mutant revealed a much more predominant

dimeric population, as compared to the wild type protein (Figure 5). Consistently

with what we found for mutant K649A, this was correlated with higher

exoribonucleolytic activity of RNase J D185A (Figure 4A-B and Table 1),

suggesting again that the structural destabilization of the monomer leads to an

increased amount of dimer with an enhanced enzymatic activity.

Analysis of the consequences of RNase J mutations on *H. pylori*

morphology

We next wanted to examine the consequences of the mutations affecting

RNase J oligomerization and/or acetylation state in live *H. pylori* cells. Since

RNase J is an essential protein, strains carrying mutations in the acetylated

residues identified above were constructed with the following strategy. *H. pylori*

strain B128 was transformed with plasmid pILL2157 derivatives either

expressing wild type or mutant variants of RNase J under the control of an

IPTG-inducible promoter. Then, the chromosomal wild type copy of RNase J

was deleted by homologous recombination and replacement with a kanamycin

resistance cassette.

In *B. subtilis*, mutants deficient in RNase J1 present an altered cell

morphology[27]. Therefore, the morphology of our *H. pylori* RNase J mutant

strains was examined by phase contrast microscopy and quantified with the

MicrobeJ ImageJ plug-in[28]. The wild type B128 strain presents cells that are

relatively heterogeneous in shape and with an average length of 2.15±0.67 µm.

We observed that, in a strain expressing wild type RNase J from the system

described above, the *H. pylori* cells were longer in the absence of IPTG [lower

amount of RNase J as compared to the WT strain[8]], than upon induction with 1

mM IPTG (where the expression of RNase J is induced). The cell length was

2.44±0.61 µm without IPTG and 2.05±0.51 µm with IPTG (Figure 6). This

indicates that in *H. pylori*, RNase J is also regulating a factor required to

maintain wild type cell morphology. Interestingly, we noted that the observed

phenotype is not dependent on the activity of RNase J alone but rather on the

RNA degradosome complex, as IPTG induction of RNase J WT in a strain

lacking RhpA fails to complement this phenotype (average length of 3.41±0.83

µm) (Figure 6A).

When we performed these tests with strains that expressed different RNase

J variants, we identified two groups of mutations. In the first group, the RNase J

variants behaved similarly to the wild type enzyme, as their expression upon

IPTG addition resulted in a reduction of bacterial length. This was the case for

RNase J K323A, K547A and K649A and, to a lesser extent, RNase J K634A. In

the second group, that includes variants K134A, K140A, K337A, K397A and

K511A, the strains presented a strong growth defect and substantial cell lysis

and cellular debris were observed. In these cases, the cells were short with a

tendency towards a coccoid morphology that is typical of *H. pylori* cultures

under stress[29]. This phenotype was particularly evident in the case of cells

expressing variant K397A. For these cultures, addition of IPTG favored bacterial

growth but failed to restore the typical helical form of healthy *H. pylori* cells,

suggesting that RNase J function is strongly impaired.

These results show that RNase J activity is important for the correct

regulation of growth and cell morphology in *H. pylori* and that several of the

residues that we found to be acetylated are important for RNase J function *in*

vivo.

RNase J is acetylated through a range of mechanisms in *H. pylori*

Several pathways contribute to acetylation in bacteria[30]. We tried to elucidate which are the enzymes responsible for the different acetylation events that occur on the RNase J of *H. pylori*. Therefore, we used the *H. pylori* strain expressing an RNase J-FLAG fusion from the chromosome to introduce deletions in genes that we predicted as acetylase candidates. The RNase J-FLAG protein was enriched from these strains and analyzed by MS as explained above. The tests were performed from three biological replicates. The acetylase candidate genes were *hpb8_615* [henceforth referred to as *rimI*, as it presents homology to *rimI* from *E. coli*[31]] and *hpb8_1270* [that contains two Gcn5-acetyltransferase (GNAT) domains]. Non-enzymatic acetylation by acetyl-CoA and acetyl-phosphate also occurs in bacteria[30,32]. The impact of this pathway was assessed in *H. pylori* by constructing a mutant in the *pta-ackA* operon that is in charge of acetyl-phosphate synthesis and metabolism. As shown in Figure 7, we found that several pathways influence the acetylation of RNase J lysine residues. RimI is partially responsible for the acetylation of RNase J residues K323, K397 and K649. The HPB8_1270 protein is partially responsible for the acetylation of residues K323, K397, K511 and K649. Acetyl phosphate-mediated non-enzymatic acetylation participated in the acetylation of K134, K323, K397 and K511 and K649. For other residues, namely K140, K337, K547 and K634, no change in acetylation levels was detected in the mutants, suggesting that other acetylation mechanisms are likely at play in *H. pylori*. These results suggest a high redundancy of the different acetylation mechanisms that act on RNase J, particularly on residues K323, K397 and K649.

456   Thus, multiple mechanisms participate in the acetylation of RNase J on

457   different residues.

Discussion

In this study, we defined the structure of the major *H. pylori* ribonuclease RNase J and its *in vitro* oligomerization state alone and in the RNA degradosome with its RhpA RNA helicase partner. We also identify *in vivo* acetylation patterns of RNase J and demonstrate that these impact on oligomerization and activity *in vitro* and *in vivo*.

We found, by AUC, that RNase J is in equilibrium between monomeric and dimeric states. Earlier studies showed that *D. radiodurans* RNase J is also distributed between monomers and dimers[13]. *Bsu*RNase J1 forms dimers and tetramers *in vitro*[14] and, when mixed with *Bsu*RNase J2, heterotetramers[10]. Similarly, RNase J1 from *S. aureus* forms homodimers and homotetramers *in vitro*[33]. RNase J of *S. pyogenes* crystallizes as tetramers that are composed of a dimer-of-dimers[12]. In the present study, at the studied *H. pylori* RNase J concentrations, no trace of tetramerization could be seen by AUC. It is likely that the observed tetrameric structure in the X-ray structure is due to crystal packaging and that it does not reflect the arrangement of RNase J in solution, as the dimer seems to be the active form. In addition, RhpA alone behaves as a monomer in solution by AUC. Together, RNase J and RhpA form a complex that is compatible with a 1:1 stoichiometry.

Acetylation has emerged in recent years as a major post-translational regulatory mechanism in prokaryotes[16], and a few examples of its influence in RNases and RNA-degrading enzymes have been described, including *E. coli* RNase R[17] and RNase II[18] and *B. subtilis* CshA[19]. Here, we found that, in *H. pylori*, as many as 9 lysine residues of RNase J were consistently acetylated, although probably not at the same time, which constitutes the first evidence of

protein acetylation in *H. pylori*. One of these residues is K649, which we also

found to be involved in the oligomerization state of RNase J by BACTH in *E.*

coli. Interestingly, this residue is located at the C-terminal domain of RNase J

(Supp Figure S2). This region was also shown to be important for the

oligomerization and exoribonucleolytic activity of the RNase J of *D.*

radiodurans[13]. Therefore, we examined the role of K649 in RNase J

oligomerization (by AUC) and exoribonuclease activity. Three mutants were

analyzed, RNase J K649A that abolishes the side chain of the lysine residue,

K649R that mimics a non-acetylated state while maintaining the positive charge

and K649Q that mimics an acetylated state[26]. Whereas the K649R variant

maintains a sedimentation profile similar to that of the wild type enzyme,

variants lacking a positive charge in K649, either K649Q or K649A, had an

increased proportion of dimer or displayed no monomeric form, respectively.

Interestingly, we observed that the K649R variant is less active than the wild

type RNase J whereas variants K649Q and K649A are more active than the

K649R variant.

Our structural data indicated that residue K649 is not located at the

interaction surface between the subunits of the dimer, but that it rather plays a

structural role in the integrity of the overall structure. We conclude that

elimination of the positive charge at residue 649 leads to a higher stabilization

of the dimeric species. In agreement with this hypothesis, we observed that

D185A, another mutation in a structurally important residue, had the same

effect and favors the dimeric species. Interestingly, in *D. radiodurans*, the

corresponding residue (D61) had already been identified as key for RNase J

dimerization[13]. Electrostatic interactions of charged side chains on the surface

of proteins can strongly contribute to the fold stability[23–25]. We propose that the

network of surface electrostatic interactions stabilizes the CTD of RNase J,

which in turn supports the homodimer. The CTD is the primary self-

complementary interface of the dimer, and the interface between the core

(amino acids 137-586) and between the dimer-of-dimers formed by the core

both have marginal stability in isolation. Thus, changes in the charge, through

acetylation, can thereby impact on stability of the CTD fold and oligomerization

state.

The question naturally arises of how the catalytic activity of RNase J is

affected by its oligomerization state. A clue is suggested in the overlay of the

structure of *H. pylori* RNase J with the *S. coelicolor* homologue, in which there

is a bound RNA substrate engaged at the active site (Pei et al., 2015) (Figure

1D). There are marked differences in the positions of loops at the entrance to

the catalytic site that would obstruct the RNA from following the same route in

the *H. pylori* enzyme, but the nucleic acid could approach the site from the

opposite rim of the entrance. The exposed peptide loop regions are positioned

to accommodate the CTD. It is possible that changes in the oligomerization

state, through changes in CTD stability, will impact on the presentation of those

loops and access to the active site in a gating mechanism.

Altogether, our results suggest that mutations that favor the dimeric RNase J

conformation also increase the activity of RNase J of *H. pylori in vitro*. As the

charge of residue 649 is important for dimerization, acetylation of K649 might tilt

the balance of the RNase J species towards the dimeric form and potentially

increase its activity. In this way, acetylation on K649 can modulate the activity of

RNase J in *H. pylori* cells. Formation of the RNase J-RhpA complex is not

affected by mutations or modifications of residue K649 in the ribonuclease, and

thus such mutations do not significantly alter the activity of this complex. Earlier

studies indicate that RNase J and the RNase J-RhpA complex can act on

different transcripts[2,8,34]. It can be envisaged that acetylation of K649 would

therefore have a differential impact on the subset of free RNase J that is not

part of a degradosome assembly.

Furthermore, we addressed the role of the acetylated residues that we

identified in *H. pylori* cells. It has been described that *B. subtilis* cells lacking

RNase J1 show morphology defects with a tendency to filament[27]. We thus

evaluated the morphology of *H. pylori* cells expressing RNase J variants, in

which each acetylated lysine was replaced by an alanine residue. We observed

that cells with a reduced concentration of wild type RNase J are longer than

those in which RNase J expression is induced. This revealed that RNase J,

most probably indirectly, is involved in the control of *H. pylori* length.

Interestingly, this phenotype does not depend on the activity of RNase J alone,

but rather on the RNA degradosome complex, since the overexpression of

RNase J WT in a Δ*rhpA* strain does not reduce cell length. Given that we

previously showed that about 80% of *H. pylori* mRNAs are at least two times

more abundant in an RNase J-depleted mutant, it is difficult to define the target

mRNA that is causing the observed elongation phenotype[8]. One possibility

would be the overexpression of MreC (9.3-fold upregulated in an RNase J-

depleted strain), that has been shown to cause cell filamentation in *H. pylori*[35].

Alanine replacement of the different acetylated positions of RNase J leads to

different phenotypes, showing that acetylation is indeed important for RNase J

activity in *H. pylori*. Variant K649A behaves like the wild type enzyme in this

test, in agreement with the notion that the morphology phenotype depends on

the activity of the whole RNA degradosome complex, which is not affected by

whether there is a charge in this position or not. The phenotype of the K134A

mutant suggests that it is affected in RNase J activity; interestingly this residue

is located within the N-terminal extension of *Hp*RNase J (Supp Figure S2), that

we previously found to be very important for efficient RNase J activity[8].

However, it is not known how these acetylated residues impact RNase J

properties. They could affect its stability, activity, oligomerization, degradosome

complex formation, compartmentalization, binding of other partners or RNA, or

a combination of these.

We found that several mechanisms participate in the acetylation of the

different residues of RNase J. Our results indicate that HPB8_1270 and the *H.*

pylori RimI homologue indeed function as acetylases. In addition, we uncovered

a role of non-enzymatic acetylation by acetyl phosphate that is produced and

metabolized by the Pta and AckA enzymes. The *E. coli* RimI protein is a

multifunctional acetyltransferase, that can perform ε-lysine acetylation and Nα

acetylation on the L31 and S18 ribosomal proteins, respectively[31,36], with high

substrate specificity. In contrast, RimI from *Mycobacterium tuberculosis* is an

Nα acetylase with more relaxed substrate specificity *in vitro*[37]. Here, we found

that the *H. pylori* RimI targets K323, K397 and K649 of RNase J. HPB8_1270 is

a protein belonging to the DUF2156 family, containing two GNAT domains. It

presents homology with the BT_3689 protein from *Bacteroides*

thetaiotaomicron, which contains an acetyl-CoA molecule in its active site (PDB

2HQY), suggesting its function as an acetyltransferase. Here, we found that

HPB8_1270 is involved in the acetylation of K323, K397, K511 and K649,

showing that in *H. pylori* this protein is most likely an acetyltransferase. Lastly,

acetyl phosphate-mediated acetylation, which is responsible for most

acetylation events in *E. coli* and *B. subtilis*[30,32], was important for the acetylation

of K134, K323, K397, K511 and K649.

These data demonstrate that multiple mechanisms can target the same

lysine residue and that there is a high redundancy in the functions of these

acetylation mechanisms. Acetylation at residue K649 was strongly dependent

on three such mechanisms; this redundancy could testify of a critical importance

of acetylation at this position. In addition, other acetylation mechanisms could

occur in *H. pylori* that might account for some of these acetylation events. More

work is needed, in particular using multiple mutants to determine how many of

these mechanisms can target the different residues and the influence of the

different acetylation mechanisms on the activity and regulation of RNase J and

the RNA degradosome.

Using Clustal Omega 1.2.4[38], we performed an alignment of the RNase J

proteins of *H. pylori* and of the closely related organism *Campylobacter jejuni*,

but also of the more distant *B. subtilis*, *S. aureus*, *S. pyogenes* and *D.*

radiodurans. Interestingly, this revealed that most of the acetylated lysines are

conserved in *H. pylori* and *C. jejuni* RNase J proteins, and some of them are

also conserved in more distant organisms (Supp Figure S5). The case of K511

is particularly interesting, as it is conserved in *S. pyogenes*, *B. subtilis*, *H. pylori*

and *C. jejuni*, and in the case of *S. aureus* this position is a Q residue which

might behave as a "permanently-acetylated" lysine. In *H. pylori*, the mutant

expressing RNase J K511A variant presents a strong growth defect. The

relevance of lysine acetylation in the regulation of the activity of the RNase J

proteins from other organisms remains to be addressed.

In conclusion, we have established that RNase J is acetylated on multiple

lysine residues through different mechanisms, and that its acetylated residues

affect the activity of RNase J and phenotypes such as cell morphology.

Furthermore, our data suggests that acetylation on K649 can influence the

oligomeric state and activity of RNase J, and that the dimeric form could be the

most active one. Thus, acetylation is an important player in the modulation of

the activity of RNase J and of the RNA degradosome of *H. pylori*.

Materials and methods

Bacterial strains and growth conditions

618
619
The bacterial strains used in this study are summarized in Supp Table S4. *H.*
pylori strains were derivatives of strain B128[39,40]. Plasmids (Supp Table S4)
were constructed and amplified using *E. coli* XL1-Blue (Agilent Technologies).
H. pylori strains were grown on Blood agar base 2 (Oxoid) plates supplemented
with 10% defibrinated horse blood and with an antibiotic-antifungal cocktail
composed of 2.5 µg/mL amphotericin B, 0.31 µg/mL polymyxin B, 6.25 µg/mL
trimethoprim and 12.5 µg/mL vancomycin. *H. pylori* mutants were selected
using 20 µg/mL kanamycin, 10 µg/mL apramycin or 6 µg/mL chloramphenicol.
For liquid cultures, we used Brucella broth supplemented with 10% fetal calf
serum (FCS) (Eurobio), the antibiotic-antifungal cocktail and, when necessary,
chloramphenicol to select for plasmid-containing bacteria. *H. pylori* cells were
grown at 37ºC under a microaerophilic atmosphere (6% O_2, 10% CO_2, 84% N_2)
using an Anoxomat (Mart Microbiology) atmosphere generator.

Plasmid preparation and transformations

634
A NucleoBond Xtra midi kit (Macherey-Nagel) and a QIAamp DNA minikit
(Qiagen) were used for plasmid preparations for *H. pylori* transformation and
genomic DNA extractions, respectively. For plasmid preparations for *E. coli*
transformation and sequencing, a QIAprep Spin Miniprep kit (Qiagen) was
used. PCR was carried out with either DreamTaq DNA polymerase
(Thermofisher) or with Q5 High-Fidelity DNA polymerase (NEB) when the
product required a high-fidelity polymerase. A list of the oligonucleotides that
were used can be found in Supp Table S5.

Construction of H. pylori mutants and fusions

The cassette for the construction of the *H. pylori* RNase J-FLAG strain was constructed by the isothermal assembly technique[41], by assembling in frame the last 500 bp of the *rnj* gene followed by the FLAG-tag, the apramycin resistance gene and the 500 bp immediately downstream from *rnj*. The resulting fragment was PCR-amplified and 1 µg of the product was introduced into *H. pylori* by natural transformation. Transformants were selected on plates with apramycin. Strains carrying the pILL2157 plasmid with the wild type *rnj* gene[2] and its derivates with the different *rnj* mutants (constructed following standard procedures with the primers compiled in Supp Table S5) were transformed in the same manner with 1.5 µg of the plasmids and selected on plates with chloramphenicol. The resulting strains were then used to transform by cassettes constructed in a similar manner to delete the *rnj* gene and replace it by a kanamycin resistance gene [as in [2]]. The transformants were selected on plates containing kanamycin, chloramphenicol and 1 mM IPTG to allow the expression of the copy of *rnj* from the plasmid.

Plasmids for BACTH were constructed from plasmids pUT18 and pKNT25[42] by inserting wild type or mutant variants of the *rnj* gene in the *Pst*I and *Kpn*I restriction sites following standard techniques using plasmids indicated in Supp Table S4.

Deletion of the genes of interest, construction of fusions and/or insertion of cassettes was verified by PCR and sequencing of the region of interest.

Purification and crystallography of RNase J

A vector co-expressing residues 21-465 of RhpA with a C-terminal 3C-

HALO-his6 tag and residues 139-691 of RNase J (C1hRSEj) (from *H. pylori*

strain 26695 gene HP1430, UniProt P56185) with a C-terminal hexa-histidine

tag was generated by the Oxford Protein Production Facility at Harwell. Cells

were grown in terrific broth supplemented with 0.5 M NaCl and 1 mM betain, 30

µg/mL kanamycin, 50 µg/mL carbenicillin and 20 µg/mL chloramphenicol. Cells

were grown at 37^{o}C until optical density at 600 nm reached 0.3, then moved to

20^{o}C and induced with 0.2 mM IPTG (isopropyl β-D-1-thiogalactopyranoside)

for 24 hours. Cells were harvested by centrifugation, then suspended in 50 mM

Tris pH 7.4, 50 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic

acid) pH 7.4, 500 mM NaCl, 1 mM TCEP (tris(2-carboxyethyl)phosphine),

0.05% v/v Triton X-100, 5% v/v glycerol and 10 mM imidazole and stored

frozen. The suspension was thawed, passed 5 times through a cell disruptor

(Avestin Emulsiflex C5 homogeniser, 1000 bars) and clarified by centrifugation

at 37,000 g for 30 minutes, 5^{o}C. The supernatant was applied to a HiTrap Ni

chelating column, washed with lysis buffer containing 0.01% v/v Triton X-100,

and 15 mM imidazole, then eluted with a gradient in imidazole. The optimal

fractions were pooled and cleaved with 3CP-GST protease cleavage, using 300

µl of 0.85 mg/mL protease to cleave C1/RSE protein isolated from 1 L culture,

at 4^{o}C overnight. This step removes HALO-Chis-tag on RhpA. The 3CP-cleaved

protein was loaded onto HALO-RESIN and the flow-through collected, which

contained the RNase J/helicase complex. The sample was concentrated with a

30kD MWCO centrifugal concentrator and applied to a S200 SEC column

equilibrated in 20 mM Tris-HCl pH 7.4, 20 mM HEPES pH 7.4, 150 mM NaCl,

5% glycerol, 0.1% v/v Triton X-100, 1 mm TCEP. The sample was concentrated

with a 30kD MWCO centrifugal concentrator to optical density at 280 nm of 3.38

and used in crystallization screens. Crystals were found in PEG I screen 0.1 M

HEPES pH 7.5, 25% wt/v polyethylene glycol 1000.

Diffraction data were collected at I24 at Diamond Light source at wavelength

0.9778 Å using a Pilatus6M detector. Crystals are in space group $P4_122$ with

cell dimensions a=b=158 c=214.26 Å. The structure was solved by molecular

replacement using *Streptomyces* RNase J (PDB 3BK2) for with PHASER and

the CCP4 suite[43,44]. There was no space in the lattice for the RhpA protein. The

model of RNase J 139-691 was built using COOT[45] and refined with PHENIX[46].

A homology model for the C-terminal domain, residues 587 to 691, was

prepared using PHYRE2[47], and the structure of the full-length protein was

predicted using the ROBETTA server[21]. The C-terminal domain from the

ROBETTA model was docked into the map and fitted as a rigid body and then

linked to the model of the well resolved core (139-586). As the density for the C-

terminal domain was poorly resolved due to disorder, the side chains were

removed. A final cycle of refinement using the core only (137-586) and not the

disordered C-terminal domain were used to provide the refinement parameters

summarized in the supplementary Table S1. A monomer of RNase J 139-691

occupies the asymmetric unit, and the tetramer is generated through

crystallographic symmetry. Estimates of buried surface area were made using

the PISA program in the CCP4 suite[48]. The model and structure factors have

been deposited in the PDB with accession code 7PCR.

Bacterial two-hybrid (BACTH)

Bacterial two hybrid assays were carried out in *E. coli* strain BTH101 as

previously described[42]. Briefly, strains carrying derivatives of the two vectors,

pUT18 and pKNT25, were grown overnight in 1 mL LB with 40 µg/mL

kanamycin, 100 µg/mL ampicillin and 0.1 mM IPTG. The OD at 600 nm of the

resulting cultures was measured in a TECAN plate reader and the beta-

galactosidase activity was calculated by mixing 500 µL of the cultures with 500

µL of buffer Z, 100 µL of chloramphenicol and 50 µL of 0.1% SDS, vortexing the

cells and then adding 200 µL of 4 mg/mL ONPG. The reactions were incubated

at 28ºC until they turned yellow and the reactions were stopped by adding 500

µL of 1 M Na_2CO_3. Samples were centrifuged for 5 min at 14,000 rpm and the

OD at 420 and 550 nm of the upper fraction was measured in a TECAN plate

reader. The beta-galactosidase activity (Miller units) was calculated as

described[49].

Protein expression and purification

Wild type and variant versions of RNase J and RhpA were cloned into a

pET28a+ vector in *E. coli* XL1-Blue and then transformed in *E. coli* Bli5 for

expression. The strains were grown in 800 mL cultures in LB with 40 µg/mL

kanamycin and 25 µg/mL chloramphenicol and, once they reached an OD of

0.5-0.8, were induced with 1 mM IPTG. The cultures were incubated at 30ºC

and 160 rpm during 3 h. The cells were collected by centrifugation at 4000 rpm

in an Avanti J-E centrifuge (Beckman Coulter) with a JA-14 rotor at 4ºC during

30 min and pellets were frozen at -80ºC until further processing.

Pellets were thawed on ice and resuspended on lysis buffer containing 25

mM Tris-HCl pH 7.8 for RNase J and pH 8 for RhpA, 500 mM NaCl, 10%

glycerol, 20 mM imidazole, 1 mM dithiothreitol (DTT), 50 U benzonase and

protease inhibitor cocktail (c0mplete EDTA-free, Roche). Cells were lysed by

sonication and centrifuged for 30 min at 4,000 rpm in a 5810R centrifuge

(Eppendorf) at 4ºC to eliminate cellular debris. The supernatant was then

incubated overnight at 4ºC and under agitation with 2 mL Ni-NTA resin (50%

v/v, Thermofisher) pre-equilibrated with lysis buffer.

Resins were packed onto Poly-Prep Chromatography columns (Bio-Rad) and

washed with 10 volumes of 25 mM Tris-HCl at the appropriate pH, 300 mM

NaCl, 1 mM DTT and 20 mM imidazole, 10 volumes of the same buffer with 50

mM imidazole and 10 volumes of the same buffer with 100 mM imidazole. The

elution was carried out with an elution buffer containing 25 mM Tris-HCl at the

appropriate pH, 300 mM NaCl, 1 mM DTT and 500 mM imidazole, in 1 mL

fractions. Aliquots of the resulting fractions were run on a 4-20% Mini-Protean

TGX stain-free precast protein gel (Bio-Rad) at 200V for 30 min and stained by

Imperial protein stain (ThermoFisher). After destaining, the corresponding

purest fractions were chosen, mixed and concentrated using Vivaspin 20 tubes

(MWCO 30,000 Da, GE Healthcare) down to 1 mL volumes. The buffer was

replaced using G-25 columns (PD MidiTrap G-25, GE Healthcare) with 25 mM

Tris-HCl, 300 mM NaCl, 1 mM DTT and 10% glycerol. Samples were aliquoted

and frozen at -80ºC.

Analytical ultracentrifugation

Samples of purified proteins were thawed on ice and the buffer was replaced

using G-25 columns (PD MiniTrap G-25, GE Healthcare) with 25 mM Tris-HCl

pH 8, 300 mM NaCl and 0.5 mM Tris(2-carboxyethyl)phosphine (TCEP, Sigma).

Appropriate amounts of the different proteins were mixed and diluted in the

same buffer with a final volume of 300 µL that was charged into 2 sector

analytical ultracentrifugation cells. The concentration of the proteins was 3 mM

when there was only one protein, 3 mM:3 mM when two proteins were in

equimolar amounts and 1 mM:5 mM when the proteins were mixed at a 1:5

ratio. The centrifugation was performed for 16 h on an Optima analytical

ultracentrifuge (Beckman-Coulter) with an 8-hole AnTi50 rotor, at 42,000 rpm

and 20ºC, measuring the optical density at 230 nm in order to determine the

sedimentation profile of the protein or complex. The data was analyzed with the

Sedfit software (NIH) using a diffusion deconvoluted continuous sedimentation

coefficient distribution c(s) with one discrete component model[50]. The model

was chosen in order to take in consideration the signal due to TCEP absorption.

The presented graphs correspond to a resolution of 300 points with a

confidence interval of 0.9. For the individual proteins, the sedimentation

coefficient independent of the concentration (S_0) was calculated.

The partial specific volume ($\bar{v}$) of the proteins was estimated by the software

Sednterp[51] based on the amino acid sequence. This software was also used for

the calculation of the density and viscosity of the buffer used throughout the

experiments. Calculation of sedimentation coefficients for protein structures was

performed with the program Hydropro[52].

Circular dichroism

The near-UV absorption spectra (195-260 nm) were measured, in the same

buffer as for the AUC experiments, by circular dichroism with a Circular

Dichroism Spectrometer Model 215 (Aviv instruments) using 0.2 mm optical

path cells with 5 mM of RNase J WT and K649A mutant. The proportions of the

different secondary structure elements were determined with the BeStSel

software [53,54] based on the amino acid sequence.

Ribonuclease activity tests

The endoribonuclease and exoribonuclease activities were measured using

6-carboxyfluorescein (6FAM)-labeled RNA substrates either in 5' or in 3' as

listed in Supp Table S5. 500 nM RNA substrate were incubated with 500 nM

protein in a total volume of 10 µL with a buffer containing 50 mM Tris-HCl pH

7.8, 100 mM NaCl, 8 mM $MgCl_2$, 0.1 mM and 0.1 mg/mL bovine serum albumin

(BSA). The reactions were carried out at 37ºC for 0-30 min for the

exoribonuclease activity tests and 0-75 min for endoribonuclease activity tests.

Reactions were stopped by adding 98% formamide and 10 mM ethylene

diamine tetraacetic acid (EDTA) and by incubating for 15 min at 95 ºC.

The degradation products were run on a Mini-Protean TBE-Urea gels (Bio-

Rad) that was pre-run for 60 min at 200 V. The samples were run for 5 min at

50 V and then for 30 min at 200 V. Gels were visualized on a Chemidoc MP

Imaging System (Bio-Rad) with 0.2 s exposure at 488 nm. The gels were then

stained with ethidium bromide to reveal the molecular weight marker (ssRNA

molecular weight marker, Takara). Experiments were performed in triplicate and

images were quantified ImageLab (Bio-Rad, version 6.1). Decay coefficients, λ,

were calculated with the formula $N(t) = N_0 e^{-\lambda t}$, where *N(t)* is the amount of

substrate at time *t* and N_0 is the initial amount. The half-lives, $t_{1/2}$ were then

calculated with the formula $t_{1/2} = \frac{Ln(2)}{\lambda}$.

Phase contrast microscopy

Phase contrast microscopy was performed with an Axio Observer

microscope (Zeiss) equipped with an Axiocam camera with X100 magnification.

Image acquisition was performed with the axiovision software. Images were

cropped and adjusted using FiJi software. The length of wild type cells and that

of cells expressing wild type or mutant versions of RNase J from a plasmid, with

or without 1 mM IPTG, was measured by using the MicrobeJ plugin of ImageJ

[28] with the following parameters: cell area above 0.7 μm^2, minimal length of 1

μm and cell width between 0.4-1 μm.

Pulldown and mass spectrometry

H. pylori cultures of strains expressing FLAG-tagged RNase J were grown in

15 mL of Brucella medium until an $OD_{600\ nm}$ of 0.6-1.2 for exponential phase

cultures. Cells were pelleted by centrifugation for 10 min at 4 ºC and 4,000 rpm

and resuspended in PBS to an $OD_{600\ nm}$ of 10. Cells were lysed by sonication

and cell debris were eliminated by centrifugation for 10 min at 13,000 rpm and 4

ºC. The supernatant constitutes the total extract fraction.

Pulldowns were performed using 2 µL of monoclonal anti-FLAG M2

antibodies produced in mice (F3165, Sigma) per reaction and 50 µL of magnetic

protein G-coated Dynabeads according to the supplier's instructions

(Dynabeads Protein G for Immunoprecipitation, 10003D, ThermoFisher).

Proteins were eluted in the denaturing mode and run in a 4-20% Mini-Protean

TGX stain-free precast protein gel (Bio-Rad) at 200V for 30 min and stained by

Imperial protein stain (ThermoFisher) and destained.

Each band corresponding to RNase J-FLAG was cut and washed several

times in 50 mM ammonium bicarbonate (ABC)-acetonitrile (ACN) (1:1) for 15

min at 37 °C. Disulfide bonds were reduced with 10 mM DTT and cysteines

alkylated with 55 mM chloroacetamide. Trypsin (V5111, Promega) digestion

was performed overnight at 37 °C in 50 mM ABC. Peptides were extracted from

the gel by two incubations in 50 mM ABC-ACN-formic acid (FA) (50:50:0.5) for

15 min at 37 °C. After ACN evaporation in a Speed-Vac, resulting peptides

were desalted with stage-tip [55] using C18 Empore disc and eluted with 80%

ACN, 0.1% FA. Peptides were resuspended in 2% ACN, 0.1% FA prior to liquid

chromatography coupled to mass spectrometry (LC-MS/MS) analysis.

For LC-MS/MS, a nanochromatographic system (Proxeon EASY-nLC 1200 -

ThermoFisher Scientific) was coupled on-line to a Q Exactive[TM] Plus Mass

Spectrometer (ThermoFisher Scientific) using an integrated column oven

(PRSO-V1 - Sonation GmbH, Biberach, Germany). For each sample, peptides

were loaded on an in-house packed 26 cm nano-HPLC column (75 µm inner

diameter) with C18 resin (1.9 µm particles, 100 Å pore size, Reprosil-Pur Basic

C18-HD resin, Dr. Maisch GmbH, Ammerbuch-Entringen, Germany) after an

equilibration step in 100 % solvent A (H_2O, 0.1% FA).

Peptides were eluted with a multi-step gradient using 2 to 7 % solvent B

(80% ACN, 0.1% FA) during 5 min, 7 to 23% during 70 min, 23 to 45% during

30 min and 45 to 95% during 5 min at a flow rate of 300 nL/min over 132 min.

Column temperature was set to 60°C.

MS data were acquired using Xcalibur software using a data-dependent Top

10 method with a survey scans (300-1700 m/z) at a resolution of 70,000 and a

MS/MS scans (fixed first mass 100 m/z) at a resolution of 17,500. The AGC

target and maximum injection time for the survey scans and the MS/MS scans

were set to 3E6, 20 ms and 1E6, 60 ms respectively. The isolation window was

set to 1.6 m/z and normalized collision energy fixed to 28 for HCD

fragmentation. We used a minimum AGC target of 1E4 for an intensity threshold

of 1.7E5. Unassigned precursor ion charge states as well as 1, 7, 8 and >8

charged states were rejected and peptide match was disable. Exclude isotopes

was enabled and selected ions were dynamically excluded for 30 seconds.

Raw data were analyzed using MaxQuant software version 1.6.6.0[56] using

the Andromeda search engine[57]. The MS/MS spectra were searched against a

UniProt *Helicobacter pylori* [strain B8[39,40]] database (download in 24/06/2019,

1,719 entries) and tagged RNase J. Usual known mass spectrometry

contaminants and reversed sequences of all entries were included.

Andromeda searches were performed choosing trypsin as specific enzyme

with a maximum number of five missed cleavages. Possible modifications

included carbamidomethylation (Cys, fixed), oxidation (Met, variable), Nter

acetylation (variable) and K acetylation (Lys, variable) for in-gel peptides. The

mass tolerance in MS was set to 20 ppm for the first search then 4.5 ppm for

the main search and 20 ppm for the MS/MS. Maximum peptide charge was set

to seven and five amino acids were required as minimum peptide length.

A false discovery rate (FDR) cutoff of 1 % was applied at the peptide and

protein levels.

Raw files and MaxQuant files (msms.txt) were processed in Skyline-daily

20.1.9.268[58] to generate MS1-XIC and perform peak integration for each K-

acetylated peptide. Normalization across samples was performed by using the

mean of the ratios from non-modified peptides (Supp Table S2).

The mass spectrometry data have been deposited to the

ProteomeXchange Consortium via de PRIDE partner repository[59] with the

dataset identifier PXD027670.

Statistical analyses

Statistical analysis was performed using Prism 9. We applied one-way

ANOVA with Dunnett's multiple comparisons test in Figure 2A, one-way ANOVA

with Sidak's multiple comparisons test in Figure 6B and two-way ANOVA with

Dunnett's multiple comparisons test in Figure 7.

Acknowledgements

We would like to thank Sébastien Brûlé and Bruno Baron for their help with AUC and circular dichroism experiments, and Frédéric Fischer for bioinformatical analysis regarding protein HPB8_1270. We thank Diane Soussan for her help in preliminary experiments of this project and the members of the Unité Pathogenèse de *Helicobacter*. ATA is part of the Pasteur - Paris University (PPU) International PhD Program. This project has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie grant agreement N° 665807, and from the Institut Carnot Pasteur Microbes & Santé. MBL was a master student of the UTC (Université de Technologie de Compiègne). Support was provided by "Fondation pour la Recherche Médicale" for the grant DBF20161136767 to HDR and the Pasteur-Weizmann Consortium of "The Roles of Noncoding RNAs in Regulation of Microbial Life Styles and Virulence" to HDR. BFL and XYP are supported by a Wellcome Trust Investigator award (200873/Z/16/Z). The authors would like to thank the DIM 1HEALTH region Ile-de-France for funding the Centrifection project that has allowed the Optima ultracentrifuge investment. We thank the staff of the OPPL at Harwell Campus, UK for preparing vectors, screening expression conditions for the recombinant proteins, and helpful discussions.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1. Structure of RNase J from *H. pylori.* Crystals were prepared from *H. pylori* RNase J residues 139-691, encompassing the beta-CASP, metallo-beta-lactamase and C-terminal domains. (A) View of the tetramer, which is a dimer-of-dimers. A monomer occupies the asymmetric unit, and the tetramer is generated through crystallographic symmetry. (B) Overlay of the predicted structure from Robetta server (beige) and the refined crystal structure (blue). The individual beta-CASP and metallo-beta-lactamase domains overlay well, but there are noted differences in the orientation of the three loop regions 174-183, 431-441 and 575-586 at the entrance to the active site (shifts indicated by orange arrows). The predicted structure is closer to the orientation seen in *Streptomyces coelicolor* RNase J, shown in overlay with *H. pylori* enzyme in the left panel of (C). The *S. coelicolor* enzyme structure (beige) is in complex with RNA, and the path of the RNA substrate along a channel to the active site is shown in the right panel. The *H. pylori* RNase J (blue) cannot bind RNA with the same orientation seen for the *S. coelicolor* enzyme due to steric clashes with the loops (left panel). The two zinc ions at the active site of the *S. coelicolor* enzyme are indicated in the right panel.

Figure 2. **The solution oligomerization state of RNase J, RhpA and the degradosome assembly.** The profiles are from analytical ultracentrifugation sedimentation of (A) RNase J, (B) RhpA and (C) the degradosome complex at different ratios of RNase J and RhpA. The profiles were recorded at 230 nm wavelength.

Figure 3. **Point mutations modify the oligomeric state of RNase J.** (A)
BACTH two hybrid experiments to asses homo-oligomerization of WT and
mutant RNase J proteins. Differences in the beta-galactosidase activity units
are indicated with ns (non-significant) ** p-value<0.005, *** p-value<0.0005, ****
p-value<0.0001. (B) Analytical ultracentrifugation sedimentation profiles of
RNase J WT and variant K649A at 230 nm wavelength. (C) Analytical
ultracentrifugation sedimentation profiles of RNase J WT and variants K649R
and K649Q at 230 nm wavelength.

Figure 4. Exoribonuclease activity of RNase J wild type and mutants. (A)
Exoribonuclease activity of RNase J WT and different variants using a 24-nt
RNA substrate labeled with 6-carboxyfluorescein (6-FAM) at its 5'-end over the
course of 30 min. (B) Exoribonuclease activity of RNase J WT and different
variants using a 24-nt RNA substrate labeled with 6-FAM at its 3'-end over the
course of 30 min.

Figure 5. Impact of D185A mutation on RNase J oligomeric state. Analytical
ultracentrifugation sedimentation profiles of RNase J WT and variant D185A,
recorded at 230 nm wavelength.

Figure 6. *Helicobacter pylori* **cell morphology is affected by RNase J**
abundance and point mutations at lysine residues that are acetylation
targets. (A) Phase contrast microscopy images of *H. pylori* Δ*rnj* B128 strains
expressing WT or mutant versions of RNase J from plasmid pILL2157 without

(top panels) or with IPTG induction (bottom panels). The phenotype of a *H.*
pylori ∆rnj-∆rhpA strains expressing WT RNase J from a plasmid is also shown.
The scale bar represents 2 µm. (B) Dot plots showing the distribution of the cell
lengths of wild type and the different mutants without or with IPTG. Horizontal
lines indicate the mean length and the error bars are the standard deviation. ns
(non-significant), ***p-value<0.0005, ****p-value<0.0001.

Figure 7. **RNase J peptides containing acetylated residues extracted from**
different *H. pylori* strains. The normalized intensities were obtained by mass
spectrometry from strains expressing FLAG-tagged RNase J in a wild-type
background or in a mutant strain lacking different acetylation mechanism. ns
(non-significant) p-value>0.05, *p-value<0.05, **p-value<0.005, ***p-
value<0.0005, ****p-value<0.0001. Error bars represent the standard deviation.

Table

Table 1. Half-lives of fluorescent RNA substrates in the presence of
different variants of RNase J alone or in complex with RhpA. The half-live
values in brackets are estimated values extrapolated from the decay slope
when the half-lives exceeded the time of the assay, over 30 min for the two first
and the last columns and over 75 min for the third column.

1001

RNase J variant	Half-life $\pm$ SD (min)			
	5'-labeled substrate (exo)	3'-labeled substrate (exo)	3'-labeled substrate with hairpins (endo)	3'-labeled substrate with hairpins + RhpA (complex)
WT	18.77 $\pm$ 3.16	7.37 $\pm$ 3.36	31.36 $\pm$ 17.57	6.29 $\pm$ 2.92
D299A	>30 [165.04 $\pm$ 93.36]	>>>30	>75 [231.05 $\pm$ 163.38]	n.d.
K649A	4.52 $\pm$ 0.52	2.75 $\pm$ 0.82	10.3 $\pm$ 0.56	6.95 $\pm$ 0.49
K649R	>30 [222.80 $\pm$ 175.05]	>30 [36.61 $\pm$ 12.04]	>75 [81.83 $\pm$ 6.81]	27.70 $\pm$ 5.4
K649Q	>30 [56.80 $\pm$ 13.02]	14.82 $\pm$ 9.11	>75 [92.52 $\pm$ 26.85]	>30 [37.33 $\pm$ 28.89]
D185A	9.14 $\pm$ 4.83	4.56 $\pm$ 1.15	28.6 $\pm$ 9.67	n.d.

1002

1003

References

1. Tejada-Arranz, A., de Crécy-Lagard, V. & de Reuse, H. Bacterial RNA Degradosomes: Molecular Machines under Tight Control. *Trends Biochem. Sci.* **45**, 42–57 (2020).

2. Redko, Y. *et al.* A minimal bacterial RNase J-based degradosome is associated with translating ribosomes. *Nucleic Acids Res.* **41**, 288–301 (2013).

3. Amieva, M. & Peek, R. M. Pathobiology of *Helicobacter pylori*–Induced Gastric Cancer. *Gastroenterology* **150**, 64–78 (2016).

4. Sharma, C. M. *et al.* The primary transcriptome of the major human pathogen Helicobacter pylori. *Nature* **464**, 250–255 (2010).

5. Eisenbart, S. K. *et al.* A Repeat-Associated Small RNA Controls the Major Virulence Factors of *Helicobacter pylori*. *Mol. Cell* **80**, 210-226 (2020).

6. Tejada-Arranz, A. & De Reuse, H. Riboregulation in the Major Gastric Pathogen *Helicobacter pylori*. *Front. Microbiol.* **12**, 712804 (2021).

7. El Mortaji, L. *et al.* The sole DEAD-box RNA helicase of the gastric pathogen *Helicobacter pylori* is essential for colonization. *MBio* **9**, e02071-17 (2018).

8. Redko, Y. *et al.* RNase J depletion leads to massive changes in mRNA abundance in *Helicobacter pylori*. *RNA Biol.* **13**, 243–253 (2016).

9. Tejada-Arranz, A. *et al.* The RNase J-Based RNA Degradosome Is Compartmentalized in the Gastric Pathogen *Helicobacter pylori*. *MBio* **11**, e01173-20 (2020).

10. Mathy, N. *et al. Bacillus subtilis* ribonucleases J1 and J2 form a complex with altered enzyme behaviour. *Mol. Microbiol.* **75**, 489–498 (2010).

11. Hausmann, S. *et al.* Both exo- and endo-nucleolytic activities of RNase J1 from *Staphylococcus aureus* are manganese dependent and active on triphosphorylated 5′-ends. *RNA Biol.* **14**, 1431–1443 (2017).

12. Pei, X.-Y., Bralley, P., Jones, G. H. & Luisi, B. F. Linkage of catalysis and 5' end recognition in ribonuclease RNase J. *Nucleic Acids Res.* **43**, 8066–76 (2015).

13. Zhao, Y. *et al.* Structural insights into catalysis and dimerization enhanced exonuclease activity of RNase J. *Nucleic Acids Res.* **43**, 5550–5559

(2015).

14.   Newman, J. A. *et al.* Unusual, Dual Endo- and Exonuclease Activity in the

Degradosome Explained by Crystal Structure Analysis of RNase J1.

Structure **19**, 1241–1251 (2011).

15.   Wolfe, A. J. Bacterial protein acetylation: new discoveries unanswered

questions. *Curr. Genet.* **62**, 335–341 (2016).

16.   Carabetta, V. J. & Cristea, I. M. Regulation, Function, and Detection of

Protein Acetylation in Bacteria. *J. Bacteriol.* **199**, e00107-17 (2017).

17.   Liang, W., Malhotra, A. & Deutscher, M. P. Acetylation Regulates the

Stability of a Bacterial Protein: Growth Stage-Dependent Modification of

RNase R. *Mol. Cell* **44**, 160–166 (2011).

18.   Song, L., Wang, G., Malhotra, A., Deutscher, M. P. & Liang, W.

Reversible acetylation on Lys501 regulates the activity of RNase II.

Nucleic Acids Res. **44**, 1979–1988 (2016).

19.   Ogura, M. & Asai, K. Glucose Induces ECF Sigma Factor Genes, sigX

and sigM, Independent of Cognate Anti-sigma Factors through

Acetylation of CshA in Bacillus subtilis. *Front. Microbiol.* **7**, 1918 (2016).

20.   Callebaut, I., Moshous, D., Mornon, J. P. & De Villartay, J. P. Metallo-β-

lactamase fold within nucleic acids processing enzymes: The β-CASP

family. *Nucleic Acids Res.* **30**, 3592–3601 (2002).

21.   Baek, M. *et al.* Accurate prediction of protein structures and interactions

using a three-track neural network. *Science* **373**, 871-876 (2021).

22.   Nooren, I. M. A. & Thornton, J. M. Diversity of protein-protein interactions.

EMBO Journal **22**, 3486–3492 (2003).

23.   Perutz, M. F. & Raidt, H. Stereochemical basis of heat stability in bacterial

ferredoxins and in haemoglobin A2. *Nature* **255**, 256–259 (1975).

24.   Perutz, M. F. Electrostatic effects in proteins. *Science* **201**, 1187–1191

(1978).

25.   Zhou, H. X. & Pang, X. Electrostatic Interactions in Protein Structure,

Folding, Binding, and Condensation. *Chemical Reviews* **118**, 1691–1741

(2018).

26.   Kamieniarz, K. & Schneider, R. Tools to Tackle Protein Acetylation.

Chem. Biol. **16**, 1027–1029 (2009).

27.   Figaro, S. *et al. Bacillus subtilis* mutants with knockouts of the genes

encoding ribonucleases RNase Y and RNase J1 are viable, with major

defects in cell morphology, sporulation, and competence. *J. Bacteriol.*

195, 2340–8 (2013).

28.   Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high

throughput bacterial cell detection and quantitative analysis. *Nat.*

Microbiol. **1**, 16077 (2016).

29.   El Mortaji, L. *et al.* A peptide of a type I toxin-antitoxin system induces

Helicobacter pylori morphological transformation from spiral shape to

coccoids. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 31398–31409 (2020).

30.   Christensen, D. G. *et al.* Post-translational Protein Acetylation: An Elegant

Mechanism for Bacteria to Dynamically Regulate Metabolic Functions.

Front. Microbiol. **10**, 1604 (2019).

31.   Christensen, D. G. *et al.* Identification of Novel Protein Lysine

Acetyltransferases in *Escherichia coli*. *MBio* **9**, e01905-18 (2018).

32.   Christensen, D. G. *et al.* Mechanisms, Detection, and Relevance of

Protein Acetylation in Prokaryotes. *MBio* **10**, 02708–18 (2019).

33.   Hausmann, S. *et al.* Both exo- and endo-nucleolytic activities of RNase J1

from *Staphylococcus aureus* are manganese dependent and active on

triphosphorylated 5′-ends. *RNA Biol.* **14**, 1431–1443 (2017).

34.   Tejada-Arranz, A. *et al.* RNase R is associated in a functional complex

with the RhpA DEAD-box RNA helicase in *Helicobacter pylori*. *Nucleic*

Acids Res. **49**, 5249–5264 (2021).

35.   El Ghachi, M. *et al.* Characterization of the elongasome core PBP2: MreC

complex of *Helicobacter pylori*. *Mol. Microbiol.* **82**, 68–86 (2011).

36.   Isono, K. & Isono, S. Ribosomal protein modification in *Escherichia coli*. II.

Studies of a mutant lacking the N-terminal acetylation of protein S18. *Mol.*

Gen. Genet. **177**, 645–51 (1980).

37.   Pathak, D., Bhat, A. H., Sapehia, V., Rai, J. & Rao, A. Biochemical

evidence for relaxed substrate specificity of Nα-acetyltransferase

(Rv3420c/rimI) of *Mycobacterium tuberculosis*. *Sci. Rep.* **6**, 28892 (2016).

38.   Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple

sequence alignments using Clustal Omega. *Mol. Syst. Biol.* **7**, 539 (2011).

39.   Farnbacher, M. *et al.* Sequencing, annotation, and comparative genome

analysis of the gerbil-adapted *Helicobacter pylori* strain B8. *BMC*

1106 *Genomics* **11**, 335 (2010).

1107 40. McClain, M. S., Shaffer, C. L., Israel, D. A., Peek, R. M. & Cover, T. L.
1108 Genome sequence analysis of *Helicobacter pylori* strains associated with
1109 gastric ulceration and gastric cancer. *BMC Genomics* **10**, 3 (2009).

1110 41. Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several
1111 hundred kilobases. *Nat. Methods* **6**, 343–345 (2009).

1112 42. Karimova, G., Pidoux, J., Ullmann, A. & Ladant, D. A bacterial two-hybrid
1113 system based on a reconstituted signal transduction pathway. *Proc. Natl.*
1114 *Acad. Sci. U. S. A.* **95**, 5752–6 (1998).

1115 43. Winn, M. D. *et al.* Overview of the CCP4 suite and current developments.
1116 *Acta Crystallogr. D. Biol. Crystallogr.* **67**, 235–42 (2011).

1117 44. McCoy, A. J. *et al.* Phaser crystallographic software. *J. Appl. Crystallogr.*
1118 **40**, 658–674 (2007).

1119 45. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and
1120 development of Coot. *Acta Crystallogr. D. Biol. Crystallogr.* **66**, 486–501
1121 (2010).

1122 46. Afonine, P. V *et al.* Towards automated crystallographic structure
1123 refinement with phenix.refine. *Acta Crystallogr. D. Biol. Crystallogr.* **68**,
1124 352–67 (2012).

1125 47. Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. E.
1126 The Phyre2 web portal for protein modeling, prediction and analysis. *Nat.*
1127 *Protoc.* **10**, 845–858 (2015).

1128 48. Krissinel, E. & Henrick, K. Inference of Macromolecular Assemblies from
1129 Crystalline State. *J. Mol. Biol.* **372**, 774–797 (2007).

1130 49. Miller, J. H. *Experiments in molecular genetics*. Cold Spring Harbor
1131 Laboratory (1972).

1132 50. Schuck, P. Size-distribution analysis of macromolecules by sedimentation
1133 velocity ultracentrifugation and Lamm equation modeling. *Biophys. J.* **78**,
1134 1606–1619 (2000).

1135 51. Laue, T., Shah, B., Ridgeway, T. & Pelletier, S. Computer-aided
1136 interpretation of analytical sedimentation data for proteins. in *Analytical*
1137 *Ultracentrifugation in Biochemistry and Polymer Science* (eds. Harding, S.
1138 E., Rowe, A. & Horton, J.) 90–125 Cambridge: Royal Society of
1139 Chemistry, (1992).

1140   52.   Ortega, A., Amorós, D. & García de la Torre, J. Prediction of
1141   hydrodynamic and other solution properties of rigid proteins from atomic-
1142   and residue-level models. *Biophys. J.* **101**, 892–8 (2011).

1143   53.   Micsonai, A. *et al.* Accurate secondary structure prediction and fold
1144   recognition for circular dichroism spectroscopy. *Proc. Natl. Acad. Sci. U.*
1145   *S. A.* **112**, E3095-103 (2015).

1146   54.   Micsonai, A. *et al.* BeStSel: a web server for accurate protein secondary
1147   structure prediction and fold recognition from the circular dichroism
1148   spectra. *Nucleic Acids Res.* **46**, W315–W322 (2018).

1149   55.   Rappsilber, J., Mann, M. & Ishihama, Y. Protocol for micro-purification,
1150   enrichment, pre-fractionation and storage of peptides for proteomics using
1151   StageTips. *Nat. Protoc.* **2**, 1896–906 (2007).

1152   56.   Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform
1153   for mass spectrometry-based shotgun proteomics. *Nat. Protoc.* **11**, 2301–
1154   2319 (2016).

1155   57.   Cox, J. *et al.* Andromeda: a peptide search engine integrated into the
1156   MaxQuant environment. *J. Proteome Res.* **10**, 1794–805 (2011).

1157   58.   MacLean, B. *et al.* Skyline: an open source document editor for creating
1158   and analyzing targeted proteomics experiments. *Bioinformatics* **26**, 966–8
1159   (2010).

1160   59.   Perez-Riverol, Y. *et al.* The PRIDE database and related tools and
1161   resources in 2019: improving support for quantification data. *Nucleic*
1162   *Acids Res.* **47**, D442–D450 (2019).

1163

1164

A

C-terminal domain

H. pylori principal dimer

B

H. pylori
loop 575-586

H. pylori
loop 174-183

S. coelicolor and
ROBETTA
corresponding loop

H. pylori
loop 431-444

metallo-beta-lactamase domain

beta-CASP domain

C

H. pylori
loop 431-444

S. coelicolor
corresponding loop

S. coelicolor active site and access

Detected acetylated residue

WT exponential phase

Δ*rimI*   Δ*hpb8_1270*   Δ*pta-ackA*