NAD kinase promotes *Staphylococcus aureus* pathogenesis by supporting production of virulence factors and protective enzymes

Clarisse Leseigneur[1], Laurent Boucontet[2], Magalie Duchateau[3], Javier Pizarro-Cerda[1], Mariette Matondo[3], Emma Colucci-Guyon[2], Olivier Dussurget[1]*

[1]Institut Pasteur, Université Paris Cité, CNRS UMR6047, Unité de Recherche *Yersinia*, F-75015 Paris, France ; [2]Institut Pasteur, Université Paris Cité, CNRS UMR3738, Unité Macrophages et Développement de l'Immunité, F-75015 Paris, France ; [3]Institut Pasteur, Université Paris Cité, CNRS USR2000, Unité de Spectrométrie de Masse pour la Biologie, Plateforme de protéomique, F-75015 Paris, France.

*For correspondence: olivier.dussurget@pasteur.fr

Abstract

Nicotinamide adenine dinucleotide phosphate (NADPH) is the primary electron donor for reductive reactions that are essential for the biosynthesis of major cell components in all organisms. Nicotinamide adenine dinucleotide kinase (NADK) is the only enzyme that catalyzes synthesis of NADP(H) from NAD(H). While the enzymatic properties and physiological functions of NADK have been thoroughly studied, the role of NADK in bacterial pathogenesis remains unknown. Here, we used CRISPR interference to knockdown NADK gene expression to address the role of this enzyme in *Staphylococcus aureus* pathogenic potential. We find that NADK inhibition drastically decreases mortality of zebrafish infected with *S. aureus*. Further, we show that NADK promotes *S. aureus* survival in infected macrophages by protecting bacteria from antimicrobial defense mechanisms. Proteome-wide data analysis revealed that production of major virulence associated factors is sustained by NADK. We demonstrate that NADK is required for expression of the quorum-sensing response regulator AgrA, which controls critical *S. aureus* virulence determinants. These findings support a key role for NADK in bacteria survival within innate immune cells and the host during infection.

Introduction

Nicotinamide adenine dinucleotide (NAD^+), its phosphorylated form ($NADP^+$) and their reduced equivalents (NADH and NADPH) are essential cofactors shared by all living organisms, including bacteria. While NAD^+ and NADH are important for cellular energy metabolism, inflammation and senescence (Chini et al., 2021; Covarrubias et al., 2021), $NADP^+$ and NADPH are key cofactors in central metabolism, being involved in tricarboxylic acid (TCA) cycle, pentose phosphate pathway as well as *de novo* synthesis of fatty acids, cholesterol, amino acids and nucleotides (Chandel, 2021). NADPH also provides the reducing power necessary for the restoration of antioxidative defense systems of the cell (Chandel, 2021). Like NAD(H), growing evidence suggests that NADP(H) has a broader role. In particular, the $NADP^+$ derivative nicotinic acid adenine dinucleotide phosphate (NAADP), which is the major intracellular calcium mobilizing molecule, links NADP(H) metabolism with calcium homeostasis and signaling, development and differentiation (Galione and Chuang, 2020).

Whereas there are two known multi-step pathways for NAD^+ biosynthesis, a single enzyme is responsible for the phosphorylation of NAD^+/NADH into $NADP^+$/NADPH: the NAD kinase (NADK) (McGuinness and Butler, 1985; Chini et al., 2021). NADK activity was first reported in the late 30's (Vestin, 1937; Von Euler and Adler, 1938), and the enzyme was purified from yeast by Kornberg in 1950 (Kornberg, 1950). It is only in 2000 that the genes encoding NADK were identified in *Micrococcus flavus* and *Mycobacterium tuberculosis* (Kawai et al., 2000). Since then, NADK genes have been identified in all living organisms, except the intracellular parasitic bacteria *Chlamydia* spp. on the basis of genome annotation (Grose et al., 2006; Fisher et al. 2013). Given the vital role of NADPH, notably during oxidative stress, NADK genes have been shown to be essential for growth of several bacteria, such as *Mycobacterium tuberculosis* (Sassetti et al., 2003)*, Bacillus subtilis* (Kobayashi et al., 2003), *Salmonella enterica* (Grose et al., 2006) and *Staphylococcus aureus* (Chaudhuri et al., 2009;

Gelin et al., 2020). NADKs essentiality may account for the poor characterization of their role in prokaryotes. In contrast, NADKs have been extensively studied in plants (Li et al., 2018), especially in *Arabidopsis thaliana* which possesses three NADK encoding genes (Chai et al., 2006; Turner et al., 2004). NADK has a key role in photosynthesis, plant cell metabolism, intracellular redox balance (Li, 2018), and response to stresses as shown during exposure to aluminum in wheat (Ślaski, 1995), upon cold-shock in green bean leaves (Ruiz et al., 2002), upon treatment with NaCl, ionizing radiations or oxidative stress in *Arabidopsis thaliana* (Chai et al., 2006; Berrin et al., 2005). In plants, NADKs are regulated by calcium and calmodulin (Tai et al., 2019). In contrast, in prokaryotes allosteric regulation of NADK activity by NADPH or NADH is the main mechanism identified so far (Grose et al., 2006).

The importance of microbial metabolic adaptation during infection (Eisenreich et al., 2010; Richardson, 2019; Teoh et al., 2021) and the unknown contribution of NADK to bacterial pathogenesis prompted us to investigate the role of NADK in *Staphylococcus aureus* pathogeny. While being carried as a commensal by around one third of the human population, *S. aureus* is a leading cause of infectious diseases, ranging from mild skin and soft tissue infections to life-threatening endocarditis and bacteremia (Turner et al., 2019). Worryingly, *S. aureus* can develop resistance to virtually all antibiotic classes available (Vestergaard et al., 2019), limiting dramatically therapeutic options for some patients. Therefore, WHO included *S. aureus* in the list of high priority pathogens to promote research and development of new antibiotics and emphasized the need for innovation and diversification in the choice of targets (https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed).

In this study, we used a genetic approach to modulate NADK activity and analyze *S. aureus* behavior in the infected host, in infected cells and in response to stresses mimicking those encountered during infection. We establish the first link between NADK activity and *S.*

aureus pathogenic potential, as NADK inhibition led to an increased survival of the host in a

zebrafish infection model. We further demonstrate that inhibition of NADK leads to a dramatic

decrease of *S. aureus* survival in macrophages, highlighting the enzyme importance during

interactions with immune cells. We further show that exposures to oxidative and envelope-

targeting antimicrobial compounds significantly decrease bacterial survival upon inhibition of

NADK activity. Proteomic analysis indicated that the production of protective enzymes,

virulence factors and the quorum-sensing response regulator AgrA relies on NADK. Thus, our

results reveal a key role for NADK in bacterial pathogenesis.

Results

NADK contributes to *S. aureus* virulence in zebrafish

As most prokaryotes, *S. aureus* possesses a single NADK encoded by the highly conserved *ppnK* gene. To investigate the role of NADK in *S. aureus* pathogenesis, we followed a genetic approach to modulate the expression of *ppnK*. We used the *S. aureus* NADK sgRNA strain (Gelin et al., 2020), in which the levels of *ppnK* expression were decreased by CRISPR interference based on plasmid pSD1 (Zhao et al., 2017). Knockdown of NADK expression in this strain was analyzed by RT-PCR, immunoblotting, and aerobic growth in BHI at 37°C (Figure supplement 1). Growth of the *S. aureus*/sgRNA strain was affected as previously reported (Gelin et al., 2020), attesting the efficiency of the knockdown. We investigated whether NADK inhibition might influence *S. aureus* pathogenesis in a zebrafish infection model. This model allows short kinetics of infection (Prajsnar et al., 2008) and has been previously used to study *S. aureus* infection dynamics (Prajsnar et al., 2012). In addition to being optically transparent, zebrafish larvae innate immune system is the only one operating during the first days of development, facilitating the study of interactions between bacteria and neutrophils or macrophages. We first determined the stability of the pSD1 plasmid during infection. Zebrafish larvae were infected at 60 hours post fertilization (hpf) by intravenous injections of either *S. aureus* containing the empty vector pSD1 or the knockdown NADK sgRNA strain. Bacteria recovered from infected larvae were plated onto BHI agar with and without chloramphenicol. There was no significant difference between the number of bacteria growing on BHI alone and BHI supplemented with chloramphenicol up to 48 hours post-infection (hpi), irrespective of the strain (Figure supplement 2A). Thus, plasmids were maintained in our experimental infection conditions. We next monitored zebrafish survival upon infection with 10^4 bacteria. *S. aureus*/pSD1 infection led to ≈ 75% fish mortality at 24 hpi (Figure 1A). Strikingly, only one larva out of 48 died when bacterial NADK was inhibited

(Figure 1A). Decreased fish mortality upon *S. aureus*/NADK sgRNA infection was also

observed 48 and 72 hpi (Figure 1A). These results indicate that NADK inhibition leads to a

decreased bacterial virulence. We then determined the bacterial burden of fish larvae 24 and 48

hpi. Interestingly, numbers of *S. aureus*/pSD1 and *S. aureus*/NADK sgRNA bacteria at both

time points varied depending on the viability of the fish (Figure 1B). As observed previously

by Prasjnar et al. (Prajsnar et al., 2012), some fishes were able to control the infection with 10^4

bacteria but a high bacterial burden was associated with larval death in most cases (Figure 1B).

Early mortality was observed in zebrafish in which *S. aureus*/pSD1 developed to high numbers

compared to larvae infected with the NADK knockdown strain. Together, these results suggest

that NADK contributes to *S. aureus* pathogenic potential, which depends on both bacterial and

host factors. To investigate the impact of NADK inhibition on phagocyte populations, we

infected intravenously transgenic zebrafish lines with *S. aureus*/pSD1 or *S. aureus*/NADK

sgRNA strains, and observed bacteria, macrophages and neutrophils at time 0, 6, 12 and 24 hpi.

S. aureus/pSD1 bacterial burden was higher than that of *S. aureus*/NADK sgRNA strain in head

and caudal regions observed in fish of comparable viability 12 hours (Figure 1C) and 24 hpi

(Figure supplement 2B). We also noticed a strong depletion of neutrophils in larvae infected

with *S. aureus*/pSD1, while this neutropenia was more limited upon infection with NADK

knockdown bacteria (Figures 1C, 1D and 1E). This correlated with virulence differences of the

two strains, given the key role of neutrophils in *S. aureus* infection (Pollitt et al., 2018; Prajsnar

et al., 2012). In contrast, the number of macrophages in larvae did not significantly decreased

upon infection with either strains (Figure supplement 2C). Altogether, our results suggest that

NADK is important for *S. aureus* pathogenesis.

Figure 1 NADK promotes *S. aureus* virulence in zebrafish. (A) Survival of zebrafish larvae uninfected (CTRL)

or intravenously injected at 60 hpf with 10^4 *S. aureus* containing the empty vector (pSD1) or the NADK

knockdown strain (NADK sgRNA) between 0 and 72 hpi (n=48). (B) Bacterial burden in zebrafish larvae upon

intravenous injection with 10^4 *S. aureus* containing the empty vector (pSD1) or the NADK knockdown strain

(NADK sgRNA). For each strain, CFU were determined in living larvae (open circles) or dead larvae (filled

circles) 0, 24 and 48 hpi. (C) Representative fluorescence confocal images of transgenic *mpx:GFP* zebrafish larvae

uninfected (CTRL), or intravenously injected with 10^4 *S. aureus* containing the empty vector (pSD1) or the NADK

knockdown strain (NADK sgRNA) at 12 hpi. Maximum intensity Z-projection images (2-μm serial optical

sections) of bacteria (red) and neutrophils (green). Scale bars, 25 μm. Insets are shown at higher magnification on

the right panels for head and caudal regions. Scale bars, 10 μm. Neutrophils containing NADK knockdown

bacteria can be seen in the caudal inset. (D) Representative fluorescence confocal images of transgenic *mpx:GFP*

zebrafish larvae uninfected (CTRL), or intravenously injected with 10^4 *S. aureus* containing the empty vector

(pSD1) or the NADK knockdown strain (NADK sgRNA) at 24 hpi, showing neutrophils (green). White arrows

indicate the injection site. Scale bars, 500 μm (E) Number of neutrophils in uninfected zebrafish larvae (CTRL)

or in zebrafish larvae intravenously injected with 10^4 *S. aureus* containing the empty vector (pSD1) or the NADK

knockdown strain (NADK sgRNA) at 24 hpi. Comparison of data was performed using one-way analysis of

variance (***p<0.001, ****p<0.0001).

NADK promotes *S. aureus* survival in macrophages

Since both neutrophils and macrophages are key immune cells specialized in the destruction of

microorganisms, we next investigated if, in addition to triggering neutrophil killing, NADK

might support *S. aureus* survival during macrophages infection. Murine RAW264.7

macrophages were infected with *S. aureus*/pSD1 or *S. aureus*/NADK sgRNA strains. Bacterial

enumerations were performed at time zero and at 1, 2, 4 and 6 hpi. No significant difference in

initial phagocytosis of the two strains could be observed, nor at early time of infection (Figure

2A). In contrast, after 2 hours of infection the percentage of intracellular bacteria was

significantly reduced upon *ppnK* knockdown, and the survival defect increased over time

(Figure 2A), indicating that bacteria were not able to survive inside macrophages without

proper NADK activity. As oxidative burst is a major antimicrobial mechanism of phagocytes

(Fang, 2004), we next investigated the role of ROS produced by macrophages in the control of

the *S. aureus*/NADK sgRNA strain using the antioxidant agent N-acetyl cysteine (NAC). NAC

partially restored bacterial survival (Figure 2A), indicating that the bacterial growth defect of

the *ppnK* knockdown strain was due to its inability to cope with oxidative stress to some extent.

Using confocal fluorescence microscopy, we then investigated bacterial intracellular

localization in macrophages 6 hours post-infection. Few *S. aureus*/NADK sgRNA bacteria were

detected compared to *S. aureus*/pSD1 (Figure 2B), and NAC treatment led to an increased

number of bacteria (Figure 2C), confirming bacterial enumeration results. Interestingly, while

all knockdown bacteria colocalized with lysosomal associated membrane protein 1 (LAMP1),

many *S. aureus*/pSD1 did not upon NAC treatment, suggesting that NADK activity might be

required for phagolysosome escape and bacterial survival in macrophages. Since *S. aureus* has

been shown to be cytotoxic to macrophages, we next wondered if the bacterial growth defect

of the NADK knockdown strain was correlated with reduced lysis of infected cells. We

measured the levels of cell death in macrophages that were uninfected, uninfected and treated

with Triton X-100 or infected with *S. aureus*/pSD1 or *S. aureus*/NADK sgRNA strains.

Detergent treatment expectedly led to massive cell lysis. Infections produced significant

macrophage death compared to uninfected cells, although to much lower levels than that

triggered by Triton X-100. Infection with the *ppnK* knockdown strain was significantly less

cytotoxic than infection with the *S. aureus*/pSD1 strain. Altogether, these results suggest than

NADK promotes survival and possibly virulence of *S. aureus* in macrophages.

Figure 2 NADK promotes *S. aureus* survival and cytotoxicity in macrophages. (A) Percentage of growth of the *S. aureus* strains containing the empty vector (pSD1) or the *ppnK* knockdown strain (NADK sgRNA) at time 0, and 1, 2, 4 and 6 hpi of RAW264.7 macrophages left untreated or treated with N-acetylcysteine (NAC). Bars indicate standard errors of the means of biological replicates (n=4). Comparison of data was performed using two-ways analysis of variance (*p<0.05, **p<0.01, ****p<0.0001). (B-C) Representative images of RAW264.7 macrophages infected with *S. aureus*/pSD1 or *S. aureus*/NADK sgRNA strains and analyzed 6 hpi by confocal fluorescence microscopy using antibodies to label *S. aureus* (Cy5, red) and LAMP-1 (FITC, green). Nuclei were labeled with DAPI (blue). B shows untreated macrophages. C shows macrophages treated with NAC. Insets are shown at higher magnification. Images are representative of three independent experiments. (D) Cell death of RAW264.7 macrophages uninfected (NI), uninfected and treated with Triton X-100 (lysis control), or 6 hpi with *S. aureus* strains containing the empty vector (pSD1) or the *ppnK* knockdown strain (NADK sgRNA). Bars indicate standard errors of the means of biological replicates. Comparison of data was performed using two-ways analysis of variance (*p<0.05, **p<0.01, ****p<0.0001). Data are representative of at least three independent experiments.

NADK protects *S. aureus* from antimicrobial defense compounds

Hydrogen peroxide is one of the reactive oxygen species generated upon activation of macrophages (Fang, 2004). We therefore compared growth of the *S. aureus* strain containing the empty vector to that of the NADK knockdown strain, upon exposure to increasing concentrations of hydrogen peroxide (H_2O_2). While the *S. aureus*/pSD1 strain resisted to high doses of H_2O_2, the *S. aureus*/NADK sgRNA strain showed increasing growth deficiency when exposed to H_2O_2 concentrations ranging from 50 to 500 μM H_2O_2 (Figure 3A). Expectedly, treatment of the *S. aureus*/pSD1 strain with catalase alone or catalase and H_2O_2 did not affect growth (Figure 3B). In contrast, addition of catalase, which catalyzes hydrogen peroxide dismutation into water and oxygen, rescued the growth defect of the *S. aureus*/NADK sgRNA strain upon H_2O_2 treatment (Figure 3B). Taken together, our results show that NADK protects *S. aureus* from toxic effects of a reactive oxygen species involved in host defense. Within phagocytes, bacteria have to cope with other mechanisms of antimicrobial defenses such as

production of lysozyme (Callewaert and Michiels, 2010; Avican et al., 2021; Fang et al., 2016).

We thus investigated the effects of NADK knockdown on *S. aureus* sensitivity to lysozyme.

Since *S. aureus* is highly resistant to lysozyme, we treated bacteria with lysozyme and the

bacteriocin lysostaphin, a combination which potentiates their antibacterial activity against *S.*

aureus (Cisani et al., 1982). At subinhibitory concentrations for the *S. aureus*/pSD1 strain, the

S. aureus/NADK sgRNA strain showed decreased viability when exposed to lysozyme and

lysostaphin (Figure 3C). Thus, NADK protects *S. aureus* from toxic effects of cell envelope-

targeting compounds.

Figure 3 NADK protects *S. aureus* from antimicrobial defense compounds. Bacterial growth was monitored at

OD_{600nm} in BHI broth at 37°C. (A) Percentage of growth of the *S. aureus* strain containing the empty vector (pSD1)

and the *ppnK* knockdown strain (NADK sgRNA) exposed for 6 hours to increasing concentrations of H_2O_2 relative

to the untreated condition. (B) Percentage of growth of the *S. aureus* strain containing the empty vector (pSD1)

and the *ppnK* knockdown strain (NADK sgRNA) exposed for 6 hours to H_2O_2 and catalase alone or in combination,

relative to the untreated condition. Data shown are representative of three independent experiments. Bars indicate

the standard error of the means of biological replicates. Comparison of data was performed using one-way analysis

of variance (ns: nonsignificant, *$p<0.05$, ***$p<0.001$, ****$p<0.0001$). (C) Bacterial cell death of the *S. aureus*

strain containing the empty vector (pSD1) and the *ppnK* knockdown strain (NADK sgRNA) untreated or exposed

for 30 min to 0.05 mg/mL lysostaphin and 5 mg/mL lysozyme. Bacterial permeability was assessed using the

CellTox Green cytotoxicity assay. Bars indicate standard errors of the means of biological replicates. Comparison

of data was performed using two-ways analysis of variance (*$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$).

Data are representative of at least three independent experiments.

NADK supports *S. aureus* protective enzymes

To elucidate mechanisms of action of NADK, unbiased mass spectrometry-based proteomics

was used to identify proteins differentially expressed upon NADK knockdown. Relative protein

abundance was compared in the *S. aureus* strain containing the empty vector and the NADK

knockdown strain grown in BHI. Interestingly, lower levels of major antioxidant enzymes were

detected in the NADK sgRNA strain (Table 1 and table supplement 1). NADK deficiency was

linked to decreased relative abundance of superoxide dismutase SodM, catalase KatA,

thioredoxin Trx2 and methionine sulfoxide reductase MsrA2. Additionally, the NADK sgRNA

strain had lower levels of the peptidoglycan *O*-acetyltransferase OatA and the glycopeptide

resistance-associated two-component system GraRS, major determinants for *S. aureus*

resistance to lysozyme and cationic antimicrobial peptides (CAMP), respectively (Bera et al.,

2005 ; Herbert et al., 2007). We then compared the proteomes of the *S. aureus* strain containing

the empty vector and the NADK knockdown strain upon hydrogen peroxide treatment. OatA,

SodM, Trx2, MrsA2, KatA and other antioxidant enzymes were more abundant in the *S.*

aureus/pSD1 strain than in the NADK deficient strain (Table supplement 2 and table

supplement 3), as it had been observed when strains were grown in BHI. To determine the

effect of NADK knockdown on *S. aureus* proteome *in vivo*, relative protein abundance was

analyzed in zebrafish infected with the *S. aureus* strain containing the empty vector or the

NADK knockdown strain. The D-alanyl carrier protein DltC, which mediates CAMP

resistance, and antioxidant enzymes, such as SodA, KatA, thiol peroxidase Tpx and alkyl

hydroperoxide reductase subunit AhpF, were detected in zebrafish infected with *S.*

aureus/pSD1 (Tables S4 and S5). In contrast, these proteins were undetectable in *S.*

aureus/NADK sgRNA-infected zebrafish. Thus, our approach identified a role for NADK in

supporting *S. aureus* protective proteins *in vitro* and *in vivo*.

Table 1. Relative abundance of a subset of protective proteins differentially expressed by *S.*
aureus pSD1 and NADK sgRNA strains.
269

Uniprot	Protein	Description	Log2R*	P**
Q2FV54	OatA	O-acetyl transferase	3.07	1.78E-05
Q2G000	Trx2	Thioredoxin 2	1.79	2.75E-05
Q2G261	SodM	Superoxide dismutase [Mn/Fe]	1.68	2.11E-06
Q2FZZ3	-	Thioredoxin domain-containing protein	1.59	7.22E-04
Q2G0D9	GraS	Sensor histidine kinase	1.31	4.00E-04
Q2G280	-	Peroxiredoxin	1.26	6.69E-07
P0A086	MsrA2	Methionine sulfoxide reductase	1.10	2.25E-05
Q2FVL7	-	Thioredoxin domain-containing protein	1.02	1.94E-03
Q2G0E0	GraR	Response regulator protein	1.00	2.23E-03
Q2FYU7	KatA	Catalase	0.59	1.00E-05

270
271   * Log2R=Log2[pSD1]/[NADK sgRNA]
272   **Adjusted p value
273

275   **NADK promotes virulence factor expression by *S. aureus***

276   To investigate additional mechanisms of action of NADK, our quantitative proteomic data were

277   analyzed further using proteomaps (Liebermeister et al., 2014). *S. aureus* proteins were

278   classified into functional groups based on KEGG annotation. Expression profiles of strains

grown in BHI were visualized by Voronoi tree maps, clustering proteins in three hierarchical categories. Top-level functional categories relying on NADK activity included biosynthesis, central carbon metabolism, other enzymes, folding, sorting and degradation, translation, transcription, and DNA maintenance (Figure 4A). Remarkably, infectious diseases, signaling molecules and interaction, and signal transduction identified at the highest level (Figure 4A), and *Staphylococcus aureus* infection, bacterial toxins and two-component system identified at the second level (Figure 4B), encompassed multiple major virulence factors such as alpha-hemolysin, gamma-hemolysin, Map, IsdA, and virulence regulators AgrA and SaeS (Figure 4C). NADK deficiency was also linked to decreased relative abundance of other virulence-related proteins, *e.g.,* phenol-soluble modulin (PSM) alpha PsmA, the lipase Lip1, leukocidin-like proteins 1 and 2, and staphopain A (Table 2). Of note, similar proteomic profiles were obtained when strains were exposed to hydrogen peroxide, confirming the importance of NADK for proper expression of virulence proteins in conditions mimicking host defenses (Figure supplement 3, Table supplement 6). Furthermore, comparative analysis of whole protein extracts from bacterial pellets and culture supernatants by immunoblotting showed a massive decrease in the levels of alpha-hemolysin secretion by the NADK deficient strain (Figures 4D and 4E). To determine if the differential protein abundance of toxins was translated into different levels of enzymatic activity, a Christie-Atkins-Munch-Peterson test was performed using *S. aureus* RN4220, *S.aureus*/pSD1, and *S. aureus*/NADK sgRNA strains (Adhikari et al. 2007). Strikingly, NADK deficiency led to a lack of hemolysis (Figure 4F). In contrast *S. aureus*/pSD1 produced a synergistic hemolysis in presence of the RN4220 strain, indicating secretion of hemolysins and phenol soluble modulins. Taken together, these data suggest that NADK is required for expression of *S. aureus* virulence determinants.

Figure 4 NADK promotes expression of *S. aureus* virulence determinants. The *S. aureus* strain containing the

empty vector (pSD1) and the *ppnK* knockdown strain (NADK sgRNA) were grown in BHI broth at 37°C. (A-C)

Whole bacterial cell lysates were analyzed by LC-MS/MS. Voronoi treemaps were generated to visualize proteins

more abundant in *S. aureus*/pSD1 than in the knockdown strain at three hierarchical levels according to KEGG

database: top level (A), second level (B), third level (C). Colors represent functional categories. Category size

based on LFQ intensity corresponds to differences in protein abundance. (D) Whole bacterial cell lysates and

culture supernatants were analyzed by immunoblotting using antibodies against alpha-hemolysin and EF-Tu. (E)

Quantification of the Hla immunoblots normalized to corresponding EF-Tu protein levels in the pellet. (F) A

Christie-Atkins-Munch-Peterson test was performed by streaking the *S. aureus* strain containing the empty vector

(pSD1) and the *ppnK* knockdown strain (NADK sgRNA) perpendicularly to the *S. aureus* RN4220 strain that

produces only beta-hemolysin (vertical streak) on sheep blood agar. Clearing zones indicate hemolysis.

Table 2. Relative abundance of a subset of virulence-related proteins differentially expressed
by *S. aureus* pSD1 and NADK sgRNA strains.
318

Uniprot	Protein	Description	Log2R*	P**
Q2FWM4	AgrA	Accessory gene regulator protein	+	NA
P0C818	PsmA4	Phenol-soluble modulin alpha 4	+	NA
Q2FVK2	HlgC	Gamma-hemolysin component C	+	NA
P0C7Y1	PsmA1	Phenol-soluble modulin alpha 1	+	NA
Q2FUU5	Lip1	Lipase	+	NA
Q2FVK1	HlgB	Gamma-hemolysin component B	7.88	3.85E-10
Q2G1X0	Hly	Alpha-hemolysin	6.99	3.85E-10
Q2FWN9	LukL2	Leukocidin-like protein 2	3.88	4.14E-08
Q2FVK3	HlgA	Gamma-hemolysin component A	3.84	1.47E-08
Q2FWP0	LukL1	Leukocidin-like protein 1	3.09	2.25E-06
Q2G2R8	SspP	Staphopain A	2.89	4.08E-07

319
320 *Log2R=Log2[pSD1]/[NADK sgRNA]; +: protein detected in pSD1 strain and not detected from NADK sgRNA strain
321 **Adjusted p value: NA: not applicable
322

NADK promotes AgrA expression by *S. aureus*

AgrA is a central regulator of *S. aureus* virulon. Therefore, the impact of NADK on AgrA

observed above may account for the positive regulation of virulence factor production. To attest

NADK-dependent regulation of virulence determinants controlled by Agr, we analyzed

transcript levels of two additional members of the Agr regulon, the positively regulated

multifunctional RNA RNAIII and the negatively regulated glutathione peroxidase gene *bsaA*.

We carried out RT-PCR on RNA purified from the *S. aureus* strain containing the empty vector

and the NADK knockdown strain during exponential growth in BHI broth at 37°C. Decreased

RNAIII transcript levels and increased *bsaA* transcript levels were detected in the NADK

knockdown strain (Figures 5A, 5B and 5D). These results are in accordance with the positive

regulatory role of NADK on AgrA, and the regulation of AgrA on RNAIII and *bsaA* expression.

To determine whether NADK sustains AgrA protein levels by promoting gene expression,

AgrA transcript levels were compared in the *S. aureus*/pSD1 and *S. aureus*/NADK sgRNA

strains. The expression of *agrA* was strongly decreased upon NADK knockdown (Figures 5C

and 5D). Thus, full NADK activity is required for optimal *agrA* expression and subsequent

virulence gene expression. To decipher the mode of action of NADK on *agrA* expression, we

next investigated the effect of the *ppnK* knockdown on *S. aureus* oxidative content, since

transcription of the *agr* system is repressed in oxidizing conditions (Baker et al., 2010; Rothfork

et al., 2004). We compared the levels of ROS in the *S. aureus*/pSD1 and NADK sgRNA strains

grown in aerobic conditions. Decreasing NADK levels led to a significant increase in ROS

content (Figure 5E), which could contribute to negative regulation of AgrA expression.

Together, our data suggest that NADK activity is important for proper *agrA* expression,

possibly by controlling the bacterial redox state, ultimately promoting *S. aureus* virulence

potential.

Figure 5 NADK is required for AgrA expression and *S. aureus* redox control. The *S. aureus* strain containing the

empty vector (pSD1) and the *ppnK* knockdown strain (NADK sgRNA) were grown aerobically in BHI broth at

37°C for 4 hours. (A-C) RNA was extracted and subjected to RT-PCR using oligonucleotides specific to *gyrA*,

RNAIII, *bsaA* and *agrA* genes, respectively. (D) Relative gene expression was normalized to *gyrA* transcript levels.

(E) *S. aureus* ROS levels were quantified using the $DCFH_2$-DA assay. Comparison of data was performed using

t-test (****p<0.0001). Data are representative of at least three independent experiments.

Discussion

The ability of *S. aureus* to survive and thrive in a wide variety of niches during infection, ranging from skin to deeper tissues and abiotic devices, mirrors highly plastic metabolic capacities allowing adaptive responses to constantly changing microenvironments (Potter et al., 2020). Metabolic pathways are not only necessary for nutrient acquisition to sustain bacterial growth, but also for regulation of virulence and are therefore central to host-pathogen interactions (Eisenreich et al., 2010; Harper et al., 2018; Richardson, 2019; Teoh et al., 2021; Tomlinson et al., 2021).

Here we show for the first time that NADK contributes to bacterial pathogenic potential. NADK was necessary for successful *S. aureus* infection of zebrafish larvae, ultimately leading to neutropenia and death. Decreasing NADK activity led to decreased host mortality without affecting bacterial burden initially, suggesting that NADK promotes virulence factor activity and/or counteracts host defenses independently of its contribution to bacterial growth capacity. Besides neutrophils, macrophages are crucial cells in the innate immune defense against infection. We uncovered that inhibition of *S. aureus* NADK activity has a major impact on bacteria interactions with macrophages, reducing dramatically bacterial survival after phagocytosis. We demonstrated that NADK contributes to resistance to ROS produced by macrophages, neutralizing an important defense mechanism against bacterial infection (Pidwill et al., 2021). Along the same lines, we showed that NADK protected *S. aureus* from hydrogen peroxide toxicity. NADK, as the only source of NADP(H), is a key component of defense against oxidative stress (Grose et al., 2006; Mailloux et al., 2011). In bacteria, it has been shown that NADK activity is increased upon exposure to oxidative stress, acting as a metabolic switch to decrease the NAD^+ pool which fuels ROS formation and increase the NADPH pool which promotes ROS scavenging activities (Singh et al., 2007). Analysis of *S. aureus* proteome revealed a drop in the levels of major antioxidant enzymes upon inhibition of *ppnK* expression.

By supporting superoxide dismutase, catalase, methionine sulfoxide reductase and other

antioxidant enzymes, NADK could prevent damages caused by ROS formed during

phagocytosis. However, antioxidant treatment of infected macrophages using NAC only

partially mitigated growth defect of the NADK knockdown strain. NADK might thus contribute

to bacterial survival within macrophages through pathways other than ROS detoxification. We

showed that inhibition of *ppnK* expression led to growth defect upon envelope stress, a

condition that is encountered in the host, particularly in macrophages where bacteria face

antimicrobial peptides and hydrolytic enzymes such as lysozyme (Rosenberger et al., 2004;

Cohn and Wiener, 1963; Callewaert and Michiels, 2010). *S. aureus* resists lysozyme through

peptidoglycan modification by the *O*-acetyltransferase OatA (Bera et al., 2005, 2006).

Regulation by the glycopeptide resistance-associated two-component system GraRS and D-

alanylation of teichoic acid by the *dlt* operon gene products also contribute to *S. aureus*

resistance to lysozyme, as well as CAMP (Herbert et al., 2007). OatA, DltA and GraRS were

less abundant when NADK was knocked down, suggesting that NADK could help *S. aureus*

resist antibacterial envelope-targeting compounds produced by macrophages and/or secreted

by the host. Strikingly, our proteomic analysis also highlighted the contribution of NADK to

production of virulence factors. Among them, PSM-alpha peptides have previously been shown

to lyse leucocytes, elicit phagosomal escape and play a key role in pathogenesis (Wang et al.,

2007; Kobayashi et al., 2011; Peschel and Otto, 2013; Grosz et al., 2014; Kitur et al., 2015).

The pore-forming toxin alpha-hemolysin has also been demonstrated to lyse leukocytes and is

required for full virulence in mouse, rabbit, and rat models of infection (Patel et al., 1987;

O'Callaghan et al., 1997; McElroy et al., 1999; Kitur et al., 2015; Seilie and Wardenburg,

2017). Gamma-hemolysin leucocidin is leukotoxic as well and contributes to *S. aureus*

bacteremia (Nilsson et al., 1999; Spaan et al., 2014). Control of these factors by NADK could

thus promote bacterial survival in macrophages and zebrafish, and trigger macrophage and

neutrophil death upon *S. aureus* infection. Remarkably, we found that inhibition of *ppnK*

expression decreases abundance of AgrA, the response regulator of *S. aureus agr* system at the

center of quorum-sensing and virulence. AgrA directly enhances the transcription of the PSM-

alpha genes and governs RNAIII-dependent expression of multiple virulence enzymes

including alpha-hemolysin and leukocidins (Queck et al., 2008). Thus, the contribution of

NADK to production of virulence factors is presumably mediated through AgrA. NADK

regulated expression of RNAIII and BsaA similarly to AgrA, corroborating this hypothesis.

Besides lowering abundance of AgrA, NADK knockdown led to decreased levels of *agrA*

transcript. Increased ROS levels detected in the knockdown strain could account for this

regulation as transcription of the *agr* operon is repressed in oxidative conditions (Rothfork et

al., 2004; Baker et al., 2010; Sun et al., 2012). Additionally, oxidation leads to formation of an

intramolecular disulfide bond between Cys-199 and Cys-228 in the DNA binding domain of

AgrA, thereby introducing a conformational change disrupting its association to promoter

region of target genes (Sun et al., 2012). A recent study confirmed oxidation sensing of the *agr*

system and revealed another mode of AgrA regulation by CoAlation of redox-active Cys199

inhibiting its DNA-binding activity (Bakovic et al., 2021).

In summary, our study shows that NADK supports production of protective enzymes,

virulence determinants and the response regulator AgrA, possibly by controlling the bacterial

redox state, ultimately promoting *S. aureus* pathogenic potential. NADK inhibition and quorum

quenching are attractive therapeutic approaches. Further investigations are now required to

fully understand the contribution of bacterial NADKs during infection processes and develop

chemical inhibitors that could be used in combination with current antibiotics to fight multidrug

resistant bacteria.

Materials and methods

Plasmids, bacterial strains, cell lines and culture conditions

Bacterial strains and plasmids used in this study are listed in Table 3. All plasmids were maintained in *Escherichia coli* TOP10. The restriction deficient *Staphylococcus aureus* strain RN4220 was transformed with plasmids isolated from *E. coli*. The *S. aureus* Xen36 strain was transformed with plasmids isolated from the RN4220 strain as previously described (Gelin et al., 2020). *E. coli* was grown in lysogeny broth (LB) medium (Difco) with shaking at 200 rpm or on LB agar plates at 37°C. *S. aureus* was grown in brain heart infusion (BHI) broth (BD) or in tryptic soy broth (TSB) (Difco) with shaking at 200 rpm or on BHI agar plates at 37°C. When required, culture medium was supplemented with antibiotics (carbenicillin 100 μg/mL for *E. coli*, chloramphenicol 15 μg/mL or anhydrotetracycline 100 ng/mL for *S. aureus*). RAW 264.7 macrophages (TIB-71, ATCC) were cultured in DMEM (Gibco) containing 10 % FCS (Gibco). Macrophages were seeded into 24-well plates at 10^5 cells per well.

Table 3 Strains and plasmids used in this study.

Strain/plasmid	Relevant characteristics	Reference
Strains		
E. coli strain		
TOP10	F⁻*mcrA Δ(mrr-hsdRMS-mcrBC) φ80lacZ ΔM15 ΔlacX74 recA1 araD139 Δ(ara-leu) 7697 galU galK λ⁻ rpsL(Str^R) endA1 nupG*	Invitrogen
S. aureus strains		
Xen36	Strain derived from a clinical isolate from a bacteremic patient	PerkinElmer, ATCC49525
Xen36/pSD1	Xen36 strain carrying plasmid pSD1	Gelin et al., 2020
Xen36/NADK sgRNA	Xen36 strain carrying plasmid pSD1 *ppnK*	Gelin et al., 2020
RN4220	Restriction-deficient strain derived from NCTC 8325-4	Kreiswirth et al., 1983
Plasmids		
pSD1	dCas9 ATc-inducible and sgRNA constitutive expression plasmid	Zhao et al., 2017
pSD1 *ppnK*	pSD1 plasmid carrying sgRNA targeting *S. aureus ppnK*	Gelin et al., 2020

Table 4 Primers used in this study

Primer	Sequence (5'-3')
RT-ppnK-F	GTGACTCCAAGTCTAATGCC
RT-ppnK-R	ATTTTTCAACTTCATGAGGTAACC
RT-gyrA-F	GTGTTATCGTTGCTCGTG
RT-gyrA-R	CGGTGTCATACCTTGTTC
RT-RNAIII-F	AGTTTCCTTGGACTCAGTGCT
RT-RNAIII-R	AGGGGCTCACGACCATACTT
RT-bsaA-F	GCGAAGAAGCAGCTCAAAAC
RT-bsaA-R	CCTTCGCGATCCACTAAAAA
RT-agrA-F	GCCCTCGCAACTGATAATCC
RT-agrA-R	CACCGATGCATAGCAGTGTTC
RT-rluA-F	CCGCGAGAAATACCGAGTGT
RT-rluA-R	TCTCCCACAATTGGATGCCC
RT-operon-F1	TGACTTGCTTAAAAAGCACACTG
RT-operon-R1	ACGAGCATTTGTCCTACTTCAGA
RT-operon-F2	AACCGTTGAAGAAACATTCGACA
RT-operon-R2	GACGCTTGTTCACCAACTTCA
RT-operon-F3	CATCGTTTGGAAAGAGCGGC
RT-operon-R3	GGCATTAGACTTGGAGTCACCT
RT-operon-F4	ACGTGTGCACGATTCTTTCAT
RT-operon-R4	ATGGCGCTCACTGTCTTCT
RT-operon-F5	AGTTCATTTGCATACGGGACG
RT-operon-R5	ACGCTCTTTTTCATCTGTGTTCA
RT-operon-F6	TAACTTGTGCGATGACGGTGG
RT-operon-R6	TTCCAATCACAATCCCCATCAA
RT-operon-F7	ACGTTGATGAATTGAAGCAAGAG
RT-operon-R7	ACTTTAGCGACACCAAAAGCA
RT-operon-F8	TCAAGTGGCGTTACAGGTGA
RT-operon-R8	TTCAAATACCGCCAACGCAT

Bacterial growth measurement

S. aureus strains were grown overnight in BHI broth at 37°C and diluted to OD_{600nm}=0.05 into BHI broth. For NADK inhibition experiments, culture medium was supplemented with 100 ng/mL anhydrotetracycline (Sigma-Aldrich) and 15 μg/mL chloramphenicol when required. For stress experiments, culture medium was supplemented with hydrogen peroxide (0 to 500 μM), catalase (10 μg/mL), lysozyme (5 mg/mL) or lysostaphin (0.05 mg/mL). Bacterial suspensions were incubated in 96-well plates with shaking at 200 rpm at 37°C. Growth was monitored with a microplate reader (Glomax Discover, Promega). Experiments were performed at least three times.

Christie-Atkins-Munch-Peterson test

S. aureus hemolytic activity was performed as described previously (Adhikari et al. 2007). Briefly, *S. aureus* RN4220 was streaked vertically at the center of a sheep blood agar plate. *S. aureus*/pSD1 and *S. aureus*/NADK sgRNA strains were then streaked perpendicularly to the RN4220 strain. The plate was incubated at 37°C for 18 hours. Blood agar plates were then photographed with a Scan500 imager (Interscience). The assay was repeated three times.

RNA extraction and RT-PCR

Bacteria grown for 4 h were collected by centrifugation 20 min at 15,000 x *g* at 4°C. Bacterial lysis was performed using Precellys lysing kit (P000914-LYSK0-A) and the Precellys program 4 (6.500 rpm for 30 sec) twice at 4°C. RNA extraction was performed in 1 mL of TRIzol (Invitrogen) according to the manufacturer's recommendations. RT-PCR was carried out with Superscript one step RT kit (Invitrogen). Primers used are indicated in Table 4. RT-PCR assays were repeated at least three times.

Immunoblotting

Bacteria grown for 4 h were collected by centrifugation 10 min at 10,000 x *g* at 4° C and resuspended in 500 µL of PBS. Bacteria were lysed using the Precellys lysing kit (P000914-LYSK0-A) and the Precellys program 4 (6,500 rpm for 30 sec) twice at 4°C. Lysates were centrifuged for 15 min at 10,000 x *g* at 4 °C and supernatants were collected for protein quantification using the Quick start Bradford protein assay kit 2 (Biorad). Samples were mixed with Laemmli buffer (Biorad) and 10% β -mercaptoethanol and denatured for 5 min at 95°C. Samples were separated onto 4−20% Miniprotean TGX stain-free precast gel (Biorad) in TGS buffer and transferred on polyvinylidene fluoride membranes. Membranes were incubated overnight at 4°C with primary antibodies diluted in 5% blotto (R114 rabbit anti-EF-Tu

489   polyclonal antibodies, 1:5000; R250 rabbit polyclonal anti-*S. aureus* NADK antibodies,

490   1:1000; anti-staphylococcal α-toxin (α-hemolysin) rabbit antiserum (Sigma-Aldrich S7531)

491   1:1000). Membranes were then incubated with 1:2500 antirabbit horseradish peroxidase-

492   conjugated antibodies (Abcam). Blots were revealed using the ECL kit (Pierce).

RAW 264.7 macrophage infection

RAW 264.7 macrophages were seeded into 24-well plates at 100,000 cells per well. *S. aureus*

Xen36/pSD1 and *S. aureus* Xen36/NADK sgRNA strains were grown to an OD_{600nm} of 0.6,

washed three times and diluted in DMEM to obtain a multiplicity of infection (MOI) of 10.

Infected macrophages were centrifuged at 500 x *g* for 5 min and incubated for 15 min at 37°C

and 5% CO_2 to synchronize phagocytosis. The medium was replaced with DMEM containing

gentamicin (20 μg/mL) for 30 min to kill extracellular bacteria. Cells were washed three times

in DPBS and incubated in DMEM supplemented with 100 ng/mL anhydrotetracycline and 10

mM N-acetyl cysteine (NAC, Sigma-Aldrich) when indicated. Cells were incubated for 6 h at

37°C and 5% CO_2. At each time points, cells were lysed in 0.2% Triton X-100 for 10 min at

37°C. The number of bacteria released from the cells was determined by plating serial dilutions

of the lysates on BHI agar plates that were incubated at 37°C for colony-forming units (CFU)

enumeration.

Immunofluorescence assay

RAW 264.7 macrophages were cultured on glass coverslips in 24-well plates. At 6 h post-

infection, cells were fixed in paraformaldehyde (PFA) 4% for 15 min, permeabilized for 4 min

with 0.1% Triton X-100 in 1% bovine serum albumin (BSA)-PBS and blocked in 1% BSA-

PBS. Fixed cells were incubated for 30 min with rabbit anti-*S. aureus* (Abcam ab20920) and

rat anti-LAMP1 1D4B (Abcam ab25245) primary antibodies. Cells were then incubated for 30

min with goat anti-rabbit Cy5 (Immunoresearch 111 175 144) or goat anti-rat FITC (Invitrogen

A11006) secondary antibodies and DAPI (10 μg/mL). Slides were mounted with Fluoromount

G (Invitrogen). Samples were observed with a Zeiss Axiovert 200M epifluorescence

microscope equipped with a Plan-apochromat objective (100X/1.4 Oil Ph3; Carl Zeiss, Inc.).

Images were acquired with a CCD camera Coolsnap, processed with Metamorph software v.6

(Molecular Devices) and analyzed with ImageJ software.

Cytotoxicity assay

Overnight cultures of *S. aureus*/pSD1 and *S. aureus*/NADK sgRNA were diluted to an OD_{600nm}

0.05 in BHI supplemented with ATc (100 ng/mL) and chloramphenicol (7 μg/mL), spread in a

96 wells plate and incubated for 6 hours at 37°C with shaking at 200 rpm. After 5h30 of growth,

lysozyme (5 mg/mL) and lysostaphin (0.05 mg/mL) were added. At t=6h, 100 μL of the CellTox

Green Reagent (1:500, G8741, Promega) were added to bacteria. After incubation at RT for 15

min, both OD_{600nm} and fluorescent signal (excitation at 475 nm, emission at 500-550 nm) were

measured with a microplate reader (Glomax Discover, Promega).

For the cytotoxicity assay on infected macrophages, culture medium was replaced by either

DPBS or by 0.2% Triton X-100 6 hours after the infection. After 10 min at 37°C, cells were

washed with DPBS, and then incubated with 100 μL of CellTox Green Reagent for 15 min at

RT. Fluorescent signal (excitation at 475 nm, emission at 500-550 nm) was measured with a

microplate reader (Glomax Discover, Promega).

ROS levels quantification

Overnight cultures of *S. aureus*/pSD1 and *S. aureus*/NADK sgRNA were diluted to an OD_{600nm}

0.05 in BHI supplemented with with ATc (100 ng/mL) and chloramphenicol (7 μg/mL), spread

in a 24 wells plate and incubated at 37°C with shaking at 200 rpm for 4h. After 4h, the OD_{600nm}

was measured and adjusted to 1, then bacteria centrifugated for 3 min at 13.000g. $DCFH_2$-DA

(5 mM) was mixed with NaOH (0.1 M) to obtain DCF. Bacterial pellets were resuspended in

100 uL of DCF and incubated for 40 min at RT protected from light in a 96-wells black plate.

Then fluorescence signal (excitation 475 nm, emission 500-550 nm) was measured with a

microplate reader (Glomax Discover, Promega).

Ethics Statement

Animal experiments were performed according to European Union guidelines for handling of

laboratory animals (http://ec.europa.eu/environment/chemicals/lab_animals/home_en.htm).

All experiments performed on larvae older than 5 days post fertilization were approved by the

Institut Pasteur Animal Care and Use Committee and the French Ministry of Education,

Research and Innovation, and registered under the reference APAFIS#31827.

Zebrafish care and maintenance

Homozygous Tg*(mfap4::mCherryF)* (ump6Tg) (Phan et al., 2018) Tg*(mpx::GFP)*[i114]

(Renshaw et al., 2006) double transgenic fishes were raised in our facility. Eggs were obtained

by natural spawning, bleached according to standard protocols, and then kept in Petri dishes

containing Volvic spring water and, from 24 h hpf onwards, 0.003% 1-phenyl-2-thiourea (PTU)

(Sigma-Aldrich) was added to prevent pigmentation. Embryos were reared at 28°C or 24°C

according to the desired speed of development; infected larvae were always kept at 28°C. All

timings in the text refer to the developmental stage at the reference temperature of 28°C. Larvae

were anesthetized with 200 μg/ml of buffered tricaine methane sulfonate (MS-222, Sigma-

Aldrich) during the injection procedure as well as during *in vivo* imaging and processing for

bacterial burden evaluation.

Zebrafish infection

The volume of injected bacterial suspension was deduced from the diameter of the drop obtained after mock microinjection, as previously described (Levraud et al., 2008). Bacteria were diluted from overnight cultures and allowed to reach exponential growth. Then, they were recovered by centrifugation, washed, and resuspended at the desired concentration in PBS. 55-60 hours post-fertilization, anesthetized zebrafish larvae were microinjected intravenously (IV) with 1 nL of bacterial suspension at the desired dose as described (Colucci-Guyon et al., 2011; Mostowy et al., 2013). Infected larvae were transferred into individual wells containing 1 mL of Volvic water and 0.003% PTU in 24-well culture plates, incubated at 28°C and regularly observed under a stereomicroscope.

Evaluation of the bacterial burden in infected zebrafish larvae

Infected zebrafish larvae were collected at 0, 24, 48 and 72 hpi and lysed for the evaluation of the bacterial burden as previously described (Boucontet et al., 2018; Mostowy et al., 2013). Each larva was placed in an individual 1.5 mL Eppendorf tube and anesthetized with tricaine (200 μg/mL), washed with 1 mL of sterile water, and placed in 150 μL of sterile water. Larvae were then homogenized using a pestle motor mixer (Argos). Each sample was transferred to an individual well of a 96-well plate and 10 X serial dilutions were performed. For CFU enumeration and to assess plasmid stability throughout the infection kinetics, serial dilutions of lysates were plated on BHI agar plates with and without chloramphenicol (15 μg/mL), that were incubated overnight at 37°C.

Zebrafish live imaging, image processing and analysis

Quantification of total macrophages and neutrophils on living transgenic reporter larvae was performed upon infection as we previously described (Mostowy et al., 2013). Briefly, bright

field, GFP and RFP images of whole living anesthetized larvae were taken using a Leica

Macrofluo™ Z16 APOA (zoom 16:1) macroscope equipped with a Leica PlanApo 2.0X lens,

and a Photometrics® CoolSNAP™ *HQ2* camera. Images were captured using the Metavue

software version 7.5.6.0 (MDS Analytical Technologies). After capture of images, larvae were

washed and transferred in a new 24-well plate filled with 1 mL of fresh water per well and

incubated at 28°C.

Whole-mount immunohistochemistry and confocal imaging

Whole larvae were fixed in 4% formaldehyde/PBS overnight at 4°C. After washes in PBSDT

(PBS, 1% DMSO, and 0.1% Tween), fixed larvae were treated with acetone at -20° for 20 min,

rinsed with 1× HBSS/0.1% Tween, permeabilized with a collagenase solution for up to 1 h

(Svoboda et al., 2001), washed in PBSDT, blocked with 10% Western Blocking Reagent

(Merck; 11921673001)/PBSDT, and then incubated for 24-48 h at 4°C with the primary

antibodies. Chicken anti-GFP (1:1000, ab13970, Abcam), and anti-*Staphylococcus aureus*

(1:500, ab20920, Abcam) antibodies were used. Following washes, samples were blocked in

10% Western blocking solution/PBSDT before overnight incubation with secondary antibodies

at 4°C with agitation. The secondary antibodies were Alexa Fluor 488-conjugated goat anti-

chicken (1:1000, A11039, Invitrogen) and Cy™5-conjugated goat anti-rabbit (1:200, 111-175-

144, Jackson Immunoresearch). Larvae were progressively embedded in glycerol (for

conservation and microscopic observation) and mounted in lateral position for imaging.

Confocal fluorescent imaging of fixed injected larvae was performed as previously described

(Colucci-Guyon et al., 2020). Briefly, injected larvae were positioned in lateral position in 35

mm glass-bottom μ-Dishes (81158, Ibidi). Larvae were immobilized in the dish using a drop of

1% low-melting-point agarose (V2111, Promega) solution in 80% glycerol. A Leica SP8

confocal microscope equipped with two PMT and one Hybrid detector, a 20X multi-IMM

(water-oil-glycerol) objective (HC PL APO CS2 20X/0.75), a X–Y motorized stage and the

LAS-X software were used to image injected larvae. All samples were acquired using the same

settings.

Total protein extraction

For *in vitro* samples, overnight cultures of *S. aureus*/pSD1 and *S. aureus*/NADK sgRNA were

diluted to an OD_{600nm} 0.05 in BHI with 200 μM H_2O_2 when indicated, spread in a 24-well plate

and incubated at 37°C with shaking at 200 rpm. After 6 h, bacteria were centrifugated for 15

min at 10,000 *g* at 4°C and the bacterial pellets were stored at -80°C. For *in vivo* samples,

zebrafish larvae were infected with *S. aureus*/pSD1 or *S. aureus*/NADK sgRNA strains (see

above). At 12 hpi, 5 larvae of each condition were isolated, anesthetized with tricaine, washed

with 1 mL of sterile water, dried, placed on dry ice and stored at -80°C. All samples were then

resuspended in 1 mL of Tris 100 mM pH 7/urea 8M. Proteins were extracted using the Precellys

kit SK38 (P000915-LYSK0-A0, Bertin), in a Cryolys homogenizer (Bertin) by 6 cycles of

agitation at 6,500 rpm for 15 s and 30 s rest.

Protein digestion and cleanup of peptides

Proteins were reduced using 5mM TCEP for 30 min at room temperature. Alkylation of the

reduced disulfide bridges was performed using 10mM iodoacetamide for 30 min at room

temperature in the dark. Proteins were then digested in two steps, first with 1 μg r-LysC Mass

Spec Grade (Promega) for 4 h at 30°C and then samples were diluted below 2 M urea with 100

mM Tris HCl pH 8.5 and 1 μg Sequencing Grade Modified Trypsin was added for the second

digestion overnight at 37°C. Proteolysis was stopped by adding formic acid (FA) at a final

concentration of 5%. The peptide samples were cleaned using C_{18} cartridges (Agilent

Technologies) in the automated AssayMAP Bravo Platform. The protocol for peptide cleanup

included in the platform was used as scaffold using the following settings. Cartridges were

primed with 100 μL of 100% acetonitrile (ACN) at 300 μL/min. Equilibration was done with

50 μL of 0.1% FA at 10 μL/min. The sample was loaded at 5 μL/min. Cup wash (25 μL) and

the internal cartridge wash (50 μL at 10 μL/min) were performed with 0.1% FA. Peptides were

eluted with 50 μL of 50% ACN/0.1% FA at 5 μL/min. Samples were dried and stored at −80

°C until further use. Peptides were resuspended in 2% acetonitrile (ACN)/0.1% FA prior to LC-

MS injection.

Liquid chromatography-mass spectrometry

LC-MS/MS analysis was performed on a Orbitrap Q Exactive™ Plus Mass Spectrometer

(Thermo Fisher Scientific) coupled with a Proxeon EASY-nLC 1200 (Thermo Fisher

Scientific). One μg of peptides was injected onto a home-made 30 cm C_{18} column (1.9 μm

particles, 100 Å pore size, ReproSil-Pur Basic C18, Dr. Maisch GmbH, Ammerbuch-Entringen,

Germany). Column equilibration and peptide loading were done at 900 bars in buffer A (0.1%

FA). Peptides were separated with a multi-step gradient from 2 to 7 % buffer B (80% ACN,

0.1% FA) in 5 min, 7 to 23 % buffer B in 70 min, 23 to 45 % buffer B in 30 min at a flow rate

of 250 nL/min. Column temperature was set to 60°C. MS data were acquired using Xcalibur

software using a data-dependent method. MS scans were acquired at a resolution of 70,000 and

MS/MS scans (fixed first mass 100 m/z) at a resolution of 17,500. The AGC target and

maximum injection time for the survey scans and the MS/MS scans were set to $3E^6$, 20ms and

$1E^6$, 60ms respectively. An automatic selection of the 10 most intense precursor ions was

activated (Top 10) with a 45 s dynamic exclusion. The isolation window was set to 1.6 m/z and

normalized collision energy fixed to 28 for HCD fragmentation. We used an underfill ratio of

1.0 % corresponding to an intensity threshold of $1.7E^5$. Unassigned precursor ion charge states

as well as 1, 7, 8 and >8 charged states were rejected and peptide match was disabled.

Data analysis

Acquired Raw files were analyzed using the MaxQuant software version 1.6.6.0 (Cox et al., 2011) using the Andromeda search engine (Cox and Mann, 2008; Tyanova et al., 2016). Sample were grouped by type of experiment. The MS/MS spectra were searched against the Staphylococcus aureus Uniprot reference proteome (2,889 entries).

All searches were performed with oxidation of methionine and protein N-terminal acetylation as variable modifications and cysteine carbamidomethylation as fixed modification. Trypsin was selected as protease allowing for up to two missed cleavages. The minimum peptide length was set to 5 amino acids and the peptide mass was limited to a maximum of 8,000 Da. The false discovery rate (FDR) for peptide and protein identification was set to 0.01. The main search peptide tolerance was set to 4.5 ppm and to 20 ppm for the MS/MS match tolerance. Second peptides were enabled to identify co-fragmentation events. One unique peptide to the protein group was required for the protein identification. A false discovery rate cut-off of 1 % was applied at the peptide and protein levels. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD032968.

The statistical analysis of the proteomics data was performed as follows: Five biological replicates were acquired per condition. To highlight significantly differentially abundant proteins between two conditions, differential analyses were conducted through the following data analysis pipeline: (1) deleting the reverse and potential contaminant proteins; (2) keeping only proteins with at least three quantified values in one of the two compared conditions to limit misidentifications and ensure a minimum of replicability; (3) log2-transformation of the remaining intensities of proteins; (4) normalizing the intensities by median centering within conditions thanks to the *normalizeD* function of the R package DAPAR (Wieczorek et al.,

2017), (5) putting aside proteins without any value in one of both compared conditions: as they

are quantitatively present in a condition and absent in another, they are considered as

differentially abundant proteins and (6) performing statistical differential analysis on them by

requiring a minimum fold-change of 2.5 between conditions and by using a LIMMA t test

(Smyth, 2005 ; Ritchie et al., 2015) combined with an adaptive Benjamini-Hochberg correction

of the p values thanks to the *adjust.p* function of the R package cp4p (Giai Gianetto et al., 2016).

The robust method of Pounds and Cheng was used to estimate the proportion of true null

hypotheses among the set of statistical tests (Pounds and Cheng, 2006). The proteins associated

with an adjusted p value inferior to an FDR level of 1% have been considered as significantly

differentially abundant proteins. Finally, the proteins of interest are therefore the proteins that

emerge from this statistical analysis supplemented by those being quantitatively absent from

one condition and present in another.

Statistical analysis

Results are expressed as means ± SEM of at least 3 replicates. Statistical analysis was performed

using GraphPad Prism (GraphPad Prism® 9.1.2. Software). One-way, two-ways ANOVA or t-

test were used to compare data. Differences between groups were considered significant when

the *p* value was lower than 0.05. Survival data were plotted using the Kaplan-Meier estimator

and log-rank (Mantel-Cox) tests were performed to assess differences between groups.

Aknowlegments

Authors are deeply indebted to Liliane Assairi, Sylvie Pochet and Gilles Labesse who led us into the world of NAD kinases. We are grateful to past and present members of the NADK consortium, all members of the *Yersinia* research unit and the French national reference center for plague and other yersiniosis, as well as Frédéric Barras and Daniel Ladant for helpful discussions and insightful comments. We thank Changlong Zhao for providing pSD1. We also wish to thank the members of the fish facility team Yohann Rolin and Noël Aimar for their excellent care of the fish.

This work was funded by the Agence Nationale de la Recherche (grant ANR-17-CE18-0011-02) and supported by Institut Pasteur, Université Paris Cité and CNRS. The *Yersinia* Research Unit is member of the Laboratory of Excellence Integrative Biology of Emerging Infectious Diseases (ANR-10-LBX-62-IBEID). We are grateful for support for equipment from the French government Programme d'investissements d'avenir France BioImaging (ANR-10-INSB-04-01).

Competing interests

The authors declare no conflict of interest.

References

Adhikari RP, Arvidson S, Novick RP. 2007. A nonsense mutation in *agrA* accounts for the defect in *agr* expression and the avirulence of *Staphylococcus aureus* 8325-4 traP::kan. *Infect Immun* **75**:4534-40. doi: 10.1128/IAI.00679-07.

Avican K, Aldahdooh J, Togninalli M, Mahmud AKMF, Tang J, Borgwardt KM, Rhen M, Fällman M. 2021. RNA atlas of human bacterial pathogens uncovers stress dynamics linked to infection. *Nat Commun* **12**:3282. doi:10.1038/S41467-021-23588-W

Baker J, Sitthisak S, Sengupta M, Johnson M, Jayaswal RK, Morrissey JA. 2010. Copper stress induces a global stress response in *Staphylococcus aureus* and represses *sae* and *agr* expression and biofilm formation. *Appl Environ Microbiol* **76**:150-160. doi: 10.1128/AEM.02268-09.

Baković J, Yu BYK, Silva D, Baczynska M, Peak-Chew SY, Switzer A, Burchell L, Wigneshweraraj S, Vandanashree M, Gopal B, Filonenko V, Skehel M, Gout I. 2021. Redox regulation of the quorum-sensing transcription factor AgrA by coenzyme A. *Antioxidants* **10**:841. doi: 10.3390/antiox10060841

Bera A, Biswas R, Herbert S, Götz F. 2006. The presence of peptidoglycan *O*-acetyltransferase in various staphylococcal species correlates with lysozyme resistance and pathogenicity. *Infect Imm* **74**: 4598-4604. doi:10.1128/IAI.00301-06

Bera A, Herbert S, Jakob A, Vollmer W, Götz F. 2005. Why are pathogenic staphylococci so lysozyme resistant? The peptidoglycan O-acetyltransferase OatA is the major determinant for lysozyme resistance of *Staphylococcus aureus*. *Mol Microbiol* **55**:778-87. doi: 10.1111/j.1365-2958.2004.04446.x

Berrin JG, Pierrugues O, Brutesco C, Alonso B, Montillet JL, Roby D, Kazmaier M. 2005. Stress induces the expression of AtNADK-1, a gene encoding a NAD(H) kinase in

Arabidopsis thaliana. Mol Genet Genomics **273**:10–19. doi:10.1007/s00438-005-1113-1

Boucontet L, Passoni G, Thiry V, Maggi L, Herbomel P, Levraud J-P, Colucci-Guyon E. 2018.

A model of superinfection of virus-infected zebrafish larvae: increased susceptibility to

bacteria associated with neutrophil death. *Front Immunol* **9**:1084.

doi:10.3389/FIMMU.2018.01084

Callewaert L, Michiels CW. 2010. Lysozymes in the animal kingdom. *J Biosci.* **35**:127-160.

doi10.1007/s12038-010-0015-5

Chai M-F, Wei P-C, Chen Q-J, An R, Chen J, Yang S, Wang X-C. 2006. NADK3, a novel

cytoplasmic source of NADPH, is required under conditions of oxidative stress and

modulates abscisic acid responses in *Arabidopsis. Plant J* **47**:665–674. doi:10.1111/j.1365-

313X.2006.02816.x

Chandel NS. 2021. Carbohydrate metabolism. *Cold Spring Harb Perspect Biol* **13**:1–15.

doi:10.1101/CSHPERSPECT.A040568

Chaudhuri RR, Allen AG, Owen PJ, Shalom G, Stone K, Harrison M, Burgis TA, Lockyer M,

Garcia-Lara J, Foster SJ, Pleasance SJ, Peters SE, Maskell DJ, Charles IG. 2009.

Comprehensive identification of essential *Staphylococcus aureus* genes using Transposon-

Mediated Differential Hybridisation (TMDH). *BMC Genomics* **10**:291. doi:10.1186/1471-

2164-10-291

Chini CCS, Zeidler JD, Kashyap S, Warner G, Chini EN. 2021. Evolving concepts in NAD$^+$

metabolism. *Cell Metab.* **33**:1076-1087. doi:10.1016/j.cmet.2021.04.003

Cisani G, Varaldo PE, Grazi G, Soro O. 1982. High-level potentiation of lysostaphin anti-

staphylococcal activity by lysozyme. *Antimicrob Agents Chemother* **21**:531-535. doi:

10.1128/AAC.21.4.531.

777 Cohn ZA, Wiener E. 1963. The particulate hydrolases of macrophages. II. Biochemical and

778  morphological response to particle ingestion. *J Exp Med* **118**:1009-20. doi:

779  10.1084/jem.118.6.1009

780 Colucci-Guyon E, Rifflet A, Saint-Auret S, da Costa A, Boucontet L, Laval T, Prehaud C,

781  Blanchard N, Levraud JP, Boneca IG, Demangel C, Guenin-Macé L. 2020. Spatiotemporal

782  analysis of mycolactone distribution in vivo reveals partial diffusion in the central nervous

783  system. *PLoS Negl Trop Dis*. **14**:e0008878. doi: 10.1371/journal.pntd.0008878

784 Colucci-Guyon E, Tinevez J-Y, Renshaw SA, Herbomel P. 2011. Strategies of professional

785  phagocytes in vivo: unlike macrophages, neutrophils engulf only surface-associated

786  microbes. *J Cell Sci* **124**:3053–3059. doi:10.1242/JCS.082792

787 Covarrubias AJ, Perrone R, Grozio A, Verdin E. 2021. NAD$^+$ metabolism and its roles in

788  cellular processes during ageing. *Nat Rev Mol Cell Biol*. **22**:119–141. doi:10.1038/s41580-

789  020-00313-x

790 Cox J, Mann M. 2008. MaxQuant enables highpeptide identification rates, individualized

791  p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol*

792  **26**:1367–1372.

793 Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen JV, Mann M. 2011. Andromeda: A

794  peptide search engine integrated into the MaxQuant environment. *J Proteome Res* **10**:1794-

795  1805.

796 Eisenreich W, Dandekar T, Heesemann J, Goebel W. 2010. Carbon metabolism of intracellular

797  bacterial pathogens and possible links to virulence. *Nat Rev Microbiol* **8**:401–412.

798  doi:10.1038/NRMICRO2351

799 Fang FC. 2004. Antimicrobial reactive oxygen and nitrogen species: concepts and

800  controversies. *Nat Rev Microbiol* **2**:820-832. doi:10.1038/nrmicro1004

801 Fang FC, Frawley ER, Tapscott T, Vázquez-Torres A. 2016. Bacterial stress responses during

host infection. *Cell Host Microbe* **20**:133–143. doi:10.1016/J.CHOM.2016.07.009

Fisher DJ, Fernandez RE, Maurelli AT. 2013. *Chlamydia trachomatis* transports NAD via the

Npt1 ATP/ADP translocase. *J Bacteriol* **195**:3381-3386. doi:10.1128/JB.00433-13

Galione A, Chuang K-T. 2020. Pyridine nucleotide metabolites and calcium release from

intracellular stores. *Adv Exp Med Biol* **1131**:371–394. doi:10.1007/978-3-030-12457-1_15

Gelin M, Paoletti J, Huteau V, Nahori M, Leseigneur C, Dugue L, Clement D, Pons J, Assairi

L, Pochet S, Labesse G, Dussurget O. 2020. From substrate to fragments to inhibitor active

in   vivo   against   *Staphylococcus   aureus*.   *ACS   Infect   Dis*   **6**:422–435.

doi:10.1021/acsinfecdis.9b00368

Giai Gianetto Q, Combes F, Ramus C, Bruley C, Coute Y, Burger T. 2016. Calibration plot for

proteomics: a graphical tool to visually check the assumptions underlying FDR control in

quantitative experiments. *Proteomics* **16**: 29–32.

Grose H, Joss L, Velick SF, Roth JR. 2006. Evidence that feedback inhibition of NAD kinase

controls   responses   to   oxidative   stress.   *Proc   Natl   Acad   Sci   USA*   **103**:7601–7606.

doi:10.1073/pnas.0602494103

Grosz M, Kolter J, Paprotka K, Winkler AC, Schäfer D, Chatterjee SS, Geiger T, Wolz C,

Ohlsen K, Otto M, Rudel T, Sinha B, Fraunholz M. 2014. Cytoplasmic replication of

Staphylococcus aureus upon phagosomal escape triggered by phenol-soluble modulin alpha.

Cell Microbiol **16**:451-65. doi: 10.1111/cmi.12233

Harper L, Balasubramanian D, Ohneck EA, Sause WE, Chapman J, Mejia-Sosa B, Lhakhang

T, Heguy A, Tsirigos A, Ueberheide B, Boyd JM, Lun DS, Torres VJ. 2018. *Staphylococcus*

aureus responds to the central metabolite pyruvate to regulate virulence. *mBio* **9**:2272–2289.

doi:10.1128/mBio.02272-17

Herbert S, Bera A, Nerz C, Kraus D, Peschel A, Goerke C, Meehl M, Cheung A, Götz F. 2007.

Molecular basis of resistance to muramidase and cationic antimicrobial peptide activity of

lysozyme in staphylococci. *PLoS Pathog* **3**:e102. doi:10.1371/journal.ppat.0030102

Kawai S, Mori S, Mukai T, Suzuki S, Yamada T, Hashimoto W, Murata K. 2000. Inorganic

polyphosphate/ATP-NAD kinase of *Micrococcus flavus* and *Mycobacterium tuberculosis*

H37Rv. *Biochem Biophys Res Commun* **276**:57–63. doi:10.1006/bbrc.2000.3433

Kitur K, Parker D, Nieto P, Ahn DS, Cohen TS, Chung S, Wachtel S, Bueno S, Prince A. 2015.

Toxin-induced necroptosis is a major mechanism of *Staphylococcus aureus* lung damage.

PLoS Pathog. **11**:e1004820. doi: 10.1371/journal.ppat.1004820.

Kobayashi K, Ehrlich SD, Albertini A, Amati G, Andersen KK, Arnaud M, Asai K, Ashikaga

S, Aymerich S, Bessieres P, Boland F, Brignell SC, Bron S, Bunai K, Chapuis J, Christiansen

LC, Danchin A, Débarbouillé M, Dervyn E, Deuerling E, Devine K, Devine SK, Dreesen O,

Errington J, Fillinger S, Foster SJ, Fujita Y, Galizzi A, Gardan R, Eschevins C, Fukushima

T, Haga K, Harwood CR, Hecker M, Hosoya D, Hullo MF, Kakeshita H, Karamata D,

Kasahara Y, Kawamura F, Koga K, Koski P, Kuwana R, Imamura D, Ishimaru M, Ishikawa

S, Ishio I, le Coq D, Masson A, Mauël C, Meima R, Mellado RP, Moir A, Moriya S,

Nagakawa E, Nanamiya H, Nakai S, Nygaard P, Ogura M, Ohanan T, O'Reilly M, O'Rourke

M, Pragai Z, Pooley HM, Rapoport G, Rawlins JP, Rivas LA, Rivolta C, Sadaie A, Sadaie

Y, Sarvas M, Sato T, Saxild HH, Scanlan E, Schumann W, Seegers JFML, Sekiguchi J,

Sekowska A, Séror SJ, Simon M, Stragier P, Studer R, Takamatsu H, Tanaka T, Takeuchi

M, Thomaides HB, Vagner V, van Dijl JM, Watabe K, Wipat A, Yamamoto H, Yamamoto

M, Yamamoto Y, Yamane K, Yata K, Yoshida K, Yoshikawa H, Zuber U, Ogasawara N.

2003. Essential *Bacillus subtilis* genes. *Proc Natl Acad Sci USA* **100**:4678–4683.

doi:10.1073/pnas.0730515100

Kobayashi SD, Malachowa N, Whitney AR, Braughton KR, Gardner DJ, Long D, Bubeck

Wardenburg J, Schneewind O, Otto M, Deleo FR. 2011. Comparative analysis of USA300

virulence determinants in a rabbit model of skin and soft tissue infection. *J Infect Dis*

204:937-41. doi: 10.1093/infdis/jir441

Kornberg A. 1950. Enzymatic synthesis of triphosphopyrine nucleotide. *J Biol Chem* **182**:805–

813.

Kreiswirth BN, Löfdahl S, Betley MJ, O'Reilly M, Schlievert PM, Bergdoll MS, Novick RP.

1983. The toxic shock syndrome exotoxin structural gene is not detectably transmitted by a

prophage. *Nature* **305**:709–712. doi:10.1038/305709a0

Levraud J-P, Colucci-Guyon E, Redd MJ, Lutfalla G, Herbomel P. 2008. *In vivo* analysis of

zebrafish innate immunity. *Methods Mol Biol* **415**:337–363. doi:10.1007/978-1-59745-570-

1_20

Liebermeister W, Noor E, Flamholz A, Davidi D, Bernhardt J, Milo R. 2014. Visual account

of protein investment in cellular functions. *Proc Natl Acad Sci USA* **111**:8488-8493.

doi.org/10.1073/pnas.1314810111

Li B Bin, Wang X, Tai L, Ma TT, Shalmani A, Liu WT, Li WQ, Chen KM. 2018. NAD kinases:

Metabolic targets controlling redox co-enzymes and reducing power partitioning in plant

stress and development. *Front Plant Sci*. **9**:379. doi:10.3389/fpls.2018.00379

Li W, Wu H, Li M, San KY. 2018. Effect of NADPH availability on free fatty acid production

in *Escherichia coli*. *Biotechnol Bioeng* **115**:444–452. doi:10.1002/bit.26464

McElroy MC, Harty HR, Hosford GE, Boylan GM, Pittet JF, Foster TJ. 1999. Alpha-toxin

damages the air-blood barrier of the lung in a rat model of *Staphylococcus aureus*-induced

pneumonia. *Infect Immun* **67**:5541-4. doi: 10.1128/IAI.67.10.5541-5544.1999

Mailloux RJ, Lemire J, Appanna VD. 2011. Metabolic networks to combat oxidative stress in

Pseudomonas fluorescens. *Antonie van Leeuwenhoek* **99**:433–442. doi:10.1007/s10482-

010-9538-x

McGuinness ET, Butler JR. 1985. NAD^+ kinase - a review. *Int J Biochem* **17**:1–11.

Mostowy S, Boucontet L, Mazon Moya MJ, Sirianni A, Boudinot P, Hollinshead M, Cossart P,

Herbomel P, Levraud JP, Colucci-Guyon E. 2013. The zebrafish as a new model for the *in*

vivo study of *Shigella flexneri* interaction with phagocytes and bacterial autophagy. *PLoS*

Pathog **9**:12–16. doi:10.1371/journal.ppat.1003588

Nilsson IM, Hartford O, Foster T, Tarkowski A. 1999. Alpha-toxin and gamma-toxin jointly

promote *Staphylococcus aureus* virulence in murine septic arthritis. *Infect Immun.* **67**:1045-

9. doi: 10.1128/IAI.67.3.1045-1049.1999. PMID: 10024541; PMCID: PMC96427.

O'Callaghan RJ, Callegan MC, Moreau JM, Green LC, Foster TJ, Hartford OM, Engel LS, Hill

JM. 1997. Specific roles of alpha-toxin and beta-toxin during *Staphylococcus aureus* corneal

infection. *Infect Immun* **65**:1571-8. doi: 10.1128/iai.65.5.1571-1578.1997

Patel AH, Nowlan P, Weavers ED, Foster T. 1987. Virulence of protein A-deficient and alpha-

toxin-deficient mutants of *Staphylococcus aureus* isolated by allele replacement. *Infect*

Immun **55**:3103-10. doi: 10.1128/iai.55.12.3103-3110.1987

Peschel A, Otto M. 2013. Phenol-soluble modulins and staphylococcal infection. *Nat Rev*

Microbiol **11**:667-73. doi: 10.1038/nrmicro3110

Phan QT, Sipka T, Gonzalez C, Levraud J-P, Lutfalla G, Nguyen-Chi M. 2018. Neutrophils use

superoxide to control bacterial infection at a distance. *PLoS Pathog* **14**:e1007157.

doi:10.1371/JOURNAL.PPAT.1007157

Pidwill GR, Gibson JF, Cole J, Renshaw SA, Foster SJ. 2021. The role of macrophages in

Staphylococcus aureus infection. *Front Immunol* **11**:3506. doi:10.3389/fimmu.2020.620339

Pollitt EJG, Szkuta PT, Burns N, Foster SJ. 2018. *Staphylococcus aureus* infection dynamics.

PLoS Pathog **14**:e1007112. doi:10.1371/journal.ppat.1007112

Potter AD, Butrico CE, Ford CA, Curry JM, Trenary IA, Tummarakota SS, Hendrix AS, Young

JD, Cassat JE. 2020. Host nutrient milieu drives an essential role for aspartate biosynthesis

during invasive *Staphylococcus aureus* infection. *Proc Natl Acad Sci USA* **117**:12394–

12401. doi:10.1073/pnas.1922211117

Pounds S, Cheng C. 2006. Robust estimation of the false discovery rate. *Bioinformatics*
22:1979-87.

Prajsnar TK, Cunliffe VT, Foster SJ, Renshaw SA. 2008. A novel vertebrate model of
Staphylococcus aureus infection reveals phagocyte-dependent resistance of zebrafish to
non-host specialized pathogens. *Cell Microbiol* **10**:2312–2325. doi:10.1111/j.1462-
5822.2008.01213.x

Prajsnar TK, Hamilton R, Garcia‑Lara J, McVicker G, Williams A, Boots M, Foster SJ,
Renshaw SA. 2012. A privileged intraphagocyte niche is responsible for disseminated
infection of *Staphylococcus aureus* in a zebrafish model. *Cell Microbiol* **14**:1600–1619.
doi:10.1111/j.1462-5822.2012.01826.x

Queck SY, Jameson-Lee M, Villaruz AE, Bach TH, Khan BA, Sturdevant DE, Ricklefs SM,
Li M, Otto M. 2008. RNAIII-independent target gene control by the *agr* quorum-sensing
system: insight into the evolution of virulence regulation in *Staphylococcus aureus*. *Mol Cell*
32:150-8. doi: 10.1016/j.molcel.2008.08.005

Renshaw SA, Loynes CA, Trushell DMI, Elworthy S, Ingham PW, Whyte MKB. 2006. A
transgenic zebrafish model of neutrophilic inflammation. *Blood* **108**:3976–3978.
doi:10.1182/BLOOD-2006-05-024075

Richardson AR. 2019. Virulence and metabolism. *Microbiol Spectr* **7**.
doi:10.1128/microbiolspec.gpp3-0011-2018

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. Limma powers
differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids*
Res **43**:e47. doi: 10.1093/nar/gkv007

Rosenberger CM, Gallo RL, Finlay BB. 2004. Interplay between antibacterial effectors: a
macrophage antimicrobial peptide impairs intracellular *Salmonella* replication. *Proc Natl*
Acad Sci USA **101**:2422-7. doi:10.1073/pnas.0304455101

Rothfork JM, Timmins GS, Harris MN, Chen X, Lusis A, Otto M, Cheung AL, Greshan HD.
2004. Inactivation of a bacterial virulence pheromone by phagocyte-derived oxidants: new
role for the NADPH oxidase in host defense. *Proc Natl Acad Sci USA* **101**:13867-72. doi:
10.1073/pnas.0402996101

Ruiz JM, Sánchez E, García PC, López-Lefebre LR, Rivero RM, Romero L. 2002. Proline
metabolism and NAD kinase activity in greenbean plants subjected to cold-shock.
Phytochemistry **59**:473–478. doi:10.1016/S0031-9422(01)00481-2

Sassetti CM, Boyd DH, Rubin EJ. 2003. Genes required for mycobacterial growth defined by
high density mutagenesis. *Mol Microbiol* **48**:77–84. doi:10.1046/j.1365-2958.2003.03425.x

Seilie ES, Bubeck Wardenburg J. 2017. *Staphylococcus aureus* pore-forming toxins: the
interface of pathogen and host complexity. *Semin Cell Dev Biol* **72**:101-116. doi:
10.1016/j.semcdb.2017.04.003

Singh R, Mailloux RJ, Puiseux-Dao S, Appanna VD. 2007. Oxidative stress evokes a metabolic
adaptation that favors increased NADPH synthesis and decreased NADH production in
Pseudomonas fluorescens. *J Bacteriol* **189**:6665–6675. doi:10.1128/JB.00555-07

Ślaski JJ. 1995. NAD^+ kinase activity in root tips of nearly isogenic linesof wheat (*Triticum*
aestivum L.) that differ in their tolerance to aluminium. *J Plant Physiol* **145**:143–147.
doi:10.1016/S0176-1617(11)81861-4

Smyth, GK. 2005. Limma: Linear models for microarray data. In: Gentleman, R., Carey, V.J.,
Huber, W., Irizarry, R.A., Dudoit, S. (eds) Bioinformatics and computational biology
solutions using R and bioconductor. Statistics for biology and health. Springer, New York,
NY. https://doi.org/10.1007/0-387-29362-0.23s

Spaan AN, Vrieling M, Wallet P, Badiou C, Reyes-Robles T, Ohneck EA, Benito Y, de Haas

CJ, Day CJ, Jennings MP, Lina G, Vandenesch F, van Kessel KP, Torres VJ, van Strijp JA,

Henry T. 2014. The staphylococcal toxins γ-haemolysin AB and CB differentially target

phagocytes by employing specific chemokine receptors. *Nat Commun* **5**:5438. doi:

10.1038/ncomms6438

Sun F, Liang H, Kong X, Xie S, Cho H, Deng X, Ji Q, Zhang H, Alvarez S, Hicks LM, Bae T,

Luo C, Jiang H, He C. 2012. Quorum-sensing *agr* mediates bacterial oxidation response via

an intramolecular disulfide redox switch in the response regulator AgrA. *Proc Natl Acad Sci*

USA **109**:9095-100. doi: 10.1073/pnas.1200603109

Svoboda KR, Linares AE, Ribera AB. 2001. Activity regulates programmed cell death of

zebrafish Rohon-Beard neurons. *Development* **128**:3511-20. doi: 10.1242/dev.128.18.3511

Tai L, Li B-B, Nie X-M, Zhang P-P, Hu C-H, Zhang L, Liu W-T, Li W-Q, Chen K-M. 2019.

Calmodulin is the fundamental regulator of NADK-mediated NAD signaling in plants. *Front*

Plant Sci **10**:681. doi:10.3389/fpls.2019.00681

Teoh WP, Chen X, Laczkovich I, Alonzo F 3rd. 2021. *Staphylococcus aureus* adapts to the host

nutritional landscape to overcome tissue-specific branched-chain fatty acid requirement.

Proc Natl Acad Sci USA **118**. doi:10.1073/PNAS.2022720118

Tomlinson KL, Lung TWF, Dach F, Annavajhala MK, Gabryszewski SJ, Groves RA, Drikic

M, Francoeur NJ, Sridhar SH, Smith ML, Khanal S, Britto CJ, Sebra R, Lewis I, Uhlemann

AC, Kahl BC, Prince AS, Riquelme SA. 2021. *Staphylococcus aureus* induces an itaconate-

dominated immunometabolic response that drives biofilm formation. *Nat Commun* **12**.

doi:10.1038/s41467-021-21718-y

Turner NA, Sharma-Kuinkel BK, Maskarinec SA, Eichenberger EM, Shah PP, Carugati M,

Holland TL, Fowler VG. 2019. Methicillin-resistant *Staphylococcus aureus*: an overview of

basic and clinical research. *Nat Rev Microbiol* **17**:203-218. doi:10.1038/s41579-018-0147-

4

Turner WL, Waller JC, Vanderbeld B, Snedden WA. 2004. Cloning and characterization of two

NAD kinases from *Arabidopsis*. Identification of a calmodulin binding isoform. *Plant*

Physiol **135**:1243–1255. doi:10.1104/pp.104.040428

Tyanova S, Temu T, Cox J. 2016. The MaxQuant computational platform for mass

spectrometry-based shotgun proteomics. *Nat Protoc* **11**:2301-2319.

Vestergaard M, Frees D, Ingmer H. 2019. Antibiotic resistance and the MRSA problem.

Microbiol Spectr **7**. doi:10.1128/microbiolspec.GPP3-0057-2018

Vestin R. 1937. Enzymic change from codehydrogenase I to codehydrogenase II.

Naturwissenschaften **25**:667–8.

Von Euler H, Adler E. 1938. Enzymic intertransformation of codehydrogenase I and

codehydrogenase II. *Z Physiol Chem* **252**:41–8.

Wang R, Braughton KR, Kretschmer D, Bach THL, Queck SY, Li M, Kennedy AD, Dorward

DW, Klebanoff SJ, Peschel A, DeLeo FR, Otto M. 2007. Identification of novel cytolytic

peptides as key virulence determinants for community-associated MRSA. *Nat Med* **13**:1501-

14. doi: 10.1038/nm1656

Wieczorek S, Combes F, Lazar C, Giai Gianetto Q, Gatto L, Dorffer A, Hesse AM, Couté Y,

Ferro M, Bruley C, Burger T. 2017. DAPAR & ProStaR: software to perform statistical

analyses in quantitative discovery proteomics. *Bioinformatics* **33**: 135-136.

Zhao C, Shu X, Sun B. 2017. Construction of a gene knockdown system based on catalytically

inactive ("dead") Cas9 (dCas9) in *Staphylococcus aureus*. *Appl Environ Microbiol* **83**.

doi:10.1128/AEM.00291-17

Figure supplement 1 NADK is important for *S. aureus* growth. (A) Total RNAs of *S. aureus* containing the empty

vector (pSD1) or the NADK knockdown strain (NADK sgRNA) were analyzed by RT-PCR using oligonucleotides

specific to the target gene *ppnK* or the control gene *gyrA*. (B) Bacterial protein extracts of *S. aureus* containing the

empty vector (pSD1) or the NADK knockdown strain (NADK sgRNA) were analyzed by immunoblotting using

antibodies against the target protein NADK or against the control protein EF-Tu. (C) Bacterial growth of *S. aureus*

containing the empty vector (pSD1) or the NADK knockdown strain (NADK sgRNA) monitored for 6 hours at

OD_{600nm} in BHI broth at 37°C upon ATc induction. Data shown are representative of three independent

experiments. The bar indicates the standard error of the means of biological replicates. Comparison of data was

performed using a t test (*$p<0.05$).

1007

Figure supplement 2 NADK contributes to *S. aureus* virulence in zebrafish. (A) The number of pSD1 (grey) or

NADK sgRNA (blue) *S. aureus* was monitored in zebrafish larvae after intravenous injection with 5.10^3 bacteria.

For each strain, CFU were determined 0, 6, 24 and 48 hpi by plating fish lysates on BHI agar (open circles) and

BHI agar supplemented with chloramphenicol (triangles). (B) Representative fluorescence confocal images of

transgenic *mpx:GFP* zebrafish larvae fixed and labeled with specific antibodies to detect GFP or *S. aureus,*

uninfected (CTRL), or intravenously injected with 10^4 *S. aureus* containing the empty vector (pSD1) or the NADK

knockdown strain (NADK sgRNA) at 0 or 24 hpi. Maximum intensity Z-projection images (2-μm serial optical

sections) of bacteria (red) and neutrophils (green) in the caudal and head regions. Scale bars, 25 μm. (C)

Representative fluorescence confocal images of transgenic *mfap4::mCherryF* zebrafish larvae fixed and labeled

with specific antibodies to detect mCherry or *S. aureus*, uninfected (CTRL), or intravenously injected with 10^4 *S.*

aureus containing the empty vector (pSD1) or the NADK knockdown strain (NADK sgRNA) at 24 hpi, showing

macrophages (red). White arrows indicate the injection site. Scale bars, 500 μm.

1020

1021

Figure supplement 3 NADK promotes expression of *S. aureus* virulence determinants. The *S. aureus* strain

containing the empty vector (pSD1) and the *ppnK* knockdown strain (NADK sgRNA) were grown at 37°C in BHI

broth supplemented with H_2O_2. (A-C) Whole bacterial cell lysates were analyzed by LC-MS/MS. Voronoi

treemaps were generated to visualize proteins more abundant in *S. aureus*/pSD1 than in the knockdown strain at

three hierarchical levels according to KEGG database: top level (A), second level (B), third level (C). Colors

represent functional categories. Category size based on LFQ intensity corresponds to differences in protein

abundance.

1029

1031

1032

Figure supplement 4 *S. aureus ppnK* gene is part of the *relQ* operon. (A) Genomic region surrounding the *ppnK* gene of *S. aureus* Xen36 between nucleotides 947626 and 955045. Bars and numbers represent the regions amplified by RT-PCR shown in B. (B) Ethidium bromide staining of DNA in an agarose gel following RT-PCR amplification using total RNA from *S. aureus* Xen36 and primers located in regions indicated in A. (C) Ethidium bromide staining of DNA in an agarose gel following RT-PCR amplification using total RNA from *S. aureus* Xen36/pSD1 or Xen36/NADK sgRNA and oligonucleotides specific to the target gene *rluA* or the control gene *gyrA*, respectively.

1040

1041

1042 **Table supplement 1.** Proteins more abundant in the *S. aureus* pSD1 strain than in the NADK
1043 sgRNA strain after growth in BHI broth (Excel file).

1044
1045

1046 **Table supplement 2.** Relative abundance of a subset of protective proteins differentially
1047 expressed by *S. aureus* pSD1 and NADK sgRNA strains exposed to hydrogen peroxide.
1048

Uniprot	Protein	Description	Log2R*	P**
Q2FV54	OatA	O-acetyl transferase	3.40	6.57E-05
Q2G261	SodM	Superoxide dismutase [Mn/Fe]	2.49	1.66E-07
Q2G000	Trx2	Thioredoxin 2	2.25	1.00E-04
Q2G0E0	GraR	Response regulator protein	2.00	7.15E-06
Q2FZZ3	-	Thioredoxin domain-containing protein	1.86	3.25E-04
Q2G280	-	Peroxiredoxin	1.67	7.37E-08
P0A086	MsrA2	Methionine sulfoxide reductase	1.37	1.27E-04
Q2FYU7	KatA	Catalase	1.17	1.83E-08
Q2FVL7	-	Thioredoxin domain-containing protein	1.02	2.22E-03

1049
1050 * Log2R=Log2[pSD1]/[NADK sgRNA]
1051 **Adjusted p value

1052
1053

1054 **Table supplement 3.** Proteins more abundant in the *S. aureus* pSD1 strain than in the NADK
1055 sgRNA strain after growth in BHI broth supplemented with hydrogen peroxide (Excel file).

1056
1057

1058 **Table supplement 4.** Relative abundance of a subset of protective proteins differentially
1059 expressed by *S. aureus* pSD1 and NADK sgRNA strains upon zebrafish infection.
1060

Uniprot	Protein	Description	Log2R*	P**
Q2G000	-	Thioredoxin domain-containing protein	+	NA
P0A0J3	SodA	Superoxide dismutase [Mn]	+	NA
Q2FYU7	KatA	Catalase	+	NA
Q2FKL3	Tpx	Thiol peroxidase	+	NA
Q2FZW4	DltC	D-alanyl carrier protein	+	NA
O05204	AhpF	Alkyl hydroperoxide reductase subunit F	+	NA
Q2FXI6	-	Thioredoxin domain-containing protein	+	NA
Q2G280	-	Peroxiredoxin	+	NA

1061
1062 * Log2R=Log2[pSD1]/[NADK sgRNA]; +: protein detected in pSD1 strain and not detected from NADK sgRNA strain
1063 **Adjusted p value: NA: not applicable
1064

1065

1066 **Table supplement 5.** Proteins more abundant in the *S. aureus* pSD1 strain than in the NADK
1067 sgRNA strain upon zebrafish infection (Excel file).

1068
1069
1070

Table supplement 6. Relative abundance of a subset of virulence-related proteins differentially expressed by *S. aureus* pSD1 and NADK sgRNA strains exposed to hydrogen peroxide.

Uniprot	Protein	Description	Log2R*	P**
Q2FVK1	HlgB	Gamma-hemolysin component B	+	NA
Q2FWM8	Hld	Delta-hemolysin	+	NA
Q2FWM4	AgrA	Accessory gene regulator protein	+	NA
Q2FVK2	HlgC	Gamma-hemolysin component C	+	NA
P0C818	PsmA4	Phenol-soluble modulin alpha 4	+	NA
P0C7Y1	PsmA1	Phenol-soluble modulin alpha 1	+	NA
Q2FUU5	Lip1	Lipase	+	NA
Q2G1X0	Hly	Alpha-hemolysin	7.99	2.92E-09
Q2FVK3	HlgA	Gamma-hemolysin component A	5.59	1.60E-06
Q2FWP0	LukL1	Leukocidin-like protein 1	3.35	7.16E-06
Q2G2R8	SspP	Staphopain A	3.27	1.17E-06
Q2FWN9	LukL2	Leukocidin-like protein 2	2.98	2.56E-07

*Log2R=Log2[pSD1]/[NADK sgRNA]; +: protein detected in pSD1 strain and not detected from NADK sgRNA strain
**Adjusted p value: NA: not applicable