Pseudomonas aeruginosa C-terminal processing protease CtpA assembles into a

hexameric structure that requires activation by a spiral-shaped lipoprotein

binding partner

Hao-Chi Hsu[a], Michelle Wang[b], Amanda Kovach[a], Andrew J. Darwin[b,#], Huilin Li[a,#]

[a]Department of Structural Biology, Van Andel Institute, Grand Rapids, Michigan, USA

[b]Department of Microbiology, New York University Grossman School of Medicine, New

York, New York, USA

Running head: Structural analysis of the CtpA-LbcA system

Address correspondence to Andrew J. Darwin, andrew.darwin@med.nyu.edu or Huilin

Li, Huilin.Li@vai.org

Abstract: 172

Text: 4,994 (excluding Abstract, Importance, References, Tables and Figure legends)

ABSTRACT

Pseudomonas aeruginosa CtpA is a carboxyl terminal-processing protease that partners with the outer membrane lipoprotein LbcA to degrade at least five cell wall-associated proteins, four of which are cell wall hydrolases. This activity plays an important role in supporting *P. aeruginosa* virulence in a mouse model of acute pneumonia. However, almost nothing is known about the molecular mechanisms underlying CtpA and LbcA function. Here, we used structural analysis to show that CtpA alone assembles into an inactive hexamer comprising a trimer of dimers, which limits its substrate access and prevents nonspecific degradation. The adaptor protein LbcA is a right-handed open spiral with 11 tetratricopeptide repeats, which might wrap around a substrate to deliver it to CtpA for degradation. By structure-guided mutagenesis and functional assays, we also showed that the interfaces of the CtpA trimer-of-dimers, and an N-terminal helix of LbcA, are important for LbcA-mediated substrate degradation by CtpA both *in vitro* and *in vivo*. This work improves our understanding of the molecular mechanism of a CTP within the C-terminal processing peptidase-3 group.

IMPORTANCE

Carboxyl-terminal processing proteases (CTPs) are found in all three domains of life. In

bacteria, some CTPs have been associated with virulence, raising the possibility that

they could be theraputic targets. However, relatively little is known about their molecular

mechanisms of action. In *Pseudomonas aeruginosa,* CtpA supports virulence by

working in complex with the outer membrane lipoprotein LbcA to degrade cell wall

hydrolases. Here, we report structure-function analyses of CtpA and LbcA, which

reveals that CtpA assembles into an inactive hexamer comprising a trimer of dimers.

LbcA is monomeric, with an N-terminal region important for binding to and activating

CtpA, followed by a spiral structure composed of 11 tetratricopetide repeats, which

could wrap around a substrate for delivery to CtpA. This work provides the first structure

of a CTP-3 group member, revealing a unique mutimeric arrangement and insight into

how this important proteolytic system functions.

INTRODUCTION

A eubacterial cell is protected by a mesh-like cell wall of peptidoglycan (PG), which is composed of linear glycan strands with peptide side chains that cross-link to each other through peptide bonds (1). To accommodate growth, these cross-links must be cleaved so that nascent PGs can be inserted into the network (1-3). Several PG endopeptidases carry out this hydrolysis, including MepS, MepM, and MepH in *Escherichia coli* (4, 5). However, if not tightly regulated, their activity could lead to rupture of the PG sacculus and cell death. One way to control endopeptidase activity is through the relatively recently discovered carboxyl-terminal processing (CTP) protease system. CTPs belong to the S41 family of serine proteases (6). All CTPs have a PDZ domain — named because it was first noted in: postsynaptic density protein of 95 kDa, Drosophila disc large tumor suppressor, and zonula occludens-1 protein — which plays roles in substrate recognition and protease regulation (7, 8). CTPs work within the cell envelope of Gram-negative and Gram-positive bacteria and have been linked to virulence (9-14).

The *E. coli* CTP Prc partners with the lipoprotein Nlpl to cleave the PG endopeptidase MepS (5). Prc is a bowl-shaped monomer, and the Nlpl adaptor forms a homodimer that binds to two separate molecules of Prc (15). In *Bacillus subtilis*, the CtpB protease processes and activates the intramembrane protease 4FA-4FB complex, thereby regulating spore formation (16, 17). CtpB has N-terminal and C-terminal dimerization domains, plus a cap domain preceding the protease core domain. Structural analysis revealed that CtpB assembles a dimeric self-compartmentalizing ring structure (18). The substrate peptide enters the proteolytic site via a narrow tunnel that is largely sequestered by the PDZ domain and becomes exposed only in the presence

of a substrate. Therefore, the CtpB protease is reversibly activated by the substrate C-

terminal peptide. In contrast to Prc and CtpB, *P. aeruginosa* CtpA has been assigned to

the C-terminal processing peptidase-3 group (19). No structural studies have been

reported so far for this group.

Pseudomonas aeruginosa is an opportunistic human pathogen. It is one of the

leading causes of sepsis in intensive care units, and outbreaks of multidrug-resistant

strains have been reported in hospitals (20-22). In contrast to *E. coli*, in which the only

CTP present is Prc, *P. aeruginosa* has both the C-terminal processing peptidase-1

group member Prc, and the C-terminal processing peptidase-3 group member CtpA (10,

19). We reported previously that CtpA is required for normal function of the type 3

secretion system and for virulence in a mouse model of acute pneumonia (10). *P.*

aeruginosa CtpA has 39% amino acid sequence identity with *B. subtilis* CtpB, which

suggests that they might share a similar fold. However, CtpA has a much longer C-

terminal region. This extended C-terminus might alter the oligomerization mode relative

to that of CtpB (18). However, it is not known if or how CtpA assembles into a self-

compartmentalizing structure to prevent nonspecific proteolysis. Unlike CtpB, CtpA is

not directly activated by a protein substrate. Instead, it requires the adaptor protein

LbcA, the <u>l</u>ipoprotein <u>b</u>inding partner of <u>C</u>tpA, for activity *in vivo* and *in vitro* (23).

LbcA is predicted to be an outer membrane lipoprotein, and to contain 11

tetratricopeptide repeats (TPRs). The TPR motif is a degenerate 34-amino-acid

sequence that mediates protein-protein interactions (24-26). LbcA promotes CtpA

protease activity, and *ctpA* and *lbcA* null mutants share common phenotypes, such as a

defective type 3 secretion system and accelerated surface attachment (10, 23, 27). Five

LbcA-CtpA substrates have been reported to date, and four of them are predicted to be

PG cross-link hydrolases: the LytM/M23 family peptidases MepM and PA4404, and the

NlpC/P60 family peptidases PA1198 and PA1199 (23, 27). Therefore, it appears that

LbcA interacts with CtpA to assemble an active proteolytic complex, which controls the

activity of these enzymes by degrading them. However, the molecular mechanisms

underlying CtpA and LbcA function are unknown. Here, we describe structural and

functional analyses of CtpA and LbcA. We show that CtpA alone assembles as an

inactive trimer of dimers and that LbcA is a right-handed spiral that might wrap around a

substrate protein. Structure-guided mutagenesis confirms the functional importance of

CtpA interfaces and identifies the interface between CtpA and LbcA.

RESULTS

CtpA assembles a trimer-of-dimers hexamer in solution. CtpA is located in the

periplasm, tethered to the outer membrane via its interaction with LbcA (23). Its N-

terminal 23 residues are a type I signal sequence, which is followed by a 14-residue

region that is enriched in alanine, glycine, and proline that contributes to a bacterial low

complexity region and is predicted to be disordered (28). Therefore, we removed these

37 amino acids to produce a truncated CtpA protein for structural and functional studies

(Supplemental Table 1, Fig. 1a). For simplicity, we refer to this ΔN37 construct as CtpA

throughout the main text. The CtpA crystals diffracted X-rays to 3.5-Å resolution. We

first used molecular replacement with the *B. subtilis* CtpB as a search model to solve

the CtpA structure, but this did not lead to a satisfactory solution. We then produced

selenomethionine-substituted CtpA crystals for SAD-based phasing. The derivatized

crystals diffracted to 3.3 Å, leading to successful structural solution (Supplemental Table

2). CtpA formed a hexamer that comprised a trimer of dimers (Fig. 1b). This oligomer

state is consistent with an estimated mass of a hexamer from the gel filtration profile

(Fig. 1c). As expected, each CtpA protomer consists of an N-terminal dimerization

region (NDR), a PDZ domain, a cap domain, a protease core domain, and a C-terminal

dimerization region (CDR) (Fig. 1d). The PDZ domain is partially disordered, but we

were able to generate a homolog PDZ model based on the CtpB structure and

accurately docked the domain guided by two anomalous density peaks from

selenomethionine residues 153 and 160 of the domain (Supplemental Figure 1). The

loop connecting S10 and H6 (aa's 378–411) within the CDR was disordered, and the

last two residues (435-436) were not resolved.

The CtpA hexamer alone in the absence of LbcA is in an inactive

configuration. The CtpA core region contains the protease active site. There is a

narrow tunnel dividing the core domain into the upper cap and lower body regions (Fig.

1d). In CtpB this tunnel was suggested to guide the substrate peptide into the proteolytic

site (18). The cap of CtpA is a four-stranded β-sheet (S4, S5, S8, S9) located below the

NDR. The body region is composed of a three-helix bundle (H3, H4, H5) and a five-

stranded β-sheet (S1, S2, S3, S6, S7). The CtpA catalytic Ser-302 is located at the end

of the narrow tunnel between the cap and the body region.

The CtpA hexamer is in an inactive conformation. The PDZ domain is in a

position that blocks the narrow substrate peptide tunnel leading to Ser-302. Furthermore,

the catalytic triad Ser-302–Lys-327–Gln-331 is well beyond hydrogen-bonding distance:

Ser-302 and Lys-327 are 4.2 Å apart and Lys-327 and Gln-331 are 9.1 Å apart (Fig. 1e).

We solved the crystal structure of CtpA that was catalytically inactive due to an S302A

substitution and found that it was also in the inactive configuration (Supplemental Figure

2). For comparison, the catalytic triad Ser-309–Lys-334–Gln-338 in the protease-active

B. subtilis CtpB are all within hydrogen-bonding distance (Fig. 1e). By superimposing

the two proteases, we found that transition to the active form requires the CtpA cap

domain to clamp down toward the catalytic site by 11 Å and also a large-scale

movement of the associated PDZ domain by 12 Å (Fig. 1f, Supplemental Video 1). The

CtpA structure is consistent with our previous observation that purified wild-type CtpA

alone was inactive in degrading its substrates (23). The addition of purified LbcA was

able to activate the protease of wild-type CtpA but not of the mutant CtpA(S302A).

Therefore, unlike CtpB, which fluctuates between an active and inactive form in solution

and can be activated by a protein substrate (18), the CtpA hexamer is locked in an

inactive configuration and requires LbcA for activation (23).

The C-terminal dimerization interface is important for full CtpA activity. *P.*

aeruginosa CtpA dimerizes via its NDR in the same way as *B. subtilis* CtpB (Fig. 2a).

However, the two CDRs of CtpA are far apart in this dimer, and six unhinged CDRs of

three CtpA dimers interact to form the triangular trimer-of-dimer complex (Figs. 1b, 2a).

In contrast, CtpB forms a parallel homodimer by both head-to-head N-terminal and tail-

to-tail C-terminal interactions (Fig. 2b) (18). The CtpA NDR is composed of the H1 and

H2 helices. The NDRs of two CtpA monomers form a domain-swapped, intermolecular

4-helix bundle, in which the NDR of one protomer reaches over to contact the cap

domain of the other protomer (Fig. 2a, 2c). The four-helix bundle is at the vertex of the

triangular complex. The NDR–NDR dimerization is driven by hydrophobic interactions.

Specifically, Leu-69, Ala-73, and Met-77 are at the intersecting point of the two crossed

H2's. Met-77 hydrophobically interacts with Ala-73 and Leu-69 of the second protomer

(Fig. 2c). H1 of protomer 1 is nearly parallel to H2 of protomer 2, with the H1 residues

Leu-46, Phe-49, Val-52, Leu-53, and Val-56 hydrophobically interacting with the H2

residues Leu-70, Ile-74, Met-77, Leu-78, and Leu-81. The Leu-46 and Phe-49 of the two

H1's are also within the van der Waals distance. The hydrophobic NDR–NDR

interaction of the CtpA dimer resembles that in the CtpB dimer, consistent with the

conserved sequence in this region (Fig. 2e).

The C-terminal dimerization interface of CtpA involves the β-strand S10 and the

helix H6 (Fig. 2d). The two S10 β-strands (Arg-370 to Glu-376) form an intermolecular,

antiparallel β-sheet that reinforces the dimer interface. The two H6's (Tyr-418 to Gly-435)

are orthogonal to each other but form a short leucine zipper in the middle section

mediated by Leu-426 and Leu-430. The CtpA C-terminal dimer interface is not like that

of CtpB, which dimerizes via interaction in a different region. Indeed, the C-terminal

sequence of *P. aeruginosa* CtpA is not conserved in *B. subtilis* CtpB (Fig. 2e), which is

consistent with their different oligomeric assembly. To investigate the functional

importance of the unique C-terminal dimerization interface of CtpA, we constructed a

mutant with a partially disrupted H6 helix by removing the last six residues Ser-431 to

Asn-436; CtpA(ΔC6). We found that CtpA(ΔC6) eluted from a gel filtration column as a

dimer (Fig. 3a), so the C-terminal truncation prevented CtpA hexamer formation.

We next asked whether the C-terminal dimerization interface, present in the

intact CtpA hexamer but disrupted in the CtpA(ΔC6) dimer, is required for normal CtpA

function. In an *in vitro* assay, we found that the CtpA(ΔC6) dimer was less active than

the CtpA hexamer in degrading the model substrate PA1198, in reactions that also

contained LbcA (compare lanes 3 and 4 in Fig. 3b).

To further investigate the functional significance of the leucine zipper in H6, we

generated two CtpA double-mutant constructs with L426K/L430K and L426A/L430A

substitutions that disrupted the leucine zipper interaction. The two purified mutant

proteins were mainly dimeric and failed to assemble into a hexamer, based on their gel

filtration profiles (Fig. 3a). Their LbcA-activated *in vitro* protease activity toward PA1198

was lower relative to wild-type CtpA (Fig. 3b, compare lanes 5 and 6 to lane 3).

However, like the CtpA(ΔC6) mutant, the leucine zipper mutants retained significant

protease activity relative to the protease dead mutant CtpA(S302A) (Fig. 3b, compare

lanes 5 and 6 with lane 2).

We extended our analysis by constructing plasmids encoding derivatives of full-

length CtpA with the ΔC6, L426K/L430K (LLKK), or L426A/L430A (LLAA) mutations.

After introducing these plasmids into a *P. aeruginosa* Δ*ctpA* mutant, immunoblot

analysis showed that the steady-state levels of the mutant proteins were similar to the

wild-type (Fig. 3c). Therefore, the mutations did not affect CtpA stability *in vivo*.

However, PA1198 accumulated in the presence of these three mutants relative to the

wild-type, although not as much as in the presence of CtpA(S302A) (Fig. 3c). This

suggests that the protease activity of the three mutants was reduced, but not abolished,

which is consistent with the *in vitro* analysis (Fig. 3b). Therefore, our studies suggest

that the hexameric assembly is important for proper CtpA function.

LbcA forms a spiral that could wrap around CtpA substrates. During the

maturation of the lipoprotein LbcA, its N-terminal 16 residues are removed, exposing

Cys-17 (Fig. 4a). Then Cys-17 is lipidated so that the N-terminus can be anchored in

the outer membrane from the periplasmic side (29-31). LbcA is able to bind to CtpA and

its substrates independently (27). Therefore, to begin to understand how LbcA might be

capable of these separate interactions, we tried to solve the LbcA crystal structure. We

were able to crystallize LbcA with its N-terminal 48 residues removed (ΔN48). The

crystal structure was solved to 3.5-Å resolution by the SAD method with

selenomethionine-derivatized LbcA(ΔN48) crystals (Fig. 4a-d). This structure contains

only α-helices and connecting loops, with a total of 29 α-helices. We resolved all 11

predicted tetratricopeptide repeats, comprising 22 α-helices from helix-7 (H7) to H28.

The sequences of the 34-residue TPRs are largely conserved (Fig. 4b). Gly/Ala at

position 8 and Ala at positions 20 and 27 are most conserved in the TPR family,

although none of these are invariant. Interestingly, TPR1, TPR5, and TPR10 contain

only 33 amino acids, but they all contain the signature residues Gly/Ala at position 8 and

Ala at position 20, and TPR1 and TPR10 also contain a signature Leu at position 24.

Within each TPR, the first helix (TPR-A) lines the inner surface and the second helix

(TPR-B) lines the outer surface of the ring structure. Alignment of the 11 TPRs in

LbcA(ΔN48) showed that more-conserved hydrophobic residues are concentrated in the

TPR-A helix and more-conserved charged residues are distributed in TPR-B helix and

in the turns connecting TPR-A and TPR-B, suggesting that the inner and the outer

surfaces of the ring may have distinct functions.

The 11 TPRs form a notched ring, with an outer diameter of about 6 nm and an

inner diameter of about 3 nm (Fig. 4c). It is a right-handed spiral structure, because the

first TPR is slightly above, and the last TPR is below the ring. The first 4 α-helices at the

N terminus form an elongated extension that partially caps the TPR ring. Helices H5

and H6 serve as a hinge that links the N-terminal helical extension and the TPR spiral.

However, the loop connecting H4 and H5 (aa's 163-171) has a high crystallographic B-

factor (150-200 $Å^2$) that is indicative of flexibility. Therefore, this loop and the H5-H6

hinge are likely to allow relative motion of the NT helical extension with respect to the

TPR spiral. The TPR ring is capped at the end by the single short α-helix, H29.

LbcA(ΔN48) purified as a monomer in solution (Fig. 5a-b). However, four

LbcA(ΔN48) molecules formed an interlocked tetramer as a dimer of dimers in the

crystal (Fig. 5c-d). The chamber of the TPR spiral of the first LbcA is occupied by the

H3-H4 of a second LbcA molecule on the top and by the H1-H2 of a third LbcA

molecule on the side (Fig. 5d). Therefore, there is a 4-helix bundle inside the first LbcA

TPR spiral. We suggest that this bundle may mimic a substrate and that the LbcA spiral

may wrap around a substrate to target it to CtpA for degradation (Fig. 5e). In this

scenario, the conserved and hydrophobic residues of the TPR-A helices lining the inner

surface of the TPR spiral may participate in binding that substrate.

The LbcA H1 is essential for binding to and activating the CtpA protease.

To identify the binding interface between CtpA and LbcA, we produced a series of N-

and/or C-terminal-truncated LbcA mutants and did a CtpA pulldown assay (Fig. 6a). We

found that removing the C-terminal five residues of LbcA did not affect the LbcA–CtpA

interaction. All constructs containing the complete H1-H29 region pulled down CtpA.

However, LbcA with either H1 truncation (His$_6$-LbcA$\triangle$N84) or H1-H4 truncation (His$_6$-

LbcA$\triangle$N165) failed to pull down CtpA. This result suggests that the LbcA NT helical

extension, in particular H1, is required for LbcA binding to CtpA, and that LbcA H1

directly participates in the binding.

We then carried out both *in vitro* and *in vivo* assays to monitor degradation of the

PA1198 substrate (23). We first incubated purified CtpA with separately purified LbcA

proteins and PA1198. As expected, PA1198 was degraded in the presence of His$_6$-

LbcA$\triangle$N48, which had the intact NT helical extension (Fig. 6b). However, PA1198 was

not degraded in the presence of His$_6$-LbcA($\triangle$N84) (missing H1) or His$_6$-LbcA($\triangle$N165)

(missing H1-H4). In fact, the outcomes of reactions with these two truncations were

indistinguishable from those containing no LbcA, or those using the protease dead

CtpA(S302) (Fig. 6b). In a $\triangle$*lbcA* mutant *P. aeruginosa* strain, plasmids encoding LbcA

with in frame deletions equivalent to the $\triangle$N84 or $\triangle$N165 constructs failed to activate

CtpA, as revealed by the accumulation of PA1198, which was present at a similar level

as in a strain without any LbcA (Fig. 6c). Therefore, both pulldowns and *in vitro* and *in*

vivo activity assays pinpointed the LbcA H1 as a key binding element for activation of

the CtpA protease.

DISCUSSION

The LbcA–CtpA system supports the function of the *P. aeruginosa* type 3 secretion system, is required for virulence in a mouse model of acute infection, and affects surface attachment (10, 23). Furthermore, four CtpA substrates are PG cross-link hydrolases, which means that the LbcA–CtpA system affects the integrity of a crucial cell envelope component and perhaps the most important antibiotic target, the cell wall. Therefore, this system could be an effective antibiotic target, and the structural analysis reported here may aid the development of such antibiotics. Here, we have shown that CtpA assembles as a hexamer. However, the hexamer alone is inactive, because the catalytic triad Ser302–Lys327–Gln331 is not in hydrogen-bonding distance. We also found that the CtpA partner protein LbcA has an N-terminal helical region that is needed to bind to CtpA, and a large spiral cavity that has the capacity to wrap around a substrate for delivery to CtpA.

How the interaction with LbcA converts CtpA into an active protease is currently unclear. Elucidation of the activation mechanism requires the determination of the CtpA-LbcA complex. However, our previous experiments suggested that most, if not all, CtpA in the cell is bound to LbcA (23). CtpA fractionates with the membrane fraction in *lbcA*$^+$ cells, but is in the soluble periplasmic fraction in Δ*lbcA* cells, and when CtpA or LbcA is purified from *P. aeruginosa*, they copurify with a lot of the other one (23). Also, the pulldown assays done here show that LbcA and CtpA can interact in the absence of substrate (Fig. 6a). Therefore, it is possible that the LbcA-CtpA complex fluctuates between inactive and active states, and perhaps the presence of a substrate would

stabilize the enzyme-adaptor-substrate in the active state for productive substrate

hydrolase degradation.

The PDZ domain is the C-terminal peptide substrate-binding element of CTPs

(32). The ability of PDZ domains to move in order to accommodate the incoming C-

terminal substrate peptide partially accounts for the substrate delivery–based activation

mechanism of the CTPs. The PDZ domain is the most flexible region in the CtpB

structure and acts as an inhibitor by blocking the substrate peptide binding in the

inactive form, but moves away to form a narrow tunnel for a substrate peptide in the

active form (18). Similarly, the PDZ domain of CtpA is highly flexible and largely invisible

in the crystal structure of the inactive CtpA hexamer, suggesting a CtpA activation

mechanism similar to that of CtpB. In the case of Prc from *E. coli*, the PDZ domain acts

as an activator rather than an inhibitor, but its movement is still a key feature of the

activation mechanism (8).

P. aeruginosa CtpA requires the partner lipoprotein LbcA for activation and

targeted proteolysis (23). This is a variation in the general substrate activation

mechanism. In this regard, the LbcA–CtpA system is analogous to the *E. coli* NlpI–Prc

system, in which the TPR-containing adaptor NlpI plays the dual function of delivering

the PG hydrolase MepS to the Prc protease for degradation, and activating the protease

(8, 15). Despite this functional similarity, significant differences exist between these two

systems. The two proteases are in different C-terminal processing peptidase families; *E.*

coli Prc is much larger than *P. aeruginosa* CtpA; and Prc functions as a monomer, not a

hexamer. Therefore, the Prc N-terminal and C-terminal helical domains are not involved

in oligomerization; instead, they wrap around the protease core, perhaps to limit

substrate access to the protease. NlpI has a total of 14 helices with four TPRs,

compared with the 29 helices and 11 TPRs of LbcA. Although NlpI and LbcA both

contain TPRs, the primary sequences of the proteins are not similar. Finally, NlpI

functions as a dimer, unlike the monomeric state of LbcA.

The LbcA–CtpA system contributes to the virulence of *P. aeruginosa*. This makes

understanding the structure and function of this proteolytic system broadly significant.

Future challenges include understanding exactly how LbcA activates CtpA, and how

protein substrates, not just peptides, are specifically recognized by the multitude of

CTPs and/or their adaptor proteins for tightly regulated proteolysis.

MATERIALS AND METHODS

Purification of CtpA. DNA encoding amino acid 38 to the C-terminus of CtpA

was sucloned into plasmid pET15b between the NdeI and XhoI sites. The plasmid

encoded N-terminal His$_6$-tagged CtpA(ΔN37). Similar plasmids encoding CtpA(ΔC6),

CtpA-L426K L430K or CtpA-L426A L430A were constructed using reverse primers that

incoprporated the C-terminal mutations. *E. coli* BL21(DE3) transformants were grown at

37°C to OD$_{600}$ = 0.6-0.7 before being induced with 0.5 mM IPTG and incubated at 16°C

overnight. Cells were harvested by centrifugation and lysed by passing through a

microfluidizer cell disruptor in 10 mM potassium phosphate, pH 8.0, 10 mM imidazole,

0.25 M NaCl. The homogenate was clarified by centrifuging at 27,000 x *g* and the

supernatant was applied to a HiTrap-Ni column (GE Healthcare) pre-equilibrated with

the lysis buffer. Proteins were eluted with a 10–300 mM imidazole gradient in 10 mM

potassium phosphate, pH 8.0, containing 0.25 M NaCl. Fractions containing His$_6$-CtpA

were collected. The N-terminal His tag was removed using thrombin (0.5 units/mg) by

dialyzing against 20 mM Tris, pH 8.0, 150 mM NaCl overnight at 4 °C. Untagged CtpA

was further purified with HiTrap-Q in 10 mM Tris, pH 8.0, and a 50–500 mM NaCl

gradient and polished by gel filtration in 10 mM Tris, pH 8.0, and 150 mM NaCl using

Superdex 200 prep grade column (16 x 600 mm, GE Healthcare).

Purification of LbcA. DNA encoding amino acid 49 to the C-terminus of LbcA

(LbcA[ΔN48]) was subcloned into the NdeI and HindIII sites of plasmid pET24b.

Plasmids encoding N-terminal His$_6$-tagged LbcA proteins were constructed similarly by

subcloning fragments into the NdeI and BamHI sites of plasmid pET15b. *E. coli*

BL21(DE3) transformants were grown at 37°C to OD$_{600}$ = 0.5 before being induced with

0.5 mM IPTG and incubated at 37°C for another 3 h. His$_6$-tagged LbcA protein was

purified with HiTrap-Ni in 10 mM potassium phosphate, pH 8.0, 0.25 M NaCl, and a 10–

300 mM imidazole gradient, followed by HiTrap-Q in 10 mM Tris, pH 8.0, and a 50–500

mM NaCl gradient. The final polish of LbcA was done in a Superdex 200 prep-grade

column preequilibrated with 10 mM Tris, pH 8.0, and 150 mM NaCl.

Protein crystallization and structural solution. CtpA(ΔN37) was crystallized at

20°C by the sitting-drop vapor diffusion method using 0.1 M sodium acetate, pH 4.0,

and 0.6 M ammonium dihydrogen phosphate at a concentration of 33 mg/mL. Diffraction

data were collected at the Lilly Research Laboratories Collaborative Access Team

(LRL-CAT) beamline of the Advanced Photon Source (APS) at Argonne National

Laboratory and processed with Mosflm software. Se-derived crystals diffracted to 3.3 Å,

which was better than the native crystals, so the Se-derived data were used to solve the

CtpA structure. Autosol of Phenix was used to locate Se sites and the resulting map

was used to build the initial model. However, the map quality from the SAD method was

not good enough to build the CtpA model due to a low anomalous signal (only 4 high-

occupancy Se were identified in one asymmetric unit). In combination with molecular

replacement, the structure of CtpA(ΔN37) was determined by combining MR-SAD using

the core domain of *B. subtilis* CtpB (PDB ID 4C2C) as the search model. The structure

of CtpA(ΔN37, S302A) was solved by PHASER of Phenix using CtpA(ΔN37) as the

search model.

LbcA(ΔN48) was crystallized at 20°C by the sitting-drop vapor diffusion method

using 0.1 M sodium acetate, pH 4.6, and 1.9 M ammonium dihydrogen phosphate at a

concentration of 45 mg/mL. Diffraction data were collected at the Life Sciences

Collaborative Access Team (LS-CAT) beamline of APS and were processed with

Mosflm. The best data set was from Se-derived crystals which diffracted to 3.5 Å, so the

Se-derived diffraction data were used to solve the LbcA structure. Se sites were

determined by the SAD method using Autosol of Phenix (33). All models of CtpA and

LbcA were built in Coot (34) and refined with Phenix.refine (35).

The crystal structures of CtpA, CtpA(S302A), and LbcA have been deposited in

the Protein Data Bank under accession codes 7RPQ, 7RQH, and 7RQF, respectively.

***In vitro* proteolysis assay.** CtpA and LbcA proteins were purified as described

above. To purify His_6-PA1198, *E. coli* strain M15 [pREP4] (Qiagen) containing

pAJD2948 was grown in 500 mL of LB broth at 37°C to an OD_{600} of 0.6-1.0. Protein

production was induced by adding 1 mM IPTG and incubation at 37 °C for 3 h, after

which cells were harvested by centrifugation. His_6-PA1198 was purified under native

conditions by nickel–nitrilotriacetic acid–agarose affinity chromatography in buffer

containing 50 mM NaH_2PO_4 and 300 mM NaCl, as recommended by the manufacturer

(Qiagen). Protein was eluted in 50 mM NaH_2PO_4, 300 mM NaCl, 50 mM imidazole, pH 8.

Assays were done as described previously (23, 36). Reactions were terminated by

adding SDS-PAGE sample buffer and incubating at 90°C for 10 min. Samples were

separated by SDS-PAGE and stained with ProtoBlue Safe (National Diagnostics). For

experiments with mutant CtpA proteins, ImageJ analysis was used to determine the

densities of the His_6-PA1198 bands. The amount of His_6-PA1198 degraded in each

reaction was determined by comparing His_6-PA1198 band density to that in the reaction

with inactive CtpA(S302A). Average values from two independent experiments are

reported.

In vivo **CtpA activity assay**. Wild-type *ctpA* was amplified from *P. aeruginosa*

PAK chromosomal DNA using a primer that annealed about 30 bp upstream of the *ctpA*

start codon and a reverse primer that annealed immediately downstream of the stop

codon. A fragment encoding CtpA(△C6) was generated with a reverse primer that

annealed downstream of codon 430 and incorporated a stop codon. Fragments

encoding CtpA-L426K L430K or CtpA-L426A L430A were generated with reverse

primers that incorporated the mutagenic mismatches. A fragment encoding CtpA(S302A)

was generated by amplifying *ctpA* from *P. aeruginosa* strain AJDP1140 DNA using the

same primer pairs used to amplify wild-type *ctpA*. All fragments were cloned into

pHERD26T using EcoRI and XbaI restriction sites added by the amplification primers.

Fragments encoding LbcA(△N84) and LbcA(△N165) were generated by

amplifying one fragment with a forward primer that annealed ~ 40 bp upstream of the

lbcA start codon and a reverse primer that annealed at codon 31, and a second

fragment with a forward primer that annealed at codon 85 or codon 166 and a reverse

primer that annealed at the *lbcA* stop codon. The forward primer for the second

fragments had a 5′ tail complementary to the end of the first fragment (codons 25-31).

The first fragment and second fragments were joined by splicing overlap extension PCR

(37). The resulting fragments encoded LbcA with internal deletions that removed

codons 32-84 (△N84) or codons 32-165 (△N165). Codons 1-31 were retained, encoding

the signal sequence followed by 15 amino acids corresponding to the N-terminus of

mature wild-type LbcA, to ensure normal signal sequence processing, lipidation, and

outer membrane trafficking. A fragment encoding wild-type *lbcA* was generated by

amplifying *lbcA* from *P. aeruginosa* DNA using a forward primer that annealed ~ 40 bp

upstream of the start codon and a reverse primer that annealed at the *lbcA* stop codon.

All fragments were cloned into pHERD26T using EcoRI and HindIII restriction sites

added by the amplification primers.

Plasmids were introduced into $\Delta ctpA$ or $\Delta lbcA$ mutants by electroporation (38).

Saturated cultures were diluted into 5 mL of LB broth containing 75 µg/mL tetracycline,

in 18-mm diameter test tubes, at OD_{600} of 0.05. Cultures were grown on a roller drum at

37 °C for 5 h. Cells were harvested by centrifugation and resuspended in SDS-PAGE

sample buffer at equal concentrations based on culture OD_{600}. Samples were separated

by SDS-PAGE and transferred to nitrocellulose by semi-dry electroblotting.

Chemiluminescent detection followed incubation with polyclonal antisera against CtpA,

LbcA, or PA1198, and then goat anti-rabbit IgG (Sigma) horseradish peroxidase

conjugate.

ACKNOWLEDGEMENTS

Diffraction data for CtpA were collected at the Lilly Research Laboratories Collaborative

Access Team (LRL-CAT) beamline and diffraction data of LbcA were collected at the

Life Sciences Collaborative Access Team (LS-CAT) beamline of the Advanced Photon

Source (APS). The APS is a U.S. Department of Energy (DOE) Office of Science User

Facility operated for the DOE Office of Science by Argonne National Laboratory under

Contract No. DE-AC02-06CH11357. Cryo-EM data were collected at the David Van

Andel Advanced Cryo-Electron Microscopy Suite at Van Andel Institute. We thank

Dolonchapa Chakraborty for supplying plasmid pAJD2948 and His$_6$-PA1198 protein,

Gongpu Zhao and Xing Meng for help with data collection, and David Nadziejka for

technical editing of the manuscript. This study was supported by U.S. National Institutes

of Health grant R01 AI136901 (to A.D.) and by Van Andel Institute (to H.L.). The content

is solely the responsibility of the authors and does not necessarily represent the official

views of the National Institutes of Health.

REFERENCES

1.   Vollmer W, Blanot D, de Pedro MA. 2008. Peptidoglycan structure and architecture. FEMS Microbiol Rev 32:149-67.

2.   Pazos M, Peters K, Vollmer W. 2017. Robust peptidoglycan growth by dynamic and variable multi-protein complexes. Curr Opin Microbiol 36:55-61.

3.   Vollmer W. 2012. Bacterial growth does require peptidoglycan hydrolases. Mol Microbiol 86:1031-5.

4.   Park SH, Kim YJ, Lee HB, Seok YJ, Lee CR. 2020. Genetic Evidence for Distinct Functions of Peptidoglycan Endopeptidases in *Escherichia coli*. Front Microbiol 11:565767.

5.   Singh SK, Parveen S, SaiSree L, Reddy M. 2015. Regulated proteolysis of a cross-link-specific peptidoglycan hydrolase contributes to bacterial morphogenesis. Proc Natl Acad Sci U S A 112:10956-61.

6.   Rawlings ND, Barrett AJ, Bateman A. 2010. MEROPS: the peptidase database. Nucleic Acids Res 38:D227-33.

7.   Ye F, Zhang M. 2013. Structures and target recognition modes of PDZ domains: recurring themes and emerging pictures. Biochem J 455:1-14.

8.   Chueh CK, Som N, Ke LC, Ho MR, Reddy M, Chang CI. 2019. Structural Basis for the Differential Regulatory Roles of the PDZ Domain in C-Terminal Processing Proteases. mBio 10.

9.   Bandara AB, Sriranganathan N, Schurig GG, Boyle SM. 2005. Carboxyl-terminal protease regulates *Brucella suis* morphology in culture and persistence in macrophages and mice. J Bacteriol 187:5767-75.

10.   Seo J, Darwin AJ. 2013. The *Pseudomonas aeruginosa* periplasmic protease CtpA can affect systems that impact its ability to mount both acute and chronic infections. Infect Immun 81:4561-70.

11.   Deng CY, Deng AH, Sun ST, Wang L, Wu J, Wu Y, Chen XY, Fang RX, Wen TY, Qian W. 2014. The periplasmic PDZ domain-containing protein Prc modulates full virulence, envelops stress responses, and directly interacts with dipeptidyl peptidase of *Xanthomonas oryzae* pv. oryzae. Mol Plant Microbe Interact 27:101-12.

12.   Carroll RK, Rivera FE, Cavaco CK, Johnson GM, Martin D, Shaw LN. 2014. The lone S41 family C-terminal processing protease in *Staphylococcus aureus* is localized to the cell wall and contributes to virulence. Microbiology (Reading) 160:1737-1748.

13.   Bandara AB, DeShazer D, Inzana TJ, Sriranganathan N, Schurig GG, Boyle SM. 2008. A disruption of *ctpA* encoding carboxy-terminal protease attenuates *Burkholderia mallei* and induces partial protection in CD1 mice. Microb Pathog 45:207-16.

14.   Nash ZM, Cotter PA. 2019. Regulated, sequential processing by multiple proteases is required for proper maturation and release of *Bordetella* filamentous hemagglutinin. Mol Microbiol 112:820-836.

15.   Su MY, Som N, Wu CY, Su SC, Kuo YT, Ke LC, Ho MR, Tzeng SR, Teng CH, Mengin-Lecreulx D, Reddy M, Chang CI. 2017. Structural basis of adaptor-mediated protein degradation by the tail-specific PDZ-protease Prc. Nat Commun 8:1516.

16. Campo N, Rudner DZ. 2006. A branched pathway governing the activation of a
developmental transcription factor by regulated intramembrane proteolysis. Mol
Cell 23:25-35.
17. Rudner DZ, Losick R. 2002. A sporulation membrane protein tethers the pro-
sigmaK processing enzyme to its inhibitor and dictates its subcellular localization.
Genes Dev 16:1007-18.
18. Mastny M, Heuck A, Kurzbauer R, Heiduk A, Boisguerin P, Volkmer R, Ehrmann
M, Rodrigues CD, Rudner DZ, Clausen T. 2013. CtpB assembles a gated
protease tunnel regulating cell-cell signaling during spore formation in *Bacillus*
subtilis. Cell 155:647-58.
19. Hoge R, Laschinski M, Jaeger KE, Wilhelm S, Rosenau F. 2011. The subcellular
localization of a C-terminal processing protease in *Pseudomonas aeruginosa*.
FEMS Microbiol Lett 316:23-30.
20. Paterson DL. 2006. The epidemiological profile of infections with multidrug-
resistant *Pseudomonas aeruginosa* and *Acinetobacter species*. Clin Infect Dis 43
Suppl 2:S43-8.
21. Tumbarello M, De Pascale G, Trecarichi EM, Spanu T, Antonicelli F, Maviglia R,
Pennisi MA, Bello G, Antonelli M. 2013. Clinical outcomes of *Pseudomonas*
aeruginosa pneumonia in intensive care unit patients. Intensive Care Med
39:682-92.
22. Crivaro V, Di Popolo A, Caprio A, Lambiase A, Di Resta M, Borriello T, Scarcella
A, Triassi M, Zarrilli R. 2009. *Pseudomonas aeruginosa* in a neonatal intensive
care unit: molecular epidemiology and infection control measures. BMC Infect
Dis 9:70.
23. Srivastava D, Seo J, Rimal B, Kim SJ, Zhen S, Darwin AJ. 2018. A Proteolytic
Complex Targets Multiple Cell Wall Hydrolases in *Pseudomonas aeruginosa*.
mBio 9.
24. Graham JB, Canniff NP, Hebert DN. 2019. TPR-containing proteins control
protein organization and homeostasis for the endoplasmic reticulum. Crit Rev
Biochem Mol Biol 54:103-118.
25. Zeytuni N, Zarivach R. 2012. Structural and functional discussion of the tetra-
trico-peptide repeat, a protein interaction module. Structure 20:397-405.
26. Blatch GL, Lassle M. 1999. The tetratricopeptide repeat: a structural motif
mediating protein-protein interactions. Bioessays 21:932-9.
27. Chakraborty D, Darwin AJ. 2021. Direct and indirect interactions promote
complexes of the lipoprotein LbcA, the CtpA protease and its substrates, and
other cell wall proteins in *Pseudomonas aeruginosa*. J Bacteriol
doi:10.1128/JB.00393-21:JB0039321.
28. Ntountoumi C, Vlastaridis P, Mossialos D, Stathopoulos C, Iliopoulos I,
Promponas V, Oliver SG, Amoutzias GD. 2019. Low complexity regions in the
proteins of prokaryotes perform important functional roles and are highly
conserved. Nucleic Acids Res 47:9998-10009.
29. Konovalova A, Silhavy TJ. 2015. Outer membrane lipoprotein biogenesis: Lol is
not the end. Philos Trans R Soc Lond B Biol Sci 370:20150030.
30. Lewenza S, Mhlanga MM, Pugsley AP. 2008. Novel inner membrane retention
signals in *Pseudomonas aeruginosa* lipoproteins. J Bacteriol 190:6119-25.

31. Narita S, Tokuda H. 2007. Amino acids at positions 3 and 4 determine the
membrane specificity of *Pseudomonas aeruginosa* lipoproteins. J Biol Chem
282:13372-8.
32. Clausen T, Kaiser M, Huber R, Ehrmann M. 2011. HTRA proteases: regulated
proteolysis in protein quality control. Nat Rev Mol Cell Biol 12:152-62.
33. Adams PD, Afonine PV, Bunkoczi G, Chen VB, Echols N, Headd JJ, Hung LW,
Jain S, Kapral GJ, Grosse Kunstleve RW, McCoy AJ, Moriarty NW, Oeffner RD,
Read RJ, Richardson DC, Richardson JS, Terwilliger TC, Zwart PH. 2011. The
Phenix software for automated determination of macromolecular structures.
Methods 55:94-106.
34. Casanal A, Lohkamp B, Emsley P. 2020. Current developments in Coot for
macromolecular model building of Electron Cryo-microscopy and
Crystallographic Data. Protein Sci 29:1069-1078.
35. van Zundert GCP, Moriarty NW, Sobolev OV, Adams PD, Borrelli KW. 2021.
Macromolecular refinement of X-ray and cryoelectron microscopy structures with
Phenix/OPLS3e for improved structure and ligand quality. Structure 29:913-921
e4.
36. Chung S, Darwin AJ. 2020. The C-terminus of substrates is critical but not
sufficient for their degradation by the *Pseudomonas aeruginosa* CtpA protease. J
Bacteriol doi:10.1128/JB.00174-20.
37. Heckman KL, Pease LR. 2007. Gene splicing and mutagenesis by PCR-driven
overlap extension. Nat Protoc 2:924-932.
38. Choi KH, Kumar A, Schweizer HP. 2006. A 10-min method for preparation of
highly electrocompetent *Pseudomonas aeruginosa* cells: application for DNA
fragment transfer between chromosomes and plasmid transformation. J Microbiol
Methods 64:391-397.
39. Strom MS, Lory S. 1986. Cloning and expression of the pilin gene of
Pseudomonas aeruginosa PAK in *Escherichia coli*. J Bacteriol 165:367-372.
40. Studier FW, Rosenberg AH, Dunn JJ, Dubendorff JW. 1990. Use of T7 RNA
polymerase to direct expression of cloned genes. Methods in Enzymology
185:60-89.
41. Qiu D, Damron FH, Mima T, Schweizer HP, Yu HD. 2008. PBAD-based shuttle
vectors for functional analysis of toxic and highly regulated genes in
Pseudomonas and *Burkholderia* spp. and other bacteria. Appl Environ Microbiol
74:7422-7426.
577

 FIGURES AND LEGENDS

FIG 1 Overall structure of CtpA. (a) Top: CtpA domain organization. The dashed lines
indicate the disordered regions (PDZ and the S10-H6 connecting loop) that are not
resolved in the crystal structure. NDR, N-terminal dimerization region; CDR, C-terminal
dimerization region. The sequence ranges of the two CtpA constructs used this study
are shown in the lower panels. (b) Cartoon and transparent surface views of CtpA
hexamer. The domains are colored according to the depiction in panel a. (c) SDS-PAGE
analysis and gel filtration profile of CtpA). (d) A CtpA subunit in cartoon view. Secondary
structural elements in CtpA are labeled, except in the PDZ domain. The two dashed
arrows indicate the mobile PDZ domain in the CtpA hexamer. (e) Comparison of the
catalytic triads of Pa CtpA and Bs CtpB (PDB ID 4C2E). The S8 and S9 labels refer to
β-strands 8 and 9 in the cap region. (f) Superposition of the core domains of inactive Pa
CtpA (green) and active Bs CtpB (cyan). Catalytic Ser-302 in CtpA and Ser-309 in CtpB
are in red sticks. Red arrows indicate lifted-up (CtpA) and clamped-down positions
(CtpB) of the cap subdomains.
594

FIG 2 Different oligomerization modes of Pa CtpA and Bs CtpB. (a) A CtpA dimer extracted from the CtpA hexamer. (b) Bs CtpB dimer (PDB ID 4C2E). (c) The N-terminal dimerization interface of CtpA involves hydrophobic interactions between two H2 (left) and between H1 and H2 (right). (d) The C-terminal dimerization interface of CtpA involves a short leucine zipper-like interaction between two H6 helices and antiparallel β-sheet formation between two S10. (e) Alignment of the conserved NDR sequence and divergent CDR between Pa CtpA and Bs CtpB.

Fig. 3. **Full protease activity of CtpA requires C-terminal dimerization region. a**)
Elution profiles of wild-type and mutant CtpA proteins. **b**) Substrate degradation assay
in vitro. His$_6$-PA1198 served as the substrate for the assay. Gels from a single
experiment are shown, but the amount of PA1198 degradation is the average from two
independent experiments, determined as described in Methods. The number below
each lane is the percentage of remaining PA1198 after 3 h, relative to the first lane
using the inactive CtpA. **c**) Substrate degradation *in vivo*. Plasmid-encoded protease-
dead CtpA(S302A), wild-type (WT), ΔC6, L426K/L430K (LLKK), L426A/L430A (LLAA)
were produced in a *P. aeruginosa* Δ*ctpA* strain. None = empty plasmid vector control.
The CtpA proteins and accumulation of the PA1198 substrate were detected by
immunoblot analysis with polyclonal antisera.

FIG 4 Overall structure of LbcA. (a) Domain organization of LbcA. The TPR motifs are shown in magenta. Also shown are two LbcA constructs used in this study. (b) Sequence alignment of the 11 TPRs of LbcA. The signature residues of TPR are marked in the red rectangles. The highly homologous hydrophobic residues are in green boxes and the highly homologous charged residues are in the blue boxes. (c) The crystal structure of LbcA(ΔN48) contains an N-terminal extension with 4 helices and a C-terminal superhelix composed of 11 TPRs. The thick gray curve in the left panel follows the right-handed spiral feature of the TPRs.

FIG 5 LbcA is a monomer in solution but assembles a tetramer in crystal. (a)
Coomassie blue stained SDS-PAGE gel of the purified of LbcAΔN48. (b) Superdex-200
elution profile of LbcAΔN48. LbcA was eluted from gel filtration column at a volume
corresponding to the monomeric state. (c) LbcA forms an intertwined tetramer in the
crystal lattice. The four protomers are individually colored. (d) This LbcA tetramer view
shows that the helices H1 and H4 of protomer 2 and helices H2 and H3 of protomer 3
form a 4-helix bundle inside the super helical coil of protomer 1 in the crystal lattice. (e)
H2H3 of a second LbcA (magenta cartoon and transparent surface) and H1H4 of a third

LbcA (light blue cartoon and transparent surface) insert into the first LbcA spiral, likely
mimicking the substrate binding by the first LbcA spiral.
646

647
648

FIG 6 The LbcA N-terminal extension is essential for the protease activity of CtpA. (a) CtpA pull-
down assay using various N-terminal and C-terminal deletion mutants of LbcA. Lane 2 is the
CtpA input control (C). Lane 3 is the background binding of CtpA to Ni beads. (b) *In vitro*
substrate (PA1198) degradation assay. (c) Substrate degradation *in vivo*. Plasmids encoding
wild-type LbcA (WT), LbcA(ΔN84), or LbcA(ΔN165) were transformed into a *P. aeruginosa*
Δ*lbcA* mutant. None = empty plasmid vector control. The LbcA proteins and accumulation of the
PA1198 substrate were detected by immunoblot analysis with polyclonal antisera.

Supplemental Table 1. Strains and plasmids used in this study

Nam	Genotype/Features	Reference or Source
***P. aeruginosa* strains**		
PAK	wild-type PAK strain	(39)
AJDP730	PAK Δ*ctpA*	(23)
AJDP1091	PAK Δ*lbcA::aacC1*	(23)
AJDP1140	PAK *ctpA-S302A*	(23)
***E. coli* strain**		
BL21(DE3)	F$^-$ *ompT gal [dcm] [lon] hsdS$_B$ (r$_B^-$ m$_B^-$; E. coli* B strain) λDE3	(40)
Plasmids		
pET15b	Amp^r, pMB1 *ori*, *T7p* expression vector	Novagen
pET24b	Kan^r, pMB1 *ori*, *T7p* expression vector	Novagen
pHERD26T	Tet^r, pMB1 *ori*, *araBp* expression vector	(41)
pQE-30	Amp^r, Col E1 *ori*, *T5p* expression vector	Qiagen
pAJD2948	*T5p-his$_6$*-PA1198 in pQE-30	(27)
pAJD3037	*araBp-ctpA* in pHERD26T	This study
pAJD3038	*araBp-ctpAΔC6* in pHERD26T	This study
pAJD3039	*araBp-ctpA-L426K L430K* in pHERD26T	This study
pAJD3045	*araBp-ctpA-S302A* in pHERD26T	This study
pAJD3063	*araBp-ctpA-L426A L430A* in pHERD26T	This study
pAJD3109	*araBp-lbcA* in pHERD26T	This study
pAJD3110	*araBp-lbcAΔN84* in pHERD26T	This study
pAJD3111	*araBp-lbcAΔN165* in pHERD26T	This study
pET15b_CtpAΔN37	*T7p-his$_6$-ctpAΔN37* in pET15b	This study

683	pET15b_CtpAΔN37ΔC6	*T7p-his₆-ctpAΔN37ΔC6* in pET15b	This study
684	pET15b_CtpAΔN37_S302A	*T7p-his₆-ctpAΔN37_S302A* in pET15b	This study
685	pET15b_CtpAΔN37_L426K/L430K	*T7p-his₆-ctpAΔN37_L426K L430K* in pET15b	This study
686	pET15b_CtpAΔN37_L426A/L430A	*T7p-his₆-ctpAΔN37_L426A L430A* in pET15b	This study
687	pET24b_LbcAΔN48	*T7p-lbcAΔN48-his₆* in pET24b	This study
688	pET15b_LbcAΔN31	*T7p-his₆-lbcAΔN31* in pET15b	This study
689	pET15b_LbcAΔN48	*T7p-his₆-lbcAΔN48* in pET15b	This study
690	pET15b_LbcAΔN84	*T7p-his₆-lbcAΔN84* in pET15b	This study
691	pET15b_LbcAΔN165	*T7p-his₆-lbcAΔN165* in pET15b	This study
692	pET15b_LbcAΔN48ΔC5	*T7p-his₆-lbcAΔN48ΔC5* in pET15b	This study

694 **Supplemental Table 2. Crystallographic data collection and refinement statistics**

	CtpAΔN37_Se	CtpA(ΔN37, S302A)	LbcAΔN48_Se
Data collection			
Wavelength (Å)	0.97931	1.07803	0.97872
Space group	H3	H3	P 32 2 1
Cell dimensions			
a, b, c (Å)	187.49, 187.49, 132.01	189.65, 189.65, 131.10	120.70, 120.70, 221.94
α, β, γ (°)	90.00, 90.00, 120.00	90.00, 90.00, 120.00	90.00, 90.00, 120.00
Resolution (Å)	93.75 – 3.30 (3.48 -3.30)*	94.82 – 3.2 (3.37 – 3.2)*	94.57 – 3.50 (3.69 – 3.50)*
R_{merge} (%)	16.7 (91.2)	8.9 (142)	18.3 (226)
$I / \sigma I$	10.8 (2.9)	15.6 (2.0)	13.2 (2.0)
Total reflections	301572 (44008)	297808 (44906)	536619 (78420)
Completeness (%)	100 (100)	99.6 (99.9)	100 (100)
Redundancy	11.6 (11.4)	10.3 (10.6)	22.1 (22.5)
Refinement			
Resolution (Å)	39.83 – 3.30	43.03 – 3.2	60.39 – 3.50
No. reflections	26026	28849	24277
R_{work} / R_{free}	0.2375 / 0. 2636	0.2422/0.2654	0.2470 / 0.2806
No. of non-hydrogen atoms	3933	3931	8352
Macromolecule	3933	3931	8352
Ligand	0	0	0
Water	0	0	0
B-factors	91.58	126.34	135.73
Macromolecule	91.58	126.34	135.73
Ligand			
Water			
R.m.s. deviations			
Bond lengths (Å)	0.003	0.003	0.003
Bond angles (°)	0.757	0.739	0.608
Ramachandran statistics (%)			
Favored	96.83	97.02	97.89
Allowed	3.17	2.98	2.11
Outliers	0	0	0

*Values in parentheses are for the last (highest) resolution shell.

Supplemental Figure 1. Se-Met peaks superimposed with PDZ homolog model. (A)

The overall electron density map of one asymmetric unit obtained from the Se-

derivatized CtpA hexamer crystal. The 2mFo–DFc electron density map is rendered at

1σ threshold and shown as gray meshes. The non-PDZ domains are shown in green

and the PDZ domains are show in magenta. The positions of Se are in orange. (B)

Enlarged view of the left PDZ region in (A).

Supplemental Figure 2. Crystal structure of CtpA(S302A) showing only a monomer. (A)

overall structure of CtpA(S302A) in ribbons, with the catalytic triad residues in sticks and

the point mutation S302A in red sticks. (B) Enlarged view of the catalytic triad in the

CtpA(S302A) crystal structure. The structure is also in the inactive configuration.

Supplemental Video 1. CtpA hexamer structure. Rotation of the hexamer, then

transition to a monomer to show the domain structure, then transition to morphing

between inactive and active computational model (based on CtpB).