1   Development of motor neurons and motor activity in zebrafish requires F-actin

2   nucleation by Fmn2b

Dhriti Nagar[1], Blake Carrington[2], Shawn M Burgess[3] and Aurnab Ghose[1*]

[1] Indian Institute of Science Education and Research (IISER) Pune, Dr Homi Bhabha
Road, Pune 411008, INDIA

[2] Zebrafish Core, Translational and Functional Genomics Branch, National Human
Genome Research Institute, National Institutes of Health, Bethesda, Maryland, USA

[3] Translational and Functional Genomics Branch, National Human Genome Research
Institute (NHGRI), National Institutes of Health (NIH), Bethesda, MD, USA

16   * Correspondence: Dr. Aurnab Ghose, Indian Institute of Science Education and
17   Research (IISER) Pune, Dr Homi Bhabha Road, Pune 411008, INDIA.

18   Email: aurnab@iiserpune.ac.in

ABSTRACT

Background

Cytoskeletal remodelling plays a pivotal role in the establishment of neuronal connectivity during development and in plasticity in adults. Mutations in the cytoskeleton regulatory protein Formin-2 (Fmn2) are associated with neurodevelopmental disorders like intellectual disability, though its function in neuronal morphogenesis has not been characterised *in vivo*.

Results

Here we develop a loss-of-function model for *fmn2b*, the zebrafish orthologue of Fmn2, using CRISPR/Cas9-mediated gene editing. *fmn2b* mutants display motor deficits starting from the earliest motor responses in the embryo. We find that *fmn2b* is expressed in motor neurons and its loss reduces motor neuron innervation of the axial muscles without affecting myotome integrity. The translocation of caudal primary (CaP) motor neuron outgrowth is compromised in *fmn2b* mutants, while rostral primary (RoP) motor neurons have missing soma or stall at the horizontal myoseptum. Strikingly, axon collateral branching of the motor neurons is severely compromised and results in reduced synaptic coverage of the myotome. Rescue experiments identify the requirement for Fmn2-mediated actin nucleation for motor neuron outgrowth and arborisation.

Conclusions

The zebrafish loss-of-function model of Fmn2 reveals the specific requirement of F-actin polymerisation by Fmn2 in neuromuscular development. It also underscores the role of Fmn2 in motor neuropathies, especially as a proportion of individuals harbouring mutations in Fmn2 present with hypotonia.

43   **Keywords**

44   Fmn2b, motor behaviour, motor neurons, axonal branching, actin nucleation

INTRODUCTION

Developing neurons migrate to their predetermined targets, undergo axonogenesis, pathfinding and eventually arborize to make functional synapses. Cytoskeletal remodelling allows dynamic interactions of the developing neurons with the environment during outgrowth, pathfinding and arborization (Dickson, 2002; Lowery and Van Vactor, 2009; Gordon-Weeks and Fournier, 2014; Flynn and Bradke, 2020) aided by several actin remodelling proteins (Dent et al., 2011; Kessels et al., 2011; Lewis et al., 2013; Coles and Bradke, 2015; Armijo-Weingart and Gallo, 2017).

Remodelling of actin filaments allows protrusive structures to probe the environment for cues in the form of growth cone filopodia and axonal and dendritic branches (Lowery and Van Vactor, 2009; Lewis et al., 2013; Armijo-Weingart and Gallo, 2017; Menon and Gupton, 2018).

In zebrafish, motor neurons exit perpendicular to the spinal cord and traverse the myotomes to arborize and make synapses onto the axial muscles (Myers et al., 1986; Westerfield et al., 1986). Adequate innervation of muscles by the motor neurons is ensured by hemisegment specific outgrowth, pathfinding and arborization. Precise neuronal morphogenesis of motor neurons is critical for achieving context-specific synaptic drive at the level of muscles, enabling the organism to execute motor behaviours.

Actin nucleating proteins are a major class of actin remodelling factors and have been implicated in neuronal morphogenesis. The actin nucleator Cobl localizes to regions with high actin dynamics and is implicated in the maintenance of neurite branching (Ahuja et al., 2007). Actin nucleators mediate the local rearrangement of actin filaments via the formation of dynamic structures like F-actin patches and F-actin blobs

69 and regulate the initiation of protrusions and branching (Ketschek and Gallo, 2010;

70 Spillane et al., 2011; Spillane and Gallo, 2014; Nithianandam and Chien, 2018; Kundu

71 et al., 2020). Reports have characterized the role of Dishevelled associated activator

72 of morphogenesis (Daam) formins in *Drosophila* axonal growth (Matusek et al., 2008;

73 Prokop et al., 2011) and Daam1a in habenular morphogenesis and IPN connectivity

74 in zebrafish (Colombo et al., 2013). Another study identified Formin-3 (Form3) in the

75 maintenance of complex dendritic arbours of nociceptive sensory neurons in

76 *Drosophila* via microtubule stabilization and adequate organelle trafficking (Das et al.,

77 2021). However, the function of formin family of actin nucleators in axonal branching

78 *in vivo* remains elusive.

79 Formin-2 (Fmn2) is part of the FMN family of formins and is known to be involved in

80 neuronal development. Biallelic mutations in Fmn2 cause neurodevelopmental defects

81 resulting in intellectual disability and muscle hypotonia in humans (Law et al., 2014).

82 Primary hippocampal neurons from Fmn2 null mice show defective dendritic spine

83 morphogenesis (Law et al., 2014) and there is increasing evidence implicating copy

84 number variations (CNVs) and single nucleotide polymorphisms (SNPs) in Fmn2 in

85 sporadic Amyotrophic Lateral Sclerosis (SALS) (Schymick et al., 2007; Pamphlett et

86 al., 2011; Morello et al., 2018). Fmn2 has previously been characterized in outgrowth,

87 pathfinding (Sahasrabudhe et al., 2016; Ghate et al., 2020; Kundu et al., 2021) and

88 axonal branching (Kundu et al., 2020) in chick spinal primary neurons *in vitro*.

89 Depletion of Fmn2 causes reduced growth cone motility and mechanotransduction in

90 chick spinal primary neurons (Sahasrabudhe et al., 2016; Ghate et al., 2020). Fmn2

91 regulates the stabilization of actin patches required for the initiation of axonal

92 branching in cultured chick spinal neurons by insulating the actin patch from actin

93 depolymerizing factors like ADF (Kundu et al., 2020). Morpholino-mediated transient

knockdown of Fmn2b, the zebrafish ortholog of Fmn2, causes impaired outgrowth of

the spiral fiber neuron commissures in the hindbrain and results in diminished short

latency escape response (Nagar et al., 2021).

In this study, we generate *fmn2b* mutants using the CRISPR-Cas9 based gene editing

to assess the contribution of Fmn2b in neural circuit development and behaviour and

to improve our understanding of the mechanistic underpinnings of the

neurodevelopmental disorders associated with Fmn2.

Motor neurons are the final, executory neurons driving motor responses. In zebrafish,

the motor neurons receive synaptic inputs from reticulospinal neurons, including the

Mauthner cell and several hindbrain interneurons, which regulate motor neuron

function in response to stimuli (Bernhardt et al., 1990; Hale et al., 2001; McLean and

Fetcho, 2008). Three subtypes of primary motor neurons are present on either side of

the spinal cord – the Caudal primary (CaP), the Middle primary (MiP) and the Rostral

primary (RoP) – with distinct morphology and muscle targets (Eisen et al., 1986; Myers

et al., 1986; Bernhardt et al., 1990; Melançon et al., 1997; Stifani, 2014). The

myotomes are also innervated by secondary motor neurons which develop after the

primary motor neurons and have less complex arborization fields (Pike et al., 1992;

Beattie et al., 1997; Menelaou and Svoboda, 2009). Motor neurons innervate the

target muscles to form structurally and functionally distinct motor units and regulate

the overall synaptic drive of the muscles during motor behaviours (Saint-Amant and

Drapeau, 1998; Drapeau et al., 2002; Bagnall and McLean, 2014; Bello‑Rojas et al.,

2019).

This study uncovers the role of the zebrafish orthologue of Formin-2, Fmn2b, in

regulating motor neuron outgrowth and branching *in vivo* and the consequent effects

on motor behaviour.

RESULTS

Generating fmn2b CRISPR mutants

Zebrafish has two paralogs of Formin-2, *fmn2a* and *fmn2b*. *fmn2b*, located on chromosome 12, has higher sequence homology with human Fmn2 compared to *fmn2a* and has the characteristic Formin Homology 1 (FH1), Formin Homology 2 (FH2) and Formin Spire Interaction (FSI) domains (Nagar et al., 2021). In contrast, the FH1 domain of *fmn2a* is truncated. As observed in humans, rodents and chicks (Leader and Leder, 2000; Sahasrabudhe et al., 2016), *fmn2b* mRNA is enriched in the zebrafish nervous system whereas *fmn2a* shows no detectable expression in the nervous system of zebrafish (Nagar et al., 2021). In this study, we used the CRISPR-Cas9 technology to generate mutants of *fmn2b*, the functional ortholog of mammalian Fmn2 in zebrafish.

To generate *fmn2b* knockout zebrafish, sgRNAs targeting the exon 1 of *fmn2b* were designed using the CRISPRscan tool (Moreno-Mateos et al., 2015) and cross verified for high scores using the ZebrafishGenomics track (LaFave et al., 2014) in UCSC genome browser. Exon 1 was targeted to ensure that the resulting mutant would lack the functional domains.

The embryos injected with sgRNA and Cas9 mRNA were raised till sexual maturity. Individual embryos from the injected clutch were outcrossed with wildtype fish of the opposite sex to obtain eggs that were genotyped using fluorescent PCR (Varshney et al., 2016). Multiple founder lines with different allelic mutations were identified for *fmn2b* by fluorescent-PCR. The F1 progeny obtained from the outcross of these founder lines was raised to adulthood and the siblings were in-crossed. The F2 progeny were raised to adulthood and genotyped using the caudal fin clip method and

subsequent Sanger sequencing of the genomic locus flanking the sgRNA-1 target

sequence to determine whether they were heterozygous or homozygous for the

inherited mutant allele.

The sgRNA caused various indels near the intended target around 696 bp into the

fmn2b cDNA sequence in exon 1, corresponding to 232nd amino acid position in the

1454 amino acid long Fmn2b protein sequence. Two alleles causing a 7 bp

(*fmn2b$^{\Delta7/\Delta7}$*) and a 4 bp (*fmn2b$^{\Delta4/\Delta4}$*) deletion, respectively, were selected and

maintained as homozygous mutants (Figure 1 A-B). The deletions result in premature

stop codons at the 280th or the 281st amino acid in the *fmn2b$^{\Delta7/\Delta7}$*and *fmn2b$^{\Delta4/\Delta4}$*

mutants, respectively (Figure 1 C). The two homozygous mutant alleles were in-

crossed to obtain heteroallelic *fmn2b* mutants (*fmn2b$^{\Delta4/\Delta7}$*) and used in subsequent

experiments.

fmn2b mutants exhibit Spontaneous Tail Coiling (STC) and Touch-Evoked

Escape Response (TEER) deficits

The first motor behaviour to appear in the developing zebrafish embryo is spontaneous

tail coiling (STC) which starts around 17 hpf and persists till 27 hpf (Saint-Amant and

Drapeau, 1998; Brustein et al., 2003). We recorded the spontaneous tail coiling in

fmn2b$^{+/+}$ and *fmn2b$^{\Delta4/\Delta7}$* embryos at 22 hpf (Movie 1). The frequency of spontaneous

tail coiling was reduced in *fmn2b$^{\Delta4/\Delta7}$* mutant embryos (3.168 ± 0.314 min^{-1}) as

compared to the wild-type *fmn2b$^{+/+}$* embryos (5.003 ± 0.203 min^{-1}) (Figure 2 A).

However, the maximum amplitude of tail coiling did not change significantly in the

fmn2b$^{\Delta4/\Delta7}$ embryos (Figure 2 B). The decrease of coiling frequency but not the

magnitude of coiling indicates that the defects are likely due to deficits in the motor

neuron or the neuromuscular junction (NMJ) function rather than loss of muscle

integrity.

As the embryo develops, the motor circuits mature to mediate touch-evoked escape

responses (TEER) beginning at 48 hpf. TEER is typically manifested as a fast C-bend

escape followed by a swimming bout. We performed the TEER assay on $fmn2b^{+/+}$ and

$fmn2b^{\Delta4/\Delta7}$ embryos at 60 hpf (Movie 2) and recorded their response at 200 fps. Both

the distance traversed (13.55 ± 1.06 mm) and the average speed (34.26 ± 2.785 mm/s)

of touch-evoked swimming was reduced in 60 hpf $fmn2b^{\Delta4/\Delta7}$ embryos as compared

to the distance covered (65.31 ± 4.93 mm) and swim speed (84.37 ± 3.253) of $fmn2b^{+/+}$

embryos (Figure 2 C, D, E). However, the maximum instantaneous speed attained

during the entire swim bout was not significantly different between $fmn2b^{+/+}$ (199.5 ±

7.039 mm/s) and $fmn2b^{\Delta4/\Delta7}$ embryos (183.2 ± 6.245 mm/s) (Figure 2 F).

The comparable maximum tail coiling amplitudes and maximum instantaneous speeds

observed in $fmn2b^{+/+}$ and $fmn2b^{\Delta4/\Delta7}$ embryos suggest abnormal motor neuron

development or NMJ function and is unlikely to involve deficits in the musculature.

Consistent with this, phalloidin staining of the $fmn2b^{\Delta4/\Delta7}$ embryos revealed no

structural deformities in the axial muscles compared to the $fmn2b^{+/+}$ embryos (Figure

S1).

fmn2b mRNA is expressed in the spinal cord and motor neurons of zebrafish

embryos

The time window of STC and TEER in zebrafish embryos coincides with the outgrowth

and pathfinding of primary motor neurons (Myers et al., 1986). *fmn2b* mRNA was

found to be expressed in the spinal cord 48 hpf onwards suggesting a possible role in

the development of spinal neurons (Figure 3 A, B). Previous reports have also reported

Fmn2 mRNA expression in the spinal cord of chick, mouse and human embryos

(Leader and Leder, 2000; Sahasrabudhe et al., 2016). To explicitly test the expression

of *fmn2b* mRNA in the motor neurons, GFP positive motor neurons were isolated using

Fluorescence-activated cell sorting (FACS) from 24 hpf and 60 hpf *Tg(mnx1:GFP)*

embryos and the expression of *fmn2b* evaluated by RT-PCR. *fmn2b* transcript was

expressed in motor neurons expressing GFP driven by the Tg(mnx1:GFP) isolated

from both 24 hpf and 60 hpf embryos (Figure 3 C). Therefore, *fmn2b* is expressed in

the spinal cord and in the motor neurons of zebrafish embryos and could mediate

motor neuron development and function.

CaP motor neuron outgrowth is slower in fmn2b mutants

To test the hypothesis that the motor defects in *fmn2b* mutants may arise due to

deficits in motor neuron morphogenesis, the $fmn2b^{\Delta4/\Delta7}$ mutant line was crossed with

the *Tg(mnx1:GFP)* line to obtain homozygous *fmn2b* mutants in the transgenic

background of GFP-labelled motor neurons. The development of motor neurons was

observed from 22 hpf to 26 hpf using time-lapse confocal imaging of the GFP-labelled

neurons (Movie 3). The growth cones of the pioneering caudal primary (CaP) motor

neurons were tracked using the Manual Tracking plugin in ImageJ, and the average

translocation speed was calculated. The $fmn2b^{\Delta4/\Delta7}$ embryos showed slow outgrowth

of the CaP neurons (Figure 3 D, E). The growth cone translocation speed of $fmn2b^{\Delta4/\Delta7}$

embryos (22.1 ± 1.575 µm/h) was slower than the $fmn2b^{+/+}$ embryos (28.34 ± 1.962

µm/h) (Figure 3 F).

Primary motor neuron outgrowth and branching defects in fmn2b mutants

In addition to the slow outgrowth of CaP motor neurons in *fmn2b* mutants, collateral

branches extended by the motor neurons at 24 hpf (Figure 4 A, B) and 60 hpf (Figure

4 G, H) were observed using the *fmn2b* mutant line $fmn2b^{\Delta4/\Delta7}$ in the background of

the transgenic line *Tg(mnx1:GFP).* The embryos were mounted laterally to image the

motor neurons and the images traced and quantified using the NeuronJ plugin in

ImageJ. At 24 hpf, 100% of the hemisegments had the CaP motor neuron soma

present in *fmn2b*$^{+/+}$ as well *fmn2b*$^{Δ4/Δ7}$ embryos. The branch density along the fascicle

length (Figure 4 E) and the length of the motor fascicle (Figure 4 F) extended by the

primary motor neurons at 24 hpf was found to be reduced in *fmn2b*$^{Δ4/Δ7}$ embryos.

Similarly, 60 hpf *fmn2b*$^{Δ4/Δ7}$ embryos showed a reduction in the density (Figure 4 K) of

collateral branches along the motor fascicle along with a reduction in the fascicle

length (Figure 4 L).

These observations implicate *fmn2b* function in both the outgrowth of the CaP motor

neuron and collateral branching of motor neurons.

F-actin nucleation activity of Fmn2 is required for motor neuron branching

The functional domains characteristic of Formin-2 are conserved across vertebrates

and ectopic expression of full-length mouse Fmn2 (mFmn2) can rescue

neurodevelopmental defects induced by morpholino-mediated knockdown of Fmn2b

in zebrafish (Nagar et al., 2021). To assess if the expression of exogenous Fmn2 could

rescue the outgrowth and branching defects, *mFmn2* mRNA tagged with mCherry was

injected in the *fmn2b*$^{Δ4/Δ7}$ mutant embryos at the 1-cell stage. The branch density and

fascicle length defects were rescued in both 24 hpf (Figure 4 C, E, F) and 60 hpf

(Figure 4 I, K, L) mutant embryos.

F-actin nucleation is the canonical function of formins, including Fmn2, and is

mediated by the FH2 domain (Goode and Eck, 2007; Quinlan et al., 2007; Yoo et al.,

2015). The FH2 domain is conserved across phyla, including zebrafish (Nagar et al.,

2021). A conserved isoleucine residue (at position 1226 in mouse Fmn2) is known to

be critical for actin nucleation (Quinlan et al., 2007; Roth-Johnson et al., 2014; Kundu

et al., 2020). Clustal omega alignment of the conserved isoleucine residue is shown

in Figure S2. To test if actin nucleation by Fmn2b is required in zebrafish motor neuron

morphogenesis, an F-actin nucleation dead version of mFmn2 with a point mutation

converting the Isoleucine at 1226 amino acid position to Alanine (mFmn2-I1226A) was

used. The mRNA for mFmn2-I1226A tagged with mCherry was injected in the

fmn2b$^{Δ4/Δ7}$ embryos at the 1-cell stage and the motor neuron development was

analyzed.

The *mFmn2-I1226A* mRNA injected 24 hpf (Figure 4 D, E, F) and 60 hpf (Figure 4 J,

K, L) mutant embryos continue to exhibit the outgrowth and branching defects similar

to the *fmn2b*$^{Δ4/Δ7}$ embryos. Rescue of phenotype exhibited by the *fmn2b*$^{Δ4/Δ7}$ mutants

by full-length mFmn2 but not the nucleation dead version, mFmn2-I1226A points

towards the conserved function of Fmn2 in motor neuron development and highlights

the significance of F-actin nucleation by Fmn2 in outgrowth and branching of motor

neurons.

Rostral Primary (RoP) motor neuron development is compromised in fmn2b

mutants

Despite slower outgrowth rates and reduced branching in *fmn2b*$^{Δ4/Δ7}$ embryos, the

CaP fascicle eventually reaches the extremity of the ventral musculature in the

zebrafish flank. On the contrary, another class of primary motor neurons, the Rostral

primary (RoP) motor neurons which typically project to the mid-dorsal and mid-ventral

musculature (Bagnall and McLean, 2014; Bello-Rojas et al., 2019), are permanently

affected in *fmn2b*$^{Δ4/Δ7}$ embryos. The RoP soma are located at their predetermined site

adjacent to the CaP and VaP cell body cluster in the spinal cord. The cell bodies of

primary motor neurons are distinctly recognizable at 24 hpf.

On examining the 24 hpf embryos, the RoP cell body was found to be missing in *fmn2b*

mutants. Compared to 2.5% of the hemisegments in *fmn2b$^{+/+}$* embryos, 46.87% of the

hemisegments in *fmn2b$^{\Delta4/\Delta7}$* embryos did not have the RoP cell body. Injection of 1-

cell stage *fmn2b$^{\Delta4/\Delta7}$* embryos with *mFmn2* mRNA rescued this defect with only 2.85%

hemisegments lacking RoP soma. However, the actin nucleation dead *mFmn2-I1226A*

mRNA failed to rescue the loss of RoP soma in *fmn2b* mutants and 43.75% of the

hemisegments continued to show loss of RoP soma (Figure 5 A-F).

At 60 hpf, the RoP motor neurons innervate their target in the mid-dorsal and mid-

ventral musculature. The RoP motor neurons in *fmn2b$^{\Delta4/\Delta7}$* mutants showed two major

types of defects, absence of RoP outgrowth and stalling of RoP at the choice point

near the horizontal myoseptum (HMS), where the RoP axons take a sharp lateral turn.

In the *fmn2b* mutant embryos, 91.3% of the hemisegments analyzed showed little to

no lateral innervation. Of these, 44.35% of the hemisegments displayed defasciculated

axons near the horizontal myoseptum, which acts as a choice point for the RoP axons

to turn laterally. The hemisegments with defasciculated axons were classified as

stalled RoP axons. Similar defects in RoP-like secondary motor neurons have been

reported to occur due to depletion of Kif1b and Fidgetin like-1 in zebrafish (Fassier et

al., 2018; Atkins et al., 2019). However, 46.95% of hemisegments had no RoP

innervation altogether and were classified as RoP absent. Only 8.7% hemisegments

showed the stereotypical RoP innervation and were classified as RoP present. In the

fmn2b$^{+/+}$ embryos, only 3.15% of the hemisegments quantified showed RoP stalling

defects, and the rest had stereotypical innervation by the RoP neurons. The RoP

stalling and outgrowth defects were rescued by injection of full-length *mFmn2* mRNA

with only 10.16% hemisegments with RoP stalling and 8.47% hemisegments with RoP

outgrowth defects. On the other hand, RoP defects persisted in the *fmn2b$^{\Delta4/\Delta7}$*

embryos injected with *mFmn2-I1226A* mRNA, with 18.3% of hemisegments having

stalled RoP axons and 81.7% without RoP innervation (Figure 5 G-L).

Collectively, these results indicate that *fmn2b* has pleiotropic effects on RoP

development dependent on F-actin nucleation function. The deficits are at the level of

loss of RoP soma, deficits in RoP axonogenesis and stalling at the horizontal

myoseptum choice point.

Effect of overexpression of Fmn2 in motor neurons

The role of *fmn2b* in motor neuron development has been established so far by

observing *fmn2b* mutants, where the loss of Fmn2b in motor neurons causes a

reduction in outgrowth and branching. Overexpression of mouse Fmn2 was employed

in wildtype embryos to test the phenotypes in the context of Fmn2 gain of function.

The embryos overexpressing mFmn2 showed increased branching, but

overexpression of the nucleation dead mFmn2-I1226A did not cause any significant

changes and was comparable to the wild-type embryos (Figure 6 A, B, E, F). The

branch density was found to be increased in mFmn2 mRNA injected embryos at 24

hpf as well as 60 hpf but not in mFmn2-I1226A mRNA injected group (Figure 6 C, G).

Interestingly, in 24 hpf embryos, the fascicle length was not significantly different in

embryos overexpressing mFmn2 but was found to be decreased in embryos

overexpressing mFmn2-I1226A. In 60 hpf embryos, the fascicle length was slightly

increased in the mFmn2 overexpression group but remain unaffected in the mFmn2-

I1226A overexpression group (Figure 6 D, H).

However, quantification of RoP soma and RoP axon outgrowth phenotypes in the

fmn2b$^{+/+}$ embryos overexpressing mFmn2 and mFmn2-I1226A did not reveal any

significant changes (Figure S3).

Neuromuscular Junction (NMJ) development in fmn2b mutants

Primary motor neurons innervate the fast-twitch muscles and form functional synapses

by 48 hpf. To visualize the neuromuscular junctions (NMJ), double immunostaining of

whole-mount 60 hpf embryos with znp-1 antibody (presynaptic marker) and α-

bungarotoxin (post-synaptic marker) were undertaken. The co-localization of the pre-

and post-synaptic markers revealed the engaged NMJ synapses in $fmn2b^{+/+}$ and

$fmn2b^{\Delta4/\Delta7}$ embryos (Figure 7 A, B). Synapses (co-localization of znp-1 and α-

bungarotoxin signals) along the neuronal arbours were quantified using the

SynapCountJ plugin in Fiji. The number of synapses identified was normalized to the

area of the myotome corresponding to one somite and the length of the neuronal

arbour.

The $fmn2b^{\Delta4/\Delta7}$ mutants show reduced synaptic coverage of the myotomes compared

to $fmn2b^{+/+}$ embryos (Figure 7 C). This result is consistent with previous observations

of reduced outgrowth and branching density, and therefore the attenuated occupancy

of myotome area, of motor neurons.

However, the number of synapses normalized to the arbour length was found to be

comparable (Figure 7 D). Thus, while the ability of motor neurons to form synapses is

intact in fmn2b mutants, the dramatically reduced arborization results in an effective

reduction in the total number of synapses per myotome. The locomotor defects

observed in fmn2b mutants is consistent with the decrease in the total number of

synapses per myotome leading to insufficient activation of the myotome.

DISCUSSION

Mutations, including loss-of-function mutations, in Fmn2 have been associated with intellectual disability, sensory dysfunction, age-associated cognitive decline (Perrone et al., 2012; Law et al., 2014; Agís‑Balboa et al., 2017; Marco et al., 2018; Gorukmez et al., 2020). Interestingly, some affected individuals also develop hypotonia (Law et al). Further, mutations in Fmn2 have also been associated with sporadic amyotrophic lateral sclerosis in multiple studies (Schymick et al., 2007; Pamphlett et al., 2011; Morello et al., 2018). These associations suggest a possible involvement of Fmn2 in motor neuron development and function.

In this study, we generated *fmn2b* mutants using the CRISPR-Cas9 system (Figure 1) and found locomotor deficits in the homozygous mutants. *The fmn2b* mutants displayed deficits right from the earliest motor behaviours (STC at 22 hpf; Figure 2) to later stages where the development of the embryonic neural circuits is complete (TEER at 60 hpf; Figure 2). *fmn2b* mRNA expression was also detected in motor neurons and the spinal cord (Figure 3). Systematic evaluation of the primary motor neurons revealed defects in the outgrowth of the CaP motor neurons. Live imaging indicated that the growth cone translocation velocity was reduced in $fmn2b^{\Delta4/\Delta7}$ embryos (Figure 3).

Recent *in vitro* studies in chick spinal neurons implicates Fmn2 in growth cone translocation. These studies implicate Fmn2 in mediating a molecular clutch that stabilizes the contact sites between the growth cone and the extracellular matrix and regulates the generation of traction forces required for outgrowth (Sahasrabudhe et al., 2016; Ghate et al., 2020). Additionally, Fmn2 facilitates the stabilization of protrusive processes, like growth cone filopodia, by coupling exploratory microtubules

with the F-actin cytoskeleton (Kundu et al., 2021). Consistent with this, the depletion

of Fmn2 reveals altered microtubule dynamics in the growth cones of zebrafish Rohon-

Beard neurons *in vivo*. Similar mechanisms involving actin and microtubule

remodelling could be involved in the translocation of CaP growth cones in zebrafish.

Even more striking than the delayed outgrowth rates of CaP motor neurons, the

development of collateral branches which provide synaptic coverage of the entire

muscle field was severely affected (Figure 4). The staggered development of motor

neurons in two phases allowed us to selectively look at the innervation of muscles by

primary motor neurons at 24 hpf and collectively look at primary and secondary motor

neuron innervation at 60 hpf. In addition, the 60 hpf embryos have elaborate branching

and fully functional NMJ synapses allowing characterization of *fmn2b* in outgrowth,

branching and synapse formation in motor neurons.

In 24 hpf *fmn2b* mutant embryos, branch density along the fascicle extended by the

primary motor neurons and the fascicle length itself was reduced, suggesting a

decrease in the innervation of the target fast muscle fibers (Figure 4). Similarly, the 60

hpf *fmn2b*$^{\Delta4/\Delta7}$ embryos have reduced branch density and fascicle outgrowth

compared to wild-type embryos and are likely to result in reduced speed and distance

covered by *fmn2b* mutants (Figure 4). Deficits in motor neuron outgrowth and collateral

branching at both 24 hpf and 60 hpf was rescued by overexpressing mouse full length

Fmn2 in the *fmn2b* homozygous mutants (Figure 4).

Axon collateral branching is a finely regulated multi-step process initiated by the rapid

assembly of F-actin resulting in a filopodia-like protrusion from the axonal shaft

(Ketschek and Gallo, 2010; Gallo, 2011, 2016; Coles and Bradke, 2015; Ketschek et

al., 2015; Armijo-Weingart and Gallo, 2017; Menon and Gupton, 2018). Further

cytoskeleton remodelling involving both the actin and microtubule cytoskeletons

stabilize the protrusion and extend it to a collateral branch (Gallo, 2011, 2016; Armijo-

Weingart and Gallo, 2017). The formation of the lateral protrusion from the axonal

shaft is a major rate-limiting step in collateral branching and is initiated by the

development of a juxtamembrane F-actin patch. *In vitro* studies implicate several actin

binding proteins in regulating axonal F-actin patches. These include the actin

nucleating Arp2/3 complex, Drebrin, WVE-1/WAVE regulatory complex (WRC) and

cortactin (Spillane et al., 2011; Hu et al., 2012; Chia et al., 2014; Spillane and Gallo,

2014; Ketschek et al., 2016; Balasanyan et al., 2017). Recently, Fmn2 has been

implicated in collateral branch formation in chick spinal neurons *in vitro*. Fmn2 appears

to regulate the lifetime of axonal F-actin patches and consequently the probability of

branch initiation (Kundu et al., 2020).

The F-actin nucleation and elongation activity of Fmn2 has been characterized *in vitro*

(Montaville et al., 2014, 2016). Abrogation of F-actin nucleation activity of Fmn2 can

be achieved by mutating a conserved Isoleucine residue in the FH2 domain to Alanine

(Quinlan et al., 2007; Roth-Johnson et al., 2014; Kundu et al., 2020). We employed

the full-length mouse Fmn2 bearing the isoleucine to alanine mutation (mFmn2-

I1226A) in rescue experiments to test if the F-actin nucleating/elongating activity is

necessary for motor neuron development.

Expression of mFmn2-I1226A in the *fmn2b* mutants could not rescue the outgrowth

and branching defects of the motor neurons in 24 hpf and 60 hpf embryos (Figure 4).

The failure of mFmn2-I1226A in rescuing the defects underscores the significance of

the actin nucleating activity of Fmn2b in motor neuron outgrowth and branching

consistent with reports from primary neuronal cultures of chick spinal cord reported

previously (Kundu et al., 2020). Therefore, the motor neuron development in zebrafish

is dependent on the F-actin nucleation activity of Fmn2b.

Loss of fmn2b had pleiotropic effects on the development of the RoP motor neuron.

The RoP soma and RoP outgrowth were severely affected in *fmn2b* mutants. At 24

hpf, multiple hemisegments in *fmn2b* mutants did not have RoP soma at its

characteristic position in the spinal cord (Figure 5). The role of Fmn2 in the regulation

of differentiation or specification of progenitors is not formally tested despite several

studies indirectly indicating pathways involving Fmn2 in cell differentiation. In a recent

report, Fmn2 has been shown to cause neural progenitor differentiation defects Fmn2

and Flna double knockout mice in a synergistic manner (Lian et al., 2016). The

absence of RoP soma in zebrafish *fmn2b* mutants may indicate a possible role of

fmn2b in neural progenitor specification and/or differentiation. The RoP soma was

seen in the expected location in *fmn2b$^{\Delta 4/\Delta 7}$* embryos injected with mFmn2 mRNA but

the defect could not be rescued by the injection of mFmn2-I1226A mRNA.

Interestingly, the F-actin nucleating activity of Fmn2 is required for the differentiation

of motor neuron progenitors or their specification. These results open up new

possibilities to uncover the mechanistic role of Fmn2 in neural development.

In wild-type zebrafish embryos, the lateral projections from RoP motor neurons and

follower secondary motor neurons begin outgrowth later than CaP and MiP neurons

(Kuwada, 1993; Liu et al., 2016). In 60 hpf *fmn2b* mutants, the side branches of RoP

motor neurons innervating the horizontal myoseptum were either not detected (46%

embryos) or appeared to be stalled at the choice point (44% embryos), i.e., the

horizontal myoseptum (Figure 5). RoP-like secondary motor neurons, which follow the

same trajectory as RoP primary motor neurons have previously been shown to have

pathfinding and stalling defects at the horizontal myoseptum in Fidgetin like-1 and

Kif1b mutants (Fassier et al., 2018; Atkins et al., 2019). Similarly, the stalled RoP

axons appear defasciculated in 60 hpf *fmn2b* mutants

One of the factors contributing to RoP outgrowth defects leading to the absence of

RoP innervation in *fmn2b* mutants could be the lack of RoP soma as seen in 24 hpf

mutant embryos. Further, the RoP stalling defect implicates Fmn2b in axonal

pathfinding in response to guidance cues at the choice point. RoP outgrowth and

stalling together caused a noticeable reduction in the innervation of the mid-dorsal and

mid-ventral region of the axial myotome. The RoP may be more severely affected than

the CaP neurons due to their late axonogenesis concomitant with the pleiotropic

function and late expression of *fmn2b* in the spinal cord at 48 hpf.

Intriguingly, the overexpression of mouse Fmn2 in *fmn2b*$^{+/+}$ (wild-type) embryos

causes opposite effects as compared to *fmn2b* knockout manifested as

hyperbranching. This implies a significant role for *fmn2b* in regulating the collateral

branching of motor neurons. Further, overexpression of the nucleation dead version

of mFmn2 does not result in increased branch density and underscores the

involvement of actin nucleation activity of Fmn2 (Figure 6).

Analysis of NMJ synapses using znp-1 and α-bungarotoxin double staining in *fmn2b*

mutants showed no changes in the number of synapses along the total length of the

motor neuron branches but showed a reduction in the total number of synapses when

normalized to the area of the target myotome. We suggest that the Fmn2b has a

primary role in regulating motor neuron branching and not in the formation of NMJ

synapses. The behavioural defects in *fmn2b* mutants are likely due to the muscles not

receiving sufficient input due to inadequate branching (Figure 7).

In a recent study, prolonged exposure of zebrafish larvae to strong and variable water

currents caused upregulation of Fmn2b (Langebeck-Jensen et al., 2019). The rapid

upregulation of Fmn2b in response to environmental stressors involving swimming and

force generation in larvae with pre-established motor neural circuits invoke the

possible involvement of *fmn2b* in neuronal plasticity and requires systematic

investigation.

Taken together, the zebrafish model of Fmn2 loss-of-function offers unexpected

insight into motor neuron development and innervation of axial muscles. In addition to

identifying the requirement of actin polymerization by Fmn2 in motor neuron

morphogenesis and motor outputs, it highlights the central role of cytoskeleton

remodelling in motor neuron homeostasis and the development of neuropathies.

CONCLUSIONS

Motor behaviours like spontaneous tail coiling and touch evoked escape response

were adversely affected in the *fmn2b* mutants. Visualizing developing motor neurons

using the *Tg(mnx1:GFP)* in *fmn2b* mutants revealed outgrowth and branching defects

but no changes to myotome integrity. Further, we find that the F-actin nucleating

activity of Fmn2 is required for motor neuron outgrowth and collateral branching. The

reduced coverage of the axial myotomes by the motor neurons results in compromised

synaptic inputs and contributes to the motor deficits observed in the *fmn2b* mutants.

METHODS

Zebrafish maintenance and procedures

All protocols used in this study were approved by the Institutional Animal Ethics Committee and the Institutional Biosafety Committee of IISER Pune or a National Human Genome Research Institute (NHGRI/NIH) Animal Care and Use Committee approved animal study protocol. The TAB5 wildtype strain of zebrafish was used for all the experiments including generation of CRISPR mutants. The TAB5 strain was used for all the wildtype outcrosses. Breeding pairs of adult zebrafish were maintained in recirculating aquaria (Techniplast) under a 14h-10h light-dark cycle. The temperature was maintained at 28.5˚C and the pH was buffered between 7.2 to 7.8. The breeding adults were crossed to obtain embryos which were collected and grown in E3 buffer and used at different stages of development as indicated (Kimmel et al., 1995). For immunostaining and live imaging experiments, the buffer was supplemented with 0.003% Phenylthiourea (PTU; Sigma) to remove pigmentation from the skin. The transgenic line *Tg(mnx1:GFP)* was used to visualize motor neurons wherein the *mnx1* promoter specific to motor neurons drives GFP expression (Flanagan-Steet et al., 2005).

Whole mount in situ hybridization

The RNA probes and procedure used for whole mount *in situ* hybridization experiments have been described previously (Nagar et al., 2021).

Isolation of motor neurons from transgenic embryos, FACS, RT-PCR

Single cell suspension was made from around 200 *Tg(mnx1:GFP)* embryos as previously described (Bresciani et al., 2018). Briefly, the embryos were dissociated by trypsinization and filtering through a 70 μm sieve to obtain single cell suspension in

1X DMEM containing 10% FBS. The cells were sorted using a BD Biosciences

fluorescence-activated cell sorting (FACS) equipment selecting cells expressing GFP,

i.e., the motor neurons. RNA was extracted using Qiagen RNeasy Kit and cDNA was

prepared using the SuperScript IV RT Kit (ThermoFisher). The cDNA was used for

amplifying *fmn2b* transcripts in the mnx1 positive motor neurons. Primers and PCR

protocol used to test the presence of *fmn2b* transcripts in motor neurons tagged by

Tg(mnx1:GFP) were the same as the ones used for amplification of ISH probes from

cDNA.

RNA injections

Capped mRNA was synthesized using the HiScribe™ T7 ARCA mRNA Kit (with

tailing) from linearized DNA template containing T7 promoter sequence upstream of

the mFmn2-mCherry and mFmn2-I1226A-mCherry constructs. The transcribed mRNA

was purified using RNeasy MinElute Cleanup Kit (Qiagen). 150 pg of mFmn2-mCherry

and mFmn2-I1226A-mCherry mRNA was injected in the zebrafish embryos of desired

genotype at 1-cell stage.

sgRNA and Cas9 injections for CRISPR mutants

sgRNA targeting exon 1 of *fmn2b* was designed as previously described (Varshney et

al., 2016). T7 HiScribe kit (NEB) was used to transcribe the sgRNA DNA template with

the appended T7 promoter. The sgRNA was purified using ZymoResearch clean up

columns. T3 mMessage mMachine RNA synthesis kit (Ambion) was used to

synthesize capped Cas9 mRNA from the pT3TS-nCas9n plasmid (kind gift from Dr

Wenbiao Chen; Addgene plasmid # 46757). The synthesized Cas9 mRNA was

purified using the RNeasy MinElute Cleanup Kit (Qiagen). 30 pg sgRNA and 300 pg

Cas9 mRNA was injected in zebrafish embryos at 1 cell stage. The sequence of

sgRNA is given below.

Fmn2b_sgRNA : GGGCGAGAGGCCTCGGCTGG

(ENSDARG00000061778.6; 12:47451436-47451459, plus strand)

524 Genotyping CRISPR mutants

For identifying founder lines for *fmn2b* mutants, fluorescent PCR was performed and

analyzed by capillary gel electrophoresis using the ABI GeneAnalyzer 3730XL, as

previously described (Carrington et al., 2015; Varshney et al., 2015, 2016).

Identification of homozygous mutant lines was also done using Sanger sequencing.

Two zebrafish mutant lines with alleles causing a premature stop codon due to a

frameshift mutation were established. The homozygous mutants were crossed to each

other to obtain a heteroallelic mutant line for *fmn2b* to reduce the effect of any

background mutations in the two mutant lines due to unintended off-target effects.

Primer sequences for sanger sequencing of genomic DNA amplicons from crispants

are as follows:

Fmn2b_PCR_F	AAGCGTAAGAACCAGAATAAGC
Fmn2b_PCR_R	TCATCCGAATGGCTTGC

536 Whole mount immunostaining

537 PTU treated embryos were collected at desired stages and fixed in 4% formaldehyde

538 overnight at 4˚C. For staining actin structures, fixed embryos were washed with 0.5%

539 PBS-Triton, permeabilized with 2% PBS-Triton for 2 hours at room temperature and

540 then incubated in Phalloidin Alexa Fluor 568 diluted 1:50 in 2% PBS Triton overnight

541 at 4 °C in dark.

Neuromuscular junction labelling and quantification

For neuromuscular junction staining of embryos, znp-1 antibody (DSHB; 1:100) and

Tetramethyl Rhodamine labelled α-bungarotoxin (Invitrogen; 1:200) were used as pre-

synaptic and post-synaptic markers respectively. Whole mount immunostaining

procedures described in previous section were followed as described. The synapses

were counted using the SynapCountJ plugin after making the traces of arbours in

NeuronJ plugin in Fiji.

Fluorescence microscopy, live imaging and mounting procedure

All the microscopic imaging was performed on the inverted LSM 780 confocal

microscope (Zeiss) with a 25x oil immersion objective (NA 1.4). For imaging of the

fixed samples, the embryos were cleared in 50% glycerol and mounted dorsal side

down on a glass bottom petri dish using low gelling agarose (Sigma). For live imaging,

the live embryos were mounted laterally in 0.5% low melt point agarose (Sigma)

containing 0.003% MS-222 (Sigma) in a coverslip bottom 35 mm petri plate.

The growth cone of motor neurons was visualized using the *Tg(mnx1:GFP)* in wildtype

or mutant background. The embryos were imaged starting at 22 hpf for 4-6 hours every

3 minutes. The growth cone translocation was analyzed using the Manual tracking

plugin in Fiji to track the movement of the leading edge across time. The coordinates

obtained were analyzed using the ibidi Chemotaxis tool to calculate the average

growth cone translocation speed.

Behaviour experiment set up and behaviour analysis

Spontaneous tail coiling (STC) assay

Embryos 22 hpf within their chorions were transferred to a 35 mm petri plate containing

E3 buffer at of 28.5˚C. A video camera (AVT Pike, F-032B) was used to record the

spontaneous tail coiling behaviour of the embryos for a time period of 3 minutes at 15

fps. The videos were processed and analysed using a MATLAB script ZebraSTM

published previously (González-Fraga et al., 2019).

Touch evoked escape response (TEER) assay

60 hpf zebrafish embryos were housed in a 35 mm petri plate containing pre-warmed

E3 buffer at 28.5 ˚C. A tuberculin needle was repurposed by attaching a soft nylon

fiber in front of the syringe holding the needle, to deliver tactile stimuli to zebrafish

embryos. A soft touch was delivered to the head of the zebrafish once and their

behaviour was recorded using a high-speed video camera (AVT Pike F-032B) at 208

fps. The videos obtained were analyzed using the Manual tracking plugin in Fiji to mark

the trajectories of the zebrafish embryos upon receiving the tactile stimulus. The tracks

were further analyzed using the ibidi Chemotaxis tool to calculate the distance

travelled and average speed. The maximum instantaneous speed was calculated

manually from the coordinates obtained from the manual tracking output.

Figures and Statistical analysis

The analysis was performed for all the experiments in a genotype blinded manner.

Image analysis was performed using Fiji and the Figure panels were assembled using

Inkscape. The Violin plots were plotted in GraphPad Prism 8 and indicated statistical

tests were performed using GraphPad Prism 8. The data is represented as mean ±

SEM in the text. The statistical test used and the p values are indicated in the Figure

legends.

AVAILABILITY OF DATA AND MATERIALS

The data generated and analyzed in this study are included in this article and the additional information files. More information can be made available upon reasonable request to the corresponding author.

Figure 1. Generation of *fmn2b* CRISPR mutants. A) Schematic of the *fmn2b* protein indicating the functional protein domains and the site of *fmn2b* sgRNA-1. The FH1 domain spans the 841-1015 amino acids, FH2 domain spans the 1016-1430 amino acids and the FSI domain spans the 1431-1454 amino acids. **B)** Genomic locus from *fmn2b* exon 1 starting from 663 bp to 720 bp (corresponding to 221-240 amino acids). The sgRNA sequence is highlighted in yellow followed by the PAM sequence in pink. Representative chromatograms of amplicons sequenced from homozygous CRISPR mutants for *fmn2b* show two alleles with a 7 bp and a 4 bp deletion, respectively. The two alleles are denoted as *fmn2b*$^{\Delta 7}$ and *fmn2b*$^{\Delta 4}$. **C)** Schematic outlining the generation of *fmn2b* homozygous mutants. Fish homozygous for two alleles are denoted as *fmn2b*$^{\Delta 7/\Delta 7}$ and *fmn2b*$^{\Delta 4/\Delta 4}$, the heteroallelic combination is denoted as *fmn2b*$^{\Delta 4/\Delta 7}$.

Figure 2. *fmn2b* mutants exhibit motor defects. A) Spontaneous tail coiling (STC)

frequency and **B)** maximum amplitude of tail coiling in 22 hpf *fmn2b*[+/+] (n=121) and

fmn2b$^{Δ4/Δ7}$ (n=68) embryos. **C)** Trajectories of 2 dpf *fmn2b*$^{+/+}$ (grey traces; n=22) and

fmn2b$^{Δ4/Δ7}$ (pink traces; n=24) embryos in Touch Evoked escape response (TEER),

re-centered to a common origin for representation. The x and y axes in the plot

correspond to the pixel values corresponding to coordinates of the fish from

consecutively acquired frames. Quantification of **D)** distance covered per swim bout,

E) average speed during the swim bout, and **F)** maximum instantaneous swimming

speed of 2 dpf *fmn2b*$^{+/+}$ (n=22) and *fmn2b*$^{Δ4/Δ7}$ (n=24) embryos in response to a tactile

stimulus. (**** p-value <0.0001; ns- not significant; Mann-Whitney U test)

Figure 3. *fmn2b* is expressed in the spinal cord and motor neurons of zebrafish

embryos. *fmn2b* is required for motor neuron growth cone translocation

A) Lateral and **B)** dorsal views of a representative 60 hpf embryo showing *fmn2b* mRNA expression in the spinal cord. **C)** Gel showing bands for amplified β2-actin (actb2) control and *fmn2b* from cDNA obtained from FACS sorted motor neurons (mnx1:GFP positive cells) isolated from 24 hpf and 60 hpf wild-type *Tg(mnx1:GFP)* embryos. **D)** Representative montages of live imaging of growth cone translocation in *fmn2b*$^{+/+}$ and **E)** *fmn2b*$^{\Delta4/\Delta7}$ embryos in the background of *Tg(mnx1:GFP)* during the time period of 22 hpf to 26 hpf. **F)** Quantification of motor neuron growth cone translocation speed in *fmn2b*$^{+/+}$ (n=15 growth cones; N=3 embryos) and *fmn2b*$^{\Delta4/\Delta7}$ (n=16 growth cones; N=4 embryos). (* p-value = 0.0298; Mann-Whitney U test). Scale bar is equivalent to 20 µm.

Figure 4. Motor neuron outgrowth and branching defects in *fmn2b* mutants

Representative micrographs of motor neurons labelled by *Tg(mnx1:GFP)* in 24 hpf **A)** *fmn2b*$^{+/+}$ and **B)** *fmn2b*$^{\Delta4/\Delta7}$ and 60 hpf **G)** *fmn2b*$^{+/+}$ and **H)** *fmn2b*$^{\Delta4/\Delta7}$ embryos. Motor neurons visualized by *Tg(mnx1:GFP)* in *fmn2b*$^{\Delta4/\Delta7}$ mutant embryos injected at 1-cell stage with full-length mouse Fmn2 (mFmn2) mRNA at **C)** 24 hpf and **I)** 60 hpf and nucleation dead mouse Fmn2 (mFmn2-I1226A) mRNA at **D)** 24 hpf and **J)** 60 hpf. **E)** Quantification of branch density (number of branches per fascicle normalized to the fascicle length) per myotome hemisegment in 24 hpf *fmn2b*$^{+/+}$ (0.1571 ± 0.005 μm^{-1}; n=70 hemisegments) and *fmn2b*$^{\Delta4/\Delta7}$ embryos (0.0981 ± 0.005 μm^{-1}; n=118 hemisegments). The defects were rescued in *fmn2b*$^{\Delta4/\Delta7}$ embryos injected with

mFmn2 mRNA (0.2139 ± 0.012 μm^{-1}; n=51 hemisegments) but not in the embryos injected with mFmn2-I1226A mRNA (0.111 ± 0.007 μm^{-1}; n=63 hemisegments). **F)** Quantification of fascicle length extended by motor neurons per myotome hemisegment in 24 hpf $fmn2b^{+/+}$ (68.97 ± 1.55 μm; n=70 hemisegments) and $fmn2b^{\Delta4/\Delta7}$ embryos (44.98 ± 1.756 μm; n=118 hemisegments). The defects were rescued in $fmn2b^{\Delta4/\Delta7}$ embryos injected with mFmn2 mRNA (73.7 ± 1.86 μm; n=51 hemisegments) but not in the embryos injected with mFmn2-I1226A mRNA (73.7 ± 1.86 μm; n=63 hemisegments). **K)** Quantification of branch density (number of branches per fascicle normalized to the fascicle length) per myotome hemisegment in 60 hpf $fmn2b^{+/+}$ (0.3746 ± 0.014 μm^{-1}; n=37 hemisegments) and $fmn2b^{\Delta4/\Delta7}$ embryos (0.2211 ± 0.015 μm^{-1}; n=57 hemisegments). The defects were rescued in $fmn2b^{\Delta4/\Delta7}$ embryos injected with mFmn2 mRNA (0.4955 ± 0.021 μm^{-1}; n=30 hemisegments) but not in the embryos injected with mFmn2-I1226A mRNA (0.2369 ± 0.012 μm^{-1}; n=48 hemisegments). **L)** Quantification of fascicle length extended by motor neurons per myotome hemisegment in 60 hpf $fmn2b^{+/+}$ (139.6 ± 3.21 μm; n=37 hemisegments) and $fmn2b^{\Delta4/\Delta7}$ embryos (120.4 ± 1.94 μm; n=57 hemisegments). The defects were rescued in $fmn2b^{\Delta4/\Delta7}$ embryos injected with mFmn2 mRNA (162.7 ± 3.38 μm; n=30 hemisegments) but not in the embryos injected with mFmn2-I1226A mRNA (124.7 ± 2.81 μm; n=48 hemisegments). (**** p-value <0.0001; ns - not significant; Kruskal Wallis test followed by Dunn's post-hoc analysis). Scale bar is equivalent to 20 μm.

Figure 5. RoP soma and lateral innervation of the myotome by RoP are affected in *fmn2b* mutants

Representative micrographs of motor neurons labelled by *Tg(mnx1:GFP)* in 24 hpf **A)** *fmn2b*^{+/+} and **B)** *fmn2b*^{Δ4/Δ7} embryos. Representative micrographs of motor neurons labelled by *Tg(mnx1:GFP)* in 24 hpf old *fmn2b*^{Δ4/Δ7} mutant embryos injected at 1-cell

stage with **C)** full-length mouse Fmn2 (mFmn2) mRNA and **D)** nucleation dead

version, mFmn2-I1226A mRNA. The pink arrowheads indicate the RoP soma or its

expected position. **E)** Bar graph summarizing the percentage of embryos with defects

in RoP soma in 24 hpf *fmn2b*$^{+/+}$ (n=18) and *fmn2b*$^{\Delta4/\Delta7}$ embryos (n=23). The defects

were rescued in *fmn2b*$^{\Delta4/\Delta7}$ embryos injected with mFmn2 mRNA (n=12) but not in the

embryos injected with mFmn2-I1226A mRNA (n=10*).* χ2 test of independence shows

significant difference between the four groups, χ2 (df = 3) =101.8 , p < 0.0001. **F)** Violin

plots depicting the variation in data summarized in the bar graphs for 24 hpf embryos.

(**** p-value <0.0001; ns - not significant; Kruskal Wallis test followed by Dunn's post-

hoc analysis). Scale bar is equivalent to 20 μm.

Representative micrographs of motor neurons labelled by *Tg(mnx1:GFP)* in 60 hpf **G)**

fmn2b$^{+/+}$ and **H)** *fmn2b*$^{\Delta4/\Delta7}$ embryos. Representative micrographs of motor neurons

labelled by *Tg(mnx1:GFP)* in 60 hpf old *fmn2b*$^{\Delta4/\Delta7}$ mutant embryos injected at 1-cell

stage with **I)** full-length mouse Fmn2 (mFmn2) mRNA and **J)** nucleation dead version,

mFmn2-I1226A mRNA. The orange arrowheads point towards the stalled RoP and the

blue arrowheads indicate no RoP outgrowth. **K)** Bar graphs summarizing the

percentage of embryos with defects in RoP axon outgrowth in 60 hpf *fmn2b*$^{+/+}$ (n=22)

and *fmn2b*$^{\Delta4/\Delta7}$ embryos (n=26). The defects were rescued in *fmn2b*$^{\Delta4/\Delta7}$ embryos

injected with mFmn2 mRNA (n=14) but not in the embryos injected with mFmn2-

I1226A mRNA (n=13). χ2 test of independence shows significant difference between

the four groups, χ2 (df = 6) = 332.7, p < 0.0001. **L)** Violin plots depicting the variation

in data summarized in the bar graphs for 60 hpf embryos. (**** p-value <0.0001; ns -

not significant; Kruskal Wallis test followed by Dunn's post-hoc analysis) Scale bar is

equivalent to 20 μm.

Figure 6. Outgrowth and branching phenotypes upon overexpression of mouse

Fmn2 in wild-type embryos

Representative micrographs of motor neurons labelled by *Tg(mnx1:GFP)* in 24 hpf old

fmn2b$^{+/+}$ embryos injected with **A)** mFmn2 mRNA and **B)** mFmn2-I1226A mRNA. **C)**

Quantification of branch density (number of branches per fascicle normalized to the

fascicle length) along the fascicle extended by motor neurons per myotome

hemisegment in *fmn2b*$^{+/+}$ embryos injected with mFmn2 mRNA (0.2728 ± 0.0217 $^{\mu m-1}$;

n=40 hemisegments) and mFmn2-I1226A mRNA (0.1617 ± 0.007 $^{\mu m-1}$; n=40

hemisegments). **D)** Quantification of fascicle length extended by motor neurons per

myotome hemisegment in *fmn2b*$^{+/+}$ embryos injected with mFmn2 mRNA (73.7 ± 1.86

μm; n=40 hemisegments) and mFmn2-I1226A mRNA (68.3 ± 1.603 μm; n=40

hemisegments). Representative micrographs of motor neurons labelled by

Tg(mnx1:GFP) in 60 hpf old *fmn2b*$^{+/+}$ embryos injected with **E)** mFmn2 mRNA and **F)**

mFmn2-I1226A mRNA. **G)** Quantification of branch density (number of branches per

fascicle normalized to the fascicle length) along the fascicle extended by motor

neurons per myotome hemisegment in 60 hpf *fmn2b*$^{+/+}$ embryos injected with mFmn2

mRNA (0.599 ± 0.027 $^{\mu m-1}$; n=28 hemisegments) and mFmn2-I1226A mRNA (0.3481

± 0.02 $^{\mu m-1}$; n=28 hemisegments). **H)** Quantification of fascicle length extended by

motor neurons per myotome hemisegment in 60 hpf *fmn2b*$^{+/+}$ embryos injected with

mFmn2 mRNA (150.4 ± 1.536 μm; n=28 hemisegments) and mFmn2-I1226A mRNA

(145 ± 2.18 μm; n=28 hemisegments). (**** p-value <0.0001; * p-value = 0.0175; ns -

not significant; Kruskal Wallis test followed by Dunn's post-hoc analysis) Scale bar is

equivalent to 20 μm.

Figure 7. Synapse coverage in *fmn2b* mutants

Representative micrographs of 60 hpf **A)** *fmn2b*$^{+/+}$ (n=6) **B)** *fmn2b*$^{\Delta4/\Delta7}$ embryos (n=7)

co-stained with znp-1 antibody and α-bungarotoxin. Quantification of colocalization of

znp-1 and α-bungarotoxin in NMJ structures of 60 hpf embryos. **C)** Number of

synapses per unit myotome area in *fmn2b*$^{+/+}$ (0.0272 ±0.002; n=6) and *fmn2b*$^{\Delta4/\Delta7}$

embryos (0.01521 ±0.0005; n=7, **p-value=0.0012). **D)** Number of synapses per unit

axon length in *fmn2b*$^{+/+}$ (0.1903± 0.009; n=6) and *fmn2b*$^{\Delta4/\Delta7}$ embryos (0.2001± 0.005;

n=7). (ns- not significant; Mann Whitney U test) Scale bar is equivalent to 20 µm. **E)**

Depletion of Fmn2b causes outgrowth and branching defects in motor neurons in

zebrafish larvae causing defects in motor behaviours. The outgrowth and branching

defects in *fmn2b* mutants can be rescued by expression of full-length mouse Fmn2 in

the mutants highlighting the requirement and conserved function of Fmn2 in

development of motor neurons *in vivo*. The inability of mFmn2-I1226A to rescue the

defects in *fmn2b* mutants underscores the requirement of the F-actin nucleation

activity of Fmn2 in outgrowth and branching of motor neurons. The blue and orange

curved arrows indicate the rescue experiments with full length mFmn2 and nucleation

dead mFmn2-I1226A in the *fmn2b* mutant background respectively. The grey and pink

traces indicate the trajectories of wildtype and *fmn2b* mutant embryos respectively in

response to tactile stimulus. The schematic is not drawn to scale.

REFERENCES

Agís-Balboa RC et al. (2017) Formin 2 links neuropsychiatric phenotypes at young age to an increased risk for dementia. EMBO J 36:2815–2828.

Ahuja R, Pinyol R, Reichenbach N, Custer L, Klingensmith J, Kessels MM, Qualmann B (2007) Cordon-Bleu Is an Actin Nucleation Factor and Controls Neuronal Morphology. Cell 131:337–350.

Armijo-Weingart L, Gallo G (2017) It takes a village to raise a branch: Cellular mechanisms of the initiation of axon collateral branches. Mol Cell Neurosci 84:36–47.

Atkins M, Gasmi L, Bercier V, Revenu C, Del Bene F, Hazan J, Fassier C (2019) FIGNL1 associates with KIF1Bβ and BICD1 to restrict dynein transport velocity during axon navigation. J Cell Biol 218:3290–3306.

Bagnall MW, McLean DL (2014) Modular Organization of Axial Microcircuits in Zebrafish. Science (80-) 343:197–200.

Balasanyan V, Watanabe K, Dempsey WP, Lewis TL, Trinh LA, Arnold DB (2017) Structure and Function of an Actin-Based Filter in the Proximal Axon. Cell Rep 21:2696–2705.

Beattie CE, Hatta K, Halpern ME, Liu H, Eisen JS, Kimmel CB (1997) Temporal separation in the specification of primary and secondary motoneurons in zebrafish. Dev Biol 187:171–182.

Bello-Rojas S, Istrate AE, Kishore S, McLean DL (2019) Central and peripheral innervation patterns of defined axial motor units in larval zebrafish. J Comp Neurol 527:2557–2572.

Bernhardt RR, Chitnis AB, Lindamer L, Kuwada JY (1990) Identification of spinal

neurons in the embryonic and larval zebrafish. J Comp Neurol 302:603–616.

Bresciani E, Broadbridge E, Liu PP (2018) An efficient dissociation protocol for

generation of single cell suspension from zebrafish embryos and larvae.

MethodsX 5:1287–1290.

Brustein E, Saint-Amant L, Buss RR, Chong M, McDearmid JR, Drapeau P (2003)

Steps during the development of the zebrafish locomotor network. J Physiol Paris

97:77–86.

Carrington B, Varshney GK, Burgess SM, Sood R (2015) CRISPR-STAT: An easy and

reliable PCR-based method to evaluate target-specific sgRNA activity. Nucleic

Acids Res 43:e157.

Chia PH, Chen B, Li P, Rosen MK, Shen K (2014) Local F-actin network links synapse

formation and axon branching. Cell 156:208–220.

Coles CH, Bradke F (2015) Coordinating Neuronal Actin-Microtubule Dynamics. Curr

Biol 25:R677–R691.

Colombo A, Palma K, Armijo L, Mione M, Signore IA, Morales C, Guerrero N, Meynard

MM, Pérez R, Suazo J, Marcelain K, Briones L, Härtel S, Wilson SW, Concha ML

(2013) Daam1a mediates asymmetric habenular morphogenesis by regulating

dendritic and axonal outgrowth. Dev 140:3997–4007.

Das R, Bhattacharjee S, Letcher JM, Harris JM, Nanda S, Foldi I, Lottes EN, Bobo

HM, Grantier BD, Mihály J, Ascoli GA, Cox DN (2021) Formin3 directs dendritic

architecture via microtubule regulation and is required   for somatosensory

nociceptive behavior. Development.

Dent EW, Gupton SL, Gertler FB (2011) The growth cone cytoskeleton in Axon

786 outgrowth and guidance. Cold Spring Harb Perspect Biol 3:1–39.

787 Dickson BJ (2002) Molecular mechanisms of axon guidance. Science (80-) 298:1959–
788  1964.

789 Drapeau P, Saint-Amant L, Buss RR, Chong M, McDearmid JR, Brustein E (2002)
790  Development of the locomotor network in zebrafish. Prog Neurobiol 68:85–111.

791 Eisen JS, Myers PZ, Westerfield M (1986) Pathway selection by growth cones of
792  identified motoneurones in live zebra fish embryos. Nature 320:269–271.

793 Fassier C, Fréal A, Gasmi L, Delphin C, Ten Martin D, De Gois S, Tambalo M, Bosc
794  C, Mailly P, Revenu C, Peris L, Bolte S, Schneider-Maunoury S, Houart C,
795  Nothias F, Larcher J-C, Andrieux A, Hazan J (2018) Motor axon navigation relies
796  on Fidgetin-like 1-driven microtubule plus end dynamics. J Cell Biol 217:1719–
797  1738.

798 Flanagan-Steet H, Fox MA, Meyer D, Sanes JR (2005) Neuromuscular synapses can
799  form in vivo by incorporation of initially aneural postsynaptic specializations.
800  Development 132:4471–4481.

801 Flynn KC, Bradke F (2020) Role of the cytoskeleton and membrane trafficking in axon–
802  dendrite morphogenesis. In: Cellular Migration and Formation of Axons and
803  Dendrites, pp 21–56. Elsevier.

804 Gallo G (2011) The cytoskeletal and signaling mechanisms of axon collateral
805  branching. Dev Neurobiol 71:201–220.

806 Gallo G (2016) Coordination of the axonal cytoskeleton during the emergence of axon
807  collateral branches. Neural Regen Res 11:709–711.

808 Ghate K, Mutalik SP, Sthanam LK, Sen S, Ghose A (2020) Fmn2 Regulates Growth

Cone Motility by Mediating a Molecular Clutch to Generate Traction Forces.

Neuroscience 448:160–171.

González-Fraga J, Dipp-Alvarez V, Bardullas U (2019) Quantification of Spontaneous

Tail Movement in Zebrafish Embryos Using a Novel Open-Source MATLAB

Application. Zebrafish 16:214–216.

Goode BL, Eck MJ (2007) Mechanism and Function of Formins in the Control of Actin

Assembly. Annu Rev Biochem 76:593–627.

Gordon-Weeks PR, Fournier AE (2014) Neuronal cytoskeleton in synaptic plasticity

and regeneration. J Neurochem 129:206–212.

Gorukmez O, Gorukmez O, Ekici A (2020) A Novel Nonsense FMN2 Mutation in

Nonsyndromic Autosomal Recessive Intellectual Disability Syndrome. Fetal

Pediatr Pathol.

Hale ME, Ritter D a., Fetcho JR (2001) A confocal study of spinal interneurons in living

larval zebrafish. J Comp Neurol 437:1–16.

Hu J, Bai X, Bowen JR, Dolat L, Korobova F, Yu W, Baas PW, Svitkina T, Gallo G,

Spiliotis ET (2012) Septin-driven coordination of actin and microtubule remodeling

regulates the collateral branching of axons. Curr Biol 22:1109–1115.

Kessels MM, Schwintzer L, Schlobinski D, Qualmann B (2011) Controlling actin

cytoskeletal organization and dynamics during neuronal morphogenesis. Eur J

Cell Biol 90:926–933.

Ketschek A, Gallo G (2010) Nerve Growth Factor Induces Axonal Filopodia through

Localized Microdomains of Phosphoinositide 3-Kinase Activity That Drive the

Formation of Cytoskeletal Precursors to Filopodia. J Neurosci 30:12185–12197.

Ketschek A, Jones S, Spillane M, Korobova F, Svitkina T, Gallo G (2015) Nerve growth

factor promotes reorganization of the axonal microtubule array at sites of axon

collateral branching. Dev Neurobiol 75:1441–1461.

Ketschek A, Spillane M, Dun XP, Hardy H, Chilton J, Gallo G (2016) Drebrin

coordinates the actin and microtubule cytoskeleton during the initiation of axon

collateral branches. Dev Neurobiol 76:1092–1110.

Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF (1995) Stages of

embryonic development of the zebrafish. Dev Dyn 203:253–310.

Kundu T, Das SS, Kumar DS, Sewatkar LK, Ghose A (2020) Antagonistic activities of

Fmn2 and ADF regulate axonal F-actin patch dynamics and the initiation of

collateral branching. bioRxiv:2020.11.16.384099.

Kundu T, Dutta P, Nagar D, Maiti S, Ghose A (2021) Coupling of dynamic microtubules

to F-actin by Fmn2 regulates chemotaxis of neuronal growth cones. J Cell Sci.

Kuwada JY (1993) Pathway selection by growth cones in the zebrafish central nervous

system. Perspect Dev Neurobiol 1:195–203.

LaFave MC, Varshney GK, Vemulapalli M, Mullikin JC, Burgess SM (2014) A Defined

Zebrafish Line for High-Throughput Genetics and Genomics: NHGRI-1. Genetics

198:167–170.

Langebeck-Jensen K, Shahar OD, Schuman EM, Langer JD, Ryu S (2019) Larval

Zebrafish Proteome Regulation in Response to an Environmental Challenge.

Proteomics 19:1900028.

Law R et al. (2014) Biallelic truncating mutations in FMN2, encoding the actin-

regulatory protein formin 2, cause nonsyndromic autosomal-recessive intellectual

disability. Am J Hum Genet 95:721–728.

Leader B, Leder P (2000) Formin-2, a novel formin homology protein of the cappuccino

subfamily, is highly expressed in the developing and adult central nervous system.

Lewis TL, Courchet J, Polleux F (2013) Cellular and molecular mechanisms underlying

axon formation, growth, and branching. J Cell Biol 202:837–848.

Lian G, Dettenhofer M, Lu J, Downing M, Chenn A, Wong T, Sheen V (2016) Filamin

A- and formin 2-dependent endocytosis regulates proliferation via the canonical

wnt pathway. Dev 143:4509–4520.

Liu LYM, Lin MH, Lai ZY, Jiang JP, Huang YC, Jao LE, Chuang YJ (2016) Motor

neuron-derived Thsd7a is essential for zebrafish vascular development via the

Notch-dll4 signaling pathway. J Biomed Sci 23:1–11.

Lowery LA, Van Vactor D (2009) The trip of the tip: understanding the growth cone

machinery. Nat Rev Mol Cell Biol 10:332–343.

Marco EJ, Aitken AB, Nair VP, da Gente G, Gerdes MR, Bologlu L, Thomas S, Sherr

EH (2018) Burden of de novo mutations and inherited rare single nucleotide

variants in children with sensory processing dysfunction. BMC Med Genomics

11:50.

Matusek T, Gombos R, Szécsényi A, Sánchez-Soriano N, Czibula Á, Pataki C, Gedai

A, Prokop A, Raskó I, Mihály J (2008) Formin Proteins of the DAAM Subfamily

Play a Role during Axon Growth. J Neurosci 28:13310 LP – 13319.

McLean DL, Fetcho JR (2008) Using imaging and genetics in zebrafish to study

developing spinal circuits in vivo. Dev Neurobiol 68:817–834.

Melançon E, Liu DWC, Westerfield M, Eisen JS (1997) Pathfinding by identified

zebrafish motoneurons in the absence of muscle pioneers. J Neurosci 17:7796–

7804.

Menelaou E, Svoboda KR (2009) Secondary motoneurons in juvenile and adult

zebrafish: Axonal pathfinding errors caused by embryonic nicotine exposure. J

Comp Neurol 512:305–322.

Menon S, Gupton S (2018) Recent advances in branching mechanisms underlying

neuronal morphogenesis .

Montaville P, Jégou A, Pernier J, Compper C, Guichard B, Mogessie B, Schuh M,

Romet-Lemonne G, Carlier MF (2014) Spire and Formin 2 Synergize and

Antagonize in Regulating Actin Assembly in Meiosis by a Ping-Pong Mechanism.

PLoS Biol 12.

Montaville P, Kühn S, Compper C, Carlier MF (2016) Role of the C-terminal extension

of formin 2 in its activation by spire protein and processive assembly of actin

filaments. J Biol Chem 291:3302–3318.

Morello G, Guarnaccia M, Spampinato AG, La Cognata V, D'Agata V, Cavallaro S

(2018) Copy Number Variations in Amyotrophic Lateral Sclerosis: Piecing the

Mosaic Tiles Together through a Systems Biology Approach. Mol Neurobiol

55:1299–1322.

Moreno-Mateos MA, Vejnar CE, Beaudoin J-D, Fernandez JP, Mis EK, Khokha MK,

Giraldez AJ (2015) CRISPRscan: designing highly efficient sgRNAs for CRISPR-

Cas9 targeting in vivo. Nat Methods 2015 1210 12:982–988.

Myers PZ, Eisen JS, Westerfield M (1986) Development and Axonal Outgrowth of

Identified Motoneurons in the Zebrafish.

Nagar D, James TK, Mishra R, Guha S, Burgess SM, Ghose A (2021) The Formin

Fmn2b Is Required for the Development of an Excitatory Interneuron Module in

the Zebrafish Acoustic Startle Circuit. eNeuro 8:ENEURO.0329-20.2021.

Nithianandam V, Chien C-T (2018) Actin blobs prefigure dendrite branching sites. J

Cell Biol 217:3731–3746.

Pamphlett R, Morahan JM, Luquin N, Yu B (2011) Looking for differences in copy

number between blood and brain in sporadic amyotrophic lateral sclerosis.

Muscle Nerve 44:492–498.

Perrone MD, Rocca MS, Bruno I, Faletra F, Pecile V, Gasparini P (2012) De novo 911

Kb interstitial deletion on chromosome 1q43 in a boy with mental retardation and

short stature. Eur J Med Genet 55:117–119.

Pike SH, Melancon EF, Eisen JS (1992) Pathfinding by zebrafish motoneurons in the

absence of normal pioneer axons. Development 114:825–831.

Prokop A, Sánchez-Soriano N, Gonçalves-Pimentel C, Molnár I, Kalmár T, Mihály J

(2011) DAAM family members leading a novel path into formin research. Commun

Integr Biol 4:538–542.

Quinlan ME, Hilgert S, Bedrossian A, Mullins RD, Kerkhoff E (2007) Regulatory

interactions between two actin nucleators, Spire and Cappuccino. J Cell Biol

179:117–128.

Roth-Johnson EA, Vizcarra CL, Bois JS, Quinlan ME (2014) Interaction between

microtubules and the drosophila formin cappuccino and its effect on actin

assembly. J Biol Chem 289:4395–4404.

Sahasrabudhe A, Ghate K, Mutalik S, Jacob A, Ghose A (2016) Formin 2 regulates

the stabilization of filopodial tip adhesions in growth cones and affects neuronal

outgrowth and pathfinding in vivo. Dev 143:449–460.

Saint-Amant L, Drapeau P (1998) Time course of the development of motor behaviors

in the zebrafish embryo. J Neurobiol 37:622–632.

Schymick JC, Scholz SW, Fung H-C, Britton A, Arepalli S, Gibbs JR, Lombardo F,

Matarin M, Kasperaviciute D, Hernandez DG, Crews C, Bruijn L, Rothstein J,

Mora G, Restagno G, Chiò A, Singleton A, Hardy J, Traynor BJ (2007) Genome-

wide genotyping in amyotrophic lateral sclerosis and neurologically normal

controls: first stage analysis and public release of data. Lancet Neurol 6:322–328.

Spillane M, Gallo G (2014) Involvement of Rho-family GTPases in axon branching.

Small GTPases 5.

Spillane M, Ketschek A, Jones SL, Korobova F, Marsick B, Lanier L, Svitkina T, Gallo

G (2011) The actin nucleating Arp2/3 complex contributes to the formation of

axonal filopodia and branches through the regulation of actin patch precursors to

filopodia. Dev Neurobiol 71:747–758.

Stifani N (2014) Motor neurons and the generation of spinal motor neuron diversity.

Front Cell Neurosci 8:293.

Varshney GK, Carrington B, Pei W, Bishop K, Chen Z, Fan C, Xu L, Jones M, LaFave

MC, Ledin J, Sood R, Burgess SM (2016) A high-throughput functional genomics

workflow based on CRISPR/Cas9-mediated targeted mutagenesis in zebrafish.

Nat Protoc 11:2357–2375.

Varshney GK, Pei W, Lafave MC, Idol J, Xu L, Gallardo V, Carrington B, Bishop K,

Jones M, Li M, Harper U, Huang SC, Prakash A, Chen W, Sood R, Ledin J,

Burgess SM (2015) High-throughput gene targeting and phenotyping in zebrafish

using CRISPR/Cas9. Genome Res 25:1030–1042.

Westerfield M, McMurray J V., Eisen JS (1986) Identified motoneurons and their

innervation of axial muscles in the zebrafish. J Neurosci 6:2267–2277.

Yoo H, Roth-Johnson EA, Bor B, Quinlan ME (2015) Drosophila Cappuccino alleles

provide insight into formin mechanism and role in oogenesis. Mol Biol Cell

26:1875–1886.

ACKNOWLEDGEMENTS

The authors thank Dr. N. K. Subhedar (IISER Pune) for critical inputs on the

manuscript. The authors acknowledge the IISER Pune Microscopy Facility, the

National Facility for Gene Function in Health and Disease (NFGFHD) at IISER Pune

and the zebrafish facility at NHGRI Zebrafish Core, NIH for access to equipment and

infrastructure.

FUNDING

The study was supported from grants from the Council of Scientific and Industrial

Research, Govt of India (37(1689)/17/EMR-II), Department of Biotechnology, Govt. of

India (BT/PR26241/GET/119/244/2017) and intramural support from IISER Pune to

A.G.. D.N. is supported by a fellowship from the Council of Scientific and Industrial

Research, Govt of India. D.N. was awarded the Indo-U.S. GETin Internship

(2018_050) (Department of Biotechnology (DBT), Government of India and Indo US

Science and Technology Forum (IUSSTF)) which supported the CRISPR Cas9

mutagenesis work at NHGRI, NIH. The National Facility for Gene Function in Health

and Disease (NFGFHD) at IISER Pune is supported by the Department of

Biotechnology, Govt. of India (BT/INF/22/SP17358/2016). S.M.B. is supported by the

Intramural Research Program of the National Human Genome Research Institute

(ZIAHG200386-06).

AUTHOR INFORMATION

AFFILIATIONS

1. Indian Institute of Science Education and Research (IISER) Pune, Dr Homi

Bhabha Road, Pune 411008, INDIA

Dhriti Nagar and Aurnab Ghose

2. Zebrafish Core, Translational and Functional Genomics Branch, National

Human Genome Research Institute, National Institutes of Health, Bethesda,

Maryland, USA

Blake Carrington

3. Translational and Functional Genomics Branch, National Human Genome

Research Institute (NHGRI), National Institutes of Health (NIH), Bethesda, MD,

USA

Shawn M Burgess

CONTRIBUTION

Conceptualization: D.N. and A.G.; Investigation and formal analysis: D.N.;

Methodology and resources: B.C. and S.M.B.; Writing – original draft: D.N. and A.G.;

Writing – review and editing: D.N., B.C., S.M.B. and A.G; Funding Acquisition: A.G..

All authors gave final approval for publication and agreed to be held accountable for

the work performed therein.

CORRESPONDING AUTHORS

Correspondence: Dr. Aurnab Ghose, Indian Institute of Science Education and

Research (IISER) Pune, Dr Homi Bhabha Road, Pune 411008, INDIA.

Email: aurnab@iiserpune.ac.in

ETHICS DECLARATIONS

1000

Ethics approval

1001

1002   All zebrafish husbandry and experimental protocols complied with institutional

1003   guidelines and were approved by the Institute Animal Ethics Committee (IAEC) and

1004   the Institutional Biosafety Committee (IBSC), IISER Pune or a National Human

1005   Genome Research Institute (NHGRI/NIH) Animal Care and Use Committee approved

1006   animal study protocol.

Consent for publication

1007

1008   All authors gave consent for publication.

Competing interests

1009

1010   The authors declare no competing interests.