LRcell: detecting the source of differential expression at the sub-cell type level from

bulk RNA-seq data.

Wenjing Ma (wenjing.ma@emory.edu)[1], Sumeet Sharma (sumeet.sharma@emory.edu)[2],

Peng Jin (peng.jin@emory.edu)[3], Shannon L. Gourley (shannon.l.gourley@emory.edu)[4],

Zhaohui Qin (zhaohui.qin@emory.edu)[1,5,*]

* Correspondence: zhaohui.qin@emory.edu

[1] Department of Computer Science, Emory University, 400 Dowman Drive, Atlanta, GA

30322, USA

[2] Graduate Program in Neuroscience, Emory University, 1462 Clifton Road NE, Atlanta, GA

30322, USA

[3] Department of Human Genetics, Emory University, 1365 Clifton Road, Atlanta, GA 30322,

USA

[4] Department of Pediatrics, School of Medicine, Emory University, 100 Woodruff Circle,

Atlanta, GA 30322, USA; Yerkes National Primate Research Center, Atlanta, GA 30322,

USA

[5] Department of Biostatistics and Bioinformatics, Rollins School of Public Health, Emory

University, 1518 Clifton Road NE, Atlanta, GA 30322, USA

Abstract

Given most tissues are consist of abundant and diverse sub cell-types, an important yet

unaddressed problem in bulk RNA-seq analysis is to identify at which sub cell-type(s) the

differential expression occur. Single-cell RNA-sequencing (scRNA-seq) technologies can

answer the question, but they are often labor-intensive and cost-prohibitive. Here, we present

LRcell, a computational method aiming to identify specific sub-cell type(s) that drives the

changes observed in a bulk RNA-seq experiment. In addition, *LRcell* provides pre-embedded

marker genes computed from putative single-cell RNA-seq experiments as options to execute

the analyses. Using three different real datasets, we show that *LRcell* successfully identifies

known cell types involved in psychiatric disorders and *LRcell* is more sensitive than even the

leading deconvolution methods.

Keywords: Cell-type enrichment, Cell marker genes, Differential gene expression

Background

Finding genes that are differentially expressed (DE) between experimental conditions is a

powerful approach to understand the molecular basis of phenotypic variation. However, most

tissues are consisted of tens or even hundreds of diverse sub-cell types and DE may only

occur in a small subset of these sub-cell types, which are relevant to the experimental

condition. Bulk RNA-seq data alone is unable to reveal the sub-cell types that drives the DE.

The rapid development and proliferation of single cell technologies resulted in massive

accumulation of single cell transcriptomics data (scRNA-seq) from diverse tissue types.

These data reveal substantial variations in transcriptional regulation among different cell

types and offer an unprecedented close-up view of the modifications underlying important

biological processes including which cell types drive DE [1]. However, steep cost and

complicated protocols prevent the wide-spread adoption of scRNA-seq.

In this study, we propose a novel computational tool named *LRcell*. Given the result from a

bulk RNA-seq DE study, the goal of *LRcell* is to delineate which sub-cell type(s) of the tissue

underwent substantial changes between the two experimental conditions. Exploiting cell

type-specific marker genes identified from generic scRNA-seq available from publicly

available data repositories, *LRcell* achieves the goal by surveying the enrichment of marker

genes across all sub-cell types in the tissue (Figure 1). Thus, no scRNA-seq experiment

matching the bulk RNA-seq experimental condition is needed. When applying *LRcell* to a

diverse panel of bulk RNA-seq DE experiments, we successfully identify known cell types

involved in the pathogenesis of psychiatric disorders as well as produce testable new

hypotheses that have the potential to produce fresh new biological insights.

Results and Discussion

In this work, we collect and curate a compendium of marker genes from multiple published scRNA-seq datasets. We then conduct *LRcell* analysis on multiple bulk RNA-seq DE experiments to demonstrate its utility.

3.1 Marker gene collection and sources

Similar to a collection of gene set for Gene Set Enrichment Analysis (GSEA) [2], *LRcell* requires a compendium of high-quality cell type marker genes. Currently, *LRcell* package provides users with multiple pre-loaded marker gene sets from human blood, human brain and mouse brain (Figure 2.A), computed from scRNA-seq datasets. Additionally, *LRcell* package offers external cell markers collected by Molecular Signatures Database (MSigDB) [3]. The external makers all originate from human species including midbrain, cord blood, ovary and skeletal muscle. We store all cell-type specific marker gene sets into another R Bioconductor ExperimentHub package named *LRcellTypeMarkers*. Additional marker gene sets are being tested and will be added to the collection.

3.2 Microglia highly enriched in Neurodegenerative Dementia

In a recent neurodegenerative dementia study, Swarup and colleagues contrasted TPR50 mice expressing tau mutant with wild type mice using bulk RNA-seq in order to identify gene networks mediating dementia [4]. To identify the cell type(s) most involved in the condition, we apply *LRcell* to the DE gene list using pre-embedded marker genes from adult mouse Frontal Cortex (FC) region [5]. From *LRcell* result, we observed that Microglia show up as highly significant (Figure 2.B) which is concordant with previous studies [6]. Additionally, the FC_11-3.unknown and FC_11-4.unknown sub cell types also show high level of

significance. No annotation is available for these two cell clusters in the original publication.

However, pair-wise comparison of marker genes among all cell clusters reveal that these two

unknown cell clusters have considerable overlaps with the FC_11-1, which is also a

Microglia cell type (Figure S1. A), which explains the pattern we observe.

3.3 CD16+ Monocytes highly enriched in Post-Traumatic Stress Disorder (PTSD)

In a recent study, Breen and colleagues conducted a bulk whole-transcriptome study using

peripheral blood leukocytes collected from U.S. Marines, among which some developed

PTSD post-deployment [7]. Using this dataset, we generate a list of DE genes that show

significant difference between the PTSD group and the control group at the pre-deployment

timepoint.

Using human marker genes derived from a single cell transcriptomic study on PBMC [8],

LRcell analysis finds that cells annotated as CD16+ non-classical Monocytes shows up as the

most significant among all cell types in PBMC (Figure 2. E). Our finding makes biological

sense because as stated in previous studies [9], heterogeneity exists in monocytes

distinguished by CD16 surface proteins and non-classical monocytes have been validated to

regulate immune responses in trauma [10,11].

3.4 Marker genes from different region or timepoints

To apply *LRcell*, an important question is that which marker gene sets to use, i.e., how to

select single cell RNA-seq data where the source of the tissue match the tissue type profiled

in the bulk transcriptomic study. This is particularly important for complex tissues such as

brain. To address this issue, we use the neurodegenerative dementia study [4] as an example,

which contains information from four brain regions: cortex, hippocampus (HC), cerebellum

(CB) and brain stem. Brain stem is excluded from our analysis due to the lack of marker gene

information from that region of the brain.

To understand how marker genes vary across brain regions, we first define marker genes in

all regions of the brain to explore their spatial pattern (Figure S1. B; Figure S1. C). We

observe that glia cells, such as Astrocytes, from different regions have higher number of

overlapping marker genes which indicates the homogeneity of glia cells across the brain. In

contrast, neurons and interneurons share very few marker genes across different brain

regions. We then apply pre-embedded adult mouse brain marker genes from FC, HC and CB

to bulk DE genes obtained from cortex, HC and CB respectively (Figure S3). We observe

that Microglia cells are highly enriched in all three brain regions whereas Astrocytes are

particularly highly enriched in CB (Figure S3). Especially when applying CB marker genes

to CB bulk DE experiment, we notice that one sub-cell type of Astrocytes is highly enriched

compared to others. Our observations demonstrate that selected cell types are heterogeneous

spatially, meaning marker genes are highly specific not only for the cell type, but also which

region the cell belongs to. Because of this finding, it is highly desirable to run *LRcell* using

marker genes of cell types located in closely matched brain regions.

We are also curious whether marker genes curated from scRNA-seq experiments

conducted on non-normal samples is acceptable as the reference. To address this question, we

use data from the HIV vaccine study [8], we observe that the expression of cell-type specific

marker genes is mostly consistent across different timepoints within the same cell type (such

as CD8 cells), and distinct across different cell types (Figure S2.A; Figure S2.B). For

example, B sub-cell types share a considerable number of marker genes across timepoints,

while sharing fewer with other cell types. Thus, although default marker genes used in *LRcell*

are collected from control samples, we believe that marker genes identified from non-normal

samples are acceptable when scRNA-seq data from normal samples are not available.

3.5 Comparison to GSEA

GSEA is a powerful tool to determine whether a pre-defined gene set show concordant shift

in expression when comparing two biological conditions. One could potentially replace

LRcell with GSEA to identify DE-driving cell types, by treating cell-type specific marker

genes as pre-defined gene sets. To compare performance of the two methods, we repeat the

analyses done in section 3.1 and 3.2 using GSEA. The GSEA result from the

neurodegenerative dementia mouse model (Figure 2C) yields several equally significant cell

types including Astrocyte, Endothelial, Microglia, Mural, Oligodendrocyte and

Polydendrocyte. The tied significances lead to difficulties in determining which cell type(s)

potentially participated in dementia pathogenesis. Similar pattern is observed in the GSEA

result on PTSD study (Figure 2F) which shows that Monocytes, Dendritic cells and some T

sub-cell types are equally enriched. Based on the above observations, we conclude that

LRcell is more effective than GSEA to identify sub cell types that are most impacted by the

condition change in bulk DE experiments.

3.5 Comparison to MuSiC

A plethora of deconvolution methods has been developed to infer the proportions of different

sub cell types from bulk transcriptomic data [1][12][13][14][15]. By deconvoluting

individual bulk samples and comparing their proportions, one can identify both cell-type

proportion and gene expression changes. Hence it is of interest to compare *LRcell* with such a

bulk deconvolution strategy. For comparison, we apply MuSiC [13], a deconvolution method

that have shown favorable performance, to the mouse neurodegenerative dementia study [4]

on both conditions. Because some layers of neurons are predicted to have almost zero

proportion (Figure 2D) when using all 81 sub-cell types, we merge the original sub-clusters

into 15 major cell types in order to achieve a better representation. Despite this, MuSiC does

not detect significant difference in Microglia or Astrocyte in terms of their proportions

between the two conditions. When applied to the PTSD study, using the original cell cluster

annotations, MuSiC shows that most of the T sub-cell types have zero proportion and the

proportion of CD14+ Monocytes is up to 60% (Figure S2G). In contrast, *LRcell* produces

more sensible results because it is not limited by the number of cell types as it can detect the

subtle differences among sub-cell types.

3.5 Discussions

Detecting transcriptional activity changes at the individual cell type level, especially their

modifications in disease samples, is crucial for understanding the mechanisms of diseases

development. Deconvolution-based computational tools have been developed to dissect cell

type proportions from bulk gene expression profiles. The computationally inferred

proportions can then be used for comparison. However, when apply to real data, such a

strategy often encounters numerical stability problem and unable to handle large number of

sub-cell types. In this study, we propose a novel alternative strategy named *LRcell*, *LRcell*

conducts enrichment analysis of cell-type specific marker genes among the top (or bottom)

DE genes identified by bulk transcriptome studies. Cell types that show the most enrichment

is likely to play an important role in the condition alteration. When applying to real datasets,

we found *LRcell* can successfully identify the involvement of the Microglia and Astrocytes in

neurodegenerative dementia and rare Monocytes in PTSD.

In spirit, *LRcell* operates similarly as GSEA. But *LRcell* is much more sensitive to minor

differences in marker genes of sub-cell types, which indicates its potential to detect changes

in sub-cell types caused by disease conditions. Additionally, when compared to existing bulk

deconvolution methods, *LRcell* is more stable in its ability to handle the similarities among

sub-cell types. Thus, *LRcell* enables researchers to glean new biological insights from the

bulk transcriptomics experiments with no need of redo the experiment using single cell

technology. We are currently applying *LRcell* to a diverse set of clinical studies (Sharma,

personal communication) to generating more biological insights.

To enable straightforward comparison, currently, we select a fix number of 100 marker genes

from each sub-cell type. Understandably, the number of marker genes for different cell types

varies, it is desirable to allow flexibility in choosing the number of marker genes based on the

transcriptomic patterns across cell types. However, different numbers of marker genes post

challenge for conducting enrichment analyses fairly across all cell types. This will be

investigated in our future studies.

LRcell currently provides embedded marker genes from human blood, human brain and

mouse brain calculated from scRNA-seq experiments along with markers from 66 cell types

in four tissues (midbrain, cord blood, ovary, and skeletal muscle) adopted from MSigDB. We

are working to include more tissue types in the future releases of *LRcell* which will make it

more widely applicable.

Conclusions

In summary, we develop *LRcell*, an R Bioconductor package for identifying sub-cell type(s)

that drive the changes observed in bulk comparative transcriptomic studies, taking

advantages of newly emerged scRNA-seq data. The rationale of *LRcell* is that we believe

marker genes of the modifying cell types tend to be enriched toward the top (or bottom) of

the DE gene lists. We conduct comprehensive surveys applying *LRcell* across various

experimental conditions, and successfully identify cell types that play important roles in

neurodegenerative dementia and PTSD. Hence, we believe *LRcell* can provide researchers

important and new biological insights in terms of the source of the biological changes at the

sub-cell type level, without the need of conducting costly and laborious scRNA-seq

experiments.

Methods

Basic Assumptions

The goal of *LRcell* is to identify the most affected sub-cell type(s) during the transition of

experimental conditions using only bulk transcriptomic data. Based on the assumptions that

cell-type specific marker genes of key cell types tend to be over-represented among the

significant DE genes in bulk transcriptomic studies, *LRcell* can discover which cell type(s) is

involved in certain disease or condition change. In recent years, computational methods have

been developed to deconvolve bulk RNA-seq data to delineate cell-type proportion changes,

which could be borrowed to answer the same question. However, whenever there are ten or

more sub-cell types, the results from deconvolution methods become unreliable. In contrast,

LRcell enables comparison across many more cell types which is important for complex

tissues such as brain.

The development of *LRcell* is inspired by LRpath [16], which is designed for linking

experimental changes to biological pathways or a pre-defined gene set. In *LRcell*, we treat

cell-type specific marker genes as gene sets and calculate the enrichment of each cell type

when comparing two biological conditions. We believe that the most enriched cell type(s) is

highly likely to play an important role in the experimental condition change.

scRNA-seq Data Preprocessing

In this study, we include marker genes from mouse whole brain, human prefrontal cortex

(pFC) and human PBMC, along with 66 cell types' markers from four tissues (midbrain, cord

blood, ovary and skeletal muscle) adopted from MSigDB. For each scRNA-seq dataset, we

first retrieved raw read count matrix. Next, we filtered out low-quality cells and genes and

apply column-wise normalization and log transformation on the data.

The mouse whole brain scRNA-seq dataset [5] produced using the Drop-seq technology

[17], contains nine brain regions from adult mice. The data provided has already been pre-

filtered by the authors. For cell types other than neurons, we directly utilized the information

provided on the study website. For neurons and interneurons, we curated the sub-cell types

following the original study.

The human pFC scRNA-seq dataset [18], produced by 10X Genomics Chromium, is

derived from the pFC region (specifically BA9). The dataset contains two conditions: healthy

controls and major depressive disorder (MDD). We split the data matrix into two parts and

filtered out cells expressing less than 10 genes and genes expressed in less than 10 cells

respectively. We also filtered out mitochondrial, ribosomal genes, and genes from annotation

clusters (Astros_1, Mix_1, Mix_2, Mix_3, Mix_4, Mix_5, Inhib_4_SST).

The PBMC dataset [8], generated by CITE-seq technology [19], is derived from an HIV

vaccine trial study which involves eight volunteers at three timepoints: immediately before,

three days and seven days after the vaccine. The study contains 161,764 cells in total. To

accelerate the marker gene selection, we separated the count matrix according to the time

label and filtered out low-quality cells and genes (mitochondrial, ribosomal genes and those

expressed in less than 1,000 cells). The cluster annotated as "Doublet" was filtered out.

Marker Gene Selection

After obtaining the log-normalized gene expression matrix along with high-quality sub-cell

type clusters, we calculated the enrichment scores for each cluster (sub-cell type) using the

marker gene selection method described in Marques et al[20]. The cluster-specific gene

enrichment is defined as the average gene expression levels of cells in that cluster divided by

the average gene expression levels in all cells. The enrichment score is adjusted by

introducing a penalty representing the fraction of cells in that cluster expressing the marker

gene. Combined, this score allows the identification of genes with cluster-specific high

expression values to be selected as marker genes. The description below is adapted from the

original publication.

Suppose there are a total of M genes, L different clusters each with N_j cells, and the

total number of cells is N. Let $E = \{E_{ijk}\}$ represent the gene by cell read count matrix. Here

$i = 1, \dots, M, j = 1, \dots, L, k = 1, \dots, N_j$ and $N = \sum_{j=1}^{L} N_j$. The overall average expression of

the ith gene across all cells is defined as:

290
$$\bar{E}_{i..} = \frac{1}{N} \sum_{j=1}^{L} \sum_{k=1}^{N_j} E_{ijk}$$

291   The average expression of gene i in the jth cluster is defined as:

292
$$\bar{E}_{ij.} = \frac{1}{N_j} \sum_{k=1}^{N_j} E_{ijk}$$

293   The enrichment for gene i in the jth cluster as:

294
$$Enrichment_{i,j} = \frac{\bar{E}_{ij.}}{\bar{E}_{i.}}$$

295   Next, we consider the proportion expressing the gene *i* in the *j*th cluster as:

$$Prop_{i,j} = \frac{1}{N_j} \sum_{k=1}^{N_j} I(E_{ijk} > 0)$$

297   The $I(\cdot)$ is an indicator function.

298   The enrichment score for gene *i* in the *j*th cluster is computed as:

$$Score_{i,j} = Enrichment_{i,j} \times (Prop_{i,j})^{power}$$

where "power" is a hyperparameter to be tuned manually to control the penalization for the

cell cluster proportion term. The power parameter is set to 1 throughout this study. After

calculating the weighed gene enrichment scores in each cluster, we ranked genes based on the

scores and selected the top 100 genes as the marker genes for each cluster.

MSigDB Marker Genes

We downloaded cell marker gene sets from MSigDB category C8 – cell type signature gene

sets. Since not all tissue types are suitable for *LRcell*, we applied the following criteria to

select tissues: 1) non-fetal tissues; 2) have more than 8 (sub-) cell types; 3) minimum number

of marker gene greater than 50; 4) median number of marker genes greater than 80. In the

end, four tissue types, the midbrain, cord blood, ovary and skeletal muscle remain.

Bulk RNA-seq Data Preprocessing

The raw count of mouse bulk RNA-seq study on neurodegenerative dementia was

downloaded from GEO (Accession number: GSE90693). DE analysis is performed using

DESeq2 [21] to obtain DEGs in each brain region.

The raw count of bulk RNA-seq study on PTSD was downloaded from Recount2

[22]. We extracted out the experiment contrasting PTSD cases and healthy controls with

timepoint of pre-employment and performed DESeq2 to obtain DEGs.

LRcell Analysis

LRcell is inspired by LRpath, which was designed for identifying sets of predefined gene sets

that show enrichment with differentially expressed transcripts in microarray experiments.

LRcell uses logistic regression or linear regression to assess whether marker genes (as defined

in the Marker Gene Selection sub-section above) of a specific cell type are more likely to be

DE genes in a particular bulk RNA-seq study. The linear regression option is added to handle

the continuous enrichment scores of marker genes. Users can choose accordingly. To

facilitate our analysis, we assume that the major sub cell types of the tissue their marker

genes are known *a priori*.

We apply *LRcell* to each cell type independently. The required input includes a list of

DE genes ranked by the level of significance, and a set of marker genes for each cell type.

Then *LRcell* runs a logistic regression as

332
$$log \frac{\theta}{1 - \theta} = \alpha + \beta x$$

333 and

334
$$\theta = P(Y = 1)$$

In which $Y = 1$ denotes that gene is a marker gene and $Y = 0$ otherwise. Hence θ

represents the chance that the gene is a marker gene. We use $-\log (p - \text{value})$ as the

explanatory variable x. Whether a specific cell type is involved in the experimental condition

change is evaluated by testing the null hypothesis that $\beta = 0$ against the alternative that $\beta \neq$

0 using the Wald test. Typically, we run *LRcell* on all sub cell types found in the tissue to see

which sub cell type(s) drives the changes.

Similar to logistic regression, linear regression directly performs

342
$$Y = \alpha + \beta x$$

where *Y* indicates the enrichment scores of genes. Same as logistic regression, the p-value

can be obtained from testing the null hypothesis that $\beta = 0$ against the alternative that $\beta \neq 0$

using the t-test.

Once the p-values are obtained, we calculate False Discover Rate (FDR) using

p.adjust() function in R to adjust p-values with Benjamini-Hochberg method.

Input and Output

LRcell requires two inputs: (1) a ranked list of genes with DE p-values in a bulk RNA-seq

experiment; (2) sets of marker genes from all sub-cell types of the bulk tissue acquired from

scRNA-seq datasets *a priori* or from MSigDB C8—cell type signature gene sets. For those cell

markers derived from scRNA-seq datasets, we offer choices for users to choose between species

as human or mouse and the region indicates the specific brain region or PBMC. For MSigDB cell

markers, we store the marker genes into the *LRcellTypeMarkers* packages which can be easily

downloaded. When running LRcell() function, the Logistic Regression (LR) option is set as the

default, while users can also set the method option as LiR if Linear Regression is desired. For

linear regression, an enrichment score is needed as input for each gene whereas gene sets are

sufficient for logistic regression. For MSigDB cell type signature gene set, LR option is

recommended as there is no enrichment score information available. For customized input, i.e., a

scRNA-seq data, we offer a LRcell_gene_enriched_scores function which takes the read counts

matrix and cell annotation as input to generate enrichment scores for genes in each cell type. For

further subsetting, get_markergenes can be used for generating marker genes for more specific

sub cell types.

The output is a list of significance p-value (or FDR), one for each sub cell type. For

visualization, *LRcell* produces Manhattan plot, which can be drawn through

plot_manhattan_enrich function. We also provided a plot (plot_marker_dist function) indicating

where certain cell-type specific marker genes locate on the bulk DE genes. The bulk DE genes

are sorted using -log10(p-value) × sign(log2FoldChange) which could potentially give

information on both up-/down-regulated directions. More detailed information about *LRcell* is

available at http://bioconductor.org/packages/release/bioc/html/LRcell.html.

LRcell requires an R version beyond 4.1 and a prerequisite installation of BiocManager.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets analyzed during the current study are available in the Gene Expression Omnibus

(GEO) with following accession numbers: mouse whole brain [5] (GSE116470), human pFC

[18] (GSE144136), human PBMC [8] (GSE164378), neurodegenerative dementia study [4]

(GSE90696) and PTSD study [7] (GSE64814). The R package is freely available on

Bioconductor (https://doi.org/doi:10.18129/B9.bioc.LRcell) and the external marker genes

are stored in another R package named *LRcellTypeMarkers* on Bioconductor

(https://doi.org/doi:10.18129/B9.bioc.LRcellTypeMarkers).

Competing interests

The authors declare that they have no competing interests.

Author's contributions

ZQ and WM conceived the idea and supervised the study. WM realized the method and

developed the R package. WM analyzed the real data and conducted the computational

experiments. SS provided guidance on validating biological experiments and tested the

package. WM and ZQ summarized the results and wrote the manuscript. All authors have

read and approved the manuscript.

References

1. Newman AM, Steen CB, Liu CL, Gentles AJ, Chaudhuri AA, Scherer F, et al. Determining cell type abundance and expression from bulk tissues with digital cytometry. Nat Biotechnol. 2019;37:773–82.

2. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. PNAS. 2005;102:15545–50.

3. Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst. 2015;1:417–25.

4. Swarup V, Hinz FI, Rexach JE, Noguchi K, Toyoshiba H, Oda A, et al. Identification of evolutionarily conserved gene networks mediating neurodegenerative dementia. Nature medicine. Nature Publishing Group; 2019;25:152–64.

5. Saunders A, Macosko EZ, Wysoker A, Goldman M, Krienen FM, Rivera H de, et al. Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain. Cell. 2018;174:1015-1030.e16.

6. Perry VH, Nicoll JA, Holmes C. Microglia in neurodegenerative disease. Nature Reviews Neurology. Nature Publishing Group; 2010;6:193.

7. Breen MS, Maihofer AX, Glatt SJ, Tylee DS, Chandler SD, Tsuang MT, et al. Gene networks specific for innate immunity define post-traumatic stress disorder. Molecular psychiatry. Nature Publishing Group; 2015;20:1538–45.

8. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated analysis of multimodal single-cell data. Cell. 2021;184:3573-3587.e29.

9. Ong S-M, Teng K, Newell E, Chen H, Chen J, Loy T, et al. A novel, five-marker alternative to CD16–CD14 gating to identify the three human monocyte subsets. Frontiers in immunology. Frontiers; 2019;10:1761.

10. Kratofil RM, Kubes P, Deniset JF. Monocyte Conversion During Inflammation and Injury. Arterioscler Thromb Vasc Biol. 2017;37:35–42.

11. Kuan P-F, Yang X, Clouston S, Ren X, Kotov R, Waszczuk M, et al. Cell type-specific gene expression patterns associated with posttraumatic stress disorder in World Trade Center responders. Transl Psychiatry. 2019;9:1.

12. Gong T, Szustakowski JD. DeconRNASeq: a statistical framework for deconvolution of heterogeneous tissue samples based on mRNA-Seq data. Bioinformatics. 2013;29:1083–5.

13. Wang X, Park J, Susztak K, Zhang NR, Li M. Bulk tissue cell type deconvolution with multi-subject single-cell expression reference. Nat Commun. 2019;10:1–9.

14. Baron M, Veres A, Wolock SL, Faust AL, Gaujoux R, Vetere A, et al. A single-cell transcriptomic map of the human and mouse pancreas reveals inter-and intra-cell population structure. Cell systems. Elsevier; 2016;3:346-360. e4.

15. Tsoucas D, Dong R, Chen H, Zhu Q, Guo G, Yuan G-C. Accurate estimation of cell-type
composition from gene expression data. Nat Commun. 2019;10:1–9.

16. Sartor MA, Leikauf GD, Medvedovic M. LRpath: a logistic regression approach for
identifying enriched biological groups in gene expression data. Bioinformatics. 2009;25:211–
7.

17. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly parallel
genome-wide expression profiling of individual cells using nanoliter droplets. Cell. Elsevier;
2015;161:1202–14.

18. Nagy C, Maitra M, Tanti A, Suderman M, Théroux J-F, Davoli MA, et al. Single-nucleus
transcriptomics of the prefrontal cortex in major depressive disorder implicates
oligodendrocyte precursor cells and excitatory neurons. Nature neuroscience. Nature
Publishing Group; 2020;23:771–81.

19. Stoeckius M, Hafemeister C, Stephenson W, Houck-Loomis B, Chattopadhyay PK,
Swerdlow H, et al. Simultaneous epitope and transcriptome measurement in single cells.
Nature Methods. 2017;14:865–8.

20. Marques S, Zeisel A, Codeluppi S, van Bruggen D, Falcão AM, Xiao L, et al.
Oligodendrocyte heterogeneity in the mouse juvenile and adult central nervous system.
Science. American Association for the Advancement of Science; 2016;352:1326–9.

21. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15:550.

22. Collado-Torres L, Nellore A, Kammers K, Ellis SE, Taub MA, Hansen KD, et al.
Reproducible RNA-seq analysis using recount2. Nat Biotechnol. 2017;35:319–21.

Figures

Figure 1 Overview of *LRcell* workflow. (A) Bulk RNA-seq experiment design with control
and disease conditions. (B) Illustration of a marker gene list derived from scRNA-seq
experiment. (C) (upper panel) Illustration of how the distribution of marker genes on bulk DE
genes may affect the regression; (lower panel) Illustration of the FDR rate for each cell type
in which the grey cells and dark blue cells are most likely involved in the condition change.

478
Figure 2 *LRcell* datasets and *applying LRcell* to real cases. (A) summary of the all tissue
types in which marker genes have been pre-embedded in *LRcell*. (B) *LRcell* result of mapping
the bulk neurodegenerative dementia DE genes to the mouse brain frontal cortext (FC)
region . Microglia is shown as the most significant cell type. (C) GSEA result of mapping the
bulk neurodegenerative dementia DE genes using the same marker genes for *LRcell* (mouse
brain FC) as input. We see numerous dots have the same significance level which is difficult
to interpret. (D) Cell type proportions for control and disease samples calculated by MuSiC.
Each box contains 17 individuals. The x-axis is ordered by the t-test significance between the
two conditions. (E) *LRcell* result of mapping bulk PTSD DE genes to human PBMC. CD16+
Monocytes is shown as the most significant cell type. (F) GSEA result of mapping bulk
PTSD DE genes to human PBMCusing the same marker genes for *LRcell* (human PBMC) as
input. Again, we see numerous dots have the same significances. (G) Cell type proportions
for control and disease samples calculated by MuSiC. The x-axis is ordered by the t-test
significance between two conditions. As shown in the figures, MuSiC failed to identify all
cell type proportions as there are plenty of 0s starting from ASCD_pDC cell type and the
CD14+ Monocytes are wrongly inferred.

495
496
Figure S1: Marker genes overlap between different brain regions. Top 100 marker genes
are selected for each cell type, thus, the maximum overlap in these figures is 100. (A)
heatmap illustrating the overlap of marker genes among cell types within the Frontal Cortex
(FC) region derived from mouse whole brain scRNA-seq dataset. The highlighted area
describes the overlap between FC_11-3.unknown, FC_11-4.unknwon and FC_11-1.Microglia
as an illustration for the similarity between these three sub-cell types. (B) Heatmap
illustrating the overlap of marker genes among cell types within the FC and cell types within
the Cerebellum (CB). (C) Heatmap illustrating the overlap of marker genes among cell types
within the FC and cell types within the Hippocampus (HC). As shown in (B) and (C), the
overlaps between cell types belong to different brain regions are lower than overlaps between
cell types belong to the same brain region.

508
Figure S2: Marker genes overlap across time points from PBMC dataset. Top 100
marker genes are selected for each cell type and the maximum overlap in figures is 100. (A)
Heatmap illustrating the overlap of the cell-type-specific marker genes between time point 0
and time point 3. (B) Heatmap illustrating the overlap of the cell-type-specific marker genes
between time point 3 and time point 7. As shown in the figure, the diagonal pattern indicates
the marker genes selected are highly cell-type-specific and shared across time points.

515
**Figure S3. Comparison of *LRcell* analysis results when the tissue from which the DE
gene lists is obtained match or un-match the tissue type from where the marker genes
are collected.** *LRcell* result when: (A) Both DE gene list and marker genes are obtained from
HC. (B) Both DE gene list and marker genes are obtained from CB. (C) DE gene list obtained
from HC and marker genes collected from FC. (D) DE gene list obtained from CB and

marker genes collected from FC. (E) DE gene list obtained from CB and marker genes
collected from HC. (F) DE gene list obtained from FC and marker genes collected from HC.
(G) DE gene list obtained from FC and marker genes collected from CB. (H) DE gene list
obtained from HC and marker genes collected from CB. Compared to (D) and (E), which
analyze CB bulk DE genes, (B) has a higher resolution as the FDR ranges from 0 to 12.
526

A Marker genes overlap (time0 vs time3)

B Marker genes overlap (time3 vs time7)