Lung injury induces alveolar type 2 cell hypertrophy and polyploidy with

implications for repair and regeneration

Anthea Weng[1], Mariana Maciel-Herrerias[1], Satoshi Watanabe[1], Lynn Welch[1], Annette S.

Flozak[1], Rogan Grant[1], Raul Piseaux Aillon[1], Laura Dada[1], Seung Hye Han[1], Monique

Hinchcliff[3], Alexander Misharin[1], GR Scott Budinger[1], Cara J. Gottardi[1,2]*

Northwestern University, Feinberg School of Medicine

[1]Department of Pulmonary Medicine, [2]Biochemistry and Molecular Genetics

Chicago, IL 60611, USA

[3]Department of Medicine, Division of Rheumatology, Allergy and Immunology, Yale University

School of Medicine

New Haven, CT 06510 USA

Running title: Injury-induced alveolar epithelial polyploidization

Corresponding author*:

Cara J. Gottardi

303 East Superior Street

Simpson-Querrey 5-525

Chicago, IL 60611, USA

c-gottardi@northwestern.edu

312-503-4123

4378 (Abstract, Results, Discussion, Figure legends; minus Methods) words

7 Figures

Abstract

Epithelial polyploidization post-injury is a conserved phenomenon, recently shown to improve

barrier restoration during wound healing. Whether lung injury can induce alveolar epithelial

polyploidy is not known. We show that bleomycin injury induces AT2 cell hypertrophy and

polyploidy. AT2 polyploidization is also seen in short term ex vivo cultures, where AT2-to-AT1

trans-differentiation is associated with substantial binucleation due to failed cytokinesis. Both

hypertrophic and polyploid features of AT2 cells can be attenuated by inhibiting the integrated

stress response (ISR) using the small molecule ISRIB. These data suggest that AT2

polyploidization may be a feature of alveolar epithelial injury. As AT2 cells serve as facultative

progenitors for the distal lung epithelium, a propensity for injury-induced binucleation has

implications for AT2 self-renewal and regenerative potential upon re-injury, which may benefit

from targeting the ISR.

Keywords: Alveolar epithelial cell, injury, binucleation, polyploidy, cytokinesis, integrated stress

response

Introduction

Although most patients recover from severe lung injury induced by SARS-CoV-2 pneumonia or other insults, some will develop persistent and progressive lung fibrosis (1-3), highlighting a growing imperative to understand how the distal lung epithelium repairs after injury (1, 3-6). While it is clear surfactant producing-alveolar type 2 pneumocytes (AT2 cells) are progenitors that give rise to alveolar type 1 pneumocytes (AT1 cells)(7, 8), the variability between individuals in the AT2-to-AT1 cell trans-differentiation program to restore normal alveolar function is unexplained.

Pathologists have long recognized that a range of injurious stimuli lead to the accumulation of abnormally large AT2 cells, described as "hypertrophic or hyperplastic" AT2 cells (9-12). Morphologically similar cells are seen in autopsy specimens or lung explants from patients with lung injury secondary to SARS-CoV-2 pneumonia (1, 5), but mechanisms underlying this cell state and its consequences for lung repair post-injury remain puzzling. A recent convergence of single cell transcriptomic-studies from a range of mouse injury models suggests that instead of fully progressing to an AT1 cell fate, AT2 cells become stalled at an intermediate transition-state characterized by activation of cell cycle arrest pathways (p16, p21, p53),TGFβ signaling and matrix remodeling (SERPINE1)(13-16). Serially collected single cell RNA-sequencing data over the course of bleomycin induced injury shows the formation of this intermediate transition state is preceded by increased expression of genes in the integrated stress response (ISR)(15). In tissue sections, these transitional cells are characterized by elevated levels of specific intermediate filament keratins (KRT8/18 in mice; KRT17 in humans) and other structural proteins, along with an intermediate AT2-to-AT1 morphology (15, 16). These data suggest small cuboidal AT2 cells differentiate into large and flattened AT1 cells through a morphologically stressful transition that appears prone to stalling after injury. Importantly, recent evidence suggests the AT2-to-AT1 transition state is not fixed, as therapeutic targeting of p53 or ISR signaling appears to guide AT2 cells through this vulnerable trans-differentiation step (16, 17). However, the cellular processes most vulnerable to this morphogenetic transition, which drives persistence of the AT2-to-AT1 transition state in injured lungs, and can be reversed by these the therapies are unknown.

Epithelial polyploidization and consequent hypertrophy post-injury is a conserved phenomenon observed from flies to vertebrates (18). It is a process by which cells increase their DNA content and bypass mitosis. Polyploid cells are larger than their diploid counterparts because cell size

scales with DNA content (19). Cell hypertrophy is advantageous during epithelial repair, as

larger cells manifest less junctional surface area per unit of epithelium, leading to decreased

permeability in an otherwise leaky-barrier environment (20, 21). Increased DNA copies can also

favor adaptation to cell and environmental stress, in ways that are context and injury dependent

(22-24). However, these benefits of polyploidization come at a cost, as polyploid cells will less

faithfully segregate DNA during mitosis (25-27). Thus, polyploidization of differentiated cell

types—particularly those that serve also as progenitors (e.g., AT2 cells)—is likely to adversely

impact future regenerative potential of the tissue. Whether AT2 polyploidy underlies injury-

induced AT2 hypertrophy and drives persistence of the AT2-to-AT1 cell stalled state is

unknown.

In this study, we show that lung injury leads to AT2 hypertrophy and polyploidization—notably

binucleated AT2s—during the acute injury response. *Ex vivo* analysis suggests the route to

polyploidy is via failed cytokinesis during the AT2-to-AT1 flattening process. A small molecule

inhibitor of the integrated stress response, ISRIB, which we have previously shown ameliorates

fibrosis development in murine models (17), inhibits the abundance of hypertrophic, polyploid

AT2 cells. These data suggest that modulating stress signals during the morphogenetically

challenging AT2-to-AT1 transition may improve key steps through which injured AT2 cells divide

and differentiate to restore the alveolar epithelium. Implications of failing to remove and replace

polyploid AT2 cells after lung injury are discussed.

Results

Lung injury induces AT2 hypertrophy

To understand how lung injury alters the morphogenetic sequence by which AT2 cells grow,

divide and differentiate to an AT1 fate, we used a genetic lineage tracing system to permanently

label AT2 cells in $Sftpc$^Cre-ER; Lox-Stop-Lox^zsGreen mice subjected to the bleomycin model of injury

and fibrogenic repair (17) (Fig.1A). As with previous studies (9-12), we confirm that bleomycin

injury leads to zsGreen+ cells that are larger than AT2 lineage-labeled mice exposed to saline

(Fig. 1B and C, quantification from 2 mice, 5 fields of view; Frozen section image analysis

pipeline in Fig. S1A). Cell size difference may be more readily apparent in 14μm thick frozen

sections than thinner 4μm paraffin sections, likely because the former method favors cell

visualization along a greater portion of the curved alveolus (Fig. S1B). The injury-induced size

increase is not an aberrant feature of cells expressing the fluorescence lineage-label, as AT2

cells lacking zsGreen but expressing surfactant protein C (SpC) also increase in size (Fig.

S1C). Lastly, we can also detect this size increase by flow cytometric analysis of AT2 cells (Fig.

S2A-C; Median Forward Scatter Area quantification from 3 mice/condition), demonstrating the

injury-induced phenotype is not an artefact of lung fixation or processing.

Lung injury promotes AT2 cell polyploidization

As cell size can scale with DNA context (19), we sought to examine nuclear features of

hypertrophic AT2 cells. Interestingly, we found a number of lineage-labeled AT2 cells with two

nuclei (Fig. 2A), often observing 1-2 binucleated zsGreen+ cells per field of view, estimating a

rate of ~8% (Fig. 2B; see also Fig.1B inset box). To validate AT2 polyploidy via an independent

method, we sought to quantify and characterize these cells by flow cytometry paired with DNA

content analysis (Fig. S2A-E). Although we readily detected AT2 cells with 2N or 4N DNA

content, too few 4N cells survived the cytospin procedure to reliably quantify binucleation rate

differences between saline and bleomycin-treated mice (Fig. S2E). New methods will be

required to reliably isolate and quantify this apparently fragile cell state.

Of interest, AT2 binucleation appears compatible with expression of the AT1 marker, podoplanin

(Fig. 2A-a' magenta arrow), but we can also detect binucleated cells expressing the AT2

marker, surfactant protein C (SpC) (Fig. S2F). These data suggest alveolar epithelial polyploidy

may be compatible with either AT2 or AT1 cell state. To address wither AT2 polyploidy is a

typical feature of the recently described AT2-to-AT1 cell transition state (14, 15), we

interrogated Krt8 expression in our lung sections. While we can find large binucleated cells that

highly express Krt8 protein, AT2 binucleation does not exclusively correlate with a Krt8-high

state (Fig. 3). Remarkably, binucleation is a common feature of primary AT2 cultures grown on

glass or filters (Fig. 4). We are confident these *ex vivo* polyploid cells derive from *bona fide* AT2

cells, as they are uniformly Krt8-positive (Fig. 4A) and can be lineage-labeled using the *Sftpc-*

^{CreER; Lox-stop-Lox eYFP} system (Fig. 4B,C). Even AT2-derived organoids, which can show AT2-to-

AT1 cell transition state features seen in injured lungs (16), exhibit a binucleation rate similar to

in vivo estimates (Fig. 4E,F vs Fig. 2B). Mechanistically, live-imaging reveals binucleated cells

can be generated via failed cytokinesis, further suggesting that how AT2 polyploidy is generated

and maintained can be interrogated in *ex vivo* cultures (Fig. 5).

Given limitations of assessing polyploidy from frozen or thick sectioning (Fig. S1B), together

with evidence that mononuclear polyploidy also exists (28), we sought an independent method

to assess the polyploidy rate in injured AT2 cells. A recent study showed that the stochastic,

multi-color Confetti reporter system can be used to assess mononuclear polyploidy in mouse

tissues, notably liver (29). While the Confetti reporter mouse is better known for tracing clonal

cell relationships in development and injury (8, 30), this tool can also validate polyploid cells in

tissues with single cells expressing two or more fluorescent proteins (Fig. S3A). In contrast to

cell clone tracing, the lineage label must be activated by tamoxifen *after genome duplication* to

detect single cell polyploidy. By subjecting $Sftpc^{Cre-ER}$ x $Rosa^{Confetti}$ mice to the bleomycin model

of injury, we could identify large, partially flattened cells that express both green fluorescent

protein (GFP) and red fluorescent protein (RFP) lineage labels surrounding what appears as a

single nucleus of lineage-label exclusion (Fig. S3B,C). We also identified small round,

GFP/RFP-dual-positive cells that likely reflect lineage-labeled AT2 cells which have doubled

their DNA content and arrested at the G2-phase of the cell cycle (Fig. S3D). As these dual-

positive cells appear an order of magnitude more rare than the AT2 binucleation rate estimated

in Figure 1 (e.g., < 0.2% lineage-labelled AT2 cells), we may be missing polyploidization events

during the AT2-to-AT1 transition, when $Sftpc^{Cre}$-expression is likely waning. Crossing the

$Rosa^{Confetti}$-reporter strain to a Cre-driver whose expression increases during the AT2-to-AT1

transition may be required to assess the full extent of alveolar epithelial polyploidization during

the lung injury-repair sequence. Nonetheless, these data show that lung injury can generate

binucleated polyploid AT2 cells (Fig. 1), and possibly polyploid mononuclear AT2 cells (Fig.

S3C), where polyploidization likely occurs during the post-injury, reparative phase of the

bleomycin model.

Inhibiting the Integrated Stress Response attenuates AT2 hypertrophy

To determine whether injury-induced AT2 hypertrophy and polyploidy can be inhibited or

reversed, we considered small molecule approaches known to target stress-activated pathways,

recently suggested to guide injured AT2 cells through their vulnerable transition towards an AT1

fate (16, 17). The integrated stress response (ISR) is a highly conserved signaling network that

helps cells adapt to a range of environmental stresses, through activating kinases that inhibit

bulk protein translation in favor of the translation of select mRNAs encoding transcription factors

such as ATF4 that promote the transcription of molecular chaperones and other stress adaptive

genes (31). A central integrator of these diverse stress-responsive kinases is the eIF2

complex, where phosphorylation of eIF2a at a single serine residue limits translation (32). In this

way, eIF2a phosphorylation and its diverse downstream consequences comprise the ISR (33,

34). A small molecule integrated stress response inhibitor, ISRIB, was found to render cells

insensitive to eIF2α phosphorylation, thereby restoring translation capacity to normalize cell

function (35-37) and limit a range of tissue pathologies in murine disease models (38-42). Our

team recently discovered that ISRIB improves fibrosis in murine models by limiting abundance

of AT2 cells stalled at the AT2-to-AT1 transition state (17). Since bleomycin-injured AT2 cells

show activation of ISR-associated genes (17), we reason persistent activation of the ISR in AT2

cells can thwart alveolar epithelial repair. However, particular morphogenetic features of the

AT2-to-AT1 transition that drive persistence of this cell stress state remain unclear. Remarkably,

we find that bleomycin-injured *Sftpc*[Cre-ER; Lox-Stop-Lox]zsGreen mice treated with ISRIB show

complete inhibition of AT2 hypertrophy, and no evidence of binucleation (Fig. 6A,B). Although

ISRIB did not prevent AT2 cell binucleation in *ex vivo* cell cultures, ISRIB treatment did limit the

rate of binucleation, as well as formation of the largest cells, inferred from nuclear size within

these cells (Fig. 6C-F). These data suggest that the AT2 hypertrophic response to injury is

sensitive to inhibition of the ISR. Evidence for modest cell size attenuation in *ex vivo* AT2

cultures suggests that ISRIB may limit hypertrophic growth through an AT2 cell-autonomous

mechanism.

Discussion

For decades, clinical pathologists have described the development of "hypertrophic" AT2 cells after injury, but mechanisms that drive this state and degree to which it benefits and/or constrains the lung injury-repair cycle is not known. We show that some of this hypertrophic growth is associated with AT2 polyploidy, most notably binucleated AT2 cells. *Ex vivo* analysis suggests the route to polyploidy is via failed cytokinesis during the AT2-to-AT1 flattening process. Remarkably, attenuating the ISR with ISRIB limits the abundance of hypertrophic, polyploid AT2 cells after bleomycin injury and *ex vivo* culture, suggesting the effects of ISRIB may be AT2 cell-autonomous. Given recent high resolution time-series sampling of the injured lung and analysis by single cell RNA-sequencing revealing that the AT2-to-AT1 transition state expresses genes associated with the ISR (15), we suggest that lung injury persistently activates the ISR in AT2 cells as they enlarge and flatten to repair the alveolar epithelium, increasing their susceptibility to cytokinesis failure into polyploid AT2 cells. A key consequence of AT2 polyploidization will be loss of the AT2 stem cell daughter, which may compromise future regenerative potential of alveoli served by that AT2 progenitor. We propose that attenuating the ISR helps guide AT2 cells through this vulnerable cell-division/trans-differentiation morphogenetic sequence without polyploidy (Fig. 7, Model).

Evidence that AT2 binucleation does not exclusively correlate with cells expressing the highest levels of Krt8 protein is noteworthy. While elevated Krt8 expression is associated with an aberrant AT2-to-AT1 transition state (14, 15), perhaps this transition is only one of a number of transcriptionally distinct routes cells take as they repair the injured alveolar epithelium. Given the increased size and presumed fragility of polyploid alveolar cells, we suspect they are inadequately represented in current lung single cell RNA-sequencing pipelines. To understand how these cells specifically contribute to alveolar repair, it will be necessary to revise existing cell sorting pipelines to quantify the abundance and degree of alveolar epithelial polyploidy, as well as isolate these cells for downstream transcriptomic analyses to establish the distinct features of the polyploid cell state (26).

Recent injury-repair studies using live-imaging of zebrafish epicardium as a simple epithelial model system reveal that cells along a closing wound-front manifest higher tension and rates of binucleation due to failed cytokinesis (21). While one study suggests this may be due to excessive integrin-based extracellular matrix adhesive tension that ultimately interferes with

contractility of the actin-myosin cytokinetic ring (43), whether persistent mechano-activation of

cell-cell adhesive junctions also contribute remains unclear.

Although bleomycin-injury leads to binucleation in only ~5% of lineage-labeled AT2 cells, total

alveolar epithelial cell polyploidy may be much greater, particularly given evidence for

mononuclear polyploidy across tissues (28). We find that injury induces a number of

hypertrophic, SpC-expressing cells manifesting greatly enlarged single nuclei, but newer

methods will be required to distinguish true mononuclear polyploidy (i.e., $\geq$4N DNA content in a

G_1-phase cell) from AT2 cells arrested post-genome doubling at the G_2-phase (i.e., also 4N

DNA content) (44). Another limitation of this study is our reliance on the DNA-damaging agent

bleomycin as an injury model, which could bias the alveolar repair response towards AT2

polyploidy. Future work will be required to determine the generality of this paradigm, and

whether AT2 polyploidy can be induced by other forms of injury (e.g., viral or pneumonectomy

model of alveologenesis).

Conceptually, it will be important address whether AT2 polyploidization is restricted to the

transdifferentiating daughter, sparing normal ploidy of the renewed AT2 or if lung injury

promotes AT2 endocycling/endomitosis, resulting in a polyploid facultative progenitor. The

former would generate polyploid AT1 cells, which might allow AT1 cells to become larger with

excess DNA copies for environmental adaptation (18). Indeed, a recent proof-of-concept study,

which genetically induced cytokinesis failure in AT2 cells, found that binucleation does not

obviously prevent trans-differentiation towards at AT1 fate (45). These data are consistent with

our own evidence that binucleation is compatible with expression of the AT1-marker, podoplanin

(Fig. 2A and 6D). Polyploidization of AT2 cells could provide extra DNA copies for surfactant

biosynthesis, similar to how mammary epithelial cells become polyploid during lactation to

maximize milk production (23). However, the induction of AT2 polyploidy would likely

compromise its progenitor function, which requires faithful segregation of additional

chromosomes during mitosis for stem cell renewal (28). Even heart tissue, which comprises

large binucleated cardiomyocytes suited to contractility, relies on a low percentage of

mononuclear diploid progenitor cells (2-14%) to drive regeneration post-injury. Mouse strains

with a greater percentage of these mononuclear diploid cells (14%) recover from injury faster

than strains with a lower percentage (2%) (46). Given this precedence, we reason that even a

small percentage of AT2 cells becoming polyploid during an injury event (~5% per injury) could

ultimately contribute to substantial loss of AT2 progenitor renewal over a lifetime, absent

alternate mechanisms for AT2 progenitor replacement.

It will be important to understand how alveolar epithelial cells sustain the polyploid state.

Forced-induction of polyploidy in retinal pigmented epithelial cells requires activation of p53

signaling (47). Given recent evidence that forced-activation of p53 may facilitate the AT2-to-

AT1 transition during lung injury (16), it is possible that induced polyploidy plays a beneficial role

in the earliest stages of alveolar epithelial repair. However, evidence that the integrated stress

response inhibitor, ISRIB, attenuates formation or duration of hypertrophic alveolar epithelial

cells (this study) and promotes lung repair in the bleomycin model (17), raises the possibility

that maintaining polyploid cells for too long may be detrimental.

The lung needs to repair in a manner that does not massively increase cell numbers (48).

Epithelial polyploidization post-injury may be a way to repair such tissues where excessive

cellularity may be problematic. Epithelial polyploidization consequent to injury is a process with

short-term benefits (e.g., larger cells with enhanced barrier function/reduced leak, increased

DNA copies for adaptation to stressful environments), but long-term consequences (e.g.,

reduced stem cell maintenance due to loss of self-renewed mononuclear AT2). Our data raise

the possibility that modulators of the ISR may be leveraged to restore AT2 progenitor function

during injury. Future work will be required to understand the means by which ISRIB limits

formation of or guides the transition of hypertrophic, polyploid AT2 cells towards an AT1

phenotype, and degree to which induced polyploidy contributes to alveolar epithelial cell

heterogeneity or adaptation in lung disease.

298 **Materials and Methods**

300 **Key resources Table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rat anti-mouse I-A/I-E (MHC II), BUV395	BD Biosciences	Cat#743876, RRID:AB_2741827
Rat anti-mouse CD45, APC	BioLegend	Cat#102410
Rat anti-mouse CD31, APC	BioLegend	Cat#103112
Rat anti-mouse CD326 (Ep-CAM), PE/Cy7	BioLegend	Cat#118216
Rat anti-mouse CD11c, PE/Cy7	BioLegend	Cat#117318
Propidium Iodide	ThermoFisher Scientific	Cat#P1304MP
Biotin Rat Anti-mouse CD16/32	BD Biosciences	Cat#553143
Biotin Rat Anti-mouse TER119/Erythroid Cells	BD Biosciences	Cat#553672
Biotin Rat Anti-mouse CD45	BD Biosciences	Cat#553078
Biotin Rat Anti-mouse CD90	BD Biosciences	Cat#554896
Anti-Prosurfactant Protein C (proSP-C)	Millipore Sigma	Cat#AB3786
Anti-β-catenin	Millipore Sigma	Cat#06-734
Anti-β-catenin	BD Biosciences	Cat#610153
Podoplanin	DSHB (University of Iowa)	Cat#8.1.1, RRID:AB_531893
Krt8	DSHB (University of Iowa)	Cat#TROMA-I, RRID:AB_531826
Alexa Fluro 680 Phalloidin	Invitrogen	Cat#A22286
Goat anti-rabbit Alexa Fluor 488	Invitrogen	Cat#A11008
Goat anti-mouse Alexa Fluor 568	Invitrogen	Cat#A11004
Goat anti-hamster Alexa Fluor 594	Invitrogen	Cat#A21113
Goat anti-rat Alexa Fluor 546	Invitrogen	Cat#A11081
Chemicals, Peptides, and Recombinant Proteins		
Bleomycin	Fresenius Kabi	Cat#0703-3154-01
0.9% saline chloride	BBraun	Cat#R5201-01
ISRIB	AdooQ BioScience	Cat#A14302
Tamoxifen	Sigma	Cat#T5648
Corn oil	Sigma	Cat#C8267
Vybrant DyeCycle Violet Stain	Invitrogen	Cat#V35003
Hoechst 33342	ThermoFisher Scientific	Cat#62249
Draq5	ThermoFisher Scientific	Cat#62251
ROCK1 inhibitor Y-27632	Selleckhem	S1049
ProLong Gold Antifade Mountant	ThermoFisher Scientific	Cat#P36934
Eriochrome black	Sigma	Cat#E2377
CD326 (EpCAM) Microbeads, mouse	Miltenyi Biotec	Cat#130-105-958
Matrigel	Corning	Cat#356231
Collagenase	Millipore Sigma	Cat#C0130
Elastase	Worthington	Cat#LS002292

Deoxyribonuclease (DNase) I	Sigma	Cat#D4513
Dispase	Corning	Cat#47743-724
PharmLyse (Red Blood Cell Lysis)	BD Biosciences	Cat#555899
Purified Rat Anti-mouse CD16/CD32 (Mouse FcBlock)	BD Biosciences	Cat#553142
Streptavidin MagneSphere Paramagnetic Particles	Promega	Cat#Z548C
Mouse CD45 MicroBeads	Miltenyi Biotec	Cat#130-052-301
autoMACS Running Buffer – MACS Separation	Miltenyi Biotec	Cat#130-091-221
Critical Commercial Assays		
SHIELD	LifeCanvas Technologies	Cat#EI-Z1001,#SH-BS,#SH-ES,#SH-ON
MultiMACS Cell24 Separator	Miltenyi Biotec	
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J (8-12 weeks old)	The Jackson Laboratory	JAX: 000664
Mouse: $Sftpc^{CreER}$	The Jackson Laboratory	JAX: 020940
Mouse: $R26R^{ZsGreen}$	The Jackson Laboratory	JAX: 007906
Mouse: $Sftpc^{CreER}$	International Mouse Strain Resource	MGI: 5444645
Mouse: $R26R^{eYFP}$	The Jackson Laboratory	JAX: 006148
Mouse: $R26R^{Confetti}$	The Jackson Laboratory	JAX: 013731
Software and Algorithms		
FIJI/ImageJ (version: 2.1.0/1.53c)	Schneider, CA et al., 2012	https://imagej.nih.gov/ij/
Stardist plugin	Schmit, U et al., 2018	Cell/nuclear size quantification https://arxiv.org/abs/1806.03535
Other		
Vibratome	Leica Microsystems	VT1200s
Cryostat	Reichert-Jung	Cryocut 1800
W1 Spinning Disk Confocal	Nikon Instruments	
BioStation CT or other	Nikon Instruments	
Axioplan2 epifluorescence microscope	Zeiss; 20x objective (Air)	AxioCAM HR Camera with AxioVision 4.8 software
A1R Confocal	Nikon Instruments	

Methods Details

Mice: C57BL/6 mice were obtained from Jackson laboratories (Jax stock 000664). *Sftpc*^CreER (49), Ai6-R26^zsGreen, *Sftpc*^CreERT2;R26ReYFP (kindly provided by Ed Morrisey, University of Pennsylvania)(50, 51) and *Sftpc*^CreER;R26R-Confetti (8) (Jax stocks 028054, 007906, 006148, 013731). Only male mice were used in bleomycin-injury experiments based on established injury severity in males relative to females, which improves dynamic range of assays for comparisons (52). Eight to twelve week old mice were used as young adult mice. All experimental protocols were approved by the Institutional Animal Care and Use Committee at Northwestern University (Chicago, IL, USA). All strains including wildtype mice are bred and housed at a barrier- and pathogen-free facility at the Center for Comparative Medicine at Northwestern University.

Cre recombination, bleomycin administration, ISRIB administration

For nuclear activation of Cre-ER, tamoxifen was dissolved in sterile corn oil at 10 mg/100μL and administered via oral gavage once a day for two days to induce Cre-recombination of floxed alleles for lineage-tracing. For bleomycin injury, mice were anesthetized and intubated prior to the intratracheal administration of bleomycin (0.025 units in 50μL 0.9% saline). Lungs were harvested at indicated time points for downstream analyses (e.g., fixation and frozen sectioning for immunofluorescence analysis, AT2 cell isolation by negative selection (53) or EpCam+ selection (54) and flow cytometry methods (17, 55) with modifications (below). ISRIB (AdooQ BioScience, A14302) was first reconstituted in DMSO to the concentration 4mg/mL and then further diluted using PBS to a final concentration of 0.3125 mg/mL (56). ISRIB was prepared at 2.5 mg/kg and administered daily through intraperitoneal injection in a volume of 200μL. Control animals were treated with vehicle (DMSO/PBS) alone.

Tissue preparation, AT2 isolation, flow cytometry and sorting: Tissue preparation for AT2 isolation and analyses was performed as described (17, 55), with modifications. Briefly, mice were euthanized and lungs were perfused through the right ventricle with 10ml of HBSS. Lungs were removed and infiltrated with dispase and 0.2 mg/ml DNase I dissolved in HBSS with Ca^{2+} and Mg^{2+}, using syringe with 30G needle. Lungs were chopped with scissors and incubated in DMEM containing 5% FBS for 45 minutes with agitation. The tissue suspension was filtered through a 70 μM strainer and washed in 10 mL media (DMEM; 5% FBS). The pellet was then suspended in red blood cell (RBC) lysis buffer for 1 minute, washed with media, and filtered through 40 μM strainer. The cells were incubated with EpCAM magnetic beads and AT2 were

collected using the MultiMACS Cell24 Separator (Miltenyi Biotec). Automated cell counting was

performed using Nexcelom K2 Cellometer with AO/PI reagent. Cells were incubated with

FcBlock (BD Biosciences, San Jose, California), and stained with a mixture of fluorochrome-

conjugated antibodies, as listed above. DNA content was quantified using Vybrant DyeCycle

Violet. Single color controls were prepared using BD CompBeads (BD Biosciences, San Jose,

California) and Arc beads (Invitrogen, Waltham, Massachusetts). Flow cytometry and cell

sorting were performed at the Northwestern University Robert H. Lurie Comprehensive Cancer

Center Flow Cytometry Core facility (Chicago, Illinois). Data were acquired on a custom BD

FACSymphony instrument using BD FACSDiva software (BD Biosciences, San Jose,

California). Compensation and analysis were performed using FlowJo software (TreeStar). Each

cell population was identified using sequential gating strategy. The percentage of cells in the

live/singlets gate was multiplied by the number of live cells using Cellometer K2 Image

cytometer to obtain cell counts. Cell sorting was performed using BD FACSAria III instrument

using 100μm nozzle and 40 psi pressure. *Cytospin*: Following AT2 sorting, collected cells were

spun at 300g for 10 minutes. Following resuspension in PBS, cells were plated onto slides.

Slides were spun in cytospin filters at 100g for 5 minutes followed by staining with phalloidin

(1:50) and Hoescht (1:10,000).

AT2 isolation for ex vivo two-dimensional culture: Lungs were inflated intratracheally with

Dispase and incubated for 45 minutes with agitation. Lungs were chopped and incubated in

7mL of media containing 10% FBS and DNase I (1 mg/mL) for 10 minutes. Tissue suspensions

were filtered through a 70μM strainer. The cell pellets were resuspended in RBC lysis buffer for

2 minutes. Alveolar epithelial cells were enriched through negative selection against biotinylated

CD16/32, CD45, CD90, and TER119 antibodies. The cell suspensions were then incubated at

37$°$C for 2 hours followed by gentle panning. Cells were counted and cultured in complete

media (10% FBS, 2% Penicillin and Streptomycin) onto 12mm glass coverslips. Starting 48

hours after isolation, cells received a daily media change either containing 0.025μM ISRIB or

DMSO. Cell samples were pooled from 3 mice. Rat AT2 cells were collected as previously

described (57).

AT2 organoid culture: For organoid culture, cells were prepared as previously described (8)

with modifications. Mice received tamoxifen 14 days prior to isolation. AT2 isolation was

conducted as described above. For fibroblast isolation, lungs were inflated with 1 mL of

collagenase (200 U/mL), elastase (4 U/mL), and DNase solution (0.25 mg/mL). Lungs were cut

into pieces and incubated for 30 minutes at 37°C in digestion buffer. Lungs were processed in

C-tubes (Miltenyi Biotec, 130-093-237) using GentleMACS/OctoMACS dissociator (Miltenyi

Biotec). Suspensions were filtered through 40 μM strainer and rinsed with MACS buffer. The

pellet was resuspended in RBC lysis buffer, washed with MACS, and incubated with CD45

magnetic beads to deplete CD45+ cells. CD45- cell fraction was collected using the MultiMACS

Cell24 Separator (Miltenyi Biotec). Cells were sorted as described above. Following sorting,

5×10^3 zsGreen+ AT2 were mixed with 5×10^4 fibroblasts in 50% Matrigel (Corning, 356231) and

50% organoid media (αMEM, penicillin/streptomycin, insulin/transferrin/selenium, FBS, heparin,

Amphotericin B, Rock1 inhibitor, L-glutamine) in 0.4 μM Transwell inserts. Cell/Matrigel mixtures

were solidified at 37°C for 5 minutes before 300μl of media was added under the Transwell. 24

hours after isolation, organoids received media change every other day containing either 1 μM

ISRIB or DMSO for 10 days.

Tissue sectioning, immunofluorescence and image analysis: After euthanasia and

perfusion, trachea was cannulated with Luer syringe stub blunt needle and mouse lungs were

inflated with 4% paraformaldehyde at 15 cm H_2O column pressure. Lungs were fixed in 4%

paraformaldehyde (PFA) overnight at 4°C followed by cryo-embedding in optimal cutting-

temperature compound (OCT). Sections were cut at 14μM on a cryo-vibratome. For

immunostaining, sections were blocked in 10% normal goat serum (NGS), 0.3%TritonX-100 in

PBS for 30 minutes followed by an additional 1:10 anti-mouse Fab (H+L) for 2 hours. Primary

antibodies were prepared in 2% NGS 0.3%TritonX-100 PBS and applied to sections overnight

at 4°C. Sections were washed in PBS followed by secondary antibodies for 30 minutes. Lastly,

sections were incubated in Hoechst (1:10,000) for 5 minutes followed by a brief incubation of

0.3% Eriochrome black prior to mounting. Primary antibodies were used at concentrations

described below. *For 2D culture*, cells were fixed 4 days after isolation in 4% PFA. For the

alveolar organoids, organoids were fixed 10 days after isolation. Fixed cells and organoids were

neutralized in 10 mM glycine followed by blocking in 10% normal goat serum (NGS) in

0.3%TritonX-100 PBS. Primary antibodies were prepared in 3% NGS in 0.3%TritonX-100 PBS

and applied for 1 hour at room temperature. Following PBS washes, fluorophore-conjugated

secondary antibodies were prepared at 1:300 for 30 minutes in the dark at room temperature.

Hoechst was diluted 1:10,000 and applied for 5 minutes at room temperature following

secondary antibody incubation. Coverslips and filters were mounted using ProLong Gold anti-

fade solution. Images were obtained using Zeiss Axioplan epifluorescence and Nikon W1

Spinning Disk confocal microscopes. *For nuclear area quantification*, 5 fields of view were

selected for each condition per biological replicate. Images were taken on a Zeiss microscope

and analyzed in Fiji using the Stardist plugin. Primary antibodies were as follows: SFTPC

(1:100); beta-catenin (1:100); Podoplanin (1:30); Krt8 (1:30), and F-actin (1:50). *For thick*

section imaging with tissue clearing, lungs were sliced in PBS using a vibratome (Leica

VT1200s) in 200μm thick sections. Tissue slices were cleared using the LifeCanvas

Technologies SHIELD method (Cambridge, MA, USA).

Live-cell imaging: Freshly isolated alveolar epithelial cells were plated at 500K/well in a 12 well

plate. 48 hours post-isolation, media was changed and adherent cells were imaged every 15-

minutes on a Biostation-CT system (Nikon Instruments, New York, USA) for a total of 5 hours.

Image sequences were analyzed in Fiji (ImageJ v: 2.1.0/1.53c).

Statistical analysis: Data analysis and statistical tests were performed using GraphPad Prism

software (v.9.2.0).

References

1. Bharat A, Querrey M, Markov NS, Kim S, Kurihara C, Garza-Castillon R, Manerikar A, Shilatifard A, Tomic R, Politanska Y, Abdala-Valencia H, Yeldandi AV, Lomasney JW, Misharin AV, Budinger GRS. Lung transplantation for patients with severe COVID-19. *Sci Transl Med* 2020; 12.

2. Bharat A, Querrey M, Markov NS, Kim S, Kurihara C, Garza-Castillon R, Manerikar A, Shilatifard A, Tomic R, Politanska Y, Abdala-Valencia H, Yeldandi AV, Lomasney JW, Misharin AV, Budinger GRS. Lung transplantation for pulmonary fibrosis secondary to severe COVID-19. *medRxiv* 2020.

3. Spagnolo P, Balestro E, Aliberti S, Cocconcelli E, Biondini D, Casa GD, Sverzellati N, Maher TM. Pulmonary fibrosis secondary to COVID-19: a call to arms? *Lancet Respir Med* 2020; 8: 750-752.

4. Chen J, Wu H, Yu Y, Tang N. Pulmonary alveolar regeneration in adult COVID-19 patients. *Cell Res* 2020; 30: 708-710.

5. Delorey TM, Ziegler CGK, Heimberg G, Normand R, Yang Y, Segerstolpe A, Abbondanza D, Fleming SJ, Subramanian A, Montoro DT, Jagadeesh KA, Dey KK, Sen P, Slyper M, Pita-Juarez YH, Phillips D, Biermann J, Bloom-Ackermann Z, Barkas N, Ganna A, Gomez J, Melms JC, Katsyv I, Normandin E, Naderi P, Popov YV, Raju SS, Niezen S, Tsai LT, Siddle KJ, Sud M, Tran VM, Vellarikkal SK, Wang Y, Amir-Zilberstein L, Atri DS, Beechem J, Brook OR, Chen J, Divakar P, Dorceus P, Engreitz JM, Essene A, Fitzgerald DM, Fropf R, Gazal S, Gould J, Grzyb J, Harvey T, Hecht J, Hether T, Jane-Valbuena J, Leney-Greene M, Ma H, McCabe C, McLoughlin DE, Miller EM, Muus C, Niemi M, Padera R, Pan L, Pant D, Pe'er C, Pfiffner-Borges J, Pinto CJ, Plaisted J, Reeves J, Ross M, Rudy M, Rueckert EH, Siciliano M, Sturm A, Todres E, Waghray A, Warren S, Zhang S, Zollinger DR, Cosimi L, Gupta RM, Hacohen N, Hibshoosh H, Hide W, Price AL, Rajagopal J, Tata PR, Riedel S, Szabo G, Tickle TL, Ellinor PT, Hung D, Sabeti PC, Novak R, Rogers R, Ingber DE, Jiang ZG, Juric D, Babadi M, Farhi SL, Izar B, Stone JR, Vlachos IS, Solomon IH, Ashenberg O, Porter CBM, Li B, Shalek AK, Villani AC, Rozenblatt-Rosen O, Regev A. COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular targets. *Nature* 2021.

6. McDonald LT. Healing after COVID-19: are survivors at risk for pulmonary fibrosis? *American journal of physiology* 2021; 320: L257-L265.

7. Rock JR, Hogan BL. Epithelial progenitor cells in lung development, maintenance, repair, and disease. *Annual review of cell and developmental biology* 2011; 27: 493-512.

8. Barkauskas CE, Cronce MJ, Rackley CR, Bowie EJ, Keene DR, Stripp BR, Randell SH, Noble PW, Hogan BL. Type 2 alveolar cells are stem cells in adult lung. *The Journal of clinical investigation* 2013; 123: 3025-3036.

9. Aso Y, Yoneda K, Kikkawa Y. Morphologic and biochemical study of pulmonary changes induced by bleomycin in mice. *Lab Invest* 1976; 35: 558-568.

10. Miller BE, Hoo KG. Regulation of phosphatidylcholine biosynthesis in activated alveolar type II cells. *American journal of respiratory cell and molecular biology* 1989; 1: 127-136.

11. Miller BE, Hook GE. Hypertrophy and hyperplasia of alveolar type II cells in response to silica and other pulmonary toxicants. *Environ Health Perspect* 1990; 85: 15-23.

12. Kasper M, Haroske G. Alterations in the alveolar epithelium after injury leading to pulmonary fibrosis. *Histol Histopathol* 1996; 11: 463-483.

13. Wu H, Yu Y, Huang H, Hu Y, Fu S, Wang Z, Shi M, Zhao X, Yuan J, Li J, Yang X, Bin E, Wei D, Zhang H, Zhang J, Yang C, Cai T, Dai H, Chen J, Tang N. Progressive Pulmonary Fibrosis Is Caused by Elevated Mechanical Tension on Alveolar Stem Cells. *Cell* 2020; 180: 107-121 e117.

14. Jiang P, Gil de Rubio R, Hrycaj SM, Gurczynski SJ, Riemondy KA, Moore BB, Omary MB,
Ridge KM, Zemans RL. Ineffectual AEC2-to-AEC1 Differentiation in IPF: Persistence of
KRT8(hi) Transitional State. *American journal of respiratory and critical care medicine*
2020; 201: 1443-1447.
15. Strunz M, Simon LM, Ansari M, Kathiriya JJ, Angelidis I, Mayr CH, Tsidiridis G, Lange M,
Mattner LF, Yee M, Ogar P, Sengupta A, Kukhtevich I, Schneider R, Zhao Z, Voss C,
Stoeger T, Neumann JHL, Hilgendorff A, Behr J, O'Reilly M, Lehmann M, Burgstaller G,
Konigshoff M, Chapman HA, Theis FJ, Schiller HB. Alveolar regeneration through a
Krt8+ transitional stem cell state that persists in human lung fibrosis. *Nat Commun* 2020;
11: 3559.
16. Kobayashi Y, Tata A, Konkimalla A, Katsura H, Lee RF, Ou J, Banovich NE, Kropski JA,
Tata PR. Persistence of a regeneration-associated, transitional alveolar epithelial cell
state in pulmonary fibrosis. *Nature cell biology* 2020; 22: 934-946.
17. Watanabe S, Markov NS, Lu Z, Piseaux Aillon R, Soberanes S, Runyan CE, Ren Z, Grant
RA, Maciel M, Abdala-Valencia H, Politanska Y, Nam K, Sichizya L, Kihshen HG, Joshi
N, McQuattie-Pimentel AC, Gruner KA, Jain M, Sznajder JI, Morimoto RI, Reyfman PA,
Gottardi CJ, Budinger GRS, Misharin AV. Resetting proteostasis with ISRIB promotes
epithelial differentiation to attenuate pulmonary fibrosis. *Proc Natl Acad Sci U S A* 2021;
118.
18. Schoenfelder KP, Fox DT. The expanding implications of polyploidy. *J Cell Biol* 2015; 209:
485-491.
19. Amodeo AA, Skotheim JM. Cell-Size Control. *Cold Spring Harb Perspect Biol* 2016; 8:
a019083.
20. Cohen E, Allen SR, Sawyer JK, Fox DT. Fizzy-Related dictates A cell cycle switch during
organ repair and tissue growth responses in the Drosophila hindgut. *Elife* 2018; 7.
21. Cao J, Wang J, Jackman CP, Cox AH, Trembley MA, Balowski JJ, Cox BD, De Simone A,
Dickson AL, Di Talia S, Small EM, Kiehart DP, Bursac N, Poss KD. Tension Creates an
Endoreplication Wavefront that Leads Regeneration of Epicardial Tissue. *Dev Cell* 2017;
42: 600-615 e604.
22. Herrtwich L, Nanda I, Evangelou K, Nikolova T, Horn V, Sagar, Erny D, Stefanowski J,
Rogell L, Klein C, Gharun K, Follo M, Seidl M, Kremer B, Munke N, Senges J, Fliegauf
M, Aschman T, Pfeifer D, Sarrazin S, Sieweke MH, Wagner D, Dierks C, Haaf T, Ness
T, Zaiss MM, Voll RE, Deshmukh SD, Prinz M, Goldmann T, Holscher C, Hauser AE,
Lopez-Contreras AJ, Grun D, Gorgoulis V, Diefenbach A, Henneke P, Triantafyllopoulou
A. DNA Damage Signaling Instructs Polyploid Macrophage Fate in Granulomas. *Cell*
2016; 167: 1264-1280 e1218.
23. Rios AC, Fu NY, Jamieson PR, Pal B, Whitehead L, Nicholas KR, Lindeman GJ, Visvader
JE. Essential role for a novel population of binucleated mammary epithelial cells in
lactation. *Nat Commun* 2016; 7: 11400.
24. Wang J, Batourina E, Schneider K, Souza S, Swayne T, Liu C, George CD, Tate T, Dan H,
Wiessner G, Zhuravlev Y, Canman JC, Mysorekar IU, Mendelsohn CL. Polyploid
Superficial Cells that Maintain the Urothelial Barrier Are Produced via Incomplete
Cytokinesis and Endoreplication. *Cell Rep* 2018; 25: 464-477 e464.
25. Quinton RJ, DiDomizio A, Vittoria MA, Kotynkova K, Ticas CJ, Patel S, Koga Y,
Vakhshoorzadeh J, Hermance N, Kuroda TS, Parulekar N, Taylor AM, Manning AL,
Campbell JD, Ganem NJ. Whole-genome doubling confers unique genetic vulnerabilities
on tumour cells. *Nature* 2021; 590: 492-497.
26. Fox DT, Soltis DE, Soltis PS, Ashman TL, Van de Peer Y. Polyploidy: A Biological Force
From Cells to Ecosystems. *Trends Cell Biol* 2020; 30: 688-694.
27. Gjelsvik KJ, Besen-McNally R, Losick VP. Solving the Polyploid Mystery in Health and
Disease. *Trends Genet* 2019; 35: 6-14.

28. Ovrebo JI, Edgar BA. Polyploidy in tissue homeostasis and regeneration. *Development*
2018; 145.
29. Matsumoto T, Wakefield L, Tarlow BD, Grompe M. In Vivo Lineage Tracing of Polyploid
Hepatocytes Reveals Extensive Proliferation during Liver Regeneration. *Cell Stem Cell*
2020; 26: 34-47 e33.
30. Nabhan AN, Brownfield DG, Harbury PB, Krasnow MA, Desai TJ. Single-cell Wnt signaling
niches maintain stemness of alveolar type 2 cells. *Science* 2018; 359: 1118-1123.
31. Emanuelli G, Nassehzadeh-Tabriz N, Morrell NW, Marciniak SJ. The integrated stress
response in pulmonary disease. *Eur Respir Rev* 2020; 29.
32. Sudhakar A, Ramachandran A, Ghosh S, Hasnain SE, Kaufman RJ, Ramaiah KV.
Phosphorylation of serine 51 in initiation factor 2 alpha (eIF2 alpha) promotes complex
formation between eIF2 alpha(P) and eIF2B and causes inhibition in the guanine
nucleotide exchange activity of eIF2B. *Biochemistry* 2000; 39: 12929-12938.
33. Pakos-Zebrucka K, Koryga I, Mnich K, Ljujic M, Samali A, Gorman AM. The integrated
stress response. *EMBO Rep* 2016; 17: 1374-1395.
34. Costa-Mattioli M, Walter P. The integrated stress response: From mechanism to disease.
Science 2020; 368.
35. Sidrauski C, McGeachy AM, Ingolia NT, Walter P. The small molecule ISRIB reverses the
effects of eIF2alpha phosphorylation on translation and stress granule assembly. *Elife*
2015; 4.
36. Sidrauski C, Tsai JC, Kampmann M, Hearn BR, Vedantham P, Jaishankar P, Sokabe M,
Mendez AS, Newton BW, Tang EL, Verschueren E, Johnson JR, Krogan NJ, Fraser CS,
Weissman JS, Renslo AR, Walter P. Pharmacological dimerization and activation of the
exchange factor eIF2B antagonizes the integrated stress response. *Elife* 2015; 4:
e07314.
37. Zyryanova AF, Kashiwagi K, Rato C, Harding HP, Crespillo-Casado A, Perera LA,
Sakamoto A, Nishimoto M, Yonemochi M, Shirouzu M, Ito T, Ron D. ISRIB Blunts the
Integrated Stress Response by Allosterically Antagonising the Inhibitory Effect of
Phosphorylated eIF2 on eIF2B. *Mol Cell* 2021; 81: 88-103 e106.
38. Anand AA, Walter P. Structural insights into ISRIB, a memory-enhancing inhibitor of the
integrated stress response. *FEBS J* 2020; 287: 239-245.
39. Chou A, Krukowski K, Jopson T, Zhu PJ, Costa-Mattioli M, Walter P, Rosi S. Inhibition of the
integrated stress response reverses cognitive deficits after traumatic brain injury. *Proc*
Natl Acad Sci U S A 2017; 114: E6420-E6426.
40. Krukowski K, Nolan A, Frias ES, Boone M, Ureta G, Grue K, Paladini M-S, Elizarraras E,
Delgado L, Bernales S, Walter P, Rosi S. Small molecule cognitive enhancer reverses
age-related memory decline in mice. *bioRxiv* 2020: 2020.2004.2013.039677.
41. Rabouw HH, Langereis MA, Anand AA, Visser LJ, de Groot RJ, Walter P, van Kuppeveld
FJM. Small molecule ISRIB suppresses the integrated stress response within a defined
window of activation. *Proc Natl Acad Sci U S A* 2019; 116: 2097-2102.
42. Watanabe S, Markov NS, Lu Z, Aillon RP, Soberanes S, Runyan CE, Ren Z, Grant RA,
Maciel M, Abdala-Valencia H, Politanska Y, Nam K, Sichizya L, Kihshen HG, Joshi N,
McQuattie-Pimentel AC, Morimoto RI, Reyfman PA, Budinger GRS, Misharin AV.
Resetting proteostasis with ISRIB prevents pulmonary fibrosis. *bioRxiv* 2020:
2020.2002.2026.965566.
43. Uroz M, Garcia-Puig A, Tekeli I, Elosegui-Artola A, Abenza JF, Marin-Llaurado A, Pujals S,
Conte V, Albertazzi L, Roca-Cusachs P, Raya A, Trepat X. Traction forces at the
cytokinetic ring regulate cell division and polyploidy in the migrating zebrafish
epicardium. *Nat Mater* 2019; 18: 1015-1023.
44. Zielke N, Edgar BA. FUCCI sensors: powerful new tools for analysis of cell proliferation.
Wiley Interdiscip Rev Dev Biol 2015; 4: 469-487.

45. Liberti DC, Kremp MM, Liberti WA, 3rd, Penkala IJ, Li S, Zhou S, Morrisey EE. Alveolar
epithelial cell fate is maintained in a spatially restricted manner to promote lung
regeneration after acute injury. *Cell Rep* 2021; 35: 109092.
46. Patterson M, Barske L, Van Handel B, Rau CD, Gan P, Sharma A, Parikh S, Denholtz M,
Huang Y, Yamaguchi Y, Shen H, Allayee H, Crump JG, Force TI, Lien CL, Makita T,
Lusis AJ, Kumar SR, Sucov HM. Frequency of mononuclear diploid cardiomyocytes
underlies natural variation in heart regeneration. *Nat Genet* 2017; 49: 1346-1353.
47. Potapova TA, Seidel CW, Box AC, Rancati G, Li R. Transcriptome analysis of tetraploid
cells identifies cyclin D2 as a facilitator of adaptation to genome doubling in the presence
of p53. *Mol Biol Cell* 2016; 27: 3065-3084.
48. Leach JP, Morrisey EE. Repairing the lungs one breath at a time: How dedicated or
facultative are you? *Genes Dev* 2018; 32: 1461-1471.
49. Rock JR, Barkauskas CE, Cronce MJ, Xue Y, Harris JR, Liang J, Noble PW, Hogan BL.
Multiple stromal populations contribute to pulmonary fibrosis without evidence for
epithelial to mesenchymal transition. *Proc Natl Acad Sci U S A* 2011; 108: E1475-1483.
50. Zacharias WJ, Frank DB, Zepp JA, Morley MP, Alkhaleel FA, Kong J, Zhou S, Cantu E,
Morrisey EE. Regeneration of the lung alveolus by an evolutionarily conserved epithelial
progenitor. *Nature* 2018; 555: 251-255.
51. Chapman HA, Li X, Alexander JP, Brumwell A, Lorizio W, Tan K, Sonnenberg A, Wei Y, Vu
TH. Integrin alpha6beta4 identifies an adult distal lung epithelial population with
regenerative potential in mice. *The Journal of clinical investigation* 2011; 121: 2855-
2862.
52. Redente EF, Jacobsen KM, Solomon JJ, Lara AR, Faubel S, Keith RC, Henson PM,
Downey GP, Riches DW. Age and sex dimorphisms contribute to the severity of
bleomycin-induced lung injury and fibrosis. *American journal of physiology* 2011; 301:
L510-518.
53. Chen J, Chen Z, Narasaraju T, Jin N, Liu L. Isolation of highly pure alveolar epithelial type I
and type II cells from rat lungs. *Lab Invest* 2004; 84: 727-735.
54. Jansing NL, McClendon J, Kage H, Sunohara M, Alvarez JR, Borok Z, Zemans RL. Isolation
of Rat and Mouse Alveolar Type II Epithelial Cells. *Methods Mol Biol* 2018; 1809: 69-82.
55. Misharin AV, Morales-Nebreda L, Mutlu GM, Budinger GR, Perlman H. Flow cytometric
analysis of macrophages and dendritic cell subsets in the mouse lung. *American journal
of respiratory cell and molecular biology* 2013; 49: 503-510.
56. Halliday M, Radford H, Sekine Y, Moreno J, Verity N, le Quesne J, Ortori CA, Barrett DA,
Fromont C, Fischer PM, Harding HP, Ron D, Mallucci GR. Partial restoration of protein
synthesis rates by the small molecule ISRIB prevents neurodegeneration without
pancreatic toxicity. *Cell Death Dis* 2015; 6: e1672.
57. Flozak AS, Lam AP, Russell S, Jain M, Peled ON, Sheppard KA, Beri R, Mutlu GM,
Budinger GR, Gottardi CJ. Beta-catenin/T-cell factor signaling is activated during lung
injury and promotes the survival and migration of alveolar epithelial cells. *J Biol Chem*
2010; 285: 3157-3167.
612
613
614

Acknowledgements

This work relied on the following Northwestern University services and core facilities: Flow Cytometry (NCI CA060553, 1S10OD011996, 1S10OD026814); Center for Advanced Microscopy (NCI CCSG P30 CA060553, NCRR 1S10 RR031680, 1S10OD021704); BioCryo facility of Northwestern University's NUANCE Center (NSF ECCS-2025633, DMR-1720139).

Funding: MH is supported by AR073270, AM is supported by HL135124, HL153312, AG049665 and AI135964; GRSB is supported by ES13995, HL071643, AG049665; CJG is supported by HL134800, AR073270 and GM129312. All authors declare no competing financial interests.

Author contributions: AW, MMH, SW, ASF, LW and RPA designed and conducted experiments; AW, MMH, ASF, AM and CJG analyzed results; RG wrote a macro for image analysis; AM advised and analyzed flow cytometry, SH, LD, MH, AM, GRSB advised and discussed results; CJG designed and supervised study, performed analysis and wrote manuscript. MH, AM, GRSB and CJG provided funding for project.

Figures

Figure 1: **Lung injury induces AT2 cell hypertrophy.** (**A**) Schematic of experimental design.

Young adult *Sftpc*^{CreER}; Lox-Stop-Lox^{ZsGreen} mice (3-5 months) received 10mg of tamoxifen via

oral gavage 22 days prior to administration of bleomycin (0.025 units, i.t.). Lungs from saline

(control) and bleomycin-treated mice were harvested on Day 10, fixed and processed for frozen

sectioning and immunofluorescence analysis. (**B**) Representative lung sections showing native

GFP fluorescence from AT2-lineage label and immunofluorescence staining of AT1 cells by

podoplanin (Pdpn, magenta). Scale bar = 50μm. Note lineage-labeled AT2 cells appear larger

after bleomycin injury. (**C**) Graph shows quantification of AT2 cell area and perimeter using

image analysis in FIJI (workflow in Fig. S1A). Data points are from two mice, 5 fields of view per

condition. Mean +/- SD. p < 0.0001 by t-test. To ensure robustness and increase N's, injury-

induced cell size increase also validated by flow cytometry in Fig. S2.

Figure S1: **AT2 cell hypertrophy by frozen- and thick-section image analysis**. (**A**) Black

and white mask of images in Figure 1 for quantification of lineage-labelled AT2 cell area and

perimeter (black objects, green arrow). (**B**) Injury-induced AT2 hypertrophy readily observed by

frozen sections, which are ~4-fold thicker than typical paraffin sections. (**C**) Hypertrophic AT2

phenotype is not an artefact of lineage-labeling AT2 cells. Confocal image of Vibratome-cut

200μm thick lung section from a C57BL/6 mouse with clearing (SHIELD method)

immunofluorescence detection of SpC (red) and nuclei (Draq5; cyan). Note bleomycin-treated

AT2 cells are clearly larger, even 21 days post-bleomycin instillation.

Figure 2: **Lung injury induces polyploidization of lineage-labeled and SpC+ AT2 cells**. (**A**)
Schematic of experimental design as in Fig.1. Representative confocal images of lung sections
from bleomycin-treated mice; two independent fields of view from low and high-magnification
insets (boxed region) are shown (a' and a"). Native GFP fluorescence from AT2-lineage label
and immunofluorescence staining of AT1 cells by podoplanin (Pdpn, magenta). DNA labeling by
Hoechst (gray). Scale bar = 50μm and 10 μm for inset. Asterisks show suspected binucleated
AT2 cells. Arrows show plan of optical z-section, revealing that nuclei appear to touch in these
cells. Magenta arrow in a' indicates podoplanin signal in binucleated cells. SpC+ binucleated
AT2 cells are shown in Fig. S2F. (**B**) Graph shows quantification of AT2 cell binucleation. Data
points are from two mice, 5 fields of view per condition. Mean +/- SD. p < 0.0001 by t-test.
Injury-induced binucleation was also interrogated by flow cytometry in Fig. S2A-E.

Figure S2: **Quantification of injury-induced AT2 hypertrophy and binucleation by flow**

cytometry. (**A**) Schematic of experimental design. (**B**) Young adult mice (3-5 months) were

administered saline alone or 0.025 units of bleomycin and lungs were harvested and digested

for flow cytometry on Day 12 with representative flow plots gated on CD45-/Epcam+ AT2 cells

with 2N (diploid, red) or 4N DNA content (cyan) (Methods). (**C**) Quantification of AT2 cell size

(median Forward Scatter Area). (**D**) Percent 2N versus 4N AT2 cells. N = 3 mice. p = 0.0008

and 0.0039 by ANOVA with Tukey's. (**E**) Validation of bleomycin-induced AT2 cell size increase

and binucleation by cytospin analysis. Cellular F-actin stained with phalloidin (green) to

distinguish *bona fide* binucleation, with absence of F-actin-rich cell-cell junction between nuclei.

(**F**) Binucleated AT2 cells can be SpC+. Confocal z-stack image of Vibratome-cut 200μm thick

lung section from a C57BL/6 mouse with clearing (SHIELD method) immunofluorescence

detection of SpC (red) and nuclei (Draq5; cyan). Double arrowhead implicates two binucleated

AT2 cells in the same field of view.

Figure 3: **AT2 cell binucleation is not universally associated with Krt8-high expression**

phenotype. (**A**) Schematic of experimental design as in Fig.1. Confocal image of lung section

from bleomycin-treated mice; same image as Figure 2A, now including Krt8

immunofluorescence channel (cyan). Asterisk denotes binucleated AT2 cell with normal levels

of Krt8 protein. Native GFP fluorescence from AT2-lineage label and immunofluorescence

staining of AT1 cells by podoplanin (Pdpn, magenta). DNA labeling by Hoechst (gray). (**B**)

Confocal image of binucleated Krt8-high cell (asterisk) lacking AT2 lineage-label. Scale bar =

50μm and 25 μm for inset. (**C**) Confocal image of lineage-labeled binucleated AT2 cell adjacent

to lineage negative mononuclear Krt8-high cell (z-stack not shown). Scale bar = 25μm.

Figure S3: Identification of rare ≥4N AT2 cells via stochastic multi-color Confetti-labeling

system. (**A**) Schematic adapted from (29) using *Sftpc*[Cre-ER; R26R-Confetti] strain (8). (**B**) Tamoxifen

activation of *Sftpc*-targeted lineage-label at days 6-7 post-bleomycin injury and harvested Day

21. (**C**) Confocal image of thick-section cleared (SHIELD) lung, maximum z-projection of 3

slices showing rare GFP/RFP dual-positive cells, suggesting mononuclear polyploidy from

partial nuclear exclusion of signal. The enlarged/flattened size of this cell (>25μm) is also

compatible with 4N polyploidy. (**D**) Lower magnification view maximum z-projection (40 planes)

shows that most lineage labeled cells are either GFP or RFP positive, with very few GFP/RFP-

dual-positive cells (see inset showing gray cell, which reflects overlay of GFP/RFP). As this cell

is ~10μm, it likely reflects a G2-phase AT2 cell with 4N DNA content. Asterisk validates

bleomycin injury showing RFP lineage label along flattening AT1 cells.

Figure 4: **Binucleation is a feature of AT2 cell cultures and organoids.** (**A**) Confocal images

of mouse AT2 cells grown 4 days on glass and rat AT2 cells grown 10 days on filters. Cells

were fixed and stained with Hoechst (DNA in cyan or gray), phalloidin (F-actin, green), Krt8

(green) or β-catenin (red) as indicated. Dotted lines (white) outline binucleated cells. (**B**)

Schematic for activation of AT2-lineage label before AT2 isolation. (**C**) Confocal image of

lineage-labeled AT2 cell (yellow) grown 3 days on glass. Arrow indicates binucleated, AT2-

derived cell. F-actin in cyan. (**D**) Schematic for activation of AT2-lineage label before AT2

isolation for organoid growth. (**E**) AT2 cell (green), F-actin (magenta) and DNA (gray) with

lineage-labeled binucleated cell (arrow). (**F**) Rate of AT2 binucleation in organoids. Scale bars =

50μm.

Figure 5: **AT2 cell binucleate via failed cytokinesis.** Murine AT2 cells were isolated and

phase-contrast live-imaged every 15-minutes (Methods). Time stamp and evident cell-cycle

phases are shown in cyan. Arrow indicated dividing cell.

Figure 6: **ISRIB limits AT2 hypertrophy and polyploidy cell-autonomously**. (**A**) Schematic

for AT2 lineage-trace before bleomycin-injury with and without ISRIB therapy. (**B**) Confocal

images of fixed frozen lung sections with native zsGreen and immunofluorescence of AT1

marker, podoplanin. Scale 50μm. (**C**) Scatter plot of zsGreen object sizes by FIJI as described

in Figure S1. Graph identical to Fig.1C with ISRIB data added. Significance by ANOVA. (**D**)

Epifluorescence overlay images of mouse AT2 cell Day2-cultures incubated with Vehicle/ISRIB

(0.025μm) through Days 4-5. Arrows (magenta) show podoplanin+ binucleated cells that are

peripheral to large flat bi-/tri-nucleated cells with diminishing podoplanin and high levels of β-

catenin (white arrow). Scale 50μm. (**E**) Nuclear area (DNA) measurements by ImageJ as proxy

for cell size allowed for unbiased analysis. N = 6829 (Vehicle) and 7387 (ISRIB) nuclei

measured. Data averaged from 3 independent AT2 isolations for cells quantified at 48 and 72-

hours +/- ISRIB. p < 0.0001**** by Mann-Whitney test. (**F**) Histogram of data in **E**. (**G**)

Binucleation events quantified as from 44 (Vehicle) and 43 (ISRIB) fields of view. p = 0.035* by

Mann-Whitney test.

Figure 7: **Model for alveolar epithelial stress-induced polyploidization in lung injury and**

repair. (**A**) AT2 "aberrant" repair (dotted blue line) due to persistent ISR, AT2 cytokinesis failure

and stalled AT2-to-AT1 cells (right). (**B**) ISRIB therapy (dotted pink line) restores ISR to levels

for normal AT2-to-AT1 differentiation (dotted gray line).