Long-term Transverse Imaging of the Hippocampus with Glass Microperiscopes

William T Redman[1], Nora S Wolcott[2], Luca Montelisciani[3], Gabriel Luna[4], Tyler D Marks[4], Kevin K Sit[5], Che-Hang Yu[6], Spencer L Smith[4,6], and Michael J Goard[2,4,5,*]

[1]Interdepartmental Graduate Program in Dynamical Neuroscience,
 University of California, Santa Barbara,
 Santa Barbara, CA 93106, USA

[2]Department of Molecular, Cellular, and Developmental Biology,
 University of California, Santa Barbara,
 Santa Barbara, CA 93106, USA

[3]Cognitive and Systems Neuroscience Group,
University of Amsterdam,
Amsterdam, NL 1098 XH

[4]Neuroscience Research Institute,
 University of California, Santa Barbara,
 Santa Barbara, CA 93106, USA

[5]Department of Psychological and Brain Sciences,
 University of California, Santa Barbara,
 Santa Barbara, CA 93106, USA

[6]Department of Electrical and Computer Engineering,
 University of California, Santa Barbara,
 Santa Barbara, CA 93106, USA

*Correspondence to: michael.goard@lifesci.ucsb.edu

ABSTRACT

The hippocampus consists of a stereotyped neuronal circuit repeated along the septal-temporal axis. This transverse circuit contains distinct subfields with stereotyped connectivity that support crucial cognitive processes, including episodic and spatial memory. However, comprehensive measurements across the transverse hippocampal circuit *in vivo* are intractable with existing techniques. Here, we developed an approach for two-photon imaging of the transverse hippocampal plane in awake mice via implanted glass microperiscopes, allowing optical access to the major hippocampal subfields and to the dendritic arbor of pyramidal neurons. Using this approach, we tracked dendritic morphological dynamics on CA1 apical dendrites and characterized spine turnover. We then used calcium imaging to quantify the prevalence of place and speed cells across subfields. Finally, we measured the anatomical distribution of spatial information, finding a non-uniform distribution of spatial selectivity along the DG-to-CA1 axis. This approach extends the existing toolbox for structural and functional measurements of hippocampal circuitry.

INTRODUCTION

The hippocampus is critical for episodic and spatial memory formation[1–3], but the neural computations underlying these functions are not well understood. The trisynaptic circuit linking entorhinal cortex (EC) to dentate gyrus (DG), DG to CA3, and CA3 to CA1 is believed to endow the hippocampus with its functional capabilities. Since the circuit was first described in the anatomical studies of Ramon y Cajal over a century ago[4], considerable work has focused on each of the major hippocampal subfields (CA1-3 and DG) to identify their roles in hippocampal processing. The resulting body of literature has indicated that the subfields have related, but distinct roles in pattern separation and completion[5–14], response to novelty[15–19] and the encoding of social variables[20–22]. Additionally, there appear to be differences between the subfields in the stability of their place fields[19,23,24]. Although this work has increased our understanding of each of the subfields individually, it is not yet clear how neuronal activity is coordinated across the hippocampus.

This lack of knowledge comes, in part, from the technological limitations that prevent the recording of neuronal populations across hippocampal subfields in the same animal. Historically, electrophysiology has been the principal tool used to study the hippocampus. Electrophysiological recordings have the advantage of high temporal resolution and can directly measure spiking, but they are typically limited to small numbers of neurons in particular subfields. Additionally, localization of recorded neurons within the hippocampus is approximate,

requiring post-hoc histological analysis to estimate the position of the electrode tracks, with the

distances between the recorded neurons and the electrode sites being poorly defined. In recent

years, calcium imaging approaches (i.e. single photon mini-endoscopes and two-photon

microscopy) have been used to record hippocampal activity, allowing for the simultaneous

measurement of large numbers of neurons with known spatial relationships[24–29]. However, these

approaches require aspiration of the overlying neocortex and are generally restricted to a single

subfield for each animal.

Taken together, current experimental techniques are limited in their ability to: (1) record

response dynamics and coordination across the hippocampus, (2) identify and distinguish

between different neural subtypes, (3) allow for the chronic recording of cells across subfields,

and (4) resolve key cellular structures, such as apical dendrites.

To address these challenges, we have developed a procedure for transverse imaging of

the trisynaptic hippocampal circuit using chronically implanted glass microperiscopes. As has

been found with previous studies using implanted microprisms in cortex[30–32], the neural tissue

remained intact and healthy for prolonged periods of time (up to 10 months), and both dendritic

structure and calcium activity could be repeatedly measured in awake mice. Optical modeling

and point spread function measurements showed that axial resolution is decreased compared to

traditional cranial windows, but is sufficient to image individual apical dendritic spines in

hippocampal neurons several millimeters below the pial surface. Using this approach, we

quantified spine turnover in CA1 apical dendrites across days. We then measured functional

responses from CA1, CA3, and DG in head-fixed mice as they explored a floating carbon fiber

arena[33,34]. We found neurons in all regions whose activity met criteria to be considered place

cells and speed cells. Further, we found a non-uniform distribution of spatial information across

the extent of the DG-to-CA1 axis, supporting findings from earlier electrophysiological studies

that found similar heterogeneity[35–37]. Taken together, this approach adds to the existing

neurophysiological toolkit and will enable chronic structural and functional measurements

across the entire transverse hippocampal circuit.

RESULTS

Optical access to the transverse hippocampus using implanted microperiscopes

To image the transverse hippocampal circuit using two-photon (2P) imaging, we developed a

surgical procedure for chronically implanting a glass microperiscope into the septal

(dorsomedial) end of the mouse hippocampus (Fig. 1A; Fig. S1; see Methods). For imaging CA1

only, we used a 1 mm x 1 mm x 2 mm microperiscope ($v1_{CA1}$; Fig. 1B, left), and for imaging the

entire transverse hippocampus (CA1-CA3, DG), we used a 1.5 mm x 1.5 mm x 2.5 mm

microperiscope (v2$_{HPC}$; Fig. 1B, right). The microperiscope hypotenuse was coated with

enhanced aluminum in order to reflect the imaging plane orthogonally onto the transverse plane

(Fig. 1C, D). To insert the microperiscope, we made an incision through the dura and tissue

(Fig. S1A), then lowered the tip of the microperiscope into the incision (Fig. S1B), pushing the

cortical tissue medially (Fig. S1C). Although this approach eliminated the need for the aspiration

of cortical tissue typically performed prior to hippocampal imaging[25], it nonetheless results in

severed connections and compressed tissue medial to the implant. Since the septal end of the

hippocampus was affected by the implant, we used immunohistochemistry to quantify the effect

of microperiscope implantation on astrocyte proliferation, as a function of distance from the

imaging face of the microperiscope (Fig. 1E; Fig. S2). Similar to previous research using

microprism implants[31], we found an increase in the prevalence of astrocytes close to the

microperiscope face, but the density decreased with distance and was indistinguishable from

the control hemisphere beyond 300 μm from the microperiscope face (Fig. 1F).

Characterizing optical properties of microperiscope

Use of the microperiscope requires imaging through several millimeters of glass, which could

cause beam clipping or optical aberrations, potentially resulting in decreased signal intensity

and optical resolution. However, the extent to which imaging through glass prisms affects optical

signal quality is not well understood. To determine how our approach affects 2P image

properties, we modeled the expected optical properties and compared it to the experimentally-

determined signal intensity and point spread function measurements using fluorescent

microspheres (Fig. 2A). To characterize signal intensity, we measured the minimum laser power

necessary for saturation at the center of the microsphere, holding other imaging parameters

constant (Fig. 2B). We found that signal intensity imaged through the microperiscope was

reduced compared to a standard coverslip (v1$_{CA1}$ microperiscope: 59.0% $\pm$ 6.6%; v2$_{HPC}$

microperiscope: 67.2% $\pm$ 8.5%; mean $\pm$ s.d.; Fig. 2C). To measure resolution, we imaged small

diameter (0.2 μm) fluorescent microspheres and determined the point spread function (PSF),

measured as the full width at half maximum (FWHM) of the lateral and axial microsphere

profiles. Compared to a standard cranial window using a 0.15 mm coverslip. We found that the

FWHM was higher for both lateral (coverslip: 0.6 μm; v1$_{CA1}$ microperiscope: 1.0 μm; v2$_{HPC}$

microperiscope: 0.8 μm; Fig. 2D-G) and axial (coverslip: 3.4 μm; v1$_{CA1}$ microperiscope: 11.7 μm;

v2$_{HPC}$ microperiscope: 9.5 μm; Fig. 2D-G) dimensions. Optical modeling indicated that the

decrease in resolution is predominantly due to clipping of the excitation beam, resulting in a

reduction of the functional numerical aperture of the imaging system (theoretical axial PSF

FWHM of v1$_{CA1}$ microperiscope: 10.9 μm; v2$_{HPC}$ microperiscope: 7.7 μm; see Methods) rather

than optical aberrations. As a result, use of adaptive optics did not significantly improve the axial

resolution, though adaptive optics did improve the signal intensity by 40-80% (data not shown).

We next measured whether the optical properties changed across the microperiscope imaging

plane, measuring the lateral and axial microsphere FWHM across the horizontal extent of the

microperiscope (Fig. 2H). We found that the resolution was mostly uniform, with no significant

differences observed in lateral or axial resolution as a function of horizontal position (Fig. 2I;

lateral: $F_{(14, 593)} = 0.53$, $p = 0.91$; axial: $F_{(14, 593)} = 0.91$, $p = 0.55$; one-way ANOVA). Finally,

we measured resolution as a function of distance from the face of the microperiscope (Fig. 2J).

The lateral FWHM did not change as a function of distance from the microperiscope face (Fig.

2K; $F_{(19, 109)} = 0.53$, $p = 0.26$; one-way ANOVA). The axial FWHM slightly increased with

greater distances from the microperiscope face (Fig. 2K; $F_{(19, 109)} = 2.51$, $p = 0.002$; one-way

ANOVA), though none of the individual positions were significantly different when corrected for

multiple comparisons ($p > 0.05$ for all positions, Tukey-Kramer *post hoc* test).

Taken together, these results indicate that imaging through the microperiscope lowers

signal intensity and axial resolution. Despite these effects, the quality is still sufficient to image

individual HPC neurons (Fig. 1C) and sub-micron morphological structures (Fig. 3) with laser

power well below the photodamage threshold.

Imaging spines on the apical dendrites of CA1

Dendritic spines are highly dynamic and motile structures that serve as the postsynaptic sites of

excitatory synapses in the hippocampus[38,39]. Previous *in vitro* studies suggest a role for dendritic

spines in structural and functional plasticity, but the transient and dynamic nature of these

structures make them ideally suited for being studied *in vivo*[40–42]. Although existing techniques

allow imaging of spines on the basal CA1 dendrites near the surface of the hippocampus[42–44],

as well as transverse offshoots of apical dendrites[45,46], imaging hippocampal neurons along the

major somatodendritic axis has not previously been possible. Using the microperiscope, we

were able to track apical dendritic spine dynamics along the dendrite for CA1-3 neurons in

awake mice, though we focused on CA1 dendrites in this study.

In order to visualize apical dendrites in CA1 neurons, we implanted a cohort of Thy1-

GFP-M mice, sparsely expressing GFP in a subset of pyramidal neurons[47], with v1$_{CA1}$

microperiscopes (Fig. 3A; n = 7 mice). We focused on CA1 apical dendrites, but both apical and

basal dendrites could be imaged in CA1-3 neurons. Although it is theoretically possible to image

DG dendritic structures using the microperiscope, the Thy1-GFP-M mouse line has dense

expression throughout DG (Fig. 1C, D), which prevented clear identification of distinct

processes. To resolve individual spines along the dendrite, high-resolution images were taken

from several axial planes spanning the segment, and a composite image was generated using a

weighted average of individual planes (Fig. 3B; see Methods). We reduced noise by filtering and

binarizing, and isolated dendrites of interest for tracking across days (Fig. 3C, D; Fig. S3A; see

Methods).

Previous studies have shown that dendritic spines fall into four major morphological

subtypes: filopodium, thin, mushroom, and stubby[48–50]. We found that our resolution was

sufficient to classify dendritic spines into their relative subtypes and evaluate density and

turnover based on these parameters (Fig. 3E, F; Fig. S3B). Consistent with previous studies, we

found a non-uniform distribution of dendritic spines: 30.4% thin, 41.0% stubby, 26.2%

mushroom, and 2.3% filopodium ($F(3,100) = 51.47$, $p < 0.0001$, one-way ANOVA; Fig. 3E). The

low proportion of filopodium found in this and previous 2P imaging and histological studies[51,52]

(2-3%), as compared to electron microscopy studies[53,54] (~7%), may result from the narrow

width of these structures causing them to fall below the detection threshold[42,44,45,55]. Consistent

with previous work[56–58], we found that particular classes, such as filopodium, had a high

turnover rate, while other classes were more stable across sessions ($F(3,100) = 7.17$, $p < 0.001$,

one-way ANOVA; Fig. 3F).

We found that total turnover dynamics reflect 15.0% $\pm$ 2.0% spine addition and 13.0% $\pm$

1.9% spine subtraction across consecutive days, representing a high degree of instability (Fig.

3H, I). We computed the survival fraction, a measure indicating the fraction of spines still

present from day one[41,42,44]. Although daily spine addition and subtraction was approximately

15.0% (Fig. 3H, I), the survival fraction curve yields a more conservative estimate of turnover

dynamics (Fig. 3G). Across 10 days, we found a 23.5% net loss in original spines, indicating

that most spine turnover takes place within an isolated population of transient spines. Both our

cumulative turnover and survival fraction results were generally similar to previous findings from

basal dendrites in CA1[42–44], indicating that apical and basal dendrites exhibit similar spine

dynamics. It should be noted that some previous studies used super-resolution techniques to

detect smaller spines and reduce optical merging. Thus, our results may suggest an inflated

degree of stability due to resolution limitations that prevent capture of filipodia and other small

spine structures[42,44,45,55]. Despite these limitations, we found that we could identify the same

dendritic processes over long time periods (up to 150 days; Fig. S4), allowing for longitudinal

experiments tracking isolated dendritic structures over long-time intervals.

Recording place and speed cells in CA1, CA3, and DG

Much of the experimental work testing the hypothesized roles of CA1, CA3, and DG neurons

has come from place cell (PC) recordings[7–9,13,59,60]. While the results of these studies have been

instrumental, it has not been possible to measure activity throughout the transverse

hippocampal circuit in the same animals. We therefore investigated the ability of our

microperiscope to record from PCs in each of the hippocampal subfields during exploration of a

spatial environment.

To measure functional responses, we implanted v2$_{HPC}$ microperiscopes in transgenic

mice expressing GCaMP6s in glutamatergic neurons[61,62] (see Methods). As with the Thy1-GFP-

M mice, we were able to image neurons from CA1-CA3, and DG in the same animal (Fig. 4A, B;

Video S1). In some cases, depending on microperiscope placement, we were able to record

from all three areas simultaneously (Fig. S5). In all HPC subfields, we found normal calcium

dynamics with clear transients (Fig. 4C, D; Videos S2, S3). The imaging fields remained stable,

and the same field could be imaged over 100 days later (Fig. 4E). In addition, microperiscope

implantation allowed measurement of neural responses from mossy cells in the dentate gyrus

(Figs. S5, S6A) and, depending on microperiscope placement, simultaneous imaging of deep-

layer cortical neurons in parietal cortex (Fig. S6B).

As 2P microscopy generally requires the animal to be head-fixed, the behavioral assays

used to probe PC activity are limited. Previous work has made use of virtual reality (VR)[24,25],

however it remains unclear how similar rodent hippocampal activity in real world environments

is to that in VR[63,64]. To study PC activity in a physical environment, our head-fixed mice explored

a carbon fiber arena that was floated using an air table[33] (Fig. 5A; see Methods). The mice were

thus able to navigate the physical chamber by controlling their movement relative to the floor.

Although this approach lacks the vestibular information present in real world navigation, it

captures somatosensory and proprioceptive information missing from virtual

environments[34,64,65]. Moreover, recent work has found that place field width and single cell

spatial information using this approach is comparable to the responses of free foraging

animals[34]. For measurement of place coding, we allowed mice to navigate a curvilinear track

over the course of 20-40 minutes (Video S4). As found previously[34], using a curvilinear track

allowed for robust sampling of the spatial environment and improved place field localization,

though they could also be measured in the open field. In order to measure spatial properties of

the hippocampal neurons independent of reward, we relied on exploration rather than active

reward administration for sampling of the environment.

To characterize place fields, we recorded from neurons in CA1, CA3, and DG (CA1: $n = $

1001; CA3: $n = 832$; and DG: $n = 463$) in transgenic mice with panexcitatory expression of

GCaMP6s (Fig. 5B; $n = 8$ mice). We found PCs by assessing lap-by-lap reliability and

goodness-of-fit to a Gaussian in all three subfields (Fig. 5C; false positive rate in shuffled data =

0%; see Methods). The distribution was in general agreement with previous 2P imaging

experiments in mice[19,24] (Fig. 5D, G; CA1: 26.0%; CA3: 18.9%; DG: 14.0%), with fields that

spanned the entirety of the track (Fig. 5F). We found that place field widths were comparable

across the three regions (Fig. 5H; CA1: mean (median) $\pm$ s.e.m = 18.8 (19.4) $\pm$ 0.4 cm; CA3:

18.6 (18.8) $\pm$ 0.5 cm; DG: 18.9 (19.5) $\pm$ 0.8 cm) and that spatial information was highest in

CA1, followed by CA3, then DG (Fig. 5I; CA1: mean (median) $\pm$ s.e.m. = 0.95 (0.87) $\pm$ 0.03

bits/inferred spike; CA3: 0.89 (0.68) $\pm$ 0.05 bits/inferred spike; DG: 0.79 (0.61) $\pm$ 0.08

bits/inferred spike). The place field width and spatial information of the CA1 place cells were

similar to those found in a previous study using the same floating chamber design with a

traditional hippocampal window implant[34].

To determine if neurons close to the microperiscope face showed unusual response

properties, we imaged functional responses as a function of distance. Place cell width did not

vary significantly as a function of depth of the imaging plane from the face of the microperiscope

(Fig. S7A; $F(4, 241) = 1.6$, $p = 0.17$, one-way ANOVA). Spatial information did vary as a function

of depth ($F(4, 1872) = 15.9$, $p = 8.6 \times 10^{-13}$, one-way ANOVA), but there was no systematic

effect (Fig. S7B). Decay constant for the fitted transients[66] did have a significant relationship

with imaging depth ($F(4, 1872) = 29.8$, $p = 4.5 \times 10^{-24}$), which may be due to aberrant activity in a

small subset of neurons close (≤ 130 μm) to the microperiscope face (Fig. S7C).

As cells responsive to speed have recently been found in the medial EC[67,68] and

CA1[68,69], we also identified neurons as speed cells (SCs) if their activity was significantly related

to running speed[67] (see Methods; average speed per recording: CA1, 71.3 $\pm$ 33.7 mm/s; CA3,

76.2 $\pm$ 32.1; DG, 101.1 $\pm$ 49.0; Two-sample Kolmogorov-Smirnov test: CA1-CA3, $p = 0.90$;

CA3-DG, $p = 0.53$; CA1-DG, $p = 0.32$). We found SCs in CA1, CA3, and DG, with all areas

having cells that showed both increased and decreased activity with higher running speeds (Fig.

5E). Speed cells were most abundant in DG (Fig. 5G; CA1: 9.4%; CA3: 13.5%; DG: 30.2%),

consistent with recent work that found it was possible to decode the speed of freely moving

animals from the activity of DG, but not CA1[60].

Distribution of PC properties along the DG-to-CA1 axis

Recent work has suggested that PC properties are heterogeneously distributed along the extent

of the DG-to-CA1 axis. In particular, place field width and spatial information have been found to

vary among different subregions of CA3, with distal CA3 (dCA3) and CA2 having lower spatial

information and larger place fields than medial CA3 (mCA3)[35–37], and proximal CA3 (pCA3)

having values most similar to DG[13]. Such distributions could be supported by known anatomical

gradients in connectivity of CA3[70–74]. However, given that these studies required separate

animals for the recording of each location along the DG-to-CA1 axis, and that electrophysiology

has limited spatial resolution along the transverse axis, we used our microperiscopes to

measure these properties throughout the DG-to-CA1 axis in the same mice with high spatial

resolution.

Using the v2$_{HPC}$ microperiscope, we simultaneously imaged from several hundred cells

(mean $\pm$ s.d.; 215 $\pm$ 74 neurons per recording; total: 1075 neurons) extending from pCA3 to

pCA1 (Fig. 6A). Recordings from distinct imaging planes in different mice ($n = 5$) were

compared by calculating the distance of individual cells from the inflection point of the DG-to-

CA1 transverse axis (Fig. 6A; see Methods). In agreement with previous studies[35–37], we found

a non-uniform distribution of spatial information along this axis (Fig. 6B, C; $F(4, 1047) = 5.10$, p

$= 4.6 \times 10^{-4}$; General Linear F-test against a flat distribution with the same mean). In particular,

we found that pCA3 cells had spatial information that was closer in value to those in DG than

mCA3 (Fig. 6B, C; Fig. 5I), and that mCA3 had spatial information that was greater than

dCA3/CA2 (Fig. 6B, C). We found that place field widths were narrowest in mCA3, although we

failed to find a statistically significant non-uniform distribution with respect to place field width

across the extent of the DG-to-CA1 axis (Fig. S8; $F(4, 206) = 1.63$, $p = 0.17$, General Linear F-

test against a flat distribution with the same mean).

DISCUSSION

The microperiscope hippocampal imaging procedure we developed allows researchers, for the

first time, to chronically image neuronal structure and functional activity throughout the

transverse hippocampal circuit in awake, behaving mice. This approach builds on microprism

procedures developed for imaging cortical columns[30–32], allowing multiple hippocampal subfields

in the same animal to be accessed optically. Using the microperiscope, we were able to resolve

spines on the apical dendrites of CA1 pyramidal cells and track them across time. Additionally,

we were able to characterize place cells (PCs) and speed cells (SCs) in all three hippocampal

subfields and investigate their anatomical distribution across subfields.

Comparison to other methods

Over the past five decades, electrophysiology has been the principal tool used to study the hippocampus. Electrophysiological recordings have much higher temporal resolution than calcium imaging and can directly measure spikes, but have limited spatial resolution. Our approach allows large scale imaging of neurons across multiple hippocampal subfields, with known spatial and morphological relationships. In addition, our approach allows genetically-controlled labeling of particular cell types, imaging of cellular structures such as dendrites and spines, and unequivocal tracking of functional and structural properties of the same cells across time, none of which are possible with existing electrophysiological approaches.

Several approaches making use of optical imaging have been developed for use in the hippocampus. These include gradient index (GRIN) lenses[75,76], and cannulas that can be combined with both one-photon head-mounted microendoscopes[26–28] and 2P imaging[24,25,29]. These methods have been limited to horizontal imaging planes, making it difficult to image CA3 and DG, and intractable to image all three subfields in the same animal. While all of these methods cause damage to the brain, and some require the aspiration of the overlying cortex, the damage is largely restricted to superficial hippocampus. This contrasts with the implantation of our microperiscope, which is inserted into the septal end of the hippocampus and necessarily causes some damage to the structure. Despite this, we find normal response properties (Fig. 4, Fig. S7), including selectivity for location and speed, in CA1-CA3 and DG (Fig. 5). Damage in the direction orthogonal to the transverse axis caused by the implantation of the microperiscope is similar to that caused by microprisms in cortex[31], as glial markers were found to decay to baseline levels with increased distance from the face of the microperiscope (Fig. 1E, F). We showed that, following successful implantation, the imaging fields were stable, and the same cells could be imaged up to 3 months later (Fig. 4E). However, we emphasize that tissue damage caused by the microperiscope assembly to the hippocampus should be taken into consideration when planning experiments and interpreting results, and that imaging planes should be at least 150 μm away from microperiscope face to avoid aberrant responses.

We found that imaging through the microperiscope had some effect on signal intensity and axial resolution. The effect depended not only on the path length through the glass, but also on the geometry of the microperiscope. For example, the larger v2$_{HPC}$ microperiscope had slightly better optical properties than the smaller v1$_{CA1}$ microperiscope (Fig. 2), likely due to reduced beam clipping. The optical properties should be considered along with the potential tissue damage during the planning of experiments. Moreover, further customizing the

microperiscope geometry for particular applications may help optimize signal quality and
minimize damage to neural structures.
To assess the ability of our optical and behavioral experimental methods to capture
spatial coding, we chose to measure place fields in the absence of reward, which is known to
modulate hippocampal activity[77]. Recent work has found that place cells are significantly less
stable in the absence of reward[78,79], which may explain the lower lap-wise reliability in our
experiments compared to similar experiments using the same behavioral apparatus[34]. However,
we found similar percentages of place fields as other optical imaging studies, despite using a
strict criteria for place cell inclusion (see Methods), and these properties did not change
systematically as a function of distance from the face of the periscope (Fig. S7A, B). This
provides further support that the functional properties of the hippocampal circuitry remained
intact when using the microperiscope. However, we note that it is possible that the place cell
characteristics we measured might be altered in the presence of reward, and that this decision
should be taken into account when interpreting the results.
319
Structural and functional properties along the transverse hippocampal circuit
We utilized the microperiscope in two experiments that would not have been possible with
existing methods: (1) we tracked the spines along apical CA1 dendrites *in vivo* (Fig. 3); (2) we
simultaneously recorded from PCs along the extent of the DG-to-CA1 transverse axis (Fig. 6).
Several studies have tracked the spines of basal CA1 dendrites *in vivo* by imaging the
dorsal surface of the hippocampus[42–44,46,55]. However, imaging spines along the major dendritic
axis has not been accomplished *in vivo*. Given that these spines make up the majority of the
input to CA1 pyramidal neurons, there is a significant need for understanding their dynamics,
and how the dynamics differ as a function of position along the somatodendritic axis. Using the
microperiscope, we tracked isolated apical dendrites for up to 10 consecutive days (Fig. 3D).
We found considerable addition and subtraction across days, indicating dynamic turnover in
apical dendrites (Fig 3H, I). However, consistent with previous studies in basal spines[42–44], we
found the majority of spines (76.5%) survived throughout the imaging period (Fig. 3G). This high
survival fraction suggests that the daily turnover rates we observe are reflective of a distinct pool
of unstable transient spines, while the majority of spines are moderately stable, and remain
present over a longer timescale. It should be noted that some prior results were obtained using
super-resolution microscopy techniques that are more suited to capturing the dynamics of
filipodia and small spines, and thus our results may reflect an inflated degree of stability[42,44,46].
The microperiscope is in principle compatible with super-resolution microscopy, and future

imaging studies that combine the two could characterize apical dendritic spine dynamics with

improved accuracy[42,44,46]. Taken together, our results add to the increasingly appreciated idea

that, even in the absence of salient learning and reward signals, dendritic spines are dynamic

and unstable structures[42,80,81]. Understanding the nature and timescales of these dynamics has

significant implications for the reported instability of PCs[23,24,27,82–84].

Previous studies have found gradients of connectivity in CA3[70–74], suggesting that there

may be functional gradients as well. Electrophysiological recordings have indeed found that

spatial information and place field width vary as a function of distance along the DG-to-CA1

axis[35–37], and that the most proximal part of CA3 is functionally more similar to DG than to the

rest of CA3[13]. This has led to a more nuanced understanding of the hippocampal circuit, with

coarse anatomical subdivisions having finer functional subdivisions. However, given that these

previous studies relied on recordings in different animals for each location along the DG-to-CA1

axis, and given that electrodes have poor spatial resolution in the transverse axis, the results

must be cautiously interpreted. Using the microperiscope, we imaged several hundred neurons

in multiple mice along the extent of the DG-to-CA1 axis (Fig. 6A). The location of each neuron

relative to the CA3 inflection point could be easily identified, allowing for unequivocal

characterization of spatial information and place field width along the DG-to-CA1 axis across

animals. Similar to the previous studies[35–37], we found a non-uniform distribution of spatial

information (Fig. 6B, C). Place field width also appeared to vary along the DG-to-CA1 axis,

though we failed to find statistically significant non-uniformity with respect to place field width

(Fig. S8B). Indeed, we found that spatial information, but not place field width, differed

significantly between CA1, CA3, and DG (Fig. 5H, I). Given the high spatial resolution and ability

to simultaneously record from cells across the DG-to-CA1 axis using this approach, our results

strengthen the hypothesis of distributed spatial coding across, and within, hippocampal

subfields.

Future applications

This paper explored a few possible uses for the microperiscope in interrogating the hippocampal

circuit. However, there are a number of candidate applications we did not pursue that could

reveal novel insight into hippocampal function. Here, we describe several of these, chosen to

highlight the utility of the microperiscope in addressing questions that are challenging or

intractable with existing approaches: (1) The microperiscope allows for the recording from

multiple hippocampal subfields simultaneously (Fig 6; Fig. S5), allowing investigation of

interactions between neurons in different subfields during behavior. This could also be useful for

determining the effect of neuron- or subfield-specific optogenetic manipulations on downstream

subfields. (2) The microperiscope enables the investigation of local circuits by allowing

morphological and genetic identification of different cell types. This includes identifying

genetically-distinct hippocampal neurons (e.g. CA2 neurons, specific interneuron subtypes), as

well was identifying particular cell types by position or morphological characteristics (e.g. mossy

cells in DG; Fig. S6A). As these distinct cell types play important roles in hippocampal

function[20,85–89], having access to them will enable a greater understanding of the hippocampal

circuit. (3) The microperiscope can be combined with retrograde-transported viruses to allow

projection-based cell identification, making it possible to identify neurons that project to specific

downstream targets. Since hippocampal neurons that project to different brain regions have

been found to exhibit distinct functional properties (e.g. neurons in ventral CA1 that project to

the nucleus accumbens shell have been implicated in social memory[21]), the union of these tools

will be a powerful means for understanding hippocampal outputs. (4) Finally, the microperiscope

provides access to the entire dendritic tree of pyramidal neurons (Figs. 1C, 3A, 5B; Video S2),

giving optical access to dendritic signaling over a much larger spatial extent than has been

previously possible[90–92]. By sparsely expressing calcium or glutamate sensors in hippocampal

pyramidal neurons, spines throughout the somatodendritic axis could be imaged, allowing

determination of how place field responses arise from the responses of individual spines, similar

to experiments in visual cortex investigating the cellular origin of orientation tuning from synaptic

inputs[93,94].

Combined with existing electrophysiological and imaging approaches, imaging of the

transverse hippocampal circuit with microperiscopes will be a powerful tool for investigating

hippocampal circuitry, structural dynamics, and function.

A

B

v1$_{CA1}$ (CA1)

1.0 mm
2.0 mm
135°
45° 1.41 mm
imaging plane

v2$_{HPC}$ (CA1-3, DG)

1.5 mm
2.5 mm
135°
1.0 mm
45° 2.12 mm
imaging plane

C   Thy1-GFP-M

CA1

CA3

DG

100 μm

D

CA1

50 μm

CA3

DG

E

600 μm

S100β (astrocyte somata)
GFAP (astrocyte processes)

F

Atsrocyte density (S100β)

250%
150%
50%
0%
-50%

% change wrt control

0   200   400   600   800   1000
Distance from prism face (μm)

Figure 1. Implanted microperiscopes allow imaging of the hippocampal transverse plane.

(A) Three-dimensional schematic[95] illustrating microperiscope implantation and light path for hippocampal imaging.

(B) Schematics[96] showing the imaging plane location of v1$_{CA1}$ (1 mm imaging plane, 2 mm total length) and v2$_{HPC}$ (1.5 mm imaging plane, 2.5 mm total length) microperiscopes.

(C) Tiled average projection of the transverse imaging plane using the v2$_{HPC}$ microperiscope implant in a Thy1-GFP-M transgenic mouse. Scale bar = 100 μm.

(D) Enlarged images of hippocampal subfields (CA1, CA3, DG) corresponding to the rectangles in (C). Scale bar = 50 μm.

(E) Example histological section stained for astrocyte cell bodies (S100β, red) and processes (GFAP, green). Scale bar = 600 μm.

(F) Quantification of astrocyte density as a function of distance from the microperiscope face, normalized to the density in the unimplanted contralateral hemisphere (n = 2 mice; 288 and 316 days post-implant; mean $\pm$ bootstrapped s.e.m.).

Figure 2: Optical characterization of cranial window and microperiscopes.

(A) Schematic of signal yield and resolution characterization experiments to allow comparison of microperiscopes (v1$_{CA1}$ and v2$_{HPC}$) and standard coverslip window (0.15 mm thickness) using 0.2 μm fluorescent microspheres imaged with a 16x/0.8 NA objective.

(B) Example microspheres imaged through the window, v1$_{CA1}$ microperiscope, and v2$_{HPC}$ microperiscope. Laser power was gradually increased to the minimum level necessary to saturate the centroid of the microsphere with all other imaging parameters held constant.

(C) Distribution of signal intensity values (n = 12 microspheres for each condition), measured as the reciprocal of the minimum laser intensity for saturation, and normalized to the mean intensity through the window (v1$_{CA1}$: 59.0% ± 6.6%; v2$_{HPC}$: 67.2% ± 8.5%, mean ± s.d.).

(D) Average X-Y profile (left) and X-Z profile (right) of fluorescent microspheres (n = 58 microspheres) imaged through the window.

(E) Average X-Y profile (left) and X-Z profile (right) of fluorescent microspheres (n = 48 microspheres) imaged through the v2HPC microperiscope (2.5 mm path length through glass).

(F) Plot of normalized fluorescence intensity profile of X dimension (lateral resolution; FWHM = 0.6 µm) and Z dimension (axial resolution; FWHM = 3.4 μm) through the centroid of the microsphere (n = 58 microspheres) imaged through the window.

(G) Plot of normalized fluorescence intensity profile of X dimension (lateral resolution; v1$_{CA1}$ FWHM = 1.0 μm, orange; v2$_{HPC}$ FWHM = 0.8 μm, blue) and Z dimension (axial resolution; v1$_{CA1}$ FWHM = 11.7 μm; v2$_{HPC}$ FWHM = 9.5 μm) through the centroid of the microsphere (n = 28 microspheres for v1$_{CA1}$, n = 48 microspheres for v2$_{HPC}$).

(H) Schematic of the experiment used to characterize lateral and axial resolution across the horizontal axis of the microperiscope. Version v2$_{HPC}$ used for larger horizontal range.

(I) Lateral (green) and axial (magenta) FWHM (mean ± s.e.m.) of average microsphere profile as a function of horizontal distance from the center of the microperiscope. FWHM values normalized to the mean at the center of the imaging axis (-100 to +100 μm). There was no effect of horizontal position on lateral FWHM ($F(14, 593)$ = 0.53, p = 0.91) or axial FWHM ($F(14, 593)$ = 0.91, p = 0.55; one-way ANOVA).

(J) Schematic of the experiment used to measure resolution as a function of distance from the microperiscope face. Version v1$_{CA1}$ used for larger working distance.

(K) Lateral (green) and axial (magenta) FWHM (mean ± s.e.m.) of average microsphere profile as a function of distance from the face of the microperiscope, with FWHM values normalized to the closest position (25 μm). There was no effect of distance of imaging depth on lateral FWHM ($F(19, 109)$ = 0.53, p = 0.26). There was a difference for axial FWHM ($F(19, 109)$ = 2.51, p = 0.002; one-way ANOVA), though no individual position was significantly different from the first position when corrected for multiple comparisons (Tukey-Kramer *post hoc* test).

Figure 3. Chronic imaging of spine morphology in CA1 apical dendrites.

(A) Average projection of CA1 neurons sparsely expressing a GFP reporter (Thy1-GFP-M) imaged through the v1$_{CA1}$ microperiscope. Scale bar = 100 μm.

(B) Weighted projection (see Methods) of the apical dendrites shown in the dashed box of (A). Scale bar = 10 μm.

(C) Filtered and binarized image (Figure S2A; see Methods) of the dendrites in (B) to allow for the identification and classification of individual dendritic spines. Scale bar = 10 μm.

(D) Tracking CA1 dendritic spines over consecutive days on a single apical dendrite. Arrowheads indicate subtracted spines and circles indicate added spines. Colors indicate spine type of added and subtracted spines: filopodium (magenta), thin (grey), stubby (blue), and mushroom (green).

(E) Average number of spines per 10 μm section of dendrite, for each of the four classes of spine (n = 26 dendrites from 7 mice); one-way ANOVA, $F(3,100)$ = 51.47, ****p < 0.0001. Error bars indicate the interquartile range (75th percentile minus 25th percentile) and circle is median.

(F) Percent spine turnover across days in each spine type (n = 26 dendrites from 7 mice); one-way ANOVA, $F(3,100)$ = 7.17, ***p < 0.001. Error bars indicate the interquartile range (75th percentile minus 25th percentile) and circle is median.

(G) Spine survival fraction across processes (n = 26 dendrites from 7 mice) recorded over 10 consecutive days.

(H) Percent of spines added between days over 10 days of consecutive imaging.

(I) Percent of spines subtracted between days over 10 days of consecutive imaging.

A GCaMP6s

CA1

DG

CA3

100 µm

B CA1

CA3

DG

50 µm

C CA1   CA3   DG

200%⌐   100%⌐   200%⌐
DF/F   25 s   DF/F   25 s   DF/F   25 s

D median = 3.5 inf. spikes/s   2.2 inf. spikes/s   1.8 inf. spikes/s

% cells

8   16

0   15   0   15   0   15
Avg. inf. spike/s.

E CA2/3 (Day 196)   CA2/3 (Day 307)

100 µm

Figure 4. Microperiscope imaging of calcium dynamics across subfields in awake mice.

(A) Example tiled image (as in Fig. 1C) for a single panexcitatory transgenic GCaMP6s mouse (Slc17a7-GCaMP6s). Scale bar = 100 μm.

(B) Enlarged images of each hippocampal subfield (CA1, CA3, DG) corresponding to the rectangles in (A). Scale bar = 50 μm.

(C) Example GCaMP6s normalized fluorescence time courses (% DF/F) for identified cells in each subfield.

(D) Distribution of average inferred spiking rate during running for each hippocampal subfield (CA1, CA3, DG; see Methods). Median is marked by black arrowhead.

(E) Example average projection of CA2/3 imaging plane 196 days post implantation (left) and 111 days later. Image was aligned using non-rigid registration (see Methods) to account for small tissue movements. Magenta arrowheads mark example vasculature and blue arrowheads mark example neurons that are visible in both images. Scale bar = 100 μm.

A

resonant-galvo scan mirrors

2-photon collection

2-photon excitation

Ti:sapphire laser

pockels

pre-chirp

PMT

PMT

B Maximum projections

CA1

CA3

DG

50 µm

100 µm

50 µm

C

1 norm. resp.

1 norm. resp.

1 norm. resp.

Laps

0   Position (cm)   61

r² = 0.98

r² = 0.90

r² = 0.95

CA1

CA3

DG

D Place Cells

35% DF/F

11% DF/F

28% DF/F

21% DF/F

9% DF/F

18% DF/F

20% DF/F

35% DF/F

16% DF/F

9% DF/F

21% DF/F

18% DF/F

E Speed Cells

0   Position (cm)   61

7% DF/F

14% DF/F

4% DF/F

12% DF/F

2% DF/F

7% DF/F

5   Speed (mm/s)   214

5   214

5   206

5   206

5   201

14   278

F

Cross-val. CA1 resp.

Cross-val. CA3 resp.

Cross-val. DG resp.

1.0 Normalized DF/F

Place cells

0   Position (cm)   61

G

CA1 (n = 1001)

23%

6%

3%

68%

CA3 (n = 832)

16%

11%

3%

70%

DG (n = 463)

10%

26%

4%

60%

Place cells   Speed cells   Neither   Conjunctive

H

NS

NS   NS

Place field width (cm)

CA1   CA3   DG

I Place cells only

*

NS   NS

Spatial information (bits/inf. spike)

4.0

3.0

2.0

1.0

0.0

CA1   CA3   DG

Figure 5. Prevalence of place and speed cells across hippocampal subfields.

(A) Schematic of air-lifted carbon fiber circular track (250 mm outer diameter) that the mice explored during imaging (Video S4). Four sections of matched visual cues lined the inner and outer walls.

(B) Example maximum projections of GCaMP6s-expressing neurons in each subfield. Scale bar = 50, 100, 50 μm respectively.

(C) Lap-by-lap activity and Gaussian fits, with r^2 value reported, of example place cells for each hippocampal subregion to illustrate our process of identifying place cells (see Methods).

(D) Examples of mean normalized calcium response (% DF/F) versus position along the circular track for four identified place cells (PCs) in each subregion. Shaded area is s.e.m.

(E) Examples of mean calcium response (% DF/F) versus speed along the circular track for two identified speed cells (SCs) in each subregion. Shaded area is s.e.m.

(F) Cross-validated average responses (normalized % DF/F) of all PCs found in each subfield, sorted by the location of their maximum activity. Responses are plotted for even trials based on peak position determined on odd trials to avoid spurious alignment.

(G) Distribution of cells that were identified as PCs, SCs, conjunctive PC+SCs, and non-coding in each subregion.

(H) Distribution of place field width for all place cells in each subregion. Error bars indicate the interquartile range (75th percentile minus 25th percentile) and circle is median. Two-sample Kolmogorov-Smirnov test: CA1-CA3, $p = 0.67$; CA3-DG, $p = 0.70$; CA1-DG, $p = 0.51$. NS, not significant ($p > 0.05$).

(I) Distribution of spatial information (bits per inferred spike) for all place cells in each subregion. Error bars and circle are the same as in (H). Two-sample Kolmogorov-Smirnov test: CA1-CA3, $p = 0.05$; CA3-DG, $p = 0.36$; CA1-DG, $p = 0.01$. *$p < 0.05$.

Figure 6. Spatial information of neurons varies along the DG-to-CA1 axis.

(A) Maximum projection of an example DG-to-CA1 axis recording. Approximate locations of CA3 and CA1 subfields are labeled. Inflection point labeled with red line. Scale bar = 100 μm.

(B) Spatial information (bits/inferred spike), pseudo-colored on a logarithmic scale, for each neuron, overlaid on the maximum projection in (A).

(C) Spatial information, as a function of distance along the DG-to-CA1 axis (pCA3 to dCA1), calculated with a sliding window; mean $\pm$ bootstrapped s.e.m.; real data (blue), shuffled control (black). Red lines indicate values that are outside the shuffled distribution ($p < 0.05$). A general Linear F-test against a flat distribution with the same mean revealed significant non-uniformity: $F(4, 1047) = 5.10$, $p = 4.6 \times 10^{-4}$ (cells with distance greater than 600 or less than -600 were not included in the statistical analysis).

SUPPLEMENTAL FIGURES

Figure S1: Overview of microperiscope implantation surgery.

(A) Schematic view from front (top row) and top (bottom row) of mouse brain indicating placement of the incision. The incision is made using a diamond microknife prior to the implantation of microperiscope. The microknife bevel faces medial to leave a flat surface for the microperiscope face.

(B) Schematic view from front (top row) and top (bottom row) of the brain during microperiscope implantation. The microperiscope is attached to the coverglass prior to implantation using optical glue, and the assembly is lowered into the incision using a blunt 18 gauge syringe mounted on a manipulator and held using a vacuum line. Once the assembly has been cemented to the skull, the vacuum is released, and the manipulator removed. The microperiscope faces lateral for imaging of the transverse plane of the dorsomedial hippocampus.

(C) Post implantation view of microperiscope assembly, with identified hippocampal subregions.

Figure S2. Additional visualization of astrocyte immunohistochemistry.

(A) Mosaic confocal image of entire slice from mouse implanted with v2$_{HPC}$ microperiscope. Note that tissue shrinkage around implant margins increases apparent implant size. Green: GFAP (astrocyte processes); Red: S100β (astrocyte cell bodies); scale bar = 1mm.

(B) Higher zoom image from cyan box in (A). Scale bar = 500 μm.

(C) Higher zoom image from orange box in (B). Scale bar = 100 μm.

(D) Higher zoom image from cyan box in (C). Note co-staining of individual astrocytes with GFAP and S100β; S100β used for all quantification (Figure 1F) due to improved localization of cell bodies. Scale bar = 50 μm.

A

Figure S3: Dendritic morphology image processing pipeline and spine classification.

(A) Steps in dendritic morphology image processing pipeline (see Methods). Weighted mean image is obtained by acquiring average projections from 4 planes spaced by 3 μm and weighting the images around the brightest part of the dendrite. Next, the mean image is high-pass filtered to reveal fine structures. The image is then binarized using a global threshold, to filter out less prominent dendrites. Morphological opening is applied to remove any elements of the high-pass filtered image that survive the binarization process that are too small to be the principal dendrite. A semi-automated manual curation process is then used to add back any individual spines that were lost during binarization and remove any unwanted dendritic segments. Cyan arrowheads indicate manually added spines. To isolate the spines, the image is skeletonized and structures that protrude laterally from the dendritic shaft and have a total area of >1 μm^2 are identified as spines. Scale bar = 10 μm.

(B) Mean profile of all filopodium, mushroom, stubby, and thin spines (n = 47, 431, 695, and 495, respectively), computed by averaging across the registered spines of each morphological classification. Spines were classified according to spine length, neck length, neck width, head circularity, and eccentricity (see Methods).

Figure S4: Long-term imaging of the same dendrite.

(A) Example weighted average projection of a dendritic process imaged 17 days after microperiscope implantation.

(B) Same as (A), imaged 167 days after microperiscope implantation (150 days later). Arrowheads mark examples of spines that are present in locations along the dendrite at day 17, but not day 167 ("lost", magenta), present in locations at day 167, but not 17 ("gained", green), and present in locations at both days 17 and 167 ("preserved", white). Note that the dendrite was not imaged continuously, so we cannot be certain the preserved spines were there during the entire duration.

(C) Images from (A) and (B) overlaid, using rigid registration.

(D) Images from (A) and (B) overlaid, using nonrigid registration (see Methods). Scale bar = 10 μm.

Figure S5. Simultaneous imaging of all three hippocampal subfields.

Maximum projection, and example GCaMP6s fluorescence time courses for identified cells, from a recording in which CA1 (n = 55 cells), CA3 (n = 158), and DG (n = 28) were simultaneously recorded from the same image plane. Scale bar = 100 μm.

Figure S6. Additional applications for microperiscope imaging.

(A) Two maximum projections of imaging planes in which both the granular cell layer and the hilus in the DG could be imaged simultaneously. Example putative mossy cells indicated with cyan arrowheads. Scale bar = 50 μm.

(B) Maximum projection from an imaging session in which both CA1 and L6 of the neocortex are visible, allowing simultaneous imaging of neurons in CA1 and deep layers of the parietal cortex. Scale bar = 100 μm.

Figure S7. Place cell properties as a function of distance from the face of the microperiscope.

(A) Place field width as a function of imaging depth relative to the microperiscope face. Error bars indicate the interquartile range (75th percentile minus 25th percentile) and circle is median ($F_{(4, 241)}$ = 1.6, p = 0.17, one-way ANOVA).

(B) Same as (A), but for spatial information of place cells ($F_{(4, 1872)}$ = 15.9, p = 8.6x10^{-13}, one-way ANOVA; Tukey-Kramer *post hoc* test for depth vs. 30 μm: 80 μm, p = 1.1x10^{-3}; 130 μm, p = 0.34; 180 μm, p = 9.92x10^{-9}; 230 μm, p = 0.15). *p < 0.05 Tukey-Kramer *post hoc* test.

(C) Same as (A), but for decay rate[66] of place cells ($F_{(4, 1872)}$ = 29.8, p = 4.5x10^{-24}; Tukey-Kramer *post hoc* test for depth vs. 30 μm: 80 μm, p = 6.70x10^{-3}; 130 μm, p = 9.38x10^{-7}; 180 μm, p = 9.92x10^{-9}; 230 μm, p = = 9.92x10^{-9}). *p < 0.05 Tukey-Kramer *post hoc* test.

Figure S8. Place field width does not significantly vary along the DG-to-CA1 axis.

(A) Place field width, pseudo-colored for each place cell, overlaid on the maximum projection from Fig. 6A. Any cell that was not a place cell is not pseudo-colored. Scale bar = 100 μm.

(B) Place field width, as a function of distance along the DG-to-CA1 axis (pCA3 to pCA1), of real data (blue) vs. shuffled control (black). Shaded area is s.e.m. Red line indicates individual positions with place field values that were outside of the shuffled distribution ($p < 0.05$). A general Linear F-test, against a flat distribution with the same mean, revealed no significant non-uniformity: $F(4, 206) = 1.63$, $p = 0.17$ (cells with distance greater than 600 or less than -600 were not included in the statistical analysis).

SUPPLEMENTAL VIDEOS

Video S1. Demonstration of two-photon imaging of a Slc17a7-GCaMP6s mouse through the microperiscope. Recording starts in the superficial cortex in front of microperiscope, then moves through the microperiscope to the hippocampus, zooming in on subfields CA1, CA3, and DG.

Video S2. Calcium activity in subfield CA1 in a Slc17a7-GCaMP6s mouse imaged through the microperiscope.

Video S3. Calcium activity in subfield DG in a Slc17a7-GCaMP6s mouse imaged through the microperiscope.

Video S4. Mouse running in the floating chamber circular track. Ambient light is higher than usual levels for improved video quality.

METHODS

Animals

For dendritic morphology experiments, Thy1-GFP-M (Jax Stock #007788) transgenic mice ($n =$ 7) were used for sparse expression of GFP throughout the forebrain. For forebrain-wide calcium indicator expression, Emx1-IRES-Cre (Jax Stock #005628) x ROSA-LNL-tTA (Jax Stock #011008) x TITL-GCaMP6s (Jax Stock #024104) triple transgenic mice ($n = 2$) or Slc17a7-IRES2-Cre (Jax Stock #023527) x TITL2-GC6s-ICL-TTA2 (Jax Stock #031562) double transgenic mice ($n = 6$) were bred to express GCaMP6s in excitatory neurons. For imaging experiments, 8-51 week old (median 17 weeks) mice of both sexes (6 males and 9 females) were implanted with a head plate and cranial window and imaged starting 2 weeks after recovery from surgical procedures and up to 10 months after microperiscope implantation. The animals were housed on a 12 hr light/dark cycle in cages of up to 5 animals before the implants, and individually after the implants. All animal procedures were approved by the Institutional Animal Care and Use Committee at University of California, Santa Barbara.

Surgical Procedures

All surgeries were conducted under isoflurane anesthesia (3.5% induction, 1.5 - 2.5% maintenance). Prior to incision, the scalp was infiltrated with lidocaine (5 mg kg-1, subcutaneous) for analgesia and meloxicam (2 mg kg-1, subcutaneous) was administered preoperatively to reduce inflammation. Once anesthetized, the scalp overlying the dorsal skull was sanitized and removed. The periosteum was removed with a scalpel and the skull was abraded with a drill burr to improve adhesion of dental acrylic.

      For hippocampal imaging, we used two types of custom-designed glass microperiscope (Tower Optical). The first (v1$_{CA1}$), for imaging the upper part of the hippocampus (CA1/CA2) consisted of a 1 x 1 x 1 mm square base and a 1 mm right angle prism, for a total length of 2 mm on the longest side (Fig. 1B, left). The second (v2$_{HPC}$), for imaging the entire transverse circuit (CA1-3, DG) had a 1.5 x 1.5 x 1.0 mm (L x W x H) square base and a 1.5 mm right angle prism, for a total length of 2.5 mm on the longest side (Fig. 1B, right). The hypotenuse of the right angle prisms were coated with enhanced aluminum for internal reflectance. The microperiscope was attached to a 5 mm diameter coverglass (Warner Instruments) with a UV-cured optical adhesive (Norland, NOA61). Prior to implantation, the skull was soaked in sterile saline and the cortical vasculature was inspected to ensure that no major blood vessels crossed the incision site. If the cortical vasculature was suitable, a 3-4 mm craniotomy was made over the implantation site (centered at 2.2 mm posterior, 1.2-1.7 mm lateral to Bregma). For the

smaller microperiscope (v1$_{CA1}$), a 1 mm length anterior-to-posterior incision centered at -2.1 mm

posterior, 1.2 mm lateral to Bregma was then made through the dura, cortex, and mediodorsal

tip of the hippocampus to a depth of 2.2 mm from the pial surface with a sterilized diamond

micro knife (Fine Science Tools, #10100-30) mounted on a manipulator (Fig. S1A). For the

larger microperiscope (v2$_{HPC}$), two overlapping 1.0 mm length anterior-to-posterior incisions

were made centered at -1.8 mm posterior / 1.7 mm lateral and -2.4 mm posterior / 1.7 mm

lateral to Bregma to a depth of 2.7 mm, with a total anterior-to-posterior incision length of 1.6

mm. Note that placements in the regions shown in Fig. 1B required incision coordinates slightly

posterior to those indicated on the atlas. Care was taken not to sever any major cortical blood

vessels. Gelfoam (VWR) soaked in sterile saline was used to remove any blood from the

incision site. Once the incision site had no bleeding, the craniotomy was submerged in cold

sterile saline, and the microperiscope was lowered into the cortex using a manipulator, with the

imaging face of the microperiscope facing lateral (Fig. S1B). Once the microperiscope assembly

was completely lowered through the incision until the coverglass was flush with the skull, the

edges of the window were sealed with silicon elastomer (Kwik-Sil, World Precision Instruments),

then with dental acrylic (C&B-Metabond, Parkell) mixed with black ink (Fig. S1C). Care was

taken that the dental cement did not protrude over the window, as it could potentially scratch the

objective lens surface. Given the working distance of the objective used in this study (3 mm),

the smaller microperiscope (v1$_{CA1}$) implant enabled imaging from 2250 - 2600 μm below the

coverglass surface, corresponding to approximately 150 - 500 μm into the lateral hippocampus

(the 150 μm of tissue nearest to the microperiscope face was not used for imaging). The larger

microperiscope (v2$_{HPC}$) implant enabled imaging from 2650 - 2850 μm below the coverglass

surface, corresponding to approximately 150 - 350 μm into the lateral hippocampus (the 150 μm

of tissue nearest to the microperiscope face was not used for imaging). The microperiscope

implantations were stable for up to ten months following the surgery (Fig. 4E).

After microperiscope implantation, a custom-designed stainless steel head plate

(eMachineShop.com) was affixed using dental acrylic (C&B-Metabond, Parkell) mixed with

black ink. After surgery, mice were administered carprofen (5 - 10 mg kg-1, oral) every 24 hr for

3 days to reduce inflammation. Microperiscope designs and head fixation hardware are

available on our institutional lab web site (https://goard.mcdb.ucsb.edu/resources).

Point Spread Function Measurements

To measure empirical point spread functions, fluorescent microspheres (0.2 μm yellow-green

fluorescent microspheres; ThermoFisher F8811) were embedded 1:2000 in 0.5% agar and

placed under the cranial window or on the face of the microperiscope. Image stacks were taken

through the microspheres (0.07 μm per pixel in XY; 0.5 μm per plane in Z) located 50-100 μm

from the microperiscope face (except for the microsphere-microperiscope distance comparisons

in Fig. 2J-K, which ranged from 25-500 μm from the microperiscope face). Candidate

microspheres were initially isolated using the FindCircles function in the Image Segmenter app

(MATLAB image processing toolbox). Only microspheres that were >25 pixels (1.7 μm) away

from nearest neighbor microspheres and were completely contained within the Z-stack were

used for further analysis. We registered isolated microspheres at their centroids and measured

the full width at half maximum (FWHM) of the average XY and XZ profiles to determine the

lateral and axial resolution, respectively.

Since the geometry of the microperiscope limits the angle of the focusing light cone

through the microperiscope, it predominately determines the functional numerical aperture at

the imaging plane. Based on the microperiscope geometry, we calculated the effective NA of

the v1$_{CA1}$ microperiscope and v2$_{HPC}$, and used it to calculate the theoretical lateral and axial

resolution according to the following formulae[97]:

478
$$\omega_{XY} = \begin{cases} \dfrac{0.320\,\lambda}{\sqrt{2}\,NA} & NA < 0.7 \\[2mm] \dfrac{0.325\,\lambda}{\sqrt{2}\,NA^{0.91}} & NA > 0.7 \end{cases}$$

479
$$\omega_Z = \frac{0.532\,\lambda}{\sqrt{2}}\left[\frac{1}{n - \sqrt{n^2 - NA^2}}\right].$$

where ω_{XY} and ω_Z are the theoretical 1/e widths of the lateral and axial point spread function, λ

is the wavelength, and NA is the numerical aperture. Note that the FWHM was calculated by

multiplying the 1/e width (ω) by $2\sqrt{\ln(2)}$.

To perform aberration correction with adaptive optics, a deformable mirror (Multi-3.5,

Boston Micromachines Corporation) was set at a plane conjugate to the raster scanning mirrors

and the back aperture of the objective lens in the two-photon imaging system. Fluorescent

microspheres (0.2 μm) were imaged, and the standard deviation of the image brightness was

maximized under different configurations of the deformable mirror. Twelve selected Zernike

modes are applied and modulated sequentially over a total of three rounds. The 12 zernike

modes are: 1) oblique astigmatism, 2) vertical astigmatism, 3) vertical trefoil, 4) vertical coma, 5)

horizontal coma, 6) oblique trefoil, 7) oblique quadrafoil, 8) oblique secondary astigmatism, 9)

primary spherical, 10) vertical secondary astigmatism, 11) vertical quadrafoil, 12) secondary

spherical. For each Zernike mode, 21 steps of amplitudes were scanned through, and images

were acquired for each step. The amplitude that resulted in the largest standard deviation was

saved and set as the starting point of the DM configuration for the scanning of the next Zernike

mode. The brightness, the lateral resolution, and the axial resolution are compared with and

without the application of the deformable mirror correction.

Air-floated Chamber

For measurement of spatial responses, mice were head-fixed in a floating carbon fiber

chamber[33] (Mobile Homecage, NeuroTar, Ltd). The chamber base was embedded with magnets

to allow continual tracking of the position and angular displacement of the chamber. Behavioral

data was collected via the Mobile HomeCage motion tracking software (NeuroTar, versions

2.2.0.9, 2.2.014, and 2.2.1.0 beta 1). During imaging experiments, image acquisition was

triggered using a TTL pulse from the behavioral software to synchronize the timestamps from

the 2P imaging and chamber tracking.

A custom carbon fiber arena (250 mm diameter) was lined with four distinct visual

patterns (5.7 cm tall, 18.1 cm wide) printed on 7 mil waterproof paper (TerraSlate) with black

rectangles (5.7 cm tall and 1.5 cm wide) placed in between the four patterns. A circular track

(Fig. 5A; Video S4) was made by adding a removable inner circle (14 cm in diameter and 4.2

cm tall) with visual cues that were matched to the outer wall printed on 7 mil waterproof paper.

The resulting circumference, along the middle of the circular track, is 61.26 cm. Transparent

tactile stickers (Dragon Grips) were placed on the arena floors to give differential tactile stimuli

along the track. In between each recording and or behavioral session, the arena walls and floors

were thoroughly cleaned.

Mice were acclimated to the arena by the following steps: 1) On the first day the mice

were placed into the chamber and allowed to freely explore without head fixation for 15 - 20

minutes. A piece of plexiglass with holes drilled through was placed on top of the arena to keep

the mice from climbing out. 2) On the second day, the mice were head-fixed to a crossbar

extending over the floating chamber (Fig. 5A) and allowed to freely explore the floating chamber

freely for 15 minutes. Air flow (3 - 6.5 psi) was adjusted to maximize steady walking/running. On

subsequent days, the head fixation time was increased by increments of 5 minutes, as long as

the mice showed increased distance walked and percent time moving. This was continued until

the mice would explore for 30 - 40 minutes and run for greater than 15% of the time. 3) Mice

were head-fixed in the floated chamber for 20 minutes with a custom light blocker attached to

their headplate. 4) Mice were head-fixed and placed on the 2P microscope to allow habituation

to the microscope noise. 5) After mice were fully habituated, 20 - 40 minute duration recording

sessions on the 2P microscope were performed.

If at any point during the above acclimation protocol the mouse significantly decreased
distance traveled or percentage of time moving, then the mouse was moved back to the
previous step.

Custom software was written to process the behavioral data output by the Mobile
HomeCage motion tracking software. Because the Mobile HomeCage motion tracking software
sampling rate was faster than the frame rate of our 2P imaging, all behavioral variables (speed,
location, polar coordinates, and heading) that were captured within the acquisition of a single 2P
frame were grouped together and their median value was used in future analysis. For the polar
angle (which we used as the location of the mouse in 1D track), the median was computed
using an open source circular statistics toolbox (CircStat 2012a) written for MATLAB[98]. We
removed any time points when the mouse was not moving, as is standard for measurement of
place fields[25]. This helps separate processes that are related to navigation from those that are
related to resting state. To do this, we smoothed the measured instantaneous speed and kept
time periods > 1 s that had speeds greater than 20 mm/s (adding an additional 0.5 s buffer on
either side of each time period).

Two-photon Imaging

After recovery from surgery and behavioral acclimation, GFP or GCaMP6s fluorescence was
imaged using a Prairie Investigator 2P microscopy system with a resonant galvo scanning
module (Bruker). For fluorescence excitation, we used a Ti:Sapphire laser (Mai-Tai eHP,
Newport) with dispersion compensation (Deep See, Newport) tuned to λ = 920 nm. Laser power
ranged from 40-80 mW at the sample depending on GCaMP6s expression levels.
Photobleaching was minimal (<1% min^{-1}) for all laser powers used. For collection, we used
GaAsP photomultiplier tubes (H10770PA-40, Hamamatsu). A custom stainless-steel light
blocker (eMachineShop.com) was mounted to the head plate and interlocked with a tube around
the objective to prevent light from the environment from reaching the photomultiplier tubes. For
imaging, we used a 16x/0.8 NA microscope objective (Nikon) to collect 760 x 760 pixel frames
with field sizes of 829 x 829 μm or 415 x 415 μm. Images were collected at 20 Hz and stored at
10 Hz, averaging two scans for each image to reduce shot noise.

For imaging spines across days, imaging fields on a given recording session were
aligned based on the average projection from a reference session, guided by stable structural
landmarks such as specific neurons and dendrites. Physical controls were used to ensure

precise placement of the head plate, and data acquisition settings were kept consistent across

sessions. Images were collected once every day for 5 - 10 days.

Two-photon Post-processing

Images were acquired using PrairieView acquisition software (Bruker) and converted into multi-

page TIF files.

For spine imaging, registration and averaging was performed for each z-plane spanning

the axial width of the dendrite to ensure all spines were captured across z-planes. The resulting

projections were weighted according to a Gaussian distribution across planes. Non-rigid

registration was used to align dendritic segments across consecutive recording sessions. The

registered images underwent high-pass filtering to extract low amplitude spine features using

code adapted from Suite2P's enhanced mean image function[99] (Fig. S3A). The resulting ROIs

were binarized using Otsu's global threshold method for spine classification (Fig. S3A). In most

cases, the global threshold successfully isolated the single most prominent dendrite. In fields

with higher background dendrites that were not desired, these extraneous dendrites were

manually excluded. To identify spines that fall below the global threshold, the user manually

specifies incrementally lower thresholds from which to select spines that were excluded in the

initial binarization. (Fig. S3A). Spines above the global threshold with an area of $>1\ \mu m^2$ were

included in our analysis. To classify each spine as one of the four major morphological classes,

we performed the following steps. First, we found the base of the spine by identifying the region

closest to the dendritic shaft. Second, we calculated the length of the spine by taking the

Euclidean distance between the midpoint of the spine base and the most distant pixel. Third,

this vector was divided evenly into three segments to find the spine head, neck, and base areas

respectively. Finally, spines were classified in the four categories, considering the following

threshold parameters (Fig. S3B): stubby (neck length $< 0.2\ \mu m$ and aspect ratio < 1.3), thin

(neck length $> 0.2\ \mu m$, spine length $< 0.7\ \mu m$, head circularity $< 0.8\ \mu m$), mushroom (neck

length $> 0.2\ \mu m$, head circularity $> 0.8\ \mu m$), and filopodium (neck length $> 0.2\ \mu m$, spine length

$< 0.8\ \mu m$, aspect ratio >1.3).

For calcium imaging sessions, the TIF files were processed using the Python

implementation of Suite2P[99]. We briefly summarize their pipeline here. First, TIFs in the image

stack undergo rigid registration using regularized phase correlations. This involves spatial

whitening and then cross-correlating frames. Next, regions of interest (ROIs) are extracted by

clustering correlated pixels, where a low-dimensional decomposition is used to reduce the size

of the data. The number of ROIs is set automatically from a threshold set on the pixel

correlations. We manually checked assigned ROIs based on location, morphology and DF/F

traces.

Since the hippocampal pyramidal cells are densely packed and the microperiscope

reduces the axial resolution, we perform local neuropil subtraction using custom code

(https://github.com/ucsb-goard-lab/two-photon-calcium-post-processing) to avoid neuropil

contamination. The corrected fluorescence was estimated according to

$$F_{corrected}\,(n)\;=\;F_{soma}\,(n)\;-\;\alpha\,(F_{neuropil}(n)\;-\;\bar{F}_{neuropil}),$$

where $F_{neuropil}$ was defined as the fluorescence in the region $<30~\mu m$ from the ROI border

(excluding other ROIs) for frame n. $\bar{F}_{neuropil}$ is $F_{neuropil}$ averaged over all frames. α was

chosen from [0, 1] to minimize the Pearson's correlation coefficient between $F_{corrected}$ and $F_{neuropil}$.

The ΔF/F for each neuron was then calculated as

$$\frac{\Delta F}{F}\;=\;\frac{F_n - F_0}{F_0},$$

where F_n is the corrected fluorescence ($F_{corrected}$) for frame n and F_0 is defined as the first mode

of the corrected fluorescence density distribution across the entire time series.

We deconvolved this neuropil subtracted ΔF/F to obtain an estimate for the

instantaneous spike rate, which we used (only) for the computation of neurons' spatial

information (see below). This inferred spike rate was obtained via a MATLAB implementation of

a sparse, nonnegative deconvolution algorithm (OASIS) used for Ca^{2+} recordings[66]. We used an

auto-regressive model of order 2 for the convolution kernel. Any cells that had spike rate > 10

spikes/sec. or <1 spikes/sec. were manually checked and were removed from consideration if

their traces appeared too noisy or sparse. Such cells were not considered in future analysis and

were not included in the total number of cells recorded from.

Spine Imaging Data Analysis

After nonrigid registration, high pass filtering, and binarization of the dendritic segment,

individual spines were extracted based on standard morphological criteria[100]. Spines projecting

laterally from the dendritic segment were extracted and analyzed as individual objects, as

described previously (Fig. S3). The sum of the members of each spine class, as well as the total

number of all spines, was recorded for each session. Spine totals (S_{total}) were then broken down

into 10 μm sections of the dendritic segment ($S_{section}$) using the following calculation:

621
$$S_{section}(n) = \frac{S_{total}(n)}{\left(D_{length} \times \frac{F_{\mu m}}{F_{pixels}}\right)} \times 10,$$

622
where length of the dendritic segment, D_{length}, was determined by skeletonizing the dendritic

shaft to 1 pixel in diameter, then taking the area of the pixels. F_{pixels} is the FOV in pixels, which

here was 760 x 760 at 16x magnification, and $F_{\mu m}$ is the FOV in microns, which was 52 x 52 μm.

Turnover was estimated at 24 h increments; turnover here is defined as the net change

in spines per day for each morphological class (Fig. 3F). To determine which specific spines

were involved in turnover across days, segments recorded 24 h apart were aligned and overlaid

using a custom MATLAB interface, which allowed the user to manually select new or removed

spines. Percent addition/subtraction $S_{a/s}$ was calculated as:

631
$$S_{a/s} = \frac{N_{a/s}(t)}{N(t)} \times 100,$$

632
where $N_{a/s}(t)$ is spines that have been added or subtracted and $N(t)$ is the total average number

of spines. To account for variance in spine classification across days, turnover of specific

classes of spines was normalized to total cumulative turnover per day.

To calculate the survival fraction curve $S(t)$, we determined which spines were present at

time t_n that were not present at time t_0[42,44]. The dendritic segment from t_0 was transparently

overlaid with segments from t_n, and replacement spines that were present in t_0 but not t_n were

manually identified. Survival fraction was quantified as

640
$$S(t) = \frac{N_r(t_n)}{N(t_0)} \times 100,$$

641
where $N_r(t_n)$ are the total spines at t_n that were also found in t_0, and $N(t_0)$ are the total number of

spines that were present in t_0. Survival fraction, as well as % addition and subtraction, was

calculated in 10 μm sections to control for segment length.

Dendrite registration (Fig. S4) was performed using built-in registration functions in the

MATLAB image processing toolbox (rigid registration: imregister, non-rigid registration:

imregdemons for displacement field estimation and warping).

Calcium Imaging Data Analysis

For calcium imaging experiments during exploration of the air-floated chamber, processed and

synchronized behavioral data and 2P imaging data were used to identify place and speed cells

as follows.

First, the 1D track was divided into 72 equal bins (each ~0.85 cm in length). Activity as a

function of position (we refer to these as spatial tuning curves) was computed for each lap, with

activity divided by occupancy of each binned location. We observed that in certain cases, the

mice traversed the track at high speeds. To avoid misattribution of slow calcium signals to

spatial bins (which were relatively small due to our small track), any lap where the average

instantaneous speed was greater than 180 mm/s was removed and not considered for further

analysis (an average of 7% of laps were removed). To assess the consistency of spatial coding

of each cell, we randomly split the laps into two groups and computed the correlation coefficient

between the averaged spatial tuning curves. We then did the same for shuffled data in which

each lap's spatial tuning curve was circularly permuted by a random number of bins. Note that

this was done for each lap, to avoid trivial effects that might emerge from circularly permuting

data that was stereotyped along the track. This was performed 500 times, and the distribution of

actual correlation coefficient values was compared to the distribution of circularly shuffled values

using a two-sample Kolmogorov-Smirnov test (α = 0.01). The distribution also had to pass a

Cohen's D analysis, having a score of greater than 0.5. A cell that passed these tests was

considered a "consistent" cell.

To identify a neuron as a place cell, the neuron had to pass the consistency test, in

addition to being well fit by a Gaussian function, $R_{DF/F} = A_0 + A\,e^{\left(\frac{(X-B)}{C}\right)^2}$, with FWHM = $2C\sqrt{\ln 2}$.

Note that in this convention, $C^2 = 2\,\sigma^2$. Specifically, we required that: 1) the adjusted R^2 >

0.375; 2) 2.5cm < FWHM < 30.6 cm (50% of track length – results in Fig. 5H did not significantly

depend on this threshold); 3) A > 0; 4) A/A_0 > 0.50. Cells that met these conditions were

characterized as place cells; with place fields at the location of maximal activity and width

defined as the FWHM. Note that these criteria are somewhat strict compared to traditional place

field criteria. When tested with data in which individual laps were time shuffled, the approach

yielded a false positive rate of 0%.

Speed cells were identified using a standard process developed for identification of

speed cells in medial entorhinal cortex and hippocampus[67–69]. We computed the Pearson

correlation of each cells' DF/F trace with the mouse's speed across the experiment. This value

is considered as a "speed score". We then circularly shuffled the DF/F 100 times (making sure

that the amount shuffled was greater than 10 frames to ensure that the shuffled distribution did

not have artificially high correlations). Cells whose speed score was greater than 99%, or less

than 1%, of the shuffled distribution were considered speed cells.

684   To compute the spatial information[101] of cell j (SI$_j$), we used the following formula,

685
$$SI_j = \frac{1}{\bar{a}_j} \sum_{k=1}^{72} p(k) a_j(k) \log_2 \left[\frac{a_j(k)}{\bar{a}_j} \right]$$

where $\bar{a}_j$ is the mean inferred spike rate of cell j, $a_j(k)$ is the mean inferred spike rate of

cell j at position bin k, and $p(k)$ is the probability of being at position bin k. We divide by $\bar{a}_j$ to

have SI in units of bits/inferred spike. To align recordings where we recorded along the CA1-DG

axis, we found the inflection point of the axis and then computed the distance of each cell to that

point. To do this, we performed the following steps. 1) We extracted the position of each

identified cell using Suite2P's centroid output. 2) We then fit a function of the form $a(x - b)^2 + c$

to the cell positions by rotating the field-of-view from 0 to 180 degrees and finding the rotation

that maximized the R^2 value of the fit. 3) We determined the inflection point as the peak of the

curve and de-rotated the fit to determine the inflection point and curve in the original

coordinates. The distance of each cell to the inflection point was found by finding the point along

the fit curve that had the minimal distance to the cell's centroid.

Immunohistochemistry

Samples were perfusion fixed using 4% paraformaldehyde in 0.1M sodium cacodylate buffer

(pH = 7.4) for 10 mins, and then immersion fixed overnight at 4ºC. Next, sections were rinsed in

cold PBS 5 x 5 mins and 1 x 1hr. Whole brains were then embedded in 10% low-melting

agarose. Subsequently, 100 μm coronal sections were cut using a vibratome (Leica, Lumberton,

NJ). Sections were then blocked overnight in normal donkey serum (Jackson ImmunoResearch;

West Grove, PA) diluted 1:20 in PBS containing 0.5% bovine serum albumin, 0.1% Triton-X

100, and 0.1% sodium azide, hereafter, PBTA at 4ºC. Next, primary antibodies anti-GFAP

(1:500; abcam; ab53554), anti-S100β (1:1000; DAKO; Z0311) were diluted in PBTA and

incubated overnight at 4ºC. Then, sections were rinsed 5 x 5 mins and 1 x 1hr before

corresponding secondary antibodies along with the nuclear stain Hoechst33342 (1:5000;

Molecular Probes; H-3570) were incubated overnight at 4ºC. Lastly, secondary antibodies were

rinsed, and sections mounted using Vectashield (Vector laboratories Inc; H-1200) and sealed

under #0 coverslips.

High resolution wide-field mosaics of brain samples were then imaged with a 20x oil
immersion lens and an Olympus Fluoview 1000 laser scanning confocal microscope (Center
Valley, PA) at a pixel array of 800 x 800 and then registered using the bio-image software
Imago (Santa Barbara CA).

We then calculated glial cell density as a function of distance from the microperiscope
face. First, each mosaic was rotated so that the medial-lateral axis of the brain sample was
aligned to be parallel with the horizontal axis of the image. Then each mosaic was cropped to
remove extraneous pixels outside of the imaged brain slice. Next, a line denoting the face of the
microperiscope was manually drawn parallel to the dorsal-ventral axis aligned with the location
of the microperiscope face. We then used a custom cell-counting algorithm that identified
potential regions of interest (ROIs). We limited the ROIs to be within the hippocampal formation
in the brain slices. The Euclidean distance between the closest point on the defined
microperiscope face and each ROI's centroid was calculated. Afterward, a similar procedure
was performed on the contralateral side of the brain slice, with a mock "microperiscope face"
defined at symmetric coordinates to the true microperiscope face, to serve as a control. These
steps were repeated for each channel of the mosaic.

After extracting each ROIs distance from the microperiscope face, we counted the
number of cells in each 50 µm distance bin. To account for basal glial cell density, we calculated
the percent change of glial cell density on the microperiscope side with respect to the control
side. This procedure was repeated 1000 times using randomly sampled distances, with
replacement, to bootstrap the sample variance.

Statistical Information

Violin plots were made using an open-source MATLAB package[102]. Statistical tests for spine
morphological types were calculated using a one-way ANOVA. Reliability across laps was
tested with a two-sample Kolmogorov-Smirnov test. Comparisons between model fits for spatial
distribution of spatial information and place field width used a General Linear F-test.

DATA AVAILABILITY

Microperiscope designs can be found on our institutional lab website
(https://goard.mcdb.ucsb.edu/resources). Microsphere data from Fig. 2, spine imaging data from
Fig. 3, and neuronal response data from Figs. 5-6 are available on Dryad (DOI: TBA, final
version will be uploaded upon manuscript acceptance). Code for point spread function

calculations, spine analysis, and place/speed cell identification is available on Github (DOI: TBA, final version will be uploaded upon manuscript acceptance).

REFERENCES

1. O'Keefe, J. & Nadel, L. *The Hippocampus as a Cognitive Map*. (Oxford University Press,
1978).

2. Squire, L. R. Memory and the Hippocampuss: A Synthesis From Findings With Rats,
Monkeys, and Humans. *Psychological Review* **99**, 195–231 (1992).

3. Tonegawa, S., Liu, X., Ramirez, S. & Redondo, R. Memory Engram Cells Have Come of
Age. *Neuron* vol. 87 918–931 (2015).

4. Ramon y Cajal, S. *Histologie du système nerveux de l'homme et des vertèbres*. (Maloine,
1911).

5. Gilbert, P. E., Kesner, R. P. & Lee, I. Dissociating hippocampal subregions: A double
dissociation between dentate gyrus and CA1. *Hippocampus* **11**, 626–636 (2001).

6. Lee, I., Yoganarasimha, D., Rao, G. & Knierim, J. J. Comparison of population coherence
of place cells in hippocampal subfields CA1 and CA3. *Nature* **430**, 456–459 (2004).

7. Leutgeb, S., Leutgeb, J. K., Treves, A., Moser, M. B. & Moser, E. I. Distinct ensemble
codes in hippocampal areas CA3 and CA1. *Science (1979)* **305**, 1295–1298 (2004).

8. Leutgeb, J. K. *et al.* Progressive transformation of hippocampal neuronal representations
in "morphed" environments. *Neuron* **48**, 345–358 (2005).

9. Leutgeb, J. K., Leutgeb, S., Moser, M. B. & Moser, E. I. Pattern separation in the dentate
gyrus and CA3 of the hippocampus. *Science (1979)* **315**, 961–966 (2007).

10. McHugh, T. J. *et al.* Dentate gyrus NMDA receptors mediate rapid pattern separation in
the hippocampal network. *Science (1979)* **317**, 94–99 (2007).

11. Nakashiba, T. *et al.* Young dentate granule cells mediate pattern separation, whereas old
granule cells facilitate pattern completion. *Cell* **149**, 188–201 (2012).

12. Kheirbek, M. A. *et al.* Differential control of learning and anxiety along the dorsoventral
axis of the dentate gyrus. *Neuron* **77**, 955–968 (2013).

13. Neunuebel, J. P. & Knierim, J. J. CA3 retrieves coherent representations from degraded
input: Direct evidence for CA3 pattern completion and dentate gyrus pattern separation.
Neuron **81**, 416–427 (2014).

14. Rennó-Costa, C., Lisman, J. E. & Verschure, P. F. M. J. A Signature of Attractor
Dynamics in the CA3 Region of the Hippocampus. *PLoS Computational Biology* **10**,
e1003641 (2014).

15. Frank, L. M., Stanley, G. B. & Brown, E. N. Hippocampal plasticity across multiple days of
exposure to novel environments. *Journal of Neuroscience* **24**, 7681–7689 (2004).

16. Karlsson, M. P. & Frank, L. M. Network dynamics underlying the formation of sparse,
informative representations in the hippocampus. *Journal of Neuroscience* **28**, 14271–
14281 (2008).

17.   Kemere, C., Carr, M. F., Karlsson, M. P. & Frank, L. M. Rapid and continuous modulation
of hippocampal network state during exploration of new places. *PLoS One* **8**, e73114
(2013).

18.   Larkin, M. C., Lykken, C., Tye, L. D., Wickelgren, J. G. & Frank, L. M. Hippocampal
output area CA1 broadcasts a generalized novelty signal during an object-place
recognition task. *Hippocampus* **24**, 773–783 (2014).

19.   Dong, C., Madar, A. D. & Sheffield, M. E. J. Distinct place cell dynamics in CA1 and CA3
encode experience in new environments. *Nature Communications* **12**, 1–13 (2021).

20.   Hitti, F. L. & Siegelbaum, S. A. The hippocampal CA2 region is essential for social
memory. *Nature* **508**, 88–92 (2014).

21.   Okuyama, T., Kitamura, T., Roy, D. S., Itohara, S. & Tonegawa, S. Ventral CA1 neurons
store social memory. *Science (1979)* **353**, 1536–1541 (2016).

22.   Meira, T. *et al.* A hippocampal circuit linking dorsal CA2 to ventral CA1 critical for social
memory dynamics. *Nature Communications* **9**, 1–14 (2018).

23.   Mankin, E. A. *et al.* Neuronal code for extended time in the hippocampus. *Proc Natl Acad*
Sci U S A **109**, 19462–19467 (2012).

24.   Hainmueller, T. & Bartos, M. Parallel emergence of stable and dynamic memory engrams
in the hippocampus. *Nature* **558**, 292–296 (2018).

25.   Dombeck, D. A., Harvey, C. D., Tian, L., Looger, L. L. & Tank, D. W. Functional imaging
of hippocampal place cells at cellular resolution during virtual navigation. *Nature*
Neuroscience **13**, 1433–1440 (2010).

26.   Ghosh, K. K. *et al.* Miniaturized integration of a fluorescence microscope. *Nature*
Methods **8**, 871–878 (2011).

27.   Ziv, Y. *et al.* Long-term dynamics of CA1 hippocampal place codes. *Nature Neuroscience*
16, 264–266 (2013).

28.   Cai, D. J. *et al.* A shared neural ensemble links distinct contextual memories encoded
close in time. *Nature* **534**, 115–118 (2016).

29.   Sheintuch, L. *et al.* Tracking the Same Neurons across Multiple Days in Ca2+ Imaging
Data. *Cell Reports* **21**, 1102–1115 (2017).

30.   Chia, T. H. & Levene, M. J. Microprisms for in vivo multilayer cortical imaging. *Journal of*
Neurophysiology **102**, 1310–1314 (2009).

31.   Andermann, M. L. *et al.* Chronic Cellular Imaging of Entire Cortical Columns in Awake
Mice Using Microprisms. *Neuron* **80**, 900–913 (2013).

32.   Low, R. J., Gu, Y. & Tank, D. W. Cellular resolution optical access to brain regions in
fissures: Imaging medial prefrontal cortex and grid cells in entorhinal cortex. *Proc Natl*
Acad Sci U S A **111**, 18739–18744 (2014).

33.   Kislin, M. *et al.* Flat-floored air-lifted platform: A new method for combining behavior with
microscopy or electrophysiology on awake freely moving rodents. *Journal of Visualized*
Experiments (2014) doi:10.3791/51869.

34.   Go, M. A. *et al.* Place Cells in Head-Fixed Mice Navigating a Floating Real-World
Environment. *Frontiers in Cellular Neuroscience* **15**, 618658 (2021).

35.   Lee, H., Wang, C., Deshmukh, S. S. & Knierim, J. J. Neural Population Evidence of
Functional Heterogeneity along the CA3 Transverse Axis: Pattern Completion versus
Pattern Separation. *Neuron* **87**, 1093–1105 (2015).

36.   Lu, L., Igarashi, K. M., Witter, M. P., Moser, E. I. & Moser, M. B. Topography of Place
Maps along the CA3-to-CA2 Axis of the Hippocampus. *Neuron* **87**, 1078–1092 (2015).

37.   Mankin, E. A., Diehl, G. W., Sparks, F. T., Leutgeb, S. & Leutgeb, J. K. Hippocampal CA2
Activity Patterns Change over Time to a Larger Extent than between Spatial Contexts.
Neuron **85**, 190–201 (2015).

38.   Engert, F. & Bonhoeffer, T. Dendritic spine changes associated with hippocampal long-
term synaptic plasticity. *Nature* **399**, 66–70 (1999).

39.   Harris, K. M. Structure, development, and plasticity of dendritic spines. *Current Opinion in*
Neurobiology **9**, 343–348 (1999).

40.   Yuste, R. & Bonhoeffer, T. Morphological changes in dendritic spines associated with
long-term synaptic plasticity. *Annual Review of Neuroscience* vol. 24 1071–1089 (2001).

41.   Yang, G., Pan, F. & Gan, W. B. Stably maintained dendritic spines are associated with
lifelong memories. *Nature* **462**, 920–924 (2009).

42.   Attardo, A., Fitzgerald, J. E. & Schnitzer, M. J. Impermanence of dendritic spines in live
adult CA1 hippocampus. *Nature* **523**, 592–596 (2015).

43.   Mizrahi, A., Crowley, J. C., Shtoyerman, E. & Katz, L. C. High-Resolution In Vivo Imaging
of Hippocampal Dendrites and Spines. *Journal of Neuroscience* **24**, 3147–3151 (2004).

44.   Pfeiffer, T. *et al.* Chronic 2P-STED imaging reveals high turnover of dendritic spines in
the hippocampus in vivo. *Elife* **7**, (2018).

45.   Gu, L. *et al.* Long-Term In Vivo Imaging of Dendritic Spines in the Hippocampus Reveals
Structural Plasticity. *Journal of Neuroscience* **34**, 13948–13953 (2014).

46.   Ulivi, A. F. *et al.* Longitudinal Two-Photon Imaging of Dorsal Hippocampal CA1 in Live
Mice. *JoVE (Journal of Visualized Experiments)* **2019**, e59598 (2019).

47.   Feng, G. *et al.* Imaging neuronal subsets in transgenic mice expressing multiple spectral
variants of GFP. *Neuron* **28**, 41–51 (2000).

48.   Chang, F. L. F. & Greenough, W. T. Transient and enduring morphological correlates of
synaptic activity and efficacy change in the rat hippocampal slice. *Brain Research* **309**,
35–46 (1984).

49. Rodriguez, A., Ehlenberger, D. B., Dickstein, D. L., Hof, P. R. & Wearne, S. L. Automated
three-dimensional detection and shape classification of dendritic spines from
fluorescence microscopy images. *PLoS ONE* **3**, 1997 (2008).

50. Son, J., Song, S., Lee, S., Chang, S. & Kim, M. Morphological change tracking of
dendritic spines based on structural features. *Journal of Microscopy* **241**, 261–272
(2011).

51. Lendvai, B., Stern, E. A., Chen, B. & Svoboda, K. Experience-dependent plasticity of
dendritic spines in the developing rat barrel cortex in vivo. *Nature* **404**, 876–881 (2000).

52. Risher, W. C., Ustunkaya, T., Alvarado, J. S. & Eroglu, C. Rapid golgi analysis method for
efficient and unbiased classification of dendritic spines. *PLoS ONE* **9**, e107591 (2014).

53. Fiala, J. C., Feinberg, M., Popov, V. & Harris, K. M. Synaptogenesis via dendritic filopodia
in developing hippocampal area CA1. *Journal of Neuroscience* **18**, 8900–8911 (1998).

54. Stewart, M. G. *et al.* Chemically induced long-term potentiation increases the number of
perforated and complex postsynaptic densities but does not alter dendritic spine volume
in CA1 of adult mouse hippocampal slices. *European Journal of Neuroscience* **21**, 3368–
3378 (2005).

55. Castello-Waldow, T. P. *et al.* Hippocampal neurons with stable excitatory connectivity
become part of neuronal representations. *PLOS Biology* **18**, e3000928 (2020).

56. Grutzendler, J., Kasthuri, N. & Gan, W. B. Long-term dendritic spine stability in the adult
cortex. *Nature* **420**, 812–816 (2002).

57. Trachtenberg, J. T. *et al.* Long-term in vivo imaging of experience-dependent synaptic
plasticity in adult cortex. *Nature* **420**, 788–794 (2002).

58. Holtmaat, A. J. G. D. *et al.* Transient and persistent dendritic spines in the neocortex in
vivo. *Neuron* **45**, 279–291 (2005).

59. van Dijk, M. T. & Fenton, A. A. On How the Dentate Gyrus Contributes to Memory
Discrimination. *Neuron* **98**, 832-845.e5 (2018).

60. Stefanini, F. *et al.* A Distributed Neural Code in the Dentate Gyrus and in CA1. *Neuron*
107, 703-716.e4 (2020).

61. Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*
499, 295–300 (2013).

62. Daigle, T. L. *et al.* A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced
Brain-Cell-Type Targeting and Functionality. *Cell* **174**, 465-480.e22 (2018).

63. Aronov, D. & Tank, D. W. Engagement of Neural Circuits Underlying 2D Spatial
Navigation in a Rodent Virtual Reality System. *Neuron* **84**, 442–456 (2014).

64. Aghajan, Z. M. *et al.* Impaired spatial selectivity and intact phase precession in two-
dimensional virtual reality. *Nature Neuroscience* **18**, 121–128 (2015).

65. Ravassard, P. *et al.* Multisensory control of hippocampal spatiotemporal selectivity.
Science (1979) **340**, 1342–1346 (2013).

66. Friedrich, J., Zhou, P. & Paninski, L. Fast online deconvolution of calcium imaging data.
PLoS Computational Biology **13**, (2017).

67. Kropff, E., Carmichael, J. E., Moser, M. B. & Moser, E. I. Speed cells in the medial
entorhinal cortex. *Nature* **523**, 419–424 (2015).

68. Iwase, M., Kitanishi, T. & Mizuseki, K. Cell type, sub-region, and layer-specific speed
representation in the hippocampal–entorhinal circuit. *Scientific Reports* **10**, 1–23 (2020).

69. Góis, Z. H. T. D. & Tort, A. B. L. Characterizing Speed Cells in the Rat Hippocampus.
Cell Reports **25**, 1872-1884.e4 (2018).

70. Claiborne, B. J., Amaral, D. G. & Cowan, W. M. A light and electron microscopic analysis
of the mossy fibers of the rat dentate gyrus. *Journal of Comparative Neurology* **246**, 435–
458 (1986).

71. Ishizuka, N., Weber, J. & Amaral, D. G. Organization of intrahippocampal projections
originating from CA3 pyramidal cells in the rat. *Journal of Comparative Neurology* **295**,
580–623 (1990).

72. Li, X. -G, Somogyi, P., Ylinen, A. & Buzsáki, G. The hippocampal CA3 network: An in vivo
intracellular labeling study. *Journal of Comparative Neurology* **339**, 181–208 (1994).

73. Ishizuka, N., Cowan, W. M. & Amaral, D. G. A quantitative analysis of the dendritic
organization of pyramidal cells in the rat hippocampus. *Journal of Comparative Neurology*
362, 17–45 (1995).

74. Witter, M. P. Intrinsic and extrinsic wiring of CA3: Indications for connectional
heterogeneity. *Learning and Memory* vol. 14 705–713 (2007).

75. Levene, M. J., Dombeck, D. A., Kasischke, K. A., Molloy, R. P. & Webb, W. W. In Vivo
Multiphoton Microscopy of Deep Brain Tissue. *Journal of Neurophysiology* **91**, 1908–
1912 (2004).

76. Barretto, R. P. J. *et al.* Time-lapse imaging of disease progression in deep brain areas
using fluorescence microendoscopy. *Nature Medicine* **17**, 223–229 (2011).

77. Gauthier, J. L. & Tank, D. W. A Dedicated Population for Reward Coding in the
Hippocampus. *Neuron* **99**, 179-193.e7 (2018).

78. Krishnan, S., Cherian, C. & Sheffield, M. E. J. Changing reward expectation transforms
spatial encoding and retrieval in the hippocampus. *bioRxiv* (2020)
doi:10.1101/2020.09.13.295469.

79. Pettit, N. L., Yuan, X. C. & Harvey, C. D. Hippocampal place codes are gated by
behavioral engagement. *Nature Neuroscience 2022 25:5* **25**, 561–566 (2022).

80. Chambers, A. R. & Rumpel, S. A stable brain from unstable components: Emerging
concepts and implications for neural computation. *Neuroscience* vol. 357 172–184
(2017).

81.   Mongillo, G., Rumpel, S. & Loewenstein, Y. Intrinsic volatility of synaptic connections — a
919   challenge to the synaptic trace theory of memory. *Current Opinion in Neurobiology* vol. 46
920   7–13 (2017).

921   82.   Kentros, C. G., Agnihotri, N. T., Streater, S., Hawkins, R. D. & Kandel, E. R. Increased
922   attention to spatial context increases both place field stability and spatial memory. *Neuron*
923   **42**, 283–295 (2004).

924   83.   Kinsky, N. R., Sullivan, D. W., Mau, W., Hasselmo, M. E. & Eichenbaum, H. B.
925   Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long
926   Timescales. *Current Biology* **28**, 3578-3588.e6 (2018).

927   84.   Mau, W., Hasselmo, M. E. & Cai, D. J. The brain in motion: How ensemble fluidity drives
928   memory-updating and flexibility. *Elife* **9**, 1–24 (2020).

929   85.   Wilson, M. A. & McNaughton, B. L. Dynamics of the hippocampal ensemble code for
930   space. *Science (1979)* **261**, 1055–1058 (1993).

931   86.   Danielson, N. B. *et al.* In Vivo Imaging of Dentate Gyrus Mossy Cells in Behaving Mice.
932   *Neuron* **93**, 552-559.e4 (2017).

933   87.   GoodSmith, D. *et al.* Spatial Representations of Granule Cells and Mossy Cells of the
934   Dentate Gyrus. *Neuron* **93**, 677-690.e5 (2017).

935   88.   Senzai, Y. & Buzsáki, G. Physiological Properties and Behavioral Correlates of
936   Hippocampal Granule Cells and Mossy Cells. *Neuron* **93**, 691-704.e5 (2017).

937   89.   Mongillo, G., Rumpel, S. & Loewenstein, Y. Inhibitory connectivity defines the realm of
938   excitatory plasticity. *Nature Neuroscience* **21**, 1463–1470 (2018).

939   90.   Sheffield, M. E. J. & Dombeck, D. A. Calcium transient prevalence across the dendritic
940   arbour predicts place field properties. *Nature* **517**, 200–204 (2015).

941   91.   Sheffield, M. E. J., Adoff, M. D. & Dombeck, D. A. Increased Prevalence of Calcium
942   Transients across the Dendritic Arbor during Place Field Formation. *Neuron* **96**, 490-
943   504.e5 (2017).

944   92.   Adoff, M. D. *et al.* The functional organization of excitatory synaptic input to place cells.
945   *Nature Communications* **12**, (2021).

946   93.   Jia, H., Rochefort, N. L., Chen, X. & Konnerth, A. Dendritic organization of sensory input
947   to cortical neurons in vivo. *Nature* **464**, 1307–1312 (2010).

948   94.   Wilson, D. E., Whitney, D. E., Scholl, B. & Fitzpatrick, D. Orientation selectivity and the
949   functional clustering of synaptic inputs in primary visual cortex. *Nature Neuroscience* **19**,
950   1003–1009 (2016).

951   95.   Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference
952   Atlas. *Cell* **181**, 936-953.e20 (2020).

953   96.   Paxinos, G. & Franklin, K. B. J. *The Mouse Brain in Stereotaxic Coordinates*. (Academic
954   Press, 2001).

97.   Zipfel, W. R., Williams, R. M. & Webb, W. W. Nonlinear magic: Multiphoton microscopy in
the biosciences. *Nature Biotechnology* vol. 21 1369–1377 (2003).

98.   Berens, P. CircStat : A MATLAB Toolbox for Circular Statistics. *Journal of Statistical
Software* **31**, 1–21 (2009).

99.   Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon
microscopy. *bioRxiv* 061507 (2016) doi:10.1101/061507.

100. Holtmaat, A. & Svoboda, K. Experience-dependent structural synaptic plasticity in the
mammalian brain. *Nature Reviews Neuroscience* vol. 10 647–658 (2009).

101. Skaggs, W., McNaughton, B. & Gothard, K. An Information-Theoretic Approach to
Deciphering the Hippocampal Code. *Advances in Neural Information Processing Systems*
5, 1030--1037 (1993).

102. Bechtold, B. Violin Plots for Matlab, Github Project. (2016).

ACKNOWLEDGEMENTS

We would like to thank Caleb Kamere and Eliott Levy for comments on the manuscript. This work was supported by NSF (M.J.G., S.L.S., NeuroNex #1707287), NIH (M.J.G., S.L.S., R01 RF1NS121919), the Larry Hillblom foundation (M.J.G.), and the Whitehall Foundation (M.J.G.).

AUTHOR CONTRIBUTIONS

W.T.R. and M.J.G. designed the experiments; M.J.G. performed the surgical implants; W.T.R., N.S.W., L.M., and M.J.G. conducted the imaging experiments and analyzed the data; G.L. and T.D.M. performed immunohistochemistry and K.K.S. analyzed the resulting images; C.H.Y. and S.L.C. performed optical modeling; W.T.R., N.S.W., and M.J.G. wrote the manuscript.

COMPETING INTERESTS

The authors declare no competing financial interests.

MATERIALS AND CORRESPONDENCE

Correspondence and material requests should be directed to: michael.goard@lifesci.ucsb.edu.