Title

Dynamics of cortical contrast adaptation predict perception of signals in noise.

Authors

Christopher F. Angeloni[1,2], Wiktor Młynarski[3,4], Eugenio Piasini[5], Aaron M. Williams[2,6], Katherine C. Wood[2], Linda Garami[2], Ann M. Hermundstad[7], Maria N. Geffen[2,6,8]*

[1]Psychology Graduate Group, University of Pennsylvania, Philadelphia, PA, USA
[2]Department of Otorhinolaryngology, University of Pennsylvania, Philadelphia, PA, USA
[3]Faculty of Biology, Ludwig Maximilian University of Munich, Germany
[4]Bernstein Center for Computational Neuroscience, Munich, Germany
[5]International School for Advanced Studies (SISSA), Trieste, Italy
[6]Neuroscience Graduate Group, University of Pennsylvania, Philadelphia, PA, USA
[7]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA, USA
[8]Department of Neuroscience, Department of Neurology, University of Pennsylvania, Philadelphia, PA, USA

*corresponding author: mgeffen@pennmedicine.upenn.edu

Abstract

Neurons throughout the sensory pathway adapt their responses depending on the statistical structure of the sensory environment. Contrast gain control is a form of adaptation in the auditory cortex, but it is unclear whether the dynamics of gain control reflect efficient adaptation, and whether they shape behavioral perception. Here, we trained mice to detect a target presented in background noise shortly after a change in the contrast of the background. The observed changes in cortical gain and behavioral detection followed the dynamics of a normative model of efficient contrast gain control; specifically, target detection and sensitivity improved slowly in low contrast, but degraded rapidly in high contrast. Auditory cortex was required for this task, and cortical responses were not only similarly affected by contrast but predicted variability in behavioral performance. Combined, our results demonstrate that dynamic gain adaptation supports efficient coding in auditory cortex and predicts the perception of sounds in noise.

Introduction

As we navigate the world around us, the statistics of the environment can change dramatically. In order to maintain stable percepts, the nervous system adapts to persistent statistical properties of sensory inputs. The efficient coding hypothesis postulates that the nervous system accomplishes this by matching the limited dynamic range of individual neurons to the statistics of incoming sensory signals[1], thereby efficiently encoding information within many types of environments[2–4]. Indeed, adaptation to environmental statistics has been found in many sensory modalities and species[5–13]. In the auditory system, neurons exhibit contrast gain control, adapting the gain of their response function to match the variability in level (contrast) of the incoming sounds[14–19]. Yet, it is unknown whether and how the dynamics of contrast gain control in the auditory system inform perception, as the dynamics of neuronal adaptation have not previously been measured simultaneously with behavior. The goal of our study was to test if the dynamics of contrast gain control in auditory cortex predict changes in the perception of targets embedded in noise.

Contrast gain control has been proposed as a mechanism for creating sensory representations of sounds that are invariant to background noise[20]. In the ferret auditory pathway, representations of sounds become more noise-tolerant in higher auditory areas, and the amount of noise-tolerance correlates with the strength of adaptation to the stimulus statistics, suggesting that level adaptation and gain control enhance the encoding of stimuli embedded in stationary noise[16]. Additionally, recent psychophysical studies found that perception in noise is altered by efficient adaptation to stimulus statistics. In humans, target level discriminability is greater in low contrast than in high contrast, an effect consistent with gain control observed in primary auditory cortex[19]. Similar relationships between gain control and behavioral percepts of sound location have also been found in ferrets[10] and guinea pigs[21]. However, in these previous studies, the measurement of psychophysical performance and neuronal gain control were performed separately in human and animal subjects, respectively, so it remains unclear how variability in adaptation over sessions or subjects relates to behavioral performance. Additionally, these previous studies focused primarily on behavioral performance after complete adaptation to stimulus contrast, whereas recent work has highlighted the dynamical nature of efficient adaptation[22,23]. In the current study, we aimed to assess whether and how the dynamics of contrast gain control reflected efficient adaptation and predicted behavioral performance across subjects.

First, we established a normative framework to model the neuronal dynamics of gain control and predict how efficient contrast adaptation should affect the detection of signals in noise. We then derived a procedure for estimating moment-to-moment changes in neuronal gain based on generalized linear models (GLM) and found that the dynamics of gain control in auditory cortex were asymmetric, as observed in the normative model. Next, to test the role of contrast gain control in auditory behavior, we trained mice to detect targets after a change in background contrast. We found that contrast-induced changes in behavioral target detection threshold, sensitivity, and behavioral adaptation dynamics followed the normative model predictions. Furthermore, we found that auditory cortex was necessary for detection in the presence of a background, but not for detection in silence, suggesting a distinct role of auditory cortex in separating targets from the background. Building on this finding, we found that the dynamics of cortical encoding of targets resembled the normative model predictions and observed behavioral adaptation, and that population activity in the auditory cortex predicted individual variability in behavioral performance. Finally, we estimated cortical gain during the task, finding that variability in neuronal gain predicted variability in task performance. Combined, our results identify a predictive relationship between efficient adaptation via gain control and detection of signals in noise, and provide a normative framework to predict the dynamics of behavioral performance in changing sensory environments.

Results

Target-in-background detection task and normative model for task predictions.

To assess how behavioral performance is impacted by adaptation to stimulus contrast, we first devised a GO/NO-GO task in which mice were trained to detect targets embedded in switching low and high contrast backgrounds. During each trial, the mouse was presented with dynamic random chords (DRCs) of one contrast, which switched after 3 s to the other contrast. At variable delays after the contrast switch, broad-band target chords were superimposed on the background chords, and mice were trained to lick for a water reward upon hearing the target (henceforth, we refer to high-to-low contrast trials as "low contrast" and low-to-high contrast trials as "high contrast", referring to the contrast where mice detected targets). Target trials were interleaved with background-only trials, in which the mouse was trained to withhold licking, but would receive a 7s timeout for licking after the contrast switch (Figure 1a,b). To assess behavioral sensitivity to targets, we parametrically varied target level in each contrast, and to assess behavioral adaptation, we parametrically varied target timing (Figure 1c). This stimulus design allowed us to quantitatively test how the dynamics of adaptation to background contrast affect perception.

To estimate the optimal time course of contrast gain control and predict its impact on target detection, we developed a normative model of behavioral performance constrained by efficient neuronal coding. In this model, we simulated a neuron designed to encode stimuli with minimal error. To efficiently exploit its finite dynamic range, the model neuron estimated the contrast of the recent stimuli, and adjusted the gain of its nonlinearity to minimize the error in estimated contrast (Figure 1d, panels 1-3; *Methods*)[22,23]. Adding targets at different levels and times relative to contrast transitions allowed us to probe the sensitivity of the model neuron to targets of varying strength over the time course of adaptation (Supplementary Figure 1c,d). When varying target strength and measuring the "psychometric" performance of the model (Equation 1, *Methods*), we found decreased detection thresholds and steeper slopes in low contrast relative to high contrast (Figure 1e). When varying target timing, two factors affected target discriminability: 1) The change in the stimulus distribution after the contrast switch; 2) The effect of gain adaptation on responses to the background (Figure 1f,g; Supplementary Figure 1c,d). Recapitulating previous results[14], the dynamics of the model responses to targets were asymmetric across each contrast (Figure 1f). These dynamics were well characterized by a single effective timescale, which we quantified by fitting an exponential (Equation 2) to each transition. The normative model presented three primary predictions for target detection behavior: When adapted to low contrast, 1) target detection thresholds will be lower and 2) model psychometric functions will have steeper slopes, and, 3) target detectability during the adaptation period will be asymmetric: rapidly decreasing after a switch to high contrast, and slowly increasing after a switch to low contrast (Figure 1h).

Estimated cortical gain dynamics follow a normative model of gain control.

Previous work on contrast gain control used static models, measuring steady-state gain after the neuron fully adapted to the new stimulus[14,16,17,19], but see[15,24]. We developed a model to estimate the gain of neurons in auditory cortex over time following a contrast transition. This generalized linear model (GLM) was fit to data recorded from the auditory cortex of a naive mouse (n = 97 neurons) presented with 3 s alternations of low and high contrast DRCs (Figure 2a,b).

The inference model is a GLM with dynamic gain control (GC-GLM) that decomposes the relationship between spiking activity (y_t) and the presented sounds into a stimulus component (x_t), contrast component ($\overline{\sigma}/\sigma_t$), and an interaction between the stimulus and the contrast ($x_t * \overline{\sigma}/\sigma_t$, where $\overline{\sigma}$ is an arbitrary constant, defined as the contrast at which the gain is 1; *Methods*). We calculated a gain index (w_t) from the fitted model parameters (Figure 2b), which quantified whether gain control estimated by the model was optimal given the background contrast levels. For the contrast levels used in this study, w_t = 1.5 indicates an optimal increase in gain during low contrast, w_t = 0.5 indicates an optimal decrease in gain during high contrast, and w_t = 1.0 indicates no gain control (*Methods*). To validate the model and gain index we simulated neurons with defined temporal trajectories of gain control and found that the model accurately recovered the ground-truth simulated gain dynamics (Supplementary Figure 2; *Supplementary Information*). For comparison, we also fit previously described linear-nonlinear (LN) models to each neuron[14,16,17,19], one with a static output nonlinearity (static-LN), and one with a contrast-dependent, or gain-controlled output nonlinearity (GC-LN, Figure 2c; representative neuron: Figure 2d-g). In this neuron, the fits of both the GC-LN model and GC-GLM exhibited contrast gain control, as characterized by high gain in low contrast and low gain in high contrast (Figure 2f and g, respectively), suggesting that both models capture similar contrast-driven changes in cortical gain.

Figure 1. Target in background detection task and normative model predictions.

a, Experimental setup. **b,** GO/NO-GO task design. Spectrograms are plotted for example NO-GO and GO trials with transitions from low to high contrast (top row) and high to low contrast (bottom row), with waveforms plotted below each spectrogram (color bar indicates the sound level). Below the example trials, the timing of the response window, schematic licks, and responses to licks are plotted. For NO-GO trials, licks in the response window received a timeout. For GO trials, licks in the response window were rewarded with 5μL of water. **c,** Example target parameters. *Top*: varied target levels, with level in dB SNR indicated by the color bar. *Bottom*: varied target times, where each arrow indicates a potential target delay. **d,** Normative model of efficient gain control. Target and background distributions for each contrast are indicated in the left panel. (1) Target stimuli are indicated with circles, while the background stimulus is indicated by a line. The stimulus response in a given time window is transformed by an adapting nonlinearity to generate spikes. (2) The spiking responses are decoded to update an estimate of the stimulus variance. (3) The gain of the nonlinearity is adjusted to optimally predict the variance of the next timestep. *Inset*: Example spike distributions of the model neuron in low and high contrast for targets (dark histograms), and background (light histograms). **e,** Model target-from-background discriminability as a function of contrast and target level. Circles indicate model performance overlaid with logistic function fits (solid lines) and thresholds (dashed lines). **f,** Model discriminability over time in low and high contrast. Circles indicate model performance overlaid with exponential function fits. **g,** Model gain dynamics over time in each contrast. **h,** From top to bottom: model predictions for target detection thresholds, slopes, and adaptation time constants in each contrast. Source data are provided as a Source Data file.

Qualitatively, GC-GLM outperformed standard LN models, primarily by capturing the adaptation dynamics after the transition (Figure 2d, middle panel), allowing us to analyze the gain control index (w_t) as a function of time (Figure 2d, bottom panel; Figure 2g). To test whether the GC-GLM could better account for the data than standard models, we compared cross-validated correlations of the model predictions with the trial-averaged PSTH for each neuron, finding a significant effect of model type on the correlations (Kruskal-Wallis test: $H(2) = 93.61$, p = 6.70e-21). GC-GLM correlation was significantly higher (Median (Mdn) = 0.75, Inter-Quartile Range (IQR) = 0.24) than the GC-LN model (Mdn = 0.54, IQR = 0.49, p = 4.41e-6, post-hoc sign-rank test) and the static-LN model (Mdn = 0.25, IQR = 0.73, p = 9.56e-10, post-hoc sign-rank test). Consistent with previous studies[14,15], we also found that the GC-LN model outperformed the static-LN model (p = 3.50e-6, Figure 2h). To compare the GC-GLM to an existing dynamical model of contrast gain control, we also fit a recent model of dynamic contrast adaptation[19] to our data (Supplementary Figure 3a). Across the neurons in this dataset, we found that the GC-GLM was a better predictor of neuronal activity than the existing model (Supplementary Figure 3b).

We then quantified whether the GC-GLM reliably detected gain control in the neuronal population. Here, we defined steady-state gain control by calculating the change in w_t between high (w_H) and low (w_L) contrast after the gain has stabilized (1 s after the contrast switch). Based on our definition of w_t, $w_H - w_L = -1$ if gain control is optimal (*Methods*). Across all neurons, we found significant gain control (Mdn: -0.10, IQR: 0.35, sign-rank test: $Z = -2.90$, p = 0.004; Figure 2i). To further validate the GLM estimates of gain, we compared the GC-GLM gain control indices at steady-state to those of the GC-LN model and found a significant relationship (linear regression: $F(1,95) = 12.20$, p = 7.33e-4, R^2 = 0.11; Figure 2j). Together, these results demonstrate that the GC-GLM model better accounts for the neuronal data by incorporating the dynamics of gain control and conclude that this method captures a similar estimate of steady-state gain control when compared to standard models.

Finally, we analyzed the dynamics of gain control by fitting w_t after each contrast switch with an exponential function (Figure 2g). In neurons with gain control ($w_t < 0$ at steady state), the average time course of w_t was asymmetric across contrast transition types, rapidly decreasing after a switch to high contrast, and slowly increasing after a switch to low contrast (n = 45 neurons; Figure 2k). Within this same population, we quantified the timescale of adaptation to each contrast using the time constant (τ) of each exponential fit, finding significantly longer time constants in low contrast (Mdn = 0.29, IQR = .39) relative to high contrast (Mdn = 0.048, IQR = 0.094; sign-rank test: $Z = 4.52$, p = 6.16e-6; Figure 2l). This asymmetry in gain adaptation reflected the dynamics of the normative model (Figure 1g), prior electrophysiological studies[14], and the behavior of previously described optimal variance estimators[25].

Mouse behavioral detection is modulated by background contrast.

We next tested whether the asymmetry in gain control in cortex was reflected in behavioral sensitivity to targets in background noise. Mice were initially trained in a simple version of the GO/NO-GO task where they were required to lick in response to a target and withhold licks on trials without a target (Figure 1b, 3a). Out of the 25 mice trained, 24 mice learned this task reliably, typically reaching criterion performance of 80% correct within 2-3 weeks in either contrast (Figure 3b). False alarm rates were significantly higher in high contrast than in low contrast (Supplementary Figure 4a), suggesting that detection is more difficult in high contrast, which we discuss next.

By varying the level of presented targets, we collected psychometric curves for each mouse in each contrast. To isolate the effect of stimulus contrast on behavioral performance, we included sessions from mice that were exposed to similar target levels in low and high contrast (Figure 3c; see Supplementary Figure 4 and Supplementary Results for results using different target ranges). In this cohort, targets were easier to detect in low contrast: we observed significantly lower detection thresholds in low contrast (Mean (M) = 8.79, standard deviation (std) = 3.13) as compared to high contrast (M = 15.39, std = 3.27; paired t-test: $t(10) = -4.20$, p = 0.0057, Figure 3d). Furthermore, psychometric slopes were significantly steeper in low contrast (M = 0.040, std = 0.0048) compared to high contrast (M = 0.036, std = 0.0026; paired t-test: $t(10) = 3.037$, p = 0.023; Figure 3e). Combined, these results demonstrate that targets were easier to detect in low contrast, as predicted from the normative model (Figure 1h).

Figure 2. Cortical adaptation to sound contrast is asymmetric.

a, Schematic[26] of acute recordings from auditory cortex. **b,** Schematic of the linear-nonlinear (LN) model, with a static (grey) or gain-controlled (blue, red) nonlinearity. **c,** A Poisson GLM for estimating gain dynamics. Note that the model estimates gain as the interaction between the stimulus and contrast. **d,** *Top*: Spike raster for a representative unit. Blue and red horizontal bars indicate low and high contrast periods of the trial, respectively. *Middle*: the spike rate of the neuron is overlaid with the predictions from a static LN model, LN model with gain control, or GLM with gain control. *Bottom*: gain index, w_t, estimated from the GLM parameters (*Methods*). Dashed lines indicate optimal and no gain control (*Methods*). Orange trace indicates the gain dynamics of the neuron. **e,** Spectro-temporal receptive field (STRF) fit to this neuron. Color bar indicates the strength of the filter response. **f,** Nonlinearities fit to the STRF prediction in low and high contrast. **g**, Dashed blue and red lines indicate the gain index of the example cell in low and high contrast. The overlaid solid lines are exponential fits to the data. **h,** Cross-validated Pearson correlation coefficients between the trial averaged model predictions and spike rates for each model (n=97 neurons; colors as in d). Error bars indicate 95[th] percentiles around the median. Asterisks indicate the results of two-way sign-rank tests. **i,** Distribution of gain control estimated by the GLM. Orange line indicates the median. Asterisks indicate the results of a two-way sign-rank test ($p = 0.0040$). **j,** Gain control estimates for each neuron from the GC-GLM and the GC-LN model (black dots) overlaid with the best linear fit (black line) and 95% confidence interval of the fit (grey area). Asterisks indicate whether the linear model was significantly different from a constant model ($p = 7.33e-4$). **k,** Gain index for all of the neurons with gain control (n=45). Light lines are the average ±SEM, while the dark lines are exponential fits to the average. **l,** Adaptation time constants from gain-controlled neurons after a switch to low (blue dots) and high contrast (red dots). Asterisks indicate the results of a two-way sign-rank test ($p = 6.16e-6$). Source data are provided as a Source Data file.

To assess behavioral adaptation to the background contrast, we presented targets at threshold level
at variable delays following the contrast transition. We observed behavioral time courses consistent with the
normative model and with gain measured in auditory cortex: after a switch of the stimulus to high contrast,
detection rates decreased quickly over time, but after a switch to low contrast detection, rates increased slowly
over time (Figure 3f). In high contrast, the first significant drop in performance occurred between the first two
time points, while in low contrast the first significant increase in performance occurred between the first and
third time points (Figure 3f, Supplementary Table 1). Indeed, fitting exponential functions to performance over
time revealed that behavioral adaptation was significantly faster in high contrast (median and interquartile
range of time constant: Mdn = 0.023, IQR = 0.082) compared to low contrast (Mdn = 0.13, IQR = 0.13; sign-
rank test: Z = 2.75, p = 0.0060; Figure 3g).
To directly compare the predictions of the normative model presented in Figure 1 to behavioral
performance, we computed a contrast modulation index to measure the percent change in behavioral and
model parameters between high and low contrast (CMI: Equation 3, *Methods*). To assess whether the model
prediction was within the range of expected behavioral values, we computed the 95% confidence intervals of
the behavioral CMI values using a bootstrap procedure. We found that the CMI values of the normative
predictions fell within the range of expected CMI values for behavioral thresholds and adaptation times. As
observed in behavior, the model predicted a decrease in slope in high contrast; however, the magnitude of
the predicted decrease was larger than the range of observed slope CMI values (Figure 3h). Taken together,
the behavioral results were qualitatively consistent with all three predictions from the normative model (Figure
1h): 1) Detection thresholds are lower in low contrast; 2) Psychometric slopes are steeper in low contrast; 3)
Performance decreases rapidly in high contrast and increases gradually in low contrast.
210
Auditory cortex is necessary for detection in background noise.
Whereas gain control is present in many areas along the auditory pathway, it is strongest in auditory
cortex[16,19]. As such, we hypothesized that auditory cortex supports the detection of sounds in the presence of
background noise. To test whether auditory cortex is required for task performance, we inactivated auditory
cortex using the GABA-A receptor agonist muscimol. We validated that muscimol disrupts cortical coding of
target sounds by applying muscimol topically to the cortical surface during passive playback of the behavioral
stimuli, finding near complete suppression of target responses (Supplementary Figure 5a-f, *Supplementary
Information*).
To test if inactivation of auditory cortex affects behavioral performance, we repeated the same
experiments in behaving mice, administering muscimol or saline bilaterally through chronically implanted
cannulae (4 mice; Figure 4a). We found a profound decrease in the response rates to targets and background
in both contrasts (Figure 4b). We quantified these effects on the psychometric curve using a three-way
ANOVA with cortical intervention (muscimol or saline), contrast, and target level as factors. We found
significant main effects of cortical intervention ($F(1,307)$ = 278.63, p = 3.83e-44), contrast ($F(1,307)$ = 4.39,
p = 0.037) and level ($F(6,307)$ = 40.90, p = 7.54e-36). Post-hoc tests showed that muscimol application
significantly decreased hit rates in both contrasts by 31.45% (95% CI: [27.76, 35.14], p = 1.060e-10), whereas
an increase in background contrast significantly decreased hit rates in both intervention conditions by 3.95%
(95% CI: [2.57, 7.64], p = 0.036). Furthermore, we observed significant interactions between target level and
cortical intervention ($F(6,307)$ = 14.11, p = 4.47e-14), and between target level and contrast ($F(6,307)$ = 2.97,
p = 7.87e-3), but we did not observe a significant interaction between contrast and cortical intervention,
suggesting that muscimol has the same effect in low and high contrast. To quantify the effects of muscimol
on behavioral performance, we extracted response rates to the maximum target level, false alarm rates,
thresholds, and slopes of psychometric functions fit to each session, and found that muscimol significantly
reduced every measure of behavioral performance, with the exception of behavioral threshold (Figure 4c,
Supplementary Table 1). From these results, we can conclude that auditory cortex is necessary for detecting
targets in background, regardless of background contrast.
A potential alternative effect of muscimol is a general loss of function that is not specific to hearing
target sounds. To control for this, we devised an alternative to the detection in background task where mice
detected targets in silence (Figure 4d). To ensure equivalency between the two tasks, we took the highest-
level target trials in the target-in-background task (25dB SNR in high contrast) and removed the background
noise during the target detection period (Figure 4e, bottom). In this new task, mice heard the exact same
targets as in the target-in-background task except that the background noise pre- and post-target
presentation was removed. This allowed us to test whether auditory cortex is specifically required for target
detection in the presence of a background (*Methods*).
245

Figure 3. Target detection performance is consistent with the normative model predictions.

a, Schematic of the behavioral setup and task outcomes. **b,** Performance as a function of task exposure. Individual traces indicate performance of mice first trained to detect targets in low contrast (blue) or high contrast (red). Thick lines are a 7-day running average across mice in each training group. The dashed red trace near 0.5 was the performance of a mouse who failed to learn the task in high contrast and was excluded from further analysis. **c,** Individual psychometric curves in low and high contrast (n=12 mice), overlaid with psychometric fits to the average. Dashed vertical lines indicate detection thresholds. **d,** Detection thresholds from the data plotted in c (n=12 mice; two-way paired t-test: p = 0.0057). **e,** Slopes of the psychometric functions plotted in c (n=12 mice; two-way paired t-test: p = 0.023). Error bars indicate ±SEM over mice. **f,** Detection performance of threshold level targets presented at different delays from the contrast switch (circles indicate mean ±SEM). Performance is overlaid with exponential fits in each contrast. Horizontal lines above the plot indicate significant sign-rank tests after false-discovery correction. **g,** Adaptation time constants from exponential functions fit to individual mice (n=21). Asterisks indicate the result of a two-way sign-rank test (p = 0.0060). Error bars indicate ±SEM over mice. **h,** Comparison of average normative model predictions (red lines) to the data (grey distributions) for psychometric thresholds, slopes and adaptation time. The y-axis is the contrast modulation index (CMI, *Methods*) of the values in each plot. Data distributions were estimated using bootstrapping of the mean (threshold, slope) or median (τ) CMI (10,000 samples with replacement). Source data are provided as a Source Data file.

To assess behavioral performance in this new task, we modulated detection difficulty by attenuating
the level of each target. As observed previously, inactivation of auditory cortex impaired detection in high
contrast (Figure 4e, top). However, cortical inactivation had little effect on behavioral performance in silence
(Figure 4e, bottom). We quantified these effects on behavior using a three-way ANOVA with cortical
intervention (muscimol or saline), task (detection in background or silence), and target level as factors (n =
26 sessions from 2 mice). We found significant main effects of intervention ($F(1,181)$ = 62.83, p = 3.62e-13),
task ($F(1,181)$ = 6.82, p = 9.86e-3), and level ($F(6,181)$ = 46.16, p = 1.69e-32). Muscimol significantly reduced
hit rates by 20.2% (95% CI: [15.19, 25.17], p = 1.060e-10, post-hoc). Hit rates for targets presented in silence
were significantly elevated by 6.65% relative to targets presented in background (95% CI: [1.65, 11.64], p =
0.0090). Furthermore, we found significant interactions between cortical intervention and task type ($F(1,181)$
= 6.36, p = 0.013), intervention and level ($F(6,181)$ = 3.47, p = 2.98e-3), and level and task type ($F(6,181)$ =
8.47, p = 5.43e-8). As before, we parameterized behavioral performance by fitting the performance on each
session with a psychometric curve, and we extracted the response rates to the maximum target level, false
alarm rates, response rates at threshold level, and slopes of psychometric functions. During the target-in-
background task, we found significant effects of muscimol on the response rates at maximum level and
threshold, a moderate effect on psychometric slope, and no effect on false alarm rate. However, muscimol
application had no significant effect on any of these measures in the target-in-silence task (Figure 4f,
Supplementary Table 1). Taken together, these results show that while both cortical inactivation and the
presence or absence of background noise affected behavioral performance, these effects interacted:
muscimol had a larger effect on performance when background noise was present.
Combined, our findings demonstrate that the auditory cortex is specifically required for detection in the
presence of background noise, but not in silence. Our next goal was to test whether neuronal activity in AC is
predictive of behavioral performance.
272
Cortical codes predict individual behavioral performance.
To better understand how representations in the auditory cortex could give rise to behavior, we
chronically recorded from populations of neurons in auditory cortex of 12 mice while they performed the
psychometric task (Figure 5a, Supplementary Figure 6). When using microdrives with drivable tetrodes, we
lowered the tetrodes a small distance at the end of each session to record from new populations of cells. In
the 242 sessions analyzed, we recorded from 18±15 neurons simultaneously (mean±standard deviation), with
a maximum of 73 and minimum of 3 neurons simultaneously recorded. For the following analyses, we only
included neurons with spike rates greater than 1 Hz, realistic spike waveforms (*Methods*), and with significant
responses to targets (AUC value significantly greater than 0.5 at two or more levels, Figure 5b, inset;
Methods). Following these selection criteria, 12±9 neurons were included in analysis for each session.
To quantify the representations of targets and background in the neuronal population (example
responses in Figure 5b,c), we adapted a population vector approach[27] to generate a target-from-noise
discriminability metric using population activity (*Methods*). This method allowed us to project trial distributions
in n-dimensional neuronal space along a single dimension that separated target and background trials (Figure
5d, left panel). We then estimated the criterion projection value that best predicted whether each trial
contained a target or just background[28] (Figure 5d, right).
This population decoding method allowed us to estimate neurometric functions to directly compare to
psychometric functions for each mouse (Figure 5e). On average, neurometric and psychometric functions
were qualitatively similar (Figure 5f) and highly correlated across sessions (Figure 5g). To assess the
relationship between neurometric and behavioral performance, we first considered the potential variables that
were likely involved to modulate behavior, namely: the level of the presented targets, the contrast of the
background, and neuronal variability that could fluctuate from session to session. To isolate the effects of
each of these variables, we used a linear model to predict behavioral performance on each session using
target level, contrast, and the neurometric performance as predictors ($F(4,794)$ = 593.0, p = 3.15e-202, R^2 =
0.69, Figure 5i, top).
298

Figure 4. Inactivation of auditory cortex selectively disrupts detection of targets in background sounds.

a, Schematic of chronic muscimol and saline application in behaving mice. Legend indicates the three potential background conditions in the task. **b,** Behavioral performance on individual sessions (light traces) as a function of contrast (red, blue) and muscimol or saline application (dashed and solid lines, respectively) for n=44 sessions. -Inf indicates performance on background-only trials. Dots indicate the average performance across sessions ±SEM, overlaid with logistic fits to the data. **c,** Effects of muscimol and contrast on multiple behavioral measures. Bars indicate the mean performance across sessions ±SEM, while dots indicate performance on individual sessions (n=23 muscimol sessions and 21 saline sessions). Asterisks indicate the significance of two-way rank-sum tests (Supplementary Table 1). **d,** Example spectrograms and waveforms with a target presented in high contrast (top) and the same stimulus when the target was presented in silence (bottom). Color bar indicates the sound level, with black indicating silence. **e,** Behavioral performance in high contrast (top) and in silence (bottom) for n=26 sessions. Error bars are ±SEM across sessions. Formatting as in b. **f,** The effect of muscimol on multiple behavioral measures when targets were presented in high contrast (red bars; n=10 sessions) or in silence (black bars; n=16 sessions). Error bars are ±SEM across sessions. Asterisks indicate the significance of two-way rank-sum tests. Formatting as in c. In all plots: ns, not significant; †p<0.1; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001. Detailed statistical results for c and f are in Supplementary Table 1. Source data are provided as a Source Data file.

As expected, we found that basic stimulus features (target level and background contrast) were strong
predictors of behavioral performance, such that performance was higher for increased target levels (b = 0.14,
95% CI [0.12, 0.15], p = 4.15e-65), while performance was lower for increased contrast (b = -0.050, 95% CI
[-0.071, -0.028], p = 5.67e-6). These results confirm our previous results on the behavioral impacts of stimulus
contrast (Figure 3). Crucially, we also observed that population activity was predictive of behavioral
performance at a session level, such that an increase in neurometric performance was accompanied by an
increase in behavioral performance (b = 0.016, 95% CI [0.0017, -0.031], p = 0.029). Finally, we found that the
relationship between behavioral and neuronal performance was mediated by stimulus contrast (b = 0.026,
95% CI [0.0042, 0.048], p = 0.020; Figure 5i, bottom). Taken together, these results suggest that session-to-
session fluctuations in cortical population activity resulted in corresponding fluctuations in behavior, and that
this relationship was affected by stimulus contrast.
To control for the effect of target level on the relationship between behavioral and neuronal
performance, we first visualized the data by regressing out the effect of level on the model terms and
behavioral performance: this visualization demonstrates the positive relationship between the two terms
(Figure 5g, inset), as captured by the significant neurometric coefficient in Figure 5i. To further control for the
effect of level, we split the data into low, medium and high target levels (Figure 5h). In all level-splits, the
relationship between neurometric and behavioral performance was significant (Supplementary Table 1).
These findings suggest that regardless of the target level, neuronal performance predicted behavioral
performance.
Next, we assessed whether neurometric performance was affected by contrast similarly to behavioral
performance. To isolate the effect of contrast on neurometric curves and best compare to psychometric curves
presented in Figure 3, we selected only recording sessions with matched target ranges (n = 117) and fit each
neurometric curve with a logistic function. We found that neurometric function thresholds increased in high
contrast (*Mdn[IQR]* of low contrast sessions (n=82): 7.19[4.83], high contrast sessions (n = 35): 13.47[8.88];
rank-sum: Z = 4.69, p = 1.34e-6, Figure 5j), and that neurometric slopes decreased in high contrast (low
contrast: 0.036[0.014], high contrast: 0.030[0.021]; Z = 1.88, p = 0.029, Figure 5k). These findings
corroborated the observed changes in psychometric functions (Figure 3) and the predictions of the normative
model (Figure 1).
Combined, these results demonstrate that parameters of neurometric and psychometric functions are
affected by contrast as predicted by a normative model of gain control. We also find that individual variation
in behavioral performance is predicted by population activity in auditory cortex, independently of the effect of
contrast and target level, which further supports the role of auditory cortex in the detection of signals in noise.
333
Dynamics of cortical target detection during adaptation.
We next measured how cortical discriminability evolved as a function of time and contrast in sessions
where mice were presented with targets at threshold level at different offsets relative to the contrast switch.
In line with our behavioral results (Figure 3f), we found that in high contrast the first significant drop in cortical
discriminability occurred between the first two target times, while in low contrast the first significant drop
occurred between the first and third target times (n = 43 recording sessions; Supplementary Table 1; Figure
5l, left). To quantify the speed of neuronal adaptation, we fit the average neuronal discrimination time course
for each mouse with an exponential function (n = 12 mice). Consistent with the normative model (Figure 1f-
h), and with gain control dynamics estimated from cortical activity (Figure 2k,l) and behavior (Figure 3f,g), we
found asymmetric adaptation in the neuronal responses, with larger adaptation time constants in low contrast
(*Mdn* = 0.14, *IQR* = 0.21) relative to high contrast (*Mdn* = 0.033, *IQR* = 0.16; sign-rank test: p = 0.016; Figure
5l, right).
346
Cortical gain predicts behavioral performance.
Our results so far demonstrate that population codes in auditory cortex are not only shaped by stimulus
features, but also predict variation in behavioral performance. To assess the role of cortical gain in behavior,
we leveraged the design of the background sounds to estimate spectrotemporal receptive fields (STRFs) and
nonlinearities of neurons recorded during task performance. For each neuron, we fit a model with a static
nonlinearity (static-LN) or a model with gain control (GC-LN; Figure 6a-d). We then pooled the neurons
recorded across sessions including only neurons with strong stimulus responses in both contrasts (*Methods*).
To ensure that observed changes in gain were not due to changes in STRF shape during the task, we first
compared STRFs estimated from each task epoch (ie. low and high contrast) and found that STRF properties
were stable across the trial (Supplementary Figure 7, *Supplemental Methods*).

Figure 5. Cortical responses to targets predict behavioral performance and exhibit contrast adaptation.

a, Schematic of chronic recordings during behavior. **b,** Spike raster from a neuron recorded during the task with average PSTHs. *Inset*: area under the ROC curve (AUC) for each level ±bootstrapped 95% CI. **c,** Neurograms of populations of simultaneously recorded neurons in low (left) and high contrast (right) sessions. Color bars indicate the difference in firing rate between the noise-only condition and each target level. The arrow indicates the cell in b. **d,** *Upper left*: schematic for estimating the population coding direction (CD). *Lower right*: probability distributions of CD projections for noise-only trials (grey) and the highest-level target trials (blue). The vertical red line indicates the criterion for the decoder. **e,** Neurometric performance and behavioral performance in representative low contrast (top) and high contrast (bottom) sessions. Dark dots are decoder performance, while light dots indicate behavioral performance, overlaid by logistic function fits. The arrow indicates the decoder performance for the distributions in d. **f,** Average psychometric and neurometric performance ±SEM over sessions. Formatting as in e. **g,** Behavioral performance as a function of neurometric performance for n=102 sessions. Each point is performance at a single target level in a single session. Point color indicates contrast, while point size indicates target level. Dots joined by grey lines were collected in the same session. *Inset*: Same data as in g, with the effect of target level regressed out (*!L*). The black line indicates the model fit and 95% CI. **h,** Behavioral performance as a function of neural performance split by target levels. Text indicates the correlation coefficient and its significance value. **i,** *Top:* the linear model used to predict behavior. *Bottom:* model coefficients ±standard error for target level (*L*), contrast (*C*), neurometric performance (*N*) terms, and the interaction between the latter two terms (*C*N*). Asterisks indicate significant predictors (n=102 sessions). **j,** Neurometric thresholds ±SEM for sessions in low (n=82) and high contrast (n=35). Asterisks indicate significant two-way rank-sum test (*p* = 1.34e-6). **k,** Same as j, but for neurometric slopes (two-way rank-sum test: *p* = 0.029). **l,** *Left*: Neurometric performance ±SEM as a function of target delay over sessions (n=87). Formatting and statistical tests are the same as Figure 3f. *Right*: Adaptation time constants ±SEM fit to the neurometric response from n=12 mice. Asterisk indicates a significant two-way sign-rank test (*p* = 0.016). Source data are provided as a Source Data file.

First, we compared the cross-validated performance of the static-LN model versus the GC-LN model. We found higher correlations using the GC-LN model (Mdn = 0.82, IQR = 0.17) relative to the static-LN model (Mdn = 0.67, IQR = 0.12; n = 2,792 neurons; sign-rank test: Z = -36.74, p = 1.88e-295; Supplementary Figure 8a). We also found significantly higher gain in low contrast (Mdn = 0.10, IQR = 0.13) than in high contrast (Mdn = 0.041, IQR = 0.023; sign-rank test: Z = 37.92, p = 1.070e-314; Figure 6e, inset). These results demonstrate that LN models can more accurately predict cortical activity when incorporating contrast gain control, and confirm previous reports of robust gain control in mouse auditory cortex[17–19].

Based on our previous results, we expected that the strength of gain control in auditory cortex would predict target detectability. When fitting the GC-LN model, we separately estimated neuronal gain during the adaptation period of the trial and the target period of the trial (defined as the time periods before and after the contrast switch, respectively; Figure 6b). To quantify the effects of contrast and trial period on gain, we performed a two-way ANOVA, with gain as the dependent variable, and contrast, trial period, and their interaction as factors (n = 2,262 neurons, after excluding outliers, see *Methods*). As expected, we found a significant main effect of contrast (F(1,4523) = 431.03, p = 1.60e-91). Furthermore, there was a significant main effect of trial period (F(1,4523) = 35.79, p = 2.36e-9) and a significant interaction between contrast and trial period (F(1,4523) = 77.91, p = 1.51e-18). Post-hoc tests revealed that, in low contrast, gain during the target period significantly increased (0.032, 95% CI: [0.024, 0.040], p = 3.77e-9), but did not significantly change in high contrast (0.0062, 95% CI: [-0.017, 0.014], p = 0.18; Figure 6e). These findings indicate that neuronal gain is not only sensitive to stimulus contrast, but also increases during the target period of the trial, specifically in low contrast.

To visualize the gross relationship between gain and behavioral performance, we first averaged the gain of stimulus-responsive neurons during the target period of the trial in each session (n = 168 sessions across 13 mice). We then selected only low contrast sessions and split the data by the median gain in the target period, computing the average psychometric curves for sessions in the bottom versus the top 50th percentile (Figure 6f, inset). We observed that sessions with high gain had steeper slopes and lower thresholds than sessions with low gain (Figure 6f). To quantify the relationship between gain and task performance, we fit a mixed-effects model using contrast and gain during the target period as fixed effects, mouse identity as a random effect, and either psychometric slopes or thresholds as the dependent variable. This approach allowed us to separate the neuronal and behavioral impact of contrast gain control from the effect of session-to-session fluctuations in gain. We tested whether gain and contrast were significant predictors of behavioral performance by comparing the full model to null models excluding either gain or contrast. We found that the model including gain was a better predictor of behavioral threshold than was the null model (Likelihood Ratio Test: χ^2(1) = 5.82, p = 0.016), indicating that thresholds decreased by 3.046 dB SNR ±1.24 (SEM) for every 10% increase in gain. Using a similar procedure, we found that contrast was also a significant predictor of behavioral threshold (Likelihood Ratio Test: χ^2(1) = 5.84, p = 0.038), with the step from high to low contrast inducing a decrease in behavioral thresholds of 3.27 dB SNR ±1.33 (Figure 6g).

We applied the same analysis to test the effects of contrast and gain on psychometric slope (Figure 6f), again finding that gain significantly predicted psychometric slopes (Likelihood Ratio Test: χ^2(1) = 6.96, p = 0.0083), such that the psychometric slope increased by 0.16 dB/PC ±0.060 for every 100% increase in gain. However, contrast did not significantly improve the fit of this model (Likelihood Ratio Test: χ^2(1) = 2.28, p = 0.13; Figure 6h). This result is not entirely unexpected, given that we observed no effect of contrast on psychometric slopes when comparing across sessions with different target distributions (Supplementary Figure 2b), which is true of the sessions used in this analysis.

Our findings suggest that the relationship between gain and behavioral performance is shaped by two sources: contrast-induced gain control and fluctuations in gain from session to session. To further disentangle the relationship between these two sources of behavioral modulation, we repeated the mixed effects models, this time using gain during the adaptation period as the predictor of interest. We hypothesized that gain in this period should not be predictive of behavioral performance, as there were no targets presented during this portion of the trial. We found that this was the case; we did not observe any predictive relationship between gain estimated in this period and behavioral performance (Supplementary Figure 8b-d; Supplementary Table 1). In summary, we found that cortical gain was modulated by both stimulus contrast and trial period, increasing when contrast is low and when mice were detecting targets. Furthermore, we found that behavioral performance was predicted by both the stimulus contrast and by session-to-session changes in cortical gain during target detection.

Figure 6. Cortical gain predicts session-to-session variability in behavioral performance.

a, Schematic of the LN model fit to neuronal responses during the behavioral task. **b,** Spike raster of a representative unit recorded during the behavioral task, sorted by the background scene presented in each trial. Below is the average spike rate (black trace) overlaid with the LN model fit with a static (grey) or gain-controlled nonlinearity (green). **c,** STRF for the representative cell. **d,** The fitted nonlinearities in low and high contrast periods of the trials. **e,** Gain distributions across all recorded neurons as a function of contrast and trial period (A: adaptation, T: target). Gain was significantly larger in the target period during low contrast (two-way ANOVA: p = 3.77e-9) but not in high contrast (p = 0.18) Inset: the average gain in low and high contrast for all cells, dashed lines indicate the median of each distribution. Gain was higher in low contrast (two-way ranksum: p = 1.60e-91. **f,** Psychometric curves split by the median gain (n=107 sessions). Light colored dots indicate performance across sessions with low gain, dark colored dots indicate performance on sessions with high gain. Error bars indicate ±SEM. *Inset*: the distribution of gain values on the same sessions. The dashed red line indicates the median used to split the data. **g,** Relationship between session-to-session changes in gain and behavioral thresholds (n=124 sessions). Each dot is a session, with the color indicating the contrast in which targets were presented. The grey line is the linear best fit. Black asterisks indicate whether gain is a significant predictor of the psychometric threshold, while magenta asterisks indicate whether contrast was a significant predictor of behavioral thresholds (Supplementary Table 1). **h,** Same formatting as g, but plotting the relationship between gain and psychometric slope (n=124 sessions). Source data are provided as a Source Data file.

413
414

Discussion

Auditory environments exhibit complex statistical properties that change over time. Changes in the dynamic range, or contrast, of acoustic inputs poses a challenge to the auditory system, which is composed of neurons with limited dynamic range. The efficient coding hypothesis predicts that as stimulus contrast changes, neurons should adjust their gain in order to match their limited dynamic range to that of the stimulus distribution[1]. Multiple studies have demonstrated that neurons throughout the auditory pathway exhibit such contrast gain control[14,16,19]. Additionally, contrast gain control is theoretically beneficial for separating signals from background noise[16,20], implicating that this phenomenon should underlie perception in noisy environments. Whereas recent work has demonstrated a link between contrast gain control and changes in behavioral discrimination in humans[19,21], it is unclear how cortical gain directly relates to behavior, as neuronal responses and behavior were not observed simultaneously. Finally, recent work has highlighted the dynamic nature of efficient neuronal codes[22,23], and it is unclear whether the observed neuronal dynamics reflect what is predicted to be theoretically efficient, and how the observed dynamics affect behavior.

Our study addressed these open questions by developing a normative framework to predict how efficient gain adaptation should affect behavior, and then testing those predictions using a combination of behavior, chronic recordings, and cortical manipulations. First, we developed a normative model based on efficient coding[22,23], which predicted that: 1) Detection thresholds of targets should be lower in low contrast than in high contrast; 2) Sensitivity to changes in target level should be greater in low contrast relative to high contrast; and 3) Detection should change asymmetrically over time: increasing slowly after a switch to low contrast, but decreasing rapidly after a switch to high contrast (Figure 1). Then, we derived a form of Poisson GLM to confirm that gain control dynamics in auditory cortex are indeed asymmetric (Figure 2). To behaviorally test the predictions of the normative model and GLM, we trained mice to detect a target embedded in dynamic random chords while manipulating the contrast of the background between high and low contrast. As predicted by the normative model, mice had lower detection thresholds and were more sensitive to changes in target level in low contrast. Behavioral adaptation was also asymmetric, decreasing rapidly after a switch to high contrast, and increasing slowly after a switch to low contrast, in agreement with our model (Figure 3). Furthermore, we found that AC is necessary for this detection-in-background task (Figure 4). When recording in AC, we found that the variability in neuronal population responses significantly predicted session-by-session variability in behavior above and beyond variability induced by stimulus features, and we demonstrated that target discriminability adapted asymmetrically, as predicted (Figure 5). Finally, we found that the amount of cortical gain also predicted behavioral performance on a session-to-session basis, independently of the effect of contrast (Figure 6). Taken together, these results support our hypothesis that the dynamics of gain control support efficient coding in auditory cortex and predict the perception of targets embedded in background sounds.

The role of cortex in behavior.

The role of auditory cortex in behavior has been subject of debate. A number of prior studies found that auditory cortex was not required for relatively simple behavioral tasks such as frequency discrimination or detection[29,30]. Rather, many studies found that auditory cortex is primarily involved in more complex behaviors, such those requiring temporal expectation[31], localization[32], or discrimination of more complex sounds[33–35]. Consistent with previous findings[36], we found that AC inactivation selectively impaired detection of targets in a noisy background, but did not impair detection of targets in silence (Figure 4). Furthermore, neuronal activity in AC predicted variability in behavioral performance (Figures 5, 6). This set of results establishes that AC is necessary for the detection of targets in background noise and supports the more general notion that AC is required for more difficult auditory tasks.

While the previous work demonstrates the necessity of auditory cortex in behavioral performance, the brain areas and mechanisms supporting the transformation from stimulus to decision are an active field of study[37,38]. By recording neuronal activity during the task, we were able to leverage behavioral variability to show that task performance covaried with representations of targets within small neuronal populations (Figure 5), and with cortical gain (Figure 6). There is a large body of literature relating cortical codes to behavioral variability: early studies in the visual system suggested that information from relatively small numbers of neurons was sufficient to match or outperform animal behavior in psychophysical tasks[39–41] and that behavioral choice can be predicted from activity in sensory areas[28,41]. These accounts suggested that variability in bottom-up sensory encoding drives the variability in behavioral output. However, more recent work suggests that variability in sensory areas is driven by top-down influences[42–45], which are modulated by attention and learning[46–49]. A recent study imaging tens of thousands of neurons in the visual cortex supports

this notion, finding that cortical representations had higher acuity than behaving mice, yet did not correlate
with behavioral performance, suggesting that perceptual discrimination depends on post-sensory brain
regions[50].
Our results suggest that bottom-up adaptation to stimulus statistics shapes behavioral output: We
observed asymmetric time courses of target discrimination following a change in contrast (Figure 3). The
asymmetric adaptation in behavioral performance was consistent with predictions derived from a normative
account of contrast gain control (Figure 1), resembled contrast gain adaptation in auditory cortex in the
absence of behavior (Figure 2), and was evident in patterns of target-driven activity in auditory cortex during
task performance (Figure 5). Indeed, there have been other studies demonstrating that individual differences
in sensory-guided behaviors are reflected in cortical activity[51,52], are bidirectionally modulated by cortical
manipulation[53,54], and can be predicted from tuning properties in auditory cortex[55,56]. While our results cannot
rule out top-down input as the causal driver of sensory decisions, they do support the notion that the sensory
information upon which decisions are made is shaped by neuronal adaptation, which thereby affects
behavioral outcomes.
485
Roles of gain in the auditory system.
Neurons throughout the auditory system adapt to the statistics of the acoustic environment, including
the frequency of stimuli over time[57,58], more complex sound patterns[24,59], and task-related or rewarded
stimuli[60–65]. In this study, we focused on contrast gain control as a fundamental statistical adaptation that
relates to efficient coding[14,17–19]. Previous work identified that responses to targets presented after opposing
changes in contrast were asymmetric[14], a result consistent with optimal estimation of stimulus variance[25]. We
observed similar asymmetric dynamics in a normative model of contrast gain control (Figure 1), which
suggests that previous observations of asymmetric adaptation are a result of efficient encoding of the
background stimulus.
Inspired by this and other previous work[60], we intentionally designed our stimuli using unbiased white-
noise backgrounds, which allowed us to fit encoding models to our data. In this study, we developed an
application of Poisson GLM that allowed us to quantify gain dynamics directly from responses to continuous
DRC stimuli. This approach allowed us to verify that gain adaptation in auditory cortex is asymmetric (Figure
2), and we found that this model better predicted neuronal activity when compared to previous models (Figure
2, Supplementary Figure 3). While we focused on the multiplicative influence of contrast in this study, the
GLM presented here could in theory be applied to any other time-varying signal that modulates neuronal gain,
such as movement[66,67], arousal[68,69], or experimental interventions such as optogenetics[70–73]
Additionally, previous studies found that contrast gain control was predictive of behavioral
performance[10,19]. Here, we extended these findings by examining the dynamics of this process and found that
behavioral detection of targets also adapted asymmetrically (Figure 3). In addition to confirming previously
reported effects of contrast on psychometric curves, these results suggest that the dynamics of contrast gain
control influenced task performance. Pursuing this further, we estimated cortical gain as mice performed the
task, and discovered a predictive relationship between stimulus contrast, gain in auditory cortex, and
behavioral performance (Figure 6). These results suggest two sources of cortical gain modulation: 1) Bottom-
up adaptation to stimulus contrast (ie. contrast gain control), and 2) session-to-session modulation of gain.
Previous studies have demonstrated this latter phenomenon, suggesting that top-down gain modulation
underlies attention[42,43,74] and the maintenance of optimal behavioral states[68,69]. Our results suggest that both
contrast gain control and session-to-session fluctuations in gain modulate behavior, providing a starting point
for dissecting the neuronal mechanisms underlying these two forms of gain modulation.
515
Cellular mechanisms of gain control.
While this work and other studies have established contrast gain control as a fundamental property of
the auditory system, the neuronal mechanisms driving gain adaptation at a cellular level remain unclear.
Previous work theorized that cortical feedback was responsible for contrast gain control in early areas[16,20], but
recent experiments disproved this hypothesis[19]. In the current study, we have likely recorded from a mixed
population of excitatory and inhibitory neurons, the latter of which exhibit specific roles in adaptation[75,76]. While
specific inhibitory neuronal subtypes facilitate divisive or subtractive control of excitatory responses in
visual[70,71] and auditory cortex[72,73], the role of these interneurons in contrast gain control has been
inconclusive[18]. Additionally, our results highlight the distinction between stimulus driven gain control and
stimulus-invariant gain control which covaried with behavior from session to session (Figure 6). Whether these
two forms of gain control share common neuronal substrates is unclear. By combining cell-specific

optogenetic methods with behavioral tasks, future studies may explore and test the causal role of local circuits
and top-down modulation in gain control and behavior.
529

530   *The missing link between efficient coding and behavior.*
531   Combined, our results develop a framework and provide support for the role of contrast gain control
in behavior. The efficient coding hypothesis has emerged as one of the leading principles in computational
neuroscience that has shaped our understanding of neuronal coding, architecture, and evolution[1,77–80], and
prior research found that human behavior follows principles of efficiency[19,81]. Here, we focused on a well-
studied form of efficient coding, contrast gain control, and developed a framework to link the dynamics of
efficient neuronal coding with behavioral performance. While the mechanisms of contrast gain control in
auditory cortex remain unclear, this study highlights potential top-down and bottom-up influences on cortical
gain, which may or may not share common neuronal substrates. We believe that the theoretical frameworks
and modelling methods presented here will be broadly applicable in future studies of neuronal gain control, a
fundamental function of the nervous system.
541
542

Methods

Animals.

All experiments were performed in adult male (n = 19) and female (n = 19) C57BL/6 (Stock No. 000664) or B6.CAST-*Cdh23^{Ahl/+}* (Stock No. 002756) mice (The Jackson Laboratory; age 12-15 weeks; weight 20-30g). Some of the mice used in these experiments were crossed with other cell-type specific -cre lines, as detailed in Supplementary Table 2. All mice were housed with, at most, five mice per cage, at 28°C on a 12-h light:dark cycle with food provided ad libitum, and a restricted water schedule (see *Water restriction*). All experiments were performed during the animals' dark cycle. All experimental procedures were in accordance with NIH guidelines and approved by the Institutional Animal Care and Use Committee at the University of Pennsylvania.

Data availability.

All data including spike times from electrophysiological recordings is available on DRYAD: https://doi.org/10.5061/dryad.6djh9w120.

Surgery.

Mice were anesthetized under isoflurane (1-3%). Prior to implantation, all mice were administered subcutaneous doses of buprenorphine (Buprenex, 0.05-0.1 mg/kg) for analgesia, dexamethasone (0.2 mg/kg) to reduce brain swelling, and bupivicane (2 mg/kg) for local anesthesia. In mice implanted with microdrives, two ground screws attached to ground wires were implanted in the left frontal lobe and right cerebellum, with an additional skull screw implanted over the left cerebellum to provide additional support. A small craniotomy was performed over the target stereotactic coordinates relative to bregma, -2.6mm anterior, -4.3mm lateral. Either custom 16-channel microdrives, 32-, or 64-channel shuttle drives (cite) holding tetrode bundles of polyimide-coated nichrome wires were chronically implanted over auditory cortex, and tetrodes were lowered 800um below the pial surface. The exposed tetrodes were covered with GelFoam (Pfizer) or sterile silicone lubricant and sealed with Kwik-Cast (World Precision Instruments). The plastic body of the microdrive and a custom stainless-steel headplate were secured to the skull using dental cement (C&B Metabond) and acrylic (Lang Dental). Mice undergoing only behavioral experiments were implanted with two skull screws in the cerebellum, and a headplate was mounted on the skull as previously described. An antibiotic (Baytril, 5mg/kg) and analgesic (Meloxicam, 5mg/kg) were administered daily (for 3 days) during recovery.

Water restriction.

Following surgical recovery (3 days post-operation), each mouse's weight was monitored for three additional days to establish a baseline weight. Over the next seven days, mice were water deprived, beginning with a daily ration of 120µL/g and gradually decreasing their ration to 40-50µL/g. During the task, if mice did not receive their full ration, the remainder of their ration was provided in their home cage. Mouse weight relative to baseline was monitored during all stages of water restriction. Additional health signs were used to determine a health score and subsequent treatment plan if a mouse lost more than 20% of baseline weight[82] as approved by the Institutional Animal Care and Use Committee at the University of Pennsylvania.

Behavioral apparatus.

During the GO/NO-GO task, the mouse was head-fixed in a custom-built, acoustically isolated chamber. A capacitive touch sensor (AT42QT1010, SparkFun) soldered to a lick spout monitored lick activity. Water rewards were dispensed from a gravity fed reservoir, controlled by a solenoid valve (161T011, Neptune Research) calibrated to deliver approximately 4-5µL of water per reward[83]. Low-level task logic – such as lick detection, reward and timeout delivery, and task timing intervals – was directly controlled by an Arduino Uno microprocessor running custom, low-latency software routines. High-level task logic, such as trial randomization, stimulus buffering and presentation, and online data collection and analysis were controlled by custom MATLAB (r2019a, Mathworks) software communicating with the Arduino over a USB serial port. Acoustic waveforms were generated in MATLAB and converted to analog signals via a soundcard (Lynx E44, Lynx Studio Technology, Inc.) or a National Instruments card (NI PCIe-6353) and delivered through an ultrasonic transducer (MCPCT-G5100-4139, Multicomp). The transducer was calibrated to have a flat frequency response between 3 kHz and 80 kHz using a 1/4-inch condenser microphone (Brüel & Kjær) positioned at the expected location of the mouse's ear[84,85]. During electrophysiological recording sessions,

598	licks were detected using an optical interrupt sensor (EE-SX771, Omron Automation), to prevent lick-related
599	electrical artifacts introduced by contact with a capacitive sensor.
600	

Behavioral timeline.
Each mouse underwent four stages in the behavioral task: 1) water restriction and habituation, 2)
behavioral training, 3) psychometric testing, and, 4) offset testing. During the induction of water restriction,
mice were habituated to head-fixation in the behavioral chambers and received water through the lick spout,
getting a drop of water for licks separated by more than 2 s. After the mouse began to receive its entire ration
by licking in the booth, behavioral training was initiated (typically after 1 week). Each mouse was initially
trained and tested in one contrast condition (see *Stimuli*), with the initial training condition counterbalanced
across mice. Behavioral performance was monitored during training, and mice were considered trained after
completing at least three consecutive sessions with over 80% percent correct. After completing training,
behavioral thresholds were measured during at least three sessions in which psychometric stimuli were
presented (see *Stimuli*). After estimating the behavioral threshold for each mouse, offset stimulus sets were
generated using threshold-level targets. After completion of at least three sessions in the offset task, each
mouse was then retrained on the remaining contrast condition. Upon reaching the training criterion of 80% in
the new contrast condition, mice were then tested in the psychometric and offset tasks as previously
described. For mice in electrophysiological experiments, this sequence of training and testing was continued
until the recording site yielded less than three units, or until the mouse stopped performing in the task.

617
Stimuli.
All stimuli were created in MATLAB and sampled at 192 kHz or 200 kHz and 32-bit resolution. A set
of dynamic random chords (DRCs) were created with different contrasts, similarly to those described in
previous studies[14,17,19]. To construct a DRC, amplitude modulated pure tones were generated at multiple
frequencies and then superimposed to create a chord. In some experiments, 34 frequencies were sampled
between 4 and ~40kHz in 1/10 octave steps, in the remaining experiments, 33 frequencies were sampled
between 4 and 64kHz in 1/8 octave steps. The amplitude envelope of each tone was generated as follows:
every 25 ms, amplitudes for each frequency were sampled from a uniform distribution with a mean of 50 dB
and a width of ±5 dB in low contrast or ±15 dB in high contrast. Between each 20 ms chord, the amplitude
envelope of each frequency band was linearly ramped over 5 ms to the amplitude value for the next chord,
such that the total duration of each chord and its ramp was 25 ms. To synthesize the stimuli, amplitude
envelopes were multiplied by a sine wave of their respective frequencies, and summed to produce the final
waveform. Each time a set of DRCs was generated, 5 unique random number generator seeds were used to
restrict the background noise to 5 distinct scenes (see raster in Figure 6 for an example of spike-locking to
the repeated scenes).
In all stages of behavioral training and testing, stimuli created for each trial consisted of a DRC
background containing a change in contrast, and the presence or lack of a target at a delay after the change
in contrast. Each trial began with 3 seconds of DRC background from one contrast, followed by a switch to
the other contrast. Targets consisted of a fixed chord composed of 17 frequencies pseudo-randomly sampled
from the frequencies contained in the DRC background, such that the target frequencies were uniformly
distributed across the frequency range of the background. To add targets to the background noise, the target
amplitude at each target frequency was simply added to a single chord in the amplitude envelope of the
background, and linearly ramped: this procedure ensured that target timing was perfectly aligned to changes
in the background noise, removing asynchronous timing cues that could be used to detect the target. Target
amplitudes are described in values of signal-to-noise ratio (SNR) relative to the average level of the
background noise (ie. a 50 dB target embedded in 50 dB background would have an SNR of 0 dB). See
Supplementary Table 3 for SNRs used for each mouse. In all trials, targets were embedded after a change in
the background contrast, with a delay and level dependent on the current training or testing stage.

646
Efficient coding model.
We simulated a model neuron that encodes incoming stimuli via an adapting neuronal nonlinearity.
Stimuli were drawn from a Gaussian distribution whose mean μ was fixed over time but whose standard
deviation σ_t could switch over time between a low and a high value ($\sigma_t = \sigma^L$ and $\sigma_t = \sigma^H$, respectively). At
each time t, a stimulus s_t was drawn from the distribution $p(s_t|\sigma_t) = \mathcal{N}(s_t; \mu, \sigma_t^2)$, transformed via a
saturating nonlinearity of the form $1/\left(1 + e^{-k(s_t - s_0)}\right)$, distorted by Gaussian noise with variance σ_n^2, and
finally discretized into N discrete levels to generate a response r_t. This discrete response was linearly

decoded to extract an estimate $\hat{s}_t$ of the current stimulus: $\hat{s}_t = p_1 r_t + p_0$. The recent history of L stimulus
estimates was used to update an estimate $\hat{\sigma}_t$ of the underlying standard deviation: $\hat{\sigma}_t = \text{std}(\hat{s}_{t-L+1}:\hat{s}_t)$. The
estimate $\hat{\sigma}_t$ was then used to select the parameters of the encoder (k, s_0) and the decoder (p_1, p_0) on the
next timestep. The encoding and decoding parameters were chosen to minimize the expected error in
decoding stimuli given the neuron's current estimate of the underlying standard deviation:
$k, s_0, p_1, p_0 = \text{argmin}\langle(\hat{s}_t - s_t)^2\rangle_{p(s_t|\hat{\sigma}_t)}$[22,23].

The parameters of the encoder and decoder were adapted based on a background stimulus with a
mean μ_B that was fixed over time and a standard deviation σ_B that switched between low and high values
σ_B^L and σ_B^H, respectively. We used this adapting nonlinearity to determine how well this model neuron could
discriminate target stimuli from this background. Target stimuli were sampled from a Gaussian distribution
with a fixed mean μ_T and with a variance σ_T that was scaled in proportion to the variance of the background
$(\sigma_T^L = f\sigma_B^L$ and $\sigma_T^H = f\sigma_B^H$, respectively). At each timestep, we computed the Bhattacharyya coefficient (BC)
of the response distributions produced by background versus target stimuli: $BC = \sum \sqrt{p(r_B)p(r_T)}$. We used
$1 - BC$ as our measure of discriminability.

We simulated the behavior of this model using a background "probe" stimulus whose standard
deviation switched every T timesteps. We simulated N_c cycles of this probe stimulus, where each cycle
consisted of T timesteps in the low state, followed by T timesteps in the high state. This yielded timeseries
of the gain k and offset s_0 of the adapting nonlinearity, as well as distributions of the neuronal response to
the background and target stimuli at each timepoint following a switch in standard deviation. We averaged
the gain and offset across cycles to obtain the average properties of the encoder at each timepoint following
a switch. We used the distribution of responses to target and background stimuli, measured across cycles,
to compute the discriminability at each timepoint following a switch. All simulations were performed with the
following values: $T = 50$, $N_c = 1,000$, $\mu_B = 0$, $\mu_T = 0$ to 3 in 0.25 steps, $\sigma_B^L = 1$, $\sigma_B^H = 3$, $f = 0.25$, $\sigma_n^2 =$
0.01, $N = 15$, $L = 12$. For Figure 1g, model discriminability in each contrast was fit with a logistic function to
estimate the sensitivity and threshold of the model. To approximate the stimulus conditions used in the
offset task, the target thresholds for each contrast were then used to select target levels to plot
discriminability over time ($\mu_T^L = 1.50$, $\mu_T^H = 2.25$; Figure 1f).
To compare behavioral performance to the model, we quantified the "psychometric performance" of
the model by fitting model discriminability over time with a logistic function:
683

$$y = \gamma + (1 - \gamma - \lambda) * \frac{1}{1 + e^{\alpha - \beta x}} \tag{1}$$

684
where α is the x-offset of the function, β determined the sensitivity of the function, γ determined the guess
rate (lower bound), λ determined the lapse rate (upper bound) and x was stimulus level. α/β determined the
threshold of this function, defined as the level corresponding to the steepest part of the curve. This function
was fit to behavioral or neuronal performance using constrained gradient descent ($fmincon$ in MATLAB)
initialized with a 10x10 grid-search of parameters α and β.
To characterize adaptation time constants, adaptation curves were fit with an exponential function

$$y = a + b * e^{-\frac{t}{\tau}} \tag{2}$$

691
692   where a determined the y-offset of the function, b was a multiplicative scaling factor, and τ was the
693 time constant of the exponential in units of time t. This function was fit to behavioral or neuronal responses
694 using constrained gradient descent initialized with a 10x10x10 grid search across all three parameters.
695   To measure the effect of stimulus contrast on parameters of interest, we computed a contrast
696 modulation index (CMI), which measured the relative change in the parameter between low and high contrast:
697

$$CMI = \frac{x_H - x_L}{x_L} \tag{3}$$

698
699 where x is the parameter in question and subscripts H and L denote the value of that parameter in high and
700 low contrast, respectively. The resulting index is 0 if there is no change, 1 if x_H is two times larger than x_L,
701 and is -0.5 if x_L is two times x_H.
702

703
704 *Behavioral task.*
705    We used a GO/NO-GO task to measure the detectability of targets in background. In this task, each
trial consisted of a noise background with a contrast shift, along with the presence or absence of a target after
the change in contrast. Mice were trained to lick when they detected a target (hit), or to withhold licking in the
absence of a target (correct reject). This behavior was reinforced by providing a 4-5 µL water reward when
the mouse licked correctly (hit), and by initiating a 7-10 s timeout when the mouse licked in the absence of a
target (false alarm). Any licks detected during the timeout period resulted in the timeout being reset. In a
subset of mice, we introduced an additional trial abort period coincident with the first part of the contrast
background, before the contrast switch. Any licks detected in this abort period resulted in the trial being
repeated after a 7-10 s timeout, until the mouse withheld from licking during this period. In this task, misses
and correct rejects were not rewarded or punished. Trials were separated by a minimum 1.5s inter-trial-interval
(ITI). To discourage spontaneous licking, licks were monitored during this period, and if any licks occurred the
ITI timer was reset.
717    To prevent mice from predicting the target, we varied the timing of the target relative to the contrast
shift. This required a method for estimating hit rates and false alarm rates at different times during each trial,
and to reward and punish the animal during these times in an unbiased manner. To approach this issue, we
considered licks only during a 1 s response window after a target presentation (eg. if a target was presented
500 ms post-contrast-switch, the response window persisted from 500 to 1500 ms post-contrast-switch). To
apply this method to background-only trials, in which no targets were presented, we considered background
trials to be target trials containing infinitely small target amplitudes. For each background trial, we assigned a
response window with equiprobable delay matched to the target conditions and considered only licks within
those "target" response windows. Thus, over the course of a session, we randomly sampled lick probabilities
in background trials during the same temporal windows as those considered during target trials. Using this
scheme, we treated target and background-only trials identically, and estimated hit rates and false alarm rates
over time in an unbiased manner.
729    Each mouse performed three stages in the behavioral task: training, psychometric testing, and offset
testing. During the training task, trials consisted of two types, background-only trials or target trials presented
with equal probability. To facilitate learning, we selected target SNRs at the highest end of the range described
previously: in low contrast training sessions, targets were 16 dB SNR, and in high contrast training sessions,
targets were 20 dB SNR. To prevent response bias as a function of target timing, we randomly varied the
target delay between 250, 500, 750 and 1000 ms after the contrast change in each trial. During the
psychometric testing task, there were 7 trial types consisting of background-only trials and target trials
spanning six different SNRs (Supplementary Table 3). Based on behavioral piloting, we presented high SNR
trials with a greater probability, to ensure that mice were consistently rewarded during the task. In low and
high contrast psychometric sessions, the probability of a background trial was 0.4, the probability of the four
lowest target SNRs was 0.05 each, and the probability of the two highest target SNRs was 0.2 each. As in
training, target timing was varied randomly between 250, 500, 750 and 1000 ms after the contrast change in
each trial. After completing at least three sessions of the psychometric task, stimuli were generated for the
offset testing task. This task consisted of 15 unique trial types: 3 target levels (background trials, threshold
target trials, and high SNR target trials), and 5 target delays relative to the contrast change (25, 75, 225, 475,
975 ms delay). Threshold target amplitudes were determined individually for each mouse by fitting
performance averaged over several sessions with a psychometric function, and extracting the level at which
the slope of the psychometric curve was steepest. Based on behavioral piloting, background trials, threshold
target trials, and high SNR target trials were presented with probabilities of 0.4, 0.2, and 0.4, respectively.
Target delay on each trial was selected with equal probability. In all behavioral stages, trial order was
pseudorandomly generated, such that there were no more than three target or background trials in a row.
750    A subset of mice (n = 2), were presented targets in silence (Figure 4). To generate this stimulus set
without changing the basic structure of the task or stimuli, we simply took the spectrograms of all stimuli
containing 25 dB SNR targets from the low-to-high contrast stimulus sessions, and set the stimulus power
flanking each target to zero. This manipulation was only performed in the target period, and the low contrast
adaptation period of the trials remained the same. Thus, the targets and adaptation periods were identical to
those presented in the target-in-background task. To vary the difficulty of the task, the level of the target was
attenuated using the following values: -75, -60, -45, -30, -15, and 0 dB attenuation relative to the 25 dB SNR
target. Mice were previously trained in the target-in-background task prior to performing the target in silence
task. Before psychometrically varying the target attenuation, mice were trained in the new task to criterion

performance. Mice generalized very rapidly to the new task, reaching 97% and 94% training accuracy on the
first day of exposure to targets in silence (mice CA124 and CA125, respectively).
761
Chronic muscimol application.
A separate cohort of mice (n = 4) were bilaterally implanted with 26 GA guide cannulae (PlasticsOne,
C315GMN-SPC mini, cut 5 mm below pedestal) in auditory cortex. The surgery was performed as described
above with the following modifications. After the skull was leveled using a stereotax, two small craniotomies
were performed -2.6 mm anterior, ±4.3 mm lateral from bregma, over auditory cortex. The guide cannulae
and dummy infusion cannulae (PlasticsOne, C315DCMN-SPC mini, cut to fit 5 mm C315GMN with a 0.5 mm
projection depth) were sterilized in an autoclave. The dummy cannulae were partially screwed into the guide
cannulae and placed in a stereotaxic clamp. After zeroing the tip of the guide cannula to the brain surface,
the cannula was lowered to 500 µm below the cortical surface. This depth was chosen because the infusion
cannulae (PlasticsOne, C315LIMN-SPC mini) project 500 µm from the end of the guide cannulae when
completely inserted, leading to a final depth of 1000 µm – the location of auditory cortex. The dummy cannulae
were then fully inserted and this procedure was repeated for the next cortical hemisphere.
Prior to injecting, two injection syringes (Hamilton Syringe, 10µL Gaslight #1701) and tubing (C313CT
tubing 023x050 PE50) were backfilled with mineral oil. Sterilized infusion cannulae were then attached to
each syringe and ~500nL of muscimol (diluted with 1x PBS to .25 mg/mL; Sigma Aldrich, M1523) or 0.9%
sterile saline was drawn up into the injection cannulae using a dual injector (Harvard Apparatus, Pump 11
Pico Plus Elite). The mouse was then headfixed and the dummy cannulae were removed and sterilized. The
loaded infusion cannulae were then screwed all the way into the guide cannulae and 400 nL of muscimol or
saline was infused bilaterally at a rate of 250 nL/minute. The infusion cannulae were then replaced with the
dummy cannulae and the mouse rested in its home cage for 30-45 minutes before beginning the behavioral
session.
783
Acute electrophysiological recordings.
For acute recordings used to fit the GC-GLM model (Figure 2), neuronal signals were recorded from
n = 1 awake, untrained mouse. Prior to the recording session, the mouse was anesthetized and a headpost
and ground pin were implanted on the skull (see *Surgery*). On the day of the recording, the mouse was briefly
anesthetized with 3% isoflurane and a small craniotomy was performed over auditory cortex using a dental
drill or scalpel (~1 mm x 1 mm craniotomy centered approximately 1.25 mm anterior to the lambdoid suture
along caudal end of the squamosal suture). A 32 channel silicon probe (Neuronexus) was then positioned
perpendicularly to the cortical surface and lowered at a rate of 1-2 µm/s to a final depth of 800-1200 µm. As
the probe was lowered, trains of brief noise bursts were repeated, and if stimulus locked responses to the
noise bursts were observed, the probe was determined to be in auditory cortex. The probe was then allowed
to settle for up to 30 minutes before starting the recording. Neuronal signals were amplified and digitized with
an Intan headstage (RHD 32ch) and recorded by an openEphys acquisition board[86] at a rate of 30 kHz.
For this experiment, the mouse was presented with 3 s DRCs alternating between low and high
contrast (uniform distribution with a mean of 50 dB and a width of ±5 dB in low contrast or ±15 dB in high
contrast at a chord rate of 25 ms, as described in *Stimuli*). In order to accurately fit the GLM in an unbiased
manner, these stimuli were highly random, composed of 100 unique chord patterns for each contrast
(Supplementary Figure 2i,j**).** For each of the two recording sites, 5 repeats of this stimulus set were played.
801
Behavioral electrophysiological recordings.
Neuronal signals were acquired from awake, behaving mice as they performed the psychometric and
offset testing tasks described previously. Chronically implanted, 16-, 32-, or 64-channel microdrives[86,87] were
connected to one or two 32 channel Intan amplifier headstages. Amplified signals were recorded at 30 kHz
using an openEphys acquisition board via an SPI cable, where the signals were digitized.
For all recordings, broadband signals were filtered between 500 and 6000 Hz, offset corrected, and
re-referenced to the median across all active channels. The preprocessed data was then sorted using
KiloSort[88] or KiloSort2 and the resulting clustering was manually corrected in phy2 according to community-
developed guidelines. The resulting units were labelled as single units if they exhibited a clear refractory
period and did not need to be split. Splitting assessments were made through manual examination of principle
component features for the two best channels of a cluster. If two noticeable clusters in feature space were
evident in a unit, the unit was either manually split, or classified as a multiunit.
814

Generalized linear model.
To justify the form of GLM used here, we discuss a how a model neuron could implement gain control
in the simplest terms, and then structure our inference model to extract the parameters of this model neuron.
We will assume that the activity of the model neuron is driven by three sources: 1) stimulus drive, 2) stimulus
contrast, and 3) the multiplicative interaction between the two, which we use to define the gain (for a formal
definition of this forward model and the inference model, see *Supplementary Information*).
As discussed previously, the stimulus used in our experiments is composed of many frequencies that change
in loudness in discrete time steps:
823

$$X_{t,f} \sim \mathcal{U}(\mu,\, \sigma_t)$$

825
826 where $X_{t,f}$ is the stimulus spectrogram that varies as a function of time t and frequency f. Each time and
827 frequency bin of X is sampled from a uniform distribution defined by an average value μ and contrast σ_t.
828   We assume that the hypothetical neuron responds selectively at some frequency and time lag, defined
829 by a filter, or STRF $\beta_{h,f}$ with history h and frequency f components. Given β, we can define the stimulus drive
830 x_t as

$$x_t = X_t \beta \tag{4}$$

831
where at each time t, X_t is a row vector of size F frequencies times H lags (ie. the "unrolled" lagged stimulus
spectrogram) and β is the STRF unrolled to a single column vector of the same size.
In the spirit of efficient coding theory, and as shown in previous work, we assume that the gain g of the neuron
should be inversely proportional to the contrast, such that $g(\sigma) \propto 1/\sigma$ (ie. when contrast is low gain should be
high, and vice-versa). We also define "neutral" gain to be the average of the gain of the neuron in low and
high contrast. Putting these two features together, we can summarize the gain of the neuron as
838

$$g(\sigma) = \frac{\overline{\sigma}}{\sigma_t} \tag{5}$$

839
where $\overline{\sigma}$ is the harmonic mean of the contrast in the low and high conditions (see *Supplementary Information*).
In the case of a 3-fold change in contrast, this function constrains the gain of the neuron between 0.5 and 1.5,
with a neutral value of 1. As mentioned previously, we consider gain to be the multiplicative interaction
between the stimulus drive and the contrast, such that the contribution of gain control to the response of the
neuron is related to $x_t \cdot \frac{\overline{\sigma}}{\sigma_t}$.
To summarize, we considered a hypothetical neuron driven by the stimulus according to a STRF β
and by the interaction between the stimulus drive and the contrast $x_t \cdot \frac{\overline{\sigma}}{\sigma_t}$. To infer the relative weights of each
of these components of the neuronal response, we defined a Poisson GLM with an intercept term and the
following predictors:

$$x_t, \qquad \frac{\overline{\sigma}}{\sigma_t}, \qquad x_t \cdot \frac{\overline{\sigma}}{\sigma_t} \tag{6}$$

849
850   In other words, the model is composed of a stimulus predictor x_t, a contrast predictor $\overline{\sigma}/\sigma_t$, and their
851 interaction. Therefore, the GLM models the firing rate λ at time t as a Poisson distribution with the following
852 mean:

$$\lambda_t = exp\left[\beta_0 + x_t \beta_1 + x_t \cdot \frac{\overline{\sigma}}{\sigma_t}\beta_2 + \frac{\overline{\sigma}}{\sigma_t}\beta_3\right] \tag{7}$$

853
854 where $\beta_0 \ldots \beta_3$ are the parameters to be inferred. Based on our behavioral data (Figure 3) and the predictions
855 of the efficient coding model (Figure 1), we expected the influence of contrast on neuronal gain to be
856 asymmetric and smooth. To enable the GLM to capture both of these qualities, we first defined the contrast
857 predictors from a set of cubic B-spline temporal basis functions, then defined separate contrast predictors for
858 transitions to low and high contrast. Incorporating these changes, we can redefine equation 4 above as

$$ln\lambda_t = \beta_0 + x_t\beta_1 + x_t \circ C'_t\beta_2 + C'_t\beta_3 \tag{8}$$

where ∘ denotes element-by-element "broadcasting" multiplication and C_t' is a matrix of contrast predictors
$\bar{\sigma}/\sigma_t$ convolved with a set of basis functions and separated by contrast transitions (see *Supplementary*
Information). For the sake of clarity, note that in the expression above, β_0 is a number, x is a column vector
of length T, β_1 is a number, C' is a T-by-$2B$ matrix, and β_2 and β_3 are column vectors of length $2B$, where B
is the number of splines.

So far, we outlined a hypothetical neuron which implements gain control, and a GLM with which we
can approximate the behavior of this neuron. Next, we describe how to use the fitted parameters to quantify
the gain of the neuron. Conceptually, an increase or decrease in the gain of the neuron is analogous to more
or less sensitivity to small changes in the stimulus. Based on this intuition, we focused on how the response
of the neuron (as modelled by a fitted GLM) is expected to change between conditions where the gain is
expected to contribute (ie. in the presence of gain control) and where it is not (ie. in the absence of gain
control, where gain is "neutral"). Following this logic, we derived a definition for gain w_t as the ratio between
the sensitivity of the fitted model with changes in contrast, compared to the sensitivity of the same model
when the contrast is at a reference value, which we defined previously as $w_t = 1$:

$$w_t = \frac{\beta_1 + C_t'\beta_2}{\beta_1 + C^0\beta_2} \tag{9}$$

875 where w_t is the estimated gain at time t, and C^0 is a reference contrast design matrix identical to C_t' except
876 that all non-zero elements are set to 1 (see *Supplementary Information* for full derivation of w_t).

877   To fit the model, we implemented a two-step procedure. In the first step, the STRF β of the neuron
878 was estimated according to the model

$$\ln \lambda_t = \alpha + X_t\beta \tag{10}$$

For the second step, we calculated the stimulus drive as described in equation 4, and then fit equation 8 to
the data for each neuron using glmfit in MATLAB (v2018a). This entire fitting procedure was 10-fold cross-
validated with folds stratified across trials of each contrast and run in parallel on a high performance computing
cluster using custom BASH scripts (v3.2.57(1)). In the first step, we fit the STRF β with F frequency bins
according to the stimulus spectrogram (F = 33 or 34, see *Stimuli*) and a history window of 300 ms (H = 12).
When fitting the full model, we defined the contrast design matrix C_t' to capture 1000 ms of contrast history
around each transition (H' = 40), convolved with a set of B-spline temporal basis functions[89] (here, we used
B-splines with a degree of 3 and 3 equally-spaced knots, constrained to go smoothly to zero at the longest
lag, which implied that B = 4).

To validate the model, we first simulated neurons according to the forward model outlined above
(Supplementary Figure 2a) while varying the amount of gain control and the temporal trajectory of gain in
different simulation runs. We found that the GLM accurately predicted the STRF shape, spike rates and gain
trajectories across a variety of simulation parameters (Supplementary Figure 2c, e-h). For a detailed
description and discussion of the simulation results, see *Supplementary Information* and Extended Data Table
4.

Behavioral and neuronal detection performance.
To calculate performance in the target-in-background detection task we adopted commonly used
signal detection theory methods[39,90] to estimate the ability of an ideal observer to discriminate between two
sensory distributions: in our case, a distribution for target trials and a distribution for background trials. When
analyzing behavior, we computed the percent correct performance of an ideal observer[91] as a function of the
probability of hits and false alarms:
902

$$pc = z^{-1}\left(\frac{z(H) - z(FA)}{\sqrt{2}}\right) \tag{11}$$

904 where z^{-1} is cumulative probability of the normal distribution (*normcdf* in MATLAB), z is the inverse of the
905 normal distribution (ie. the z-score, *norminv* in MATLAB), H is the hit rate, and FA is the false alarm rate. For
906 psychophysical performance, hit rates and false alarm rates near 0 and 1 were adjusted using the log-linear
907 rule[92], to reduce biases in performance estimation resulting from low numbers of trials.

To calculate neuronal performance in the same reference frame as the behavior, we employed similar ideal observer techniques. First, neuronal responses (either spike rates of single units, or population projection values), were averaged in a 100ms window post target onset (for background trials, this window was randomly chosen on each trial to coincide with target presentation times on target trials). Then, using the distributions of responses during target and background trials, we computed receiver-operating-characteristic curves and took the area under the curve (AUC) as the percent correct of an ideal observer discriminating between the target and background distributions. To determine whether the AUC value for a given set of trial distributions was significantly different from chance, we performed a bootstrap procedure where we sampled from all the trials with replacement 500 times and recomputed AUC for each sample. If the 95% confidence intervals for this bootstrapped distribution did not include chance (.5), we defined that AUC value as significant. For population analyses which generated single-trial predictions, neuronal hit and false alarm rates were transformed to percent correct as described above.

Population response metrics.
On sessions where three or more neurons were simultaneously recorded, we used a population vector technique[27] to estimate the ability of neuronal populations to discriminate targets from background. First, spike rates in each trial were averaged in a 100ms window post-target onset. Then, using a leave-one-out procedure, we computed a trial averaged population vector for target trials, v_T, and a separate average population vector for background trials, v_B. We then estimated the coding direction in high dimensional neuronal space that best separated the target and background responses: $CD = v_T - v_B$. The held out trial was then projected along this dimension, by taking the population response vector on that trial v_{trial} and projecting it along the estimated coding direction using the dot product: $projection\ value = v_{trial} * CD$. This procedure was repeated holding out each trial, and estimating the coding direction from the remaining trials. For psychometric testing sessions, the target responses from the two loudest target levels were used to estimate coding direction, and in offset testing sessions the target responses from the high SNR target trials were used. After computing projections for every trial, the resulting matrix was normalized between 0 and 1.

Population classifier.
Based on previously described methods[28], we used a criterion-based decision rule to estimate how a hypothetical down-stream neuron may read out the neuronal activity of a population of neurons. As before, trial distributions of neuronal responses to targets or background were created from the average activity in a 100ms window post-target. Then, we sampled 100 criterion values between the minimum and maximum response, and for each criterion estimated the proportion of correct trials under two decision rules: 1) report target present if the response is greater than the criterion, or, 2) report target present if the response is less than the criterion. By assessing these two decision rules, neurons that were suppressed by target presence were treated equally to neurons that were enhanced by target presence. Finally, we chose the criterion and decision rule that yielded the highest proportion of correct trials, and computed neuronal hit rates and false alarm rates for each target level, and background-only trials. These hit rates and false alarm rates were then transformed to percent correct according to Equation 8.

Linear-nonlinear model.
First, we selected only neurons in the dataset which had reliable responses to stimulus repeats. To determine response reliability, we computed a noise ratio (NR) for each neuron, which describes the amount of variability in the response due to noise versus the amount of variability in the response driven by the stimulus[93,94]. Values approaching 0 indicate increasingly reliable responses to the stimulus, so for the remaining analyses, we included neurons with NR < 100.
The linear nonlinear model was composed of a spectrotemporal receptive field (STRF) and a set of rectifying nonlinearities. The STRF β was fit using normalized reverse correlation

$$\beta = [XX^T]^{-1}X\lambda \tag{1}$$

where X is the stimulus design matrix X_t defined in equation 4 and λ is the spike count in each 25 ms bin of the DRC stimulus. When defining X, we used a history window of 300 ms ($H = 12$) and frequency bins corresponding to the frequencies composing the dynamic random chords (see *Stimuli*). After fitting the STRF,

we fit the nonlinearities of the neuron. This two-step fitting procedure was repeated using 10 fold cross-validation, as described below.

For each fold, we selected 90% of the trials for training, leaving the remaining 10% to be held out for testing. Within each trial, we excluded neuronal responses around transitions from silence, or transitions in contrast, to prevent the model from overfitting strong transients in the neuronal response. Additionally, we excluded neuronal responses within a 50ms window after target presentation, to prevent overfitting of target responses. Given these exclusion criteria, we calculated the duration of stimulus sampled in the target period for each trial, and, for each trial, sampled the same duration of stimulus within the adaptation period. This procedure ensured that the model was fit to the same amount of high and low contrast stimulation per trial, to minimize overfitting to one contrast condition. Then, a stimulus design matrix X was defined using these stimulus periods, and the STRF was fit using equation 12. We tested whether STRF properties were affected by stimulus contrast, and found STRFs to be largely stable when estimated separately for each contrast (*Supplementary Information* and Supplementary Figure 7). Therefore, we used both periods of contrast to estimate β.

Using the STRF fit to the training data, we computed the linear drive x_t by convolving the STRF with the lagged spectrogram of the training stimulus (equation 4). For the GC-LN model we separated the linear predictions into low and high contrast periods, while for the static-LN model all matched time points were used. We generated a histogram of the linear prediction values (50 bins), and for each bin, computed the mean spike rate of the neuron when the linear prediction fell within those bin edges (Figure 6d, scatter points). The resulting set of linear prediction values and average spike rates were fit with an exponential function:

$$y = a + be^{c(x-d)} \qquad (2)$$

where a determined the minimum firing rate, b was a multiplicative scaling factor, c determined the gain of the exponent, and d determined the x-offset, or firing threshold of the neuron. This function was fit to each cell using constrained gradient descent ($fmincon$ in MATLAB), using a 10x10 grid search for parameters b and c. The gain for each neuron was defined as c. This entire process was repeated for each cross-validation fold, and the final parameter estimates for the STRF and nonlinearities were taken as the average over the 10 runs.

To determine the relationship between neuronal gain and behavioral performance, we computed the average neuronal gain across all noise responsive neurons (NR < 100) in each session for the adaptation and target periods in the trial. We then compared the session-averaged gain values to the fitted thresholds and slopes of the psychometric curve across sessions using the mixed-effects linear models outlined in the main text.

Inclusion criteria.

Unless otherwise noted, behavioral sessions in which the false alarm rate exceeded 20% were discarded from analysis. One mouse (ID: CA122) had consistently high false alarm rates in the high contrast condition, so we excluded high contrast sessions from this mouse from all analyses. For Figures 5 and 6, we removed neurons with low spike rates (<1Hz) and noise-like or inverted (ie. upward inflected) spike waveforms. To determine waveform quality, we computed the width of each waveform at half of the minimum value (FWHM) and its correlation with the average waveform over all neurons. Neurons whose waveforms had outlier FWHM values ($isoutlier$ in MATLAB), negative correlations, or were not significantly correlated with the average (Bonferoni corrected $p > 5.85e-6$) were removed from further analysis. For Figure 5f-l, sessions with stable population decoding performance were included (defined as sessions where more than half of the target types elicited significant population AUC values, as determined by the bootstrap procedure described previously). For Figure 6e-h, only neurons with noise ratios less than 100 were included in all analyses.

Data Availability.

The data generated in this study, including sorted spike times from electrophysiological recordings and behavioral data, have been deposited on Dryad: https://doi.org/10.5061/dryad.6djh9w120. Every Figure and Supplementary Figure has associated data with the following exceptions. Figure 1, Supplementary Figure 1, and Supplementary Figure 2 are the results of simulations and thus have no associated raw data. Simulation parameters can be found in the code. There are no restrictions on data availability. A Source Data file has been provided with this manuscript.

Code Availability.

Code for behavioral and neural analysis: https://github.com/geffenlab/contrast_behavior. Code for the GC-GLM model and simulations: https://github.com/geffenlab/contrast_glm.

Acknowledgements.

This work was supported by NIDCD R01DC015527, R01DC014479, NINDS R01NS113241 to MNG and NIDCD F31DC016524 to CFA. AMW was supported by NIDCD T32DC016903. The authors would also like to thank Dr. Yale Cohen, Dr. Nicole Rust, and Dr. David Brainard for their helpful feedback on this study and Dashiell Ridolfi-Starr, Karmi Oxman, Nitay Caspi, Tyler Ling, and Margaret DenBleyker for experimental assistance.

Author Contributions Statement.

CFA and MNG conceived of the behavioral and neural recording experiments. WM and AMH conceived of and implemented the normative gain control model. EP and CFA conceived of and implemented the GC-GLM model simulations and fitting procedures. AMW performed the acute muscimol recordings (Supplementary Figure 5). CFA, KCW and LG created the behavioral apparatus and performed behavioral experiments. CFA performed the surgeries, electrophysiological behavior experiments and muscimol behavior experiments. KCW performed acute electrophysiology experiments (Figure 2). CFA analyzed the data. CFA and MNG wrote the paper.

Competing Interest Statement.

The authors declare no competing interests.

1.   Barlow, H. B. Possible Principles Underlying the Transformations of Sensory Messages. in *Sensory*
Communication vol. 6 216–234 (2013).
2.   Brenner, N., Bialek, W. & De Ruyter Van Steveninck, R. Adaptive rescaling maximizes information
transmission. *Neuron* **26**, 695–702 (2000).
3.   Bharioke, A. & Chklovskii, D. B. Automatic Adaptation to Fast Input Changes in a Time-Invariant
Neural Circuit. *PLoS Comput Biol* **11**, 1004315 (2015).
4.   Borst, A. & Theunissen, F. E. Information theory and neural coding. *Nature Neuroscience* vol. 2 947–
957 (1999).
5.   Baccus, S. A. & Meister, M. Fast and slow contrast adaptation in retinal circuitry. *Neuron* **36**, 909–919
(2002).
6.   Dean, I., Harper, N. S. & McAlpine, D. Neural population coding of sound level adapts to stimulus
statistics. *Nat. Neurosci.* **8**, 1684–1689 (2005).
7.   Lesica, N. A. *et al.* Adaptation to Stimulus Contrast and Correlations during Natural Visual
Stimulation. *Neuron* **55**, 479–491 (2007).
8.   Gutnisky, D. A. & Dragoi, V. Adaptive coding of visual information in neural populations. *Nature* **452**,
220–224 (2008).
9.   Wen, B., Wang, G. I., Dean, I. & Delgutte, B. Dynamic range adaptation to sound level statistics in the
auditory nerve. *J. Neurosci.* **29**, 13797–13808 (2009).
10.   Dahmen, J. C., Keating, P., Nodal, F. R., Schulz, A. L. & King, A. J. Adaptation to Stimulus Statistics
in the Perception and Neural Representation of Auditory Space. *Neuron* **66**, 937–948 (2010).
11.   Wen, B., Wang, G. I., Dean, I. & Delgutte, B. Time course of dynamic range adaptation in the auditory
nerve. *J. Neurophysiol.* **108**, 69–82 (2012).
12.   Clarke, S. E., Longtin, A. & Maler, L. Contrast coding in the electrosensory system: Parallels with
visual computation. *Nature Reviews Neuroscience* vol. 16 733–744 (2015).
13.   Clemens, J., Ozeri-Engelhard, N. & Murthy, M. Fast intensity adaptation enhances the encoding of
sound in Drosophila. *Nat. Commun.* **9**, 1–15 (2018).
14.   Rabinowitz, N. C., Willmore, B. D. B., Schnupp, J. W. H. & King, A. J. Contrast Gain Control in
Auditory Cortex. *Neuron* **70**, 1178–1191 (2011).
15.   Rabinowitz, N. C., Willmore, B. D. B. B., Schnupp, J. W. H. H. & King, A. J. Spectrotemporal Contrast
Kernels for Neurons in Primary Auditory Cortex. *J. Neurosci.* **32**, 11271–11284 (2012).
16.   Rabinowitz, N. C., Willmore, B. D. B., King, A. J. & Schnupp, J. W. H. Constructing Noise-Invariant
Representations of Sound in the Auditory Pathway. *PLoS Biol.* **11**, e1001710 (2013).
17.   Cooke, J. E., King, A. J., Willmore, B. D. B. & Schnupp, J. W. H. Contrast gain control in mouse
auditory cortex. *J. Neurophysiol.* **120**, 1872–1884 (2018).
18.   Cooke, J. E. *et al.* Contrast gain control occurs independently of both parvalbumin-positive
interneuron activity and shunting inhibition in auditory cortex. *J. Neurophysiol.* **123**, 1536–1551
(2020).
19.   Lohse, M., Bajo, V. M., King, A. J. & Willmore, B. D. B. Neural circuits underlying auditory contrast
gain control and their perceptual implications. *Nat. Commun.* **11**, 1–13 (2020).
20.   Willmore, B. D. B., Cooke, J. E. & King, A. J. Hearing in noisy environments: Noise invariance and
contrast gain control. *J. Physiol.* **592**, 3371–3381 (2014).
21.   Maier, J. K. *et al.* Adaptive coding is constrained to midline locations in a spatial listening task. *J.*
Neurophysiol. **108**, 1856–1868 (2012).
22.   Młynarski, W. F. & Hermundstad, A. M. Adaptive coding for dynamic sensory inference. *Elife* **7**,
(2018).
23.   Młynarski, W. F. & Hermundstad, A. M. Efficient and adaptive sensory codes. *Nat. Neurosci.* **24**, 998–
1009 (2021).
24.   Pennington, J. R. & David, S. V. Complementary effects of adaptation and gain control on sound
encoding in primary auditory cortex. *eNeuro* **7**, 1–17 (2020).
25.   DeWeese, M. & Zador, A. Asymmetric Dynamics in Optimal Variance Adaptation. *Neural Comput.* **10**,
1179–1202 (1998).
26.   Paxinos, G. & Franklin, K. B. J. *Paxinos and Franklin's the Mouse Brain in Stereotaxic Coordinates*.
Academic Press vol. 2nd Editio (Academic Press, 2001).
27.   Li, N., Daie, K., Svoboda, K. & Druckmann, S. Robust neuronal dynamics in premotor cortex during
motor planning. *Nature* **532**, 459–464 (2016).
28.   Christison-Lagay, K. L., Bennur, S. & Cohen, Y. E. Contribution of spiking activity in the primary

095   auditory cortex to detection in noise. *J. Neurophysiol.* **118**, 3118–3131 (2017).

29. Talwar, S. K., Musial, P. G. & Gerstein, G. L. *Role of mammalian auditory cortex in the perception of elementary sound properties. Journal of Neurophysiology* vol. 85 www.jn.org (2001).

30. Gimenez, T. L., Lorenc, M. & Jaramillo, S. Adaptive categorization of sound frequency does not require the auditory cortex in rats. *J. Neurophysiol.* **114**, 1137–1145 (2015).

31. Jaramillo, S. & Zador, A. M. The auditory cortex mediates the perceptual effects of acoustic temporal expectation. in *Nature Neuroscience* vol. 14 246–253 (2011).

32. Wood, K. C., Town, S. M., Atilgan, H., Jones, G. P. & Bizley, J. K. Acute inactivation of primary auditory cortex causes a sound localisation deficit in ferrets. *PLoS One* **12**, (2017).

33. Kato, H. K., Gillet, S. N. & Isaacson, J. S. Flexible Sensory Representations in Auditory Cortex Driven by Behavioral Relevance. *Neuron* **88**, 1027–1039 (2015).

34. Ceballo, S., Piwkowska, Z. & Bourg, J. Targeted Cortical Manipulation of Auditory Perception In Brief. *Neuron* **104**, 1168-1179.e5 (2019).

35. Li, Z. *et al.* Corticostriatal control of defense behavior in mice induced by auditory looming cues. *Nat. Commun.* **12**, 1–13 (2021).

36. Town, S., Wood, K. & Bizley, J. Signal processing in auditory cortex underlies degraded speech sound discrimination in noise. *bioRxiv* 833558 (2019) doi:10.1101/833558.

37. Musall, S., Urai, A. E., Sussillo, D. & Churchland, A. K. Harnessing behavioral diversity to understand neural computations for cognition. *Current Opinion in Neurobiology* vol. 58 229–238 (2019).

38. Shadlen, M. N. & Kiani, R. Decision making as a window on cognition. *Neuron* vol. 80 791–806 (2013).

39. Newsome, W. T., Britten, K. H. & Movshon, J. A. Neuronal correlates of a perceptual decision. *Nature* **341**, 52–54 (1989).

40. Britten, K. H. *et al.* The analysis of visual motion: a comparison of neuronal and psychophysical performance. *J. Neurosci.* **12**, 4745–4765 (1992).

41. Shadlen, M. N., Britten, K. H., Newsome, W. T. & Movshon, J. A. A computational analysis of the relationship between neuronal and behavioral responses to visual motion. *J. Neurosci.* **16**, 1486–1510 (1996).

42. Nienborg, H. & Cumming, B. G. Decision-related activity in sensory neurons reflects more than a neurons causal effect. *Nature* **459**, 89–92 (2009).

43. Cumming, B. G. & Nienborg, H. Feedforward and feedback sources of choice probability in neural population responses. *Current Opinion in Neurobiology* vol. 37 126–132 (2016).

44. Tsunada, J., Liu, A. S. K., Gold, J. I. & Cohen, Y. E. Causal contribution of primate auditory cortex to auditory perceptual decision-making. *Nat. Neurosci.* **19**, 135–142 (2015).

45. Steinmetz, N. A., Zatka-Haas, P., Carandini, M. & Harris, K. D. Distributed coding of choice, action and engagement across the mouse brain. *Nature* **576**, 266–273 (2019).

46. Cohen, M. R. & Newsome, W. T. Context-Dependent Changes in Functional Circuitry in Visual Area MT. *Neuron* **60**, 162–173 (2008).

47. Cohen, M. R. & Newsome, W. T. Estimates of the contribution of single neurons to perception depend on timescale and noise correlation. *J. Neurosci.* **29**, 6635–6648 (2009).

48. Ni, A. M., Ruff, D. A., Alberts, J. J., Symmonds, J. & Cohen, M. R. Learning and attention reveal a general relationship between population activity and behavior. *Science (80-.).* **359**, 463–465 (2018).

49. Downer, J. D., Niwa, M. & Sutter, M. L. Task engagement selectively modulates neural correlations in primary auditory cortex. *J. Neurosci.* **35**, 7565–7574 (2015).

50. Stringer, C., Michaelos, M., Tsyboulski, D., Lindo, S. E. & Pachitariu, M. High-precision coding in visual cortex. *Cell* (2021) doi:10.1016/j.cell.2021.03.042.

51. Hires, S. A., Gutnisky, D. A., Yu, J., O'Connor, D. H. & Svoboda, K. Low-noise encoding of active touch by layer 4 in the somatosensory cortex. *Elife* **4**, (2015).

52. Hobbs, J. A., Towal, R. B. & Hartmann, M. J. Z. Spatiotemporal patterns of contact across the rat vibrissal array during exploratory behavior. *Front. Behav. Neurosci.* **9**, 356 (2016).

53. Aizenberg, M. & Geffen, M. N. Bidirectional effects of aversive learning on perceptual acuity are mediated by the sensory cortex. *Nat. Neurosci.* **16**, 994–6 (2013).

54. Aizenberg, M., Mwilambwe-Tshilobo, L., Briguglio, J. J., Natan, R. G. & Geffen, M. N. Bidirectional Regulation of Innate and Learned Behaviors That Rely on Frequency Discrimination by Cortical Inhibitory Neurons. *PLoS Biol.* **13**, e1002308 (2015).

55. Briguglio, J. J., Aizenberg, M., Balasubramanian, V. & Geffen, M. N. Cortical neural activity predicts

sensory acuity under optogenetic manipulation. *J. Neurosci.* **38**, 2094–2105 (2018).

56.   Wood, K. C., Angeloni, C. F., Oxman, K., Clopath, C. & Geffen, M. N. Neuronal activity in sensory
cortex predicts the specificity of learning. *bioRxiv* 2020.06.02.128702 (2020)
doi:10.1101/2020.06.02.128702.

57.   Ulanovsky, N., Las, L. & Nelken, I. Processing of low-probability sounds by cortical neurons. *Nat*
Neurosci **6**, 391–398 (2003).

58.   Natan, R. G., Carruthers, I. M., Mwilambwe-Tshilobo, L. & Geffen, M. N. Gain Control in the Auditory
Cortex Evoked by Changing Temporal Correlation of Sounds. *Cereb. Cortex* **27**, 2385–2402 (2017).

59.   Espejo, M. L., Schwartz, Z. P. & David, S. V. Spectral tuning of adaptation supports coding of sensory
context in auditory cortex. *PLoS Comput. Biol.* **15**, e1007430 (2019).

60.   Fritz, J., Shamma, S., Elhilali, M. & Klein, D. Rapid task-related plasticity of spectrotemporal receptive
fields in primary auditory cortex. *Nat. Neurosci.* **6**, 1216–1223 (2003).

61.   Mesgarani, N., Fritz, J. & Shamma, S. A computational model of rapid task-related plasticity of
auditory cortical receptive fields. *J. Comput. Neurosci.* **28**, 19–27 (2010).

62.   David, S. V., Fritz, J. B. & Shamma, S. A. Task reward structure shapes rapid receptive field plasticity
in auditory cortex. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 2144–9 (2012).

63.   Yin, P., Fritz, J. B. & Shamma, S. A. Rapid spectrotemporal plasticity in primary auditory cortex
during behavior. *J. Neurosci.* **34**, 4396–408 (2014).

64.   Niwa, M., Johnson, J. S., O'Connor, K. N. & Sutter, M. L. Active engagement improves primary
auditory cortical Neurons' ability to discriminate temporal modulation. *J. Neurosci.* **32**, 9323–9334
(2012).

65.   Fritz, J. B., Elhilali, M. & Shamma, S. A. Adaptive changes in cortical receptive fields induced by
attention to complex sounds. *J. Neurophysiol.* **98**, 2337–46 (2007).

66.   Schneider, D. M., Nelson, A. & Mooney, R. A synaptic and circuit basis for corollary discharge in the
auditory cortex. *Nature* **513**, 189–94 (2014).

67.   Schneider, D. M., Sundararajan, J. & Mooney, richard. A cortical filter that learns to suppress the
acoustic consequences of movement. *Nature* (2018) doi:10.1038/s41586-018-0520-5.

68.   McGinley, M. J., David, S. V. & McCormick, D. A. Cortical Membrane Potential Signature of Optimal
States for Sensory Signal Detection. *Neuron* **87**, 179–192 (2015).

69.   Reimer, J. *et al.* Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex.
Nat. Commun. **7**, 1–7 (2016).

70.   Atallah, B. V., Bruns, W., Carandini, M. & Scanziani, M. Parvalbumin-Expressing Interneurons
Linearly Transform Cortical Responses to Visual Stimuli. *Neuron* **73**, 159–170 (2012).

71.   Wilson, N. R., Runyan, C. A., Wang, F. L. & Sur, M. Division and subtraction by distinct cortical
inhibitory networks in vivo. *Nature* **488**, 343–348 (2012).

72.   Seybold, B. A., Phillips, E. A. K., Schreiner, C. E. & Hasenstaub, A. R. Inhibitory Actions Unified by
Network Integration. *Neuron* **87**, 1181–1192 (2015).

73.   Phillips, E. A. K. & Hasenstaub, A. R. Asymmetric effects of activating and inactivating cortical
interneurons. *Elife* **5**, e18383 (2016).

74.   Reynolds, J. H. & Heeger, D. J. The Normalization Model of Attention. *Neuron* **61**, 168–185 (2009).

75.   Natan, R. G. *et al.* Complementary control of sensory adaptation by two types of cortical
interneurons. *Elife* **4**, 163–174 (2015).

76.   Natan, R. G., Rao, W. & Geffen, M. N. Cortical Interneurons Differentially Shape Frequency Tuning
following Adaptation. *Cell Rep.* **21**, 878–890 (2017).

77.   Attneave, F. Some informational aspects of visual perception. *Psychol. Rev.* **61**, 183–193 (1954).

78.   Simoncelli, E. P. & Olshausen, B. A. Natural image statistics and neural representation. *Annual*
Review of Neuroscience vol. 24 1193–1216 (2001).

79.   Simoncelli, E. P. Vision and the statistics of the visual environment. *Current Opinion in Neurobiology*
vol. 13 144–149 (2003).

80.   Młynarski, W., Hledík, M., Sokolowski, T. R. & Tkačik, G. Statistical analysis and optimality of neural
systems. *Neuron* **109**, 1227-1241.e5 (2021).

81.   Wei, X.-X. & Stocker, A. A. A Bayesian observer model constrained by efficient coding can explain
'anti-Bayesian' percepts. *Nat. Neurosci.* **18**, 1509–1517 (2015).

82.   Guo, Z. V. *et al.* Procedures for behavioral experiments in head-fixed mice. *PLoS One* **9**, (2014).

83.   Isett, B. R., Feasel, S. H., Lane, M. A. & Feldman, D. E. Slip-Based Coding of Local Shape and
Texture in Mouse S1. *Neuron* **97**, 418-433.e5 (2018).

84.   Carruthers, I. M., Natan, R. G. & Geffen, M. N. Encoding of ultrasonic vocalizations in the auditory
cortex. *J Neurophysiol* **109**, 1912–1927 (2013).
85.   Carruthers, I. M. *et al.* Emergence of invariant representation of vocalizations in the auditory cortex. *J.*
Neurophysiol. jn.00095.2015 (2015) doi:10.1152/jn.00095.2015.
86.   Voigts, J. *et al.* An easy-to-assemble, robust, and lightweight drive implant for chronic tetrode
recordings in freely moving animals. *J. Neural Eng.* **17**, 26044 (2020).
87.   Voigts, J., Siegle, J., Pritchett, D. L. & Moore, C. I. The flexDrive: An ultra-light implant for optical
control and highly parallel chronic recording of neuronal ensembles in freely moving mice. *Front.*
Syst. Neurosci. **7**, 8 (2013).
88.   Pachitariu, M., Steinmetz, N., Kadir, S., Carandini, M. & Harris, K. *Fast and accurate spike sorting of*
high-channel count probes with KiloSort. Advances in Neural Information Processing Systems vol. 29
(2016).
89.   Eilers, P. H. C. & Marx, B. D. Flexible smoothing with B-splines and penalties. *Stat. Sci.* **11**, 89–102
(1996).
90.   Stanislaw, H. & Todorov, N. Calculation of signal detection theory measures. *Behav. Res. Methods,*
Instruments, Comput. **31**, 137–149 (1999).
91.   Rocchi, F. & Ramachandran, R. Neuronal adaptation to sound statistics in the inferior colliculus of
behaving macaques does not reduce the effectiveness of the masking noise. *J. Neurophysiol.* **120**,
2819–2833 (2018).
92.   Hautus, M. J. Corrections for extreme proportions and their biasing effects on estimated values of d′.
Behav. Res. Methods, Instruments, Comput. **27**, 46–51 (1995).
93.   Sahani, M. & Linden, J. F. How linear are auditory cortical responses? in *Advances in Neural*
Information Processing Systems 109–116 (2003). doi:10.1124/dmd.105.005157.concerning.
94.   Sahani, M. & Linden, J. F. *Evidence optimization techniques for estimating stimulus-response*
functions. Advances in Neural Information Processing Systems https://papers.nips.cc/paper/2294-
evidence-optimization-techniques-for-estimating-stimulus-response-functions.pdf (2003).
233
234