1  # Dynamics of cortical contrast adaptation predict perception of signals in noise.
2

3  Christopher F. Angeloni[1,2], Wiktor Młynarski[3], Eugenio Piasini[4], Aaron M. Williams[2,4], Katherine C. Wood[2],
4  Linda Garami[2], Ann Hermundstad[5], Maria N. Geffen[2,4]
5
6  [1]Psychology Graduate Group, University of Pennsylvania, Philadelphia, PA, USA
7  [2]Department of Otorhinolaryngology, University of Pennsylvania, Philadelphia, PA, USA
8  [3]Institute of Science and Technology Austria, Klosterneuburg, Austria
9  [4]International School for Advanced Studies, Trieste, Italy
10  [5]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA, USA

**Abstract**

Neurons throughout the sensory pathway adapt their responses depending on the statistical structure of the sensory environment. This adaptation is thought to be efficient, resulting in neuronal codes that maximize information about the stimulus. Contrast gain control is a form of efficient adaptation in the auditory cortex and is believed to be crucial for enhancing the detection of signals embedded in background noise. However, it is unclear whether the dynamics of contrast gain control reflect efficient adaptation, and how they inform behavioral perception in noise. Here, we trained mice to detect a target presented in background noise shortly after a change in the contrast of the background. The observed changes in cortical gain and behavioral detection followed the dynamics of a normative model of efficient contrast gain control; specifically, target detection and sensitivity to target level improved in low contrast backgrounds relative to high contrast backgrounds. Additionally, the time course of target detectability adapted asymmetrically depending on contrast, decreasing rapidly after a transition to high contrast, and increasing slowly after a transition to low contrast. Auditory cortex was required for detection of targets in background noise and cortical neuronal responses exhibited the patterns of target detectability observed during behavior and in the normative model. Furthermore, variability in cortical gain predicted behavioral performance beyond the effect of stimulus-driven gain control. Combined, our results demonstrate that dynamic gain adaptation supports efficient coding in auditory cortex and predicts behavioral performance in a target-in-background detection task.

**Introduction**

As we navigate the world around us, the statistics of the environment can change dramatically. In order to maintain stable percepts, the nervous system adapts to persistent statistical properties of sensory inputs. The efficient coding hypothesis postulates that the nervous system accomplishes this by matching the limited dynamic range of individual neurons to the statistics of incoming sensory signals[1], thereby efficiently encoding information within many types of environments[2–4]. Indeed, adaptation to environmental statistics has been found in many sensory modalities and species[5–13]. In the auditory system, neurons exhibit contrast gain control, adapting the gain of their response function to match the variability in level (contrast) of the incoming sounds[14–19]. Yet, it is unknown whether and how the dynamics of contrast gain control in the auditory system inform perception, as the dynamics of neuronal adaptation have not previously been measured simultaneously with behavior. The goal of our study was to test whether the dynamics of contrast gain control in auditory cortex predict changes in the perception of targets embedded in noise.

Contrast gain control has been proposed as a mechanism for creating sensory representations of sounds that are invariant to background noise[20]. In the ferret auditory pathway, representations of sounds become more noise-tolerant in higher auditory areas, and the amount of noise-tolerance correlates with the strength of adaptation to the stimulus statistics, suggesting that level adaptation and gain control enhance the encoding of stimuli embedded in stationary noise[16]. Additionally, recent psychophysical studies found that perception in noise is altered by efficient adaptation to stimulus statistics. In humans, target level discriminability is greater in low contrast than in high contrast, an effect consistent with gain control observed in primary auditory cortex[19]. Similar relationships between gain control and behavioral percepts of sound location have also been found in ferrets[10] and guinea pigs[21]. However, in these previous studies, the measurement of psychophysical performance and neuronal gain control were performed separately in human and animal subjects, respectively, so it remains unclear how variability in adaptation over sessions or subjects relates to behavioral performance. Additionally, these previous studies focused primarily on behavioral performance after complete adaptation to stimulus contrast, whereas recent work has highlighted the dynamical nature of efficient adaptation[22,23]. In the current study, we aimed to assess whether and how the dynamics of contrast gain control reflected efficient adaptation and predicted behavioral performance across subjects.

First, we established a normative framework to model the neuronal dynamics of gain control and predict how efficient contrast adaptation should affect the detection of signals in noise. We then derived a novel procedure for estimating moment-to-moment changes in neuronal gain based on generalized linear models (GLM) and found that the dynamics of gain control in auditory cortex were asymmetric, as observed in the normative model. Next, to test the role of contrast gain control in auditory behavior, we trained mice to detect targets after a change in background contrast. We found that contrast-induced changes in behavioral target detection threshold, sensitivity, and behavioral adaptation dynamics followed the normative model predictions. Furthermore, we found that auditory cortex was necessary for detection in the presence of a background, but not for detection in silence, suggesting a distinct role of auditory cortex in separating targets from the background. Building on this finding, we found that the dynamics of cortical encoding of targets resembled the normative model predictions and observed behavioral adaptation, and that population activity in the auditory cortex predicted individual variability in behavioral performance. Finally, we estimated cortical gain during the task, finding that variability in neuronal gain predicted variability in task performance. Combined, our results identify a predictive relationship between efficient adaptation via gain control and detection of signals in noise, and provide a normative framework to predict the dynamics of behavioral performance in changing sensory environments.

**Results**

*A novel target-in-background detection task and normative model for task predictions.*

To assess how behavioral performance is impacted by adaptation to stimulus contrast, we first devised a GO/NO-GO task in which mice were trained to detect targets embedded in switching low and high contrast backgrounds. During each trial, the mouse was presented with dynamic random chords (DRCs) of one contrast, which switched after 3 s to the other contrast. At variable delays after the contrast switch, broadband target chords were superimposed on the background chords, and mice were trained to lick for a water reward upon hearing the target (henceforth, we refer to high-to-low contrast trials as "low contrast" and low-to-high contrast trials as "high contrast", referring to the contrast where mice detected targets). Target trials were interleaved with background-only trials, during which the mouse was trained to withhold licking, but would receive a 7s timeout for licking after the contrast switch (Figure 1a,b). To assess behavioral sensitivity to targets, we parametrically varied target level in each contrast, and to assess behavioral adaptation, we parametrically varied target timing (Figure 1c). This stimulus design allowed us to quantitatively test whether and how the dynamics of adaptation to background contrast affect perception.

To estimate the optimal time course of contrast gain control and predict its impact on target detection, we developed a normative model of behavioral performance constrained by efficient neuronal coding. In this model, we simulated a neuron designed to encode stimuli with minimal error. To efficiently exploit its finite dynamic range, the model neuron estimated the contrast of the recent stimuli, and adjusted the gain of its nonlinearity to minimize the error in estimated contrast (Figure 1d, panels 1-3; *Online Methods*)[22,23]. Adding targets at different levels and times relative to contrast transitions allowed us to probe the sensitivity of the model neuron to targets of varying strength over the time course of adaptation (Extended Data Figure 1c,d). When varying target strength and measuring the "psychometric" performance of the model (Equation 1, *Online Methods*), we found decreased detection thresholds and steeper slopes in low contrast relative to high contrast (Figure 1e). When varying target timing, two factors affected target discriminability: 1) A change in the stimulus distribution after the contrast switch; 2) The effect of gain adaptation on responses to the background (Figure 1f,g; Extended Data Figure 1c,d). Recapitulating previous results[14], the dynamics of the model responses to targets were asymmetric across each contrast (Figure 1f). These dynamics were well characterized by a single effective timescale, which we quantified by fitting an exponential (Equation 2) to each transition. The normative model presented three primary predictions for target detection behavior: When adapted to low contrast, 1) target detection thresholds will be lower and 2) model psychometric functions will have steeper slopes, and, 3) target detectability during the adaptation period will be asymmetric: rapidly decreasing after a switch to high contrast, and slowly increasing after a switch to low contrast (Figure 1h).

*Estimated cortical gain dynamics follow a normative model of gain control.*

Previous work on contrast gain control used static models, measuring steady-state gain after the neuron fully adapted to the new stimulus[14,16,17,19], but see[15,24]. We developed a model to estimate the gain of neurons in auditory cortex over time following a contrast transition. This generalized linear model (GLM) was fit to data recorded from the auditory cortex of a naive mouse (n = 97 neurons) presented with 3 s alternations of low and high contrast DRCs (Figure 2a,b).

The inference model is a GLM with dynamic gain control (GC-GLM) that decomposes the relationship between spiking activity ($y_t$) and the presented sounds into a stimulus component ($x_t$), contrast component ($\bar{\sigma}/\sigma_t$), and an interaction between the stimulus and the contrast ($x_t * \bar{\sigma}/\sigma_t$, where $\bar{\sigma}$ is an arbitrary constant, defined as the contrast at which the gain is 1; *Online Methods*). We calculated a gain index ($w_t$) from the fitted model parameters (Figure 2b), which quantified whether gain control estimated by the model was optimal given the background contrast levels. For the contrast levels used in this study, $w_t$ = 1.5 indicates an optimal increase in gain during low contrast, $w_t$ = 0.5 indicates an optimal decrease in gain during high contrast, and $w_t$ = 1.0 indicates no gain control (*Online Methods*). To validate the model and gain index we simulated neurons with defined temporal trajectories of gain control and found that the model accurately recovered the ground-truth simulated gain dynamics (*Extended Data Figure 2*; *Supplementary Information*). For comparison, we also fit previously described linear-nonlinear (LN) models to each neuron[14,16,17,19], one with a static output nonlinearity (static-LN), and one with a contrast-dependent, or gain-controlled output nonlinearity (GC-LN, Figure 2c; representative neuron: Figure 2d-g). In this neuron, the fits of both the GC-LN model and GC-GLM exhibited contrast gain control, as characterized by high gain in low contrast and low gain in high contrast (Figure 2f and g, respectively), suggesting that both models capture similar contrast-driven changes in cortical gain.

**Figure 1. Target in background detection task and normative model predictions.**

**a,** Experimental setup. **b,** GO/NO-GO task design. Spectrograms are plotted for example NO-GO and GO trials with transitions from low to high contrast (top row) and high to low contrast (bottom row), with waveforms plotted below each spectrogram (color bar indicates the sound level). Below the example trials, the timing of the response window, schematic licks, and responses to licks are plotted. For NO-GO trials, licks in the response window received a timeout. For GO trials, licks in the response window were rewarded with 5uL of water. **c,** Example target parameters. *Top*: varied target levels, with level in dB SNR indicated by the color bar. *Bottom*: varied target times, where each arrow indicates a potential target delay. **d,** Normative model of efficient gain control. Target and background distributions for each contrast are indicated in the left panel. (1) Target stimuli are indicated with circles, while the background stimulus is indicated by a line. The stimulus response in a given time window is transformed by an adapting nonlinearity to generate spikes. (2) The spiking responses are decoded to update an estimate of the stimulus variance. (3) The gain of the nonlinearity is adjusted to optimally predict the variance of the next timestep. *Inset*: Example spike distributions of the model neuron in low and high contrast for targets (dark histograms), and background (light histograms). **e,** Model target-from-background discriminability as a function of contrast and target level. Circles indicate model performance overlaid with logistic function fits (solid lines) and thresholds (dashed lines). **f,** Model discriminability over time in low and high contrast. Circles indicate model performance overlaid with exponential function fits. **g,** Model gain dynamics over time in each contrast. **h,** From top to bottom: model predictions for target detection thresholds, slopes, and adaptation time constants in each contrast.

Qualitatively, GC-GLM outperformed standard LN models, primarily by capturing the adaptation dynamics after the transition (Figure 2d, middle panel), allowing us to analyze the gain control index ($w_t$) as a function of time (Figure 2d, bottom panel; Figure 2g). To test whether the GC-GLM could better account for the data than standard models, we compared cross-validated correlations of the model predictions with the trial-averaged PSTH for each neuron, finding a significant effect of model type on the correlations (n = 97 neurons; Kruskal-Wallis test: $H(2)$ = 93.61, p = 6.70e-21). GC-GLM correlation was significantly higher (Median ($Mdn$) = 0.75, Inter-Quartile Range ($IQR$) = 0.24) than the GC-LN model ($Mdn$ = 0.54, $IQR$ = 0.49, $p$ = 4.41e-6, post-hoc sign-rank test) and the static-LN model ($Mdn$ = 0.25, $IQR$ = 0.73, $p$ = 9.56e-10, post-hoc sign-rank test). Consistent with previous studies[14,15], we also found that the GC-LN model outperformed the static-LN model ($p$ = 3.50e-6, Figure 2h). To compare the GC-GLM to an existing dynamical model of contrast gain control, we also fit a recent model of dynamic contrast adaptation[19] to our data (Extended Data Figure 3a). Across the neurons in this dataset, we found that the GC-GLM was a better predictor of neuronal activity than the existing model (Extended Data Figure 3b).

We then quantified whether the GC-GLM reliably detected gain control in the neuronal population. Here, we defined steady-state gain control by calculating the change in $w_t$ between high ($w_H$) and low ($w_L$) contrastafter the gain has stabilized (1 s after the contrast switch). Based on our definition of $w_t$, $w_H - w_L$ = $-1$ if gain control is optimal (*Online Methods*). Across all neurons, we found significant gain control ($Mdn$: -0.10, $IQR$: 0.35, sign-rank test: $Z$ = -2.90, $p$ = 0.004; Figure 2i). To further validate the GLM estimates of gain, we compared the GC-GLM gain control indices at steady-state to those of the GC-LN model and found a significant relationship (linear regression: $F(1,95)$ = 12.20, $p$ = 7.33e-4, $R^2$ = 0.11; Figure 2j). Together, these results demonstrate that the GC-GLM model better accounts for the neuronal data by incorporating the dynamics of gain control and conclude that this method captures a similar estimate of steady-state gain control when compared to standard models.

Finally, we analyzed the dynamics of gain control by fitting $w_t$ after each contrast switch with an exponential function (Figure 2g). In neurons with gain control ($w_t < 0$ at steady state), the average time course of $w_t$ was asymmetric across contrast transition types, rapidly decreasing after a switch to high contrast, and slowly increasing after a switch to low contrast (n = 45 neurons; Figure 2k). Within this same population, we quantified the timescale of adaptation to each contrast using the time constant ($\tau$) of each exponential fit, finding significantly longer time constants in low contrast ($Mdn$ = 0.29, $IQR$ = .39) relative to high contrast ($Mdn$ = 0.048, $IQR$ = 0.094; sign-rank test: $Z$ = 4.52, $p$ = 6.16e-6; Figure 2l). This asymmetry in gain adaptation reflected the dynamics of the normative model (Figure 1g), prior electrophysiological studies[14], and the behavior of previously described optimal variance estimators[25].

*Mouse behavioral detection is modulated by background contrast.*

We next tested whether the asymmetry in gain control in cortex was reflected in behavioral sensitivity to targets in background noise. Mice were initially trained in a simple version of the GO/NO-GO task where they were required to lick in response to a target and withhold licks on trials without a target (Figure 1b, 3a). Out of the 25 mice trained, 24 mice learned this task reliably, typically reaching criterion performance of 80% correct within 2-3 weeks in either contrast (Figure 3b). False alarm rates were significantly higher in high contrast than in low contrast (Extended Data Figure 4a), suggesting that detection is more difficult in high contrast, which we discuss next.

By varying the level of presented targets, we collected psychometric curves for each mouse in each contrast. To isolate the effect of stimulus contrast on behavioral performance, we included sessions from mice that were exposed to similar target levels in low and high contrast (n = 11 mice; Figure 3c; see Extended Data Figure 4 and Supplementary Results for results using different target ranges). In this cohort, targets were easier to detect in low contrast: we observed significantly lower detection thresholds in low contrast (Mean ($M$) = 8.79, standard deviation ($std$) = 3.13) as compared to high contrast ($M$ = 15.39, $std$ = 3.27; paired t-test: $t(10)$ = -4.20, $p$ = 0.0057, Figure 3d). Furthermore, psychometric slopes were significantly steeper in low contrast ($M$ = 0.040, $std$ = 0.0048) compared to high contrast ($M$ = 0.036, $std$ = 0.0026; paired t-test: $t(10)$ = 3.037, $p$ = 0.023; Figure 3e). Combined, these results demonstrate that targets were easier to detect in low contrast, as predicted from the normative model (Figure 1h).

**Figure 2. Cortical adaptation to sound contrast is asymmetric.**

**a,** Schematic of acute recordings from auditory cortex. **b,** Schematic of the linear-nonlinear (LN) model, with a static (grey) or gain-controlled (blue, red) nonlinearity. **c,** A Poisson GLM for estimating gain dynamics. The model decomposed the spiking response, $y_t$, of the neuron into a stimulus component, $x_t$, a contrast component, $\bar{\sigma}/\sigma_t$, and an interaction between the stimulus and contrast, which determined the gain estimate. **d,** *Top*: Spike raster for a representative unit. Blue and red horizontal bars indicate low and high contrast periods of the trial, respectively. *Middle*: the spike rate of the neuron is overlaid with the predictions from a static LN model, LN model with gain control, or GLM with gain control. Colors are indicated in the legend. *Bottom*: gain index, $w_t$, estimated from the GLM parameters (*Methods*). Dashed black lines indicate gain index values for optimal gain control and grey dashed lines indicate gain index values for no gain control. Orange trace indicates the gain dynamics of the representative neuron. **e,** Spectro-temporal receptive field (STRF) fit to this neuron. Color bar indicates the strength of the filter response. **f,** Nonlinearities fit to the STRF prediction in low and high contrast. **g,** Dashed blue and red lines indicate the gain index of the example cell in low and high contrast. The overlaid solid lines are exponential fits to the data. **h,** Cross-validated Pearson correlation coefficients between the trial averaged model predictions and spike rates for each model (colors as in **d**). Errorbars indicate 95th percentiles around the median. Asterisks indicate the results of sign-rank tests. **i,** Distribution of gain control estimated by the GLM. Orange line indicates the median. Asterisks indicate the results of a sign-rank test. **j,** Gain control estimates for each neuron from the GC-GLM or the GC-LN model (black dots) overlaid with the best linear fit and 95% confidence interval of the fit. Asterisks indicate whether the linear model was significantly different from a constant model. **k,** Gain index for all of the neurons with gain control (n = 45). Light lines are the average ±SEM, while the dark lines are exponential fits to the average. **l,** Adaptation time constants from gain-controlled neurons after a switch to low (blue dots) and high contrast (red dots). Asterisks indicate the results of a sign-rank test. In all plots: ns, not significant; †p<0.1; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001.

189   To assess behavioral adaptation to the background contrast, we presented targets at threshold level
190 at variable delays following the contrast transition. We observed behavioral time courses consistent with the
191 normative model and with gain measured in auditory cortex: after a switch of the stimulus to high contrast,
192 detection rates decreased quickly over time, but after a switch to low contrast detection, rates increased slowly
193 over time (Figure 3f). In high contrast, the first significant drop in performance occurred between the first two
194 time points, while in low contrast the first significant increase in performance occurred between the first and
195 third time points (Figure 3f, Extended Data Table 1). Indeed, fitting exponential functions to performance over
196 time revealed that behavioral adaptation was significantly faster in high contrast (median and interquartile
197 range of time constant: *Mdn* = 0.023, *IQR* = 0.082) compared to low contrast (n = 21 mice, *Mdn* = 0.13, *IQR*
198 = 0.13; rank-sum test: *Z* = 2.75, *p* = 0.0060; Figure 3g).
199   To directly compare the predictions of the normative model presented in Figure 1 to behavioral
200 performance, we computed a contrast modulation index for the behavioral and model parameters (CMI:
201 Equation 3, *Online Methods*). To assess whether the model prediction was within the range of expected
202 behavioral values, we computed the 95% confidence intervals of the behavioral CMI values using a bootstrap
203 procedure. We found that the CMI values of the normative predictions fell within the range of expected CMI
204 values for behavioral thresholds and adaptation times. As observed in behavior, the model predicted a
205 decrease in slope in high contrast; however, the magnitude of the predicted decrease was larger than the
206 range of observed slope CMI values (Figure 3h). Taken together, the behavioral results were qualitatively
207 consistent with all three predictions from the normative model (Figure 1h): 1) Detection thresholds are lower
208 in low contrast; 2) Psychometric slopes are higher in low contrast; 3) Performance decreases rapidly in high
209 contrast and increases gradually in low contrast.
210
211 *Auditory cortex is necessary for detection in background noise.*
212   Whereas gain control is present in many areas along the auditory pathway, it is strongest in auditory
213 cortex[16,19]. As such, we hypothesized that auditory cortex supports the detection of sounds in the presence of
214 background noise. To test whether auditory cortex is required for task performance, we inactivated auditory
215 cortex using the GABA-A receptor agonist muscimol. We validated that muscimol disrupts cortical coding of
216 target sounds by applying muscimol topically to the cortical surface during passive playback of the behavioral
217 stimuli, finding near complete suppression of target responses (Extended Data Figure 5a-f, *Supplementary*
218 *Information*).
219   To test whether inactivation of auditory cortex affects behavioral performance, we repeated the same
220 experiments in behaving mice, administering muscimol or saline bilaterally through chronically implanted
221 cannulae (n = 44 sessions from 4 mice; Figure 4a). We found a profound decrease in the response rates to
222 targets and background in both contrasts (Figure 4b). We quantified these effects on the psychometric curve
223 using a three-way ANOVA with cortical intervention (muscimol or saline), contrast, and target level as factors.
224 We found significant main effects of cortical intervention ($F(1,307)$ = 278.63, *p* = 3.83e-44), contrast ($F(1,307)$
225 = 4.39, *p* = 0.037) and level ($F(6,307)$ = 40.90, *p* = 7.54e-36). Post-hoc tests showed that muscimol application
226 significantly decreased hit rates in both contrasts by 31.45% (95% CI: [27.76, 35.14], *p* = 1.060e-10), whereas
227 an increase in background contrast significantly decreased hit rates in both intervention conditions by 3.95%
228 (95% CI: [2.57, 7.64], *p* = 0.036). Furthermore, we observed significant interactions between target level and
229 cortical intervention ($F(6,307)$ = 14.11, *p* = 4.47e-14), and between target level and contrast ($F(6,307)$ = 2.97,
230 *p* = 7.87e-3), but we did not observe a significant interaction between contrast and cortical intervention,
231 suggesting that muscimol has the same effect in low and high contrast. To quantify the effects of muscimol
232 on behavioral performance, we extracted response rates to the maximum target level, false alarm rates,
233 thresholds, and slopes of psychometric functions fit to each session, and found that muscimol significantly
234 reduced every measure of behavioral performance, with the exception of behavioral threshold (Figure 4c,
235 Extended Data Table 1). From these results, we can conclude that auditory cortex is necessary for detecting
236 targets in background, regardless of background contrast.
237   A potential alternative effect of muscimol is a general loss of function that is not specific to hearing
238 target sounds. To control for this, we devised an alternative to the detection in background task where mice
239 detected targets in silence (Figure 4d). To ensure equivalency between the two tasks, we took the highest-
240 level target trials in the target-in-background task (25dB SNR in high contrast) and removed the background
241 noise during the target detection period (Figure 4e, bottom). In this new task, mice heard the exact same
242 targets as in the target-in-background task except that the background noise pre- and post-target presentation
243 was removed. This allowed us to test whether auditory cortex is specifically required for target detection in
244 the presence of a background (*Online Methods*).
245

**Figure 3. Target detection performance is consistent with the normative model predictions.**

**a,** Schematic of the behavioral setup and task outcomes. **b,** Performance as a function of task exposure. Individual traces indicate performance of mice first trained to detect targets in low contrast (blue) or high contrast (red). Thick lines are a 7-day running average across mice in each training group. The dashed red trace near 0.5 was the performance of a mouse who failed to learn the task in high contrast and was excluded from further analysis. **c,** Individual psychometric curves in low and high contrast, overlaid with psychometric fits to the average. Dashed vertical lines indicate detection thresholds. **d,** Detection thresholds from the data plotted in **c**. **e,** Slopes of the psychometric functions plotted in **c**. In **d, e** asterisks indicate the significance of paired t-tests. **f,** Detection performance of threshold level targets presented at different delays from the contrast switch (circles indicate mean ±SEM). Performance is overlaid with exponential fits in each contrast. Horizontal lines above the plot indicate significant sign-rank tests after false-discovery correction. **g,** Adaptation time constants from exponential functions fit to individual mice. Asterisks indicate the result of a sign-rank test. **h,** Comparison of normative model predictions (grey) to the data (black) for psychometric thresholds, slopes and adaptation time. The y-axis is the contrast modulation index (CMI, *Methods*) of the value in each plot, with error bars indicating the 95% confidence interval of the data determined via bootstrap. In all plots: ns, not significant; †$p<0.1$; *$p<0.05$; **$p<0.01$; ***$p<0.001$; ****$p<0.0001$.

To assess behavioral performance in this new task, we modulated detection difficulty by attenuating
the level of each target. As observed previously, inactivation of auditory cortex impaired detection in high
contrast (Figure 4e, top). However, cortical inactivation had little effect on behavioral performance in silence
(Figure 4e, bottom). We quantified these effects on behavior using a three-way ANOVA with cortical
intervention (muscimol or saline), task (detection in background or silence), and target level as factors (n =
26 sessions from 2 mice). We found significant main effects of intervention ($F(1,181) = 62.83$, $p = 3.62e-13$),
task ($F(1,181) = 6.82$, $p = 9.86e-3$), and level ($F(6,181) = 46.16$, $p = 1.69e-32$). Muscimol significantly reduced
hit rates by 20.2% (95% CI: [15.19, 25.17], $p = 1.060e-10$, post-hoc). Hit rates for targets presented in silence
were significantly elevated by 6.65% relative to targets presented in background (95% CI: [1.65, 11.64], $p =$
0.0090). Furthermore, we found significant interactions between cortical intervention and task type ($F(1,181)$
= 6.36, $p = 0.013$), intervention and level ($F(6,181) = 3.47$, $p = 2.98e-3$), and level and task type ($F(6,181) =$
8.47, $p = 5.43e-8$). As before, we parameterized behavioral performance by fitting the performance on each
session with a psychometric curve, and we extracted the response rates to the maximum target level, false
alarm rates, response rates at threshold level, and slopes of psychometric functions. During the target-in-
background task, we found significant effects of muscimol on the response rates at maximum level and
threshold, a moderate effect on psychometric slope, and no effect on false alarm rate. However, muscimol
application had no significant effect on any of these measures in the target-in-silence task (Figure 4f, Extended
Data Table 1). Taken together, these results show that while both cortical inactivation and the presence or
absence of background noise affected behavioral performance, these effects interacted: muscimol had a
larger effect on performance when background noise was present.
Combined, our findings demonstrate that the auditory cortex is specifically required for detection in the
presence of background noise, but not in silence. Our next goal was to test whether neuronal activity in AC is
predictive of behavioral performance.
272
*Cortical codes predict individual behavioral performance.*
To better understand how representations in the auditory cortex could give rise to behavior, we
chronically recorded from populations of neurons in auditory cortex of 12 mice while they performed the
psychometric task (Figure 5a, Extended Data Figure 6). When using microdrives with drivable tetrodes, we
lowered the tetrodes a small distance at the end of each session to record from new populations of cells. In
the 242 sessions analyzed, we recorded from 18±15 neurons simultaneously (mean±standard deviation), with
a maximum of 73 and minimum of 3 neurons simultaneously recorded. For the following analyses, we only
included neurons with spike rates greater than 1 Hz, realistic spike waveforms (*Online Methods*), and with
significant responses to targets (AUC value significantly greater than 0.5 at two or more levels, Figure 5b,
inset; *Online Methods*). Following these selection criteria, 12±9 neurons were included in analysis for each
session.
To quantify the representations of targets and background in the neuronal population (example
responses in Figure 5b,c), we adapted a population vector approach[26] to generate a target-from-noise
discriminability metric using population activity (*Online Methods*). This method allowed us to project trial
distributions in $n$-dimensional neuronal space along a single dimension that separated target and background
trials (Figure 5d, left panel). We then estimated the criterion projection value that best predicted whether each
trial contained a target or just background[27] (Figure 5d, right).
This population decoding method allowed us to estimate neurometric functions to directly compare to
psychometric functions for each mouse (Figure 5e). On average, neurometric and psychometric functions
were qualitatively similar (Figure 5f) and highly correlated across sessions (Figure 5g). To assess the
relationship between neurometric and behavioral performance, we first considered the potential variables that
were likely involved to modulate behavior, namely: the level of the presented targets, the contrast of the
background, and neuronal variability that could fluctuate from session to session. To isolate the effects of
each of these variables, we used a linear model to predict behavioral performance on each session using
target level, contrast, and the neurometric performance as predictors ($F(4,794) = 593.0$, $p = 3.15e-202$, $R^2 =$
0.69, Figure 5i, top).
299

**Figure 4. Inactivation of auditory cortex selectively disrupts detection of targets in background sounds.**

**a,** Schematic of chronic muscimol and saline application in behaving mice. Legend indicates the three potential background conditions in the task. **b,** Behavioral performance on individual sessions (light traces) as a function of contrast (red, blue) and muscimol or saline application (dashed and solid lines, respectively). -Inf indicates performance on background-only trials. Dots indicate the average performance across mice and sessions ±SEM, overlaid with logistic fits to the data. **c,** Effects of muscimol and contrast on multiple behavioral measures. Bars indicate the mean performance, while dots indicate performance on individual sessions. Asterisks indicate the significance of rank-sum tests. **d,** Example spectrograms and waveforms with a target presented in high contrast (top) and the same stimulus when the target was presented in silence (bottom). Color bar indicates the sound level, with black indicating silence. **e,** Behavioral performance in high contrast (top) and in silence (bottom). Formatting as in **b**. **f,** The effect of muscimol on multiple behavioral measures when targets were presented in high contrast (red bars) or in silence (black bars). Formatting as in **c**. In all plots: ns, not significant; †p<0.1; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001.

As expected, we found that basic stimulus features (target level and background contrast) were strong predictors of behavioral performance, such that performance was higher for increased target levels ($b$ = 0.14, 95% CI [0.12, 0.15], $p$ = 4.15e-65), while performance was lower for increased contrast ($b$ = -0.050, 95% CI [-0.071, -0.028], $p$ = 5.67e-6). These results confirm our previous results on the behavioral impacts of stimulus contrast (Figure 3). Crucially, we also observed that population activity was predictive of behavioral performance at a session level, such that an increase in neurometric performance was accompanied by an increase in behavioral performance ($b$ = 0.016, 95% CI [0.0017, -0.031], $p$ = 0.029). Finally, we found that the relationship between behavioral and neuronal performance was mediated by stimulus contrast ($b$ = 0.026, 95% CI [0.0042, 0.048], $p$ = 0.020; Figure 5i, bottom). Taken together, these results suggest that session-to-session fluctuations in cortical population activity resulted in corresponding fluctuations in behavior, and that this relationship was mediated by stimulus contrast.

To control for the effect of target level on the relationship between behavioral and neuronal performance, we first visualized the data by regressing out the effect of level on the model terms and behavioral performance: this visualization demonstrates the positive relationship between the two terms (Figure 5g, inset), as captured by the significant neurometric coefficient in Figure 5i. To further control for the effect of level, we split the data into low, medium and high target levels (Figure 5h). In all level-splits, the relationship between neurometric and behavioral performance was significant (Extended Data Table 1). These findings suggest that regardless of the target level, neuronal performance predicted behavioral performance.

Next, we assessed whether neurometric performance was affected by contrast similarly to behavioral performance. To isolate the effect of contrast on neurometric curves and best compare to psychometric curves presented in Figure 3, we selected only recording sessions with matched target ranges (n = 117) and fit each neurometric curve with a logistic function. We found that neurometric function thresholds increased in high contrast (*Mdn[IQR]* of low contrast sessions (n=82): 7.19[4.83], high contrast sessions (n = 35): 13.47[8.88]; rank-sum: $Z$ = 4.69, $p$ = 1.34e-6, Figure 5j), and that neurometric slopes decreased in high contrast (low contrast: 0.036[0.014], high contrast: 0.030[0.021]; $Z$ = 1.88, $p$ = 0.029, Figure 5k). These findings corroborated the observed changes in psychometric functions (Figure 3) and the predictions of the normative model (Figure 1).

Combined, these results demonstrate that parameters of neurometric and psychometric functions are affected by contrast as predicted by a normative model of gain control. We also find that individual variation in behavioral performance is predicted by population activity in auditory cortex, independently of the effect of contrast and target level, which further supports the role of auditory cortex in the detection of signals in noise.

*Dynamics of target detection during adaptation.*

We next measured how cortical discriminability evolved as a function of time and contrast in sessions where mice were presented with targets at threshold level at different offsets relative to the contrast switch. In line with our behavioral results (Figure 3f), we found that in high contrast the first significant drop in cortical discriminability occurred between the first two target times, while in low contrast the first significant drop occurred between the first and third target times (n = 43 recording sessions; Extended Data Table 1; Figure 5l, left). To quantify the speed of neuronal adaptation, we fit the average neuronal discrimination time course for each mouse with an exponential function (n = 8 mice). Consistent with the normative model (Figure 1f-h), and with gain control dynamics estimated from cortical activity (Figure 2k,l) and behavior (Figure 3f,g), we found asymmetric adaptation in the neuronal responses, with larger adaptation time constants in low contrast (*Mdn* = 0.14, *IQR* = 0.21) relative to high contrast (*Mdn* = 0.033, *IQR* = 0.16; sign-rank test: $p$ = 0.016; Figure 5l, right).

*Cortical gain predicts behavioral performance.*

Our results so far demonstrate that population codes in auditory cortex are not only shaped by stimulus features, but also predict variation in behavioral performance. To assess the role of cortical gain in behavior, we leveraged the design of the background sounds to estimate spectrotemporal receptive fields (STRFs) and nonlinearities of neurons recorded during task performance. For each neuron, we fit a model with a static nonlinearity (static-LN) or a model with gain control (GC-LN; Figure 6a-d). We then pooled the neurons recorded across sessions including only neurons with strong stimulus responses in both contrasts (*Online Methods*). To ensure that observed changes in gain were not due to changes in STRF shape during the task, we first compared STRFs estimated from each task epoch (ie. low and high contrast) and found that STRF properties were stable across the trial (Extended Data Figure 7, *Supplemental Methods*).

**Figure 5. Cortical responses to targets predict behavioral performance and exhibit contrast adaptation.**

**a,** Schematic of chronic recordings during behavior. **b,** Spike raster from a neuron recorded during the task. Trials are sorted by target level, as indicated in the legend. Below is the average spike rate in each condition, smoothed with a 5ms gaussian window. Inset is the area under the ROC curve (AUC) as a function of level, with error bars indicating bootstrapped 95% confidence intervals. **c,** Neurograms of populations of simultaneously recorded neurons for a low contrast (left) and high contrast (right) behavioral session. Color bars indicate the difference in firing rate between the noise-only condition and each target level on the x-axis for each cell on the y-axis. The arrow indicates the cell in b. **d,** *Upper left*: schematic for estimating the direction (CD) that best separates population responses to targets (blue dots) and noise (grey dots). *Lower right*: probability distributions of population projections for noise-only trials (grey) and the highest level target trials (blue). The vertical red line indicates the criterion for the population decoder. **e,** Neurometric performance compared to behavioral performance in representative low contrast (left) and high contrast (right) sessions. In both plots, dark dots indicate the decoder performance, while light dots indicate behavioral performance on the same session, overlaid by logistic function fits. The arrow in the left panel is the decoder performance for the distributions plotted in d. **f,** Average psychometric and neurometric performance ±SEM over all trials. Solid lines indicate logistic fits to the average. As in **c**, light shades indicate behavioral performance and dark shades indicate neurometric performance. **g,** Behavioral performance as a function of neurometric performance for n=133 sessions. Each point is performance at a single target level in a single session. Point color indicates whether the session was a low contrast session (blue) or high contrast session (red), while point size indicates the target level. Dots joined by grey lines were collected in the same behavioral sessions. Inset: Same data as in **g**, with the effect of target level regressed out (*!L*). The x-axis is the adjusted model prediction and the y-axis is the adjusted behavioral percent correct. The black line indicates the model fit and 95% confidence interval. **h,** Behavioral performance as a function of neural performance split by target levels. Text indicates the correlation coefficient and its significance value. **i,** Top: schematic of the linear model used to predict behavior. Bottom: model coefficients and standard error for target level (*L*), contrast (*C*), neurometric performance (*N*), and the interaction between the latter two terms (*C\*N*). Asterisks indicate significant predictors. **j,** Neurometric thresholds for n=113 sessions in low and high contrast. Asterisks indicate significant rank-sum test across contrast. **k,** Same as j, but for neurometric slopes. **l,** Left: Neurometric performance ±SEM as a function of target delay. Formatting and statistical tests are the same as Figure 3f. Right: Adaptation time constants of the neurometric response for n = 8 mice. Asterisk indicates a significant rank-sum test. In all plots: ns, not significant; †p<0.1; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001.

First, we compared the cross-validated performance of the static-LN model versus the GC-LN
model. We found higher correlations using the GC-LN model (*Mdn* = 0.82, *IQR* = 0.17) relative to the static-
LN model (*Mdn* = 0.67, *IQR* = 0.12; n = 2,792 neurons; sign-rank test: *Z* = -36.74, *p* = 1.88e-295; Extended
Data Figure 8a). We also found significantly higher gain in low contrast (*Mdn* = 0.10, *IQR* = 0.13) than in
high contrast (*Mdn* = 0.041, *IQR* = 0.023; sign-rank test: *Z* = 37.92, *p* = 1.070e-314; Figure 6e, inset). These
results demonstrate that LN models can more accurately predict cortical activity when incorporating contrast
gain control, and confirm previous reports of robust gain control in mouse auditory cortex[17–19].
Based on our previous results, we expected that the strength of gain control in auditory cortex would
predict target detectability. When fitting the GC-LN model, we separately estimated neuronal gain during the
adaptation period of the trial and the target period of the trial (defined as the time periods before and after
the contrast switch, respectively; Figure 6b). To quantify the effects of contrast and trial period on gain, we
performed a two-way ANOVA, with gain as the dependent variable, and contrast, trial period, and their
interaction as factors (n = 2,262 neurons, after excluding outliers, see *Online Methods*). As expected, we
found a significant main effect of contrast ($F(1,4523) = 431.03$, *p* = 1.60e-91). Furthermore, there was a
significant main effect of trial period ($F(1,4523) = 35.79$, *p* = 2.36e-9) and a significant interaction between
contrast and trial period ($F(1,4523) = 77.91$, *p* = 1.51e-18). Post-hoc tests revealed that, in low contrast,
gain during the target period significantly increased (0.032, 95% CI: [0.024, 0.040], *p* = 3.77e-9), but did not
significantly change in high contrast (0.0062, 95% CI: [-0.017, 0.014], *p* = 0.18; Figure 6e). These findings
indicate that neuronal gain is not only sensitive to stimulus contrast, but also increases during the target
period of the trial, specifically in low contrast.
To visualize the gross relationship between gain and behavioral performance, we first averaged the
gain of stimulus-responsive neurons during the target period of the trial in each session (n = 168 sessions
across 13 mice). We then selected only low contrast sessions and split the data by the median gain in the
target period, computing the average psychometric curves for sessions in the bottom versus the top 50th
percentile (Figure 6f, inset). We observed that sessions with high gain had steeper slopes and lower
thresholds than sessions with low gain (Figure 6f). To quantify the relationship between gain and task
performance, we fit a mixed-effects model using contrast and gain during the target period as fixed effects,
mouse identity as a random effect, and either psychometric slopes or thresholds as the dependent variable.
This approach allowed us to separate the neuronal and behavioral impact of contrast gain control from the
effect of session-to-session fluctuations in gain. We tested whether gain and contrast were significant
predictors of behavioral performance by comparing the full model to null models excluding either gain or
contrast. We found that the model including gain was a better predictor of behavioral threshold than was the
null model (Likelihood Ratio Test: $\chi^2(1) = 5.82$, *p* = 0.016), indicating that thresholds decreased by 3.046 dB
SNR ±1.24 (SEM) for every 10% increase in gain. Using a similar procedure, we found that contrast was also
a significant predictor of behavioral threshold (Likelihood Ratio Test: $\chi^2(1) = 5.84$, *p* = 0.038), with the step
from high to low contrast inducing a decrease in behavioral thresholds of 3.27 dB SNR ±1.33 (Figure 6g).
We applied the same analysis to test the effects of contrast and gain on psychometric slope (Figure
6f), again finding that gain significantly predicted psychometric slopes (Likelihood Ratio Test: $\chi^2(1) = 6.96$, *p*
= 0.0083), such that the psychometric slope increased by 0.16 dB/PC ±0.060 for every 100% increase in gain.
However, contrast did not significantly improve the fit of this model (Likelihood Ratio Test: $\chi^2(1) = 2.28$, *p* =
0.13; Figure 6h). This result is not entirely unexpected, given that we observed no effect of contrast on
psychometric slopes when comparing across sessions with different target distributions (Extended Data
Figure 2b), which is true of the sessions used in this analysis.
Our findings suggest that the relationship between gain and behavioral performance is shaped by two
sources: contrast-induced gain control and fluctuations in gain from session to session. To further disentangle
the relationship between these two sources of behavioral modulation, we repeated the mixed effects models,
this time using gain during the adaptation period as the predictor of interest. We hypothesized that gain in this
period should not be predictive of behavioral performance, as there were no targets presented during this
portion of the trial. We found that this was the case; we did not observe any predictive relationship between
gain estimated in this period and behavioral performance (Extended Data Figure 8b-d; Extended Data Table
1). In summary, we found that cortical gain was modulated by both stimulus contrast and trial period,
increasing when contrast is low and when mice were detecting targets. Furthermore, we found that behavioral
performance was predicted by both the stimulus contrast and by session-to-session changes in cortical gain
during target detection.
413

**Figure 6. Cortical gain predicts session-to-session variability in behavioral performance.**

**a,** Schematic of the LN model fit to neuronal responses during the behavioral task. **b,** Spike raster of a representative unit recorded during the behavioral task, sorted by the background scene presented in each trial. Below is the average spike rate (black trace) overlaid with the LN model fit with a static (grey) or gain-controlled nonlinearity (green). **c,** STRF for the representative cell. **d,** The fitted nonlinearities in low and high contrast periods of the trials. **e,** Gain distributions across all recorded neurons as a function of contrast (indicated by the color) and whether the gain was estimated during the adaptation period (A) or the target period (T) of the trial (see labels in **b**). Inset: the average gain in low and high contrast for all cells, dashed lines indicate the median of each distribution. Asterisks indicate the results of rank-sum tests. **f,** Psychometric curves split by the median gain. Light colored dots indicate performance across sessions with low gain, dark colored dots indicate performance on sessions with high gain. Error bars indicate ±SEM. *Inset*: the distribution of gain values on the same sessions. The dashed red line indicates the median used to split the data. **g,** Relationship between session to session changes in gain and behavioral thresholds. Each dot is a session, with the color indicating the contrast in which targets were presented. The grey line is the linear best fit. Black asterisks indicate whether gain is a significant predictor of the psychometric threshold, while magenta asterisks indicate whether contrast was a significant predictor of behavioral thresholds. **h,** Same formatting as **g**, but plotting the relationship between gain and psychometric slope. In all plots: ns, not significant; †p<0.1; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001.

414
415

**Discussion**

Auditory environments exhibit complex statistical properties that change over time. Changes in the dynamic range, or contrast, of acoustic inputs poses a challenge to the auditory system, which is composed of neurons with limited dynamic range. The efficient coding hypothesis predicts that as stimulus contrast changes, neurons should adjust their gain in order to match their limited dynamic range to that of the stimulus distribution[1]. Multiple studies have demonstrated that neurons throughout the auditory pathway exhibit such contrast gain control[14,16,19]. Additionally, contrast gain control is theoretically beneficial for separating signals from background noise[16,20], implicating that this phenomenon should underlie perception in noisy environments. Whereas recent work has demonstrated a link between contrast gain control and changes in behavioral discrimination in humans[19,21], it is unclear how cortical gain directly relates to behavior, as neuronal responses and behavior were not observed simultaneously. Finally, recent work has highlighted the dynamic nature of efficient neuronal codes[22,23], and it is unclear whether the observed neuronal dynamics reflect what is predicted to be theoretically efficient, and how the observed dynamics affect behavior.

Our study addressed these open questions by developing a normative framework to predict how efficient gain adaptation should affect behavior, and then testing those predictions using a combination of behavior, chronic recordings, and cortical manipulations. First, we developed a normative model based on efficient coding[22,23], which predicted that: 1) Detection thresholds of targets should be lower in low contrast than in high contrast; 2) Sensitivity to changes in target level should be greater in low contrast relative to high contrast; and 3) Detection should change asymmetrically over time: increasing slowly after a switch to low contrast, but decreasing rapidly after a switch to high contrast (Figure 1). Then, we used a novel form of a Poisson GLM to confirm that gain control dynamics in auditory cortex are indeed asymmetric (Figure 2). To behaviorally test the predictions of the normative model and GLM, we trained mice to detect a target embedded in dynamic random chords while manipulating the contrast of the background between high and low contrast. As predicted by the normative model, mice had lower detection thresholds and were more sensitive to changes in target level in low contrast. Behavioral adaptation was also asymmetric, decreasing rapidly after a switch to high contrast, and increasing slowly after a switch to low contrast, in agreement with our model (Figure 3). Furthermore, we found that AC is necessary for this detection-in-background task (Figure 4). When recording in AC, we found that the variability in neuronal population responses significantly predicted session-by-session variability in behavior above and beyond variability induced by stimulus features, and we demonstrated that target discriminability adapted asymmetrically, as predicted (Figure 5). Finally, we found that the amount of cortical gain also predicted behavioral performance on a session-to-session basis, independently of the effect of contrast (Figure 6). Taken together, these results support our hypothesis that the dynamics of gain control support efficient coding in auditory cortex and predict the perception of targets embedded in background sounds.

*The role of cortex in behavior.*

The role of auditory cortex in behavior has been subject of debate. A number of prior studies found that auditory cortex was not required for relatively simple behavioral tasks such as frequency discrimination or detection[28,29]. Rather, many studies found that auditory cortex is primarily involved in more complex behaviors, such those requiring temporal expectation[30], localization[31], or discrimination of more complex sounds[32–34]. Consistent with previous findings[35], we found that AC inactivation selectively impaired detection of targets in a noisy background, but did not impair detection of targets in silence (Figure 4). Furthermore, neuronal activity in AC predicted variability in behavioral performance (Figures 5, 6). This set of results establishes that AC is necessary for the detection of targets in background noise and supports the more general notion that AC is required for more difficult auditory tasks.

While the previous work demonstrates the necessity of auditory cortex in behavioral performance, the brain areas and mechanisms supporting the transformation from stimulus to decision are an active field of study[36,37]. By recording neuronal activity during the task, we were able to leverage behavioral variability to show that task performance covaried with representations of targets within small neuronal populations (Figure 5), and with cortical gain (Figure 6). There is a large body of literature relating cortical codes to behavioral variability: early studies in the visual system suggested that information from relatively small numbers of neurons was sufficient to match or outperform animal behavior in psychophysical tasks[38–40] and that behavioral choice can be predicted from activity in sensory areas[27,40]. These accounts suggested that variability in bottom-up sensory encoding drives the variability in behavioral output. However, more recent work suggests that variability in sensory areas is driven by top-down influences[41–44], which are modulated by attention and learning[45–48]. A recent study imaging tens of thousands of neurons in the visual cortex supports

472     this notion, finding that cortical representations had higher acuity than behaving mice, yet did not correlate
473     with behavioral performance, suggesting that perceptual discrimination depends on post-sensory brain
474     regions[49].
475           Our results suggest that bottom-up adaptation to stimulus statistics shapes behavioral output: We
observed asymmetric time courses of target discrimination following a change in contrast (Figure 3). The
asymmetric adaptation in behavioral performance was consistent with predictions derived from a normative
account of contrast gain control (Figure 1), resembled contrast gain adaptation in auditory cortex in the
absence of behavior (Figure 2), and was evident in patterns of target-driven activity in auditory cortex during
task performance (Figure 5). Indeed, there have been other studies demonstrating that individual differences
in sensory-guided behaviors are reflected in cortical activity[50,51], are bidirectionally modulated by cortical
manipulation[52,53], and can be predicted from tuning properties in auditory cortex[54,55]. While our results cannot
rule out top-down input as the causal driver of sensory decisions, they do support the notion that the sensory
information upon which decisions are made is shaped by neuronal adaptation, which thereby affects
behavioral outcomes.
486
487     *Roles of gain in the auditory system.*
488           Neurons throughout the auditory system adapt to the statistics of the acoustic environment, including
the frequency of stimuli over time[56,57], more complex sound patterns[24,58], and task-related or rewarded
stimuli[59–64]. In this study, we focused on contrast gain control as a fundamental statistical adaptation that
relates to efficient coding[14,17–19]. Previous work identified that responses to targets presented after opposing
changes in contrast were asymmetric[14], a result consistent with optimal estimation of stimulus variance[25]. We
observed similar asymmetric dynamics in a normative model of contrast gain control (Figure 1), which
suggests that previous observations of asymmetric adaptation are a result of efficient encoding of the
background stimulus.
496           Inspired by this and other previous work[59], we intentionally designed our stimuli using unbiased white-
noise backgrounds, which allowed us to fit encoding models to our data. In this study, we developed a novel
application of Poisson GLM that allowed us to quantify gain dynamics directly from responses to continuous
DRC stimuli. This approach allowed us to verify that gain adaptation in auditory cortex is asymmetric (Figure
2), and we found that this model better predicted neuronal activity when compared to previous models (Figure
2, Extended Data Figure 3). While we focused on the multiplicative influence of contrast in this study, the GLM
presented here could in theory be applied to any other time-varying signal that modulates neuronal gain, such
as movement[65,66], arousal[67,68], or experimental interventions such as optogenetics[69–72]
504           Additionally, previous studies found that contrast gain control was predictive of behavioral
performance[10,19]. Here, we extended these findings by examining the dynamics of this process and found that
behavioral detection of targets also adapted asymmetrically (Figure 3). In addition to confirming previously
reported effects of contrast on psychometric curves, these results suggest that the dynamics of contrast gain
control influenced task performance. Pursuing this further, we estimated cortical gain as mice performed the
task, and discovered a predictive relationship between stimulus contrast, gain in auditory cortex, and
behavioral performance (Figure 6). These results suggest two sources of cortical gain modulation: 1) Bottom-
up adaptation to stimulus contrast (ie. contrast gain control), and 2) session-to-session modulation of gain.
Previous studies have demonstrated this latter phenomenon, suggesting that top-down gain modulation
underlies attention[41,42,73] and the maintenance of optimal behavioral states[67,68]. Our results suggest that both
contrast gain control and session-to-session fluctuations in gain modulate behavior, providing a starting point
for dissecting the neuronal mechanisms underlying these two forms of gain modulation.
516
517     *Cellular mechanisms of gain control.*
518           While this work and other studies have established contrast gain control as a fundamental property of
the auditory system, the neuronal mechanisms driving gain adaptation at a cellular level remain unclear.
Previous work theorized that cortical feedback was responsible for contrast gain control in early areas[16,20], but
recent experiments disproved this hypothesis[19]. In the current study, we have likely recorded from a mixed
population of excitatory and inhibitory neurons, the latter of which exhibit specific roles in adaptation[74,75]. While
specific inhibitory neuronal subtypes facilitate divisive or subtractive control of excitatory responses in
visual[69,70] and auditory cortex[71,72], the role of these interneurons in contrast gain control has been
inconclusive[18]. Additionally, our results highlight the distinction between stimulus driven gain control and
stimulus-invariant gain control which covaried with behavior from session to session (Figure 6). Whether these
two forms of gain control share common neuronal substrates is unclear. By combining cell-specific

optogenetic methods with behavioral tasks, future studies may explore and test the causal role of local circuits
and top-down modulation in gain control and behavior.
530
*The missing link between efficient coding and behavior.*
Combined, our results develop a framework and provide support for the role of contrast gain control
in behavior. The efficient coding hypothesis has emerged as one of the leading principles in computational
neuroscience that has shaped our understanding of neuronal coding, architecture, and evolution[1,76–79], and
prior research found that human behavior follows principles of efficiency[19,80]. Here, we focused on a well-
studied form of efficient coding, contrast gain control, and developed a framework to link the dynamics of
efficient neuronal coding with behavioral performance. While the mechanisms of contrast gain control in
auditory cortex remain unclear, this study highlights potential top-down and bottom-up influences on cortical
gain, which may or may not share common neuronal substrates. We believe that the theoretical frameworks
and modelling methods presented here will be broadly applicable in future studies of neuronal gain control, a
fundamental function of the nervous system.
542
543

**Online Methods**

*Animals.*

All experiments were performed in adult male (n = 19) and female (n = 19) C57BL/6 (Stock No. 000664) or B6.CAST-*Cdh23*$^{Ahl/+}$ (Stock No. 002756) mice (The Jackson Laboratory; age 12-15 weeks; weight 20-30g). Some of the mice used in these experiments were crossed with other cell-type specific -cre lines, as detailed in Extended Data Table 2. All mice were housed with, at most, five mice per cage, at 28°C on a 12-h light:dark cycle with food provided ad libitum, and a restricted water schedule (see *Water restriction*). All experiments were performed during the animals' dark cycle. All experimental procedures were in accordance with NIH guidelines and approved by the Institutional Animal Care and Use Committee at the University of Pennsylvania.

*Surgery.*

Mice were anesthetized under isoflurane (1-3%). Prior to implantation, all mice were administered subcutaneous doses of buprenorphine (Buprenex, 0.05-0.1 mg/kg) for analgesia, dexamethasone (0.2 mg/kg) to reduce brain swelling, and bupivicane (2 mg/kg) for local anesthesia. In mice implanted with microdrives, two ground screws attached to ground wires were implanted in the left frontal lobe and right cerebellum, with an additional skull screw implanted over the left cerebellum to provide additional support. A small craniotomy was performed over the target stereotactic coordinates relative to bregma, -2.6mm anterior, -4.3mm lateral. Either custom 16-channel microdrives, 32-, or 64-channel shuttle drives (cite) holding tetrode bundles of polyimide-coated nichrome wires were chronically implanted over auditory cortex, and tetrodes were lowered 800um below the pial surface. The exposed tetrodes were covered with GelFoam (Pfizer) or sterile silicone lubricant and sealed with Kwik-Cast (World Precision Instruments). The plastic body of the microdrive and a custom stainless-steel headplate were secured to the skull using dental cement (C&B Metabond) and acrylic (Lang Dental). Mice undergoing only behavioral experiments were implanted with two skull screws in the cerebellum, and a headplate was mounted on the skull as previously described. An antibiotic (Baytril, 5mg/kg) and analgesic (Meloxicam, 5mg/kg) were administered daily (for 3 days) during recovery.

*Water restriction.*

Following surgical recovery (3 days post-operation), each mouse's weight was monitored for three additional days to establish a baseline weight. Over the next seven days, mice were water deprived, beginning with a daily ration of 120uL/g and gradually decreasing their ration to 40-50uL/g. During the task, if mice did not receive their full ration, the remainder of their ration was provided in their home cage. Mouse weight relative to baseline was monitored during all stages of water restriction. Additional health signs were used to determine a health score and subsequent treatment plan if a mouse lost more than 20% of baseline weight, as described by previously published methods[81] and approved by the Institutional Animal Care and Use Committee at the University of Pennsylvania.

*Behavioral apparatus.*

During the GO/NO-GO task, the mouse was head-fixed in a custom-built, acoustically isolated chamber. A capacitive touch sensor (AT42QT1010, SparkFun) soldered to a lick spout monitored lick activity. Water rewards were dispensed from a gravity fed reservoir, controlled by a solenoid valve (161T011, Neptune Research) calibrated to deliver approximately 4-5uL of water per reward[82]. Low-level task logic – such as lick detection, reward and timeout delivery, and task timing intervals – was directly controlled by an Arduino Uno microprocessor running custom, low-latency software routines. High-level task logic, such as trial randomization, stimulus buffering and presentation, and online data collection and analysis were controlled by custom MATLAB (Mathworks) software communicating with the Arduino over a USB serial port. Acoustic waveforms were generated in MATLAB and converted to analog signals via a soundcard (Lynx E44, Lynx Studio Technology, Inc.) or a National Instruments card (NI PCIe-6353) and delivered through an ultrasonic transducer (MCPCT-G5100-4139, Multicomp). The transducer was calibrated to have a flat frequency response between 3 kHz and 80 kHz using a 1/4-inch condenser microphone (Brüel & Kjær) positioned at the expected location of the mouse's ear, as described previously[83,84]. During electrophysiological recording sessions, licks were detected using an optical interrupt sensor (EE-SX771, Omron Automation), to prevent lick-related electrical artifacts introduced by contact with a capacitive sensor.

*Behavioral timeline.*

Each mouse underwent four stages in the behavioral task: 1) water restriction and habituation, 2)
behavioral training, 3) psychometric testing, and, 4) offset testing. During the induction of water restriction,
mice were habituated to head-fixation in the behavioral chambers and received water through the lick spout,
getting a drop of water for licks separated by more than 2 s. After the mouse began to receive its entire ration
by licking in the booth, behavioral training was initiated (typically after 1 week). Each mouse was initially
trained and tested in one contrast condition (see *Stimuli*), with the initial training condition counterbalanced
across mice. Behavioral performance was monitored during training, and mice were considered trained after
completing at least three consecutive sessions with over 80% percent correct. After completing training,
behavioral thresholds were measured during at least three sessions in which psychometric stimuli were
presented (see *Stimuli*). After estimating the behavioral threshold for each mouse, offset stimulus sets were
generated using threshold-level targets. After completion of at least three sessions in the offset task, each
mouse was then retrained on the remaining contrast condition. Upon reaching the training criterion of 80% in
the new contrast condition, mice were then tested in the psychometric and offset tasks as previously
described. For mice in electrophysiological experiments, this sequence of training and testing was continued
until the recording site yielded less than three units, or until the mouse stopped performing in the task.
615
*Stimuli.*
All stimuli were created in MATLAB and sampled at 192 kHz or 200 kHz and 32-bit resolution. A set
of dynamic random chords (DRCs) were created with different contrasts, similarly to those described in
previous studies[14,17,19]. To construct a DRC, amplitude modulated pure tones were generated at multiple
frequencies and then superimposed to create a chord. In some experiments, 34 frequencies were sampled
between 4 and ~40kHz in 1/10 octave steps, in the remaining experiments, 33 frequencies were sampled
between 4 and 64kHz in 1/8 octave steps. The amplitude envelope of each tone was generated as follows:
every 25 ms, amplitudes for each frequency were sampled from a uniform distribution with a mean of 50 dB
and a width of ±5 dB in low contrast or ±15 dB in high contrast. Between each 20 ms chord, the amplitude
envelope of each frequency band was linearly ramped over 5 ms to the amplitude value for the next chord,
such that the total duration of each chord and its ramp was 25 ms. To synthesize the stimuli, amplitude
envelopes were multiplied by a sine wave of their respective frequencies, and summed to produce the final
waveform. Each time a set of DRCs was generated, 5 unique random number generator seeds were used to
restrict the background noise to 5 distinct scenes (see raster in Figure 6 for an example of spike-locking to
the repeated scenes).
In all stages of behavioral training and testing, stimuli created for each trial consisted of a DRC
background containing a change in contrast, and the presence or lack of a target at a delay after the change
in contrast. Each trial began with 3 seconds of DRC background from one contrast, followed by a switch to
the other contrast. Targets consisted of a fixed chord composed of 17 frequencies pseudo-randomly sampled
from the frequencies contained in the DRC background, such that the target frequencies were uniformly
distributed across the frequency range of the background. To add targets to the background noise, the target
amplitude at each target frequency was simply added to a single chord in the amplitude envelope of the
background, and ramped as described previously: this procedure ensured that target timing was perfectly
aligned to changes in the background noise, removing asynchronous timing cues that could be used to detect
the target. Target amplitudes are described in values of signal-to-noise ratio (SNR) relative to the average
level of the background noise (ie. a 50 dB target embedded in 50 dB background would have an SNR of 0
dB). See Extended Data Table 3 for SNRs used for each mouse. In all trials, targets were embedded after a
change in the background contrast, with a delay and level dependent on the current training or testing stage.
644
*Efficient coding model.*
We simulated a model neuron that encodes incoming stimuli via an adapting neuronal nonlinearity.
Stimuli were drawn from a Gaussian distribution whose mean $\mu$ was fixed over time but whose standard
deviation $\sigma_t$ could switch over time between a low and a high value ($\sigma_t = \sigma^L$ and $\sigma_t = \sigma^H$, respectively). At
each time $t$, a stimulus $s_t$ was drawn from the distribution $p(s_t|\sigma_t) = \mathcal{N}(s_t; \mu, \sigma_t^2)$, transformed via a
saturating nonlinearity of the form $1/(1 + e^{-k(s_t - s_0)})$, distorted by Gaussian noise with variance $\sigma_n^2$, and
finally discretized into $N$ discrete levels to generate a response $r_t$. This discrete response was linearly
decoded to extract an estimate $\hat{s}_t$ of the current stimulus: $\hat{s}_t = p_1 r_t + p_0$. The recent history of $L$ stimulus
estimates was used to update an estimate $\hat{\sigma}_t$ of the underlying standard deviation: $\hat{\sigma}_t = \text{std}(\hat{s}_{t-L+1}; \hat{s}_t)$. The
estimate $\hat{\sigma}_t$ was then used to select the parameters of the encoder ($k, s_0$) and the decoder ($p_1, p_0$) on the
next timestep. The encoding and decoding parameters were chosen to minimize the expected error in

decoding stimuli given the neuron's current estimate of the underlying standard deviation:

$k, s_0, p_1, p_0 = \text{argmin}\langle(\hat{s}_t - s_t)^2\rangle_{p(s_t|\hat{\sigma}_t)}$[22,23].

The parameters of the encoder and decoder were adapted based on a background stimulus with a
mean $\mu_B$ that was fixed over time and a standard deviation $\sigma_B$ that switched between low and high values
$\sigma_B^L$ and $\sigma_B^H$, respectively. We used this adapting nonlinearity to determine how well this model neuron could
discriminate target stimuli from this background. Target stimuli were sampled from a Gaussian distribution
with a fixed mean $\mu_T$ and with a variance $\sigma_T$ that was scaled in proportion to the variance of the background
($\sigma_T^L = f\sigma_B^L$ and $\sigma_T^H = f\sigma_B^H$, respectively). At each timestep, we computed the Bhattacharyya coefficient ($BC$)
of the response distributions produced by background versus target stimuli: $BC = \sum \sqrt{p(r_B)p(r_T)}$. We used
$1 - BC$ as our measure of discriminability.

We simulated the behavior of this model using a background "probe" stimulus whose standard
deviation switched every $T$ timesteps. We simulated $N_c$ cycles of this probe stimulus, where each cycle
consisted of $T$ timesteps in the low state, followed by $T$ timesteps in the high state. This yielded timeseries
of the gain $k$ and offset $s_0$ of the adapting nonlinearity, as well as distributions of the neuronal response to
the background and target stimuli at each timepoint following a switch in standard deviation. We averaged
the gain and offset across cycles to obtain the average properties of the encoder at each timepoint following
a switch. We used the distribution of responses to target and background stimuli, measured across cycles,
to compute the discriminability at each timepoint following a switch. All simulations were performed with the
following values: $T = 50$, $N_c = 1,000$, $\mu_B = 0$, $\mu_T = 0$ to 3 in 0.25 steps, $\sigma_B^L = 1$, $\sigma_B^H = 3$, $f = 0.25$, $\sigma_n^2 =$
$0.01$, $N = 15$, $L = 12$. For Figure 1g, model discriminability in each contrast was fit with a logistic function to
estimate the sensitivity and threshold of the model. To approximate the stimulus conditions used in the
offset task, the target thresholds for each contrast were then used to select target levels to plot
discriminability over time ($\mu_T^L = 1.50$, $\mu_T^H = 2.25$; Figure 1f).

To compare behavioral performance to the model, we quantified the "psychometric performance" of
the model by fitting model discriminability over time with a logistic function:
681

$$y = \gamma + (1 - \gamma - \lambda) * \frac{1}{1 + e^{\alpha - \beta x}} \tag{1}$$

682
where $\alpha$ is the x-offset of the function, $\beta$ determined the sensitivity of the function, $\gamma$ determined the guess
rate (lower bound), $\lambda$ determined the lapse rate (upper bound) and $x$ was stimulus level. $\alpha/\beta$ determined the
threshold of this function, defined as the level corresponding to the steepest part of the curve. This function
was fit to behavioral or neuronal performance using constrained gradient descent ($fmincon$ in MATLAB)
initialized with a 10x10 grid-search of parameters $\alpha$ and $\beta$.

To characterize adaptation time constants, adaptation curves were fit with an exponential function

$$y = a + b * e^{-\frac{t}{\tau}} \tag{2}$$

689
690    where $a$ determined the y-offset of the function, $b$ was a multiplicative scaling factor, and $\tau$ was the
691    time constant of the exponential in units of time $t$. This function was fit to behavioral or neuronal responses
692    using constrained gradient descent initialized with a 10x10x10 grid search across all three parameters.

693    To measure the effect of stimulus contrast on parameters of interest, we computed a contrast
694    modulation index (CMI), which measured the relative change in the parameter between low and high contrast:
695

$$CMI = \frac{x_H - x_L}{x_L} \tag{3}$$

696
697    where $x$ is the parameter in question and subscripts $H$ and $L$ denote the value of that parameter in high and
698    low contrast, respectively. The resulting index is 0 if there is no change, 1 if $x_H$ is two times larger than $x_L$,
699    and is -0.5 if $x_L$ is two times $x_H$.

700
701
*Behavioral task*.
We used a GO/NO-GO task to measure the detectability of targets in background. In this task, each
trial consisted of a noise background with a contrast shift, along with the presence or absence of a target after
the change in contrast. Mice were trained to lick when they detected a target (hit), or to withhold licking in the

absence of a target (correct reject). This behavior was reinforced by providing a 4-5 uL water reward when
the mouse licked correctly (hit), and by initiating a 7-10 s timeout when the mouse licked in the absence of a
target (false alarm). Any licks detected during the timeout period resulted in the timeout being reset. In a
subset of mice, we introduced an additional trial abort period coincident with the first part of the contrast
background, before the contrast switch. Any licks detected in this abort period resulted in the trial being
repeated after a 7-10 s timeout, until the mouse withheld from licking during this period. In this task, misses
and correct rejects were not rewarded or punished. Trials were separated by a minimum 1.5s inter-trial-interval
(ITI). To discourage spontaneous licking, licks were monitored during this period, and if any licks occurred the
ITI timer was reset.

To prevent mice from predicting the target, we varied the timing of the target relative to the contrast
shift. This required a method for estimating hit rates and false alarm rates at different times during each trial,
and to reward and punish the animal during these times in an unbiased manner. To approach this issue, we
considered licks only during a 1 s response window after a target presentation (eg. if a target was presented
500 ms post-contrast-switch, the response window persisted from 500 to 1500 ms post-contrast-switch). To
apply this method to background-only trials, in which no targets were presented, we considered background
trials to be target trials containing infinitely small target amplitudes. For each background trial, we assigned a
response window with equiprobable delay matched to the target conditions and considered only licks within
those "target" response windows. Thus, over the course of a session, we randomly sampled lick probabilities
in background trials during the same temporal windows as those considered during target trials. Using this
scheme, we treated target and background-only trials identically, and estimated hit rates and false alarm rates
over time in an unbiased manner.

Each mouse performed three stages in the behavioral task: training, psychometric testing, and offset
testing. During the training task, trials consisted of two types, background-only trials or target trials presented
with equal probability. To facilitate learning, we selected target SNRs at the highest end of the range described
previously: in low contrast training sessions, targets were 16 dB SNR, and in high contrast training sessions,
targets were 20 dB SNR. To prevent response bias as a function of target timing, we randomly varied the
target delay between 250, 500, 750 and 1000 ms after the contrast change in each trial. During the
psychometric testing task, there were 7 trial types consisting of background-only trials and target trials
spanning six different SNRs (Extended Data Table 3). Based on behavioral piloting, we presented high SNR
trials with a greater probability, to ensure that mice were consistently rewarded during the task. In low and
high contrast psychometric sessions, the probability of a background trial was 0.4, the probability of the four
lowest target SNRs was 0.05 each, and the probability of the two highest target SNRs was 0.2 each. As in
training, target timing was varied randomly between 250, 500, 750 and 1000 ms after the contrast change in
each trial. After completing at least three sessions of the psychometric task, stimuli were generated for the
offset testing task. This task consisted of 15 unique trial types: 3 target levels (background trials, threshold
target trials, and high SNR target trials), and 5 target delays relative to the contrast change (25, 75, 225, 475,
975 ms delay). Threshold target amplitudes were determined individually for each mouse by fitting
performance averaged over several sessions with a psychometric function, and extracting the level at which
the slope of the psychometric curve was steepest. Based on behavioral piloting, background trials, threshold
target trials, and high SNR target trials were presented with probabilities of 0.4, 0.2, and 0.4, respectively.
Target delay on each trial was selected with equal probability. In all behavioral stages, trial order was
pseudorandomly generated, such that there were no more than three target or background trials in a row.

A subset of mice (n = 2), were presented targets in silence (Figure 4). To generate this stimulus set
without changing the basic structure of the task or stimuli, we simply took the spectrograms of all stimuli
containing 25 dB SNR targets from the low-to-high contrast stimulus sessions, and set the stimulus power
flanking each target to zero. This manipulation was only performed in the target period, and the low contrast
adaptation period of the trials remained the same. Thus, the targets and adaptation periods were identical to
those presented in the target-in-background task. To vary the difficulty of the task, the level of the target was
attenuated using the following values: -75, -60, -45, -30, -15, and 0 dB attenuation relative to the 25 dB SNR
target. Mice were previously trained in the target-in-background task prior to performing the target in silence
task. Before psychometrically varying the target attenuation, mice were trained in the new task to criterion
performance. Mice generalized very rapidly to the new task, reaching 97% and 94% training accuracy on the
first day of exposure to targets in silence (mice CA124 and CA125, respectively).

759
760 *Chronic muscimol application*.

761        A separate cohort of mice (n = 4) were bilaterally implanted with 26 GA guide cannulae (PlasticsOne, C315GMN-SPC mini, cut 5 mm below pedestal) in auditory cortex. The surgery was performed as described previously with the following modifications. After the skull was leveled using a stereotax, two small craniotomies were performed -2.6 mm anterior, ±4.3 mm lateral from bregma, over auditory cortex. The guide cannulae and dummy infusion cannulae (PlasticsOne, C315DCMN-SPC mini, cut to fit 5 mm C315GMN with a 0.5 mm projection depth) were sterilized in an autoclave. The dummy cannulae were partially screwed into the guide cannulae and placed in a stereotaxic clamp. After zeroing the tip of the guide cannula to the brain surface, the cannula was lowered to 500 µm below the cortical surface. This depth was chosen because the infusion cannulae (PlasticsOne, C315LIMN-SPC mini) project 500 µm from the end of the guide cannulae when completely inserted, leading to a final depth of 1000 µm – the location of auditory cortex. The dummy cannulae were then fully inserted and this procedure was repeated for the next cortical hemisphere.

772        Prior to injecting, two injection syringes (Hamilton Syringe, 10µL Gaslight #1701) and tubing (C313CT tubing 023x050 PE50) were backfilled with mineral oil. Sterilized infusion cannulae were then attached to each syringe and ~500nL of muscimol (diluted with 1x PBS to .25 mg/mL; Sigma Aldrich, M1523) or 0.9% sterile saline was drawn up into the injection cannulae using a dual injector (Harvard Apparatus, Pump 11 Pico Plus Elite). The mouse was then headfixed and the dummy cannulae were removed and sterilized. The loaded infusion cannulae were then screwed all the way into the guide cannulae and 400 nL of muscimol or saline was infused bilaterally at a rate of 250 nL/minute. The infusion cannulae were then replaced with the dummy cannulae and the mouse rested in its home cage for 30-45 minutes before beginning the behavioral session.

*Acute electrophysiological recordings.*

783        For acute recordings used to fit the GC-GLM model (Figure 2), neuronal signals were recorded from n = 1 awake, untrained mouse. Prior to the recording session, the mouse was anesthetized and a headpost and ground pin were implanted on the skull (see *Surgery*). On the day of the recording, the mouse was briefly anesthetized with 3% isoflurane and a small craniotomy was performed over auditory cortex using a dental drill or scalpel (~1 mm x 1 mm craniotomy centered approximately 1.25 mm anterior to the lambdoid suture along caudal end of the squamosal suture). A 32 channel silicon probe (Neuronexus) was then positioned perpendicularly to the cortical surface and lowered at a rate of 1-2 µm/s to a final depth of 800-1200 µm. As the probe was lowered, trains of brief noise bursts were repeated, and if stimulus locked responses to the noise bursts were observed, the probe was determined to be in auditory cortex. The probe was then allowed to settle for up to 30 minutes before starting the recording. Neuronal signals were amplified and digitized with an Intan headstage (RHD 32ch) and recorded by an openEphys acquisition board[85] at a rate of 30 kHz.

794        For this experiment, the mouse was presented with 3 s DRCs alternating between low and high contrast (uniform distribution with a mean of 50 dB and a width of ±5 dB in low contrast or ±15 dB in high contrast at a chord rate of 25 ms, as described in *Stimuli*). In order to accurately fit the GLM in an unbiased manner, these stimuli were highly random, composed of 100 unique chord patterns for each contrast (Extended Data Figure 2i,j**)**. For each of the two recording sites, 5 repeats of this stimulus set were played.

*Behavioral electrophysiological recordings*.

801        Neuronal signals were acquired from awake, behaving mice as they performed the psychometric and offset testing tasks described previously. Chronically implanted, 16-, 32-, or 64-channel microdrives[85,86] were connected to one or two 32 channel Intan amplifier headstages. Amplified signals were recorded at 30 kHz using an openEphys acquisition board via an SPI cable, where the signals were digitized.

805        For all recordings, broadband signals were filtered between 500 and 6000 Hz, offset corrected, and re-referenced to the median across all active channels. The preprocessed data was then sorted using KiloSort[87] or KiloSort2 and the resulting clustering was manually corrected in phy2 according to community-developed guidelines. The resulting units were labelled as single units if they exhibited a clear refractory period and did not need to be split. Splitting assessments were made through manual examination of principle component features for the two best channels of a cluster. If two noticeable clusters in feature space were evident in a unit, the unit was either manually split, or classified as a multiunit.

*Generalized linear model.*

814        To justify the form of GLM used here, we discuss a how a model neuron could implement gain control in the simplest terms, and then structure our inference model to extract the parameters of this model neuron. We will assume that the activity of the model neuron is driven by three sources: 1) stimulus drive, 2) stimulus

817  contrast, and 3) the multiplicative interaction between the two, which we use to define the gain (for a formal
818  definition of this forward model and the inference model, see *Supplementary Information*).
819  As discussed previously, the stimulus used in our experiments is composed of many frequencies that change
820  in loudness in discrete time steps:
821
822
$$X_{t,f} \sim \mathcal{U}(\mu, \sigma_t)$$
823
824  where $X_{t,f}$ is the stimulus spectrogram that varies as a function of time $t$ and frequency $f$. Each time and
825  frequency bin of $X$ is sampled from a uniform distribution defined by an average value $\mu$ and contrast $\sigma_t$.
826  We assume that the hypothetical neuron responds selectively at some frequency and time lag, defined
827  by a filter, or STRF $\beta_{h,f}$ with history $h$ and frequency $f$ components. Given $\beta$, we can define the stimulus drive
828  $x_t$ as

$$x_t = X_t\beta \qquad (4)$$

829
where at each time $t$, $X_t$ is a row vector of size $F$ frequencies times $H$ lags (ie. the "unrolled" lagged stimulus
spectrogram) and $\beta$ is the STRF unrolled to a single column vector of the same size.
In the spirit of efficient coding theory, and as shown in previous work, we assume that the gain $g$ of the neuron
should be inversely proportional to the contrast, such that $g(\sigma) \propto 1/\sigma$ (ie. when contrast is low gain should be
high, and vice-versa). We also define "neutral" gain to be the average of the gain of the neuron in low and
high contrast. Putting these two features together, we can summarize the gain of the neuron as
836

$$g(\sigma) = \frac{\overline{\sigma}}{\sigma_t} \qquad (5)$$

837
where $\overline{\sigma}$ is the harmonic mean of the contrast in the low and high conditions (see *Supplementary Information*).
In the case of a 3-fold change in contrast, this function constrains the gain of the neuron between 0.5 and 1.5,
with a neutral value of 1. As mentioned previously, we consider gain to be the multiplicative interaction
between the stimulus drive and the contrast, such that the contribution of gain control to the response of the
neuron is related to $x_t \cdot \frac{\overline{\sigma}}{\sigma_t}$.
To summarize, we considered a hypothetical neuron driven by the stimulus according to a STRF $\beta$
and by the interaction between the stimulus drive and the contrast $x_t \cdot \frac{\overline{\sigma}}{\sigma_t}$. To infer the relative weights of each
of these components of the neuronal response, we defined a Poisson GLM with an intercept term and the
following predictors:

$$x_t, \qquad \frac{\overline{\sigma}}{\sigma_t}, \qquad x_t \cdot \frac{\overline{\sigma}}{\sigma_t} \qquad (6)$$

847
848  In other words, the model is composed of a stimulus predictor $x_t$, a contrast predictor $\overline{\sigma}/\sigma_t$, and their
849  interaction. Therefore, the GLM models the firing rate $\lambda$ at time $t$ as a Poisson distribution with the following
850  mean:

$$\lambda_t = exp\left[\beta_0 + x_t\beta_1 + x_t \cdot \frac{\overline{\sigma}}{\sigma_t}\beta_2 + \frac{\overline{\sigma}}{\sigma_t}\beta_3\right] \qquad (7)$$

851
852  where $\beta_0 \dots \beta_3$ are the parameters to be inferred. Based on our behavioral data (Figure 3) and the predictions
853  of the efficient coding model (Figure 1), we expected the influence of contrast on neuronal gain to be
854  asymmetric and smooth. To enable the GLM to capture both of these qualities, we first defined the contrast
855  predictors from a set of cubic B-spline temporal basis functions, then defined separate contrast predictors for
856  transitions to low and high contrast. Incorporating these changes, we can redefine equation 4 above as

$$ln\lambda_t = \beta_0 + x_t\beta_1 + x_t \circ C_t'\beta_2 + C_t'\beta_3 \qquad (8)$$

857
where $\circ$ denotes element-by-element "broadcasting" multiplication and $C_t'$ is a matrix of contrast predictors
$\overline{\sigma}/\sigma_t$ convolved with a set of basis functions and separated by contrast transitions (see *Supplementary*
*Information*). For the sake of clarity, note that in the expression above, $\beta_0$ is a number, $x$ is a column vector

of length $T$, $\beta_1$ is a number, $C'$ is a $T$-by-$2B$ matrix, and $\beta_2$ and $\beta_3$ are column vectors of length $2B$, where $B$
is the number of splines.
So far, we outlined a hypothetical neuron which implements gain control, and a GLM with which we
can approximate the behavior of this neuron. Next, we describe how to use the fitted parameters to quantify
the gain of the neuron. Conceptually, an increase or decrease in the gain of the neuron is analogous to more
or less sensitivity to small changes in the stimulus. Based on this intuition, we focused on how the response
of the neuron (as modelled by a fitted GLM) is expected to change between conditions where the gain is
expected to contribute (ie. in the presence of gain control) and where it is not (ie. in the absence of gain
control, where gain is "neutral"). Following this logic, we derived a definition for gain $w_t$ as the ratio between
the sensitivity of the fitted model with changes in contrast, compared to the sensitivity of the same model
when the contrast is at a reference value, which we defined previously as $w_t = 1$:

$$w_t = \frac{\beta_1 + C_t'\beta_2}{\beta_1 + C^0\beta_2} \tag{9}$$

872
873      where $w_t$ is the estimated gain at time $t$, and $C^0$ is a reference contrast design matrix identical to $C_t'$ except
874      that all non-zero elements are set to 1 (see *Supplementary Information* for full derivation of $w_t$).
875          To fit the model, we implemented a two-step procedure. In the first step, the STRF $\beta$ of the neuron
876      was estimated according to the model

$$\ln \lambda_t = \alpha + X_t\beta \tag{10}$$

877
For the second step, we calculated the stimulus drive as described in equation 4, and then fit equation 8 to
the data for each neuron using $\mathrm{glmfit}$ in MATLAB. This entire fitting procedure was 10-fold cross-validated
with folds stratified across trials of each contrast. In the first step, we fit the STRF $\beta$ with $F$ frequency bins
according to the stimulus spectrogram ($F$ = 33 or 34, see *Stimuli*) and a history window of 300 ms ($H$ = 12).
When fitting the full model, we defined the contrast design matrix $C_t'$ to capture 1000 ms of contrast history
around each transition ($H'$ = 40), convolved with a set of B-spline temporal basis functions[88] (here, we used
B-splines with a degree of 3 and 3 equally-spaced knots, constrained to go smoothly to zero at the longest
lag, which implied that $B$ = 4).
To validate the model, we first simulated neurons according to the forward model outlined above
(Extended Data Figure 2a) while varying the amount of gain control and the temporal trajectory of gain in
different simulation runs. We found that the GLM accurately predicted the STRF shape, spike rates and gain
trajectories across a variety of simulation parameters (Extended Data Figure 2c, e-h). For a detailed
description and discussion of the simulation results, see *Supplementary Information* and Extended Data Table
4.
892
*Behavioral and neuronal detection performance.*
To calculate performance in the target-in-background detection task we adopted commonly used
signal detection theory methods[38,89] to estimate the ability of an ideal observer to discriminate between two
sensory distributions: in our case, a distribution for target trials and a distribution for background trials. When
analyzing behavior, we computed the percent correct performance of an ideal observer[90] as a function of the
probability of hits and false alarms:
899

$$pc = z^{-1}\left(\frac{z(H) - z(FA)}{\sqrt{2}}\right) \tag{11}$$

900
where $z^{-1}$ is cumulative probability of the normal distribution ($\mathrm{normcdf}$ in MATLAB), $z$ is the inverse of the
normal distribution (ie. the z-score, $\mathrm{norminv}$ in MATLAB), $H$ is the hit rate, and $FA$ is the false alarm rate. For
psychophysical performance, hit rates and false alarm rates near 0 and 1 were adjusted using the log-linear
rule[91], to reduce biases in performance estimation resulting from low numbers of trials.
To calculate neuronal performance in the same reference frame as the behavior, we employed similar
ideal observer techniques. First, neuronal responses (either spike rates of single units, or population projection
values), were averaged in a 100ms window post target onset (for background trials, this window was randomly
chosen on each trial to coincide with target presentation times on target trials). Then, using the distributions
of responses during target and background trials, we computed receiver-operating-characteristic curves and
took the area under the curve (AUC) as the percent correct of an ideal observer discriminating between the

target and background distributions. To determine whether the AUC value for a given set of trial distributions was significantly different from chance, we performed a bootstrap procedure where we sampled from all the trials with replacement 500 times and recomputed AUC for each sample. If the 95% confidence intervals for this bootstrapped distribution did not include chance (.5), we defined that AUC value as significant. For population analyses which generated single-trial predictions, neuronal hit and false alarm rates were transformed to percent correct as described above.

*Population response metrics.*

On sessions where three or more neurons were simultaneously recorded, we used a population vector technique[26] to estimate the ability of neuronal populations to discriminate targets from background. First, spike rates in each trial were averaged in a 100ms window post-target onset. Then, using a leave-one-out procedure, we computed a trial averaged population vector for target trials, $v_T$, and a separate average population vector for background trials, $v_B$. We then estimated the coding direction in high dimensional neuronal space that best separated the target and background responses: $CD = v_T - v_B$. The held out trial was then projected along this dimension, by taking the population response vector on that trial $v_{trial}$ and projecting it along the estimated coding direction using the dot product: $projection\ value = v_{trial} * CD$. This procedure was repeated holding out each trial, and estimating the coding direction from the remaining trials. For psychometric testing sessions, the target responses from the two loudest target levels were used to estimate coding direction, and in offset testing sessions the target responses from the high SNR target trials were used. After computing projections for every trial, the resulting matrix was normalized between 0 and 1.

*Population classifier.*

Based on previously described methods[27], we used a criterion-based decision rule to estimate how a hypothetical down-stream neuron may read out the neuronal activity of a population of neurons. As before, trial distributions of neuronal responses to targets or background were created from the average activity in a 100ms window post-target. Then, we sampled 100 criterion values between the minimum and maximum response, and for each criterion estimated the proportion of correct trials under two decision rules: 1) report target present if the response is greater than the criterion, or, 2) report target present if the response is less than the criterion. By assessing these two decision rules, neurons that were suppressed by target presence were treated equally to neurons that were enhanced by target presence. Finally, we chose the criterion and decision rule that yielded the highest proportion of correct trials, and computed neuronal hit rates and false alarm rates for each target level, and background-only trials. These hit rates and false alarm rates were then transformed to percent correct according to Equation 8.

*Linear-nonlinear model.*

First, we selected only neurons in the dataset which had reliable responses to stimulus repeats. To determine response reliability, we computed a noise ratio (NR) for each neuron, which describes the amount of variability in the response due to noise versus the amount of variability in the response driven by the stimulus[92,93]. Values approaching 0 indicate increasingly reliable responses to the stimulus, so for the remaining analyses, we included neurons with NR < 100.

The linear nonlinear model was composed of a spectrotemporal receptive field (STRF) and a set of rectifying nonlinearities. The STRF β was fit using normalized reverse correlation

$$\beta = [XX^T]^{-1}X\lambda \tag{1}$$

where $X$ is the stimulus design matrix $X_t$ defined in equation 4 and $\lambda$ is the spike count in each 25 ms bin of the DRC stimulus. When defining $X$, we used a history window of 300 ms ($H$ = 12) and frequency bins corresponding to the frequencies composing the dynamic random chords (see *Stimuli*). After fitting the STRF, we fit the nonlinearities of the neuron. This two-step fitting procedure was repeated using 10 fold cross-validation, as described below.

For each fold, we selected 90% of the trials for training, leaving the remaining 10% to be held out for testing. Within each trial, we excluded neuronal responses around transitions from silence, or transitions in contrast, to prevent the model from overfitting strong transients in the neuronal response. Additionally, we excluded neuronal responses within a 50ms window after target presentation, to prevent overfitting of target

responses. Given these exclusion criteria, we calculated the duration of stimulus sampled in the target period for each trial, and, for each trial, sampled the same duration of stimulus within the adaptation period. This procedure ensured that the model was fit to the same amount of high and low contrast stimulation per trial, to minimize overfitting to one contrast condition. Then, a stimulus design matrix $X$ was defined using these stimulus periods, and the STRF was fit using equation 12. We tested whether STRF properties were affected by stimulus contrast, and found STRFs to be largely stable when estimated separately for each contrast (*Supplementary Information* and Extended Data Figure 7). Therefore, we used both periods of contrast to estimate β.

Using the STRF fit to the training data, we computed the linear drive $x_t$ by convolving the STRF with the lagged spectrogram of the training stimulus (equation 4). For the GC-LN model we separated the linear predictions into low and high contrast periods, while for the static-LN model all matched time points were used. We generated a histogram of the linear prediction values (50 bins), and for each bin, computed the mean spike rate of the neuron when the linear prediction fell within those bin edges (Figure 6d, scatter points). The resulting set of linear prediction values and average spike rates were fit with an exponential function:

$$y = a + be^{c(x-d)} \tag{2}$$

where $a$ determined the minimum firing rate, $b$ was a multiplicative scaling factor, $c$ determined the gain of the exponent, and $d$ determined the x-offset, or firing threshold of the neuron. This function was fit to each cell using constrained gradient descent ($fmincon$ in MATLAB), using a 10x10 grid search for parameters $b$ and $c$. The gain for each neuron was defined as $c$. This entire process was repeated for each cross-validation fold, and the final parameter estimates for the STRF and nonlinearities were taken as the average over the 10 runs.

To determine the relationship between neuronal gain and behavioral performance, we computed the average neuronal gain across all noise responsive neurons (NR < 100) in each session for the adaptation and target periods in the trial. We then compared the session-averaged gain values to the fitted thresholds and slopes of the psychometric curve across sessions using the mixed-effects linear models outlined in the main text.

*Inclusion criteria*.

Unless otherwise noted, behavioral sessions in which the false alarm rate exceeded 50% were discarded from analysis. One mouse (ID: CA122) had consistently high false alarm rates in the high contrast condition, so we excluded high contrast sessions from this mouse from all analyses. For Figures 5 and 6, we removed neurons with low spike rates (<1Hz) and noise-like or inverted (ie. upward inflected) spike waveforms. To determine waveform quality, we computed the width of each waveform at half of the minimum value (FWHM) and its correlation with the average waveform over all neurons. Neurons whose waveforms had outlier FWHM values ($isoutlier$ in MATLAB), negative correlations, or were not significantly correlated with the average (Bonferoni corrected $p > 5.85e-6$) were removed from further analysis. For Figure 5f-l, sessions with stable population decoding performance were included (defined as sessions where more than half of the target times elicited significant population AUC values, as determined by the bootstrap procedure described previously). For Figure 6e-h, only neurons with noise ratios less than 100 were included in all analyses.

**Data Availability.**

All data including sorted spike times from electrophysiological recordings and behavioral data are available on Dryad: https://doi.org/10.5061/dryad.6djh9w120. Every Figure and Extended Data Figure have associated data with the following exceptions. Figure 1, Extended Data Figure 1, and Extended Data Figure 2 are the results of simulations and thus have no associated raw data. Simulation parameters can be found in the code. There are no restrictions on data availability.

**Code Availability.**

Code for behavioral and neural analysis: https://github.com/chris-angeloni/contrast_behavior. For the GC-GLM model and simulations: https://github.com/chris-angeloni/contrast_glm.

**Acknowledgements.**

018    This work was supported by NIDCD R01DC015527, R01DC014479, NINDS R01NS113241 to
019    MNG and NIDCD F31DC016524 to CFA. CFA and AMW were supported by NIDCD T32DC016903. The
020    authors would also like to thank Dr. Yale Cohen, Dr. Nicole Rust, and Dr. David Brainard for their helpful
021    feedback on this study and Dashiell Ridolfi-Starr, Karmi Oxman, Nitay Caspi, Tyler Ling, and Margaret
022    DenBleyker for experimental assistance.

1. Barlow, H. B. Possible Principles Underlying the Transformations of Sensory Messages. in *Sensory Communication* vol. 6 216–234 (2013).

2. Brenner, N., Bialek, W. & De Ruyter Van Steveninck, R. Adaptive rescaling maximizes information transmission. *Neuron* **26**, 695–702 (2000).

3. Bharioke, A. & Chklovskii, D. B. Automatic Adaptation to Fast Input Changes in a Time-Invariant Neural Circuit. *PLoS Comput Biol* **11**, 1004315 (2015).

4. Borst, A. & Theunissen, F. E. Information theory and neural coding. *Nature Neuroscience* vol. 2 947–957 (1999).

5. Baccus, S. A. & Meister, M. Fast and slow contrast adaptation in retinal circuitry. *Neuron* **36**, 909–919 (2002).

6. Dean, I., Harper, N. S. & McAlpine, D. Neural population coding of sound level adapts to stimulus statistics. *Nat. Neurosci.* **8**, 1684–1689 (2005).

7. Lesica, N. A. *et al.* Adaptation to Stimulus Contrast and Correlations during Natural Visual Stimulation. *Neuron* **55**, 479–491 (2007).

8. Gutnisky, D. A. & Dragoi, V. Adaptive coding of visual information in neural populations. *Nature* **452**, 220–224 (2008).

9. Wen, B., Wang, G. I., Dean, I. & Delgutte, B. Dynamic range adaptation to sound level statistics in the auditory nerve. *J. Neurosci.* **29**, 13797–13808 (2009).

10. Dahmen, J. C., Keating, P., Nodal, F. R., Schulz, A. L. & King, A. J. Adaptation to Stimulus Statistics in the Perception and Neural Representation of Auditory Space. *Neuron* **66**, 937–948 (2010).

11. Wen, B., Wang, G. I., Dean, I. & Delgutte, B. Time course of dynamic range adaptation in the auditory nerve. *J. Neurophysiol.* **108**, 69–82 (2012).

12. Clarke, S. E., Longtin, A. & Maler, L. Contrast coding in the electrosensory system: Parallels with visual computation. *Nature Reviews Neuroscience* vol. 16 733–744 (2015).

13. Clemens, J., Ozeri-Engelhard, N. & Murthy, M. Fast intensity adaptation enhances the encoding of sound in Drosophila. *Nat. Commun.* **9**, 1–15 (2018).

14. Rabinowitz, N. C., Willmore, B. D. B., Schnupp, J. W. H. & King, A. J. Contrast Gain Control in Auditory Cortex. *Neuron* **70**, 1178–1191 (2011).

15. Rabinowitz, N. C., Willmore, B. D. B. B., Schnupp, J. W. H. H. & King, A. J. Spectrotemporal Contrast Kernels for Neurons in Primary Auditory Cortex. *J. Neurosci.* **32**, 11271–11284 (2012).

16. Rabinowitz, N. C., Willmore, B. D. B., King, A. J. & Schnupp, J. W. H. Constructing Noise-Invariant Representations of Sound in the Auditory Pathway. *PLoS Biol.* **11**, e1001710 (2013).

17. Cooke, J. E., King, A. J., Willmore, B. D. B. & Schnupp, J. W. H. Contrast gain control in mouse auditory cortex. *J. Neurophysiol.* **120**, 1872–1884 (2018).

18. Cooke, J. E. *et al.* Contrast gain control occurs independently of both parvalbumin-positive interneuron activity and shunting inhibition in auditory cortex. *J. Neurophysiol.* **123**, 1536–1551 (2020).

19. Lohse, M., Bajo, V. M., King, A. J. & Willmore, B. D. B. Neural circuits underlying auditory contrast gain control and their perceptual implications. *Nat. Commun.* **11**, 1–13 (2020).

20. Willmore, B. D. B., Cooke, J. E. & King, A. J. Hearing in noisy environments: Noise invariance and contrast gain control. *J. Physiol.* **592**, 3371–3381 (2014).

21. Maier, J. K. *et al.* Adaptive coding is constrained to midline locations in a spatial listening task. *J. Neurophysiol.* **108**, 1856–1868 (2012).

22. Młynarski, W. F. & Hermundstad, A. M. Adaptive coding for dynamic sensory inference. *Elife* **7**, (2018).

23. Młynarski, W. F. & Hermundstad, A. M. Efficient and adaptive sensory codes. *Nat. Neurosci.* **24**, 998–1009 (2021).

24. Pennington, J. R. & David, S. V. Complementary effects of adaptation and gain control on sound encoding in primary auditory cortex. *eNeuro* **7**, 1–17 (2020).

25. DeWeese, M. & Zador, A. Asymmetric Dynamics in Optimal Variance Adaptation. *Neural Comput.* **10**, 1179–1202 (1998).

26. Li, N., Daie, K., Svoboda, K. & Druckmann, S. Robust neuronal dynamics in premotor cortex during motor planning. *Nature* **532**, 459–464 (2016).

27. Christison-Lagay, K. L., Bennur, S. & Cohen, Y. E. Contribution of spiking activity in the primary auditory cortex to detection in noise. *J. Neurophysiol.* **118**, 3118–3131 (2017).

28. Talwar, S. K., Musial, P. G. & Gerstein, G. L. *Role of mammalian auditory cortex in the perception of*

*elementary sound properties. Journal of Neurophysiology* vol. 85 www.jn.org (2001).

29.   Gimenez, T. L., Lorenc, M. & Jaramillo, S. Adaptive categorization of sound frequency does not
require the auditory cortex in rats. *J. Neurophysiol.* **114**, 1137–1145 (2015).

30.   Jaramillo, S. & Zador, A. M. The auditory cortex mediates the perceptual effects of acoustic temporal
expectation. in *Nature Neuroscience* vol. 14 246–253 (2011).

31.   Wood, K. C., Town, S. M., Atilgan, H., Jones, G. P. & Bizley, J. K. Acute inactivation of primary
auditory cortex causes a sound localisation deficit in ferrets. *PLoS One* **12**, (2017).

32.   Kato, H. K., Gillet, S. N. & Isaacson, J. S. Flexible Sensory Representations in Auditory Cortex Driven
by Behavioral Relevance. *Neuron* **88**, 1027–1039 (2015).

33.   Ceballo, S., Piwkowska, Z. & Bourg, J. Targeted Cortical Manipulation of Auditory Perception In Brief.
*Neuron* **104**, 1168-1179.e5 (2019).

34.   Li, Z. *et al.* Corticostriatal control of defense behavior in mice induced by auditory looming cues. *Nat.*
*Commun.* **12**, 1–13 (2021).

35.   Town, S., Wood, K. & Bizley, J. Signal processing in auditory cortex underlies degraded speech
sound discrimination in noise. *bioRxiv* 833558 (2019) doi:10.1101/833558.

36.   Musall, S., Urai, A. E., Sussillo, D. & Churchland, A. K. Harnessing behavioral diversity to understand
neural computations for cognition. *Current Opinion in Neurobiology* vol. 58 229–238 (2019).

37.   Shadlen, M. N. & Kiani, R. Decision making as a window on cognition. *Neuron* vol. 80 791–806
(2013).

38.   Newsome, W. T., Britten, K. H. & Movshon, J. A. Neuronal correlates of a perceptual decision. *Nature*
**341**, 52–54 (1989).

39.   Britten, K. H. *et al.* The analysis of visual motion: a comparison of neuronal and psychophysical
performance. *J. Neurosci.* **12**, 4745–4765 (1992).

40.   Shadlen, M. N., Britten, K. H., Newsome, W. T. & Movshon, J. A. A computational analysis of the
relationship between neuronal and behavioral responses to visual motion. *J. Neurosci.* **16**, 1486–
1510 (1996).

41.   Nienborg, H. & Cumming, B. G. Decision-related activity in sensory neurons reflects more than a
neurons causal effect. *Nature* **459**, 89–92 (2009).

42.   Cumming, B. G. & Nienborg, H. Feedforward and feedback sources of choice probability in neural
population responses. *Current Opinion in Neurobiology* vol. 37 126–132 (2016).

43.   Tsunada, J., Liu, A. S. K., Gold, J. I. & Cohen, Y. E. Causal contribution of primate auditory cortex to
auditory perceptual decision-making. *Nat. Neurosci.* **19**, 135–142 (2015).

44.   Steinmetz, N. A., Zatka-Haas, P., Carandini, M. & Harris, K. D. Distributed coding of choice, action
and engagement across the mouse brain. *Nature* **576**, 266–273 (2019).

45.   Cohen, M. R. & Newsome, W. T. Context-Dependent Changes in Functional Circuitry in Visual Area
MT. *Neuron* **60**, 162–173 (2008).

46.   Cohen, M. R. & Newsome, W. T. Estimates of the contribution of single neurons to perception
depend on timescale and noise correlation. *J. Neurosci.* **29**, 6635–6648 (2009).

47.   Ni, A. M., Ruff, D. A., Alberts, J. J., Symmonds, J. & Cohen, M. R. Learning and attention reveal a
general relationship between population activity and behavior. *Science (80-. ).* **359**, 463–465 (2018).

48.   Downer, J. D., Niwa, M. & Sutter, M. L. Task engagement selectively modulates neural correlations in
primary auditory cortex. *J. Neurosci.* **35**, 7565–7574 (2015).

49.   Stringer, C., Michaelos, M., Tsyboulski, D., Lindo, S. E. & Pachitariu, M. High-precision coding in
visual cortex. *Cell* (2021) doi:10.1016/j.cell.2021.03.042.

50.   Hires, S. A., Gutnisky, D. A., Yu, J., O'Connor, D. H. & Svoboda, K. Low-noise encoding of active
touch by layer 4 in the somatosensory cortex. *Elife* **4**, (2015).

51.   Hobbs, J. A., Towal, R. B. & Hartmann, M. J. Z. Spatiotemporal patterns of contact across the rat
vibrissal array during exploratory behavior. *Front. Behav. Neurosci.* **9**, 356 (2016).

52.   Aizenberg, M. & Geffen, M. N. Bidirectional effects of aversive learning on perceptual acuity are
mediated by the sensory cortex. *Nat. Neurosci.* **16**, 994–6 (2013).

53.   Aizenberg, M., Mwilambwe-Tshilobo, L., Briguglio, J. J., Natan, R. G. & Geffen, M. N. Bidirectional
Regulation of Innate and Learned Behaviors That Rely on Frequency Discrimination by Cortical
Inhibitory Neurons. *PLoS Biol.* **13**, e1002308 (2015).

54.   Briguglio, J. J., Aizenberg, M., Balasubramanian, V. & Geffen, M. N. Cortical neural activity predicts
sensory acuity under optogenetic manipulation. *J. Neurosci.* **38**, 2094–2105 (2018).

55.   Wood, K. C., Angeloni, C. F., Oxman, K., Clopath, C. & Geffen, M. N. Neuronal activity in sensory

cortex predicts the specificity of learning. *bioRxiv* 2020.06.02.128702 (2020)
doi:10.1101/2020.06.02.128702.
56. Ulanovsky, N., Las, L. & Nelken, I. Processing of low-probability sounds by cortical neurons. *Nat*
*Neurosci* **6**, 391–398 (2003).
57. Natan, R. G., Carruthers, I. M., Mwilambwe-Tshilobo, L. & Geffen, M. N. Gain Control in the Auditory
Cortex Evoked by Changing Temporal Correlation of Sounds. *Cereb. Cortex* **27**, 2385–2402 (2017).
58. Espejo, M. L., Schwartz, Z. P. & David, S. V. Spectral tuning of adaptation supports coding of sensory
context in auditory cortex. *PLoS Comput. Biol.* **15**, e1007430 (2019).
59. Fritz, J., Shamma, S., Elhilali, M. & Klein, D. Rapid task-related plasticity of spectrotemporal receptive
fields in primary auditory cortex. *Nat. Neurosci.* **6**, 1216–1223 (2003).
60. Mesgarani, N., Fritz, J. & Shamma, S. A computational model of rapid task-related plasticity of
auditory cortical receptive fields. *J. Comput. Neurosci.* **28**, 19–27 (2010).
61. David, S. V., Fritz, J. B. & Shamma, S. A. Task reward structure shapes rapid receptive field plasticity
in auditory cortex. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 2144–9 (2012).
62. Yin, P., Fritz, J. B. & Shamma, S. A. Rapid spectrotemporal plasticity in primary auditory cortex
during behavior. *J. Neurosci.* **34**, 4396–408 (2014).
63. Niwa, M., Johnson, J. S., O'Connor, K. N. & Sutter, M. L. Active engagement improves primary
auditory cortical Neurons' ability to discriminate temporal modulation. *J. Neurosci.* **32**, 9323–9334
(2012).
64. Fritz, J. B., Elhilali, M. & Shamma, S. A. Adaptive changes in cortical receptive fields induced by
attention to complex sounds. *J. Neurophysiol.* **98**, 2337–46 (2007).
65. Schneider, D. M., Nelson, A. & Mooney, R. A synaptic and circuit basis for corollary discharge in the
auditory cortex. *Nature* **513**, 189–94 (2014).
66. Schneider, D. M., Sundararajan, J. & Mooney,  richard. A cortical filter that learns to suppress the
acoustic consequences of movement. *Nature* (2018) doi:10.1038/s41586-018-0520-5.
67. McGinley, M. J., David, S. V. & McCormick, D. A. Cortical Membrane Potential Signature of Optimal
States for Sensory Signal Detection. *Neuron* **87**, 179–192 (2015).
68. Reimer, J. *et al.* Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex.
*Nat. Commun.* **7**, 1–7 (2016).
69. Atallah, B. V., Bruns, W., Carandini, M. & Scanziani, M. Parvalbumin-Expressing Interneurons
Linearly Transform Cortical Responses to Visual Stimuli. *Neuron* **73**, 159–170 (2012).
70. Wilson, N. R., Runyan, C. A., Wang, F. L. & Sur, M. Division and subtraction by distinct cortical
inhibitory networks in vivo. *Nature* **488**, 343–348 (2012).
71. Seybold, B. A., Phillips, E. A. K., Schreiner, C. E. & Hasenstaub, A. R. Inhibitory Actions Unified by
Network Integration. *Neuron* **87**, 1181–1192 (2015).
72. Phillips, E. A. K. & Hasenstaub, A. R. Asymmetric effects of activating and inactivating cortical
interneurons. *Elife* **5**, e18383 (2016).
73. Reynolds, J. H. & Heeger, D. J. The Normalization Model of Attention. *Neuron* **61**, 168–185 (2009).
74. Natan, R. G. *et al.* Complementary control of sensory adaptation by two types of cortical
interneurons. *Elife* **4**, 163–174 (2015).
75. Natan, R. G., Rao, W. & Geffen, M. N. Cortical Interneurons Differentially Shape Frequency Tuning
following Adaptation. *Cell Rep.* **21**, 878–890 (2017).
76. Attneave, F. Some informational aspects of visual perception. *Psychol. Rev.* **61**, 183–193 (1954).
77. Simoncelli, E. P. & Olshausen, B. A. Natural image statistics and neural representation. *Annual*
*Review of Neuroscience* vol. 24 1193–1216 (2001).
78. Simoncelli, E. P. Vision and the statistics of the visual environment. *Current Opinion in Neurobiology*
vol. 13 144–149 (2003).
79. Młynarski, W., Hledík, M., Sokolowski, T. R. & Tkačik, G. Statistical analysis and optimality of neural
systems. *Neuron* **109**, 1227-1241.e5 (2021).
80. Wei, X.-X. & Stocker, A. A. A Bayesian observer model constrained by efficient coding can explain
'anti-Bayesian' percepts. *Nat. Neurosci.* **18**, 1509–1517 (2015).
81. Guo, Z. V. *et al.* Procedures for behavioral experiments in head-fixed mice. *PLoS One* **9**, (2014).
82. Isett, B. R., Feasel, S. H., Lane, M. A. & Feldman, D. E. Slip-Based Coding of Local Shape and
Texture in Mouse S1. *Neuron* **97**, 418-433.e5 (2018).
83. Carruthers, I. M., Natan, R. G. & Geffen, M. N. Encoding of ultrasonic vocalizations in the auditory
cortex. *J Neurophysiol* **109**, 1912–1927 (2013).

84.   Carruthers, I. M. *et al.* Emergence of invariant representation of vocalizations in the auditory cortex. *J.*
*Neurophysiol.* jn.00095.2015 (2015) doi:10.1152/jn.00095.2015.
85.   Voigts, J. *et al.* An easy-to-assemble, robust, and lightweight drive implant for chronic tetrode
recordings in freely moving animals. *J. Neural Eng.* **17**, 26044 (2020).
86.   Voigts, J., Siegle, J., Pritchett, D. L. & Moore, C. I. The flexDrive: An ultra-light implant for optical
control and highly parallel chronic recording of neuronal ensembles in freely moving mice. *Front.*
*Syst. Neurosci.* **7**, 8 (2013).
87.   Pachitariu, M., Steinmetz, N., Kadir, S., Carandini, M. & Harris, K. *Fast and accurate spike sorting of*
*high-channel count probes with KiloSort*. *Advances in Neural Information Processing Systems* vol. 29
(2016).
88.   Eilers, P. H. C. & Marx, B. D. Flexible smoothing with B-splines and penalties. *Stat. Sci.* **11**, 89–102
(1996).
89.   Stanislaw, H. & Todorov, N. Calculation of signal detection theory measures. *Behav. Res. Methods,*
*Instruments, Comput.* **31**, 137–149 (1999).
90.   Rocchi, F. & Ramachandran, R. Neuronal adaptation to sound statistics in the inferior colliculus of
behaving macaques does not reduce the effectiveness of the masking noise. *J. Neurophysiol.* **120**,
2819–2833 (2018).
91.   Hautus, M. J. Corrections for extreme proportions and their biasing effects on estimated values of d′.
*Behav. Res. Methods, Instruments, Comput.* **27**, 46–51 (1995).
92.   Sahani, M. & Linden, J. F. How linear are auditory cortical responses? in *Advances in Neural*
*Information Processing Systems* 109–116 (2003). doi:10.1124/dmd.105.005157.concerning.
93.   Sahani, M. & Linden, J. F. *Evidence optimization techniques for estimating stimulus-response*
*functions*. *Advances in Neural Information Processing Systems* https://papers.nips.cc/paper/2294-
evidence-optimization-techniques-for-estimating-stimulus-response-functions.pdf (2003).
94.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).
217
218

**Supplemental Experimental Procedures**

*Acute electrophysiological recordings with muscimol or saline.*
Neuronal signals were recorded from n = 2 awake, untrained mice. Prior to the recording session, each mouse was anesthetized and a headpost and ground pin were implanted on the skull (see *Surgery* in the main text). On the day of the recording, the mouse was briefly anesthetized with 3% isoflurane and a small craniotomy was performed over auditory cortex using a dental drill or scalpel (~1mm x 1mm craniotomy centered approximately 1.25mm anterior to the lambdoid suture along caudal end of the squamosal suture). A 32-channel silicon probe (Neuronexus) was then positioned perpendicularly to the cortical surface and lowered at a rate of 1-2µm/s to a final depth of 800-1200µm. As the probe was lowered, trains of brief noise bursts were repeated, and if stimulus locked responses to the noise bursts were observed, the probe was determined to be in auditory cortex. The probe was then allowed to settle for up to 30 minutes before starting the recording.
For the muscimol and saline recordings (Extended Data Figure 5), a durotomy was performed over the injection site and baseline neuronal responses to the behavioral stimuli were recorded. Then, 2.5µL of .25mg/mL muscimol or 0.9% sterile saline solution was topically applied to the surface of auditory cortex and allowed 30 minutes to penetrate the tissue. The same stimuli were then recorded again after the elapsed time. In these recordings, the same targets and DRC background presented during behavior were presented. Neuronal signals from n = 2 mice (1 mouse for muscimol application, 1 mouse for saline application) were amplified and digitized using a Cheetah Digital LYNX system (Neuronalynx) at a rate of 32kHz.

**Supplemental Results**

*The range of presented target levels shapes psychometric performance curves.*
In the experiments presented here, we utilized several sets of target levels to assess psychometric performance (for a summary of the target levels used, see Extended Data Table 3). When computing psychometric curves across all of the target conditions for each mouse (n = 25; Extended Data Figure 4b), we found that detection thresholds were lower in low contrast (Mean dB SNR (*M*) = 8.56, standard deviation (*std*) = 1.81) compared to high contrast (*M* = 14.23, *std* = 2.57; paired t-test: *t(23)* = -8.19, *p* = 2.88e-8, Extended Data Figure 4c). From our normative model, we expected psychometric slopes to decrease in high contrast. Instead, when combining sessions with different target levels, we found a significant increase in slope during high contrast (*M* = 0.056, *std* = 0.018) when compared to low contrast (*M* = 0.045, *std* = 0.0052; paired t-test: *t(23)* = -3.42, *p* = 0.0024, Extended Data Figure 4d). We hypothesized that psychometric performance was sensitive to the range of targets presented. To test this, we split the data by the range of target levels used in each session, finding that targets drawn from a narrow range resulted in steeper psychometric slopes than targets drawn from a wide range, regardless of the background contrast (Extended Data Figure 4e-h). Therefore, we concluded that the paradoxical increase in psychometric slope was due to the narrow range of targets selected for many high contrast sessions. To control for these effects and thus isolate the effect of background contrast on psychometric slope, we considered only sessions with identical ranges of targets in low and high contrast and found that slopes did indeed decrease in high contrast (Figure 3).

*Muscimol application disrupts cortical encoding of targets.*
In n = 2 awake, naïve mice, we first recorded baseline responses to the stimuli used in the psychometric task, then topically applied muscimol or saline, waited 30 minutes, and recorded stimulus responses again. After muscimol application, there was a marked decrease in neuronal responses to targets compared to the baseline recordings (Extended Data Figure 5b, left). Notably, in our saline control, we observed little to no change in neuronal responses after saline application (Extended Data Figure 5b, right). We next compared how contrast, level and muscimol or saline application changed the responses during the pre- and post-application periods, finding that muscimol significantly reduced the firing rates between pre- and post-application periods, while saline significantly increased firing rates (Extended Data Figure 5c,d, Extended Data Table 1). We speculate that the small increase in firing rate between pre- and post-saline application was due to changes in recording quality or due to neuronal drift over the ~1 hour recording session, and note that the effect size of saline pre-post application is very small ($\eta^2$ = 0.0046) compared to the effect size of

muscimol ($\eta^2$ = 0.38). We then used a three-way ANOVA to compare the effects of muscimol, contrast, and
target level on target responses in the saline and muscimol recording sessions. We found a significant main
effect of muscimol ($F(1)$ = 322.65, $p$ = 4.88e-67) and level ($F(6)$ = 15.48, $p$ = 1.98e-17), but no main effect of
contrast ($F(1)$ = 0.39, $p$ = 0.53), indicating nearly complete suppression of responses to both targets and
background in high and low contrast (Extended Data Figure 5e,f). These results confirmed that muscimol
effectively disrupts the cortical coding of our behavioral stimuli.
281
*Muscimol application does not prevent licking.*
An additional alternative effect of muscimol is a general loss of the ability to lick. To assess this, we
monitored the lick probability of the mice throughout the trial duration, and found that muscimol specifically
reduced licking responses during the period where targets were presented (rank-sum test: $Z$ = -4.23, $p$ =
2.34e-5; Extended Data Figure 5g, right panel of Extended Data Figure 5h). Mice also tended to lick
immediately after the trial onset (Extended Data Figure 5i, green trace), but we found that the lick rates under
muscimol and saline conditions were identical during this period (rank-sum test: $Z$ = 0.23, $p$ = 0.81; Extended
Data Figure 4h, left panel). These results suggest that muscimol does not impair the mouse's ability to lick in
general, but results in a specific deficit in licking in response to targets.
291
*STRF are stable across contrasts.*
To interpret the gain changes observed during the behavioral task, we first needed to eliminate the
possibility that STRF structure changed across the behavioral epochs of the task (ie, between low and high
contrast periods). To assess STRF stability during these two trial periods, we computed STRFs independently
in low and high contrast using a GLM. Next, we selected STRFs with well-defined features by computing the
SNR of each STRF. SNR was computed by calculating the ratio between the standard deviation of the STRF
coefficients in the first 100ms to the SNR of the coefficients in the remaining STRF. This metric assumes that
a STRF that is purely noise would have the same variability in the early and late phase of the STRF, while a
STRF with stronger stimulus selectivity would have greater variability during the early phase, due to a strong
response to some stimulus feature. A sampling of high SNR STRFs are plotted in Extended Data Figure 7a.
Next we selected only neurons with SNRs greater than 1.75 in both contrasts (approximately 2
standard deviations greater than the mean SNR, n = 55 neurons, Extended Data Figure 7b). Within this subset
of well-tuned neurons, we then computed the best frequency (BF) and lag by first computing the STRF
response averaged over time and frequency, respectively, and multiplied by the standard deviation of
coefficients in each bin. The BF and lag were determined to be the peak of the average frequency and time
components, respectively. Additionally, we computed the range of the STRFs. As described previously[17,19],
we found no significant change in BF (sign-rank test: $Z$ = 1.34, $p$ = 0.18, Extended Data Figure 7c) or lag ($Z$
= -1.29, $p$ = 0.20, Extended Data Figure 7d), but did observe a significant increase in STRF range ($Z$ = -5.76,
$p$ = 8.19e-9, Extended Data Figure 7e). As noted in previous work, the change in STRF range is consistent
with a change in neuronal gain. Taken together, these findings demonstrate that contrast, and thus the
different behavioral epochs of the task, did not significantly influence STRF features.
313
314    **Generalized linear model of contrast gain control dynamics**
315
316        A primary goal of the current study was to estimate the influence of stimulus contrast on neuronal gain
317    dynamics, for instance, after a switch from one contrast to another. To approach this problem, we first define
318    a model neuron with dynamic gain control.
319
320    *Forward model*
321        To best approximate the stimuli used in our experiments, we define the stimulus environment of our
322    model as an $F$-dimensional signal that evolves in discrete time steps:
323
324 $$X_{t,f} \sim \mathcal{N}(\mu,\ \sigma_t),$$
325
326    where $X_{t,f}$ is a stimulus spectrogram that varies as a function of time $t$ and frequency $f$. Each time and
327    frequency bin of $X$ is sampled from a normal distribution defined by an average value $\mu$ and contrast $\sigma_t$ at
328    time $t$.
329        To approximate the behavior of real neurons, we define a model neuron that has a two-dimensional
330    linear filter (representing the STRF of the neuron):

$$\beta_{h,f} = \mathcal{N}(m, C; h, f),$$

334 where stimulus filter $\beta_{h,f}$ is defined as a two-dimensional gaussian distribution evaluated at lag $h$ and
335 frequency $f$. The filter location in frequency-history space is defined by its mean $m$ and covariance matrix $C$.
336 The stimulus drive of the neuron at each time step, $x_t$, is then computed as the convolution of the stimulus
337 matrix and the linear filter:

$$x_t = X_t \beta \tag{3}$$

340 where $X_t$ at each time $t$ is a row vector of length $F \cdot H$ (ie. the unrolled stimulus spectrogram lagged by H lags)
341 and β is the filter, unrolled as a column vector of the same length.
342 The model neuron has a firing rate that depends only on the stimulus drive $x_t$ and the contrast $\sigma_t$ at
343 time $t$. We then assume that the number of spikes $y_t$ emitted by the neuron at each time step follow a Poisson
344 distribution:

$$y_t \sim Poisson(\lambda_t)$$

348 where $\lambda_t$ is the firing rate at time $t$, given by

$$\lambda_t = exp[a + g(\sigma_t)b(x_t - c)] \tag{4}$$

351 where $g$ is a gain control function, and $a$, $b$, and $c$ are parameters of the model. The parameter $a$ represents
352 the baseline response of the neuron, $b$ is a scaling factor of the stimulus drive, and $c$ represents the operating
353 point of the gain. We remove the obvious degeneracy in the definition of $g$ and $b$ (only their product matters)
354 by requiring that $g$ be adimensional and such that

$$\frac{1}{2}[g(\sigma_H) + g(\sigma_L)] = 1 \tag{5}$$

where $\sigma_H$ and $\sigma_L$ are the high and low contrast values. This constraint forces the neutral value of the gain,
$g = 1$ to be the midpoint between gain in the high and low contrast conditions.

*Optimal gain control*
In the spirit of the efficient coding principle, we derived a form for $g(\sigma)$ that will guarantee that, under
certain conditions, the dynamic range of the neuron will be approximately conserved under changes in
contrast. To do this, we define the dynamic range as

$$R(\sigma) = \lambda(\mu + \sigma) - \lambda(\mu - \sigma) \tag{6}$$

366 which can be rewritten using equation 15 as

$$R(\sigma) = e^a \big[ exp\big(g(\sigma)b(\mu + \sigma - c)\big) - exp\big(g(\sigma)b(\mu - \sigma - c)\big)\big]. \tag{7}$$

368 If the argument of the exponentials is not too large, we can linearize this expression to obtain

$$R(\sigma) \simeq 2e^a b\sigma \cdot g(\sigma) \tag{8}$$

370 and that $R$ is approximately independent of $\sigma$ provided that $g(\sigma) \propto 1/\sigma$. So, for our model, we set

$$g(\sigma) = \frac{\overline{\sigma}}{\sigma_t} \tag{9}$$

373 where $\overline{\sigma}$ is the harmonic mean of $\sigma_H$ and $\sigma_L$:

$$\bar{\sigma} := \left[\frac{1}{2}\left(\frac{1}{\sigma_H} + \frac{1}{\sigma_L}\right)\right]^{-1} = 2\frac{\sigma_H \sigma_L}{\sigma_H + \sigma_L} \tag{10}$$

Finally, to validate that our fitting methods are sensitive to real world neurons, which do not necessarily adjust their gain to account for changes in contrast according to the model just described, we consider an interpolation scheme that smoothly transforms a model with positive gain control to a similar model without gain control, or with "anti" gain control. To do this, we redefine $g$ as follows:

$$g(\sigma) \to \xi g(\sigma) + (1 - \xi), \qquad -1 \leq 0 \leq 1 \tag{11}$$

so that by changing $\xi$ we can control whether gain control is optimal ($\xi = 1$), non-existant ($\xi = 0$), or "anti" ($\xi = -1$).

Putting everything together, the final expression for the firing rate of the forward model is

$$\lambda_t = exp\left[a + b\left(\xi\frac{\bar{\sigma}}{\sigma_t} + (1 - \xi)\right)(x_t - c)\right] \tag{10}$$

*Generalized linear model*

The forward model developed in the previous section provides a simple approximation of the relationship between the stimulus, stimulus contrast and neuronal responses. We also note that the form of the forward model lends itself to estimation using a Poisson GLM, provided that the predictors are chosen appropriately. As such, we define the inference model as a Poisson GLM with an intercept term and the following predictors:

$$(x_t - \mu), \qquad \frac{\bar{\sigma}}{\sigma_t}, \qquad (x_t - \mu)\frac{\bar{\sigma}}{\sigma_t}$$

In other words, the model is composed of a stimulus predictor $(x_t - \mu)$, a contrast predictor $(\bar{\sigma}/\sigma_t)$, and their interaction. Therefore, the GLM models the data at time $t$ as a Poisson distribution with the following mean:

$$\lambda_t = exp\left[\beta_0 + \beta_1(x_t - \mu) + \beta_2\frac{\bar{\sigma}}{\sigma_t}(x_t - \mu) + \beta_3\frac{\bar{\sigma}}{\sigma_t}\right] \tag{12}$$

where $\beta_0 \dots \beta_3$ are the parameters to be inferred, and, as defined previously, $x_t$ is the stimulus drive of the neuron determined by its STRF.

*Model fitting*

To fit the model, we took a two-step approach. First we found the best-fit filter (STRF) for the neuron. Then, we fit the full GLM to determine how the linear drive determined by the STRF is modulated by contrast. In the first step, the linear drive is obtained by fitting the model

$$\ln\lambda_t = \alpha + X_t\beta \tag{13}$$

where $X_t$ is a design matrix defined as a function of frequency bins $f$ and history lags $h$, and $\beta$ is the fitted STRF. Stimulus drive $x_t$ is then computed as in equation 14.

We then define the full model according to equation 23,

$$\ln\lambda_t = \beta_0 + \beta_1 x_t + \sum_{i=1}^{B}\beta_{2i}\, x_t \cdot (b_i * c)(t) + \sum_{i=1}^{B}\beta_{3i}(b_i * c)(t) \tag{14}$$

where $c(t) = \bar{\sigma}/\sigma_t$ and $\{b_i\}_{i=1}^{B}$ is a set of cubic B-spline temporal basis functions. By defining a matrix $C$ as follows

$$C_{ti} = (b_i * c)(t) \tag{15}$$

414
415    we can rewrite equation 25 in a more compact form:

$$\ln \lambda = \beta_0 + x\beta_1 + x \circ C\beta_2 + C\beta_3 \tag{16}$$

416
where $\circ$ denotes element-by-element "broadcasting" multiplication.
To fit asymmetric changes in firing rate after transitions to low or high contrast, we took the simple
approach of defining separate sets of contrast predictors for each transition type. This amounted to modifying
$C$ by masking transitions to high contrast or transitions to low contrast with zeros, such that the model fit a
window $H'$ of 40 time bins around each contrast transition. To do so, we created a new matrix $C'$ by duplicating
$C$ column-wise. Then, we define the first $B$ columns as predictors for the transition to low contrast by masking
a 1 second period around each transition to high contrast with zeros. This same procedure was repeated for
the remaining columns in $C'$, instead masking out the transition to low contrast. Substituting this into equation
27, we obtain

$$\ln \lambda = \beta_0 + x\beta_1 + x \circ C'^{\beta_2} + C'^{\beta_3} \tag{17}$$

426
427    For the sake of clarity, note that in the expression above, $\beta_0$ is a number, $x$ is a column vector of length $T$, $\beta_1$
428    is a number, $C'$ is a $T$-by-$2B$ matrix, and $\beta_2$ and $\beta_3$ are column vectors of length $2B$.
429
*Defining gain*
We have outlined a forward model for simulating neuronal activity according to efficient coding of
stimulus contrast, and described an inference model (a Poisson GLM) for estimating the influence of the
stimulus, stimulus contrast, and their interaction. In this section, we describe how to use the fitted parameters
to quantify the amount of gain control in the neuron.
Conceptually, an increase or decrease in the gain of a system is analogous to more or less sensitivity
to small changes in the stimulus, dependent what is modulating the gain (in our case, the recent history of the
contrast). Based on this intuition, we focus on how the response of the neuron (as modeled by a fitted GLM)
is expected to change between conditions where the gain is expected to contribute (i.e. in the presence of
gain control) and where it is not (ie. in the absence of gain control, where gain is "neutral").
To do this, we start by considering the gradient of the link function (the log rate) at time $t$ with respect
to $X_t$:

$$\begin{aligned}
\eta_t = \nabla_{x_t} \ln\lambda_t &= \nabla_{x_t}[\beta_0 + x_t\beta_1 + x_t C_t \beta_2 + C_t \beta_3] \\
&= \nabla_{x_t}[\beta_0 + (X_t\beta)\beta_1 + (X_t\beta)C_t\beta_2 + C_t\beta_3] \\
&= \beta(\beta_1 + C_t\beta_2)
\end{aligned} \tag{18}$$

442
We can immediately read equation 29 as "the STRF of the model is modulated by a factor of $\beta_1 + C_t\beta_2$ at time
$t$", and define the gain based on this intuition, but we will take a slightly longer and more formal route to get
to the same result.
The gradient $\eta$ is a vector with the same dimensionality of $\beta_1$ and $C_t\beta_2$, and it encapsulates all
information about the sensitivity of the link function to small changes in $X_t$ at a given time. Because $X_t$ is not
a scalar (it has $H \cdot F$ components), these changes can happen along many dimensions, and the sensitivity
can be different in different directions. We can define the gain based on the sensitivity to changes in a specific
direction $v$ (assuming for concreteness that $||v|| = 1$, although this is not necessary for the derivation below).
If $X_t = r \cdot v$, where $r$ is some scalar, then

$$\frac{d\ln\lambda_t}{dr} = \langle \eta_t, v \rangle \tag{19}$$

452
453    by definition of the gradient. We can then define the gain $w$ along direction $v$ as the ratio between the
454    sensitivity of the log rate to changes along $v$ and the sensitivity one would have if the contrast $C_t$ was at some
455    reference value $C^0$ where we define $w = 1$ by construction. If we do so, we obtain
456

$$\tag{20}$$

$$\left(\frac{d\ln\lambda_t}{}\right)\left(\frac{d\ln\lambda(C = C^0)}{}\right)^{-1}$$

$$= \frac{\langle \beta, v \rangle (\beta_1 + C_t \beta_2)}{\langle \beta, v \rangle (\beta_1 + C^0 \beta_2)}$$
$$= \frac{\beta_1 + C_t \beta_2}{\beta_1 + C^0 \beta_2}$$

Note that this definition does not depend on the initial choice of $v$, or even on the specifics of the choice of
basis functions used to define $C$. In conclusion, by reasoning about the sensitivity of the response of the fitted
GLM, we define a value $w_t$ which captures the relationship between the true gain $g$ and the stimulus contrast
$c_t$.

*Simulations*
To validate our inference model, we simulated neuronal activity according to the generative model
defined in the *Forward Model* section (Extended Data Figure 2a). We were interested in capturing several
dimensions upon which the generative model could vary, namely, the amount of gain control in the simulated
neurons ξ, and the dynamics of the gain function $g$.
To parametrically control the evolution of gain over time, we simulated different temporal trajectories of gain
control, by modifying $g(\sigma_t)$ as follows

$$g(\sigma_c, \tau_c)_t = g(\sigma_{c-1}) + (g(\sigma_c) - g(\sigma_{c-1})) \cdot exp(\tau_c, t) \tag{21}$$

where the gain $g$ after a switch to contrast $\sigma_c$ transitions from the gain in the previous contrast $g(\sigma_{c-1})$ to the
gain in the current contrast $g(\sigma_c)$ according to an exponential function with time constant $\tau_c$. Note that $\tau_c$
could vary between the two contrasts to simulate asymmetric dynamics.
For each neuron, we first generated a STRF and linear drive according to equation 14 (Extended Data
Figure 2b,d). For different sets of simulated neurons, we parametrically varied the amount of gain control ξ
between -1 and 1, and varied the gain time courses to simulate three types of gain adaptation dynamics: 1)
Slow transitions to low contrast with fast transitions to high contrast, 2) Fast, symmetric transitions to each
contrast, 3) Fast transitions to low contrast and slow transitions to high contrast (Extended Data Figure 2f).
We simulated 100 neurons for each combination of ξ and τ, with other simulation parameters held constant
(Extended Data Table 4). Extended Data Figure 2e plots the average firing rates and overlaid model fits for
three sets of simulations with optimal gain control (ξ = 1) while varying τ. Importantly, the model flexibly
captured the gain dynamics in the three simulated adaptation conditions, with the gain estimate $w_t$ following
the true gain trajectory (Extended Data Figure 2f). For additional values of ξ, the model accurately predicted
the firing rates (Extended Data Figure 2g) and gain trajectories (Extended Data Figure 2h). We observed that
some combinations of ξ and τ elicited large firing rate transients, particularly in the cases where simulated
gain slowly adapted after a switch to high contrast (bottom panels in Extended Data Figure 2e, f, g, h). This
behavior is expected, as gain remains relatively high for a longer period after the switch, causing large
fluctuations in firing rate as the stimulus drive during high contrast is increased. These large firing rate
transients seemed to reduce the accuracy of gain estimate $w$, but we observed that the predicted time courses
still captured the overall asymmetries present in the underlying model.
During our behavioral recordings, we used a limited number of background noise scenes (n = 5) to
reduce the overall size of the stimulus set. However, it became clear that our model required a larger sample
of stimulus space to accurately estimate gain. To demonstrate this, we plotted the simulation results when
neurons were exposed to 100 unique background scenes (Extended Data Figure 2i) compared to simulations
where neurons were only exposed to 5 unique background scenes, as in our behavioral recordings (Extended
Data Figure 2j). We observed that with 100 scenes, estimates of $w$ were very close to the true gain values,
but were consistently underestimated in the case of 5 background scenes, even in the case of perfect gain
control. Therefore, when analyzing our behavioral recordings, we used a standard linear-nonlinear model to
estimate neuronal gain (Figure 5), as we previously found that gain estimates from the GLM were highly
correlated with gain estimated from the LN model (Figure 2i).

**Extended Data Table 1:** Statistical Comparisons.

| Comparison | Figure | Center | Spread | N | Test | Statistic | Effect Size | p-value |
|---|---|---|---|---|---|---|---|---|
| Behavior percent correct, low contrast: time 1 vs. time 2 | 2g | T1: 0.68 T2: 0.70 (median) | T1: 0.10 T2: 0.15 (IQR) | 21 mice | Two-tailed Wilcoxon sign-rank test (FDR corrected[94] for multiple comparisons) | Z = -1.93 Rank: 60 | $Z/\sqrt{n}$ = -0.42 | 0.054 |
| Behavior percent correct, low contrast: time 1 vs. time 3 | | T1: 0.68 T3: 0.82 (median) | T1: 0.10 T3: 0.092 (IQR) | | | Z = -4.01 Rank: 0 | $Z/\sqrt{n}$ = -0.88 | 5.96e-5 |
| Behavior percent correct, low contrast: time 1 vs. time 4 | | T1: 0.68 T4: 0.87 (median) | T1: 0.10 T4: 0.190 (IQR) | | | Z = -4.01 Rank: 0 | $Z/\sqrt{n}$ = -0.88 | 5.96e-5 |
| Behavior percent correct, low contrast: time 1 vs. time 5 | | T1: 0.68 T5: 0.91 (median) | T1: 0.10 T5: 0.11 (IQR) | | | Z = -4.01 Rank: 0 | $Z/\sqrt{n}$ = -0.88 | 5.96e-5 |
| Behavior percent correct, high contrast: time 1 vs. time 2 | | T1: 0.82 T2: 0.77 (median) | T1: 0.083 T2: 0.19 (IQR) | | | Z = 2.84 Rank: 181 | $Z/\sqrt{n}$ = 0.62 | 0.0045 |
| Behavior percent correct, high contrast: time 1 vs. time 3 | | T1: 0.82 T3: 0.77 (median) | T1: 0.083 T3: 0.14 (IQR) | | | Z = 2.17 Rank: 163 | $Z/\sqrt{n}$ = 0.47 | 0.030 |
| Behavior percent correct, high contrast: time 1 vs. time 4 | | T1: 0.82 T4: 0.78 (median) | T1: 0.083 T4: 0.16 (IQR) | | | Z = 3.36 Rank: 195 | $Z/\sqrt{n}$ = 0.73 | 7.80e-4 |
| Behavior percent correct, high contrast: time 1 vs. time 5 | | T1: 0.82 T5: 0.79 (median) | T1: 0.083 T5: 0.12 (IQR) | | | Z = 1.94 Rank: 157 | $Z/\sqrt{n}$ = 0.42 | 0.052 |
| ANOVA for effects of pre-post muscimol application, contrast, and level on firing rate in ACtx | S5c | n/a | n/a | 42 neurons | three-way ANOVA | $F_{pre-post}(1)$ = 812.54 $F_{contrast}(1)$ = 22.64 $F_{level}(6)$ = 21.70 | $\eta^2$ = 0.38 $\eta^2$ = 0.011 $\eta^2$ = 0.061 | 4.48e-136 2.19e06 2.77e-24 |
| ANOVA for effects of pre-post saline application, contrast, and level on firing rate in ACtx | S5d | n/a | n/a | 104 neurons | three-way ANOVA | $F_{pre-post}(1)$ = 15.40 $F_{contrast}(1)$ = 0.43 $F_{level}(6)$ = 76.067 | $\eta^2$ = 0.0046 $\eta^2$ = 1.29e-4 $\eta^2$ = 0.14 | 8.89-5 0.51 1.76e-88 |
| Percent correct max dB SNR, low contrast: muscimol vs. saline | 3c | Musc.: 0.10 Saline: 0.85 (median) | Musc.: 0.67 Saline: 0.27 (IQR) | 10 musc. sessions, 10 saline sessions (4 mice) | Two-tailed Wilcoxon rank-sum test | Z = -2.76 Rank: 68 | $Z/\sqrt{n}$ = -0.62 | 0.0058 |
| Threshold (dB SNR), low contrast: muscimol vs. saline | | Musc.: 14.78 Saline: 9.66 (median) | Musc.: 18.46 Saline: 6.88 (IQR) | | | Z = 0.72 Rank: 115 | $Z/\sqrt{n}$ = 0.16 | 0.47 |
| FA rate, low contrast: muscimol vs. saline | | Musc.: 0.026 Saline: 0.132 (median) | Musc.: 0.10 Saline: 0.85 (IQR) | | | Z = -2.91 Rank: 66 | $Z/\sqrt{n}$ = -0.65 | 0.0036 |
| Max slope (PC/dB), low contrast: muscimol vs. saline | | Musc.: 0.026 Saline: 0.072 (median) | Musc.: 0.056 Saline: 0.030 (IQR) | | | Z: -2.68 Rank: 69 | $Z/\sqrt{n}$ = -0.60 | 0.0073 |
| Percent correct max dB SNR, high contrast: muscimol vs. saline | | Musc.: 0.06 Saline: 0.80 (median) | Musc.: 0.10 Saline: 0.85 (IQR) | 13 musc. sessions, 10 saline sessions (4 mice) | | Z = -4.06 Rank: 92 | $Z/\sqrt{n}$ = -0.83 | 4.96e-5 |
| Threshold (dB SNR), high contrast: muscimol vs. saline | | Musc.: 16.77 Saline: 18.80 (median) | Musc.: 21.33 Saline: 5.89 (IQR) | | | Z = -0.35 Rank: 156 | $Z/\sqrt{n}$ = -0.071 | 0.73 |
| FA rate, low contrast: muscimol vs. saline | | Musc.: 0.027 Saline: 0.213 (median) | Musc.: 0.10 Saline: 0.85 (IQR) | | | Z = -3.19 Rank: 107 | $Z/\sqrt{n}$ = -0.65 | 0.0014 |
| Max slope (PC/dB), high contrast: muscimol vs. saline | | Musc.: 0.012 Saline: 0.058 | Musc.: 0.024 Saline: 0.018 | | | Z = -3.77 Rank: 97 | $Z/\sqrt{n}$ = -0.77 | 1.66e-4 |

| | | (median) | (IQR) | | | | | |
|---|---|---|---|---|---|---|---|---|
| Percent correct max dB SNR, target in high contrast : muscimol vs. saline | 3f | Musc.: 0.07 Saline: 0.82 (median) | Musc.: 0.51 Saline: 0.095 (IQR) | 5 musc. sessions, 5 saline sessions (2 mice) | Two-tailed Wilcoxon rank-sum test | Z = nan Rank: 15 | $Z/\sqrt{n}$ = nan | 0.0079 |
| Percent correct at threshold, target in high contrast: muscimol vs. saline | | Musc.: 0.03 Saline: 0.53 (median) | Musc.: 0.35 Saline: 0.11 (IQR) | | | Z = nan Rank: 17 | $Z/\sqrt{n}$ = nan | 0.032 |
| FA rate, target in high contrast : muscimol vs. saline | | Musc.: 0.12 Saline: 0.23 (median) | Musc.: 0.22 Saline: 0.11 (IQR) | | | Z = nan Rank: 21 | $Z/\sqrt{n}$ = nan | 0.22 |
| Max slope (PC/dB), target in high contrast : muscimol vs. saline | | Musc.: 0.038 Saline: 0.057 (median) | Musc.: 0.046 Saline: 0.012 (IQR) | | | Z = nan Rank: 19 | $Z/\sqrt{n}$ = nan | 0.095 |
| Percent correct max dB SNR, target in silence: muscimol vs. saline | | Musc.: 0.85 Saline: 0.92 (median) | Musc.: 0.23 Saline: 0.15 (IQR) | 8 musc. sessions, 8 saline sessions (2 mice) | | Z = nan Rank: 53 | $Z/\sqrt{n}$ = nan | 0.13 |
| Percent correct at threshold, target in silence : muscimol vs. saline | | Musc.: 0.11 Saline: 0.22 (median) | Musc.: 0.28 Saline: 0.22 (IQR) | | | Z = nan Rank: 55 | $Z/\sqrt{n}$ = nan | 0.20 |
| FA rate, target in silence : muscimol vs. saline | | Musc.: 0.029 Saline: 0.041 (median) | Musc.: 0.038 Saline: 0.11 (IQR) | | | Z = nan Rank: 60 | $Z/\sqrt{n}$ = nan | 0.44 |
| Max slope (PC/dB), target in silence : muscimol vs. saline | | Musc.: 0.028 Saline: 0.031 (median) | Musc.: 0.015 Saline: 0.0048 (IQR) | | | Z = nan Rank: 63 | $Z/\sqrt{n}$ = nan | 0.65 |
| Correlation coefficient between behavioral and neuronal percent correct (split by target level): Low level | 5h | n/a | n/a | 215 sessions | Pearson correlation | Pearson r = 0.31 | n/a | 3.44e-5 |
| Medium level | | n/a | n/a | 194 sessions | | Pearson r = 0.28 | n/a | 6.18e-5 |
| High level | | n/a | n/a | 203 sessions | | Pearson r = 0.15 | n/a | 0.032 |
| Neuronal percent correct, low contrast: time 1 vs. time 2 | 5l | T1: 0.79 T2: 0.83 (median) | T1: 0.15 T2: 0.22 (IQR) | 43 sessions | Two-tailed Wilcoxon sign-rank test (FDR corrected[94] for multiple comparisons) | Z = -1.12 Rank: 418 | $Z/\sqrt{n}$ = -0.17 | 0.26 |
| Neuronal percent correct, low contrast: time 1 vs. time 3 | | T1: 0.79 T3: 0.85 (median) | T1: 0.15 T3: 0.15 (IQR) | | | Z = -3.61 Rank: 198 | $Z/\sqrt{n}$ = -0.56 | 0.00031 |
| Neuronal percent correct, low contrast: time 1 vs. time 4 | | T1: 0.79 T4: 0.92 (median) | T1: 0.15 T4: 0.20 (IQR) | | | Z = -4.68 Rank: 103 | $Z/\sqrt{n}$ = -0.72 | 2.89e-6 |
| Neuronal percent correct, low contrast: time 1 vs. time 5 | | T1: 0.79 T5: 0.91 (median) | T1: 0.15 T5: 0.16 (IQR) | | | Z = -5.34 Rank: 31 | $Z/\sqrt{n}$ = -0.82 | 9.44e-8 |
| Neuronal percent correct, high contrast: time 1 vs. time 2 | | T1: 0.78 T2: 0.74 (median) | T1: 0.15 T2: 0.12 (IQR) | | | Z = 2.62 Rank: 690 | $Z/\sqrt{n}$ = 0.40 | 0.0088 |
| Neuronal percent correct, high contrast: time 1 vs. time 3 | | T1: 0.78 T3: 0.76 (median) | T1: 0.15 T3: 0.13 (IQR) | | | Z = 1.45 Rank: 593 | $Z/\sqrt{n}$ = 0.22 | 0.15 |
| Neuronal percent correct, high contrast: time 1 vs. time 4 | | T1: 0.78 T4: 0.83 | T1: 0.15 T4: 0.20 | | | Z = -0.24 Rank: 453 | $Z/\sqrt{n}$ = -0.037 | 0.81 |

| | | (median) | (IQR) | | | | | |
|---|---|---|---|---|---|---|---|---|
| Neuronal percent correct, high contrast: time 1 vs. time 5 | | T1: 0.78 T5: 0.83 (median) | T1: 0.15 T5: 0.16 (IQR) | | | Z = -2.00 Rank: 307 | $Z/\sqrt{n}$ = -0.31 | 0.045 |
| Mixed-effects model: `threshold ~ gain_target + contrast + (contrast-1|mouse)` | 6g | **Model Coefficients** Estimate ± standard error [tstat(df), p-value] Intercept: 10.97±1.33 t(120) = 8.27, p = 2.059e-13 Target gain: -30.46±12.45 t(120) = -2.45, p = 0.016 Contrast: 3.27±1.55 t(120)= 2.10, p = 0.038 | | 168 sessions | Likelihood ratio test against model without gain: `threshold ~ contrast + (contrast-1|mouse)` | $\chi^2(1)$ = 5.82 | | 0.016 |
| | | | | | Likelihood ratio test against model without contrast: `threshold ~ gain_target + (contrast-1|mouse)` | $\chi^2(1)$ = 3.71 | | 0.054 |
| Mixed-effects model: `slope ~ gain_target + contrast + (contrast-1|mouse)` | 6h | Intercept: 0.039±0.0064 t(120) = 6.23, p = 7.14e-9 Target gain: 0.16±0.060 t(120) = 2.67, p = 0.0085 Contrast: 0.0094±0.062 t(120)= 1.52, p = 0.13 | | 168 sessions | Likelihood ratio test against model without gain: `slope ~ contrast + (contrast-1|mouse)` | $\chi^2(1)$ = 6.96 | | 0.0083 |
| | | | | | Likelihood ratio test against model without contrast: `slope ~ gain_target + (contrast-1|mouse)` | $\chi^2(1)$ = 2.28 | | 0.13 |
| Mixed-effects model: `thresh ~ gain_adapt + contrast + (contrast-1|mouse)` | S8c | Intercept: 5.33±1.64 t(120) = 3.26, p = 0.0015 Adaptation gain: 56.66±35.62 t(120) = 1.59, p = 0.11 Contrast: 2.77±1.92 t(120)= 1.44, p = 0.15 | | 168 sessions | Likelihood ratio test against model without gain: `thresh ~ contrast + (contrast-1|mouse)` | $\chi^2(1)$ = 2.51 | | 0.11 |
| | | | | | Likelihood ratio test against model without contrast: `thresh ~ gain_adapt + (contrast-1|mouse)` | $\chi^2(1)$ = 2.020 | | 0.16 |
| Mixed-effects model: `slope ~ gain_adapt + contrast + (contrast-1|mouse)` | S8d | Intercept: 0.062±0.0078 t(120) = 7.98, p = 9.63e-13 Adaptation gain: -0.14±0.17 t(120) = -0.80, p = 0.43 Contrast: 0.0049±0.0084 t(120)= 0.58, p = 0.57 | | 168 sessions | Likelihood ratio test against model without gain: `slope ~ contrast + (contrast-1|mouse)` | $\chi^2(1)$ = 0.64 | | 0.43 |
| | | | | | Likelihood ratio test against model without contrast: `slope ~ gain_adapt + (contrast-1|mouse)` | $\chi^2(1)$ = 0.33 | | 0.57 |

**Extended Data Table 2:** Mouse strains and genders.

| Experiment | Figures | Strain | N [female, male] |
|---|---|---|---|
| Acute ACtx recordings | Figure 2 | CDH23 | 1 [M] |
| Behavior (no microdrive) | Figure 3 | C57BL/6 x CamKII-cre | 1 [F], 4 [M] |
| | | C57BL/6 x PV-cre | 1 [F] |
| | | CDH23 x SOM-cre | 1 [F], 1 [M] |
| Behavior (microdrive) | Figure 3, Figure 5, Figure 6 | CDH23 | 4 [F], 4 [M] |
| | | C57BL/6 x PV-cre | 1 [F] |
| | | C57BL/6  x SOM-cre | 1 [F] |
| | | CDH23 x SOM-cre | 1 [F], 2 [M] |
| | | CDH23 x CamKII-cre | 1 [F] |
| Muscimol (behavior) | Figure 4 | CDH23 | 2 [F], 2 [M] |
| Muscimol (acute recording) | Supplemental Figure 4 | CDH23 x CamKII-cre | 1 [M] |
| | | CDH23 | 1 [M] |
| Acute ACtx recordings | Supplemental Figure 5 | CDH23 x SOM-cre | 3 [F], 2 [M] |
| | | CDH23 x PV-cre | 1 [F], 1 [M] |
| | | CDH23 x VGAT | 2 [F] |
| | | **Total:** | 19 [F], 19 [M] |

**Extended Data Table 3:** Target SNRs used during psychometric testing.

| Target Levels [range] | [n]: Mouse IDs | n Sessions (total) | n High-Low Contrast Sessions | n Low-High Contrast Sessions |
|---|---|---|---|---|
| 0, 5, 10, 15, 20, 25 dB SNR [25] | [12]: CA102, CA104, CA106, CA107, CA118, CA119, CA121, CA122, CA123, CA124, CA125, CA126 | 214 | 111 | 103 |
| -5, 0, 5, 10, 15, 20 dB SNR [25] | [8]: CA102, CA104, CA106, CA107, CA118, CA119, CA121, CA122 | 31 | 31 | 0 |
| 0, 4, 8, 12, 16, 20 dB SNR [20] | [1]: CA046 | 1 | 0 | 1 |
| 5, 8, 11, 14, 17, 20 dB SNR [15] | [4]: CA118, CA119, CA121, CA122 | 68 | 52 | 16 |
| 8, 10.4, 12.8, 15.2, 17.6, 20 dB SNR [12] | [15]: CA046, CA047, CA048, CA049, CA051, CA052, CA055, CA061, CA070, CA072, CA073, CA074, CA075, CA104, CA107 | 111 | 0 | 111 |
| -4, 0, 4, 8, 12, 16 dB SNR [20] | [11]: CA051, CA052, CA055, CA061, CA070, CA072, CA073, CA074, CA075, CA102, CA106 | 91 | 91 | 0 |
| -5, -1, 3, 7, 11, 15 dB SNR [20] | [5]: CA046, CA047, CA048, CA049, CA051 | 19 | 19 | 0 |
| -75, -60, -45, -30, -15, 0 dB attenuation rel. 25dB SNR | [2]: CA124, CA125 | 20 | n/a | n/a |

**Extended Data Table 4:** GLM Simulation Parameters

| Parameter | Value |
|---|---|
| μ | 30 |
| $\sigma_L, \sigma_H$ | $[1,3]$ |
| β centroid $m$ (frequency bin $f$, history bin $h$) | $[20,2]$ |
| β covariance matrix $C$ | $\begin{bmatrix} 0.8 & 0.1 \\ 0.1 & 0.5 \end{bmatrix}$ |
| β dimensions $(F \cdot H)$ | $[33, 12]$ |
| Baseline rate $a$ | 0.1 |
| Stimulus scaling $b$ | 1 |
| Gain operating point $c$ | μ |
| Gain control ξ | $[-1.0, -0.5, 0, 0.5, 1.0]$ |
| Adaptation time constants $[\tau_L \ \tau_H]$ | $\begin{bmatrix} Slow-Fast: 0.05 & 0.5 \\ Fast-Fast: 0.5 & 0.5 \\ Fast-Slow: 0.5 & 0.05 \end{bmatrix}$ |
| Simulated background scenes | 100 *or* 5 |
| Contrast history $H'$ | 40 |
| B-spline degree, knots | $[3, 7]$ |

**Extended Data Figure 1 (related to Figure 1). Normative model responses, predictions, and example response distributions.**

**a,** The firing rate of the simulated neuron as a function of time. Traces shaded in blue or red indicate the firing rate to periods of low or high contrast background noise, respectively. The green trace indicates the model response to overlaid targets. **b,** The true contrast (labelled as variance) of the stimulus (blue, red, and dashed gray lines) along with the average model estimate of the contrast (solid black line) over time. **c,** Discriminability as a function of time and contrast, with the trace color indicating the contrast after the switch. The dashed vertical line indicates the time of the contrast switch. Open circles indicate time samples used to plot the distributions in **d**. **d,** Target (green) and background (blue or red) distributions as a function of time and contrast. The top row includes responses to targets and background in low contrast. Each column denotes a different time step relative to the change in contrast, as indicated by the column title. The bottom row is the same, but for high contrast. Arrows between **c** and **d** indicate distributions which yielded the indicated value of discriminability in the trace.

**Extended Data Figure 2 (related to Figure 2). Simulation results to validate the GC-GLM.**

**a,** Schematic of simulated neurons in the forward model. Each neuron received broadband noise inputs which changed contrast every 2s ($X_{t,f}$). A STRF modelled by a 2D-gaussian function with added noise filtered the stimulus to generate a linear response. This filter response was then modulated by a gain control function, which controlled the amount and time-course of gain control based on the stimulus contrast. This gain modulated output was then exponentiated and stochastic spikes were generated using a Poisson process. **b,** Example STRF from one simulated neuron. Colorbar indicates STRF magnitude. **c,** Model estimate of the STRF averaged across 100 simulated neurons. **d,** Example linear drive for one simulated neuron over 500 trials (ie. the filter response of the STRF convolved with the stimulus). **e,** Each panel plots the average firing rates of 100 simulated neurons (solid teal lines) and corresponding GC-GLM fits (dashed black lines) when simulating perfect gain control (GC = 1.0). Each row corresponds to 100 simulations of different gain time courses, with the top row depicting a slow transition to low contrast, with a fast transition to high contrast. The middle row plots simulations were both transitions were fast. The bottom row plots simulations where the transition to low contrast was fast, with a slow transition to high contrast. The corresponding rows of panels **f**, **g**, and **h**, are the results of simulations with the same gain time courses. **f,** Average gain time-course of the simulated neurons (solid teal lines) and the corresponding GC-GLM estimate of the gain, $w$, averaged over 100 simulations (black dashed lines). Insets of each panel depict the contrast kernels (dashed lines) and gain kernels (solid lines) estimated for each contrast. Blue lines indicate kernels after a switch to low contrast and red lines indicate kernels after a switch to high contrast. **g,** Average log firing rate for simulations with different gain time-courses and different degrees of gain control (GC value; the legend in the lower right indicates the color-GC value mapping). Each plotted line indicates the average firing rate/prediction for 100 simulations. **h,** Average

gain time-course of all simulations (solid colored lines) and the average estimates of $w$ (dashed gray lines). **i,** Simulations with 100 unique stimulus scenes, repeated 5 times each. Left panel plots the average firing rates and model fits. Right panel plots the true gain time-course (solid lines) and the average model gain estimate, $w$ (dashed lines). The shaded areas indicate 2.5 and 97.5 percentiles of the gain estimates. **j,** Simulations with 5 unique stimulus scenes, repeated 100 times each. Formatting as in **i**. For panels **e-j**, the GC value colors and line formatting are indicated in the legend on the bottom right.

**Extended Data Figure 3 (related to Figure 2). GC-GLM comparison to a dynamic contrast adaptation model.**

**a,** Example model fits to the neuron plotted in Figure 2d. Grey volume indicates the smoothed spike count, while overlaid pink and orange lines indicate fits of a previously published contrast adaptation model[19] and the GC-GLM described in this study, respectively. Inset text indicates the correlation between the model prediction and the firing rate, and the adaptation time constant for the model from Lohse et al. **b,** Model performance for each neuron quantified by cross-validated correlation between the trial averaged firing rate and the trial-averaged model prediction. Colors as in a, with green dots representing the GC-LN model correlations, for comparison. A sign-rank test between the Lohse et al. model and the GC-GLM indicated that the GC-GLM was a better fit to the data (n = 97 neurons; *median ± IQR difference in r*: 0.065 ± 0.12, Z = 7.66, *p* = 1.93e-14).

**Extended Data Figure 4 (related to Figure 3). Behavioral slopes are affected by the target level range.**

**a,** The effect of contrast on the false alarm rates in psychometric sessions (n = 25 mice). Each dot and line represent a mouse, the blue and red bars indicate the mean false alarm rate for low and high contrast ±SEM. Results of a paired t-test ($t(23)$ = -6.16, $p$ = 2.75e-6) across contrast revealed a significantly higher false alarm rate in high contrast (Mean ($M$) = 0.22, standard deviation ($std$) = 0.080) compared to low contrast ($M$ = 0.13, $std$ = 0.054). **b,** Psychometric curves for all mice (n = 25). Thin traces are curves of individual mice, overlaid with psychometric fits to the entire dataset. **c,** Comparison of psychometric thresholds for the curves plotted in **b**, formatted as in **a**. **d,** Slope of the psychometric curves plotted in **b**, formatted as in **a**. **e,** Average psychometric curves and percent correct for mice presented with a narrow range of targets (range = 15 dB SNR; Extended Data Table 3, row 4; dashed lines and open dots), and those presented with a wide range of targets (range = 25 dB SNR; Extended Data Table 3, row 1; solid lines and filled dots) in low contrast. Error bars indicate ±SEM. **f,** Psychometric slope for each mouse when low contrast targets were from narrow or wide target distributions. Each bar indicates the mean for each condition ±SEM. An unpaired t-test ($t(9)$ = 4.48, $p$ = 0.0015) indicated significantly steeper slopes in response to narrow target distributions ($M$ = 0.051, $std$ = 0.0051) compared to wide target distributions ($M$ = 0.039, $std$ = 0.0042). **g,** Average psychometric curves and percent correct for mice presented with a narrow range of targets (average of ranges = 12 and 15 dB SNR; Extended Data Table 3, rows 4 and 5; dashed lines and open dots) or wide range of targets (range = 25 dB SNR; Extended Data Table 3, row 1; solid lines and filled dots) in high contrast. **h,** Psychometric slope for each mouse when high contrast targets came from narrow or wide distributions. Formatting as in **f**. An unpaired t-test ($t(28)$ = 7.00, $p$ = 1.30e-7) indicated significantly steeper slopes in response to narrow target distributions ($M$ = 0.068, $std$ = 0.015) compared to wide target distributions ($M$ = 0.036, $std$ = 0.0070) in high contrast. In all plots: $^{ns}p$>0.1; $^{†}p$<0.1, *$p$<0.05, **$p$<0.01, ***$p$<0.001, ****$p$<0.0001.

**Extended Data Figure 5 (related to Figure 4). Confirmation of cortical inactivation with muscimol.**

**a,** Setup schematic for acute muscimol recordings in ACtx. **b,** Example spike rasters from two different neurons pre- and post-muscimol or saline application. On top of the raster is the timeline for each recording. Rasters are sorted by contrast and target level, with color indicating low or high contrast backgrounds, color shade indicating target level, and gray indicating background-only trials (-Inf). *Left panel:* spike raster of a representative neuron recorded prior to muscimol application, followed by the raster for the same neuron 30 minutes after muscimol application. *Insets:* Mean firing rate for each condition. Shade indicates target level and the scale bar indicates the firing rate. Error bars are ±SEM across trials. *Right panel:* Example neuron before and after application of saline. Formatting as in left panels. **c,** Firing rates before and after muscimol application as a function of target level and contrast. Dark dashed lines indicate spike rates recorded pre-muscimol application and light dashed lines indicate the responses post-application. **d,** Firing rates before and after saline application. As in **c**, dark lines are responses recorded prior to saline application and light lines indicate responses recorded after saline application. In **c** and **d**, blue and red plots indicate responses during low contrast and high contrast, respectively, and the circles not connected by a line and labelled "-Inf" are responses to background alone. **e,** Area under the ROC curve (AUC) averaged across neurons after drug application in muscimol and saline recording sessions in low contrast. Filled circles and solid lines are responses after saline was applied while open circles and dashed lines are responses after muscimol was applied. Error bars indicate ±SEM across neurons. **f,** Same as **e**, but for high contrast. **g,** Lick probability over time during muscimol or saline sessions. Dashed vertical lines indicate trial onset (0 s) and the contrast switch (3 s). Green traces are muscimol sessions and black traces are saline sessions. The shading around each trace indicates ±SEM across sessions. **h,** *Left:* comparison of lick probability during the adaptation period. *Right:* comparison of lick probability during the target period. Each circle indicates a session and color is as in **g**. **i,** Cumulative probability of licking throughout the trial, normalized within muscimol or saline conditions to sum to 1. Colors as in **g**, **h**. Shading indicates ±SEM across sessions. In all plots: $^{ns}p>0.1$; $^{†}p<0.1$, $^*p<0.05$, $^{**}p<0.01$, $^{***}p<0.001$, $^{****}p<0.0001$.

**Extended Data Figure 6 (related to Figure 5). Histological tracing of tetrode tracks.**

**a,b** Tetrode placement targeting A1 in two mice. The location of primary auditory cortex is indicated with white lines, and labelled A1, assessed by alignment of the slice features to a mouse atlas. Inset: the slice position in the atlas and displacement from bregma.

**Extended Data Figure 7 (related to Figure 6). STRFs are unaffected by contrast during behavioral performance.**

**a,** Example STRFs estimated in low and high contrast during behavioral performance (rows are individual neurons, columns are STRFs estimated per contrast). Inset text indicates the SNR of the STRF (*Online Methods*), and the best frequency (BF). **b,** Histogram of SNRs across all neurons and contrasts. We restricted analyses to well-defined STRFs, including only cells with STRFs which were greater than 1.72 SNR (~2std of the SNR distribution) in both low and high contrast. The SNR cutoff is indicated by the red line. **c,** Best frequency of high SNR STRFs in low and high contrast. **d,** The temporal lag of the peak STRF response in low and high contrast. **e,** The range between the minimum and maximum STRF coefficients. In c-d, the significance of a sign-rank test across contrasts is indicated in the lower right corner. In all plots: $^{ns}p>0.1$; $^{\dagger}p<0.1$, $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$.

**Extended Data Figure 8 (related to Figure 6). STRFs are unaffected by contrast, and the relationship between gain during adaptation periods in the trial and behavior.**

**a,** Correlation coefficients between the prediction of a linear-nonlinear model using STRFs estimated from the model without gain control (static-LN) versus a model with gain control (GC-LN). Each dot indicates a neuron. The red solid line indicates unity. The red "x" indicates the median correlation in each contrast. Asterisks indicate the significance of a sign-rank test. **b,** Psychometric performance in low contrast, averaged based on a median split of average cortical gain during the *adaptation* period of the trial. Light dots and lines indicate the session average and psychometric fit to sessions in the bottom 50th percentile of gain, while dark dots and lines indicate the same values for sessions in the top 50th percentile of gain. Errorbars indicate ±SEM across sessions. *Inset*: distribution of average gain in each session estimated from the adaptation period. The red dashed line indicates the median of the distribution, and the histogram bars are shaded according to whether they fall above (dark blue) or below (light blue) the median. **c,** Session-wise relationship between average gain in the adaptation period and psychometric threshold. Each dot indicates the gain and threshold for a single session, and its color indicates the contrast of the adaptation period. The gray line is the best linear fit to the data. The text in the lower right indicates the results of Likelihood Ratio Tests for models including gain as a predictor (in gray) or contrast as a predictor (in red). Full statistical results in Extended Data Table 1. Grey and black "ns" indicate that gain in the adaptation period and contrast, respectively, did not significantly predict psychometric slopes. **d,** Same as in **j**, but plotting psychometric slope as a function of gain. In all plots: $^{ns}p>0.1$; $^{†}p<0.1$, $^{*}p<0.05$, $^{**}p<0.01$, $^{***}p<0.001$, $^{****}p<0.0001$.