1 Title: **Cryptographic-like mechanism allows scatter-hoarders to**
2 **cache and retrieve their food secretly**
3

4 **Authors:**
5 Oren Forkosh[1,2]
6

7 **Affiliations:**
8 [1] Department of Cognitive and Brain Sciences, The Hebrew University of Jerusalem,
9 Jerusalem, Israel
10 [2] Department of Animal Sciences, The Hebrew University of Jerusalem, Rehovot, Israel
11

Abstract
For animals, the ability to hide and retrieve valuable information, such as the location of
food, can mean the difference between life and death. Here, we propose that to achieve
this, their brain uses spatial cells similarly to how we utilize encryption for data security.
Some animals are able to cache hundreds of thousands of food items annually by each
individual and later retrieve most of what they themselves stashed. Rather than memorizing
their cache locations as previously suggested, we propose that they use a single
cryptographic-like mechanism during both caching and retrieval. The model we developed
is based on hippocampal spatial cells, which respond to an animal's positional attention,
such as when the animal enters a specific region (place-cells) or gazes at a particular
location (spatial-view-cells). We know that the region that activates each spatial cell remains
consistent across subsequent visits to the same area but not between areas. This
remapping, combined with the uniqueness of cognitive maps, produces a persistent crypto-
hash function for both food caching and retrieval. We also show that the model stores
temporal information that helps animals in food caching order preference as previously
observed. This behavior, which we refer to as crypto-taxis, might also explain consistent
differences in decision-making when animals are faced with a large number of alternatives
such as in foraging.
30
31

Main Text

Animals have much to hide. Some species evade potential predators or prey by finding cover or by using camouflage (*1*), mimicry, and other means of disguise (*2*). Others conceal their eggs or offspring, mask an illness or an injury to avoid being targeted by predators (*3*), or stash valuable resources, such as food.

Scatter hoarding is probably the largest-scale manifestation of secretive behavior in the animal kingdom. Many species of animals engage in this behavior, which involves storing food at multiple cache sites to preserve it for times when food is scarce (*4*). Several bird species, such as the Siberian tit (Poecile cinctus), were observed to cache over 500,000 items per individual in one year (*4*). While much of the research on scatter hoarding was conducted on birds, this behavior is not specific to them (*4*): squirrels (*5*), chipmunks (*6*), and even foxes (*7*, *8*) stash food for times of need. As caching sites cannot be defended, the success of this strategy is often contingent on an animal's ability to keep the stashes away from prying eyes and hard to find (*5*).

Once the valuables are stashed, scatter hoarders are faced with the considerable challenge of retrieving the hidden items. It was shown that hoarding birds do not randomly seek out cached food (*9*); rather, they mainly retrieve that which they had themselves hidden – a remarkable feat, especially given the large number of cache sites (*10*). In the 1950s, Olof Swanberg observed that almost 90% of Eurasian nutcracker (*Nucifraga caryocatactes*) excavation holes contained cracked nutshells, indicating successful retrievals (*4*). In one elegant experiment, Stevens and Krab attached tiny magnets to the legs of marsh tits (*Poecile palustris*) (*12*). The birds were then offered serially numbered peanuts labeled with a radioactive isotope, enabling the researchers to locate each bird's caching sites using a scintillation counter (*13*). Magnet detectors were placed at each location to detect the presence of marked birds. During the experiment, the birds retrieved about 25% of their own stored food, while none of the control caches (set up by the researchers) were activated. The actual retrieval rate was probably higher, but the experiment faced some technical hurdles (*4*).

The ability to retrieve items from cache sites depends on spatial information such as visual cues. In (*14*), black-capped chickadees (*Poecile atricapillus*) were placed in an enclosure and their food-caching behavior was tracked. Object rearrangement around the enclosure greatly impaired the chickadees' ability to find their cache sites; manipulation of prominent global landmarks (large cardboard cutouts and a poster) had a much stronger effect on the

birds' retrieval performance than small proximal objects (5-cm squares). Shifting objects by
as little as 20 cm to the right significantly decreased the chickadees' ability to recover the
food. Moreover, in almost 70% of the cases, the birds searched within 5 cm of the location
implied by the more prominent landmarks, with a mean displacement of around 20 cm.
Assuming this is approximately the caching resolution, the finding indicates that a small
area of 10x10 meters can hold as much as 2,500 potential caching sites.
75
The hippocampus plays a central part in the remarkable cognitive feat of caching (*9*). This
is not surprising, as the hippocampus is known to be involved in processing spatial
information in the brain (*11*). A large subpopulation of neurons within the hippocampus in
animals such as mice, rats, and bats exhibit *place-cell* behavior; that is, they increase in their
spike rate in response to the animal's entering a specific region within a given site (usually,
but not always, one region per cell). The region activating each place cell often changes
when the animal moves to a new area, often in an unpredictable manner. However, if the
animal returns to a site previously visited, the place cell's receptive fields also return to their
previous arrangement, and this change happens practically instantaneously. This
remapping of the receptive field within a given environment is mostly insensitive to
landmark manipulations. In primates, we usually find a related type of cells referred to as
spatial view cells. These cells respond remotely when an animal is gazing at a specific region,
independently of the animal's location or head direction (*12*).
89
A well-known homolog to the mammalian hippocampus also exists in birds, with similar
involvement in spatial and episodic memory (*13*). Hippocampus size in birds was found to
correlate with birds' ability to stash food. Although the interpretation of this correlation is
under debate (*14*, *15*), animals that used more cache sites generally had a larger
hippocampus than non-caching bird species (*16*, *17*). In addition, even within the same
species, the size of the hippocampus was found to be larger in individuals dwelling in
harsher environments that makes them more dependent of the cached food (*18*). Moreover,
hippocampal neurogenesis has a seasonal element and seems to correlate with caching
activity throughout the year (*19*). For many years, the spatially responsive cells found in
avian brains were less related to a fixed position in space and more related to the challenge
the animal faced such as the position of a goal within a maze (*20*). Only very recently the
existence of place cells was demonstrated in the tufted titmouse (*Baeolophus bicolor*) (*21*).
102
Taken together with the fact that the hippocampus is involved in memory, these
observations have led researchers to hypothesize that caching requires some form of
spatial and episodic memory (*22*).

Yet as birds and other animals need an internal mechanism to guide them to stash food in
specific locations, the same mechanism can also be used to direct them to the exact same
locations while retrieving the food in that area. Such a mechanism may serve as a mnemonic
device (as suggested in (*23*)) or possibly replace the need for memory altogether. Such a
pseudo-random approach is much simpler than remembering hundreds of thousands of
stashing sites while still supporting all the existing empirical evidence. The guidance is
based on prominent landmarks in the terrain, such as trees and rocks, which are not likely
to substantially change over time, and can be used in the subsequent cache retrieval. And
we already know of a specific set of neurons to be able to do precisely this – the previously
mentioned hippocampal spatial cells. Spatial cells are unique to each individual, they assign
scores and rankings (using spike rates) to different locations within each area, they persist
over time, and remap when in the same area. We show how these properties allow animals
to find their cache sites efficiently and secretly.

Cryptography in the brain
A mechanism, or mapping, that can facilitate efficient hiding and retrieval of multiple cache
sites without relying on memory would need to have several basic properties. From a
theoretical perspective, the class of methods that achieve this is known as cryptographic-
hashing functions (or crypto-hashes, for short), which, as the name suggests, are comprised
of two components: hashing functions and cryptographic keys (*24*).

In this context, hashing refers to a class of functions that map arbitrarily complex data
(images, texts, audio files, and others) to a fixed size lower-dimensional representation.
Computer applications often use crypto-hashes to store objects into memory efficiently by
mapping them directly to a memory location (a type of a mnemonic device). In the case of
food caching, hashing can be used to map a set of landmarks within and around an area
onto a selected caching site within that area (Figure 1A). Efficient hash functions are such
that the probability of assigning different cues the same output is kept to a minimum; This
property reduces the possibility of collisions and redundancies that can occur when two
different inputs result in the same output. It also makes better use of all the available
resources – in the case of animals allowing the use of the entire area for caching.

Unlike standard hash functions, crypto-hashes incorporate an additional entity – a private
key that renders the mapping unique to the key owner (*24*). Assuming no two individuals
have the same key, it also means that the mapping will result in unique caching sites within
the same area (Figure 1B). Another valuable property of crypto-hashes is that it is often

difficult to infer the key from a small number of examples, so even if another animal finds
several caches, it will not be able to deduce the location of all others.

A straightforward and biologically plausible realization of crypto-hashing is through a
simple two-layer neural network model with sparse connectivity (Figure 1C; see
supplementary information). The first layer, the input layer, represents the visual cues
(landmarks) within a given patch of land (Figure 2A, 2B). The second layer is a 2D lattice of
spatial neurons, in which each neuron points to a specific location in a given area (Figure
2C). The firing rate of each spatial neuron corresponds to the likelihood of choosing its
particular location as a caching site. The spatial neurons are sparsely innervated by the
input neurons.

In our model, we set the number of connections to a constant, typically equivalent to the
number of landmarks the model uses (usually four). This sparse connectivity helps maintain
a low number of potential cache sites by the output layer. A simple equation can summarize
the activity of each spatial neuron

159
$$C_{x_i,y_i} = \left| \sum_{x,y} w^i_{x,y} I_{x,y} \right|$$

where C_{x_i,y_i} is the score, or spike-rate, of the spatial neuron pointing to coordinate x_i, y_i of
the output grid; $I_{x,y}$ is the input from coordinates x, y; and $w^i_{x,y}$ represents the strength of
the connectivity between $I_{x,y}$ and C_{x_i,y_i}. The strength, or weight $w^i_{x,y}$, of the connections was
assigned randomly at between one and minus-one. The inputs were also set in the range
one and minus-one, where the absolute magnitude represents the landmark's prominence
(one is very prominent and zero is designated as not noticeable). The sign represents the
landmark type, for example positive values represent "trees", and negatives represent
"rocks". The scores of the output neurons C_{x_i,y_i} effectively determine the probability of their
target area to be used as a cache site. We use the absolute value to keep the outputs
positive, although it has no computational benefit to the model. Nor is the choice to
distinguish between two types of objects by allowing negative inputs.

This neural network is a crypto-hash function, as it fulfills the three essential properties: (1)
It maps a complex terrain into a point with minimal overlapping probability across the
terrain or (2) across subjects (Figure 3A, 3B), and (3) reconstructing or decrypting the
mapping from examples is difficult. The third point stems from the fact that the connection
to each output-layer neuron is chosen randomly and independently of the others. Thus it is
effectively equivalent to holding a unique key for each neuron.

Figure 1. **Cryptographic mechanisms in the brain**. (A) Hash functions take high-dimensional objects (a map of a complex terrain, for example) and map them into a low-dimensional representation such as a point within that terrain. (B) Crypto-hash functions also include a private key that makes mapping the same object unique across individuals with different keys. (C) *Crypto-hashing in a two-layer neural network*. Each neuron $I_{x,y}$ in the input layer represents landmarks within a small square area. We assume two types of landmarks, which we refer to as 'trees' and 'rocks'. Neurons that point to trees are assigned a positive value, rocks get negative values, and if no object is within the neuron's receptive field, it is set to zero. The absolute value of $I_{x,y}$ corresponds to an object's prominence; thus, prominent trees will get +1, smaller trees – 0.5, and small rocks may be assigned a value of - 0.3. The spatial output layer is a 2D mesh that assigns a caching score to each location within the site. The higher the score, the more likely this location would be used for caching. Each neuron in the output layer is innervated by a small number of input neurons. In all of our simulations, we matched this number to the number of landmarks the model uses.

Figure 2. **Using spatial features to map cache sites without requiring brain plasticity**. (A-C) Four examples of the outcome of a crypto-hash neuronal network (in rows). (A) We simulated a terrain with four prominent landmarks by randomly choosing four cells within a 10x10 grid. The cells were assigned random values between -1 and +1, so that the absolute value represents the prominence of the spatial feature (cells with values close to +1 and -1 being the most prominent), and the sign represents the type of object. We refer to positive-valued cells as "trees" and negative cells as "rocks" for brevity. (B) The representation of the landscapes from (A) in the neural network's input layer. The size of the colored inlaid boxes represents the object's prominence and their color its sign (green for positive values or "trees", and gray for negatives or "rocks"). (C) The output of the target layer of two randomly chosen neural networks (Animal 1 and 2) in response to the inputs in (B). The output layer creates a unique probabilistic map of possible cache sites.

The result can serve as a traditional crypto-hash function by choosing the target neuron

with the highest score as the mapping outcome (see Materials an Methods). However, a

probabilistic map with multiple outcomes of varying probabilities – apart from being more

biologically feasible – also provides several benefits. The most straightforward benefit is

that it allows for an arbitrary number of caching sites within each area by choosing the

spatial neurons with the top scores.

Another valuable property of probabilistic maps is that they allow the addition of temporal

considerations into caching behavior (*22*). Assuming the order of food recovery starts with

locations that have higher scores, items with higher nutritional values or perishable items
(such as dead insects, as opposed to seeds) could be stashed in places with higher scores –
making them more likely to be recovered prior to items assigned to lower-scored locations
(Figure 3C). In addition, avoiding previously excavated sites requires only memorizing the
score of the last excavated and choosing only sites with a lower score (we refer to this
behavior as bookmarking).
195
Finally, we did not address how mappings in the model are kept allocentric and invariant to
the animal's position. A straightforward approach to achieve such invariance was
suggested in an elegant paper about geometric hashing (*25*). The method there is based on
choosing two prominent objects in the area and using them to scale and align all landmarks.
The vector connecting the two most prominent landmarks defines the direction axis and
the distance between them sets the scale unit. Using this approach, we can obtain a model
that is insensitive to affine transformations.
203
204
205

Figure 3. **Properties of a crypto-hash neural network.** The probability of overlapping sites between two random networks as a function of (A) potential cache sites and (B) the number of landmarks the network uses. The different colors correspond to the number of caches used in each area. The dotted lines are the overlap as expected from a random choice of sites. (C) *The probabilistic nature of the target layer scores allows the network to maintain temporal dynamics.* We assume that caching and

retrieval order are determined by the target-layer score, from highest to lowest. If the choice is not absolute but probabilistic, we still get the same temporal dynamic in both phases. The shaded area around each line represents the standard deviation.

METHODS
Landmarks and terrain
While the algorithm is not sensitive to the number of landmarks, for the sake of simplicity,
we assumed a fixed number of landmarks within each area - four, in this paper. These
landmarks were divided into two categories, which we refer to as "trees" and "rocks". The
locations of the landmarks were chosen randomly and uniformly from an n-by-n square grid
(in most cases we use n=10). Each landmark was then assigned a random value between
minus one and one that signifies the objects category and prominence: Cells that have
positive values are referred to as trees (values between zero and one), while rocks had
negative numbers (between zero and minus one). The absolute value of each landmark
signifies its prominence - so prominent trees have values closer to one and prominent rocks
have values closer to minus one. Zero marks a no-object. The outcome is a sparse n-by-n
matrix $S_{n \times n}$ with values that vary between minus one and one.
220
221 **Crypto-Hash Functions**
222 Hash functions map data with arbitrary dimensions to a fixed-length value (*24*). In
223 mathematical terms, a hash function $g_p(\bar{s})$ is such that
224
225 $$g_p(\bar{s}) = \bar{y}_p$$
226
227 where $\bar{s} \in S$ is a vector of arbitrary length, and $\bar{y}_p = (y_1, y_2, \ldots, y_p) \in \mathcal{Y}_p$ is a vector of a fixed-
228 length p. Since the length of $\bar{s}$ is often larger than that of $\bar{y}_p$ hash-functions can be viewed
229 as a special case of dimensionality-reduction.
230
An optimal hash function is such that the probability of mapping two inputs onto the same
output is minimal, or, equivalently, that all outputs values should have (roughly) the same
probability. This principal of uniformity can be formulated as
$$p(g_p(\bar{s}) = y) \approx \frac{1}{|\mathcal{Y}_p|}$$
where $|\mathcal{Y}_p|$ is the cardinality (number of elements or size) of the set $\mathcal{Y}_p$ of all possible output
values. Because of their uniformity, hash functions are often used in data storage and
retrieval tasks as they allow data access at nearly a constant time while requiring a storage
size that is only slightly larger than the space needed to store the data itself.
239
240 Crypto-hash functions introduce an additional term, a private-key k, to the basic hash
241 function
242

243
$$g_p(\bar{s}; k) = \bar{y}_p.$$

The key ensures that the mapping is unique, i.e. the probability that the same inputs
produce the same outputs for different keys is close to chance. Crypto-hash functions, like
hash functions in general, are deterministic, meaning that the same combination of input
and key will always produce the same output value. However, crypto-hash mappings are
also one-way-functions, meaning that they are difficult to invert; knowing an output value
gives very little information about the input or key.

Crypto-Hash Neural Network
Choice and retrieval of cache sites is based on prominent landmarks within a terrain.
Assuming $S_{n \times n}$ is the representation of the current area's terrain (see the 'simulated terrain'
section), our crypto-hash function can be defined as

(1)
$$g_p(S_{n \times n}; k) = (x, y)$$

where (x, y) is the cache coordinate within the area so that $x, y \in \{1, \dots, n\}$, and k is the crypto-
key.

A straightforward and biologically plausible to achieve this is using a neural network. We
define a two-layer network where, for simplicity, the neurons on both layers are organized
as a grid with x, y indices. The value of each neuron in the first layer $I_{x,y}$, which is the input
layer, is set according to the corresponding area tile or the (x, y)'th cell in $S_{n \times n}$. Each output
layer neuron C_{x_i, y_i} was sparsely connected to the input layer, and the weights $w_{x,y}^i$ were
randomly distributed between minus-one and one. The value of output neurons is the
absolute value of the weighted sum of their inputs

(2)
$$C_{x_i, y_i} = \left| \sum_{x,y} w_{x,y}^i I_{x,y} \right|.$$

269 In order to get a crypto-hash function in the form (1) we can take the index of the maximal
270 value or

271
$$g_p(S_{n \times n}; k) = \operatorname*{argmax}_{x', y'} C_{x', y'}.$$

272 In our case, a unique key k is obtained from the random choice of weights between the
273 neural network's layers.

275 However, as we mentioned in the text, keeping the function as a probabilistic mapping like
276 in (2) has several benefits and this is the form we use in the paper.
277
278 **DISCUSSION**
279
Although spatial cells, such as place cells have been extensively studied, the mechanism we
presented is one of the few explanations to how the brain might utilize these cells. We
proposed here that spatial cells can serve as the brain's crypto-hash functions, enabling
animals to hide food in unique cache sites and later to retrieve it. So far, there has been no
theory that explains what principles guide animals when choosing cache sites; If it was
merely a question of optimality, all animals might end up choosing the same sites, which
would lead to theft (or kleptoparasitism).
287
For brevity, we did not address how mappings in the model are kept allocentric and
invariant to the animal's position. A straightforward approach to achieve such invariance
was suggested in an elegant paper about geometric hashing (*30*). The method there is
based on choosing the two most prominent objects in the area and using them to scale and
align all landmarks. The vector connecting the two most prominent landmarks defines the
direction axis and the distance between them sets the scale unit. Using this approach, we
obtain a model that is insensitive to affine transformations (see supplementary
information).
296
While our focus here is on scattered hoarding, a similar mechanism may also be involved in
processes of decision making when the number of possibilities is large. A bee foraging for
nectar and pollen, for example, would benefit from choosing flowers less likely to have been
visited by other bees. The same function-driven mechanism (rather than memory-driven)
may also help steer migratory animals back to previously used nesting areas. And since the
hippocampus is involved in abstract knowledge in addition to spatial information (*11*), the
scope of decision-making might be much broader; Since diversity is a key characteristic of
all living system, it is tempting to think that humans' individualistic tendencies might also
be somehow related to our proposed brain circuitry.
306
While the work we presented is theoretical, it raises some obvious predictions. First, that
knowing the spiking patterns of spatial cells will enable us to determine cache site locations.
Moreover, if we know the remapping between sites well, we can use it to decrypt the internal
circuitry and predict cache sites in a new site that the animal is yet to visit. Finally, we predict

that the location of cache sites within a given area would be consistent across multiple hide
and retrieval iterations.
313
314 The instinct to choose cache sites that are both unique and obscure has a clear evolutionary
315 advantage. We therefore fondly suggest addressing this movement pattern as *cryptotaxis*
316 and the neurons involved as *crypto-cells*.
317
318

Acknowledgments
We would like to send our love and appreciation to Prof. David Sherry from Western
University for inspiring a large part of the idea for this paper with his beautiful works. We
thank Shiri Gerson for her help, wonderful comments, and lots of support. We also thank
Prof. Haim J. Wolfson for his help on the theory of hashing. Special thanks also to Prof. Anat
Barnea for her invaluable introduction to the world of scatter hoarding and the avian
hippocampus. We also thank Prof. Arnon Lotem for his insightful ideas, enthusiasm, and
encouragement.
327
328 ***Funding***
329 This work was supported by the Israeli Science Foundation (ISF; Grant 2505/20) and the Ring
330 Center for Interdisciplinary Environmental Research.

331
332 ***Competing interests***
333 Authors have no competing interests
334 ***Data and materials availability***
335 All code is available online

Bibliography

1.	M. Stevens, S. Merilaita, Animal camouflage: current issues and new perspectives. *Philosophical Transactions of the Royal Society B: Biological Sciences*. **364**, 423–427 (2009).

2.	G. D. Ruxton, W. L. Allen, T. N. Sherratt, M. P. Speed, *Avoiding attack: the evolutionary ecology of crypsis, aposematism, and mimicry* (Oxford University Press, 2019).

3.	R. K. Butler, D. P. Finn, Stress-induced analgesia. *Progress in Neurobiology*. **88**, 184–202 (2009).

4.	C. C. Smith, O. J. Reichman, The Evolution of Food Caching by Birds and Mammals. *Annual Review of Ecology and Systematics*. **15**, 329–351 (1984).

5.	L. F. Jacobs, E. R. Liman, Grey squirrels remember the locations of buried nuts. *Animal Behaviour*. **41**, 103–110 (1991).

6.	R. H. Yahner, The Adaptive Nature of the Social System and Behavior in the Eastern Chipmunk, Tamias striatus. *Behavioral Ecology and Sociobiology*. **3**, 397–427 (1978).

7.	N. Tinbergen, Von den Vorratskammern des Rotfuchses: (Vulpes vulpes L.)1. *Zeitschrift für Tierpsychologie*. **22**, 119–149 (1965).

8.	D. W. MacDonald, Food caching by red foxes and some other carnivores. *Z Tierpsychol*. **42**, 170–185 (1976).

9.	A. Brodin, The history of scatter hoarding studies. *Philos Trans R Soc Lond B Biol Sci*. **365**, 869–881 (2010).

10.	T. A. Stevens, J. R. Krebs, Retrieval of stored seeds by Marsh Tits Parus palustris in the field. *Ibis*. **128**, 513–525 (1986).

11.	T. E. J. Behrens, T. H. Muller, J. C. R. Whittington, S. Mark, A. B. Baram, K. L. Stachenfeld, Z. Kurth-Nelson, What Is a Cognitive Map? Organizing Knowledge for Flexible Behavior. *Neuron*. **100**, 490–509 (2018).

12.	E. T. Rolls, Spatial view cells and the representation of place in the primate hippocampus. *Hippocampus*. **9**, 467–480 (1999).

13.	E. D. Jarvis, O. Güntürkün, L. Bruce, A. Csillag, H. Karten, W. Kuenzel, L. Medina, G. Paxinos, D. J. Perkel, T. Shimizu, G. Striedter, J. M. Wild, G. F. Ball, J. Dugas-Ford, S. E. Durand, G. E. Hough, S. Husband, L. Kubikova, D. W. Lee, C. V. Mello, A. Powers, C. Siang, T. V. Smulders, K. Wada, S. A. White, K. Yamamoto, J. Yu, A. Reiner, A. B. Butler, Avian brains and a new understanding of vertebrate brain evolution. *Nature Reviews Neuroscience*. **6**, 151–159 (2005).

14.	T. C. Roth, A. Brodin, T. V. Smulders, L. D. LaDage, V. V. Pravosudov, Is bigger always better? A critical appraisal of the use of volumetric analysis in the study of the hippocampus. *Philos Trans R Soc Lond B Biol Sci*. **365**, 915–931 (2010).

15.	D. F. Sherry, Neuroecology. *Annu Rev Psychol*. **57**, 167–197 (2006).

16.   J. R. Krebs, D. F. Sherry, S. D. Healy, V. H. Perry, A. L. Vaccarino, Hippocampal
specialization of food-storing birds. *Proc Natl Acad Sci U S A*. **86**, 1388–1392 (1989).

17.   D. F. Sherry, A. L. Vaccarino, K. Buckenham, R. S. Herz, The hippocampal complex of
food-storing birds. *Brain Behav Evol*. **34**, 308–317 (1989).

18.   T. C. Roth, V. V. Pravosudov, Hippocampal volumes and neuron numbers increase
along a gradient of environmental harshness: a large-scale comparison. *Proc Biol Sci*. **276**,
401–405 (2009).

19.   A. Barnea, F. Nottebohm, Seasonal recruitment of hippocampal neurons in adult
free-ranging black-capped chickadees. *Proc Natl Acad Sci U S A*. **91**, 11217–11221 (1994).

20.   D. F. Sherry, S. L. Grella, M. F. Guigueno, D. J. White, D. F. Marrone, Are There Place
Cells in the Avian Hippocampus? *BBE*. **90**, 73–80 (2017).

21.   H. L. Payne, G. F. Lynch, D. Aronov, Precise spatial representations in the
hippocampus of a food-caching bird. *bioRxiv* (2020).

22.   N. S. Clayton, A. Dickinson, Episodic-like memory during cache recovery by scrub jays.
Nature. **395**, 272–274 (1998).

23.   A. Barnea, F. Nottebohm, Patterns of food storing by black-capped chickadees
suggest a mnemonic hypothesis. *Animal Behaviour*. **49**, 1161–1176 (1995).

24.   C. Paar, J. Pelzl, Understanding Cryptography: A Textbook for Students and
Practitioners (Springer Science & Business Media, 2009).

25.   H. J. Wolfson, I. Rigoutsos, Geometric hashing: an overview. *IEEE Comput. Sci. Eng.* **4**,
10–21 (1997).

399
400