1
2
3   **Nuclear SUN1 Stabilizes Endothelial Cell Junctions via Microtubules to Regulate**
4   **Blood Vessel Formation**
5
6

7   Danielle B Buglak[a], Molly R Kulikauskas[a], Ziqing Liu[b], Ariel L Gold[b], Allison P Marvin[b],
8   Andrew Burciu[b], Natalie T Tanke[a], Shea N Ricketts[c], Karina Kinghorn[a], Morgan Oatley[b],
9   Bryan N Johnson[b], Pauline Bougaran[b], Celia E Shiau[b], Stephen L Rogers[b], and Victoria
10 L Bautch[a,b,d,#]
11
12
[a]Curriculum in Cell Biology and Physiology, The University of North Carolina at Chapel
Hill, Chapel Hill, North Carolina 27599, USA.
[b]Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill,
North Carolina 27599, USA.
[c]Department of Pathology, The University of North Carolina at Chapel Hill, Chapel Hill,
North Carolina 27599, USA.
[d]McAllister Heart Institute, The University of North Carolina at Chapel Hill, Chapel Hill,
North Carolina 27599, USA
26 [#]Corresponding author: Victoria L Bautch, PhD
27   Professor of Biology
28   Department of Biology, CB#3280
29   University of North Carolina at Chapel Hill
30   Chapel Hill, NC 27599 USA
31   E-mail: bautch@med.unc.edu

37 Running head: SUN1 Regulates Endothelial Junctions and Angiogenesis
44 Abstract: 206 words
45 Text: 47,835 characters with spaces (including figure legends; excluding methods,
46 references, and supplement)
47 Figures: 7 main, 8 supplemental, 6 movies
48 References: 113
49

SUMMARY

The nuclear membrane protein SUN1 promotes blood vessel formation and barrier

function by stabilizing endothelial cell-cell junctions. Communication between SUN1 and

endothelial cell junctions relies upon proper microtubule dynamics and Rho signaling far

from the nucleus, revealing long-range cellular communication from the nucleus to the

cell periphery that is important for vascular development and function.

ABSTRACT

Endothelial cells line all blood vessels, where they coordinate blood vessel formation

and the blood-tissue barrier via regulation of cell-cell junctions. The nucleus also

regulates endothelial cell behaviors, but it is unclear how the nucleus contributes to

endothelial cell activities at the cell periphery. Here we show that the nuclear-localized

LINC complex protein SUN1 regulates vascular sprouting and barrier function via

effects on endothelial cell-cell junction morphology and function. Loss of murine

endothelial *Sun1* impaired blood vessel formation and destabilized junctions, angiogenic

sprouts formed but retracted in SUN1-depleted sprouts, and zebrafish vessels lacking

Sun1b had aberrant junctions and defective cell-cell connections. At the cellular level,

SUN1 stabilized endothelial cell-cell junctions, promoted barrier function, and regulated

contractility. Mechanistically, SUN1 depletion altered cell behaviors via the cytoskeleton

without changing transcriptional profiles. Reduced peripheral microtubule density, fewer

junction contacts and increased catastrophes accompanied SUN1 loss, and microtubule

depolymerization phenocopied effects on junctions. Depletion of GEF-H1, a

microtubule-regulated Rho activator, or the LINC complex protein nesprin-1 rescued

defective junctions of SUN1-depleted endothelial cells. Thus, endothelial SUN1

regulates peripheral cell-cell junctions from the nucleus via LINC complex-based

microtubule interactions that affect peripheral microtubule dynamics and Rho-regulated

contractility, and this long-range regulation is important for proper blood vessel

sprouting and barrier function.

Keywords: Endothelial cell, SUN1, LINC, nucleus, junctions, angiogenesis, microtubules,

contractility, Rho, GEF-H1, nesprin-1

INTRODUCTION

Blood vessels form and expand via sprouting angiogenesis, a dynamic process

whereby endothelial cells migrate from pre-existing vessels to form new conduits

(Carmeliet & Jain, 2011; Wacker & Gerhardt, 2011; Kushner & Bautch, 2013; Bautch &

Caron, 2015). During angiogenesis, endothelial cell-cell junctions destabilize and

rearrange to allow for repolarization and migration towards pro-angiogenic cues (Esser

et al., 1998; Dejana, 2004; Blum et al., 2008). Specifically, the endothelial cell adherens

junction protein VE-cadherin is required for vascular sprouting and viability (Carmeliet et

al., 1999; Montero-Balaguer et al., 2009; Sauteur et al., 2014; Szymborska & Gerhardt,

2018). As vessels mature, endothelial cell junctions stabilize and form a functional

barrier that regulates egress of fluid and oxygen; barrier dysfunction leads to increased

permeability and severe disease (Claesson-Welsh, 2015; Rho et al., 2017; Claesson-

Welsh et al., 2021). Thus, regulation of endothelial cell junction stability is important

developmentally and for vascular homeostasis.

Adherens junctions are key to integrating and regulating both external and

internal cellular inputs from multiple sources, including the microtubule and actin

cytoskeletons (Ligon et al., 2001; Shaw et al., 2007; Bellett et al., 2009; Dejana &

Vestweber, 2013; Abu Taha & Schnittler, 2014). For example, increased actomyosin

contractility destabilizes endothelial cell adherens junctions, and disorganized junctional

actin accompanies VE-cadherin loss (Huveneers et al., 2012; Sauteur et al., 2014;

Angulo-Urarte et al., 2018). VE-cadherin loss also changes microtubule dynamics, and

disruption of microtubule dynamics destabilizes junctions and barrier function

(Komarova et al., 2012). Coordination of inputs from the actin and microtubule

cytoskeletons regulates endothelial cell barrier integrity and sprouting dynamics via

small GTPases (Birukova et al., 2006; Mavria et al., 2006; Sehrawat et al., 2008, 2011;

Wimmer et al., 2012; Szymborska & Gerhardt, 2018). In particular, RhoA signaling is

regulated by microtubules via GEF-H1, a RhoGEF that is inactive while bound to

microtubules and activated upon release, leading to RhoA activation, increased

actomyosin contractility, and changes to endothelial cell barrier function (Krendel et al.,

2002; Birukova et al., 2006; Birkenfeld et al., 2008). However, how the endothelial cell

nucleus affects these processes is poorly understood.

The nucleus is usually found far from the cell periphery and junctions, yet it is

important for functions critical to angiogenesis and vascular remodeling, such as

polarity, migration, and mechanotransduction (Tkachenko et al., 2013; Guilluy et al.,

2014; Graham et al., 2018), and perturbations of some nuclear membrane proteins

affect transcriptional profiles (Li et al., 2017; May & Carroll, 2018; Carley et al., 2021).

The linker of the nucleoskeleton and cytoskeleton (LINC) complex is comprised of both

SUN (Sad1p, UNC-84) and KASH (Klarsicht, ANC-1, Syne/Nesprin Homology) proteins

(Starr & Fridolfsson, 2010) that function as a bridge between the nucleus and the

cytoskeleton, and also link through subnuclear lamin filaments to chromatin (Haque et

al., 2006). SUN proteins localize to the inner nuclear membrane and bind KASH

proteins, or nesprins, from their C-terminus and lamins at their N-terminus, thus

providing a structural link from the nuclear cortex to the cellular cytoskeleton

(Padmakumar et al., 2005; Haque et al., 2006; McGee et al., 2006; Stewart-Hutchinson

et al., 2008). Nesprins are long spectrin-rich proteins localized to the outer nuclear

envelope that bind SUN proteins via their C-terminus while N-terminally interacting

indirectly with microtubules (via various motor proteins such as dynein and kinesin) and

intermediate filaments (via plectins), and directly with actin via calponin homology

domains (Ketema et al., 2007; Meyerzon et al., 2009; Zhang et al., 2009; Fridolfsson et

al., 2010; Starr & Fridolfsson, 2010). Two mammalian SUN proteins are ubiquitously

expressed, and based on functional consequences of SUN manipulations it has been

posited that SUN1 regulates microtubule-based functions while SUN2 coordinates actin

regulation (Zhu et al., 2017). However, *in vitro* binding studies do not reveal a SUN-

nesprin specificity to account for this bias (Stewart-Hutchinson et al., 2008; Ostlund et

al., 2009), so how complexes are assembled and sorted in cells is unclear. It is well-

established that the LINC complex integrates external inputs sensed by focal

adhesions, such as substrate stiffness, to regulate transcription (Carley et al., 2022), but

how the LINC complex relays signals from the nucleus to the cell periphery is less

understood.

144   The LINC complex functions in cultured endothelial cells, as knockdown of

nesprin-3 leads to impaired endothelial polarity under flow (Morgan et al., 2011), while

nesprin-2 and lamin A regulate proliferation and apoptosis in endothelial cells exposed

to shear stress (Han et al., 2015). Depletion of nesprin-1 alters tension on the nucleus

(Chancellor et al., 2010; Anno et al., 2012), and knockdown of nesprin-1 or nesprin-2

leads to reduced collective endothelial migration (Chancellor et al., 2010; King et al.,

2014). Recent work showed compromised matrix adhesion and barrier function of

cultured endothelial cells using a dominant negative KASH (Denis et al., 2021).

However, less is understood about the roles of the SUN proteins in endothelial cell

function.

155   The LINC complex is required for viability *in vivo*. Loss of both mammalian *Sun*

genes is embryonic lethal due to impaired neuronal nuclear migration required for

proper neuronal differentiation (Lei et al., 2009; Zhang et al., 2009). Global *Sun2* loss

affects epidermal nuclear positioning and cell adhesion, leading to alopecia (Stewart et

al., 2015), while global loss of both *Sun* genes impairs epidermal differentiation due to

altered integrin signaling (Carley et al., 2021). A role for the LINC complex in

mechanotransduction *in vivo* is suggested by findings that perturbations in mechanically

active skeletal and cardiac muscle affect function (Zhang et al., 2005, 2010; Lei et al.,

2009; Banerjee et al., 2014; Stroud et al., 2017; Zhou et al., 2017; van Ingen & Kirby,

2021). However, while global deletion of multiple LINC components to disrupt the entire

complex highlight its importance (Lei et al., 2009; Zhang et al., 2009; Carley et al.,

2021), these studies do not reveal functions of individual LINC components in specific

tissues. Whether the SUN proteins cell autonomously regulate the vascular

endothelium, which is also mechanically active due to outward pressure and shear

stress from blood flow, has not been explored.

171   Mutations in *LMNA* (lamin A/C) cause a premature aging syndrome linked to

cardiovascular defects (Capell & Collins, 2006). The LINC complex protein SUN1 is

mis-expressed in this disease (Chen et al., 2012), and cellular defects due to the *LMNA*

mutation are rescued by reduced levels of SUN1 protein, highlighting a potential

function for SUN1 in the disease pathology (Chen et al., 2012; Chang et al., 2019).

Here, we present an in-depth analysis of how the LINC complex component SUN1

affects blood vessel development and function *in vivo*. We found that *Sun1* cell-

autonomously regulates blood vessel sprouting and barrier function *in vivo,* and these

effects are consistent with SUN1 regulating endothelial cell functions via adherens

junction activity. In primary endothelial cells, nuclear SUN1 coordinates peripheral

microtubule dynamics that in turn regulate peripheral RhoGEF activation, junction

stability and barrier function. Thus, nuclear SUN1 that resides far from endothelial cell

junctions regulates endothelial cell-cell communication and blood vessel sprouting via a

novel microtubule-based integration pathway from the nucleus to the cell periphery.

RESULTS

The nuclear LINC protein SUN1 regulates vascular development

The LINC complex is important for cell migration (Chancellor et al., 2010; King et

al., 2014; Denis et al., 2021), and blood vessel formation involves extensive endothelial

cell migration; thus, we hypothesized that the LINC complex regulates angiogenic

sprouting. Because mutations in endothelial cell *LMNA* causative for human

cardiovascular disease are associated with expression changes in the LINC protein

SUN1 (Chen et al., 2012), and because *Sun1* has not been functionally analyzed in the

vascular endothelium *in vivo*, we first asked whether SUN1 is required for vascular

development. Utilizing a mouse line carrying a conditional *Sun1* allele that we generated

from *Sun1^{tm1a}* "knockout first" mice **(Figure 1-figure supplement 1A-B)**, *Sun1$^{fl/fl}$* mice

were bred to *Sun1$^{fl/+}$;Cdh5CreERT2/+* mice to generate *Sun1^{iECKO}* (inducible endothelial

cell knockout) mice with both endothelial cell-selective and temporal control over *Sun1*

excision. Examination of lung DNA, which is rich in endothelial cells, revealed

appropriate excision *in vivo* **(Figure 1-figure supplement 1C)**.

The retinal vasculature of *Sun1^{iECKO}* pups injected with tamoxifen at P (postnatal

day) 1-3 and harvested at P7 had significantly reduced radial expansion relative to

littermate controls **(Figure 1A-C, Figure 1-figure supplement 1D)**, consistent with a

role for *Sun1* in vascular development. *Sun1^{iECKO}* retinas also had increased density at

the vascular front **(Figure 1-figure supplement 1E)**, consistent with defects in

sprouting that prevent expansion and thus increase density (Hellström et al., 2007;

Ricard et al., 2012; Angulo-Urarte et al., 2018). Since vessel densities in the plexus

ahead of arteries and veins exhibit heterogeneity, we measured by area and found

increased density in the plexus ahead of both arteries and veins in *Sun1^{iECKO}* retinas

(Figure 1B, D). Because adherens junction dynamics regulate vascular sprouting

(Sauteur et al., 2014), this mutant phenotype suggested that endothelial cell-cell

junctions were affected by loss of *Sun1*. VE-cadherin localization, a readout of junction

integrity (Huveneers et al., 2012; Bentley et al., 2014; Wylie et al., 2018; Vion et al.,

2020)**,** was significantly less linear and more punctate in *Sun1iECKO* vessels, indicating

increased adherens junction turnover and junction instability **(Figure 1E-F)**. Dextran

injection was used to functionally evaluate the effects of *Sun1* loss on vascular barrier

function *in vivo*, and *Sun1iECKO* mice had increased signal in the surrounding tissue

compared to controls **(Figure 1-figure supplement 1F)**, suggesting increased vessel

permeability. Together, these data indicate a specific role for SUN1 in angiogenic

sprouting and endothelial cell-cell junctions *in vivo*.

Nuclear SUN1 is required for sprouting angiogenesis

225   *Sun1* loss disrupts vascular development in the postnatal mouse retina **(Figure**

1), but this tissue is not amenable to long-term live image analysis. To query dynamic

aspects of angiogenic sprouting, which occurs via regulated changes in endothelial

adherens junction stability (Sauteur et al., 2014; Angulo-Urarte et al., 2018; Wylie et al.,

2018), we utilized a 3D sprouting model (Nakatsu & Hughes, 2008) coupled with

temporal image acquisition. Reduced levels of endothelial cell SUN1 via siRNA

knockdown (KD) **(Figure 2-figure supplement 1A-B)** led to significantly decreased

sprout length and branching **(Figure 2A-C)**, reminiscent of the decreased radial

expansion of *Sun1iECKO* retinal vessels described above. SUN1 depletion did not

significantly influence the proportion of EdU-labeled or Ki67 stained cells **(Figure 2-**

figure supplement 1C-F), indicating that the abnormal sprouting and branching is not

downstream of reduced proliferation. Live-cell imaging revealed that control sprouts

typically elongated over time, with little retraction once they extended from the bead

(Figure 2D-E, Movie 1). In contrast, SUN1 KD sprouts retracted more often, and many

mutant sprouts collapsed partially or completely **(Figure 2D-E, Movie 2)**. SUN1 KD

sprouts also showed a more diffuse VE-cadherin junction pattern **(Figure 2F-G)**, similar

to those of *Sun1^iECKO^* mice and indicative of over-activated junctions. Thus, SUN1 is

required for proper vascular sprout dynamics and morphology, and reduced sprout

length and branching are likely downstream of excess sprout retractions and perturbed

junctions in SUN1-depleted vessels.

The LINC complex is important for mechanotransduction in muscle fibers with

high mechanical loads (van Ingen & Kirby, 2021), and sprouting angiogenesis is

regulated by mechanical forces arising from blood pressure and blood flow (Huang et

al., 2003). To determine whether SUN1 also regulates sprouting dynamics under

laminar flow *in vivo*, we analyzed embryonic zebrafish using a *Tg(fli:LifeAct-GFP)*

reporter that labels the endothelial actin cytoskeleton. Zebrafish have two *Sun1* genes,

sun1a and *sun1b*; the SUN domain of *sun1b* is more homologous to human *SUN1,* and

Sun1b is more highly expressed in cardiovascular tissue, so this gene was chosen for

manipulation. Sun1b depletion in zebrafish embryos via morpholino (MO) injection led to

significantly increased numbers of shorter endothelial cell filopodia at 33-34 hpf (hours

post fertilization) in the inter-segmental vessels (ISVs) that sprout towards and connect

to the dorsal longitudinal anastomotic vessel (DLAV) **(Figure 3A-C)**. Like the

morphants, fish carrying a point mutation in the *sun1b* gene leading to a premature stop

codon (*sun1b^sa33109^*, see Methods for details) had shorter filopodia, although filopodia

numbers were unchanged in the mutant background **(Figure 3D-F)**. Because increased

filopodia are typically seen in actively migrating and sprouting endothelial cells

(DeLisser, 2011), these changes are consistent with Sun1b regulating endothelial cell

activation in developing zebrafish vessels exposed to physiological flow forces.

Next, *sun1b* morphant fish were imaged from 26-36 hpf to determine the effects

of Sun1b depletion on vascular sprouting *in vivo*. In controls, the ISVs sprouted towards

the dorsal plane and connected to the DLAV between 32-36 hpf **(Figure 3G-H, Movie**

3). In contrast, numerous ISVs either failed to reach the DLAV or made aberrant

connections in *sun1b* morphant fish **(Figure 3G-H, Movie 4)**. Staining for the tight

junction protein ZO-1 revealed less linear and more abnormally shaped junctions in

sun1b morphant fish **(Figure 3I-J)**. These results complement the 3D sprouting analysis

and show that the nuclear LINC complex protein SUN1 is important in regulating

endothelial cell sprouting dynamics and junction morphology under flow forces *in vivo*.

SUN1 stabilizes endothelial cell-cell junctions and regulates barrier function

SUN1 loss or depletion in mouse, zebrafish, and 3D sprouting models resulted in

abnormal endothelial cell-cell junctions and sprouting behaviors. Thus, we examined

more rigorously the hypothesis that SUN1 regulates endothelial cell junction stability

and morphology. Primary human umbilical vein endothelial cells (HUVEC) in confluent

monolayers had more serrated cell-cell junctions without altered levels of VE-cadherin

protein expression after SUN1 depletion, indicative of activated and destabilized

junctions **(Figure 4A, Figure 4-figure supplement 1A-B)**. Activated and destabilized

endothelial cell adherens junctions are associated with impaired vascular barrier

function, so we measured electrical resistance across endothelial monolayers using

Real Time Cell Analysis (RTCA) that provides an impedance value positively correlated

with barrier function. SUN1 KD endothelial cells had reduced electrical resistance

compared to controls **(Figure 4B-C)**, indicative of impaired barrier function and

consistent with the increased dextran permeability *in vivo*. Endothelial cells *in vivo* are

exposed to blood flow that remodels endothelial junctions (Seebach et al., 2000; Yang

et al., 2020). SUN1 depleted HUVEC exposed to laminar shear stress for 72h elongated

and aligned properly, but adherens junctions were more serrated and allowed for

significantly more matrix exposure as assessed by a biotin labeling assay (Dubrovskyi

et al., 2013) under both static and flow conditions **(Figure 4D-F, Figure 4-figure**

supplement 1C-D). These findings support the hypothesis that nuclear SUN1 regulates

vascular barrier function via effects on endothelial cell junction stability and morphology.

Dysfunctional cell-cell junctions can result from abnormal junction formation or

the inability of formed junctions to stabilize. To determine how SUN1 functionally

regulates endothelial junctions, adherens junctions were disassembled via Ca^{2+}

chelation, then reformed upon chelator removal. Junctions were measured using line

scans of VE-cadherin intensity along the cell-cell junctions **(Figure 4-figure**

supplement 1E), such that junctions with a linear VE-cadherin signal (stable) had a

higher value than those with more serrated patterns (destabilized). No significant

difference between SUN1 KD and control junctions was seen at early times post-

washout, indicating that SUN1 depletion does not affect adherens junction formation

(Figure 4G-H). However, later times post-washout revealed a significant increase in

serrated junctions and gaps between endothelial cells in SUN1 KD monolayers relative

to controls **(Figure 4G-H)**. Consistent with these findings, SUN1 KD endothelial cells

also had increased VE-cadherin internalization at steady state, consistent with actively

remodeling junctions **(Figure 4-figure supplement 1F-G)**. Thus, SUN1 is not required

to form endothelial cell-cell junctions but is necessary for proper junction maturation and

stabilization.

SUN1 regulates microtubule localization and dynamics in endothelial cells absent

effects on gene transcription

316   We next considered how SUN1 that resides in the nuclear membrane regulates

cell behaviors at the cell periphery. SUN1 may affect cell junctions directly via the

cytoskeleton, or SUN1 may indirectly affect junctions downstream of gene expression

regulation. Because SUN1 is reported to regulate gene transcription and RNA export in

other cell types (Li et al., 2017; May & Carroll, 2018), we asked whether endothelial cell

transcriptional profiles were altered by SUN1 depletion. To our surprise, RNASeq

analysis of HUVEC under both static and flow conditions revealed essentially no

significant changes in RNA profiles except for *SUN1* itself **(Table 1, Figure 5-figure**

supplement 1), while in the same experiment we documented extensive expression

changes in both up- and down-regulated genes between control HUVEC in static vs.

laminar flow conditions, as we and others have shown (Conway et al., 2010; Liu et al.,

2021; Maurya et al., 2021; Ruter et al., 2021). These findings are similar to a re-analysis

of HeLa cell data (data not shown, (Li et al., 2017)) that revealed few significant

changes in gene expression following SUN1 depletion. Thus, although the LINC

complex is important for nuclear communication that can affect gene expression, SUN1

is not required for this communication in endothelial cells.

333   We hypothesized that SUN1 directly regulates endothelial cell junction stability

via the cytoskeleton. SUN1 has a functional relationship with the microtubule

cytoskeleton (Zhu et al., 2017), and microtubule dynamics regulate endothelial cell-cell

junctions (Sehrawat et al., 2008, 2011; Komarova et al., 2012). Thus, we asked whether

the observed adherens junction defects following SUN1 depletion resulted from

changes in the microtubule cytoskeleton. Microtubule depolymerization via nocodazole

treatment phenocopied SUN1 KD and destabilized adherens junctions in control

monolayers but did not exacerbate the junction defects seen with SUN1 KD **(Figure 5A-**

B). These findings suggest that endothelial cell junction defects are downstream of

microtubule perturbations induced by SUN1 depletion.

344   We next investigated how SUN1 affects microtubule localization in endothelial

cell monolayers and found that significantly fewer microtubules reached the cell

periphery or surrounded the nucleus in SUN1 depleted cells **(Figure 5C, E, Figure 5-**

figure supplement 2A-B). The changes in peripheral microtubule localization were

accompanied by significantly fewer microtubule-junction contacts, while α-tubulin levels

around the MTOC were not affected **(Figure 5C, D, F)**. Microtubule dynamics were

assessed via tip tracking using mCherry-labeled tip protein EB3, which decorated the

microtubule lattice and concentrated at growing microtubule tips **(Figure 5G)**. This

labeling pattern can occur with EB overexpression but does not affect growth rate

(Komarova et al., 2005), so the decorated lattice was used to assess microtubule

catastrophe and shrinkage. Microtubules in SUN1 depleted cells had increased

shrinkage rates (a more negative value) coupled with increased catastrophe rate and

time spent shrinking **(Figure 5G-K, Movies 5, 6)**, consistent with elevated microtubule

depolymerization and impaired microtubule dynamics downstream of SUN1 loss, while

the microtubule growth rate was unchanged. Taken together, these results indicate that

loss of SUN1 impairs microtubule localization and dynamics, and these changes

associate with destabilized endothelial cell junctions.

SUN1 regulates endothelial cell contractility

Microtubule dynamics communicate with the actin cytoskeleton to regulate cell-

cell junctions (Verin et al., 2001; Birukova et al., 2004, 2006), and cellular changes in

actomyosin contractility contribute to junction activation (Rauzi et al., 2010; Huveneers

et al., 2012). We hypothesized that actomyosin mis-regulation induced by SUN1

depletion contributes to endothelial cell junction destabilization and found that SUN1

depletion led to ectopic stress fibers, distinct radial actin bundles at the periphery, and

increased phosphorylated myosin light chain (ppMLC), consistent with increased

actomyosin contractility **(Figure 6-figure supplement 1A-B)**. Pharmacological

blockade of myosin-II ATPase rescued the destabilized cell junctions, ectopic stress

fibers and radial actin structures seen with SUN1 depletion **(Figure 6A-B, Figure 6-**

figure supplement 1C), suggesting that SUN1 depletion increases contractility. In

contrast, thrombin treatment that induces contractility produced over-activated junctions

in controls that phenocopied SUN1 depletion but did not further activate junctions of

SUN1 KD cells, indicating that SUN1 loss induces a maximal contractile state **(Figure**

6C-D). Taken together, these data indicate that SUN1 loss results in hypercontractility

and destabilized endothelial cell-cell junctions.

SUN1 affects endothelial junctions through microtubule-associated Rho GEF-H1

To better understand the link between microtubule dynamics, actomyosin

contractility, and junction regulation in endothelial cells, we examined Rho signaling

downstream of SUN1 in endothelial cells. SUN1 silencing increases RhoA activity in

HeLa cells (Thakar et al., 2017), and pharmacological inhibition of Rho kinase (ROCK)

signaling in SUN1 depleted endothelial cells rescued the destabilized endothelial cell-

cell junctions **(Figure 6-figure supplement 2A-B)** and prevented the radial actin

structures **(Figure 6-figure supplement 1D)**. RhoA signaling and barrier function in

endothelial cells is regulated by the microtubule-associated RhoGEF, GEF-H1, which is

inactive while bound to microtubules and activated following microtubule

depolymerization and release (Krendel et al., 2002; Birukova et al., 2006; Birkenfeld et

al., 2008). We hypothesized that the impaired microtubule dynamics and Rho-

dependent hypercontractility observed following SUN1 depletion were linked via GEF-

H1. Consistent with this idea, GEF-H1 strongly localized to peripheral microtubules in

control cells but was significantly less localized and more diffuse in SUN1 depleted

endothelial cells **(Figure 6E-F)**. Furthermore, depletion of GEF-H1 in endothelial cells

that were also depleted for SUN1 rescued the destabilized cell-cell junctions observed

with SUN1 KD alone **(Figure 6G-H, Figure 6-figure supplement 2C)**, showing that

GEF-H1 is required to transmit the effects of SUN1 depletion to endothelial cell

junctions. Thus, nuclear SUN1 normally promotes microtubule-GEFH1 interactions to

regulate actomyosin contractility and endothelial cell-cell junction stability, providing a

novel linkage from the LINC complex to endothelial cell junctions via the microtubule

cytoskeleton.

SUN1 exerts its effects on endothelial junctions through nesprin-1

Since SUN1 binds nesprins to interact with the cytoskeleton, we considered

whether SUN1-nesprin interactions were involved in SUN1 regulation of microtubule

dynamics and endothelial cell junction stability. The KASH protein nesprin-1 modulates

tight junction protein localization under laminar shear stress (Yang et al., 2020) and

regulates microtubule dynamics at the nucleus in muscle syncytia (Gimpel et al., 2017).

Co-depletion of SUN1 with nesprin-1 in endothelial cells rescued the effects of SUN1

depletion on junction morphology and functional destabilization measured by matrix

biotin labeling **(Figure 7A-B, Figure 7-figure supplement 1A)**. This rescue extended to

the cytoskeleton, as co-depletion rescued both decreased peripheral microtubule

density and microtubule-GEF-H1 contacts seen with SUN1 depletion **(Figure 7-figure**

supplement 1B-E). Thus, the LINC complex protein nesprin-1 is required to transmit

the effects of SUN1 depletion to endothelial cell junctions.

DISCUSSION

The nucleus compartmentalizes and organizes genetic material, but how the

nucleus directly communicates with other organelles and the cytoskeleton to regulate

cell behaviors is poorly understood. Here, we show for the first time that the nuclear

LINC complex protein SUN1 regulates angiogenic sprouting and vascular barrier

function via long-distance regulation of endothelial cell-cell junctions *in vivo*, and that

these effects go through the microtubule cytoskeleton to regulate microtubule dynamics

and actomyosin contractility. Our data is consistent with a model in which endothelial

SUN1 stabilizes peripheral microtubules that coordinate activity of the microtubule-

associated Rho exchange factor GEF-H1. GEF-H1 becomes activated to stimulate Rho

signaling upon release from microtubules, and we posit that peripheral microtubules

normally regulate local GEF-H1 activity to maintain appropriate microtubule-junction

interactions and actomyosin contractility in endothelial cells, leading to cell-cell junctions

that remodel to support angiogenic sprouting and maintain vessel barrier integrity

(Figure 7C). Loss of SUN1 reduces peripheral microtubules, resulting in GEF-H1 over-

activation, elevated RhoA signaling, and increased contractility, leading to destabilized

endothelial cell adherens junctions that impair blood vessel formation and function in

fish and mice. The LINC complex is further implicated in endothelial junction regulation

by our finding that the KASH protein nesprin-1 is required for the effects of SUN1 loss

on microtubules, GEF-H1, and proper junction function. Thus, we describe a specific

role for SUN1 as a critical mediator of communication between the endothelial cell

nucleus and the cell periphery via microtubule regulation of GEF-H1.

We show that the LINC complex protein SUN1 is required for endothelial

adherens junction stabilization and proper blood vessel formation and function *in vivo*.

Although *Sun2* partially compensates for loss of *Sun1* developmentally (Lei et al., 2009;

Zhang et al., 2009) and thus may prevent more profound vascular effects, the significant

vascular defects in vessels lacking *Sun1* indicate non-redundant functions for SUN1 in

vascular development. The adherens junctions of expanding retinal vessels are

destabilized in mice lacking endothelial *Sun1*, and this cellular phenotype is

accompanied by increased vessel permeability, reduced radial expansion, and

increased network density, indicating that destabilized junctions contribute to the vessel

network perturbations. Mutations in other genes that affect endothelial cell junction

integrity, such as *Smad6*, *Pi3kca*, and *Yap/Taz*, also perturb retinal angiogenesis

(Angulo-Urarte et al., 2018; Neto et al., 2018; Wylie et al., 2018). Live-image analysis of

active vessel sprouting showed that SUN1 regulates filopodia dynamics and

anastomosis in zebrafish and sprout dynamics in mammalian endothelial cell sprouts,

and these vessels also had abnormal junctions in the absence of SUN1. Altered sprout

dynamics are found in other scenarios where VE-cadherin is absent or abnormal such

as Wnt inhibition, loss of VE-cadherin, PI3-kinase inhibition, and excess centrosomes

(Sauteur et al., 2014, 2017; Kushner et al., 2016; Angulo-Urarte et al., 2018; Hübner et

al., 2018; Buglak et al., 2020). The LINC complex regulates endothelial cell aggregation

into tube-like structures in Matrigel (King et al., 2014; Denis et al., 2021), consistent with

our findings that highlight a central role for SUN1 in blood vessel formation *in vivo*.

Unlike genes encoding components of junctions and signaling effectors, SUN1 functions

at significant cellular distances from cell junctions.

How does SUN1 regulate endothelial cell-cell junctions from a distance? The

importance of the LINC complex in transducing signals from the cell periphery and

outside the cell to the nucleus to affect nuclear envelope properties and gene

transcription is well-studied (Carley et al., 2022); however, how information goes from

the nucleus to the cell periphery is less well-understood. Although the LINC complex is

clearly important in nucleus-to-cytoplasm communication, most studies have globally

manipulated the LINC complex (Zhang et al., 2009; Graham et al., 2018; Carley et al.,

2021; Denis et al., 2021), and how individual components contribute is not well-

understood. Recently Ueda et al (2022) found that SUN1 regulated focal adhesion

maturation in non-endothelial cells *in vitro* via effects on the actin cytoskeleton (Ueda et

al., 2022), and we found that SUN1 regulates endothelial cell-cell junctions through

microtubules, suggesting that SUN1 is important for signaling from the nucleus to the

cell periphery by regulating cytoskeletal organization at several levels. SUN1 is thought

to be important in microtubule-associated LINC complex functions (Zhu et al., 2017),

and disruption of microtubules or their dynamics destabilizes adherens junctions in both

endothelial and non-endothelial cells (Komarova et al., 2012; Vasileva & Citi, 2018).

Microtubule plus end dynamics regulate E-cadherin accumulation at epithelial cell

adherens junctions (Stehbens et al., 2006), and dynein is thought to anchor microtubule

plus ends to junctions via the plus end binding protein EB1 (Ligon et al., 2001; Shaw et

al., 2007; Bellett et al., 2009). Microtubule minus ends also regulate protein

accumulation at junctions and are anchored to the junction through a complex involving

p120 catenin, PLEKHA7, and CAMSAP3 (Meng et al., 2008). Our data show that SUN1

depletion impairs both microtubule dynamics and microtubule localization. Specifically,

elevated rates of microtubule catastrophe and shrinkage absent changes in growth rate

likely account for reduced peripheral microtubule density and microtubule-junction

contacts. Endothelial junction destabilization was phenocopied by microtubule

depolymerization, consistent with our model that the influence of SUN1 on microtubule

dynamics and localization is important for the stabilization of endothelial cell junctions.

Here we show that SUN1 regulates endothelial cell junctions via the microtubule-

regulated Rho activator GEF-H1. GEF-H1 specifically modulates RhoA signaling at

endothelial cell adherens junctions to influence VE-cadherin internalization (Juettner et

al., 2019). Microtubule depolymerization releases GEF-H1 to activate RhoA signaling

and elevate actomyosin contractility (Krendel et al., 2002; Birkenfeld et al., 2008) in non-

endothelial cells, while direct manipulation of GEF-H1 via depletion or blockade

attenuates agonist-induced endothelial barrier dysfunction (Birukova et al., 2006). Our

work revealed that peripheral GEF-H1 localization was more diffuse following

endothelial SUN1 silencing, suggesting its activation with SUN1 loss, and Rho-kinase

inhibition rescued the SUN1 depletion-induced destabilization of endothelial cell

junctions. These findings support that altered microtubule dynamics downstream of

SUN1 depletion promote the release and over-activation of peripheral GEF-H1, and this

activation destabilizes cell junctions.

Thus, SUN1 regulation of microtubule dynamics is linked to its regulation of

endothelial cell junction stability, although exactly how SUN1 influences microtubule

dynamics and function is unclear. SUN1 resides in the nuclear envelope and alters

gene transcription in other cell types (Li et al., 2017; May & Carroll, 2018). Our

transcriptional analysis of endothelial cells under flow did not reveal significant

transcriptome changes with SUN1 depletion, indicating that SUN1 regulation of

endothelial junctions occurs via its role in LINC complex interactions with the

cytoskeleton. Nesprin-1, a KASH protein that functions in LINC complex-cytoskeletal

interactions, is shown here to mediate the effects of SUN1 loss on endothelial junctions,

suggesting that SUN1 normally sequesters nesprin-1 to prevent formation of ectopic

complexes. Since fewer microtubules surrounded the nucleus in SUN1-depleted

endothelial cells, SUN1 may normally prevent abnormal or unstable nesprin-1 LINC

complexes that promote microtubule depolymerization, and SUN1 loss allows these

complexes to form. SUN1 may compete for nesprin-1 binding with other nuclear

envelope proteins that interact with nesprin-1, such as SUN2 or nesprin-3 (Stewart-

Hutchinson et al., 2008; Taranum et al., 2012; Yang et al., 2020). This idea is consistent

with the finding that SUN1 antagonizes SUN2-based LINC complexes that promote

RhoA activity in HeLa cells, although microtubule localization changes were not

reported (Thakar et al., 2017).

Our finding that SUN1 regulates endothelial cell barrier function and blood vessel

sprouting has implications for diseases associated with aging, as vascular defects

underlie most cardiovascular disease. Children with Hutchinson-Gilford Progeria

Syndrome (HGPS) have a mutation in the *LMNA* gene encoding lamin A/C, resulting in

accumulation of an abnormal lamin protein called progerin; these patients age rapidly

and die in their early to mid-teens from severe atherosclerosis (De Sandre-Giovannoli,

2003; Eriksson et al., 2003; Olive et al., 2010). Progerin has increased SUN1 affinity

that leads to SUN1 accumulation in HGPS patient cells (Haque et al., 2010; Chen et al.,

2012, 2014; Chang et al., 2019). Endothelial cells also accumulate SUN1 in HGPS

mouse models (Osmanagic-Myers et al., 2019), and loss of *Sun1* partially rescues

progeria phenotypes in mouse models and patient cells (Chen et al., 2012; Chang et al.,

2019). Thus, nuclear membrane perturbations affecting SUN1 cause disease, and here

we find that nuclear SUN1 regulates microtubules to affect both the microtubule and

actin cytoskeletons. These effects are transmitted to endothelial cell-cell junctions far

from the site of SUN1 localization to influence endothelial cell behaviors, blood vessel

sprouting and barrier function.

MATERIALS & METHODS

Microscopy

Unless otherwise stated, all imaging was performed as follows: confocal images

were acquired with an Olympus confocal laser scanning microscope and camera

(Fluoview FV3000, IX83) using 405nm, 488nm, 561nm, and 640nm lasers and a

UPlanSApo 40x silicone-immersion objective (NA 1.25), UPlanSApo 60x oil-immersion

objective (NA 1.40), or UPlanSApo 100x oil-immersion objective (NA 1.40). Imaging

was performed at RT for fixed samples. Images were acquired with the Olympus

Fluoview FV31S-SW software and all image analysis, including Z-stack compression,

was performed in Fiji (Linkert et al., 2010; Schindelin et al., 2012). Any adjustments to

brightness and contrast were performed evenly for images in an experiment.

Mice

All experiments involving animals were performed with approval from the

University of North Carolina, Chapel Hill Institutional Animal Care and Use Committee

(IACUC). C57Bl6N-*Sun1$^{tm1a(EUCOMM)Wtsi}$*/CipheOrl mice were obtained from the European

Mouse Mutant Archive (EMMA) mouse repository. FlpO-B6N-Albino (Rosa26-FlpO/+)

mice were obtained from the UNC Animal Models Core. *Tg(Cdh5-cre/ERT2)1Rha* mice

were generated by Dr. Ralf Adams (Sörensen et al., 2009) and obtained from Cancer

Research UK. The *Sun1^{tm1a}* allele was identified via genomic PCR to amplify the LacZ

insertion (Forward: 5'- ACTATCCCGACCGCCTTACT-3'; Reverse: 5'-

TAGCGGCTGATGTTGAACTG-3'). The *Sun1^{fl}* allele was generated by breeding

Sun1^{tm1a} mice with FlpO-B6N-Albino (Rosa26-FlpO/+) mice to excise the *lacZ* insertion.

The *Sun1^{fl}* allele was identified via genomic PCR using the following primers (Forward:

5'- GCTCTCTGAAACATGGCTGA-3'; Reverse: 5'- ATCCGGGGTGTTTGGATTAT-3').

Sun1^{fl} mice were bred to *Tg(Cdh5-cre/ERT2)1Rha* mice to generate

Sun1^{fl/fl};*Cdh5CreERT2* pups for endothelial-selective and temporally controlled deletion

of exon 4 of the *Sun1* gene. The excised *Sun1* allele was identified via genomic PCR on

lung tissue using the following primers (Forward: 5'- CTTTTGGGCTGCTCTGTTGT-3';

Reverse: 5'- ATCCGGGGTGTTTGGATTAT-3'). PCR genotyping for FlpO and

Cdh5CreERT2 mice was performed with the following primers (FlpO: Forward: 5'-

TGAGCTTCGACATCGTGAAC -3'; Reverse: 5'-TCAGCATCTTCTTGCTGTGG-3')

(Cdh5CreERT2: Forward: 5'- TCCTGATGGTGCCTATCCTC-3'; Reverse: 5'-

CCTGTTTTGCACGTTCACCG-3'). Induction of Cre was performed via IP injection of

pups at P1, P2, and P3 with 50µl of 1mg/ml tamoxifen (T5648, Sigma) dissolved in

sunflower seed oil (S5007, Sigma). Littermates lacking either *Cdh5CreERT2* or the

Sun1^{fl} allele were used as controls.

Mouse retinas

Tamoxifen-injected mice were sacrificed at P7, eyes were collected, fixed in 4%

PFA for 1h at RT, then dissected and stored at 4°C in PBS for up to 2 weeks (Chong et

al., 2017). Retinas were permeabilized in 0.5% Triton X-100 (T8787, Sigma) for 1h at

RT, blocked for 1h at RT in blocking solution (0.3% Triton X-100, 0.2% BSA (A4503,

Sigma), and 5% goat serum (005-000-121, Jackson Immuno)), then incubated with VE-

cadherin antibody (anti-mouseCD144, 1:100, 550548, BD Pharmingen) in blocking

solution overnight at 4°C. Samples were washed 3X, then incubated with Isolectin B4

AlexaFluor 488 (1:100, I21411, ThermoFisher) and goat anti-rat AlexaFluor 647 (1:500,

A21247, Life Technologies) for 1h at RT. Retinas were mounted with Prolong Diamond

Antifade mounting medium (P36961, Life Technology) and sealed with nail polish.

Images were obtained using either a UPlanSApo 10x air objective (NA 0.40) or

UPlanSApo 40x silicone-immersion objective (NA 1.25). Percent radial expansion was

calculated by dividing the distance from the retina center to the vascular front by the

distance from the retina center to the edge of the tissue. Four measurements/retina

were averaged. Vascular density was measured by imaging a 350µm x 350µm ROI at

the vascular edge. Fiji was used to threshold images, and the vessel area was

normalized to the area of the ROI (n=4 ROI/retina, chosen at 2 arteries and 2 veins).

Junctions were measured by taking the ratio of the mean intensity of the junction and

the mean intensity of the area immediately adjacent to the junction. Mean intensity was

measured via line scans in Fiji. 16 junctions/retina were measured.

Retina blood vessel permeability

Tamoxifen-injected mice were anesthetized at P7 in isoflurane for 5 min. The

abdomen was opened, and the diaphragm was cut. 100µl of 5mg/ml 10kDa Dextran-

Texas Red (D1863, Invitrogen) in PBS was injected into the left ventricle of the heart.

Eyes were immediately collected and fixed in 4% PFA for 1h at RT, then dissected and

stained as described above. Leak was determined by making a mask of the vessel area

using the isolectin channel, then assessing the dextran signal outside the vessel.

3D sprouting angiogenesis assay

The 3D sprouting angiogenesis assay was performed as previously described

(Nakatsu & Hughes, 2008; Nesmith et al., 2017). 48h following siRNA knockdown,

HUVEC were coated onto cytodex 3 microcarrier beads (17048501, GE Healthcare Life

Sciences) and embedded in a fibrin matrix by combining 7µl of 50U/ml thrombin (T7201-

500UN, Sigma) with 500µl of 2.2 mg/ml fibrinogen (820224, Fisher) in a 24-well glass-

bottomed plate (662892, Grenier Bio). The matrix was incubated for 20 min at RT

followed by 20 min at 37°C to allow the matrix to solidify. EGM-2 was then added to

each well along with 200μl of normal human lung fibroblasts (CC2512, Lonza) at a

concentration of 2×10^5 cells/ml. At day 7 of sprouting, fibroblasts were removed via

trypsin treatment (5X-trypsin for 3min at 37°C), and samples were fixed in 4% PFA for

15 min at RT. 0.5% Triton X-100 in DBPS was added to the wells, and incubation was

overnight at 4°C. After rinsing 3X in DPBS, samples were blocked (5% goat serum

(005-000-121, Jackson Immuno), 1% BSA (A4503, Sigma), and 0.3% Triton X-100

(T8787, Sigma)) overnight at 4°C. Samples were rinsed 3X in DPBS then anti-VE-

cadherin antibody (1:1000, 2500S, Cell Signaling) in blocking solution was added for

24h at 4°C. Samples were rinsed 3X 10 min in 0.5% Tween 20 then washed overnight

at 4°C in 0.5% Tween. Samples were rinsed 3X in DPBS, then DAPI (0.3μM,

10236276001, Sigma) and AlexaFluor488 Phalloidin (1:50, A12379, Life Technologies)

in blocking solution were added to the wells, and incubation was overnight at 4°C prior

to rinsing 3X in DPBS. For whole bead analysis, images were acquired in the Z-plane

using a UPlanSApo 20x oil-immersion objective (NA 0.58) and processed in Fiji.

Average sprout length was measured by tracing each sprout from base (bead) to tip,

then averaging lengths per bead. Branching was measured by counting total branch

points and normalizing to total sprout length per bead using the AnalyzeSkeleton plugin

(Arganda-Carreras et al., 2010). For junctions, images were acquired in the Z-plane

using a UPlanSApo 60x oil-immersion objective (NA 1.40). Junctions were measured by

taking the ratio of the mean intensity of the junction and the mean intensity of the area

immediately adjacent to the junction. Mean intensity was measured via line scans in Fiji.

Live imaging on HUVEC sprouts was performed between days 4-6.5 of

sprouting. Images were acquired on an Olympus VivaView Incubator Fluorescence

Microscope with a UPLSAPO 20x objective (NA 0.75) and 0.5x magnification changer

(final magnification 20x) with a Hamamatsu Orca R2 cooled CCD camera at 30 min

intervals for 60h at 37°C. Images were acquired using the MetaMorph imaging software

and analyzed in Fiji. Sprouts were considered to have "retracted" if they regressed

towards the bead for at least 3 imaging frames (1.5h).

Zebrafish

All experimental Zebrafish (*Danio rerio*) procedures performed in this study were

reviewed and approved by the University of North Carolina Chapel Hill Animal Care and

Use Committee. Animals were housed in an AAALAC-accredited facility in compliance

with the *Guide for the Care and Use of Laboratory Animals* as detailed on protocols.io

(dx.doi.org/10.17504/protocols.io.bg3jjykn). *Tg(fli:LifeAct-GFP)* was a gift from Wiebke

Herzog. *sun1b^{sa33109}* mutant fish were obtained from the Zebrafish International

Resource Center (ZIRC). For genotyping, the target region of the *sun1b* gene was

amplified via genomic PCR using the following primers (Forward: 5'-

GGCTGCGTCAGACTCCATTA-3'; Reverse: 5'-TTGAGTTAAACCCAGCGCCT-3'). The

amplicon was then sequenced by Sanger sequencing (GENEWIZ) using the forward

primer. Morphant fish were obtained by injecting 2.5-5ng of non-targeting (NT) (5'-

CCTCTTACCTCAGTTACAATTTATA-3', GeneTools, LLC) or *sun1b* (5'-

CGCAGTTTGACCATCAGTTTCTACA-3', GeneTools, LLC) morpholinos into

Tg(fli:LifeAct-GFP) embryos at the 1-cell stage. Fish were grown in E3 medium at

28.5°C to 33-34 hpf.

Zebrafish imaging

Dechorionated embryos were incubated in ice cold 4% PFA at 4°C overnight or

RT for 2h. Embryos were permeabilized in 0.5% Triton X-100 in PBST (DPBS + 0.1%

Tween 20) for 1 h at RT then blocked (PBST + 0.5% Triton X-100 + 1% BSA + 5% goat

serum + 0.01% sodium azide) for 2h at RT. Anti-ZO1 primary antibody (1:500, 33-9100,

Thermo Fisher) was added overnight at 4°C. Embryos were rinsed in PBST overnight at

4°C. Goat anti-mouse AlexaFluor647 secondary antibody (1:1000, A-21236, Life

Technologies) was added overnight at 4°C. Embryos were washed 3X in PBST for 30

min then overnight in PBST at 4°C. Embryos were rinsed in PBS and a fine probe was

used to de-yolk and a small blade to separate the trunk from the cephalic region.

Samples were mounted using Prolong Diamond Antifade mounting medium (P36961,

Life Technology) and the coverslip was sealed with petroleum jelly. Imaging was at RT

using a UPlanSApo 20x oil-immersion objective (NA 0.58) or a UPlanSApo 60x oil-

immersion objective (NA 1.40) with an additional 3x magnification, for a total

magnification of 180x. Filopodia length was measured by drawing a line from the

filopodia base to the tip. Filopodia number was measured by counting the number of

filopodia and normalizing to the total vessel length. Filopodia in at least 3 areas per fish

were measured. Junctions were analyzed by drawing a line along the junction then

normalizing to the shortest distance between the two ends of the junction. At least 3

junctions were measured per fish.

For live imaging of zebrafish, fish were dechorionated and then anesthetized with

1x Tricaine in E3 for 5 min. Fish were embedded in 0.5% agarose in 1x Tricaine-E3

medium on a glass-bottomed plate in a stage-top incubator (TOKAI HIT, WSKM) at

28.5°C. Images were acquired using a UPlanSApo 40x air objective (NA 0.95) every

15min for 10-15h. ISVs that did not reach the DLAV or connected at non-consistent

intervals were considered to have a missing or aberrant DLAV connection if at least 1

ISV posterior to the scored ISV made a normal connection.

Cell culture

696   HUVEC (C2519A, Lonza) were cultured in EBM-2 (CC-3162, Lonza)

supplemented with the Endothelial Growth Medium (EGM-2) bullet kit (CC-3162, Lonza)

and 1x antibiotic-antimycotic (Gibco). Normal human lung fibroblasts (CC2512, Lonza)

were cultured in DMEM (Gibco) with 10% fetal bovine serum (FBS) and 1x antibiotic-

antimycotic (Gibco). All cells were maintained at 37°C and 5% CO_2. For contractility

inhibition experiments, HUVEC were treated with 10µM (-) Blebbistatin (B0560-1MG,

Sigma) for 15 min at 37°C or 10µM Y-27632 (10187-694, VWR) for 30 min at 37°C then

immediately fixed in 4% PFA. For induction of contractility, HUVEC were treated with

0.5U/ml thrombin (T7201-500UN, Sigma) for 10 min at 37°C. For microtubule

depolymerization, HUVEC were treated with 10µM nocodazole (M1404, Sigma) for 20

min at 37°C then immediately fixed in methanol.

siRNA knockdown

709   HUVEC were transfected with non-targeting siRNA (NT, #4390847, Life

Technologies), SUN1 siRNA #1 (439240, #s23630, Life Technologies), SUN1 siRNA #2

(439240, #s23629, Life Technologies), GEF-H1 siRNA (439240, #s17546, Life

Technologies), and/or nesprin-1 siRNA (M-014039-02-0005, Dharmacon) using

Lipofectamine 2000 (11668027, Invitrogen) or Lipofectamine 3000 (L3000015,

ThermoFisher). siRNA at 0.48µM in Opti-MEM (31985-070, Gibco) and a 1:20 dilution of

Lipofectamine in Opti-MEM were incubated separately at RT for 5 min, then combined

and incubated at RT for 15 min. HUVEC were transfected at ~80% confluency with

siRNA at 37°C for 24h, then recovered in EGM-2 for an additional 24h. HUVEC were

seeded onto glass chamber slides coated with 5µg/ml fibronectin (F2006-2MG, Sigma)

for experiments.

RNA sequencing and analysis

RNA was extracted using TRIzol (15596018, Invitrogen) from 3 biological

replicates (independent experiments) of HUVEC under static or homeostatic laminar

flow (15d/cm^2, 72hr) conditions, and KAPA mRNA HyperPrep Kit (7961901001, Roche)

was used to prepare stranded libraries for sequencing (NovaSeq S1). 2–3 × 10^7

50-bp paired-end reads per sample were obtained and mapped to human genome

GRCh38 downloaded from https://support.10xgenomics.com/single-cell-gene-

expression/software/pipelines/latest/advanced/references with STAR using default

settings (Dobin et al., 2013). Mapping rate was over 80% for all samples, and gene

expression was determined with Htseq-count using the union mode

(https://htseq.readthedocs.io/en/master/) (Putri et al., 2022). Differential expression

analysis was performed with DESeq2 (Love et al., 2014) using default settings in R, and

lists of differentially expressed genes were obtained (p adjusted < 0.1).

Immunofluorescence staining

For experiments visualizing microtubules, HUVEC were fixed in ice-cold

methanol for 10 min at 4°C. For all other experiments, HUVEC were fixed with 4% PFA

for 10 min at RT and permeabilized with 0.1% Triton X-100 (T8787, Sigma) for 10 min

at RT. Fixed HUVEC were blocked for 1h at RT in blocking solution (5% FBS, 2X

antibiotic-antimycotic (Gibco), 0.1% sodium azide (s2002-100G, Sigma) in DPBS). Cells

were incubated in primary antibody overnight at 4°C, then washed 3X for 5 min in

DPBS. Secondary antibody and DRAQ7 (1:1000, ab109202, Abcam), DAPI (0.3μM,

10236276001, Sigma), and/or AlexaFluor488 Phalloidin (1:100, A12379, Life

Technologies) were added for 1h at RT followed by 3X washes for 10 min each in

DPBS. Slides were mounted with coverslips using Prolong Diamond Antifade mounting

medium (P36961, Life Technology) and sealed with nail polish. Primary and secondary

antibodies were diluted in blocking solution. The following primary antibodies were

used: anti-VE-cadherin (1:500, 2500S, Cell Signaling), anti-SUN1 (1:500, ab124770,

Abcam), anti-Ki67 (1:500, ab15580, Abcam), anti-phospho-myosin light chain 2

(Thr18/Ser19) (1:500, 3674S, Cell Signaling), anti-alpha-tubulin (1:500, 3873S, Cell

Signaling), anti-GEF-H1 (1:500, ab155785, Abcam), and anti-SYNE1 (1:500,

HPA019113, Atlas Antibodies). The following secondary antibodies from Life

Technologies were used: goat anti-mouse AlexaFluor 488 (1:500, A11029), goat anti-

rabbit AlexaFluor 594 (1:500, A11037), goat anti-mouse 647 (1:500, A21236), and goat

anti-rabbit 647 (1:500, A21245).

Western blotting

Cells were scraped into RIPA buffer with protease/phosphatase inhibitor (5872S,

Cell Signaling) then centrifuged at 13000 rpm at 4°C for 20 min. Lysate was reduced in

sample loading buffer and dithiothreitol (R0861, Thermo Fisher) and boiled for 10 min at

100°C. Samples were stored at -20°C until use. 10μg of sample were run on a 10%

stain-free polyacrylamide gel (161-0183, BioRad) then transferred onto a PVDF

membrane on ice for 1.5h. Membranes were blocked in OneBlock (20-313,

Prometheus) for 1 h at RT then washed 3X in PBST. Anti-GEF-H1 (1:1000, ab155785,

Abcam), anti-VE-cadherin (1:14,000, 2500S, Cell Signaling), or anti-GAPDH (1:5000,

97166S, Cell Signaling) was added overnight at 4°C. Membranes were washed 3X in

PBST then donkey anti-rabbit HRP secondary antibody (1:10,000, A16035, Thermo

Fisher) was added for 1h at RT. Immobilon Forte HRP Substrate (WBLUF0100,

Millipore Sigma) was added for 30 sec. Blots were exposed for 8 sec.

EdU labeling

HUVEC were labeled with EdU using the Click-It EdU Kit 488 (Invitrogen,

C10337) and fixed according to the manufacturer's instructions. Cells positive for EdU

labeling were counted and compared to total cell number to obtain percent positive.

Junction analysis

Endothelial cell adherens junctions were quantified in monolayers using Fiji to

generate 15µm line scans of VE-cadherin signal parallel to the cell junctions. VE-

cadherin signal was integrated to obtain the area under the curve. Linear junctions with

consistent VE-cadherin signal thus had a large area under the curve, while more

serrated junctions had reduced area under the curve **(Fig S3e)**. Measurements were

performed on at least 9 cells per field of view, with 3-6 fields of view per condition.

Real time cell analysis (RTCA)

Barrier function was assessed using the xCELLigence Real-Time Cell Analyzer

(RTCA, Acea Biosciences/Roche Applied Science) to measure electrical impedance

across HUVEC monolayers seeded onto microelectrodes. HUVEC were seeded to

confluency on the microelectrodes of the E-plate (E-plate 16, Roche Applied Science).

Electrical impedance readings were taken every 2 min for 5h. The percent change in

cell index was obtained at the 5h timepoint using the following formula: $(\text{Cell Index}_{SUN1}-$

$\text{Cell Index}_{NT})/\text{ABS}(\text{Cell Index}_{NT}).$

Flow experiments

Flow experiments were performed using an Ibidi pump system (10902, Ibidi) as

described (Ruter et al., 2021). HUVEC were seeded onto fibronectin coated Ibidi slides

(either μ-Slide I Luer I 0.4 mm (80176, Ibidi) or μ-Slide Y-shaped (80126, Ibidi) in flow

medium (EBM-2 with 10% FBS and 1x Antibiotic-antimycotic). HUVEC were exposed to

laminar shear stress for 30 min at 5 dyn/cm2, followed by 30 min at 10 dyn/cm2, and

finally for 72 h at 15 dyn/cm2. Alignment was measured by taking the ratio of the

longitudinal axis to the transverse axis relative to the flow vector. At least 10 cells were

measured per condition per experiment. Vascular permeability *in vitro* was determined

using the biotin matrix labeling assay as described below.

Biotin matrix labeling assay

Labeling of biotinylated matrix was assessed as described (Dubrovskyi et al.,

2013). Briefly, fibronectin was biotinylated by incubating 0.1mg/mL fibronectin with

0.5mM EZ-Link Sulfo-NHS-LC-Biotin (A39257, ThermoFisher) for 30 min at RT. Glass

chamber slides were coated with 5μg/ml biotinylated-fibronectin and HUVEC were

seeded on top. At confluency, HUVEC were treated with 25μg/mL Streptavidin-488

(S11223, Invitrogen) for 3 min then immediately fixed. For quantification, Fiji was used

to threshold the streptavidin signal, and the streptavidin area was measured and

normalized for total area for at least 3 fields of view per experiment.

Junction reformation assay

The EDTA junction reformation assay was performed as previously described

(Wright et al., 2015). Briefly, HUVEC were treated with 3mM EDTA (EDS-100G, Sigma-

Aldrich) for 1h at 37°C. EDTA was then washed out 3X with DPBS, incubation at 37°C

in EGM-2 was continued, and cells were fixed at 0 min, 20 min, 40 min, and 60 min

intervals.

VE-cadherin internalization

VE-cadherin internalization was performed as described (Wylie et al., 2018).

Briefly, HUVEC were plated on 5µg/ml fibronectin and grown to confluency. After

overnight serum starvation (Opti-MEM (31985-070, Gibco) supplemented with 1% FBS

(F2442, Sigma), and 1x antibiotic-antimycotic (Gibco)), cells were washed with pre-

chilled PBS+ (14040182, ThermoFisher) on ice at 4°C, then incubated in ice-cold

blocking solution (EBM-2 (CC-3162, Lonza) supplemented with 0.5% BSA (A4503,

Sigma)) for 30 min at 4°C. HUVEC were then incubated with VE-cadherin BV6 antibody

(1:100, ALX-803-305-C100, Enzo) in blocking solution for 2h on ice at 4°C. Following

VE-cadherin labeling, cells were washed with PBS+ then incubated in pre-warmed

internalization medium (EBM-2) at 37°C for 1h. Finally, HUVEC were incubated in acid

wash (0.5M NaCl/0.2M acetic acid) for 4 min at 4°C to remove remaining labeled VE-

cadherin on the cell surface, then washed with PBS+ and fixed. For quantification,

internalized VE-cadherin area was measured in Fiji, then normalized to total cell area

for at least 9 cells per experiment.

Microtubule analysis

For nuclear microtubule analysis, an ROI was drawn in Fiji over the nucleus. Fiji

was used to threshold α-tubulin signal, and the area of the signal was measured within

the ROI. This was performed on 8 cells per field of view, with 3-6 fields per condition per

experiment. For high-resolution microtubule analysis, high-resolution confocal images

were acquired with a Zeiss 880 Confocal with AiryScan FAST microscope with GaAsP

detector and camera (Zeiss) using a Plan-Apo 63x oil immersion objective (NA 1.40)

and 488nm and 561nm lasers. Imaging was performed at RT, and images were

acquired with the Zeiss 880 software and with the AiryScan detector. Images were then

processed with AiryScan. All image analysis, including Z-stack compression, was

performed in Fiji. For microtubule density, an ROI was drawn at the MTOC and the cell

periphery. Fiji was used to threshold α-tubulin signal, and the area of the signal was

measured within the ROI. This was performed on at least 10 cells per condition per

experiment. For junction analysis, the number of contacts between α-tubulin and VE-

cadherin were counted and normalized to the junction length. At least 15 junctions were

measured per condition per experiment.

Microtubule tip tracking

HUVEC were infected with an EB3-mCherry Lentivirus (Kushner et al., 2014) for

24h to visualize microtubule comets. Following infection, HUVEC were incubated with

siRNAs for NT or SUN1. For live imaging, cells were incubated at 37°C in a stage-top

incubator (TOKAI HIT, WSKM). Images were acquired on an Andor XD spinning disk

confocal microscope based on a CSU-X1 Yokogawa head with an Andor iXon 897 EM-

CCD camera. A 561nm laser and FF01-607/36 emission filter were used. Images were

acquired with a 470ms exposure and 32ms readout time for 2 min (240 frames) using a

UPlanSApo 60x silicone-oil immersion objective (NA 1.30) and Metamorph software.

Microtubules at the cell periphery were tracked using the Manual Tracking plugin in Fiji

and were tracked for at least 60 frames (30 sec). Track information was acquired from

the x and y coordinates using a custom algorithm in Visual Basic in Excel provided by

Dan Buster at the University of Arizona. 10 microtubule tracks were measured per cell.

GEF-H1 analysis

For GEF-H1 localization analysis, an ROI was drawn at the periphery of the cell,

and the α-tubulin signal was used to create a mask. Mean signal intensity was then

measured for GEF-H1 within the α-tubulin mask and outside of it, and a ratio was taken.

At least 9 cells were analyzed per condition per experiment.

Statistics

Student's two-tailed unpaired *t*-test was used to determine statistical significance

in experiments with 2 groups and one-way ANOVA with Tukey's multiple comparisons

test was used in experiments with 3 groups. For thrombin, blebbistatin, Y-27632, and

nocodazole experiments, two-way ANOVA with Tukey's multiple comparisons test was

used to determine statistical significance. X^2 was used for categorical data. For box and

whisker plots, boxes represent the upper quartile, lower quartile, and median; whiskers

represent the minimum and maximum values. Statistical tests and graphs were made

using the Prism 9 software (GraphPad Software).

Data availability

The RNA-seq data that support the findings of this study are available in the

Gene Expression Omnibus (GEO) under the accession number GSE213099.

ACKNOWLEGMENTS

We thank Michelle Altemara and staff at the Zebrafish Aquaculture Core and

Kaitlyn Quigley for fish room support, Aaron Friedman and Caroline Crater for mouse

room support, and Yosuke Mukouyama, Nick Buglak, and Bautch Lab members for

critical discussion and feedback. We also thank Dan Buster for the software used in

microtubule tip tracking experiments and Angelika Noegel for sharing RNASeq data with

us. Airy Scan imaging was performed at the UNC Hooker Imaging Core Facility and

microtubule tip tracking imaging was performed with Pablo Ariel at the UNC Microscopy

Services Laboratory, supported in part by P30 CA016086 Cancer Center Core Support

Grant to the UNC Lineberger Comprehensive Cancer Center.

AUTHOR CONTRIBUTIONS

FUNDING

This work was supported by grants from the National Institutes of Health (R35

HL139950 to VLB), the Integrated Vascular Biology Training Grant (5T32HL069768-17,

DECLARATION OF INTERESTS: None.

REFERENCES

Abu Taha, A., & Schnittler, H.-J. (2014). Dynamics between actin and the VE-
cadherin/catenin complex: novel aspects of the ARP2/3 complex in regulation of
endothelial junctions. Cell Adhesion & Migration, *8*(2), 125–135.
https://doi.org/10.4161/cam.28243

Angulo-Urarte, A., Casado, P., Castillo, S. D., Kobialka, P., Kotini, M. P., Figueiredo, A.
M., Castel, P., Rajeeve, V., Milà-Guasch, M., Millan, J., et al. (2018). Endothelial
cell rearrangements during vascular patterning require PI3-kinase-mediated
inhibition of actomyosin contractility. Nature Communications, *9*(1), 4826.
https://doi.org/10.1038/s41467-018-07172-3

Anno, T., Sakamoto, N., & Sato, M. (2012). Role of nesprin-1 in nuclear deformation in
endothelial cells under static and uniaxial stretching conditions. Biochemical and
Biophysical Research Communications, *424*(1), 94–99.
https://doi.org/10.1016/j.bbrc.2012.06.073

Arganda-Carreras, I., Fernández-González, R., Muñoz-Barrutia, A., & Ortiz-De-
Solorzano, C. (2010). 3D reconstruction of histological sections: Application to
mammary gland tissue. Microscopy Research and Technique, *73*(11), 1019–1029.
https://doi.org/10.1002/jemt.20829

Banerjee, I., Zhang, J., Moore-Morris, T., Pfeiffer, E., Buchholz, K. S., Liu, A., Ouyang,
K., Stroud, M. J., Gerace, L., Evans, S. M., et al. (2014). Targeted ablation of
nesprin 1 and nesprin 2 from murine myocardium results in cardiomyopathy,
altered nuclear morphology and inhibition of the biomechanical gene response.
PLoS Genetics, *10*(2), e1004114. https://doi.org/10.1371/journal.pgen.1004114

Bautch, V. L., & Caron, K. M. (2015). Blood and lymphatic vessel formation. Cold Spring
Harbor Perspectives in Biology, *7*(3), a008268.
https://doi.org/10.1101/cshperspect.a008268

Bellett, G., Carter, J. M., Keynton, J., Goldspink, D., James, C., Moss, D. K., &
Mogensen, M. M. (2009). Microtubule plus-end and minus-end capture at adherens
junctions is involved in the assembly of apico-basal arrays in polarised epithelial
cells. Cell Motility and the Cytoskeleton, *66*(10), 893–908.
https://doi.org/10.1002/cm.20393

Bentley, K., Franco, C. A., Philippides, A., Blanco, R., Dierkes, M., Gebala, V., Stanchi,
F., Jones, M., Aspalter, I. M., Cagna, G., et al. (2014). The role of differential VE-
cadherin dynamics in cell rearrangement during angiogenesis. Nature Cell Biology,
16(4), 309–321.

Birkenfeld, J., Nalbant, P., Yoon, S.-H., & Bokoch, G. M. (2008). Cellular functions of
GEF-H1, a microtubule-regulated Rho-GEF: is altered GEF-H1 activity a crucial
determinant of disease pathogenesis? Trends in Cell Biology, *18*(5), 210–219.
https://doi.org/10.1016/j.tcb.2008.02.006

Birukova, A. A., Adyshev, D., Gorshkov, B., Bokoch, G. M., Birukov, K. G., & Verin, A.
D. (2006). GEF-H1 is involved in agonist-induced human pulmonary endothelial
barrier dysfunction. American Journal of Physiology. Lung Cellular and Molecular
Physiology, *290*(3), L540-8. https://doi.org/10.1152/ajplung.00259.2005

Birukova, A. A., Liu, F., Garcia, J. G. N., & Verin, A. D. (2004). Protein kinase A
attenuates endothelial cell barrier dysfunction induced by microtubule disassembly.
American Journal of Physiology. Lung Cellular and Molecular Physiology, *287*(1),
L86-93. https://doi.org/10.1152/ajplung.00441.2003

Blum, Y., Belting, H.-G., Ellertsdottir, E., Herwig, L., Lüders, F., & Affolter, M. (2008).
Complex cell rearrangements during intersegmental vessel sprouting and vessel
fusion in the zebrafish embryo. Developmental Biology, *316*(2), 312–322.
https://doi.org/10.1016/j.ydbio.2008.01.038

Buglak, D. B., Kushner, E. J., Marvin, A. P., Davis, K. L., & Bautch, V. L. (2020). Excess
centrosomes disrupt vascular lumenization and endothelial cell adherens junctions.
Angiogenesis, *23*(4), 567–575. https://doi.org/10.1007/s10456-020-09737-7

Capell, B. C., & Collins, F. S. (2006). Human laminopathies: nuclei gone genetically
awry. Nature Reviews. Genetics, *7*(12), 940–952. https://doi.org/10.1038/nrg1906

Carley, E., King, M. C., & Guo, S. (2022). Integrating mechanical signals into cellular
identity. Trends in Cell Biology, *32*(8), 669–680.
https://doi.org/10.1016/j.tcb.2022.02.006

Carley, E., Stewart, R. M., Zieman, A., Jalilian, I., King, D. E., Zubek, A., Lin, S.,
Horsley, V., & King, M. C. (2021). The LINC complex transmits integrin-dependent
tension to the nuclear lamina and represses epidermal differentiation. ELife, *10*.
https://doi.org/10.7554/eLife.58541

Carmeliet, P., & Jain, R. K. (2011). Principles and mechanisms of vessel normalization
for cancer and other angiogenic diseases. Nature Reviews Drug Discovery, *10*,
417–427.

Carmeliet, P., Lampugnani, M. G., Moons, L., Breviario, F., Compernolle, V., Bono, F.,
Balconi, G., Spagnuolo, R., Oosthuyse, B., Dewerchin, M., et al. (1999). Targeted
deficiency or cytosolic truncation of the VE-cadherin gene in mice impairs VEGF-
mediated endothelial survival and angiogenesis. Cell, *98*(147–157).

Chancellor, T. J., Lee, J., Thodeti, C. K., & Lele, T. (2010). Actomyosin tension exerted
on the nucleus through nesprin-1 connections influences endothelial cell adhesion,
migration, and cyclic strain-induced reorientation. Biophysical Journal, *99*(1), 115–
123. https://doi.org/10.1016/j.bpj.2010.04.011

Chang, W., Wang, Y., Luxton, G. W. G., Östlund, C., Worman, H. J., & Gundersen, G.
G. (2019). Imbalanced nucleocytoskeletal connections create common polarity
defects in progeria and physiological aging. Proceedings of the National Academy
of Sciences of the United States of America, *116*(9), 3578–3583.
https://doi.org/10.1073/pnas.1809683116

Chen, C.-Y., Chi, Y.-H., Mutalif, R. A., Starost, M. F., Myers, T. G., Anderson, S. A.,
Stewart, C. L., & Jeang, K.-T. (2012). Accumulation of the Inner Nuclear Envelope
Protein Sun1 Is Pathogenic in Progeric and Dystrophic Laminopathies. Cell, *149*(3),
565–577. https://doi.org/10.1016/j.cell.2012.01.059

Chen, Z.-J., Wang, W.-P., Chen, Y.-C., Wang, J.-Y., Lin, W.-H., Tai, L.-A., Liou, G.-G.,
Yang, C.-S., & Chi, Y.-H. (2014). Dysregulated interactions between lamin A and
SUN1 induce abnormalities in the nuclear envelope and endoplasmic reticulum in
progeric laminopathies. Journal of Cell Science, *127*(Pt 8), 1792–1804.
https://doi.org/10.1242/jcs.139683

Chong, D. C., Yu, Z., Brighton, H. E., Bear, J. E., & Bautch, V. L. (2017). Tortuous
Microvessels Contribute to Wound Healing via Sprouting Angiogenesis.
Arteriosclerosis, Thrombosis, and Vascular Biology, *37*(10), 1903–1912.
https://doi.org/10.1161/ATVBAHA.117.309993

Claesson-Welsh, L. (2015). Vascular permeability--the essentials. Upsala Journal of
Medical Sciences, *120*(3), 135–143.
https://doi.org/10.3109/03009734.2015.1064501

Claesson-Welsh, L., Dejana, E., & McDonald, D. M. (2021). Permeability of the
Endothelial Barrier: Identifying and Reconciling Controversies. Trends in Molecular
Medicine, *27*(4), 314–331. https://doi.org/10.1016/j.molmed.2020.11.006

Conway, D. E., Williams, M. R., Eskin, S. G., & McIntire, L. V. (2010). Endothelial cell
responses to atheroprone flow are driven by two separate flow components: low
time-average shear stress and fluid flow reversal. American Journal of Physiology.
Heart and Circulatory Physiology, *298*(2), H367-74.
https://doi.org/10.1152/ajpheart.00565.2009

De Sandre-Giovannoli, A. (2003). Lamin A Truncation in Hutchinson-Gilford Progeria.
Science, *300*(5628), 2055–2055. https://doi.org/10.1126/science.1084125

Dejana, E. (2004). Endothelial cell-cell junctions: happy together. Nature Reviews.
Molecular Cell Biology, *5*(4), 261–270. https://doi.org/10.1038/nrm1357

Dejana, E., & Vestweber, D. (2013). The role of VE-cadherin in vascular morphogenesis
and permeability control. Progress in Molecular Biology and Translational Science,
116, 119–144. https://doi.org/10.1016/B978-0-12-394311-8.00006-6

DeLisser, H. M. (2011). Modulators of endothelial cell filopodia: PECAM-1 joins the club.
Cell Adhesion & Migration, *5*(1), 37–41. https://doi.org/10.4161/cam.5.1.13575

Denis, K. B., Cabe, J. I., Danielsson, B. E., Tieu, K. V, Mayer, C. R., & Conway, D. E.
(2021). The LINC complex is required for endothelial cell adhesion and adaptation
to shear stress and cyclic stretch. Molecular Biology of the Cell, mbcE20110698.
https://doi.org/10.1091/mbc.E20-11-0698

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., & Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner.

Bioinformatics (Oxford, England), *29*(1), 15–21.
https://doi.org/10.1093/bioinformatics/bts635

Dubrovskyi, O., Birukova, A. A., & Birukov, K. G. (2013). Measurement of local permeability at subcellular level in cell models of agonist- and ventilator-induced lung injury. Laboratory Investigation; a Journal of Technical Methods and Pathology, *93*(2), 254–263. https://doi.org/10.1038/labinvest.2012.159

Eriksson, M., Brown, W. T., Gordon, L. B., Glynn, M. W., Singer, J., Scott, L., Erdos, M. R., Robbins, C. M., Moses, T. Y., Berglund, P., et al. (2003). Recurrent de novo point mutations in lamin A cause Hutchinson-Gilford progeria syndrome. Nature, *423*(6937), 293–298. https://doi.org/10.1038/nature01629

Esser, S., Lampugnani, M. G., Corada, M., Dejana, E., & Risau, W. (1998). Vascular endothelial growth factor induces VE-cadherin tyrosine phosphorylation in endothelial cells. Journal of Cell Science, *111 (Pt 1*, 1853–1865. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/9625748

Fridolfsson, H. N., Ly, N., Meyerzon, M., & Starr, D. A. (2010). UNC-83 coordinates kinesin-1 and dynein activities at the nuclear envelope during nuclear migration. Developmental Biology, *338*(2), 237–250. https://doi.org/10.1016/j.ydbio.2009.12.004

Gimpel, P., Lee, Y. L., Sobota, R. M., Calvi, A., Koullourou, V., Patel, R., Mamchaoui, K., Nédélec, F., Shackleton, S., Schmoranzer, J., et al. (2017). Nesprin-1α-Dependent Microtubule Nucleation from the Nuclear Envelope via Akap450 Is Necessary for Nuclear Positioning in Muscle Cells. Current Biology : CB, *27*(19), 2999-3009.e9. https://doi.org/10.1016/j.cub.2017.08.031

Graham, D. M., Andersen, T., Sharek, L., Uzer, G., Rotherberg, K., Hoffman, B. D., Rubin, J., Balland, M., Bear, J. E., & Burridge, K. (2018). Enucleated cells reveal differential roles of the nucleus in cell migration, polarity, and mechanotransduction. Journal of Cell Biology, jcb.201706097.

Guilluy, C., Osborne, L. D., Van Landeghem, L., Sharek, L., Superfine, R., Garcia-Mata, R., & Burridge, K. (2014). Isolated nuclei adapt to force and reveal a mechanotransduction pathway in the nucleus. Nature Cell Biology, *16*(4), 376–381. https://doi.org/10.1038/ncb2927

Han, Y., Wang, L., Yao, Q.-P., Zhang, P., Liu, B., Wang, G.-L., Shen, B.-R., Cheng, B., Wang, Y., Jiang, Z.-L., et al. (2015). Nuclear envelope proteins Nesprin2 and LaminA regulate proliferation and apoptosis of vascular endothelial cells in response to shear stress. Biochimica et Biophysica Acta, *1853*(5), 1165–1173. https://doi.org/10.1016/j.bbamcr.2015.02.013

Haque, F., Lloyd, D. J., Smallwood, D. T., Dent, C. L., Shanahan, C. M., Fry, A. M., Trembath, R. C., & Shackleton, S. (2006). SUN1 interacts with nuclear lamin A and cytoplasmic nesprins to provide a physical connection between the nuclear lamina and the cytoskeleton. Molecular and Cellular Biology, *26*(10), 3738–3751.

Haque, F., Mazzeo, D., Patel, J. T., Smallwood, D. T., Ellis, J. A., Shanahan, C. M., &
Shackleton, S. (2010). Mammalian SUN protein interaction networks at the inner
nuclear membrane and their role in laminopathy disease processes. The Journal of
Biological Chemistry, *285*(5), 3487–3498. https://doi.org/10.1074/jbc.M109.071910

Hellström, M., Phng, L.-K., Hofmann, J. J., Wallgard, E., Coultas, L., Lindblom, P., Alva,
J., Nilsson, A.-K., Karlsson, L., Gaiano, N., et al. (2007). Dll4 signalling through
Notch1 regulates formation of tip cells during angiogenesis. Nature, *445*(7129),
776–780. https://doi.org/10.1038/nature05571

Huang, C., Sheikh, F., Hollander, M., Cai, C., Becker, D., Chu, P.-H., Evans, S., &
Chen, J. (2003). Embryonic atrial function is essential for mouse embryogenesis,
cardiac morphogenesis and angiogenesis. Development (Cambridge, England),
130(24), 6111–6119. https://doi.org/10.1242/dev.00831

Hübner, K., Cabochette, P., Diéguez-Hurtado, R., Wiesner, C., Wakayama, Y.,
Grassme, K. S., Hubert, M., Guenther, S., Belting, H.-G., Affolter, M., et al. (2018).
Wnt/β-catenin signaling regulates VE-cadherin-mediated anastomosis of brain
capillaries by counteracting S1pr1 signaling. Nature Communications, *9*(1), 4860.
https://doi.org/10.1038/s41467-018-07302-x

Huveneers, S., Oldenburg, J., Spanjaard, E., van der Krogt, G., Grigoriev, I.,
Akhmanova, A., Rehmann, H., & de Rooij, J. (2012). Vinculin associates with
endothelial VE-cadherin junctions to control force-dependent remodeling. The
Journal of Cell Biology, *196*(5), 641–652. https://doi.org/10.1083/jcb.201108120

Juettner, V. V, Kruse, K., Dan, A., Vu, V. H., Khan, Y., Le, J., Leckband, D., Komarova,
Y., & Malik, A. B. (2019). VE-PTP stabilizes VE-cadherin junctions and the
endothelial barrier via a phosphatase-independent mechanism. The Journal of Cell
Biology, *218*(5), 1725–1742. https://doi.org/10.1083/jcb.201807210

Ketema, M., Wilhelmsen, K., Kuikman, I., Janssen, H., Hodzic, D., & Sonnenberg, A.
(2007). Requirements for the localization of nesprin-3 at the nuclear envelope and
its interaction with plectin. Journal of Cell Science, *120*(Pt 19), 3384–3394.
https://doi.org/10.1242/jcs.014191

King, S. J., Nowak, K., Suryavanshi, N., Holt, I., Shanahan, C. M., & Ridley, A. J.
(2014). Nesprin-1 and nesprin-2 regulate endothelial cell shape and migration.
Cytoskeleton (Hoboken, N.J.), *71*(7), 423–434. https://doi.org/10.1002/cm.21182

Komarova, Y. A., Huang, F., Geyer, M., Daneshjou, N., Garcia, A., Idalino, L., Kreutz,
B., Mehta, D., & Malik, A. B. (2012). VE-cadherin signaling induces EB3
phosphorylation to suppress microtubule growth and assemble adherens junctions.
Molecular Cell, *48*(6), 914–925.

Komarova, Y., Lansbergen, G., Galjart, N., Grosveld, F., Borisy, G. G., & Akhmanova,
A. (2005). EB1 and EB3 control CLIP dissociation from the ends of growing
microtubules. Molecular Biology of the Cell, *16*(11), 5334–5345.
https://doi.org/10.1091/mbc.e05-07-0614

Krendel, M., Zenke, F. T., & Bokoch, G. M. (2002). Nucleotide exchange factor GEF-H1
mediates cross-talk between microtubules and the actin cytoskeleton. Nature Cell
Biology, *4*(4), 294–301. https://doi.org/10.1038/ncb773

Kushner, E. J., & Bautch, V. L. (2013). Building blood vessels in development and
disease. Current Opinion in Hematology, *20*(3), 231–236.
https://doi.org/10.1097/MOH.0b013e328360614b

Kushner, E. J., Ferro, L. S., Liu, J.-Y., Durrant, J. R., Rogers, S. L., Dudley, A. C., &
Bautch, V. L. (2014). Excess centrosomes disrupt endothelial migration via
centrosome scattering. Journal of Cell Biology, *206*(2), 257–272.

Kushner, E. J., Ferro, L. S., Yu, Z., & Bautch, V. L. (2016). Excess centrosomes perturb
dynamic endothelial cell repolarization during blood vessel formation. Molecular
Biology of the Cell, *27*(12), 1911–1920.

Lei, K., Zhang, X., Ding, X., Guo, X., Chen, M., Zhu, B., Xu, T., Zhuang, Y., Xu, R., &
Han, M. (2009). SUN1 and SUN2 play critical but partially redundant roles in
anchoring nuclei in skeletal muscle cells in mice. PNAS, *106*(25), 10207–10212.

Li, P., Stumpf, M., Müller, R., Eichinger, L., Glöckner, G., & Noegel, A. A. (2017). The
function of the inner nuclear envelope protein SUN1 in mRNA export is regulated
by phosphorylation. Scientific Reports, *7*(1), 9157. https://doi.org/10.1038/s41598-
017-08837-7

Ligon, L. A., Karki, S., Tokito, M., & Al, E. (2001). Dynein binds to beta-catenin and may
tether microtubules at adherens junctions. Nature Cell Biology, *3*, 913–917.

Linkert, M., Rueden, C. T., Allan, C., Burel, J.-M., Moore, W., Patterson, A., Loranger,
B., Moore, J., Neves, C., MacDonald, D., et al. (2010). Metadata matters: access to
image data in the real world. Journal of Cell Biology, *189*(5), 777–782.
https://doi.org/10.1083/jcb.201004104

Liu, Z., Ruter, D. L., Quigley, K., Tanke, N. T., Jiang, Y., & Bautch, V. L. (2021). Single-
Cell RNA Sequencing Reveals Endothelial Cell Transcriptome Heterogeneity Under
Homeostatic Laminar Flow. Arteriosclerosis, Thrombosis, and Vascular Biology,
41(10), 2575–2584. https://doi.org/10.1161/ATVBAHA.121.316797

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biology, *15*(12), 550.
https://doi.org/10.1186/s13059-014-0550-8

Maurya, M. R., Gupta, S., Li, J. Y.-S., Ajami, N. E., Chen, Z. B., Shyy, J. Y.-J., Chien,
S., & Subramaniam, S. (2021). Longitudinal shear stress response in human
endothelial cells to atheroprone and atheroprotective conditions. Proceedings of the
National Academy of Sciences of the United States of America, *118*(4).
https://doi.org/10.1073/pnas.2023236118

Mavria, G., Vercoulen, Y., Yeo, M., Paterson, H., Karasarides, M., Marais, R., Bird, D.,
& Marshall, C. J. (2006). ERK-MAPK signaling opposes Rho-kinase to promote

endothelial cell survival and sprouting during angiogenesis. Cancer Cell, *9*(1), 33–
44. https://doi.org/10.1016/j.ccr.2005.12.021

May, C. K., & Carroll, C. W. (2018). Differential incorporation of SUN-domain proteins
into LINC complexes is coupled to gene expression. PloS One, *13*(5), e0197621.
https://doi.org/10.1371/journal.pone.0197621

McGee, M. D., Rillo, R., Anderson, A. S., & Starr, D. A. (2006). UNC-83 IS a KASH
protein required for nuclear migration and is recruited to the outer nuclear
membrane by a physical interaction with the SUN protein UNC-84. Molecular
Biology of the Cell, *17*(4), 1790–1801. https://doi.org/10.1091/mbc.e05-09-0894

Meng, W., Mushika, Y., Ichii, T., & Takeichi, M. (2008). Anchorage of microtubule minus
ends to adherens junctions regulates epithelial cell-cell contacts. Cell, *135*(5), 948–
959. https://doi.org/10.1016/j.cell.2008.09.040

Meyerzon, M., Fridolfsson, H. N., Ly, N., McNally, F. J., & Starr, D. A. (2009). UNC-83 is
a nuclear-specific cargo adaptor for kinesin-1-mediated nuclear migration.
Development (Cambridge, England), *136*(16), 2725–2733.
https://doi.org/10.1242/dev.038596

Montero-Balaguer, M., Swirsding, K., Orsenigo, F., Cotelli, F., Mione, M., & Dejana, E.
(2009). Stable vascular connections and remodeling require full expression of VE-
cadherin in zebrafish embryos. PloS One, *4*(6), e5772.
https://doi.org/10.1371/journal.pone.0005772

Morgan, J. T., Pfeiffer, E. R., Thirkill, T. L., Kumar, P., Peng, G., Fridolfsson, H. N.,
Douglas, G. C., Starr, D. A., & Barakat, A. I. (2011). Nesprin-3 regulates endothelial
cell morphology, perinuclear cytoskeletal architecture, and flow-induced
polarization. Molecular Biology of the Cell, *22*(22), 4324–4334.
https://doi.org/10.1091/mbc.E11-04-0287

Nakatsu, M. N., & Hughes, C. C. W. (2008). An optimized three-dimensional in vitro
model for the analysis of angiogenesis. Methods in Enzymology, *443*, 65–82.

Nesmith, J. E., Chappell, J. C., Cluceru, J. G., & Bautch, V. L. (2017). Blood vessel
anastomosis is spatially regulated by Flt1 during angiogenesis. Development
(Cambridge, England), *144*(5), 889–896. https://doi.org/10.1242/dev.145672

Neto, F., Klaus-Bergmann, A., Ong, Y. T., Alt, S., Vion, A.-C., Szymborska, A.,
Carvalho, J. R., Hollfinger, I., Bartels-Klein, E., Franco, C. A., et al. (2018). YAP
and TAZ regulate adherens junction dynamics and endothelial cell distribution
during vascular development. ELife, *7*. https://doi.org/10.7554/eLife.31037

Olive, M., Harten, I., Mitchell, R., Beers, J. K., Djabali, K., Cao, K., Erdos, M. R., Blair,
C., Funke, B., Smoot, L., et al. (2010). Cardiovascular pathology in Hutchinson-
Gilford progeria: correlation with the vascular pathology of aging. Arteriosclerosis,
Thrombosis, and Vascular Biology, *30*(11), 2301–2309.
https://doi.org/10.1161/ATVBAHA.110.209460

Osmanagic-Myers, S., Kiss, A., Manakanatas, C., Hamza, O., Sedlmayer, F., Szabo, P.
L., Fischer, I., Fichtinger, P., Podesser, B. K., Eriksson, M., et al. (2019).
Endothelial progerin expression causes cardiovascular pathology through an
impaired mechanoresponse. The Journal of Clinical Investigation, *129*(2), 531–545.
https://doi.org/10.1172/JCI121297

Ostlund, C., Folker, E. S., Choi, J. C., Gomes, E. R., Gundersen, G. G., & Worman, H.
J. (2009). Dynamics and molecular interactions of linker of nucleoskeleton and
cytoskeleton (LINC) complex proteins. Journal of Cell Science, *122*(Pt 22), 4099–
4108. https://doi.org/10.1242/jcs.057075

Padmakumar, V. C., Libotte, T., Lu, W., Zaim, H., Abraham, S., Noegel, A. A.,
Gotzmann, J., Foisner, R., & Karakesisoglou, I. (2005). The inner nuclear
membrane protein Sun1 mediates the anchorage of Nesprin-2 to the nuclear
envelope. Journal of Cell Science, *118*(Pt 15), 3419–3430.
https://doi.org/10.1242/jcs.02471

Putri, G. H., Anders, S., Pyl, P. T., Pimanda, J. E., & Zanini, F. (2022). Analysing high-
throughput sequencing data in Python with HTSeq 2.0. Bioinformatics (Oxford,
England). https://doi.org/10.1093/bioinformatics/btac166

Rauzi, M., Lenne, P.-F., & Lecuit, T. (2010). Planar polarized actomyosin contractile
flows control epithelial junction remodelling. Nature, *468*(7327), 1110–1114.
https://doi.org/10.1038/nature09566

Rho, S.-S., Ando, K., & Fukuhara, S. (2017). Dynamic Regulation of Vascular
Permeability by Vascular Endothelial Cadherin-Mediated Endothelial Cell-Cell
Junctions. Journal of Nippon Medical School = Nippon Ika Daigaku Zasshi, *84*(4),
148–159. https://doi.org/10.1272/jnms.84.148

Ricard, N., Ciais, D., Levet, S., Subileau, M., Mallet, C., Zimmers, T. A., Lee, S.-J.,
Bidart, M., Feige, J.-J., & Bailly, S. (2012). BMP9 and BMP10 are critical for
postnatal retinal vascular remodeling. Blood, *119*(25), 6162–6171.
https://doi.org/10.1182/blood-2012-01-407593

Ruter, D. L., Liu, Z., Ngo, K. M., X, S., Marvin, A., Buglak, D. B., Kidder, E. J., & Bautch,
V. L. (2021). SMAD6 transduces endothelial cell flow responses required for blood
vessel homeostasis. Angiogenesis, *24*(2), 387–398.
https://doi.org/10.1007/s10456-021-09777-7

Sauteur, L., Affolter, M., & Belting, H.-G. (2017). Distinct and redundant functions of
Esam and VE-cadherin during vascular morphogenesis. Development.
https://doi.org/10.1242/dev.140038

Sauteur, L., Krudewig, A., Herwig, L., Ehrenfeuchter, N., Lenard, A., Affolter, M., &
Belting, H.-G. (2014). Cdh5/VE-cadherin Promotes Endothelial Cell Interface
Elongation via Cortical Actin Polymerization during Angiogenic Sprouting. Cell
Reports, *9*(2), 504–513. https://doi.org/10.1016/j.celrep.2014.09.024

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-
source platform for biological-image analysis. Nature Methods, *9*(7), 676–682.
https://doi.org/10.1038/nmeth.2019

Seebach, J., Dieterich, P., Luo, F., Schillers, H., Vestweber, D., Oberleithner, H., Galla,
H. J., & Schnittler, H. J. (2000). Endothelial barrier function under laminar fluid
shear stress. Laboratory Investigation; a Journal of Technical Methods and
Pathology, *80*(12), 1819–1831. https://doi.org/10.1038/labinvest.3780193

Sehrawat, S., Cullere, X., Patel, S., Italiano, J., & Mayadas, T. N. (2008). Role of Epac1,
an exchange factor for Rap GTPases, in endothelial microtubule dynamics and
barrier function. Molecular Biology of the Cell, *19*(3), 1261–1270.
https://doi.org/10.1091/mbc.e06-10-0972

Sehrawat, S., Ernandez, T., Cullere, X., Takahashi, M., Ono, Y., Komarova, Y., &
Mayadas, T. N. (2011). AKAP9 regulation of microtubule dynamics promotes
Epac1-induced endothelial barrier properties. Blood, *117*(2), 708–718.
https://doi.org/10.1182/blood-2010-02-268870

Shaw, R. M., Fay, A. J., Puthenveedu, M. A., von Zastrow, M., Jan, Y.-N., & Jan, L. Y.
(2007). Microtubule plus-end-tracking proteins target gap junctions directly from the
cell interior to adherens junctions. Cell, *128*(3), 547–560.
https://doi.org/10.1016/j.cell.2006.12.037

Sörensen, I., Adams, R. H., & Gossler, A. (2009). DLL1-mediated Notch activation
regulates endothelial identity in mouse fetal arteries. Blood, *113*(22), 5680–5688.
https://doi.org/10.1182/blood-2008-08-174508

Starr, D. A., & Fridolfsson, H. N. (2010). Interactions between nuclei and the
cytoskeleton are mediated by SUN-KASH nuclear-envelope bridges. Annual
Review of Cell and Developmental Biology, *26*, 421–444.
https://doi.org/10.1146/annurev-cellbio-100109-104037

Stehbens, S. J., Paterson, A. D., Crampton, M. S., Shewan, A. M., Ferguson, C.,
Akhmanova, A., Parton, R. G., & Yap, A. S. (2006). Dynamic microtubules regulate
the local concentration of E-cadherin at cell-cell contacts. Journal of Cell Science,
119(Pt 9), 1801–1811. https://doi.org/10.1242/jcs.02903

Stewart-Hutchinson, P. J., Hale, C. M., Wirtz, D., & Hodzic, D. (2008). Structural
requirements for the assembly of LINC complexes and their function in cellular
mechanical stiffness. Experimental Cell Research, *314*(8), 1892–1905.
https://doi.org/10.1016/j.yexcr.2008.02.022

Stewart, R. M., Zubek, A. E., Rosowski, K. A., Schreiner, S. M., Horsley, V., & King, M.
C. (2015). Nuclear-cytoskeletal linkages facilitate cross talk between the nucleus
and intercellular adhesions. The Journal of Cell Biology, *209*(3), 403–418.
https://doi.org/10.1083/jcb.201502024

Stroud, M. J., Feng, W., Zhang, J., Veevers, J., Fang, X., Gerace, L., & Chen, J. (2017).
Nesprin 1α2 is essential for mouse postnatal viability and nuclear positioning in

skeletal muscle. The Journal of Cell Biology, *216*(7), 1915–1924. https://doi.org/10.1083/jcb.201612128

Szymborska, A., & Gerhardt, H. (2018). Hold Me, but Not Too Tight-Endothelial Cell-Cell Junctions in Angiogenesis. Cold Spring Harbor Perspectives in Biology, *10*(8). https://doi.org/10.1101/cshperspect.a029223

Taranum, S., Sur, I., Müller, R., Lu, W., Rashmi, R. N., Munck, M., Neumann, S., Karakesisoglou, I., & Noegel, A. A. (2012). Cytoskeletal interactions at the nuclear envelope mediated by nesprins. International Journal of Cell Biology, *2012*, 736524. https://doi.org/10.1155/2012/736524

Thakar, K., May, C. K., Rogers, A., & Carroll, C. W. (2017). Opposing roles for distinct LINC complexes in regulation of the small GTPase RhoA. Molecular Biology of the Cell, *28*(1), 182–191. https://doi.org/10.1091/mbc.E16-06-0467

Tkachenko, E., Gutierrez, E., Saikin, S. K., Fogelstrand, P., Kim, C., Groisman, A., & Ginsberg, M. H. (2013). The nucleus of endothelial cell as a sensor of blood flow direction. Biology Open, *2*(10), 1007–1012. https://doi.org/10.1242/bio.20134622

Ueda, N., Maekawa, M., Matsui, T. S., Deguchi, S., Takata, T., Katahira, J., Higashiyama, S., & Hieda, M. (2022). Inner Nuclear Membrane Protein, SUN1, is Required for Cytoskeletal Force Generation and Focal Adhesion Maturation. Frontiers in Cell and Developmental Biology, *10*, 885859. https://doi.org/10.3389/fcell.2022.885859

van Ingen, M. J. A., & Kirby, T. J. (2021). LINCing Nuclear Mechanobiology With Skeletal Muscle Mass and Function. Frontiers in Cell and Developmental Biology, *9*, 690577. https://doi.org/10.3389/fcell.2021.690577

Vasileva, E., & Citi, S. (2018). The role of microtubules in the regulation of epithelial junctions. Tissue Barriers, *6*(3), 1539596. https://doi.org/10.1080/21688370.2018.1539596

Verin, A. D., Birukova, A., Wang, P., Liu, F., Becker, P., Birukov, K., & Garcia, J. G. (2001). Microtubule disassembly increases endothelial cell barrier dysfunction: role of MLC phosphorylation. American Journal of Physiology. Lung Cellular and Molecular Physiology, *281*(3), L565-74. https://doi.org/10.1152/ajplung.2001.281.3.L565

Vion, A.-C., Perovic, T., Petit, C., Hollfinger, I., Bartels-Klein, E., Frampton, E., Gordon, E., Claesson-Welsh, L., & Gerhardt, H. (2020). Endothelial Cell Orientation and Polarity Are Controlled by Shear Stress and VEGF Through Distinct Signaling Pathways. Frontiers in Physiology, *11*, 623769. https://doi.org/10.3389/fphys.2020.623769

Wacker, A., & Gerhardt, H. (2011). Endothelial development taking shape. Current Opinion in Cell Biology, *23*(6), 676–685. https://doi.org/10.1016/j.ceb.2011.10.002

Wimmer, R., Cseh, B., Maier, B., Scherrer, K., & Baccarini, M. (2012). Angiogenic

sprouting requires the fine tuning of endothelial cell cohesion by the Raf-1/Rok-α
complex. Developmental Cell, *22*(1), 158–171.
https://doi.org/10.1016/j.devcel.2011.11.012

Wright, C. E., Kushner, E. J., Du, Q., & Bautch, V. L. (2015). LGN directs interphase
endothelial cell behavior via the microtubule network. PLoS ONE, *10*(9), 1–15.
https://doi.org/10.1371/journal.pone.0138763

Wylie, L. A., Mouillesseaux, K. P., Chong, D. C., & Bautch, V. L. (2018). Developmental
SMAD6 loss leads to blood vessel hemorrhage and disrupted endothelial cell
junctions. Developmental Biology, *442*(2), 199–209.
https://doi.org/10.1016/j.ydbio.2018.07.027

Yang, F., Zhang, Y., Zhu, J., Wang, J., Jiang, Z., Zhao, C., Yang, Q., Huang, Y., Yao,
W., Pang, W., et al. (2020). Laminar Flow Protects Vascular Endothelial Tight
Junctions and Barrier Function via Maintaining the Expression of Long Non-coding
RNA MALAT1. Frontiers in Bioengineering and Biotechnology, *8*, 647.
https://doi.org/10.3389/fbioe.2020.00647

Zhang, J., Felder, A., Liu, Y., Guo, L. T., Lange, S., Dalton, N. D., Gu, Y., Peterson, K.
L., Mizisin, A. P., Shelton, G. D., et al. (2010). Nesprin 1 is critical for nuclear
positioning and anchorage. Human Molecular Genetics, *19*(2), 329–341.
https://doi.org/10.1093/hmg/ddp499

Zhang, Q., Ragnauth, C. D., Skepper, J. N., Worth, N. F., Warren, D. T., Roberts, R. G.,
Weissberg, P. L., Ellis, J. A., & Shanahan, C. M. (2005). Nesprin-2 is a multi-
isomeric protein that binds lamin and emerin at the nuclear envelope and forms a
subcellular network in skeletal muscle. Journal of Cell Science, *118*(Pt 4), 673–687.
https://doi.org/10.1242/jcs.01642

Zhang, X., Lei, K., Yuan, X., Wu, X., Zhuang, Y., Xu, T., Xu, R., & Han, M. (2009).
SUN1/2 and Syne/Nesprin-1/2 complexes connect centrosome to the nucleus
during neurogenesis and neuronal migration in mice. Neuron, *64*(2), 173–187.
https://doi.org/10.1016/j.neuron.2009.08.018

Zhou, C., Li, C., Zhou, B., Sun, H., Koullourou, V., Holt, I., Puckelwartz, M. J., Warren,
D. T., Hayward, R., Lin, Z., et al. (2017). Novel nesprin-1 mutations associated with
dilated cardiomyopathy cause nuclear envelope disruption and defects in
myogenesis. Human Molecular Genetics, *26*(12), 2258–2276.
https://doi.org/10.1093/hmg/ddx116

Zhu, R., Antoku, S., & Gundersen, G. G. (2017). Centrifugal Displacement of Nuclei
Reveals Multiple LINC Complex Mechanisms for Homeostatic Nuclear Positioning.
Current Biology : CB, *27*(20), 3097-3110.e5.
https://doi.org/10.1016/j.cub.2017.08.073

1351

FIGURE LEGENDS

Figure 1. The nuclear LINC protein SUN1 regulates vascular development.

(a) Schematic of tamoxifen-induced excision of exon 4 of *Sun1* in pups from cross of

Sun1$^{fl/fl}$ X *Sun1$^{fl/+}$;Cdh5CreERT2* mice. **(b)** Representative images of P7 mouse retinas

of indicated genotypes, stained for IB4 (isolectin). Scale bar, 500µm. Inset shows

vascular plexus ahead of vein. Red line shows expansion of vascular front. Scale bar

inset, 150µm. **(c)** Quantification of vessel network radial expansion in **b**. n=186 ROIs

from 44 retinas (controls) and 63 ROIs from 16 retinas (*Sun1^{iECKO}*) from 6 independent

litters. ****, *p*<0.0001 by student's two-tailed unpaired *t*-test. **(d)** Quantification of

vascular density ahead of either arteries or veins. n=87 ROIs (controls, artery), 38 ROIs

(*Sun1^{iECKO}*, artery), 84 ROIs (controls, vein), and 37 ROIs (*Sun1^{iECKO}*, vein) from 27

retinas (controls) and 12 retinas (*Sun1^{iECKO}*) from 3 independent litters. **, *p*<0.01 by

student's two-tailed unpaired *t*-test. **(e)** Representative images of IB4 (isolectin) (green,

vessels) and VE-cadherin (white, junctions) staining in P7 retinas of indicated

genotypes. Scale bar, 50µm. **(f)** Quantification of ectopic VE-cadherin as shown in **e**.

n=144 junctions (9 retinas, controls) and 160 junctions (10 retinas, *Sun1^{iECKO}*). ****,

p<0.0001 by student's two-tailed unpaired *t*-test.

1369

Figure 2. Nuclear SUN1 is required for sprouting angiogenesis.

(a) Representative images of HUVEC with indicated siRNAs in 3D angiogenic sprouting

assay. Sprouts were stained for Phalloidin (actin) and then depth encoded such that

cooler colors are further in the Z-plane and warmer colors are closer in the Z-plane.

Scale bar, 100µm. **(b)** Quantification of average sprout length of 3D angiogenic sprouts

shown in **a**. n=42 beads (NT) and 43 beads (SUN1 KD) compiled from 5 replicates. ****,

p<0.0001 by student's two-tailed unpaired *t*-test. **(c)** Quantification of branches/mm of

3D angiogenic sprouts shown in **a**. n=41 beads (NT) and 43 beads (SUN1 KD) compiled

from 5 replicates. ****, $p<0.0001$ by student's two-tailed unpaired t-test. **(d)** Stills from

Movie S1 and Movie S2 showing sprouting dynamics of HUVEC with indicated siRNAs

over 50h. Scale bar, 50µm. **(e)** Quantification of HUVEC sprout extensions and

retractions shown in **d**. n=101 sprouts (NT) and 77 sprouts (SUN1 KD) compiled from 3

replicates. $p<0.001$ by χ^2 analysis. **(f)** Representative images of HUVEC with indicated

siRNAs and stained with indicated antibodies in the 3D sprouting angiogenesis assay.

Endothelial cells were stained for DAPI (cyan, DNA), phalloidin (green, actin), and VE-

cadherin (white, junctions). Arrows indicate normal junctions; arrowheads indicate

abnormal junctions. Scale bar, 20µm. **(g)** Quantification of ectopic VE-cadherin as

shown in **f**. n=32 junctions (NT) and 30 junctions (SUN1 KD) compiled from 2 replicates.

****, $p<0.0001$ by student's two-tailed unpaired t-test.

1389

Figure 3. SUN1 regulates actin dynamics and angiogenic sprout extension *in*

***vivo*.**

(a) Representative images of zebrafish embryos at 34 hpf with indicated morpholino

treatments; anterior to left. *Tg(fli:LifeAct-GFP)* (green, vessels). Insets show ISVs with

filopodia, outlines highlighted to show filopodia. Scale bar, 20µm. **(b)** Quantification of

filopodia number shown in **a**. n=39 ROIs (15 fish, NT) and 56 ROIs (20 fish, *sun1b* MO)

compiled from 3 replicates. **, $p<0.01$ by student's two-tailed unpaired t-test. **(c)**

Quantification of average filopodia length shown in **a**. n=39 ROIs (15 fish, NT MO) and

56 ROIs (20 fish, *sun1b* MO) compiled from 3 replicates. ****, $p<0.0001$ by student's

two-tailed unpaired t-test. **(d)** Representative images of zebrafish embryos at 34 hpf

with indicated genotypes; anterior to left. *Tg(fli:LifeAct-GFP)* (green, vessels). Insets

show ISVs with filopodia, outlines highlighted to show filopodia. Scale bar, 20µm. **(e)**

Quantification of filopodia number shown in **d**. n=27 ROIs (9 fish, $sun1b^{+/+}$) and 30 ROIs

(10 fish, $sun1b^{sa33109}$) compiled from 2 replicates. ns, not significant by student's two-

tailed unpaired t-test. **(f)** Quantification of average filopodia length shown in **d**. n=27

ROIs (9 fish, $sun1b^{+/+}$) and 30 ROIs (10 fish, $sun1b^{sa33109}$) compiled from 2 replicates.

, $p<0.01$ by student's two-tailed unpaired t-test. **(g) Stills from Movie S3 and Movie S4

showing ISV sprouting from 26 hpf to 36 hpf in zebrafish embryos with indicated

morpholino treatment; anterior to left. *Tg(fli:LifeAct-GFP)* (green, vessels). White

arrowhead points to ISV that does not extend or connect to DLAV. Yellow arrow points

to ISV that extends but does not connect to DLAV. Scale bar, 20µm. **(h)** Quantification

of ISV connection to DLAV shown in **g**. n=32 ISVs (6 fish, NT MO) and 36 ISVs (6 fish,

sun1b MO) compiled from 2 replicates. $p<0.05$ by χ^2 analysis. **(i)** Representative images

of zebrafish embryos at 34 hpf with indicated morpholino treatments; anterior to left.

Tg(fli:LifeAct-GFP) (green, vessels); ZO-1 (white junctions). Outlines highlighted to

show junction shapes. Scale bar, 10µm. **(j)** Quantification of junction morphology shown

in **i**. n=136 junctions (21 fish, NT MO) and 142 junctions (22 fish, *sun1b* MO). *******,

$p<0.001$ by student's two-tailed unpaired t-test.

1418

Figure 4. SUN1 stabilizes endothelial cell-cell junctions and regulates barrier

function.

(a) Representative images of HUVEC with indicated knockdowns in monolayers.

Endothelial cells were stained for DAPI (cyan, DNA) and VE-cadherin (white, junctions).

Insets show junctions. Scale bar, 10µm. **(b)** Representative graph of impedance

measured by RTCA. **(c)** Quantification of % change in cell index for RTCA measured at

5h. Normalized to NT cell index. n=5 replicates. *****, $p<0.05$ by student's two-tailed

unpaired t-test. **(d)** Representative images of HUVEC with indicated siRNAs plated on

biotinylated fibronectin and exposed to 15 dyn/cm^2 shear stress for 72 h then treated

with streptavidin. Endothelial cells were stained for DAPI (cyan, DNA), streptavidin

(green), and VE-cadherin (white, junctions). Arrow indicates flow direction. Insets show

junctions. Scale bar, 20μm. **(e)** Quantification of cell alignment shown in **d**. n=59 cells

(NT) and 73 cells (SUN1 KD) compiled from 3 replicates. **(f)** Quantification of

streptavidin area shown in **d**. n=15 ROIs (NT) and 15 ROIS (SUN1 KD) compiled from 3

replicates. **(g)** Representative images of HUVEC with indicated siRNAs showing

adherens following EDTA washout. Endothelial cells were stained for DAPI (cyan, DNA)

and VE-cadherin (white, junctions). Insets show junctions. Scale bar, 20μm. **(h)**

Quantification of VE-cadherin line scans at 20 min, 40 min, and 60 min post EDTA

washout in **g**. 20 min: n=31 junctions (NT) and 23 junctions (SUN1 KD); 40 min: n=49

junctions (NT) and 33 junctions (SUN1 KD); 60 min: n=33 junctions (NT) and 33

junctions (SUN1 KD) compiled from 3 replicates. ns, not significant; ****, $p<0.0001$ by

student's two-tailed unpaired t-test.

1441

Figure 5. SUN1 regulates microtubule localization and dynamics in endothelial

cells.

(a) Representative images of HUVEC with indicated siRNAs and indicated treatments.

Endothelial cells were stained for DAPI (cyan, DNA) and VE-cadherin (white, junctions).

Insets show junctions. Scale bar, 20μm. **(b)** Quantification of VE-cadherin line scans for

treatments shown in **a**. n=106 junctions (NT, vehicle), 101 junctions (NT, Nocodazole),

105 junctions (SUN1 KD, vehicle), and 96 junctions (SUN1 KD, Nocodazole) compiled

from 3 replicates. ns, not significant; ****, $p<0.0001$ by two-way ANOVA with Tukey's

multiple comparisons test. **(c)** Representative images of HUVEC with indicated siRNAs.

Endothelial cells were stained for α-tubulin (cyan, microtubules) and VE-cadherin (red,

junctions). Red insets show α-tubulin at the MTOC (microtubule organizing center),

yellow insets show α-tubulin contacts at junctions. Scale bar, 20μm. **(d)** Quantification of

α-tubulin area at the MTOC shown in **c**. n=19 cells (NT) and 10 cells (SUN1 KD)

compiled from 3 replicates. ns, not significant by student's two-tailed unpaired *t*-test. **(e)**

Quantification of peripheral α-tubulin area shown in **c**. n=39 cells (NT) and 46 cells

(SUN1 KD) compiled from 3 replicates. ****, $p<0.0001$ by student's two-tailed unpaired

t-test. **(f)** Quantification of contacts between α-tubulin and VE-cadherin shown in **c**.

n=75 junctions (NT) and 48 junctions (SUN1 KD) compiled from 3 replicates. ***,

$p<0.001$ by student's two-tailed unpaired *t*-test. **(g)** Stills from Movie S5 and Movie S6

showing microtubule growth in EB3-mCherry labeled HUVEC. White dot indicates start

of track. Yellow line indicates growth, red line indicates shrinkage. Scale bar, 10μm. **(h)**

Quantification of microtubule growth rate from EB3-mCherry microtubule tracking.

N=120 microtubules (12 cells, NT) and 117 microtubules (12 cells, SUN1 KD) compiled

from 2 replicates. ns, not significant by student's two-tailed unpaired *t*-test. **(i)**

Quantification of microtubule shrink rate from EB3-mCherry microtubule tracking. n=120

microtubules (12 cells, NT) and 117 microtubules (12 cells, SUN1 KD) compiled from 2

replicates. **, $p<0.01$ by student's two-tailed unpaired *t*-test. **(j)** Quantification of

catastrophe rate from EB3-mCherry microtubule tracking. n=120 microtubules (12 cells,

NT) and 117 microtubules (12 cells, SUN1 KD) compiled from 2 replicates. ****,

$p<0.0001$ by student's two-tailed unpaired *t*-test. **(k)** Quantification of percent of time

spent shrinking from EB3-mCherry microtubule tracking. n=120 microtubules (12 cells,

NT) and 117 microtubules (12 cells, SUN1 KD) compiled from 2 replicates. ****,

$p<0.0001$ by student's two-tailed unpaired *t*-test.

1475

Figure 6. SUN1 regulates endothelial cell contractility and exerts its effects on

junctions through the microtubule-associated GEF-H1.

(a) Representative images of HUVEC with indicated siRNAs and indicated treatments.

Endothelial cells were stained for DAPI (cyan, DNA) and VE-cadherin (white, junctions).

Insets show junctions. Scale bar, 20µm. **(b)** Quantification of VE-cadherin line scans for

treatments shown in **a**. n=159 junctions (NT, vehicle), 154 junctions (NT, blebbistatin),

151 junctions (SUN1 KD, vehicle), and 149 junctions (SUN1 KD, blebbistatin) compiled

from 3 replicates. ns, not significant; ****, $p<0.0001$ by two-way ANOVA with Tukey's

multiple comparisons test. **(c)** Representative images of HUVEC with indicated siRNAs

and indicated treatments. Endothelial cells were stained for DAPI (cyan, DNA) and VE-

cadherin (white, junctions). Insets show junctions. Scale bar, 20µm. **(d)** Quantification of

VE-cadherin line scans for treatments shown in **c**. n=75 junctions (NT, vehicle), 70

junctions (NT, thrombin), 71 junctions (SUN1 KD, vehicle), and 73 junctions (SUN1 KD,

thrombin) compiled from 3 replicates. ns, not significant; ****, $p<0.0001$ by two-way

ANOVA with Tukey's multiple comparisons test. **(e)** Representative images of HUVEC

with indicated siRNAs. Endothelial cells were stained for DAPI (blue, DNA), α-tubulin

(red, microtubules), and GEF-H1 (cyan). Insets show α-tubulin and GEF-H1

colocalization. Scale bar, 20µm. **(f)** Quantification of free GEF-H1 normalized to α-

tubulin associated GEF-H1 shown in **e**. n=30 cells (NT) and 30 cells (SUN1 KD)

compiled from 3 replicates. **, $p<0.01$ by student's two-tailed unpaired t-test. **(g)**

Representative images of HUVEC with indicated siRNAs and indicated treatments.

Endothelial cells were stained for DAPI (cyan, DNA) and VE-cadherin (white, junctions).

Insets show junctions. Scale bar, 20µm. **(h)** Quantification of VE-cadherin line scans

from knockdowns shown in **g**. n=169 junctions (NT), 166 junctions (SUN1 KD), 170

junctions (SUN1/GEF-H1 KD) compiled from 3 replicates. ns, not significant; ****,

$p < 0.0001$ by one-way ANOVA with Tukey's multiple comparisons test.

1502

Figure 7. SUN1 regulates endothelial cell junctions through nesprin-1.

(a) Representative images of HUVEC with indicated siRNAs cultured on biotinylated

fibronectin and treated with streptavidin upon confluence. Endothelial cells were stained

for DAPI (cyan, DNA), Streptavidin (green), and VE-cadherin (white, junctions). Insets

show junctions. Scale bar, 20µm. **(b)** Quantification of streptavidin area shown in **a**. n=6

ROIs (NT), 6 ROIs (SUN1 KD), and 6 ROIs (SUN1/Nesprin-1 KD) from 1 representative

replicate. *, $p < 0.05$; ****, $p < 0.0001$ by one-way ANOVA with Tukey's multiple

comparisons test. **(c)** Model describing proposed role of SUN1 in angiogenic sprouting

and endothelial cell junction stabilization.

a *Sun1^fl/fl;Cdh5CreERT2/+*

P0 P1 P2 P3 P4 P5 P6 P7

↑ ↑ ↑ Tamoxifen injection   ↑ Collect retinas

b Isolectin | Inset

Control

Sun1^iECKO

c %Radial Expansion — Control, *Sun1^iECKO* — ****

d Plexus distal to:

Artery — Vascular area/Total area — Control, *Sun1^iECKO* — **

Vein — Vascular area/Total area — Control, *Sun1^iECKO* — **

e Isolectin | VE-cadherin | Merge

Control

Sun1^iECKO

f Ectopic VE-cadherin/Junction VE-cadherin — Control, *Sun1^iECKO* — ****

a NT   SUN1 KD

Actin, Depth Encoded

b Average sprout length (μm) — NT vs SUN1 KD — ****

c Branches/mm — NT vs SUN1 KD — ****

d 0 h   10 h   20 h   30 h   40 h   50 h

NT

SUN1 KD

e n=101 sprouts

n=77 sprouts

p<0.001

Extensions
Retractions

f DAPI   Actin   VE-cadherin   Merge

NT

SUN1 KD

g Ectopic VE-cadherin/Junction VE-cadherin — NT vs SUN1 KD — ****

Table 1. SUN1 depletion does not alter endothelial gene expression.

Condition	# Upregulated DEG	# Downregulated DEG
NT_FLOW vs NT_STAT	1323	1109
SUN1_STAT vs NT_STAT	0	1
SUN1_FLOW vs NT_FLOW	1	1

Abbr: NT, non-targeting; STAT, static; DEG, differentially expressed genes.
Red numbers indicate that single downregulated gene was *SUN1*.