CsrA-controlled proteins reveal new dimensions of *Acinetobacter baumannii* desiccation tolerance

Yasuhiro Oda[1], Madelyn M. Shapiro [2], Nathan M. Lewis [3], Xuefei Zhong[4], Holly K. Huse[5], Weizhi Zhong[1], James E. Bruce[6], Colin Manoil[6], and Caroline S. Harwood[1]

[1]Department of Microbiology, University of Washington, Seattle, Washington, 98195-7735 USA'

[2] Seattle Children's Research Institute, Seattle, Washington, 98109 USA

[3] Department of Plant and Microbial Biology. University of Minnesota, St. Paul, MN. 55108 USA

[4] Analytical Chemistry Group. Regeneron Pharmaceuticals, Tarrytown, NY 10591 USA

[5] Department of Pathology. Harbor-UCLA Medical Center, Torrance, CA 90502-2059 USA

[6]Department of Genome Sciences, University of Washington, Seattle, Washington, 98195-5065 USA

Address correspondence to Caroline S. Harwood, csh5@uw.edu

Running title: *Acinetobacter baumannii* desiccation tolerance

Key words: *Acinetobacter baumannii*, desiccation, CsrA, intrinsically disordered proteins

ABSTRACT

Hospital environments are excellent reservoirs for the opportunistic pathogen *Acinetobacter*

baumannii in part because it is exceptionally tolerant to desiccation. We found that relative to

other *A. baumannii* strains, the virulent strain AB5075 was strikingly desiccation resistant at 2%

relative humidity (RH), suggesting that it's a good model for studies of the functional basis of

this trait. Consistent with results from other *A. baumannii* strains at 30% RH, we found the

global post-transcriptional regulator CsrA to be critically important for desiccation tolerance of

AB5075 at 2% RH. To identify CsrA-controlled proteins that may contribute to desiccation

tolerance we used proteomics to identify proteins that were differentially present in wild type and

csrA mutant cells. Subsequent mutant analysis revealed nine genes that were required for wild

type levels of desiccation tolerance, five of which had modest phenotypes. Catalase and a

universal stress protein gene were moderately important for desiccation tolerance and two

genes of unknown function had very strong desiccation phenotypes. The predicted amino acid

sequence of one of these genes predicts an intrinsically disordered protein. This category of

proteins is widespread in eukaryotes but less so in prokaryotes. Our results suggest there may

be mechanisms responsible for desiccation tolerance that have not previously been explored in

bacteria.

IMPORTANCE

Acinetobacter baumannii is commonly found in terrestrial environments but can cause

nosocomial infections in very sick patients. A factor that contributes to the prevalence of *A.*

baumannii in hospital settings is that it is intrinsically resistant to dry conditions. Here, we

established the virulent strain *A. baumannii* AB5075 as a model for studies of desiccation

tolerance at very low relative humidity. Our results show that this trait depends on two proteins

of unknown function, one of which is predicted to be an intrinsically disordered protein. This

category of protein is critical for the small animals named tardigrades to survive desiccation. Our

results suggest that *A. baumannii* may have novel strategies to survive desiccation that have

not previously been seen in bacteria.

INTRODUCTION

Hospital-acquired infections are an important healthcare concern and economic burden (1, 2)

and environmental persistence plays a critical role in the transmission of bacteria that cause

these infections (3–6). One such bacterium is *Acinetobacter baumannii,* an opportunistic

pathogen that infects very sick patients. It is responsible for about 2% of nosocomial infections

in the United States and Europe and the frequencies are higher in the rest of the world. *A.*

baumannii is especially problematic because on a global basis, about 45% of isolates are multi-

drug resistant (7). A factor that contributes to the prevalence of *A. baumannii* in hospital settings

is desiccation tolerance. *A. baumannii* can survive in a desiccated state on inanimate dry

surfaces for days to several months (8–10). These surfaces include materials that are often

encountered in the hospital, such as polyvinyl chloride, rubber, and stainless steel (11) .

When desiccated, bacteria must respond to diverse stresses that include accumulation of

reactive oxygen species, loss of cytoplasmic volume, and loss of cell membrane integrity (12,

13). Proteomics analysis of *A. baumannii* showed that desiccated cells had higher levels of

proteins involved in protein stabilization, antimicrobial resistance, and reactive oxygen species

detoxification (14). Attributes of *A. baumannii*, that have been shown to be associated with

desiccation tolerance include biofilm formation (15, 16) and protein aggregation (17). $LpxM_{AB}$-

dependent acetylation of lipid A is essential for survival of *A. baumannii* ATCC17978 at 40% RH

(18), and a *recA* mutant of ATCC17978, defective in DNA repair, had a pleiotropic phenotype,

including a defect in desiccation tolerance (19). *katE*, encoding catalase also contributes to

desiccation tolerance (20). Despite these observations, the number of genes identified in *A.*

baumannii that are specifically involved in desiccation tolerance is small. This could be because

A. baumannii cells have evolved modified cell structures that are both essential for viability and

important for desiccation tolerance. The genetic basis for this would be difficult to uncover in

mutant screens. It is also possible that the conditions of desiccation used, typically 30% RH in

studies to date, were not sufficiently severe to allow identification of some desiccation tolerance

genes.

With the goal of identifying new desiccation tolerance genes, we established *A. baumannii* strain

AB5075 as a model for studying desiccation tolerance under severe conditions of 2% RH. We

then followed up on a recent report showing that *csrA,* which encodes a global post-

transcriptional regulator, is important for desiccation tolerance (21) and identified nine CsrA-

controlled proteins that confer desiccation tolerance on AB5075 at 2% RH. One of these has

predicted properties that suggest new dimensions of desiccation tolerance.

RESULTS

Desiccation assay. Previous studies have shown that *A. baumannii* can survive in a desiccated

state for days to several months (8–11, 20). For these and other desiccation studies,

investigators worked with a variety of strains and usually incubated cells at either 30% RH or in

room air, which varied between 25 and 61% RH in one study (20). These differences can make

it difficult to compare desiccation phenotypes between studies. Thus we thought it important to

establish a robust desiccation assay that reduces experimental variables like choice of strains,

drying times, and RH during desiccation, using a highly desiccation resistant strain.

Following from previous reports, saturated calcium chloride hexahydrate solution placed in a

sealed plastic Snapware container caused the RH inside the container to rapidly equilibrate to

30% (9). We found that use of DRIERITE instead of calcium chloride, resulted in an RH of 2%.

To test desiccation tolerance under different conditions, we grew bacteria to a desired density in

Trypticase Soy Broth (TSB), harvested them, washed them twice, and resuspended them in

phosphate buffer to a final OD_{600} of 1. Drops of cell suspension were placed on polycarbonate

membranes and filtered to allow for rapid drying. The membranes were then placed in

uncapped 15 ml conical centrifuge tubes and incubated in desiccation containers. After various

periods of incubation, buffer was added to each centrifuge tube followed by 5 min of shaking on

a rotary shaker. Viable cell numbers were then determined by plating on TSB agar. To control

for the stress of filtration we did viable counts immediately following filtration and took this as our

"day 0" time point.

Relative desiccation tolerance of *A. baumannii* strains. As shown in Fig 1A, *A. baumannii*

strain AB5075 and *Escherichia coli* strain W3100 each survived desiccation at 30% RH far

better than *Pseudomonas aeruginosa* strain PAO1. However at 2% RH, *A. baumannii* survived

far better than either *E. coli* or *P. aeruginosa*. As has been reported (20, 22), we found that *A.*

baumannii stationary phase cells were much more tolerant to desiccation than actively growing

cells (Fig. S1) and so we routinely used stationary phase cells in our desiccation assays. We

compared the desiccaton tolerance of AB5075 to two additional frequently used laboratory

strains of *A. baumannii*, ATCC17978 and ATCC19606. We found that AB5075 was strikingly

more resistant than the others at 2% RH and somewhat mores resistant at 30% (Fig. 1B).

Given the strength of its phenotype, these findings indicate that AB5075 is a good model for

studing desiccation tolerance. This strain was isolated from a surgical wound, is multidrug

resistant and is highly virulent in an animal model (23). A comprehensive ordered mutant library

of AB5075 is available that has two to three seqenced Tn insertions in each gene and is called

the three-allelle library (24).

CsrA is critical for desiccation tolerance. We examined the contribution of the post-

transcriptional regulator CsrA to desiccation tolerance of AB5075 by constructing a *csrA*

deletion mutant (ΔcsrA). We found that the mutant grew poorly on TY agar and had an

elongated cell morphology when grown in TY broth (Fig. 2A). On agar plates, large colonies

frequently appeared on a backgound of poor growth, likely due to occurance of second site

suppressor mutations in the ΔcsrA strain. The ΔcsrA strain was also defective in growth on other

nutrient-rich media, including LB broth, nutrient broth, and TSB broth. A similar sensitivity to

growth in complex media was reported by Farrow et al. for several *A. baumannii* strains

including strain AB5075 (21). In agreement with Farrow et al., a ΔcsrA mutant grew as the wild

type in defined medium, in our case, M9 minimal medium with 10 mM succinate as a sole

carbon source (M9/succinate), and it had close to a wild type cell morphology (Fig 2A). A

Yersinia enterocolitica csrA mutant, has a growth defect in LB due to the presence of 90 mM of

NaCl (25). However, the *A. baumannii* ΔcsrA mutant was not sensitive to this level of NaCl. In

fact, the mutant grew in M9/succinate supplemented with up to 100 mM of NaCl without a

significant reduction of growth compared to the wild type.

When desiccated after growth in M9/succinate to stationary phase, the AB5075 *ΔcsrA* mutant

lost almost all viability over 6 days (Fig. 2B and Table 1). The desiccation phenotype was

complemented by expressing *csrA in trans*. ΔcsrA mutant cells incubated for 6 days after being

filtered and resuspended in PBS remained fully viable (Fig 2B).

CsrA affects mutiple cellular processes in *A. baumannii*. To identify genes whose

translation might be regulated by CsrA, we compared the proteomes of wild-type and ΔcsrA

cells (Table S2A). There were 97 proteins present at higher levels in the ΔcsrA mutant

compared to the wild type (ratio of ΔcsrA/WT ≥2.5, Table S2B). Among these were proteins for

type IV pilus assembly, synthesis of the siderophore ferric acinetobactin, and a

glutamate/aspartate transporter. The ΔcsrA mutant also had elevated levels of enzymes for the

catabolism of hydroxcinnamates, phenylacetate and quinate. Levels of an alcohol

dehydrogenase (ABUW_1621) and an aldehyde dehydrogenase (ABUW_1624) were also

elevated. The ∆*csrA* mutant was defective in pilus-mediated twiching motility as assessed by

movement aross a soft-agar plate (Fig. 3A). The mutant also had a severe growth defect when

grown on succinate in the presence of ethanol (Fig. 3B). One possible explanation for this is that

the *csrA* mutant metabolized ethanol to form toxic acetaldehyde to levels that slowed growth.

There were 106 proteins present in lower amounts in the ∆*csrA* mutant compared to the wild

type (ratio of WT/∆*csrA* ≥ 2.5, Table S2C). A large proportion of these (39%) are annotated as

hypothetical proteins. Several membrane proteins, and proteins annotated as involved in *β*-

lactam antibiotic resistance (ABUW_1194, 2619, and 3497), trehalose synthesis (ABUW_3123)

and possibly biofilm formation (ABUW_0916) were in lower abundance in the ∆*csrA* mutant

compared to wild type. As reported previously, a ∆*csrA* mutant did not form biofilms (21). This

phenotype was complemented by expressing the *csrA* gene *in trans* (Fig. 3C). The ∆*csrA*

proteome profile also suggested that CsrA is invovled in promoting the expression of proteins

invovled in oxidative stress, including peroxidase (ABUW_0628) and catalase (*katE*,

ABUW_2436). Indeed the ∆*csrA* mutant lacked detectable catalase activity (Fig. 3D).

Genes important for desiccation tolerance in *A. baumannii* AB5075. We took advantabge

of the three-allele transposon library to test how important some of the gene transcripts that

were likely to be controlled by CsrA were for desiccation tolerance (Tables 1 and S1). When

possible, we tested two different transposon mutants (transposon insertions in different

positions of the gene) for each gene. *A. baumannii* AB5075 produces opaque and translucent

colony variants that interconvert at high frequency and reflect changes in the thickness of

capsular exopolysaccharide (26). AB5075 cells with decreased capsule production are about

2.5-fold more sensitive to desiccation (27). Here, we used only opaque colonies of AB5075 and

its mutant derivatives in our desiccation assays.

We found that *katE* and *ABUW_2639* mutants were about 5-fold more sensitive to desiccation

than the wild type, whereas *ABUW_2433* and *ABUW_2437* mutants were greater than 100-fold

more sensitive to desiccation than the wild type (Table 1, Fig 4). The phentoytpes of

ABUW_2433 and *ABUW_2437* mutants could be complemented (Fig S2). The ABUW_2433

protein has 411 amino acids and is annotated as a KGG domain-containing protein. The KGG

domain comprises a small region in the N-terminus of the protein and the remainder of the

protein is annotated by InterPro as a disordered region that includes a series of AT_hook DNA

binding motifs (SMART SM00384). The full length ABUW_2433 sequence was predicted to be

intrinsically unstructured when queried with the IUPred3 tool (https://iupred.elte.hu) (28).

ABUW_2437 is annotated as an iron-containing redox enzyme or a heme-oxygenase -like

protein (Fig 4). The predicted *ABUW_2437* transcript has traits characterisitic of a target of CsrA

post-transcriptional regulation. The DNA sequence predicts a relatively long 5' untranslated

region (316 bp) and there is a predicted CsrA binding motif (GGA) in the ribosome binding site

of the transcript. *ABUW_2639* is annoated as belonging to a universal stress protein A family. It

has been shown to protect *A. baumannii* ATCC17987 from oxidative stress of hydrogen

peroxide (29).

We wondered if *ABUW_2433* and *ABUW_2437* might play a role in promoting desiccation

tolerance of the two *A. baumannii* strains, ATCC17978 and ATCC19606, that do not survive

well at 2% RH (Fig 1B). ATCC19606 has the gene region shown in Fig 4 intact, but the gene

that is homologous to *ABUW_2433*, encoding the KGG domain-containing protein, is annotated

as a pseudogene. ATCC17978 appears to be missing a gene in homologus to *ABUW_2433*.

However it has conitguous *katE* and iron-containing redox protein genes (*A1S_1386 and*

A1S_1385). Expression of the two AB5075 genes *in trans* improved the survival of the two

ATCC strains at 2% RH (Fig 5), providiing evidence that ABUW_2433 and ABUW_2437 are

generally important for desiccaton tolerance.

Other possible desiccation tolerance genes. As shown in Table 1, we identified an

additional six genes that are likely regulated by CsrA, that may have a small role in desiccation

tolerance. *ABUW_0916*, encoding a biofilm-associated protein and *otsA*, encoding tehalose-6-

phosphate synthase, are the only two of the six for which we can hypothesize some connection

to desiccation. Biofilms have been shown to be important for desiccation tolerance of bacteria

(15, 16) and trehalose has been shown to play a significant role in desiccation tolerance of

eukaryotes and bacteria (30, 31). When added extrinsically to cultures, trehalose increased the

desiccation tolerance of *A. baumannii* ATCC 19606 (22). However, a *ΔmtlD-otsB* mutant of

ATCC19606, defective in endogenous production of the compatible solutes, mannitol, and

trehalose, was not more sensitive to desiccation than the wild type (22).

DISCUSSION

Depletion of water during desiccation leads to loss of membrane integrity and accompanying

disruption of aerobic respiration results in the generation of reactive oxygen species, including

hydrogen peroxide (32). So, it makes sense that *katE*, encoding catalase, contributes to

desiccation tolerance. Proteomics analyses of *A. baumannii* showed that proteins involved in

redox defense including catalase, alkyl peroxidase reductases and superoxide dismutase were

elevated in stationary-phase cells (33), which is consistent with the observation made by many

that cells stationary-phase cells survive desiccation much better than exponentially growing

cells.

Since the desiccation-tolerance genes *ABUW_2433* and *ABUW_2437* are near or adjacent to

katE, on the genome and all are likely to be controlled by CsrA, it seemed important to consider

that they might somehow mediate oxidative stress tolerance even though the amino acid

sequences of the encoded proteins don't have motifs typically associated with reactive oxygen

species detoxification. However, we were unable to demonstrate that *ABUW_2433*::Tn and

ABUW_2437::Tn mutants were sensitive to hydrogen peroxide and paraquat - both powerful

oxidizing agents. In addition, a study that looked at effects of hydrogen peroxide exposure on

gene expression in *A. baumannii,* found that *katE* but not *ABUW_2433* or *ABUW_2437*, was

expressed at elevated levels and neither of these genes is part of the OxyR regulon that

controls the response to oxidative stress in *A. baumannii* (34). Thus, we do not think that either

of these proteins, which have the greatest defects in desiccation found to date, are likely to

function by protecting cells against oxidative stress

The predicted physical properties of ABUW_2433 provide suggestions as to how it may

function. It is an intrinsically disordered protein (IDP) that is highly hydrophilic, with 27%

positively charged amino acids residues and 31% negatively charged residues. It is also

predicted to assume a collapsed or extended conformation, likely depending on its context

(ROBETTA PFRMAT TS prediction; https://robetta.bakerlab.org). ABUW_2433 has 13 repeated

AT-hook DNA binding motifs that extend across about 70% of the protein. This motif

preferentially binds to AT-rich sequences in the minor groove of DNA. AT-hook DNA binding

motifs are found primarily in eukaryotic proteins, many of which have roles in transcriptional

regulation (35, 36). Only 8.5% of annotated AT hook DNA binding motifs are found in bacteria,

but about half of these are found in gamma proteobacteria, the group to which *A. baumannii*

belongs. We hypothesize that ABUW_2433 binds to *A. baumannii* DNA and somehow protects

it from desiccation-induced damage. IDPs are critical for the microscopic animals called

tardigrades to survive desiccation. When desiccated, some of these proteins vitrify and probably

trap desiccation sensitive molecules in a noncrystalline amorphous matrix, thereby protecting

them from denaturation or other forms of destruction (37, 38). IDPs or proteins with IDP

domains are less common in prokaryotes than in eukaryotes, but drawing from work on

eukaryotes, they are proposed to play a central role in cellular process in bacteria that may

depend on the formation of molecular condensates (39). It is possible that this is important for

the viability of desiccated *A. baumannii*.

Although not much work has been done on *Acinetobacte*r CsrA, based on what is known for

other gamma proteobacteria, we hypothesize that a set of ncRNAs that is induced by a GacSA

(ABUW_3306 and ABUW_3639) two-component regulatory system, controls the repressor

activity of CsrA by sequestering it (40). We can draw a link between the GasSA system and

CsrA because they both appear to control catabolism of the aromatic compound phenylacetate.

An *A. baumannii* ∆*gacA* mutant is unable to catabolize phenylacetate (41), and our proteomics

results suggest that CsrA acts to repress the synthesis of at least one enzyme required for

phenylacetate degradation. In fact, we have determined that the ∆*csrA* mutant grows better on

phenylacetate; with a doubling time of 39 min, than the wild type, which has doubling time of 50

min. We hypothesize that a ∆*gacA* mutant does not synthesize ncRNAs that would normally

"sponge-up" CsrA, thus allowing CsrA to bind to the 13 -gene (ABUW_2524 to ABUW_2536)

phenylacetate mRNA transcript to repress its translation. At this point we do not have a clear

understanding of the inventory of *A. baumannii* ncRNAs that may bind to CsrA, but ncRNAs are

abundant in AB5075, and several of them are expressed at extremely high levels (42). The

desiccation phenotype of CsrA appears to depend on its ability to activate translation and

although it's difficult to reconcile this activity with a model where CsrA is sequestered by

ncRNAs, it is known that ncRNA turnover can occur resulting in the release of free CsrA (43).

Most of what is known about mechanisms of CsrA action centers on its role as a repressor of

translation (44–47) and it may be of interest to probe its capability as an activator in *A.*

baumannii.

We found that *A. baumannii* AB5075 survived desiccation for six days at 2% RH much better

than two other *A. baumannii* strains that we tested, but it is important to note that most studies

of desiccation tolerance have been carried out at about 30% RH or in room air and the

emphasis has been on the number of days or months that a particular strain remains viable

when desiccated. When Farrow et al (20) tested the survival of several strains that were dried

and incubated at an RH of 25–61% (mean 46%) they found AB5075 to have an average

survival time of 90 days, whereas strains ATCC19606 and ATCC17978 had average survival

times of 3 and 34 days respectively. Even though AB5075 is tolerant to desiccation over

months at a mean RH of 46% and over days at 2% RH, we cannot necessarily conclude that the

same sets of genes are needed for desiccation tolerance under these two different conditions.

For example, Farrow et al. (20) found that the response-regulator protein BmfR was important

for desiccation tolerance of ATCC17978 in long term desiccation assays, whereas we did not

observe a role of *bmfR* in protecting AB5075 from desiccation in shorter term incubations at

2%RH (Table S1).

Here we established a robust assay for desiccation tolerance of a highly virulent strain of *A.*

baumannii and identified two genes, *ABUW_2433* and *ABUW_2437*, that are extremely

important for desiccation tolerance. Outside of the *Acinetobacter* genus, *ABUW_2437* has

homologs in *Pseudomonas stutzeri* (40% amino acid identity). However, except for a partial

homolog found in *Enterobacteriaceae* bacterium TzEc051 (99% identity over 38% of the

AB5075 protein), *ABUW_2433* does not appear outside *Acinetobacter*. This and the unusual

predicted physical properties of ABUW_2433 as anIDP, suggest that there is novel physiological

basis for desiccation tolerance in *A. baumannii* that remains to be explored.

MATERIALS AND METHODS

Bacterial strains and growth conditions. Strains used in this study are listed in Table S3A.

Strain AB5075 (AB5075-UW) was used as a wild type (24) and individual transposon mutants

were obtained from the Manoil lab comprehensive ordered transposon mutant library at the

University of Washington (24). All strains except for the $\Delta csrA$ mutant were routinely grown and

maintained in TY (10 g Tryptone, 5 g Yeast extract, and 8 g NaCl in 1000 ml) medium or BBL

Trypticase Soy Broth (TSB) media at 37°C, unless otherwise stated. The $\Delta csrA$ mutant was

grown in M9/succinate.

Desiccation assay. Strains from a frozen stock (-80°C) were streaked onto TY plates and

incubated at 37°C. Colonies (three to five) were picked and inoculated into 2 ml of TSB, and

cultures were grown overnight at 37°C with a shaking speed of 200rpm. Overnight cultures were

diluted to yield an initial OD_{600} of 0.025 in 10 ml TSB in a 50 ml Erlenmeyer flask. Cultures were

grown at 37°C with a shaking speed of 200 rpm to mid-exponential-phase (OD_{600}=0.4 to 0.6) or

stationary-phase (24 hours after inoculation). Cells were harvested by centrifugation and

washed twice with Dulbecco's phosphate-buffered saline (DPBS, Gibco), and cell density was

adjusted to OD_{600}=1 (about 5 x 10^8 cells/ml) with DPBS. Cell suspension (2 spots of 50 µl each

per membrane) was filtered onto a 0.4 µm Whatman nucleopore polycarbonate track-etched

membranes (25 mm diameter) that had been placed in Nalgene analytical filter units, and the

membranes were then placed into 15 ml uncapped centrifuge tubes. To obtain the T0 (baseline)

viable cell number, 1 ml of DPBS was immediately added to one tube and incubated for 5 min at

room temperature (24 $\pm$ 2°C) on a rotary shaker. Viable cell numbers were determined by

plating on TSB agar. For desiccation, tubes with membranes were placed in a Snapware

containers (2.3 x 6.3 x 8.4 inches) that contained DRIERITE in the lids of 50 ml centrifuge tubes

(x4, 7.5 g of DRIERITE desiccant in each lid) or saturated calcium chloride hexahydrate solution

in 5 ml beaker (x8) to yield the RHs of 2% or 30% ($\pm$ 2), respectively. The Snapware containers

were incubated at room temperature. Digital hygrometers (VWR International Ltd) were placed

in each container to monitor the RH. At desired time points, tubes containing membranes were

removed from the containers, 1 ml of DPBS was added to each tube, and incubated for 5 min at

room temperature on a rotary shaker. Viable cell counts were determined on TSB agar. For

each strain, a minimum of three biological replicates of desiccation assays were performed

except for individual transposon mutants, which were assayed twice for each allele.

Construction of the Δ*csrA* mutant. In-frame deletion of the *csrA* (*ABUW_2750*) gene was

generated by overlap extension PCR as described (48). PCR primers are listed in Table S3B.

PCR product was cloned into mobilizable suicide vector pEX2-TetRA and transformed into *E.*

coli NEB 10-beta (New England Bio Labs). The sequence-verified deletion construct was

transformed into *E. coli* S17-1, and further mobilized into *A. baumannii* strain AB5075 by

conjugation on TY agar. Single recombinant conjugants were first selected on M9/succinate

plate containing 20 µg/ml Tc, and Tc resistant colonies were further plated onto M9/succinate

plate containing 5% sucrose. Sucrose resistant and Tc sensitive colonies were screened by

colony PCR and sequencing to validate the expected chromosomal in-frame deletion of the *csrA*

gene.

To complement the Δ*csrA* mutant, the full length *csrA* gene plus the 15 bp upstream that

contains the putative ribosome binding site was PCR amplified and cloned into pMMB67EH-

TetRA. The construct was transformed into *E. coli* NEB 10-beta (New England Bio Labs). The

sequence-verified construct was transformed into *E. coli* S17-1, and further mobilized into the

Δ*csrA* mutant by conjugation on M9/succinate plates containing 20 µg/ml Tc. As a negative

control, empty vector pMMB67EH-TetRA was used. The same procedure was used to clone

ABUW_2433 and *ABUW_2437* for complementation experiments.

Phenotypic characterization of the Δ*csrA* mutant. M9/succinate was used in all experiments.

For motility assays, an overnight culture (16 to 18 hours) was diluted to yield OD_{600}=0.5, and 2

μl of sample was spotted onto the freshly prepared M9/succinate plate and incubated at 37°C

for 24 h. For biofilm assays, an overnight culture was inoculated into 100 μl of M9/succinate in

Costar vinyl 96 well "U" bottom plates (initial OD_{600}=0.05), and the plates were sealed with

Breath-Easy sealing membranes. After incubation at room temperature for 48 h, culture was

removed, the plate was rinsed with tap water twice, and 150 μl of 0.1% crystal violet solution

was added to each well. After incubating at room temperature for 15 min, crystal violet solution

was removed, the plate was rinsed with tap water 5 times, and the plate was dried at room

temperature. For catalase assays, cells were harvested at OD_{600}=0.5, supernatant was

removed, and cells were resuspended in DPBS to yield 10 mg wet cell/100 μl DPBS. 100 μl of

cell suspensions were placed in 13 x 100 mm borosilicate glass tubes. Then 100 μl of 1% Triton

X-100 and 100 μ1 of 30% hydrogen peroxide were added, mixed thoroughly, and incubated for

15 min at room temperature (49) .

Label-free protein quantification. Since the Δ*csrA* mutant had a severe growth defect on TY

medium, both the wild type and Δ*csrA* mutant were grown in M9/succinate. Two biological sample

replicates were prepared for each strain. Cells from each culture were harvested at OD_{600}=0.5 by

centrifugation, washed twice with DPBS, and cells were stored in -80°C before further analysis.

Cells were lysed in buffer containing 4% SDS, 100 mM Tris pH8.0, 10 mM DTT by heating at

95°C for 5 min. After cooling to room temperature, the lysates were sonicated with ultrasonication

probe on ice to shear DNA. Total protein concentration was determined by the BCA assay

(Thermo Pierce, Rockford, IL). 500 μg of each protein lysate was reduced and alkylated, diluted

in 8 M urea solution, and the SDS was removed with a 3kD molecular weight cutoff filter. After

buffer exchange, the protein lysates were digested with trypsin (Promega, Madison, WI) at 37°C

overnight and the digested samples were desalted with 1cc C18 Sep-Pak solid phase extraction

cartridges (Milford, MA, Waters). The eluted samples were vacuum dried and resuspended in

0.1% formic acid. Reverse phase nanoLC-MS analysis of the protein digests was carried out with

a Thermo Easy-nanoLC coupled to a Thermo Q-Exactive Plus Orbitrap mass spectrometer.

Triplicate top 20 data-dependent acquisition runs were acquired for each sample, and 1 µg of

protein digest was loaded for each run. The peptides were separated by a 50 cm x 75 µm I.D C8

column (5 µm diameter, 100 Å pore size C8 MichromMagic beads) with a 90 min 10 to 30% B

gradient (solvent A: 0.1% formic acid in water, solvent B: 0.1% formic acid in acetonitrile, flow rate

300 nl/min). The MS data acquisition parameters were set as follows: full MS scan resolution 70k,

maximum ion injection time 100 mS, AGC target 10^6, scan range of 400 to 2000 m/z; MS/MS scan

resolution 17.5 k, maximum ion injection time 100 mS, AGC target 5^4, isolation window 1.6 m/z,

HCD NCE 35 scan range of 200 to 2000 m/z; loop count 20, intensity threshold 5^3, underfill ratio

1%, dynamic exclusion 10 sec. High resolution MS^2 spectra were searched against a target-decoy

proteome database of strain AB5075 (a total of 7678 sequences) downloaded from Uniprot

(Oct17, 2017) using Comet (version 2015.02 rev. 1) (50) with following parameters: precursor

peptide mass tolerance 20 ppm, allowing for -1, 0, +1, +2, or +3 ^{13}C offsets; fragment ion mass

tolerance 0.02 Da; static modification, carbamidomethylation of cysteine (57.0215 Da); variable

modification, methionine oxidation (15.9949 Da). The search results were further processed by

PeptideProphet (51) for probability assignment to each peptide-spectrum match, and

ProteinProphet (52) for protein inference and protein probability modeling. The output pepXML

files from three technical replicates were grouped for subsequent spectral counting analysis using

Abacus (53). The pepXML and protXML files for each sample, combined ProteinProphet file from

all samples were parsed into Abacus for spectral counting of each protein. The following filters

were applied for extracting spectral counts from MS/MS datasets: (1) the minimum

PeptideProphet score the best peptide match of a protein must have maxIniProbTH=0.99; (2) The

minimum PeptideProphet score a peptide must have to be even considered by Abacus,

iniProbTH=0.50; (3) The minimum ProteinProphet score a protein group must have in the

COMBINED file, minCombinedFilePw=0.90. Spectral counts for 1616 proteins were reported

across four sample groups (two strains and two biological replicates) with estimated protein false

discovery rate of 1.94%. The protein expression fold changes between wild type AB5075 and

Δ*csrA* mutant were computed from adjusted spectral counts output from Abacus. The mass

spectometry data have been deposited to the ProteomeXchange Consortium via the PRIDE

partner repository with the dataset identifier xxxxxxxxx..

ACKNOWLEDGMENTS

We thank Dr. Hemantha Don Kulasekara and Dr. Samuel Miller (University of Washington) for

sharing the pEX2-TetRA and pMMB67EH-TetRA vectors. We thank Indranil Biswas for alerting

us to the possibility that intrinsically disordered proteins could be involved in desiccation

tolerance in *A. baumannii*.

FUNDING

This work was supported by the Functional Genomics Program, National Institute of Allergy and

Infectious Diseases under Grant 1U19AI107775-01.

409 **Table 1. Genes that contribute to desiccation tolerance of *A. baumannii* AB5075.**

Gene	Gene name	Gene annotation	Relative viability loss ratio (day 6/day 0 desiccation)[a]	Protein abundance (ratio of WT/*ΔcsrA*)
AB5075 (WT)			1.0	
ABUW_0916		Biofilm-associated protein	1.5	2.6
ABUW_2433		KGG domain-containing protein	125	4.3
ABUW_2436	*katE*	Catalase	7.5	3.2
ABUW_2437		Heme oxygenase-like protein	150	10.5
ABUW_2639		Universal stress protein family	6.0	4.4
ABUW_2724		Hypothetical protein	4.0	2.4
ABUW_2750	*csrA*	Carbon storage regulator	480	8.5
ABUW_3123	*otsA*	Trehalose-6-phosphate synthase	2.0	10.5
ABUW_3346	*acnA*	Aconitate hydratase 1	1.5	3.2

410 [a] All mutants were except the ΔcsrA strain were grown in TSB broth to the stationary phase of growth
411 prior to being desiccated. The viability loss of wt day6/day 0 was 17 for the WT grown in TSB broth. The
412 viability loss of WT grown in M9/succinate was 276
413
414

FIGURE LEGENDS

Figure 1. Desiccation tolerance of *A. baumannii*. (A) Comparison of AB5075, *E. coli* W3110,

and *P. aeruginosa* PAO1 at 30% RH or 2% RH. (B) Desiccation tolerance of *A. baumannii*

strains AB5075, ATCC17978, and ATCC19606 after 0 days (control) and 6 days of desiccation

at 30% RH or 2% RH. The cell numbers represent the total number of viable cells recovered

from each membrane. The data are the average of three or more biological replicates, and

standard deviations are shown as error bars.

Figure 2. Growth and desiccation tolerance of a Δ*csrA* mutant (A) Comparison of wild type (WT)

and a Δ*csrA* mutant on TY and M9/succinate agar and in TY and M9/succinate broth. (B)

Desiccation tolerance of WT, Δ*csrA* mutant, Δ*csrA* mutant with pMMB (empty vector), and Δ*csrA*

mutant with *csrA* expressed in trans (pAB2750) at day 0 and at day 6 of desiccation at 2% RH

or at 6 days in PBS. The cell number represents the total number of viable cells recovered from

each membrane. The data are the average of three or more biological replicates, and standard

deviations are shown as error bars.

Figure 3. Phenotypes of the Δ*csrA* mutant: (A) twitching motility; (B) growth of wild type (black

symbol) or the Δ*csrA* mutant (red symbol) in M9/succinate in the absence (closed symbol) or

presence (open symbol) of 0.5% ethanol. The data shown are representative of each strain and

condition; (C) crystal violet staining of biofilms from wild type, Δ*csrA* mutant, Δ*csrA* mutant with

pMMB (empty vector), or Δ*csrA* mutant with pAB2750; and (D) catalase activity of wild type,

Δ*csrA* mutant, Δ*csrA* mutant with pMMB (empty vector), or Δ*csrA* mutant with pAB2750.

Figure 4. *A. baumannii* genes important for desiccation tolerance. (A) Genes and (B)

Desiccation tolerance of mutants at 2% RH. The cell number represents the total number of

viable cells recovered from each membrane. The data are the average of three or more

biological replicates, and standard deviations are shown as error bars.

Figure 5. Effect of expression of *ABUW_2433* (red bars) or *ABUW_2337* (blue bars) gene *in*

trans on desiccation tolerance of *A. baumannii* (A) ATCC17978 and (B) ATCC19606 after 0

days (control) and 6 days of desiccation at 2% RH. Empty vector (pMMB, black bars) was used

as a control. The data are the average of three or more biological replicates, and standard

deviations are shown as error bars.

SUPPLEMENTAL MATERIAL

Figure S1. Desiccation tolerance of *A. baumannii* AB5075 cells from the log- and stationary-phases of growth. The cell number represents the total number of viable cells recovered from each membrane. The data are the averages of three or more biological replicates, and standard deviations are shown as error bars.

Figure S2. Complementation of desiccation phenotypes of (A) *ABUW_2433*::Tn and (B) *ABUW_2437*::Tn after 0 days (control) and 6 days of desiccation at 2% RH. Empty vector (pMMB) was used as a control. The data are the average of three or more biological replicates, and standard deviations are shown as error bars.

Table S1. Mutants from the three-allele library tested for desiccation tolerance

Table S2. (A) Label-free protein quantification of wild type and the Δ*csrA* mutant. (B) List of proteins up-regulated (ratio of Δ*csrA*/WT ≥2.5 and Δ*csrA* read count ≥7.5) in the Δ*csrA* mutant compared to wild type. (C) List of proteins down-regulated (ratio of Δ*csrA*/WT ≤0.4 and WT read count ≥7.5) in the Δ*csrA* mutant compared to wild type.

Table S3. (A) Bacterial strains used in this study. (B) Plasmids and primers used in this study

REFERENCES

1. Magill SS, Edwards JR, Bamberg W, Beldavs ZG, Dumyati G, Kainer MA, Lynfield R, Maloney M,
McAllister-Hollod L, Nadle J, Ray SM, Thompson DL, Wilson LE, Fridkin SK, Team EIPH-AI and
AUPS. 2014. Multistate Point-Prevalence Survey of Health Care–Associated Infections. New Engl J
Medicine 370:1198–1208.

2. Marchetti A, Rossiter R. 2013. Economic burden of healthcare-associated infection in US acute care
hospitals: societal perspective. J Med Econ 16:1399–1404.

3. Datta R, Platt R, Yokoe DS, Huang SS. 2011. Environmental Cleaning Intervention and Risk of
Acquiring Multidrug-Resistant Organisms From Prior Room Occupants. Arch Intern Med 171:491–494.

4. Huang SS, Datta R, Platt R. 2006. Risk of Acquiring Antibiotic-Resistant Bacteria From Prior Room
Occupants. Arch Intern Med 166:1945–1951.

5. Munoz-Price LS, Namias N, Cleary T, Fajardo-Aquino Y, DePascale D, Arheart KL, Rivera JI, Doi Y.
2013. Acinetobacter baumannii: Association between Environmental Contamination of Patient Rooms
and Occupant Status. Infect Control Hosp Epidemiology 34:517–520.

6. Thom KA, Johnson JK, Lee MS, Harris AD. 2011. Environmental contamination because of
multidrug-resistant Acinetobacter baumannii surrounding colonized or infected patients. Am J Infect
Control 39:711–715.

7. Harding CM, Hennon SW, Feldman MF. 2018. Uncovering the mechanisms of Acinetobacter
baumannii virulence. Nat Rev Microbiol 16:91–102.

8. Giannouli M, Antunes LC, Marchetti V, Triassi M, Visca P, Zarrilli R. 2013. Virulence-related traits of
epidemic Acinetobacter baumannii strains belonging to the international clonal lineages I-III and to the
emerging genotypes ST25 and ST78. Bmc Infect Dis 13:282.

9. Jawad A, Seifert H, Snelling AM, Heritage J, Hawkey PM. 1998. Survival of Acinetobacter baumannii
on Dry Surfaces: Comparison of Outbreak and Sporadic Isolates. J Clin Microbiol 36:1938–1941.

10. Antunes LCS, Imperi F, Carattoli A, Visca P. 2011. Deciphering the multifactorial nature of
Acinetobacter baumannii pathogenicity. Plos One 6:e22674.

11. Wendt C, Dietze B, Dietz E, Rüden H. 1997. Survival of Acinetobacter baumannii on dry surfaces. J
Clin Microbiol 35:1394–1397.

12. Lebre PH, Maayer PD, Cowan DA. 2017. Xerotolerant bacteria: surviving through a dry spell. Nat
Rev Microbiol 15:285–296.

13. Potts M. 1994. Desiccation tolerance of prokaryotes. Microbiol Rev 58:755–805.

14. Gayoso CM, Mateos J, Méndez JA, Fernández-Puente P, Rumbo C, Tomás M, Ilarduya ÓM de, Bou
G. 2013. Molecular Mechanisms Involved in the Response to Desiccation Stress and Persistence in
Acinetobacter baumannii. J Proteome Res 13:460–476.

15. Espinal P, Marti S, Vila J. 2012. Effect of biofilm formation on the survival of Acinetobacter
baumannii on dry surfaces. J Hosp Infect 80:56 60.

16. Orsinger-Jacobsen SJ, Patel SS, Vellozzi EM, Gialanella P, Nimrichter L, Miranda K, Martinez LR.
2013. Use of a stainless steel washer platform to study Acinetobacter baumannii adhesion and biofilm
formation on abiotic surfaces. Microbiology+ 159:2594–2604.

17. Wang X, Cole CG, DuPai CD, Davies BW. 2020. Protein Aggregation is Associated with
Acinetobacter baumannii Desiccation Tolerance. Microorg 8:343.

18. Boll JM, Tucker AT, Klein DR, Beltran AM, Brodbelt JS, Davies BW, Trent MS. 2015. Reinforcing
Lipid A Acylation on the Cell Surface of Acinetobacter baumannii Promotes Cationic Antimicrobial
Peptide Resistance and Desiccation Survival. Mbio 6:e00478 15.

19. Aranda J, Bardina C, Beceiro A, Rumbo S, Cabral MP, Barbé J, Bou G. 2011. Acinetobacter
baumannii RecA Protein in Repair of DNA Damage, Antimicrobial Resistance, General Stress Response,
and Virulence. J Bacteriol 193:3740–3747.

20. Farrow JM, Wells G, Pesci EC. 2018. Desiccation tolerance in Acinetobacter baumannii is mediated
by the two-component response regulator BfmR. Plos One 13:e0205638.

21. Farrow JM, Wells G, Palethorpe S, Adams MD, Pesci EC. 2020. CsrA supports both environmental
persistence and host-associated growth of Acinetobacter baumannii. Infect Immun
https://doi.org/10.1128/iai.00259-20.

22. Zeidler S, Müller V. 2019. The role of compatible solutes in desiccation resistance of Acinetobacter
baumannii. Microbiologyopen 8:e00740.

23. Jacobs AC, Thompson MG, Black CC, Kessler JL, Clark LP, McQueary CN, Gancz HY, Corey BW,
Moon JK, Si Y, Owen MT, Hallock JD, Kwak YI, Summers A, Li CZ, Rasko DA, Penwell WF, Honnold
CL, Wise MC, Waterman PE, Lesho EP, Stewart RL, Actis LA, Palys TJ, Craft DW, Zurawski DV. 2014.
AB5075, a Highly Virulent Isolate of Acinetobacter baumannii, as a Model Strain for the Evaluation of
Pathogenesis and Antimicrobial Treatments. Mbio 5:e01076 14.

24. Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, Huse HK, Zurawski DV,
Brittnacher MJ, Manoil C. 2015. Resources for Genetic and Genomic Analysis of Emerging Pathogen
Acinetobacter baumannii. J Bacteriol 197:2027–2035.

25. LeGrand K, Petersen S, Zheng Y, Liu KK, Ozturk G, Chen J-Y, Young GM. 2015. CsrA impacts
survival of Yersinia enterocolitica by affecting a myriad of physiological activities. Bmc Microbiol 15:31.

26. Chin CY, Tipton KA, Farokhyfar M, Burd EM, Weiss DS, Rather PN. 2018. A high-frequency
phenotypic switch links bacterial virulence and environmental survival in Acinetobacter baumannii. Nat
Microbiol 3:563–569.

27. Tipton KA, Chin C-Y, Farokhyfar M, Weiss DS, Rather PN. 2018. Role of capsule in resistance to
disinfectants, host antimicrobials and desiccation in Acinetobacter baumannii. Antimicrob Agents Ch
62:AAC.01188-18.

28. Erdős G, Pajkos M, Dosztányi Z. 2021. IUPred3: prediction of protein disorder enhanced with
unambiguous experimental annotation and visualization of evolutionary conservation. Nucleic Acids Res
49:gkab408-.

29. Elhosseiny NM, Amin MA, Yassin AS, Attia AS. 2015. Acinetobacter baumannii universal stress
protein A plays a pivotal role in stress response and is essential for pneumonia and sepsis pathogenesis.
Int J Med Microbiol 305:114–123.

30. Crowe JH, Oliver AE, Tablin F. 2002. Is There a Single Biochemical Adaptation to Anhydrobiosis?
Integr Comp Biol 42:497–503.

31. Leslie SB, Israeli E, Lighthart B, Crowe JH, Crowe LM. 1995. Trehalose and sucrose protect both
membranes and proteins in intact bacteria during drying. Appl Environ Microb 61:3592–7.

32. França MB, Panek AD, Eleutherio ECA. 2007. Oxidative stress and its effects during dehydration.
Comp Biochem Physiology Part Mol Integr Physiology 146:621–631.

33. Soares NC, Cabral MP, Gayoso C, Mallo S, Rodriguez-Velo P, Fernández-Moreira E, Bou G. 2010.
Associating Growth-Phase-Related Changes in the Proteome of Acinetobacter baumannii with Increased
Resistance to Oxidative Stress. J Proteome Res 9:1951–1964.

34. Juttukonda LJ, Green ER, Lonergan ZR, Heffern MC, Chang CJ, Skaar EP. 2018. Acinetobacter
baumannii OxyR regulates the transcriptional response to hydrogen peroxide. Infect Immun
87:IAI.00413-18.

35. Aravind L, Landsman D. 1998. AT-hook motifs identified in a wide variety of DNA-binding proteins.
Nucleic Acids Res 26:4413–4421.

36. Su L, Deng Z, Leng F. 2020. The Mammalian High Mobility Group Protein AT-Hook 2 (HMGA2):
Biochemical and Biophysical Properties, and Its Association with Adipogenesis. Int J Mol Sci 21:3710.

37. Boothby TC, Tapia H, Brozena AH, Piszkiewicz S, Smith AE, Giovannini I, Rebecchi L, Pielak GJ,
Koshland D, Goldstein B. 2017. Tardigrades Use Intrinsically Disordered Proteins to Survive
Desiccation. Mol Cell 65:975-984.e5.

38. Hesgrove C, Boothby TC. 2020. The biology of tardigrade disordered proteins in extreme stress
tolerance. Cell Commun Signal 18:178.

39. Cohan MC, Pappu RV. 2020. Making the Case for Disordered Proteins and Biomolecular
Condensates in Bacteria. Trends Biochem Sci 45:668–680.

40. Lapouge K, Schubert M, Allain FH-T, Haas D. 2008. Gac/Rsm signal transduction pathway of
gamma-proteobacteria: from RNA recognition to regulation of social behaviour. Mol Microbiol 67:241
253.

41. Cerqueira GM, Kostoulias X, Khoo C, Aibinu I, Qu Y, Traven A, Peleg AY. 2014. A global virulence
regulator in Acinetobacter baumannii and its control of the phenylacetic acid catabolic pathway. J Infect
Dis 210:46 55.

42. Weiss A, Broach WH, Lee MC, Shaw LN. 2016. Towards the complete small RNome of
Acinetobacter baumannii. Microb Genom 2:e000045.

43. Romeo T, Vakulskas CA, Babitzke P. 2013. Post-transcriptional regulation on a global scale: form
and function of Csr/Rsm systems. Environ Microbiol 15:313–324.

44. Pourciau C, Lai Y-J, Gorelik M, Babitzke P, Romeo T. 2020. Diverse Mechanisms and Circuitry for
Global Regulation by the RNA-Binding Protein CsrA. Front Microbiol 11:601352.

45. Romeo T, Babitzke P. 2018. Global Regulation by CsrA and Its RNA Antagonists. Microbiol Spectr
6.

46. Sobrero PM, Valverde C. 2020. Comparative Genomics and Evolutionary Analysis of RNA-Binding
Proteins of the CsrA Family in the Genus Pseudomonas. Frontiers Mol Biosci 7:127.

47. Renda A, Poly S, Lai Y-J, Pannuri A, Yakhnin H, Potts AH, Bevilacqua PC, Romeo T, Babitzke P.
2020. CsrA-Mediated Translational Activation of ymdA Expression in Escherichia coli. Mbio 11.

48. Horton RM, Ho SN, Pullen JK, Hunt HD, Cai Z, Pease LR. 1993. Gene splicing by overlap extension.
Methods in enzymology 217:270 279.

49. Iwase T, Tajima A, Sugimoto S, Okuda K, Hironaka I, Kamata Y, Takada K, Mizunoe Y. 2013. A
Simple Assay for Measuring Catalase Activity: A Visual Approach. Sci Rep-uk 3:3081.

50. Eng JK, Jahan TA, Hoopmann MR. 2013. Comet: An open-source MS/MS sequence database search
tool. Proteomics 13:22–24.

51. Keller A, Nesvizhskii AI, Kolker E, Aebersold R. 2002. Empirical Statistical Model To Estimate the
Accuracy of Peptide Identifications Made by MS/MS and Database Search. Anal Chem 74:5383–5392.

52. Nesvizhskii AI, Keller A, Kolker E, Aebersold R. 2003. A Statistical Model for Identifying Proteins
by Tandem Mass Spectrometry. Anal Chem 75:4646–4658.

53. Fermin D, Basrur V, Yocum AK, Nesvizhskii AI. 2011. Abacus: A computational tool for extracting
and pre-processing spectral count data for label-free quantitative proteomic analysis. Proteomics
11:1340–1345.

Figure 1. Desiccation tolerance of *A. baumannii*. (A) Comparison of AB5075, *E. coli* W3110, and *P. aeruginosa* PAO1 at 30% RH or 2% RH. (B) Desiccation tolerance of *A. baumannii* strains AB5075, ATCC17978, and ATCC19606 after 0 days (control) and 6 days of desiccation at 30% RH or 2% RH. The cell numbers represent the total number of viable cells recovered from each membrane. The data are the average of three or more biological replicates, and standard deviations are shown as the error bars.

Figure 2. Growth and desiccation tolerance of the ΔcsrA mutant (A) Comparison of wild type (WT) and the ΔcsrA mutant on TY and M9/succinate agar and in TY and M9/succinate broth. (B) Desiccation tolerance of WT, ΔcsrA mutant, ΔcsrA mutant with pMMB (empty vector), and ΔcsrA mutant with csrA expressed in trans (pAB2750) at day 0 and at day 6 of desiccation at 2% RH or at 6 days in PBS. The cell number represents the total number of viable cells recovered from each membrane. The data are the average of three or more biological replicates, and standard deviations are shown as the error bar.

Figure 3. Phenotypes of the Δ*csrA* mutant: (A) twitching motility; (B) growth of wild type (black symbol) or the Δ*csrA* mutant (red symbol) in M9/succinate in the absence (closed symbol) or presence (open symbol) of 0.5% ethanol. The data shown are the representative of each strain and condition; (C) crystal violet staining of biofilms from wild type, Δ*csrA* mutant, Δ*csrA* mutant with pMMB (empty vector), or Δ*csrA* mutant with pAB2750; and (D) catalase activity of wild type, Δ*csrA* mutant, Δ*csrA* mutant with pMMB (empty vector), or Δ*csrA* mutant with pAB2750.

Figure. 4. *A. baumannii* genes important for desiccation tolerance. (A) Genes (B) Desiccation tolerance of mutants at 2% RH. The cell number represents the total number of viable cells recovered from each membrane. The data are the average of three or more biological replicates, and standard deviations are shown as the error bar.

Figure 5. Effect of expression of ABUW_2433 (red bar) or ABUW_2337 (blue bar) gene *in trans* desiccation tolerance *A. baumannii* (A) ATCC17978 and (B) ATCC19606 after 0 days (control) and 6 days of desiccation at 2% RH. Empty vector (pMMB, black bar) was used as a control. The data are the average of three or more biological replicates, and standard deviations are shown as the error bar.

Supplementary Figure 1. Desiccation tolerance of *A. baumannii* AB5075 cells from the log-phase and stationery-phase of growth. The cell number represents a total number of viable cells recovered from each membrane. The data are the averages of three or more biological replicates, and standard deviations are shown as error bars.

Supplementary Figure 2. Complementation of desiccation phenotypes of (A) ABUW_2433::Tn and (B) ABUW_2437::Tn after 0 days (control) and 6 days of desiccation at 2% RH. Empty vector (pMMB) was used as a control. The data are the average of three or more biological replicates, and standard deviations are shown as the error bars.