1

Rampant transposition following RNAi loss causes hypermutation and antifungal drug

resistance in clinical isolates of a human fungal pathogen

Shelby J. Priest[1], Vikas Yadav[1†], Cullen Roth[2,3†], Tim A. Dahlmann[4], Ulrich Kück[4], Paul M.

Magwene[2], and Joseph Heitman[1]*

[1] Department of Molecular Genetics and Microbiology, Duke University Medical Center,
Durham, NC, USA

[2] Department of Biology, Duke University, Durham, NC, USA

[3]University Program in Genetics and Genomics, Duke University, Durham, NC, USA

[4] Allgemeine und Molekulare Botanik, Ruhr-Universität Bochum, Bochum, Germany

[†]These authors contributed equally to this work.

Short title: RNAi loss and transposition cause hypermutation in *Cryptococcus*

* Corresponding author

Email: heitm001@duke.edu (JH)

Abstract

Microorganisms survive and compete within their environmental niches and avoid

evolutionary stagnation by stochastically acquiring mutations that enhance fitness. Although

increased mutation rates are often deleterious in multicellular organisms, hypermutation can be

beneficial for microbes in the context of strong selective pressures. To explore how

hypermutation arises in nature and elucidate its consequences, we employed a collection of 387

sequenced clinical and environmental isolates of *Cryptococcus neoformans*. This fungal

pathogen is responsible for ~15% of annual AIDS-related deaths and is associated with high

mortality rates, attributable to a dearth of antifungal drugs and increasing drug resistance.

Isolates were screened for the ability to rapidly acquire antifungal drug resistance, and two

robust hypermutators were identified. Insertion of the non-LTR Cnl1 retrotransposon was found

to be responsible for the majority of drug-resistant isolates. Long-read whole-genome sequencing

revealed both hypermutator genomes have two unique features: 1) hundreds of Cnl1 copies

organized in subtelomeric arrays on both ends of almost all chromosomes, and 2) a nonsense

mutation in the first exon of *ZNF3*, a gene encoding an RNAi component involved in silencing

transposons. Quantitative trait locus mapping identified a significant genetic locus associated

with hypermutation that includes the mutant *znf3* allele, and CRISPR-mediated genome editing

of the *znf3* single-base pair nonsense mutation abolished the hypermutation phenotype and

restored siRNA production. In sum, hypermutation and drug resistance in these isolates results

from loss of RNAi combined with subsequent accumulation of a large genomic burden of a

novel transposable element in *C. neoformans*.

Introduction

Stochastic mutations and genomic rearrangements provide variation in populations for natural selection to act upon and enable evolution. However, genetic changes are a double-edged sword: too little variation can lead to evolutionary stagnation, while too much variation can lead to a lethal accumulation of deleterious mutations. Hypermutation, one extreme of this mutational spectrum, can lead to adaptation, disease, or eventual extinction if left unchecked.

Microbes are known to adopt highly mutable states that would normally be viewed as deleterious from the perspective of multicellular organisms. Studies have found that microorganisms with defects in pathways associated with maintaining genomic integrity, such as those involved in chromosome stability, DNA mismatch repair, DNA damage repair, and cell cycle checkpoints associated with recognizing DNA damage, accelerate adaptation to environmental stressors[1–3]. These defects can be beneficial in the short term, yet deleterious in the long term as mutations continue to accumulate. Defects in DNA mismatch repair resulting in increased mutation rates have been reported in fungi, including the model yeast *Saccharomyces cerevisiae*, the human pathogen *Candida glabrata*, in an outbreak strain of *Cryptococcus deuterogattii*, and in several clinical isolates of the model basidiomycete human fungal pathogen *Cryptococcus neoformans*[4–10]. Genomic stability in pathogenic *Cryptococcus* species is also significantly affected by karyotypic changes and transposable elements, both of which can mediate antifungal drug resistance[11–14].

Cryptococcus is an environmentally ubiquitous haploid basidiomycete and facultative human pathogen[15]. Approximately 95% of cryptococcal infections are attributable to the serotype A group, *C. neoformans* var. *grubii*, now known as *C. neoformans*, which is divided into four lineages: VNI, VNII, VNBI, and VNBII[16–18]. This species infects immunocompromised

individuals and accounts for ~15% of HIV/AIDS-related deaths[19]. The threat of cryptococcal

infections is exacerbated because the arsenal of antifungal drugs is limited. Amphotericin B, a

fungicidal polyene, is often used in combination with 5-flucytosine (5-FC), an antimetabolite, to

treat cryptococcal infections[20]. Unfortunately, amphotericin B and 5-FC have undesirable side

effects, and 5-FC monotherapy frequently leads to resistance[21,22]. Fluconazole is used to treat

asymptomatic patients with isolated cryptococcal antigenemia, those with disease limited to lung

nodules or central nervous system infections after clearance of cerebrospinal fluid cultures, or for

chronic maintenance therapy[20]. However, *C. neoformans* frequently develops resistance to

fluconazole via aneuploidy or mutations in the sterol biosynthesis pathway, contributing to

recurrent infections[11,23–25]. The limited number of drugs available to treat cryptococcosis,

prevalence of resistance and recurrent infections, and difficulty in developing novel antifungal

therapies combine to make *C. neoformans* drug resistance an important clinical problem.

Transposons in the C. *neoformans* H99 reference strain and the sister species

Cryptococcus deneoformans JEC21 reference strain have been characterized[13,26,27]. The genomes

of these species encode many retrotransposons with and without long-terminal repeats, known as

LTR retrotransposons and non-LTR retrotransposons, respectively, which move via a copy-and-

paste mechanism, allowing them to proliferate throughout the genome if unchecked. The most

well-characterized *Cryptococcus* LTR-retrotransposons are Tcn1 through Tcn6, which are

primarily located in centromeric regions[28]. The *C. deneoformans* JEC21 genome also encodes

three types of DNA transposons (T1, T2, and T3), as well as ~25 copies of the non-LTR

retrotransposon Cnl1 (*C. neoformans* LINE-1), which is thought to associate with telomeric

repeat sequences[13,14,29]. In the *C. neoformans* H99 genome, there are no full-length copies of

Cnl1, and DNA transposons are rare[13].

Studies have illustrated that transposon silencing in *Cryptococcus* is governed by RNAi through three primary lines of evidence: 1) siRNAs map predominantly to transposable elements, 2) RNAi mutants show increased transposon expression, and 3) spliceosomes stall on transposable element transcripts at an unusually high rate, triggering RNAi[13,30–34]. Other mechanisms thought to regulate *Cryptococcus* transposons include 5-methylcytosine DNA methylation[28,35,36] and heterochromatic marks[37]. Interestingly, the outbreak species *C. deuterogattii* is RNAi deficient because many genes encoding RNAi components are severely truncated or absent entirely[32]. This loss of RNAi has been shown to be associated with loss of all functional transposable elements, consequently shorter centromeres, and higher rates of intron retention[28,38].

Here, we identified two clinical, hypermutator *C. neoformans* isolates with significantly increased mutation rates on antifungal drug media. The majority of drug resistance in these two strains is mediated by insertions of the Cnl1 transposon into genes whose mutation confers resistance. Genetic backcrossing, quantitative trait loci mapping, and CRISPR-mediated gene editing all confirmed that a nonsense mutation in the RNAi component *ZNF3*, resulting in loss of RNAi, is the cause of hypermutation in these two strains. Small RNA sequencing confirmed the role of Znf3 in silencing Cn1l, and whole-genome sequencing revealed both hypermutator genomes encode >800 copies or fragments of Cnl1. This is the first time full-length copies of Cnl1 have been identified in *C. neoformans*, and the massive Cnl1 burden in these hypermutators is substantially higher than previously observed in any other *Cryptococcus* strain. Our results demonstrate the hypermutator phenotype described here is attributable to loss of RNAi, allowing rampant transposition of Cnl1. These transposition events lead to Cnl1 accumulation at subtelomeres and movement to novel genomic locations, which can result in drug resistance.

Results

Identification of two clinical, hypermutator *C. neoformans* isolates

To identify natural isolates of *C. neoformans* with increased mutation rates, we screened strains in the Strain Diversity Collection (SDC) for an increased ability to produce colonies resistant to various classes of antifungal drugs. The SDC contains 387 strains from all four *C. neoformans* lineages (VNI, VNII, VNBI and VNBII), including geographically diverse clinical and environmental isolates of both mating types. For each isolate in this collection, whole-genome sequencing (WGS) and phylogenetic relationships are available[18]. To screen isolates for increased mutation rates in a relatively high-throughput manner, strains were grown in liquid cultures, swabbed onto agar plates supplemented with either 5-fluorocytosine (5-FC) or a combination of FK506 and rapamycin (immunosuppressants that bind FKBP12 to form complexes that inhibit activity of calcineurin and TOR, respectively), and qualitatively evaluated for their ability to generate resistant colonies (Figure 1A)[39–42]. Strains that produced more spontaneously resistant colonies on average than the control strain H99 were categorized as hypermutator candidates. We screened 186 strains and identified 40 hypermutator candidates (Table S1). Interestingly, all but one of fourteen (93%) environmental isolates screened were identified as hypermutator candidates (as compared to only 16% (27/170) clinical isolates, *p*-value < 0.001, Fisher's exact test). Two previously identified hypermutator strains with mismatch repair defects, C23 and C45, were identified as hypermutator candidates as well[9].

We chose to focus on two clinical strains, Bt65 and Bt81, that produced the most rapamycin + FK506-resistant (R+F[R]) colonies (Figure 1A). Bt65 and Bt81 are both VNBII *MAT***a** strains isolated from different HIV-positive individuals in Botswana[18,43]. To quantify the mutation rates of Bt65 and Bt81, we performed fluctuation assays on YPD + rapamycin +

Figure 1. Hypermutation in Bt65 and Bt81 is driven primarily by the insertion of Cnl1 into *FRR1*.
(A) Generation of spontaneously resistant colonies on YPD + rapamycin + FK506 medium was utilized to identify hypermutator candidates; pictures of representative plates are shown. Strains include the phylogenetically closely related strains involved in fluctuation assay in B as well as positive (*msh2Δ*) and negative (H99) controls. **(B)** Mutation rates of closely related VNBII strains and controls on YPD + rapamycin + FK506. Bars represent the mutation rate and error bars represent 95% confidence intervals; mutation rates represent the number of mutations per cell per generation. Schematic depicts the phylogenetic relationships of all strains included in fluctuation analyses based on Desjardins et al. 2017[18]. Mutational spectra in *FRR1* in YPD + rapamycin + FK506-resistant colonies of H99, Bt65, and Bt81 as characterized by **(C)** gel electrophoresis and Sanger sequencing of *FRR1* PCR products. MicroINDELs are defined as insertions or deletions < 50 bp. All mutations are relative to the appropriate rapamycin + FK506-sensitive parental strain.

FK506, YNB + 5-fluoroorotic acid (5-FOA), and YNB + 5-FC media. Both Bt65 and Bt81

produced significantly higher mutation rates on YPD + rapamycin + FK506 compared to H99 as

well as eleven of the most closely phylogenetically related strains (Figure 1B). On 5-FC, only the

strain NRHc5014.ENR and the KN99α *msh2Δ* positive control had significantly higher mutation

rates compared to H99 (Figure S1A); on 5-FOA medium (Figure S1B), only KN99α *msh2Δ*

produced a significantly higher mutation rate.

A recent study illustrated how incubation at an elevated temperature of 37°C results in

increased mutation rates due to transposon mobilization in the closely related species *C.*

deneoformans[14]. To determine if elevated temperature contributed to hypermutation in Bt65 and

Bt81, we concurrently grew these strains as well as wild-type H99 and *msh2Δ*, *ago1Δ*, and

rdp1Δ genetic deletion mutants in the H99 genetic background overnight at 30°C and 37°C, and

then performed a fluctuation analysis on YPD + rapamycin + FK506 medium. Fluctuation

analysis revealed H99, Bt65, and Bt81 had lower mutation rates when grown overnight at 37°C

compared to growth at 30°C (Figure S2). Interestingly, all of the genetic deletion mutants

showed increased mutation rates after growth at 37°C (Figure S2). These results suggest that

unlike *C. deneoformans*, growth at 37°C reduces mutation rates in wild-type *C. neoformans*

strains and does not contribute to or exacerbate hypermutation in Bt65 and Bt81.

Characterization of mutation spectra in *C. neoformans* hypermutator strains

After quantifying the mutation rates of Bt65 and Bt81 strains on media with various

antifungal drugs, we investigated the types of mutations conferring resistance to the combination

of rapamycin and FK506. PCR amplification of the *FRR1* gene (which encodes FKBP12, the

shared target of rapamycin and FK506, and the only gene in which mutations confer resistance to

both FK506 and rapamycin) followed by gel electrophoresis revealed the expected wild-type

PCR product size (~1.2 kb) for all but two (35/37) H99 R+F^R colonies; the remaining two

produced PCR products smaller than expected, indicative of deletions (Figure S3A). In contrast,

large insertions of various sizes were observed in the majority of the Bt65 and Bt81 R+F^R

colonies (62/77 and 36/37 independent colonies, respectively) (Figure 1C, S3B, C). Only one

resistant colony derived from a non-hypermutator strain, Bt84, (1/10 independent colonies) had

an insertion in *FRR1*. No insertions in *FRR1* were observed in any of the other closely related or

control strains.

We subsequently sequenced *FRR1* in H99, Bt65, Bt81, and Bt84 R+F^R colonies to

determine the genetic changes responsible for the larger PCR products (Figure 1C). In 37 R+F^R

colonies of the H99 control strain, SNPs in *FRR1* were largely responsible for resistance (57%,

21/37 colonies), while resistance in the remaining colonies was attributable to small

insertions/deletions (microINDELs; 38%, 14/37) or large deletions (5%, 2/37). Conversely, in

the hypermutator isolate Bt65, insertions of the non-LTR retrotransposon Cnl1 were responsible

for the majority of rapamycin + FK506 resistance (77.4%, 24/31). Rapamycin + FK506

resistance in the remaining Bt65 colonies was either due to SNPs (19.4%, 6/31) or microINDELs

(3.2%, 1/31). In all sequenced PCR products from R+F^R colonies of Bt81, Cnl1 insertions were

responsible for resistance (27/27 colonies). Cnl1 insertions in Bt65 and Bt81 ranged from 54 bp

to ~3500 bp, and this range in transposon sizes is a common characteristic of non-LTR

retrotransposons. The single insertion observed in *FRR1* in the R+F^R colony of Bt84 had no

homology with any annotated *Cryptococcus* transposons but was identified as a repetitive

element by RepeatMasker and shared minor homology with a Copia-58 BG-I transposable

element.

The 5-FC- and 5-FOA-resistant colonies of Bt65, Bt81, and H99 were similarly

characterized to determine the sources of resistance to antifungal drugs with different

mechanisms of action. Resistance to 5-FOA is conferred by mutations in the *URA3* or *URA5*

genes of the uracil biosynthesis pathway[44,45]. Among the subset of H99, Bt65, and Bt81 5-FOA^R

colonies sequenced, mutations were only identified in *URA5*. In almost all colonies, resistance

was conferred by SNPs or INDELs, and only one Cnl1 insertion event was identified in a Bt81 5-

FOA^R colony (Figure S3D). We also PCR amplified genes in which mutations are known to

confer resistance to 5-FC, including *FUR1* and *UXS1*[46]. Only two of the 5-FC^R isolates analyzed

from H99, Bt65, and Bt81 produced a PCR product larger than expected for all assessed genes,

and subsequent sequencing revealed a Cnl1 insertion in *FUR1* in two independent 5-FC^R Bt81

isolates (Figure S3E).

Analysis of the Cnl1 insertions observed to confer resistance to rapamycin + FK506, 5-

FC, and 5-FOA revealed Cnl1 preferentially inserts at guanine- and cytosine-rich regions of

target genes, a known property of this element[27]. Target site duplication sequences flanking Cnl1

insertions were not present in many instances, but when present, ranged from 1 to 12 bp in

length. Cnl1 insertions ranged greatly in size, from 25-bp fragments to full-length Cnl1 copies

(3,494 bp). The smallest Cnl1 insertion (25 bp) was followed immediately by a 59-bp deletion in

FRR1. Cnl insertions in the *FRR1* gene were observed in both orientations (5' to 3' and 3' to 5'

relative to *FRR1* transcription). Of the 51 characterized Cnl1 *FRR1* insertions, 27 were in the 5'

UTR, and all but one were in the same orientation as *FRR1* transcription, 23 were in exons (7

oriented 5' to 3', 16 oriented 3' to 5'), and one insertion was in an intron of *FRR1* in the 3' to 5'

orientation, potentially disrupting splicing or transcription.

QTL mapping identifies loci that significantly contribute to the hypermutator phenotype

To determine the genetic cause of the hypermutator phenotype and rampant transposition

in Bt65 and Bt81 and to determine the genetic consequences of this phenotype, we conducted

quantitative trait locus (QTL) mapping. For this purpose, a total of 165 basidiospores were

dissected from a genetic cross between Bt65 *MAT***a** and an H99 *crg1Δ MAT*α mutant with an

enhanced mating phenotype, and 47 F_1 progeny germinated (28%).

Twenty-eight Bt65**a** x H99α F_1 progeny were selected for fluctuation analysis and whole-

genome sequencing. Aligning the paired-read Illumina sequencing data from the 28 F_1 progeny

identified 215,411 bi-allelic SNPs that were utilized for QTL mapping. For 24 of the segregants

as well as for the Bt65 and H99 *crg1Δ* parental strains, the mutation rate on rapamycin + FK506

medium served as the phenotype for association tests (Figure 2A). Across the 14 chromosomes

and bi-allelic SNP sites, two QTL with large effect (heritability = 64%) were identified at

approximately 919-1,120 kb on Chromosome 3 and 987-1,193 kb on Chromosome 11 (Figure

2B and S4). Analysis of these QTLs revealed that the SNPs in each QTL were co-segregating

and that they shared the same distributions of phenotype scores (Figure S4 and S5). The borders

of the QTL spanning Chromosomes 3 and 11 were determined by calculating 95% confidence

intervals and examining recombination break points along each chromosome. Interestingly, these

two QTLs span the chromosomal translocation between Chromosomes 3 and 11 that is unique to

H99 (Figure S4, S5, S6). We thus treated these QTLs as the same QTL for subsequent analysis.

Within the QTL there are a total of 108 and 85 genes along Chromosome 3 and

Chromosome 11, respectively, and for 82 and 77 of these genes (respectively), the published

annotation and SNP data was used to characterize differences in predicted protein sequence and

expected protein lengths between the H99 and Bt65 parental strains (Figure S4 and Table S2).

Figure 2. QTL analysis of hypermutator phenotype. (**A**) Quantification of mutation rates on YPD + rapamycin + FK506 medium – sorted smallest to largest, left to right – for F_1 progeny and parental strains, H99 *crg1Δ* and Bt65. Inheritance of the Bt65 *znf3* allele or H99 *crg1Δ ZNF3* allele in F_1 progeny is indicated above mutation rates. Colored bar plots and vertical black lines depict the mean mutation rate and associated 95% confidence intervals (CI) per segregant. Mutation rates represent the number of mutations per cell per generation. (**B**) Manhattan plot showing the strength in association (y-axis) between bi-allelic SNPs and hypermutator phenotype, across the 14 chromosomes (x-axis). Colors separate SNPs across chromosomes. The permutation-based significance threshold ($\alpha = 0.01$) is depicted with a horizontal dashed line. (**C**) Predicted *ZNF3* gene and Znf3 protein models in H99 and Bt65. A grey horizontal bar depicts the gene body in the upper panel, and larger grey rectangles represent exons; the gene is depicted 5' to 3' and is 5417 nt in length. The locations of SNPs differing

between Bt65 and H99 are shown by vertical black rungs along the gene model. Amino acids specified by mRNA codons in the indicated region of *ZNF3* exon 1 (nucleotides 364 to 411) are shown for H99 and Bt65 to illustrate the effect of the C to T mutation (nucleotide 409) predicted to cause a nonsense mutation in Bt65. The bottom panel depicts the predicted impact of the nonsense mutation on the Znf3 protein in Bt65. White rectangles along the protein schematic depict the three C2H2-type zinc finger domains of Znf3.

Among these, 71 and 60 genes along Chromosome 3 and Chromosome 11, respectively, have at

least one predicted nonsynonymous change in protein sequence, seven of which harbor a

predicted nonsense (i.e. stop-gain) or stop-loss mutation. One of these genes is *ZNF3*

(CNAG_02700), which encodes a C2H2 type zinc finger protein with three zinc finger domains.

Znf3 was previously identified as an RNAi silencing component that localizes to P-bodies and

whose mutation results in increased expression of transposable elements[32,33]. *ZNF3* is located on

Chromosome 3 and has a SNP – C to T – within the first exon in the Bt65 genetic background,

which is predicted to cause a nonsense mutation, severely truncating Znf3 from 1,561 amino

acids to only 96 amino acids (Figure 2C). In addition, this nonsense mutation may also result in

nonsense-mediated mRNA decay of the mutant *znf3* mRNA. Based on the publicly available

whole-genome sequencing of all isolates in the SDC, the *znf3* nonsense mutation in exon 1 is

unique to Bt65 and Bt81 and not present in any other strain. Another gene of known function

within the QTL encodes a long-chain acyl-CoA synthetase (CNAG_01836, Chromosome 11)

and a SNP – G to A – within the last exon of this gene is predicted to cause an early nonsense

mutation in the Bt65 background (Figure S7). Given the dramatic difference in the predicted

protein length of *ZNF3* between the H99 and Bt65 parental alleles (relative to other genes in this

QTL with predicted stop-loss or nonsense mutations), and previous studies demonstrating the

role of Znf3 in RNAi and transposon silencing, we hypothesized *ZNF3* could be the quantitative

trait gene (QTG) and the SNP leading to the predicted stop gain in the first exon could be the

quantitative trait nucleotide (QTN) underlying the hypermutation phenotype[32,33].

Few Bt81 F$_1$ progeny display a hypermutator phenotype

In addition to generating and analyzing Bt65**a** x H99α *crg1*Δ F$_1$ progeny, 42 F$_1$ progeny

were derived from a genetic cross between the other hypermutator strain, Bt81**a,** and H99α

*crg1*Δ. The *ZNF3* alleles of all 42 Bt81 F$_1$ progeny were sequenced to determine whether they

had inherited the non-functional *znf3* allele from Bt81 or the functional *ZNF3* allele from H99.

Of the 42 F$_1$ progeny, only four inherited the mutant *znf3* allele from Bt81, a significantly lower

number than would be expected based on Mendelian inheritance patterns (chi-square test, *p*-

value < 0.01). The four progeny with non-functional Bt81 *znf3* alleles had the highest mutation

rates of 18 F$_1$ progeny that were analyzed (Figure S8). The mutation rates for three of the four

znf3 progeny, however, were not significantly higher than the Bt81 F$_1$ progeny with functional

the *ZNF3* allele and were not as high as would be expected based on the results from the Bt65**a** x

H99α F$_1$ progeny (Figure 2A).

Cnl1 elements are organized into subtelomeric arrays in hypermutator genomes

For all strains in the SDC, including Bt65 and Bt81, only short-read WGS data was

available[18]. Because of the known difficulties in assembling repetitive elements, such as Cnl1,

with short-read sequencing data, we conducted long-read WGS with the Oxford Nanopore

Technologies MinION sequencing platform to generate more complete assemblies for Bt65,

Bt81, and two of the most closely phylogenetically related non-hypermutator strains, Bt89 and

Bt133. With the long-read sequencing data, we were able to assemble chromosome-level

genomes for all four strains. In the assemblies, we observed the known chromosomal

translocation between Chromosomes 3 and 11 unique to H99[47] and identified a translocation

between Chromosomes 1 and 13 unique to Bt65 and Bt81 (Figure S6). These two gross

chromosomal rearrangements explain the relatively low germination frequency (28%) of Bt65**a** x

H99α F_1 progeny because each translocation should decrease germination by ~50%.

Analysis of the genomes of Bt65 and Bt81 revealed large arrays of the Cnl1 transposon at

all but one end of each of the 14 linear chromosomes (27/28 subtelomeric regions in Bt65 and

28/28 in Bt81) (Figure 3A, 3B). The assembled Cnl1 arrays (defined as ≥ 2 Cnl1 copies) in Bt65

and Bt81 range from 5 kb to 80 kb in length. These highly repetitive arrays made it difficult and,

in some instances, impossible to confidently assemble telomeric repeat sequences at the ends of

each Bt65 and Bt81 chromosome. Using manual telomere extension via read mapping, we were

able to identify telomere repeats at only 20 chromosome ends in Bt65 and 13 in Bt81. In contrast

to Bt65 and Bt81, genome assemblies for Bt89 and Bt133 were assembled with telomere repeats

on all 28 chromosome ends without any manual extension (Figure S6). In these assemblies, some

telomeres had no copies of Cnl1 while others had Cnl1 arrays up to 30 kb in length (Figure S9).

Further analysis revealed the Bt65 genome harbors at least 414 fragments of Cnl1,

including 105 full-length copies, while the Bt81 genome appears to encode even more Cnl1

elements, with at least 449 fragments, including 147 full-length copies (Table 1). It is important

to note that due to incomplete ends for most chromosomes, it is likely Bt81 and Bt65 encode

additional copies of Cnl1 that were not assembled. The presence of long Cnl1 arrays in Bt65 and

Bt81 was surprising because the *C. neoformans* H99 reference strain encodes only 22 fragments

of Cnl1 and no full-length copies, and therefore, *C. neoformans* was not thought to harbor

functional Cnl1 elements (Figure 3C) (Table 1).

Apart from the subtelomeres, retrotransposons in *Cryptococcus* are also enriched at

centromeres, specifically the LTR retrotransposons Tcn1-Tcn6[26,28,47]. The changes in Cnl1

Figure 3. Retrotransposon content in the genomes of H99, Bt65, and Bt81. Distributions of the Tcn1 through Tcn6 LTR-retrotransposons and the Cnl1 non-LTR retrotransposon in subtelomeric and centromeric regions of **(A)** Bt65, **(B)** Bt81, and **(C)** H99 genomes depicted in Figure S6. In Bt65 and Bt81, 80 kb of subtelomeric regions are displayed, and 50 kb subtelomeric regions are displayed for H99 to show the full distribution of subtelomeric Cnl1 elements. Subtelomeric arrays of Cnl1 are depicted at the end of each chromosome in Bt65 and Bt81, while only 7 Cnl1 elements are localized subtelomerically in H99. Shading corresponds to fragments of the Cnl1 elements, and gene arrowheads indicate the direction of transcription for all retrotransposons.

303   transposon content in Bt65 and Bt81 along with a previous study establishing a link between loss

304   of RNAi and centromere length[28] motivated us to characterize the centromeres in Bt65, Bt81,

305   Bt89, and Bt133. Identification of centromeres in these four isolates revealed shorter centromeres

306   on average compared to H99: 40.3 kb in Bt65, 40 kb in Bt81, 35.6 kb in Bt89, and 41.2 kb in

Table 1. Cnl1 burden in H99, hypermutator strains, related non-hypermutator strains, and six Bt65 x H99 _crg1Δα_ F$_1$ progeny based on Nanopore sequencing data.

Strain	Hypermutator Status	Total Cnl1 burden (>50 bp)	Full-length Cnl1 copies (>99% in length)
H99	Non-hypermutator	22	0
Bt65	Hypermutator	414	105
Bt81	Hypermutator	449	147
Bt89	Non-hypermutator	261	24
Bt133	Non-hypermutator	246	23
Progeny 2	Hypermutator	212	30
Progeny 8	Hypermutator	172	40
Progeny 14	Non-hypermutator	296	68
Progeny 18	Non-hypermutator	321	88
Progeny 20	Non-hypermutator	425	136
Progeny 34	Hypermutator	187	41

Bt133, compared to an average length of 47.3 kb in H99. However, this difference did not reach

statistical significance (ANOVA, _p_-value = 0.153, Figure S10 and Table S3). Many centromeres

in the assessed VNBII isolates had undergone numerous rearrangements and several inversions

relative to one another (Figure S11). Centromeric alterations have also previously been observed

in _C. neoformans_ genetic deletion mutants lacking the canonical RNAi components Ago1 and

Rdp1[28]. Combined, these analyses suggest that while Cnl1 is more abundant in the

hypermutators Bt65 and Bt81 as well as the non-hypermutators Bt89 and Bt133, other

retrotransposons are not substantially increased in number compared to H99 (Figure 3 and S9).

Characterization of H99 _crg1Δ_ x Bt65 F$_1$ progeny genomes reveals invasion of Cnl1

elements into naïve telomeres

Expression of transposable elements, including Cnl1, has been shown to be upregulated

during sexual reproduction in _C. neoformans_[30,32,34]. To investigate how increased expression of

Cnl1 during mating impacts the genome, six of the 28 progeny utilized for QTL mapping were

also selected for long-read whole-genome sequencing on the nanopore MinION sequencing

platform: three hypermutator progeny that inherited the Bt65 *znf3* allele (P2, P8, and P34) and

three non-hypermutator progeny that inherited the H99 *ZNF3* allele (P14, P18, and P20).

Nanopore sequencing identified recombination points across the genomes of the progeny,

providing information on which genomic regions were inherited from either parent and

confirming these were F$_1$ genetic recombinants (Figure 4).

Figure 4. Genetic recombination sites and Cnl1 distribution in Bt65 x H99 F$_1$ progeny.
Recombination sites along each of the 14 chromosomes for the six Bt65**a** x H99α F$_1$ progeny for which
long-read whole-genome sequencing was conducted. Genomic loci depicted in blue were inherited from
the H99 parent, and orange genomic loci were inherited from the Bt65 parent. Cnl1 elements throughout
the F$_1$ progeny and parental genomes are indicated by black arrowheads in the upper panel. Centromeres
are indicated by dark blue boxes in only the parental genomes. Hypermutator F$_1$ progeny are indicated
with asterisks, and the *ZNF3* locus is indicated in each strain with a red arrowhead. Regions enlarged
below illustrate Cnl1 subtelomeric arrays on several chromosomes and depict examples of Cnl1 array
expansion (e.g. Chromosome 4, P18), contraction (e.g. Chromosome 1, P14), and invasion of naïve H99
subtelomeres (e.g. Chromosome 1, P8). Telomeric repeat sequences are indicated by black half circles
only in the enlarged panels.

Surprisingly, the genome assemblies for the three progeny with the functional *ZNF3*

allele inherited from H99 appear to encode more full-length Cnl1 elements and fragments

compared to the three *znf3* progeny. However, of the three *znf3* progeny, telomeric repeat

sequences were only identified at the end of two chromosomes (out of a possible 84 telomeric

ends across the three progeny, 2/84 or 2%): on one arm of Chromosome 3 in P8 and one end of

Chromosome 12 in P34 (Figure S12). This is in contrast to the 31 telomeres that were accurately

assembled across the three *ZNF3* progeny (31/84 or 37%) (Figure S12). The smaller number of

telomeres identified in the *znf3* progeny assemblies suggest there might be more Cnl1 elements

that were not accurately included, similar to the assemblies for the hypermutators Bt65 and Bt81.

Therefore, the Cnl1 burden quantified and presented in Table 1 might not accurately capture the

entire Cnl1 burden of these strains.

A previous study found *ZNF3* to be a haploinsufficient gene because no progeny isolated

from a *ZNF3* x *znf3Δ* cross showed evidence of sex-induced RNAi-mediated silencing[32]. This

haploinsufficiency allowed us to analyze Cnl1 dynamics both hypermutator and non-

hypermutator progeny and identified two additional phenomena. First, nearly all Cnl1 arrays in

the progeny show signs of expansion and contraction relative to the Bt65 parental genome,

suggesting these elements are either highly mobile during sexual reproduction, undergoing high

levels of recombination, or both. Additionally, combined analysis of subtelomeric region

inheritance patterns and Cnl1 arrays revealed Cnl1 elements are capable of invading naïve

subtelomeric regions inherited from the H99 parent, i.e. regions that previously had no Cnl1

elements/fragments (Figure 4, Figure S12). In the three *znf3* progeny, 65% (28/43) of the naïve

telomeric regions inherited from H99 acquired Cnl1 copies and arrays in many cases. In the three

ZNF3 progeny, 81% (35/44) of naïve telomeric regions inherited from H99 now had Cnl1

elements. Overall, both *ZNF3* and *znf3* F_1 progeny inherited roughly equivalent numbers of

telomeric regions from either parent and Cnl1 invaded a majority of the naïve H99 telomeres.

356	**Complementation of the nonsense mutation in *ZNF3* significantly lowers the mutation rate**

357	**and restores production of siRNAs**

358	 All of the available evidence thus far suggested the nonsense mutation in *ZNF3* unique to

359	Bt65 and Bt81 is responsible for the hypermutation phenotype, possibly due to compromised

360	RNAi silencing of Cnl1 elements. To test this hypothesis, we used CRISPR-mediated gene

361	editing to restore the functional *ZNF3* allele in Bt65. Gene editing was achieved with the

362	transient CRISPR-Cas9 coupled with electroporation (TRACE) system and the utilization of a

363	functional *ZNF3* allele from a closely related strain, Bt133, such that only the SNP responsible

364	for the nonsense mutation would be changed to the wild-type nucleotide (found in H99 and all

365	SDC isolates except Bt65 and Bt81)[48]. Following transformation and selection, we identified two

366	Bt65 transformants that had successfully integrated a single copy of the Bt133 *ZNF3* allele at the

367	endogenous *znf3* locus, Bt65+*ZNF3-1* and Bt65+*ZNF3-2*. These two independent Bt65+*ZNF3*

368	transformants were subjected to fluctuation analysis to determine if changing the single SNP

369	responsible for the *znf3* nonsense mutation restored the mutation rate to a wild-type level. On

370	YPD + rapamycin + FK506 medium, both transformants had significantly lower mutation rates

371	than Bt65, similar to those observed in H99 and Bt133 (Figure 5A).

372	 We next sequenced the sRNA repertoires of the Bt65+*ZNF3* isolates as well as Bt65,

373	Bt81, and an H99 *znf3Δ* mutant; H99 and an H99 *rdp1Δ* mutant served as positive and negative

374	controls, respectively. Analysis of the size distribution of sRNAs aligning to the *C. neoformans*

375	genome showed that Bt65, Bt81, and H99 *znf3Δ* displayed profiles similar to that of the

376	canonical RNAi mutant H99 *rdp1Δ* because they lacked the characteristic siRNA peak at 21-24

377	nt (Figure 5B). sRNA analysis also revealed that *ZNF3* complementation in Bt65 restored the 21-

378	24 nt sRNA peak. We also characterized the 5' nucleotide identity of sRNAs of different lengths

Figure 5. *ZNF3 complementation in Bt65 significantly reduces mutation rates and restores siRNA production.* **(A)** Mutation rates of H99, Bt65, Bt133, and the two independent *ZNF3* complementation mutants, Bt65 + *ZNF3-1* and Bt65 + *ZNF3-2*, on YPD + rapamycin + FK506 medium. Bars represent mutation rate (number of mutations per cell per generation) and error bars represent 95% confidence intervals. **(B)** Size distributions of sRNA reads from each indicated strain. Dashed vertical lines indicate the 21 to 24 nucleotide size range, the characteristic sizes of siRNAs produced by the RNAi pathway. **(C)** Proportion of sRNA reads (y-axis) with the indicated 5' nucleotide identity (color of stacked bar) at each sRNA read size (x-axis). siRNAs produced by the RNAi pathway characteristically have a 5' uracil nucleotide. **(D)** Quantification of sense and antisense sRNAs from Bt65, Bt65 + *ZNF3-1*, and Bt65 + *ZNF3-2* aligning to an array of subtelomeric Cnl1 elements on Chromosome 1 of Bt65. Transposable elements along the chromosome are indicated by dark grey boxes, while intergenic regions are light grey.

and found only H99 and the two Bt65+*ZNF3* transformants had a peak of 21-24 nt sRNAs with a

predominance for a 5' U, another characteristic of siRNAs produced by the RNAi pathway

(Figure 5C).

To determine how complementation of *ZNF3*, and thus restoration of RNAi, specifically

impacted silencing of Cnl1, we quantified sRNAs aligning to Cnl1 elements across the Bt65

genome (Table S4). Relative to Bt65, the normalized expression of sRNAs corresponding to

Cnl1 were increased 5.6-fold in H99 (a strain with no full-length Cnl1 elements and only 22

Cnl1 fragments), 10.6-fold in Bt65+*ZNF3-1*, and 12.8-fold in Bt65+*ZNF3-2* (Table S4). To

demonstrate the marked difference in Cnl1 sRNAs in the two Bt65+*ZNF3* transformants

compared to Bt65, sRNAs were plotted along a telomeric Cnl1 array on Chromosome 1 in Bt65

(Figure 5D). These results suggest that changing the single nucleotide responsible for the

nonsense mutation in *ZNF3* back to the wild-type nucleotide found in closely related strains as

well as H99 was able to successfully restore RNAi in Bt65. This result also provides evidence

that all other RNAi components have been largely maintained in a functional form despite the

loss of RNAi activity in Bt65.

Discussion

Transposable element mobilization can alter gene expression, gene function, and even genomic stability. In this study, we identified two clinical, hypermutator isolates of *C. neoformans*, Bt65 and Bt81, with a massive accumulation of the non-LTR retrotransposon Cnl1 at all but one subtelomeric loci of all fourteen chromosomes. Furthermore, the Cnl1 element was capable of inserting in non-subtelomeric regions of the genome, resulting in resistance to diverse classes of antifungal drugs. These findings were unprecedented as *C. neoformans* is thought to be an RNAi-proficient species[13,49]. Additionally, *C. neoformans* isolates were also thought to lack full-length copies of Cnl1, and only the sister species *C. deneoformans* has been shown to harbor full-length Cnl1 elements capable of mobilization[13,27]. These findings highlight the importance of intraspecific diversity at both the genotypic and phenotypic levels.

Following isolation, phenotyping, and genotyping of the Bt81**a** x H99α F_1 progeny, a significantly smaller number of progeny inherited the Bt81 *znf3* allele than expected. These results were surprising giving the approximately 1:1 inheritance pattern of the *ZNF3* allele from either parent in the Bt65**a** x H99α F_1 progeny. Based on long-read whole-genome sequencing, Bt81 likely has a substantially higher burden of Cnl1 than Bt65. Previous studies have shown that expression of transposons, including Cnl1, is significantly upregulated during sexual reproduction in RNAi mutants, such as H99 *znf3*Δ[30,32–34]. The higher Cnl1 burden in Bt81 combined with RNAi-deficiency and transposon upregulation during mating could lead to an increased frequency of deleterious Cnl1 insertions in progeny lacking *znf3*, and thus a biased inheritance pattern of *ZNF3*. It is also possible that a higher Cnl1 burden favored the selection of a suppressor mutation in the Bt81 F_1 progeny. Additionally, *znf3*Δ mutants are the only *C. neoformans* RNAi mutants studied thus far in which progeny from unilateral genetic crosses (i.e.

417   crosses in which only one parent lacks *ZNF3*) exhibit complete loss of RNAi-mediated

418   silencing[32], and thus loss of *ZNF3* results in haploinsufficiency. Overall, the unequal *znf3*

419   inheritance patterns in Bt81 F_1 progeny indicate a sufficiently high burden of Cnl1 elements may

420   be deleterious during sexual reproduction.

421   In Bt65, QTL mapping and genetic complementation demonstrated the hypermutator

phenotype was caused by a single SNP in the RNAi gene *ZNF3*. Changing the *znf3* nonsense

mutation to the nucleotide found in the laboratory reference strain as well as phylogenetically

closely related strains lowered the mutation rate to a wild-type level and restored RNAi function,

including expression of siRNAs corresponding to Cnl1, thus confirming the role of Znf3 in Cnl1

regulation. Although *ZNF3* complementation restored the mutation rate and siRNA production,

the identified QTL which spans Chromosomes 3 and 11 accounted for only 64% of the

hypermutator phenotype. This may suggest the existence of additional contributing loci of small

effect, such as a mutation in another RNAi component that results in a partial loss of function,

although the mapping population used here is under powered to detect such small-effect loci. It is

also possible that variation in Cnl1 burden across the F_1 progeny accounts for the remainder of

the genetic loci that contribute to the hypermutator phenotype. However, because short-read

sequencing data was utilized in variant calling, these Cnl1 loci may have been difficult to map in

the F_1 progeny or failed to pass quality criteria and were subsequently filtered out during

preprocessing of the genetic data used in QTL analysis.

436   Despite having the wild-type nucleotide in the first exon of *ZNF3*, where Bt65 and Bt81

have a nonsense mutation, both Bt89 and Bt133 (two of the most closely related strains) have a

substantial accumulation of subtelomeric Cnl1 arrays relative to the laboratory reference strain

H99, which has none. The considerable Cnl1 burden in Bt89 and Bt133 indicates that they may

descend from an ancestral strain that had lost RNAi function, but which was subsequently

restored, such as through a genetic cross in which they inherited a wild-type *ZNF3* allele. The

bias against *znf3* inheritance in Bt81 F_1 progeny and the lower but still impressive Cnl1 burden in

Bt89 and Bt133 potentially illustrate a natural example of how *C. neoformans* genomes have

struck a balance in their mutational capability, switching between high mutational capacities

during times of RNAi loss, and genomic stability when RNAi is restored, all via one SNP in only

one key RNAi gene. The expansion and contraction of Cnl1 arrays as well as the ability of Cnl1

to invade naïve subtelomeric regions inherited from the H99 parent in the Bt65 F_1 progeny

genomes was also exceptional. The observed Cnl1 subtelomeric dynamics mirror those observed

for the MoTeR transposons of the fungal plant pathogen *Magnaporthe oryzae*, which were also

recently shown to localize to highly dynamic subtelomeric arrays[50]. It is possible the Cnl1

subtelomeric arrays identified here could eventually overcome the requirement for telomerase, as

in the example of *Drosophila* telomeres, in which the functions of telomerase have been

supplanted by a telomeric retrotransposon[51,52].

The finding that only a single SNP rendered the RNAi pathway non-functional in Bt65,

and that no additional obvious mutations had occurred in other genes that might disrupt RNAi

function, suggest Bt65 represents an intermediate evolutionary stage and could illustrate the

natural consequences of relatively recent RNAi loss. Further analysis of the consequences of

RNAi loss in Bt65 (and potentially Bt81) through experimental evolution or gene regulation

analyses could shed light on the short-term consequences of RNAi loss at the genomic and

phenotypic levels. Bt65 could thus serve as an interesting intermediate comparator between

RNAi-proficient *C. neoformans* isolates and the closely related RNAi-deficient species *C.*

deuterogattii[7,32]. Instances of relatively recent loss of RNAi have also been observed in a natural

Caenorhabditis elegans isolate, which has a large deletion in a RIG-I homolog required for

RNAi and was shown to be infected with an RNA virus[53,54]. Unlike the identified *C. elegans*

virus-infected strain and several other RNAi-deficient fungal species, such as *Saccharomyces*

cerevisiae, *Ustilago maydis*, and several *Malassezia* species, we were unable to identify a

dsRNA virus in either of the two *C. neoformans* hypermutator strains identified here (see

Materials and Methods)[49,55]. It is possible though, that the hypermutators harbor other types of

mycoviruses (ssRNA, for example) that we were unable to detect through our approaches or that

the mycovirus was cured by common microbiological isolation practices[55].

The identification of this hypermutator phenotype in natural *C. neoformans* clinical

isolates has important implications for antifungal drug resistance and potentially other adaptive

consequences. Here, we showed Cnl1 insertion could confer resistance to diverse classes of

antifungal drugs, including the clinically relevant antifungal 5-FC. Insertion of Cnl1 into other

genes, particularly those involved in the sterol biosynthesis pathway, could confer resistance to

amphotericin B and fluconazole, the only other antifungal drugs effective for *C. neoformans*

treatment[23,56]. This mechanism of drug resistance also has interesting implications for a novel

antifungal approach that utilizes dsRNA to initiate RNAi silencing in fungal plant pathogens[57].

The effects of Cnl1 insertion at non-coding loci, such as promoters and 3' untranslated regions,

could also impact overall genomic stability or alter gene expression to have important

phenotypic implications for virulence, similar to the effects of the *Ac/Ds* elements of maize, the

first transposable elements discovered[58]. Alterations in gene expression might also confer

resistance to drugs for which resistance cannot be gained through loss of function mutations.

Even if full resistance isn't acquired, altered gene expression could contribute to antifungal drug

tolerance, like the tolerance observed in *Candida albicans*, which contributes to persistent

infections in immunocompetent patients[59,60].

At this stage it is difficult to know how Cnl1 transposition and accumulation may be

selected for or against over time. The subtelomeric arrays identified in Bt65 and Bt81 may

undergo cycles of amplification and recombination-mediated contraction allowing them to

exploit Cnl1 mutagenesis when under stress, similar to the retrotransposons replication cycles

observed in some plants[61,62]. Maintaining an RNAi-deficient background could also be adaptive

in the context of viral infection, as has been shown in yeast harboring the killer virus, which

outcompete neighboring uninfected strains, and in mice harboring latent herpesvirus, which are

protected from the bacterial pathogens *Listeria monocytogenes* and *Yersinia pestis*[63,64].

Conversely, the mutational impact of Cnl1 mobilization could be highly deleterious over the long

term and therefore may not represent a massive contributor to the rise of drug resistance. Natural

selection could either select for reversion to a functional RNAi-pathway through mutation of the

znf3 nonsense mutation or instead preserve loss of RNAi and eliminate all full-length

transposable elements, as has been observed in *C. deuterogattii*[28]. Future research on the

potential for Cnl1 insertion to mediate resistance to amphotericin B and fluconazole, and the

impact of hypermutation due to Cnl1 mobilization on *in vivo* drug resistance, adaptive potential,

and genomic stability over time will be of great interest.

Materials and Methods

Strains and growth

The *C. neoformans* strains described in this study are listed in Table S5. Strains were

stored at -80°C in liquid yeast extract peptone dextrose (YPD) supplemented with 15% glycerol.

Strains were inoculated on YPD agar plates, grown for three days at 30°C, and maintained at

4°C. Due to the hypermutator phenotypes associated with several of the strains in this study,

strains were not maintained on YPD agar plates for routine use for more than two weeks; fresh

cells from frozen glycerol stocks were inoculated to YPD agar plates as needed.

Screening for hypermutator candidates

Assays for the emergence of resistance (papillation assays) were conducted as previously

described[65]. In brief, ten independent overnights per strain were grown overnight at standard

laboratory conditions in 5mL liquid YPD medium. Cultures were then spun down, washed, and

concentrated in 2 mL dH$_2$O. Each culture was swabbed to a quadrant of either YPD + 100 ng/mL

rapamycin + 1 µg/mL FK506 agar medium or YNB + 100 µg/mL 5-fluorocytosine agar medium.

YPD + rapamycin + FK506 plates and YNB + 5-fluorocytosine plates were incubated for up to

seven days at 37°C and 30°C, respectively. Fisher's exact probability test was used to determine

if the associations environmental isolates and the hypermutator phenotype was statistically

significant using the VassarStats online software (http://vassarstats.net).

Fluctuation assays

Fluctuation assays were conducted as previously described[65]. Briefly, ten independent

overnights of each strain were grown overnight in 5 mL liquid YPD medium at 30°C. Cultures

were washed three times and resuspended in dH$_2$O. Cells were then plated to the appropriate

medium (100 μL 10^{-5} cells on YPD, 100 μL 10^{-2} cells on YNB + 5-FC, and 100 μL undiluted

cells on YPD + rapamycin + FK506 and YNB + 5-FOA). Mutation rates and 95% confidence

intervals were calculated using the FluCalc program[66]. For the increased temperature fluctuation

analysis, strains were grown overnight at either 30°C or 37°C before use in fluctuation assays, as

indicated. YPD + rapamycin + FK506 plates were incubated and 37°C; all other media was

incubated at 30°C. Mutation rates and confidence intervals for all fluctuation assays in this study

are provided in Table S6.

Characterizing mutation spectra

537   Following selection on antifungal drug media, resistant colonies were streak purified to

YPD medium. Genomic DNA was isolated from the purified colonies, and genes in which

mutations are known to cause resistance to the corresponding antifungal drug were PCR

amplified (*URA5* and *URA3* for 5-FOA-resistant colonies[44,45], *FRR1* for rapamycin+FK506-

resistant colonies[41,42], and *FUR1*, and *UXS1* for 5-FC-resistant colonies[46]). Oligonucleotides used

for all PCR reactions in this study are listed in Table S7. PCR products were subjected to gel

electrophoresis, products of interest were extracted from agarose gels using a QIAgen gel

extraction kit and sequenced through classical Sanger sequencing conducted at Genewiz.

Sequenced mutations, including transposon insertions, were characterized with both Sequencher

software and the Clustal Omega Multiple Sequence Alignment program[67]. Identified transposon

insertion sequences in *FRR1*, *URA5*, and *FUR1* are listed in Table S8.

Illumina sequencing

Single colonies from strains for whole-genome Illumina sequencing were inoculated in 50 mL of liquid YPD medium and grown overnight at 30°C, shaking. Cells were collected and lyophilized as previously described[65], and high molecular weight DNA was isolated following the CTAB protocol as previously described[68]. Strains were barcoded and sequencing libraries were generated with the Kapa HyperPlus library kit for 300bp inserts, pooled, and sequenced using paired-end, 2 x 150bp reads on an Illumina HiSeq 4000 platform at the Duke University Sequencing and Genomic Technologies Core facility.

Generation of F$_1$ progeny

Bt65**a** x H99α *crg1*Δ and Bt81**a** x H99α *crg1*Δ F$_1$ progeny were generated by genetically crossing either Bt65 or Bt81 with H99 *crg1*Δ on Murashige Skoog (MS) medium (Sigma) following Basic Protocol 1 as described in Sun et al. 2019[69]. Basidiospores were randomly isolated through microdissection after three weeks of incubation on MS following Basic Protocol 2 as described in Sun et al. 2019[69].

Nanopore sequencing and genome assemblies

The DNA samples for nanopore sequencing were isolated and purified using the CTAB DNA preparation protocol described previously[70]. The size estimation of the obtained DNA was done using PFGE electrophoresis and quality was determined using NanoDrop. Once the high-quality DNA was obtained, sequencing was performed using the MinION device with the MinKNOW interface. During sequencing, Bt65, Bt89, and Bt133 were multiplexed together whereas six of the Bt65**a** x H99α progeny were multiplexed for a second sequencing run. For

multiplexing, samples were barcoded using EXP-NBD103/EXP-NBD104 kits and libraries were

made using SQK-LSK109 kit as per the manufacturer's instructions. The libraries generated

were sequenced on R9.4.1 flow cell and reads were obtained in .fast5 format. These reads were

then converted to fastq format using Guppy_basecaller (v 4.2.2_linux64). The reads were de-

multiplexed using qcat (https://github.com/nanoporetech/qcat) or Guppy_barcoder (part of

Guppy_basecaller) with barcode trimming option during processing. Bt81 nanopore sequencing

was done as a standalone sample using an R9 flow cell (FLO-MN106) and basecalling was

performed during the run itself.

The sequences obtained for each sample were then assembled via Canu (v2.0 or v2.1.1)

to obtain contig-level genome assemblies. For the assembly, only >2 kb long reads were used for

the Bt65**a** x H99α F$_1$ progeny and Bt81, whereas >5 kb were used for Bt65, Bt89, and Bt133

genomes. Contigs were then assigned chromosome numbers based on their synteny with the

reference genome, H99. The numbering of chromosomes involved in translocations was assigned

based on the respective syntenic centromere. Some of the chromosomes were not fully

assembled and were broken into multiple contigs (Chr 1, Chr 2 for Bt65, Chr 2, Chr11, Chr14 for

Bt89, and Chr 2, Chr 5 for Bt133). For such cases, the respective contigs were joined artificially

and then processed by read-mapping to obtain complete collinear chromosomes. Specifically, the

contigs were stitched together in orientation as determined based on their synteny. Corrected

reads obtained from Canu were then mapped to the respective genomes and duplicated or

missing regions from the junction were identified. The chromosome sequence was then corrected

accordingly by inserting/correcting/deleting sequences and a full-length chromosome sequence

was obtained. Complete resolution of junctions was obtained for Bt65, Bt89, and Bt133 genomes

by this approach. However, some of the Bt65 F_1 progeny chromosomes could not be resolved,

probably due to hybrid origin of sequencing reads, and were left with gaps as such.

Once chromosome level genome assemblies were obtained for the Bt65, Bt81, Bt89, and

Bt133 genomes, the genome sequences were further processed to improve telomeric and

subtelomeric regions. For this purpose, the corrected reads obtained from Canu were mapped

back to the respective chromosome-level genomes using minimap2 v2.14. The obtained bam

files were then analyzed manually by IGV and consensus or, in a few cases, individual reads (up

to 30 kb) representing extra sequence beyond an assembled chromosome were extracted as sam

files. These consensus extra sequences were then added onto the chromosome sequences to

obtain longer chromosomes. In some cases, read mapping also resulted in the identification of

incorrect sequence assembly at subtelomeric regions, and in those cases, the sequence was

trimmed until a consensus sequence was observed at the end of the chromosome. Once these

corrections were made, the genome assemblies were polished via one round of nanopolish and

five rounds of pilon, except for the Bt81 genome, for which only 5X pilon polishing was

performed. As a result of these corrections and polishing, final assemblies were obtained for each

of the four isolates and are described in the study. For the Bt65 F_1 progeny genome assemblies,

the subtelomeric extension/curation was not performed, but they were polished using both

nanopolish and 5X pilon.

Centromere, telomere, and Cnl1 mapping

Centromeres in Bt65, Bt81, Bt89, and Bt133 were defined based on their synteny with

the reference H99 genome (genome assembly ASM301198v1) The final polished genomes were

used and centromere locations were identified by BLASTn analysis using H99 centromere-

flanking genes as query sequences. Once the centromere locations were defined, Tcn1-Tcn6

locations within those regions were mapped by BLASTn analysis. For the representation, only

BLAST hits longer than 400 bp were mapped. For the overlapping BLAST hits with multiple

Tcn elements, the longest and best BLAST result was used, and the rest of the matches were

discarded from further analysis. All the hits were then visualized using Geneious Prime and

maps were exported as .svg files, which were then processed using Adobe Illustrator.

For the Cnl1 mapping at the subtelomeres, the longest *CNL1* insertion sequence from the

Bt65 genome was used as the query sequence and BLASTn was performed against each genome.

BLAST hits longer than 50 bp were mapped to the respective genomes and visualized using

Geneious Prime where the hits were color-coded based on their lengths. The zoomed views for

these maps were then exported as .svg files, processed using Adobe Illustrator, and combined

with centromere Tcn mapping analysis to generate final figures.

RepeatMasker was used to annotate all transposons in the *de novo* genome assembly of

Bt65. For this purpose, RepeatMasker (v4.0.7) with Dfam (v3.3) and

RepBaseRepeatMaskerEdition-20181026 libraries was used, supplemented with RepBase EMBL

database (v26.04)[71–73]. The "-species fungi" option was used to identify all repeats in the genome

and provided additional support for the manual Tcn and Cnl1 mapping.

Synteny maps

Synteny comparisons between the genomes were performed using SyMAP v4.2 with the

H99 genome as the reference (genome assembly ASM301198v1). The synteny comparison was

conducted using default parameters and synteny block maps were exported as .svg files. The

maps were processed using Adobe Illustrator for visualization. The phylogenetic relationship as

depicted in Figure S6 was drawn based on the earlier representation[18]. The telomere and

centromere locations were marked manually based on the presence of the telomere repeat

sequence and Tcn mapping, respectively.

For the centromere comparisons, all centromere sequences along with Tcn annotations

were converted into GenBank format. The files were then used for synteny comparison via

EasyFig v2.2.3. The maps were exported as .svg files which were processed in Adobe Illustrator.

Recombination maps for Bt65 x H99 F$_1$ progeny

Six of the Bt65**a** x H99α F$_1$ progeny were sequenced with on the nanopore MinION

sequencing platform and their genomes were assembled and polished using the methods

described above. Once their genomes were assembled, recombination maps were generated by

mapping the Illumina sequence data from the parental strains to each of the progeny genomes.

For this purpose, both H99 and Bt65 Illumina reads were used from published datasets

(SRR642222 and SRR647805 for H99; SRR836876, SRR836877, SRR836878, SRR836880,

SRR836884, and SRR836885 for Bt65). Reads from all runs were merged to obtain a single file

for both H99 and Bt65. The reads were then mapped to the progeny genomes using Geneious

Prime default mapper with three iterations. Variants with 90X coverage and at least 90% variant

frequency were called from these mapped files. These variants along with coverage analysis

were then used to identify recombination sites and generate recombination maps. Cnl1 mapping

for each of progeny genome was performed as described above. The location of *ZNF3* in each

genome was identified by BLASTn analysis using H99 *ZNF3* (CNAG_02700) as the query

sequence.

Genetic variant calling and segregant filtering

Whole-genome sequencing data of 28 F_1 progeny from the Bt65**a** x H99α *crg1*Δ cross were aligned via BWA (v0.7.12-r1039)[74] to an H99 reference genome (downloaded from FungiDB [http://fungidb.org/fungidb/] on April 15th, 2020; FungiDB-46_CneoformansH99_Genome.fasta) and genetic variants between Bt65 and H99 were called using SAMtools (v0.1.19-96b5f22941)[75] and FreeBayes (v1.2.0)[76]. Approximately 300,000 raw genetic variants were identified across the segregants. The genotypic correlation between progeny, the read coverage per genetic variant, and the ratio of reads suggesting the H99 vs. Bt65 allele per variant were monitored across the genome to identify clones, progeny with aneuploid genomes, and heterozygotic diploids (respectively). Two pairs of clones were identified (Supplementary Table S9) and one segregant from each pair was retained for analysis. F_1 progeny 25 was identified as a heterozygotic diploid (Supplementary Figure S13) and removed from initial analysis. Instances of aneuploidy (and partial duplications) are observed along Chromosomes 3, 4, 11, and 13 within six segregants from this cross and for initial filtering and analysis, those with heterozygotic aneuploidy were removed from analysis (Supplementary Table S9).

Genetic variant filtering

After removing clones and samples with aneuploidy or diploidy, raw genetic variants were filtered by limiting sites to bi-allelic SNPs, called across all the progeny (100% call rate), with greater than 10X read coverage (and a maximum of 200X), a minor allele frequency of 5%, and a quality score greater than 4 (and less than 5.4). These filtering criteria were selected after examining the bivariate relationships between allele frequency, read depth, and quality scores per

chromosome (Supplementary Figure S14A and 14B). Genetic variant sites were also removed if

within one kb of the centromere along a given chromosome[28]. After filtering, a total of 215,411

bi-allelic SNPs were retained for further analysis. The median distance between contiguous SNP

sites is 45 bp, and less than 0.01% of neighboring sites had a distance larger than two kb. The

allele frequencies across the genome ranged between 25 and 75% of segregant with the Bt65

allele, except for a large portion of Chromosome 13, between 0 to 500 kb, where over 88% of

segregants inherited the Bt65 allele (Figure S14C). With these data, a Poisson regression

(methods described in Roth *et al*. 2018[77]) was used to relate the average number of crossovers

across F_1 progeny as a function of chromosome size. With this model, the estimated genome-

wide, physical-to-genetic distance in this cross is 8.14 kb/cM.

QTL mapping

For use in association tests, across 24 F_1 progeny and the two parental strains, the

215,411 bi-allelic SNPs were collapsed into 1,237 unique haplogroups made up of genetic

variants sites that co-segregated within the segregant genomes, such that, between any two

haplogroups, at least one segregant contains a change in allele (i.e. a recombination event

between the Bt65**a** and H99α *crg1*Δ genomes). Collapsing genetic variants into haplogroups

reduces the number of repeated tests in association mapping and computational costs[78]. Across

the 1,237 haplogroups, the mutation rate on rapamycin + FK506 medium was used as the

phenotype and tested for association using a Kruskal-Wallis H-test. The $-\log_{10}$ (*p*-value) from

this test across haplogroups was monitored to identify QTL. Significance thresholds were

established via 10,000 permutations with an α = 0.01 as described in Churchill and Doerge

(1994)[79], and 95% confidence intervals for the QTL locations were generated as described in

Visscher *et al.* (1996)[80]. The heritability at the peak of identified QTL was estimated using linear

regression with the model: $M = \mu + \beta I + e$, were M is the mutation rate x 10^7, e is an error term, μ

is the mean mutation rate x 10^7, I is an indicator variable for the allele at the QTL peak – coded

as 0 if from H99α *crg1*Δ or 1 if from Bt65**a** – and β is the effect of having the H99α *crg1*Δ vs.

the Bt65**a** allele at the QTL. The variation explained (R^2) from this model was used as an

estimate of the heritability.

Gene annotation and SNP effect prediction

For genes within the identified QTL spanning Chromosomes 3 and 11, the alleles

between H99 and Bt65 were imputed using filtered SNP data (described above). The published

H99 reference strain annotation (downloaded from FungiDB [http://fungidb.org/fungidb/] on

April 15th, 2020; FungiDB-46_CneoformansH99.gff) was used to predict changes in protein

sequence between the H99 and Bt65 parental backgrounds.

CRISPR-mediated genetic complementation of *ZNF3* in Bt65

To change the single nucleotide responsible for the nonsense mutation in the first exon of

ZNF3, a thymine (base 976004 of H99 Chromosome 3 (CNA3 assembly, accession

GCA_000149245.3)), to the wild-type cytosine found in H99 and other phylogenetically closely

related strains, the Transient CRISPR-Cas9 Coupled with Electroporation (TRACE) system was

used[48]. Briefly, the gene encoding Cas9 was PCR amplified from plasmid pXL1-CAS9-HYG.

The *SH1-NEO* construct encoding *NEO* (G418 resistance) targeted to a safe haven locus (SH1)

was amplified from plasmid pSDMA57[81]. For the sgRNA expression construct, the U6 promoter

was amplified from XL280α gDNA, and the sgRNA scaffold was amplified from plasmid

pYF515[82]. Overlap PCR was used to generate the sgRNA construct with the U6 promoter and

sequences targeting either SH1 or the nonsense mutation in *ZNF3*. A 2,171bp region was

amplified from Bt133, containing the wild-type C nucleotide in *ZNF3* exon 1 and no other

mutations relative to Bt65 for integration at the *ZNF3* endogenous locus in Bt65 (1,197bp

upstream of the *ZNF3* start codon to 971bp after the start codon). The *SH1-NEO* construct was

linearized with the AscI restriction enzyme (NEB). 2µg of the Bt133 *ZNF3* recombination

template, 2µg of the *SH1-NEO* linearized construct, 250ng of the *ZNF3* gRNA, 250ng of the

SH1 gRNA, and 1.5µg of Cas9 DNA were transformed simultaneously into Bt65 via

electroporation using a BIO-RAD Gene Pulser.

Transformants were selected on YPD + G418 agar plates. Successful transformants were

identified through restriction enzyme digest with BtsI-v2 (NEB), which cleaves the first exon of

Bt65 *znf3* at the nonsense mutation but does not cleave the first exon of *ZNF3* in Bt133 (or H99)

(Figure S15A). PCR was also used to ensure that no transformants had integrated copies of the

gene encoding Cas9 or the gRNA constructs and that only a single Bt133 *ZNF3* allele had

integrated correctly at the endogenous *znf3* locus (Figure S15B-F). Sanger sequencing was used

to further confirm correct replacement of the Bt65 *znf3* allele including the nonsense mutation

with the Bt133 *ZNF3* allele. No identified Bt65+*ZNF3* transformants also had a stably integrated

copy of the *NEO* gene at the safe haven locus.

sRNA isolation and sequencing

C. neoformans cells were grown overnight in 50 mL YPD medium at standard laboratory

conditions. Following culture, cells were spun down, supernatant was removed, and cells were

frozen at -80°C for one hour. Cells were then freeze dried with a Labconco Freezone 4.5

lyophilizer overnight. 70 mg of lyophilized material was used for sRNA isolation following the

mirVana miRNA Isolation Kit manufacturer's instructions. sRNA was quantified with a Qubit 3

Fluorometer and quality was verified with an Agilent Bioanalyzer using an Agilent Small RNA

Kit. sRNA libraries were prepared with a QiaPrep miRNA Library Prep Kit and 1 x 75 bp reads

were sequenced on the Illumina NextSeq 500 System at the Duke University Sequencing and

Genomic Technologies Core facility.

sRNA data processing

Initial quality control of the small RNA libraries was performed with FastQC 0.11.9[83]

followed by the removal of QIAseq library adapters (5':

GTTCAGAGTTCTACAGTCCGACGATC; 3': AACTGTAGGCACCATCAAT) with cutadapt

2.8[84]. All untrimmed reads or reads smaller than 14 nt were discarded. The surviving trimmed

reads were mapped with bowtie v1.2.3[85] to the *C. neoformans* Bt65 genome, allowing multiple

alignments but no mismatches. The resulting SAM files were converted into BAM file format

with SAMtools 1.9[75] and feature read counts of transposable elements were calculated with

BEDTools[86] using the `intersect -wa´ option and the annotations of transposable elements, which

were identified with RepeatMasker using the repbase database for *C. neoformans*[71,72].

Normalization of the read counts to reads per million (RPM) was performed, allowing the

comparison of the libraries. Furthermore, the read depth on both DNA strands was analyzed with

SAMtools and custom made perl scripts were used to calculate the read size distribution and 5'-

nucleotide preference of the small RNA reads as previously described[87,88].

Double-stranded RNA enrichment

For dsRNA enrichment, *Cryptococcus* cells were grown overnight in 50mL liquid YPD

medium at 30°C. RNA was extracted, and dsRNA was enriched as previously described[55].

dsRNA enrichment in H99, Bt65, and Bt81 did not reveal the presence of any large dsRNA

segments (Figure S16).

Data and software availability

Genetic variant filtering, QTL mapping, and SNP effect prediction was conducted in

python (anaconda 3.7.3) via custom scripts; hosted at:

https://github.com/magwenelab/Hypermutator_QTL . All sequencing data is available under

BioProject PRJNA749953.

Acknowledgments

We thank and acknowledge Blake Billmyre for initial project guidance, Shelly Clancey

for instruction in conducting fluctuation assays and dsRNA enrichment protocols, Josh Granek

for preliminary analyses of hypermutator genomes, Zanetta Chang for assistance in sRNA

isolation, Kayla Sylvester for assistance with screening of SDC isolates, and the laboratory of

Chris Holley at Duke University for the use of their Nanodrop and BioAnalyzer equipment for

preliminary sRNA analysis. We thank Mark Farman and Mostafa Rahnama for stimulating

discussion on the impacts of transposons on telomere dynamics. We thank Kevin Zhu for

assistance with figure generation. We also thank Sheng Sun, Blake Billmyre, Andy Alspaugh,

Sue Jinks-Robertson, Asiya Gusa, and Kayla Sylvester for critical reading and comments on the

manuscript. This work was funded by NIH/NIAID F31 Fellowship 1F31AI143136-02A1

awarded to S.J.P. and NIH/NIAID R37 MERIT award AI39115-23, R01 grant AI50113-16, and

R01 grant AI33654-04 awarded to J.H. J.H. is co-Director and Fellow of the CIFAR program

Fungal Kingdom: Threats & Opportunities. We also thank the Madhani Laboratory and NIH

grant R01 AI100272 for the KN99α *msh2Δ* deletion strain. T.A.D. and U.K. are funded by the

German Research Foundation (DFG) (Bonn Bad-Godesberg, Germany) (KU 517/15-1).

References

1. Sniegowski, P. D., Gerrish, P. J. & Lenski, R. E. Evolution of high mutation rates in experimental populations of *E. coli*. *Nature* **387**, 703–705 (1997).

2. Barrick, J. *et al.* Genome evolution and adaptation in a long-term experiment with *Escherichia coli*. *Nature* **461**, 1243–1247 (2009).

3. Tenaillon, O. *et al.* Tempo and mode of genome evolution in a 50,000-generation experiment. *Nature* **536**, 165–170 (2016).

4. Harfe, B. D. & Jinks-Robertson, S. Mismatch repair proteins and mitotic genome stability. *Mutat. Res.* **451**, 151–167 (2000).

5. Fisher, K. J., Buskirk, S. W., Vignogna, R. C., Marad, D. A. & Lang, G. I. Adaptive genome duplication affects patterns of molecular evolution in *Saccharomyces cerevisiae*. *PLoS Genet.* **14**, e1007396 (2018).

6. Billmyre, R. B., Croll, D. & Li, W. Highly recombinant VGII *Cryptococcus gattii* population develops clonal outbreak clusters through both sexual macroevolution and asexual microevolution. *mBio* **5**, e01494-14 (2014).

7. Billmyre, R. B., Clancey, S. A. & Heitman, J. Natural mismatch repair mutations mediate phenotypic diversity and drug resistance in *Cryptococcus deuterogattii*. *eLife* **6**, e28802 (2017).

8. Healey, K. R. *et al.* Prevalent mutator genotype identified in fungal pathogen *Candida glabrata* promotes multi-drug resistance. *Nat. Commun.* **7**, 11128 (2016).

9. Boyce, K. J. *et al.* Mismatch repair of DNA replication errors contributes to microevolution in the pathogenic fungus *Cryptococcus neoformans*. *mBio* **8**, e00595-17 (2017).

829   10.   Steenwyk, J. L. *et al.* Extensive loss of cell-cycle and DNA repair genes in an ancient

830   lineage of bipolar budding yeasts. *PLoS Biol.* **17**, e3000255 (2019).

831   11.   Sionov, E., Lee, H., Chang, Y. C. & Kwon-Chung, K. J. *Cryptococcus neoformans*

832   overcomes stress of azole drugs by formation of disomy in specific multiple

833   chromosomes. *PLoS Pathog.* **6**, e1000848 (2010).

834   12.   Kwon-Chung, K. J. & Chang, Y. C. Aneuploidy and drug resistance in pathogenic fungi.

835   *PLoS Pathog.* **8**, 8–11 (2012).

836   13.   Janbon, G. *et al.* Characterizing the role of RNA silencing components in *Cryptococcus*

837   *neoformans*. *Fungal Genet. Biol.* **47**, 1070–1080 (2010).

838   14.   Gusa, A. *et al.* Transposon mobilization in the human fungal pathogen *Cryptococcus* is

839   mutagenic during infection and promotes drug resistance *in vitro*. *Proc. Natl. Acad. Sci.*

840   *U. S. A.* **117**, 9973–9980 (2020).

841   15.   Idnurm, A. *et al.* Deciphering the model pathogenic fungus *Cryptococcus neoformans*.

842   *Nat. Rev. Microbiol.* **3**, 753–764 (2005).

843   16.   Chayakulkeeree, M. & Perfect, J. R. Cryptococcosis. *Infect. Dis. Clin. North Am.* **20**, 507–

844   544 (2006).

845   17.   Hagen, F. *et al.* Recognition of seven species in the *Cryptococcus gattii*/*Cryptococcus*

846   *neoformans* species complex. *Fungal Genet. Biol.* **78**, 16–48 (2015).

847   18.   Desjardins, C. A. *et al.* Population genomics and the evolution of virulence in the fungal

848   pathogen *Cryptococcus neoformans*. *Genome Res.* **27**, 1207–1219 (2017).

849   19.   Rajasingham, R. *et al.* Global burden of disease of HIV-associated cryptococcal

850   meningitis: an updated analysis. *Lancet Infect. Dis.* **17**, 873–881 (2017).

20.   Brouwer, A. *et al.* Combination antifungal therapies for HIV-associated cryptococcal

meningitis: a randomised trial. *Lancet* **363**, 1764–1767 (2004).

21.   Laniado-Laborín, R. & Cabrales-Vargas, M. N. Amphotericin B: side effects and toxicity.

Rev. Iberoam. Micol. **26**, 223–227 (2009).

22.   Vandeputte, P. *et al.* Molecular mechanisms of resistance to 5-fluorocytosine in laboratory

mutants of *Candida glabrata*. *Mycopathologia* **171**, 11–21 (2011).

23.   Perfect, J. R. & Cox, G. M. Drug resistance in *Cryptococcus neoformans*. *Drug Resist.*

Updat. **2**, 259–269 (1999).

24.   Nosanchuk, J. D., Cleare, W., Franzot, S. P. & Casadevall, A. Amphotericin B and

fluconazole affect cellular charge, macrophage phagocytosis, and cellular morphology of

Cryptococcus neoformans at subinhibitory concentrations. *Antimicrob. Agents Chemother.*

43, 233–239 (1999).

25.   Posteraro, B. *et al.* Identification and characterization of a *Cryptococcus neoformans* ATP

binding cassette (ABC) transporter-encoding gene, *CnAFR1*, involved in the resistance to

fluconazole. *Mol. Microbiol.* **47**, 357–371 (2003).

26.   Loftus, B. J. *et al.* The genome of the basidiomycetous yeast and human pathogen

Cryptococcus neoformans. *Science* **307**, 1321–1324 (2005).

27.   Goodwin, T. J. D. & Poulter, R. T. M. The diversity of retrotransposons in the yeast

Cryptococcus neoformans. *Yeast* **18**, 865–880 (2001).

28.   Yadav, V. *et al.* RNAi is a critical determinant of centromere evolution in closely related

fungi. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 3108–3113 (2018).

29.   Cruz, M. C. *et al.* Rapamycin antifungal action is mediated via conserved complexes with

FKBP12 and TOR kinase homologs in *Cryptococcus neoformans*. *Mol. Cell. Biol.* **19**,

4101–4112 (1999).

30.   Wang, X. *et al.* Sex-induced silencing defends the genome of *Cryptococcus neoformans*

876   via RNAi. *Genes Dev.* **24**, 2566–2582 (2010).

877   31.   Dumesic, P. A. *et al.* Stalled spliceosomes are a signal for RNAi-mediated genome

878   defense. *Cell* **152**, 957–968 (2013).

879   32.   Feretzaki, M., Billmyre, R. B., Clancey, S. A., Wang, X. & Heitman, J. Gene network

880   polymorphism illuminates loss and retention of novel RNAi silencing components in the

881   *Cryptococcus* pathogenic species complex. *PLoS Genet.* **12**, e1005868 (2016).

882   33.   Feretzaki, M. & Heitman, J. Genetic circuits that govern bisexual and unisexual

883   reproduction in *Cryptococcus neoformans*. *PLoS Genet.* **9**, e1003688 (2013).

884   34.   Wang, X., Darwiche, S. & Heitman, J. Sex-induced silencing operates during opposite-sex

885   and unisexual reproduction in *Cryptococcus neoformans*. *Genetics* **193**, 1163–1174

886   (2013).

887   35.   Huff, J. T. & Zilberman, D. Dnmt1-independent CG methylation contributes to

888   nucleosome positioning in diverse eukaryotes. *Cell* **156**, 1286–1297 (2014).

889   36.   Catania, S. *et al.* Evolutionary persistence of DNA methylation for millions of years after

890   ancient loss of a *de novo* methyltransferase. *Cell* **180**, 263–277 (2020).

891   37.   Dumesic, P. A. *et al.* Product binding enforces the genomic specificity of a yeast

892   Polycomb repressive complex. *Cell* **160**, 204–218 (2015).

893   38.   Ferrareze, P. A. G. *et al.* Application of an optimized annotation pipeline to the

894   *Cryptococcus deuterogattii* genome reveals dynamic primary metabolic gene clusters and

895   genomic impact of RNAi loss. *G3* **11**, jkaa070 (2021).

896   39.   Breuder, T., Hemenway, C. S., Movva, N. R., Cardenas, M. E. & Heitman, J. Calcineurin

897   is essential in cyclosporin A- and FK506-sensitive yeast strains. *Proc. Natl. Acad. Sci. U.*

S. A. **91**, 5372–5376 (1994).

40.   Parent, S. A. *et al.* Calcineurin-dependent growth of an FK506- and CsA-hypersensitive

mutant of *Saccharomyces cerevisiae*. *J. Gen. Microbiol.* **139**, 2973–2984 (1993).

41.   Heitman, J., Movva, N. R., Hiestand, P. C. & Hall, M. N. FK 506-binding protein proline

rotamase is a target for the immunosuppressive agent FK 506 in *Saccharomyces*

cerevisiae. *Proc. Natl. Acad. Sci. U. S. A.* **88**, 1948–1952 (1991).

42.   Heitman, J., Movva, N. R. & Hall, M. N. Targets for cell cycle arrest by the

immunosuppressant rapamycin in yeast. *Science* **253**, 905–909 (1991).

43.   Litvintseva, A. P., Thakur, R., Vilgalys, R. & Mitchell, T. G. Multilocus sequence typing

reveals three genetic subpopulations of *Cryptococcus neoformans* var. *grubii* (serotype A),

including a unique population in Botswana. *Genetics* **172**, 2223–2238 (2006).

44.   Kwon-Chung, K. J., Varma, A., Edman, J. C. & Bennett, J. Selection of *ura5* and *ura3*

mutants from the two varieties of *Cryptococcus neoformans* on 5-fluoroorotic acid

medium. *J. Med. Vet. Mycol.* **30**, 61–69 (1992).

45.   Edman, J. C. & Kwon-Chung, K. J. Isolation of the *URA5* gene from *Cryptococcus*

neoformans var. *neoformans* and its use as a selective marker for transformation. *Mol.*

Cell. Biol. **10**, 4538–4544 (1990).

46.   Billmyre, R. B., Clancey, S. A., Li, L. X., Doering, T. L. & Heitman, J. 5-fluorocytosine

resistance is associated with hypermutation and alterations in capsule biosynthesis in

Cryptococcus. *Nat. Commun.* **11**, 127 (2020).

47.   Janbon, G. *et al.* Analysis of the genome and transcriptome of *Cryptococcus neoformans*

var. grubii reveals complex RNA expression and microevolution leading to virulence

attenuation. *PLoS Genet.* **10**, e1004261 (2014).

48.   Fan, Y. & Lin, X. Multiple applications of a Transient CRISPR-Cas9 Coupled with

Electroporation (TRACE) system in the *Cryptococcus neoformans* species complex.

Genetics **208**, 1357–1372 (2018).

49.   Drinnenberg, I. A. *et al.* RNAi in budding yeast. *Science* **326**, 544–550 (2009).

50.   Rahnama, M. *et al.* Transposon-mediated telomere destabilization: a driver of genome

evolution in the blast fungus. *Nucleic Acids Res.* **48**, 7197–7217 (2020).

51.   Mason, J. A. & Biessmann, H. The unusual telomeres of *Drosophila*. *Trends Genet.* **11**,

58–62 (1995).

52.   Pardue, M. Lou & DeBaryshe, P. G. Retrotransposons provide an evolutionarily robust

non-telomerase mechanism to maintain telomeres. *Annu. Rev. Genet.* **37**, 485–511 (2003).

53.   Ashe, A. *et al.* A deletion polymorphism in the *Caenorhabditis elegans* RIG-I homolog

disables viral RNA dicing and antiviral immunity. *eLife* **2**, e00994 (2013).

54.   Félix, M.-A. *et al.* Natural and experimental infection of *Caenorhabditis* nematodes by

novel viruses related to nodaviruses. *PLoS Biol.* **9**, e1000586 (2011).

55.   Clancey, S. A., Ruchti, F., Leibundgut-Landmann, S., Heitman, J. & Ianiri, G. A novel

mycovirus evokes transcriptional rewiring in the fungus *Malassezia* and stimulates beta

interferon production in macrophages. *mBio* **11**, e01534-20 (2020).

56.   Kanafani, Z. A. & Perfect, J. R. Resistance to antifungal agents: mechanisms and clinical

impact. *Antimicrob. Resist.* **46**, 120–128 (2008).

57.   Weiberg, A. *et al.* Fungal small RNAs suppress plant immunity by hijacking host RNA

interference pathways. *Science* **342**, 118–123 (2013).

58.   McClintock, B. The origin and behavior of mutable loci in maize. *Proc. Natl. Acad. Sci.*

U. S. A. **36**, 344–355 (1950).

59.   Rosenberg, A. *et al.* Antifungal tolerance is a subpopulation effect distinct from resistance
945   and is associated with persistent candidemia. *Nat. Commun.* **9**, 2470 (2018).

946   60.   Berman, J. & Krysan, D. J. Drug resistance and tolerance in fungi. *Nat. Rev. Microbiol.*
947   **18**, 319–331 (2020).

948   61.   Devos, K. M., Brown, J. K. M. & Bennetzen, J. L. Genome size reduction through
949   illegitimate recombination counteracts genome expansion in *Arabidopsis*. *Genome Res.*
950   **12**, 1075–1079 (2002).

951   62.   Vitte, C. & Panaud, O. LTR retrotransposons and flowering plant genome size: emergence
952   of the increase/decrease model. *Cytogenet. Genome Res.* **110**, 91–107 (2005).

953   63.   Drinnenberg, I. A., Fink, G. R. & Bartel, D. P. Compatibility with killer explains the rise
954   of RNAi-deficient fungi. *Science* **333**, 1592 (2011).

955   64.   Barton, E. S. *et al.* Herpesvirus latency confers symbiotic protection from bacterial
956   infection. *Nature* **447**, 326–329 (2007).

957   65.   Priest, S. J. *et al.* Factors enforcing the species boundary between the human pathogens
958   *Cryptococcus neoformans* and *Cryptococcus deneoformans*. *PLoS Genet.* **17**, e1008871
959   (2021).

960   66.   Radchenko, E. A., McGinty, R. J., Aksenova, A. Y., Neil, A. J. & Mirkin, S. M.
961   Quantitative analysis of the rates for repeat-mediated genome instability in a yeast
962   experimental system. *Methods Mol. Biol.* **1672**, 421–438 (2018).

963   67.   Madeira, F. *et al.* The EMBL-EBI search and sequence analysis tools APIs in 2019.
964   *Nucleic Acids Res.* **47**, W636–W641 (2019).

965   68.   Pitkin, J. W., Panaccione, D. G. & Walton, J. D. A putative cyclic peptide efflux pump
966   encoded by the *TOXA* gene of the plant-pathogenic fungus *Cochliobolus carbonurn*.

967   *Microbiology* **142**, 1557–1565 (1996).

968 69. Sun, S., Priest, S. J. & Heitman, J. *Cryptococcus neoformans* mating and genetic crosses.

969   *Curr. Protoc. Microbiol.* **53**, e75 (2019).

970 70. Yadav, V., Sun, S., Coelho, M. A. & Heitman, J. Centromere scission drives chromosome

971   shuffling and reproductive isolation. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 7917–7928

972   (2020).

973 71. Bao, W., Kojima, K. K. & Kohany, O. Repbase Update, a database of repetitive elements

974   in eukaryotic genomes. *Mob. DNA* **6**, 11 (2015).

975 72. Smit, A., Hubley, R. & Green, P. RepeatMasker Open-4.0. *http://www.repeatmasker.org*

976   (2013).

977 73. Storer, J., Hubley, R., Rosen, J., Wheeler, T. J. & Smit, A. F. The Dfam community

978   resource of transposable element families, sequence models, and genome annotations.

979   *Mob. DNA* **12**, 2 (2021).

980 74. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

981   transform. *Bioinformatics* **25**, 1754–1760 (2009).

982 75. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

983   2078–2079 (2009).

984 76. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.

985   *arXiv* 1207.3907 [q-bio.GN] (2012).

986 77. Roth, C., Sun, S., Billmyre, R. B., Heitman, J. & Magwene, P. M. A high-resolution map

987   of meiotic recombination in *Cryptococcus deneoformans* demonstrates decreased

988   recombination in unisexual reproduction. *Genetics* **209**, 567–578 (2018).

989 78. Xu, S. Genetic mapping and genomic selection using recombination breakpoint data.

Genetics **195**, 1103–1115 (2013).

79.   Churchill, G. A. & Doerge, R. W. Empirical threshold values for quantitative trait

mapping. *Genetics* **138**, 963–971 (1994).

80.   Visscher, P. M., Thompson, R. & Haley, C. S. Confidence intervals in QTL mapping by

bootstrapping. *Genetics* **143**, 1013–1020 (1996).

81.   Arras, S. D. M., Chitty, J. L., Blake, K. L., Schulz, B. L. & Fraser, J. A. A genomic safe

haven for mutant complementation in *Cryptococcus neoformans*. *PLoS One* **10**, e0122916

(2015).

82.   Fang, Y., Cui, L., Gu, B., Arredondo, F. & Tyler, B. M. Efficient genome editing in the

oomycete *Phytophthora sojae* using CRISPR/Cas9. *Curr. Protoc. Microbiol.* **44**, 21A.1.1-

21A.1.26 (2017).

83.   Andrews, S. FastQC: a quality control tool for high throughput sequence data.

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (2010).

84.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal **17**, 10–12 (2011).

85.   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009).

86.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841–842 (2010).

87.   Dahlmann, T. A. & Kück, U. Dicer-dependent biogenesis of small RNAs and evidence for

microRNA-like RNAs in the penicillin producing fungus *Penicillium chrysogenum*. *PLoS*

One **10**, e0125989 (2015).

88.   Ianiri, G. *et al.* Mating-type-specific ribosomal proteins control aspects of sexual

reproduction in *Cryptococcus neoformans*. *Genetics* **214**, 635–649 (2020).

1014

 Supplementary Figures

 Supplementary Figure S1. Bt65 and Bt81 do not display a hypermutator phenotype on 5-

 FC or 5-FOA. Mutation rates of closely related VNBII strains and controls on **(A)** YNB + 5-FC

 and **(B)** YNB + 5-FOA media. Bars represent the mutation rate and error bars represent 95%

 confidence intervals; mutation rates represent the number of mutations per cell per generation.

 Schematic depicts the phylogenetic relationships of all strains included in fluctuation analyses

 based on Desjardins et al. 2017[18].

1024

1025

1026

1027

1028

1029 **Supplementary Figure S2. Growth at elevated temperature does not result in increased**

1030 **mutation rates in *C. neoformans* strains.** Fluctuation assays were used to quantify the mutation

1031 rates of strains grown overnight at 30°C or 37°C and plated on YPD + rapamycin + FK506

1032 medium. Bars indicate mean mutation rate and error bars indicate 95% confidence intervals.

1033 Mutation rates represent the number of mutations per cell per generation.

1034

1035

Supplementary Figure S3. Gel electrophoresis *FRR1*, *URA5*, and *FUR1* PCR products

from resistant colonies. Gel electrophoresis of *FRR1* PCR products from **(A)** all H99 rapamycin

+ FK506-resistant colonies and a subset of **(B)** Bt65 and **(C)** Bt81 rapamycin + FK506-resistant

colonies sequenced in Figure 1D. PCR amplification of wild-type *FRR1* in *C. neoformans*

produces a 1,165 bp electrophoretic species (primers ZC7/8). Gel electrophoresis of a subset of

(D) *URA5* PCR products from H99, Bt65, and Bt81 5-FOA-resistant colonies and **(E)** *FUR1*

PCR products from 5-FC-resistanct colonies of Bt65 and Bt81.

1043

1044

1045

Supplementary Figure S4. Protein length differences of genes within QTL. In the upper

panels, points mark the strength of association (y-axis) between bi-allelic SNP sites and

hypermutation for Chromosome 3 and Chromosome 11 (top left and right, respectively). Grey

dashed lines depict the 95% confidence intervals (CI) of the two QTL. Lower panels show the

predicted differences in lengths of proteins (y-axis) encoded by annotated genes in Bt65

compared to H99 within each 95% CI of the QTL (x-axis) on Chromosome 3 and Chromosome

11 (bottom left and right, respectively). The name of each gene with a predicted nonsense

mutation is annotated. Blue and red colors denote the gene orientation.

1054

1055

1056

1057

1058

Supplementary Figure S5. **Haplotype maps of Bt65 x H99 F₁ progeny utilized for QTL**

mapping. For the QTLs on Chromosome 3 and Chromosome 11 (left and right, respectively) the

haplotypes (x-axis) are inferred by SNP data per segregant (y-axis) and colored blue or orange if

inherited from H99 *crg1Δ* or Bt65, respectively. Segregants are sorted along the y-axis by the

quantification of their mutation rate; largest to smallest, top to bottom. Vertical red lines display

the boundaries of the QTL(s). Vertical black lines depict the approximate location of the

translocation between H99 and Bt65. The boundaries of the QTG, *ZNF3*, are depicted by vertical

green lines. Vertical white spaces indicate the approximate locations of the centromeres.

1067

1068

1069

1070

1071

1072

Supplementary Figure S6. **Large-scale genomic rearrangements between the H99, Bt65,**

Bt81, Bt89, and Bt133 genomes. Nanopore whole-genome sequencing followed by synteny

analysis was used to identify all indicated genomic rearrangements with respect to the reference

strain H99. There is a chromosomal translocation between Chromosomes 3 and 11 that is unique

to H99, and a translocation between H99 Chromosomes 1 and 13 that is unique to Bt65 and

Bt81. The phylogenetic relationships of these strains are depicted in the top schematic, telomeric

repeat sequences accurately identified in the genomic assemblies are indicated by black half

circles, and centromeres are indicated by white circles.

1081

1082

1083

1084

Supplementary Figure S7. **CNAG_01836 gene model.** A grey horizontal bar depicts the entire

gene body in the upper panel, and larger grey rectangles show locations of exons. The gene is

depicted 5' to 3', left to right, and is 1896 nt in length. The locations of SNPs differing between

Bt65 and H99 are shown by vertical black rungs along the gene model. Amino acids specified by

mRNA codons in the indicated region of CNAG_01836 Exon 7 (nucleotide 1831 to 1848) are

shown to illustrate the G to A mutation (nucleotide 1842) predicted to cause an early nonsense

mutation in Bt65. The bottom panel depicts the predicted outcome of the nonsense mutation on

the protein encoded by CNAG_01836 in Bt65.

1093

1098

Supplementary Figure S8. Mutation rates of Bt81 x H99 F₁ progeny. Fluctuation analysis

was used to quantify the mutation rates of the indicated strains on YPD + rapamycin + FK506

medium (y-axis) – sorted smallest to largest, left to right – for F₁ progeny and the parental

strains, H99α *crg1Δ* and Bt81 (x-axis). Bars indicate the mean mutation rate and error bars

represent 95% confidence intervals. Mutation rates represent the number of mutations per cell

per generation. Inheritance of the Bt81 *znf3* allele or H99 *crg1Δ ZNF3* allele in the F₁ progeny is

indicated above mutation rates.

1106

1107

1108

1109

10% [____________________] 100%
Cnl1 length percent
■ Tcn1 ■ Tcn2 ■ Tcn3 ■ Tcn4 ■ Tcn5 ■ Tcn6

1110

Supplementary Figure S9. Subtelomeric and centromeric retrotransposons in Bt89 and

Bt133. Distributions of the Tcn1-Tcn6 LTR-retrotransposons and the Cnl1 non-LTR

retrotransposon in the genomes of (A) Bt89 and (B) Bt133. 50 kb of subtelomeric regions as well

as centromeric regions are displayed for both strains. Shading corresponds to the lengths of the

Cnl1 elements, and gene arrowheads indicate the direction of transcription for all

retrotransposons.

1117

1118

1119

1120

1121

1122   **Supplementary Figure S10. Centromere lengths do not significantly differ among H99,**

1123   **Bt65, Bt81, Bt89, and Bt133.** The length of each centromere (y-axis) is plotted for each strain

1124   (x-axis). The thin horizontal black line indicates average centromere length and the thicker black

1125   error bars indicate the standard error of the mean. No significant difference was found between

1126   the average centromere length of each strain (ANOVA, *p*-value = 0.153).

1127

1128

1129

1130

1131 **Supplementary Figure S11**. **Centromeric rearrangements among hypermutators and two**

1132 **closely related strains.** Shared homology, genomic rearrangements, inversions, insertions and

1133 deletions among the centromeres of each of the 14 chromosomes are depicted for Bt65, Bt81,

1134 Bt89, and Bt133. The organization of LTR-retrotransposons Tcn1 through Tcn6 are also

1135 indicated for each centromere of each strain.

1136

1137

Supplementary Figure S12. **Distribution of Cnl1 among Bt65 x H99 F₁ progeny and parental strains.** The Cnl1 non-LTR elements identified in the nanopore-based whole-genome assemblies are depicted for H99, Bt65, three hypermutator F_1 progeny (P02, P08, and P34, all on the left), and three non-hypermutator F_1 progeny (P14, P18, and P20, all on the right). Blue and orange bars under the subtelomeric region of each chromosome indicate which parental strain the region was inherited from (orange for Bt65, blue for H99). Red asterisks indicate invasion of Cnl1 into an H99 subtelomeric region that previously had zero Cnl1 copies/fragments. Accurate assembly of telomeric repeat sequences at the end of each chromosome is indicated by a black half circle. Cnl1 length is also indicated by the shade of black for each element.

1147

1148

1149

1150

1151

Supplementary Figure S13. **Genomic diagnostic plot of Bt65 x H99 F$_1$ progeny 25.** For the

14 chromosomes (columns), the log$_2$ (read depth) (top row) and allelic read depth ratio (bottom

row) per genetic variant are shown for the progeny 25. Red and blue colors indicate the

prediction of the allele inherited (Bt65 vs. H99, respectively) at each genetic variant. Allelic read

depth ratios nearing 0.5 suggest both alleles are present for a given genetic variant. Marked

ranges on Chromosomes 3 and 11 log$_2$ (read depth) plots indicate the significant hypermutator

QTL. Black vertical lines depict the boundaries of the centromeres. The boundaries of the *MAT*

locus on Chromosome 5 are shown by vertical green lines.

1160

1161

1162

1163

1164

1165

Supplementary Figure S14. **Visualization of genetic variant filtering criteria.** (**A**) For the 14

chromosomes (columns), the quality scores of the genetic variants (x-axis) vs. the minimum read

depth across the 28 Bt65 x H99 F_1 segregants (y-axis). (**B**) the quality scores of genetic variants

(x-axis) vs. the portion of progeny with the Bt65 allele per genetic variant (y-axis) per

chromosome (columns). (**C**) The portion of progeny with the Bt65 allele per genetic variant (y-

axis) across each chromosome (x-axis). The raw genetic variants are shown in black and the

filtered SNPs, used in analysis, are shown in red.

1173

1174

1175

1176

Supplementary Figure S15. Identification and confirmation of correct Bt65+*ZNF3*

transformants. (A) BtsI restriction enzyme digestion of *ZNF3* PCR products from

nourseothricin-resistant transformants and controls (primers SJP186/187). **(B)** PCR amplification

of the ZNF3 allele using primers outside of the Bt133 *ZNF3* allele used for homologous

recombination (primers SJP208/209). PCR amplification to ensure correct integration of the **(C)**

5' and **(D)** 3' ends of the Bt133 *ZNF3* allele at the endogenous *ZNF3* locus (primers SJP208/187,

and SJP186/209, respectively). PCR to ensure neither **(E)** *CAS9* nor **(F)** the gRNA constructs

were integrated into the transformants (primers JOHE41657/45812 and JOHE50451/50452,

respectively).

1186

1187

1188

1189

1190

1191

Supplementary Figure S16. **Enrichment for dsRNA does not identify any fragments likely**

to be dsRNA mycoviruses. Pictured on the left are RNA samples following LiCl enrichment for

dsRNA run on a 1% agarose gel. Total RNA prior to dsRNA enrichment is pictured on the right

on a 1% agarose gel. *Ms+* is a *Malassezia sympodialis* strain that harbors a dsRNA virus, and

Ms- is a congenic virus-cleared strain[55]. Two biological replicates for all samples are shown and

labeled (1) and (2). The TriDye 1 kb DNA ladder (NEB) was used to estimate RNA fragment

sizes.

1199

1200   **Supplementary Table Legends**

1201   **Supplementary Table S1. Strains included in preliminary screen of SDC isolates for**

1202   **hypermutation phenotype.**

1203

1204   **Supplementary Table S2. Genetic variants and predicted changes in genes within QTL**

1205   **between H99 and Bt65.**

1206

1207   **Supplementary Table S3. (A) Centromere lengths in H99, Bt65, Bt81, Bt89, and Bt133, and**

1208   **(B) one-way ANOVA and Tukey's HSD post hoc statistical tests for differences in mean**

1209   **centromere length.**

1210

1211   **Supplementary Table S4. sRNA analysis**

1212

1213   **Supplementary Table S5. Strains used in this study.**

1214

1215   **Supplementary Table S6. Mutation rates and 95% confidence intervals for all fluctuation**

1216   **assays.**

1217

1218   **Supplementary Table S7. Oligonucleotides used in this study.**

1219

1220   **Supplementary Table S8. Cnl1 insertion sequences in PCR products.**

1221

1222   **Supplementary Table S9. Aneuploid, diploid, and clonal Bt65 x H99 F_1 progeny.**