Genetic compensation of triacylglycerol biosynthesis in the green microalga

Chlamydomonas reinhardtii

Yi-Ying Lee[a], Rudolph Park[b], Stephen M. Miller[b], and Yantao Li[a, c, 1]

[a] Institute of Marine and Environmental Technology, University of Maryland Center for

Environmental Science, Baltimore, MD 21202, USA

[b] Department of Biological Sciences, University of Maryland, Baltimore County, Baltimore, MD

21250, USA

[c] Department of Marine Biotechnology, University of Maryland Baltimore County, Baltimore,

MD 21202, USA

[1] Correspondence: Yantao Li

                701 E Pratt Street, Baltimore, MD 21202

                Phone: (410) 234-8833

                Email: **yantao@umces.edu**

Key words: genetic compensation; triacylglycerol biosynthesis; *Chlamydomonas reinhardtii*;

CRISPR; gene knockout.

Abstract

Genetic compensation through transcriptional adaptation has been proposed to explain phenotypic differences between gene knockouts and knockdowns. With the rapid development of reverse genetic tools such as CRISPR/Cas9 and RNAi in microalgae, it is increasingly important to assess whether genetic compensation affects the phenotype of engineered algal mutants. While exploring triacylglycerol (TAG) biosynthesis pathways in *Chlamydomonas reinhardtii*, it was discovered that knockout of certain genes catalyzing rate-limiting steps of TAG biosynthesis, type-2 diacylglycerol acyltransferase genes (*DGTTs*), triggered genetic compensation under abiotic stress conditions. Genetic compensation of a *DGTT1* null mutation by a related *PDAT* gene was observed regardless of the strain background or mutagenesis approach, e.g., CRISPR/Cas 9 or insertional mutagenesis. However, no compensation was found in the *PDAT* knockout mutant. The effect of *PDAT* knockout was evaluated in a Δ*vtc1* mutant, in which *PDAT* was up-regulated under stress. Knockout of *PDAT* in the Δ*vtc1* background induced a 12.8-fold upregulation of *DGTT1* and a 272.3% increase in TAG content in Δ*vtc1/pdat1* cells, while remaining viable. These data suggest that genetic compensation contributes to the genetic robustness of microalgal TAG biosynthetic pathways, maintaining lipid and redox homeostasis in the knockout mutants under abiotic stress with a mechanism distinct from that found in plants. This work demonstrates examples of genetic compensation in microalgae, implies the physiological relevance of genetic compensation in TAG biosynthesis, and provides guidance for future genetic engineering and mutant characterization efforts.

Introduction

Organisms have evolved various genetic buffering systems to maintain fitness in response to genetic perturbations, including functionally redundant genes (Tautz, 1992), protein feedback loops (Barabási & Oltvai, 2004), and acquisition of adaptive mutations (Teng et al., 2013). Such buffering or compensation systems ensure similar growth or developmental outcomes despite some genetic changes (El-Brolosy & Stainier, 2017). Genetic compensation is defined as the phenomenon whereby the effect of a deleterious mutation is buffered by the genome (Kontarakis & Stainier, 2020). Recently, genetic compensation through transcriptional adaptation, by which a homologous gene or genes are upregulated in response to an early nonsense mutation in a related gene, has been proposed as a mechanism for maintaining genetic robustness in a number of metazoans, including zebrafish, mice, and nematodes, and to explain profound differences between gene knockouts and knockdowns at certain loci (El-Brolosy et al., 2019; Z. Ma et al., 2019; Rossi et al., 2015; Serobyan et al., 2020). Likewise, comparison between gene knockouts and knockdowns in some model plant systems has revealed puzzling discrepencies that suggest transcriptional adaptation is acting to buffer the genome in these species as well (Braun et al., 2008; Chen et al., 2014; Gao et al., 2015; Rodriguez-Leal et al., 2019).

Increasingly, gene knockdown and knockout tools have been deployed in the green microalga *Chlamydomonas reinhardtii* (Chlamydomonas; (Greiner et al., 2017; Jiang, Brueggeman, Horken, Plucinak, & Weeks, 2014; Xiaobo Li et al., 2019; Molnar et al., 2009; Rohr, Sarkar, Balenger, Jeong, & Cerutti, 2004; Schroda, 2006)), but little attention has been paid as to whether these tools might for some genes lead to very different outcomes, and more broadly, whether genetic compensation occurs in green algae (El-Brolosy & Stainier, 2017;

Kontarakis & Stainier, 2020). Here we set out to address these questions as part of an ongoing

study of lipid biosynthesis in Chlamydomonas, using triacylglycerol (TAG) biosynthesis

pathways as examples.

Microalgae store TAG as a carbon and energy-rich compound in cells, whose synthesis

can be stimulated under various stress conditions (Hu et al., 2008; Lenka et al., 2016). Nutrient

deprivation such as nitrogen (N) deprivation and phosphorus (P) deprivation are most commonly

used stressors to induce TAG production. Under N deprivation, algal cell growth is almost

completely arrested while P deprivation promotes TAG production with moderate cell growth

(Iwai, Ikeda, Shimojima, & Ohta, 2014). Biosynthesis of TAG involves sequential acylation to

the precursor glycerol-3-phosphate. Two pathways are involved in the last step of TAG

biosynthesis. One pathway is a *de novo* acyl-CoA-dependent route mediated by diacylglycerol

acyltransferase (DGAT) (Ohlrogge & Browse, 1995). Three major types of DGATs including

two structurally distinctive, membrane bound type-1 and type-2 DGATs and a soluble cytosolic

type-3 DGAT have been identified in plants, algae, and other eukaryotes (Cases et al., 1998;

Lardizabal et al., 2001; Oelkers, Cromley, Padamsee, Billheimer, & Sturley, 2002). The other

TAG biosynthesis pathway is an acyl-CoA-independent route that requires membrane lipid

breakdown and is mediated by a plastidic membrane bound enzyme, phospholipid:diacylglycerol

acyltransferase (PDAT) (Dahlqvist et al., 2000; Yoon, Han, Li, Sommerfeld, & Hu, 2012).

The genome of *C. reinhardtii* harbors one type-1 *DGAT* gene (*DGAT1*), five type-2

DGAT genes (*DGTT1-5*), and one *PDAT* gene (Bagnato et al., 2017; Boyle et al., 2012; Miller et

al., 2010; Yoon et al., 2012). RNA-Seq analysis previously revealed that the abundance of *PDAT*,

DGAT1 and *DGTT1* transcripts was up-regulated, and that of *DGTT2* and *DGTT3* was moderate

but constant, and the abundance of transcripts of *DGTT4* and *DGTT5* was low to undetectable in

response to N deprivation (Boyle et al., 2012). Subsequent RT-qPCR analyses corroborated these

findings for *PDAT*, *DGAT1* and *DGTT1*, while *DGTT2-3* were found to be induced by N

deprivation (Liu, Han, Yoon, Hu, & Li, 2016; Yoon et al., 2012). Similar results were found in

response to P deprivation, where *PDAT*, *DGAT1*, and *DGTT1-4* were up-regulated during TAG

accumulation (Iwai et al., 2014). Functional studies using *in vitro* enzyme assays and gene

knockdowns have largely yielded results consistent with expectations based on these expression

data. *DGTT1-3* and *PDAT* were shown to catalyze TAG biosynthesis *in vitro* (Liu et al., 2016;

Yoon et al., 2012). Knockdown of *DGTT1*, *DGTT2* or *DGTT3* each resulted in a moderately

reduced TAG (TAG-less) phenotype under persistent stress, while knockdown of *PDAT* resulting

in a moderate TAG-less phenotype during both favorable growth and the early phase of stress

induction (Liu et al., 2016; Yoon et al., 2012), suggesting *DGTT*s and *PDAT* are functionally

related genes involved in TAG biosynthesis in Chlamydomonas. However, in some cases,

analysis of gene knockdowns can be limited by off-target effects and low efficiency of gene

silencing (Fedorov et al., 2006; Y. Ma, Creanga, Lum, & Beachy, 2006), so null mutants are

needed to better understand gene function. Among Chlamydomonas TAG biosynthesis genes, so

far only *pdat* mutants have been characterized, which had reduced TAG content under stress

(Boyle et al., 2012). However, whether the *PDAT* insertion mutations analyzed were buffered by

expression of related genes such as *DGTT*s is not known, and up to now no *DGTT* mutants have

been analyzed. So the precise contributions of these genes to TAG accumulation remains obscure.

In this work we analyzed TAG accumulation in strains bearing null mutations in several

genes including *DGTT1-3* and *PDAT*, and found phenotypes that contradicted known

knockdown phenotypes in these genes. Our data showed genetic compensation through up-

regulation of related TAG biosynthetic genes for *dgtt1* and *dgtt2* mutants resulted in a higher

TAG content under stress, which is contrary to the TAG-less phenotype in the respective

knockdown lines. Interestingly, no genetic compensation was found in the *pdat* mutant. However,

when evaluated in the Δ*vtc1* background (defective for Vacuolar Transporter Chaperone 1, a

component of the polyphosphate polymerase complex), where *PDAT* expression and function is

likely critical for storing excess energy as TAG under stress, the *PDAT* knockout resulted in

compensation by up-regulation of *DGTT1* and a higher TAG content. Our data demonstrate

genetic compensation exists in microalgae and imply the physiological importance of genetic

compensation in TAG biosynthesis under stress.

Results

Genetic compensation is present in microalgae

Type-2 DGATs are believed to be major contributors for TAG biosynthesis in algae, but

previous knockdown of Chlamydomonas type-2 *DGAT* genes *DGTT1*, *DGTT2* and *DGTT3*

caused only relatively small (~10 - 20%) suppression of TAG accumulation (Liu et al., 2016).

We set out to determine whether null mutations in these genes might have larger effects, so we

obtained *dgtt1*, *dgtt2* and *dgtt3* CLiP library knockout mutants generated by the Jonikas group

(Xiaobo Li et al., 2019; Xiaobo Li et al., 2016) and analyzed their TAG accumulation. Each

mutant strain carries a CIB1 cassette insertion that disrupts the respective *DGTT* gene. We

further examined the genotype of these mutants and found the *dgtt1* mutant

(LMJ.RY0402.223444) insertion is in the middle of intron 1, the *dgtt2* mutant

(LMJ.RY0402.213587) insertion is at the junction between exon 4 and intron 4, and the *dgtt3*

mutant (LMJ.RY0402.223565) insertion is at the beginning of exon 3 (Fig. 1A; Fig. S1). No

detectable transcripts of *DGTT1*, *DGTT2* and *DGTT3* were found in the respective mutants (Fig.

1C), suggesting all the mutants are null.

TAG biosynthesis was comparatively studied in the *dgtt1*, *dgtt2*, and *dgtt3* mutants and

their parent strain (CC5325; wild-type) under P deprivation, which was reported to induce TAG

accumulation in *C. reinhardtii* while maintaining moderate cell growth (Molnar et al., 2009).

Surprisingly, the *dgtt1* and the *dgtt2* knockout mutants had significantly higher TAG content

than the wild-type strain (t-test, p ≤ 0.05, Fig. 1B) either on a per cell basis or per dry weight

(DW) basis. The TAG content of *dgtt1* and *dgtt2* mutant cells increased by about 90% and 70%

per cell, respectively, and by 45% and 76% per unit dry weight, compared with wild-type cells at

72 h under P deprivation (Fig. 1B). These results are in stark contrast to those for the knockdown

strains, which had reduced TAG content (Liu et al., 2016), and they suggest a compensation

mechanism exists in Chlamydomonas that affects other TAG genes/enzymes in the *dgtt1* and

dgtt2 mutant. By contrast, the *dgtt3* knockout mutant contained ~40% less TAG per cell and per

dry weight compared with the wild-type (Fig. 1B), and notably it had about twice the decrease in

TAG as that caused by the *dgtt3* knockdown (Liu et al., 2016). In Chlamydomonas, *DGTTs* and

PDAT are homologs. Sequence analysis revealed identities between *PDAT* and *DGTT1*, *DGTT2*,

and *DGTT3* were 50.9%, 49.8%, and 49.2%, respectively (Table S1), all above 40% sequence

identity needed to induce genetic compensation by homologs in previous studies (Z. Ma et al.,

2019).

To explore possible genetic compensation in the knockout mutants, the expression of

TAG biosynthetic genes was examined, including that of *DGTT1*, *DGTT2*, *DGTT3*, *PDAT*, and

PGD1 (encodes an MGDG-specific lipase and presumably functions with PDAT in membrane

recycling) (X. Li et al., 2012; Liu et al., 2016; Yoon et al., 2012). In the *dgtt1* mutant, the

expression of *DGTT2*, *DGTT3* and *PGD1* remained unchanged, but the expression of *PDAT* was

upregulated by 9-fold compared with the wild-type at 24 h under P deprivation (Fig. 1C). In the

dgtt2 mutant, the expression of all tested genes was upregulated; compared with the wild-type,

the expression of *DGTT1*, *DGTT3*, *PDAT* and *PGD1* in the *dgtt2* mutant was increased by 19-

fold, 2.6-fold, 9-fold, and 7.6-fold, respectively, at 24 h under P deprivation (Fig. 1C),

correlating with higher TAG accumulation in the *dgtt2* mutant. In the *dgtt3* knockout strain, the

expression level of *PDAT* was significantly higher than for the wild type (t-test, $p \leq 0.05$; Fig.

1C), but that increase did not lead to a functional compensation for TAG accumulation (Fig. 1B).

166   Our findings suggest that a mutation leading to a premature stop codon in two different

type-2 *DGAT* genes likely activates a genetic compensatory network by upregulation of

homologous genes in Chlamydomonas. To determine whether this genetic compensation was

specific to insertional mutants in the CLiP library genetic background, or whether it might be a

more general phenomenon of null mutants with premature stop codons, we used CRISPR/Cas9

to generate a *dgtt1* mutant in a different strain background (*C. reinhardtii* CC3403). In the

resulting mutant (*CC-dgtt1*), the *DGTT1* gene carries a 1229-bp insertion corresponding to

pHS_SaCas9 sequence (Cas9 expression vector) near the beginning of exon 3 of the coding

region (Fig. 2A and Fig. S2A) that results in premature stop codons (Fig. S2B) and disrupts

expression of the *DGTT1* gene. The TAG content of *CC-dgtt1* was comparable to that of the

wild-type on a per cell basis, and 66% higher than the wild-type per unit dry weight under P

deprivation (Fig. 2B). Moreover, this phenotype was maintained under N deprivation (Fig. S3).

Similar to the situation for the *dgtt1* mutant from the CLiP library, RT-qPCR analyses revealed

that *PDAT* expression was upregulated by 6.3-fold in the *CC-dgtt1* mutant, while other TAG

biosynthesis genes (*DGTT2*, *DGTT3* and *PGD1*) remained the same as in the wild-type (Fig. 2C).

Taken together, genetic compensation of *DGTT* seems to occur in Chlamydomonas mutants

regardless of the strain background or early-frameshift mutant allele.

Compensation of *DGTT* knockouts is not reciprocal in the *pdat* mutant

185   Since *C. reinhardtii* cells are capable of compensating defects of *DGTT1* and *DGTT2*

mutations by upregulating related genes such as *PDAT*, we asked whether the mechanism is

reciprocal in the *pdat* mutant. To this end, we assessed TAG accumulation and the expression of

the *DGTT1-3* genes in a *pdat* knockout mutant (CC4502) previously described in (Boyle et al.,

2012). Unlike the *dgtt1 and dgtt2* mutants, the *pdat* mutant had reduced TAG levels (Fig. 3),

similar to the *pdat* knockdown lines (Yoon et al., 2012). The relative TAG content of the *pdat*

knockout mutant was ~30% lower than that of the wild-type strain at 48 h and 72 h under P

deprivation (Fig. 3A), similar to the decrease in TAG content previously reported for this strain

in response to N deprivation (Boyle et al., 2012). In P-deprived *pdat* cells, the expression of

DGTT1-3 were the same as or slightly lower than in the wild-type (Fig. 3B). These data suggest

no genetic compensation of TAG biosynthesis occurs in the *pdat* mutant.

Compensation by upregulation of *DGTT1* in a *Δvtc1/pdat1* double mutant

198   Our previous work suggests *PDAT* primarily regulates TAG biosynthesis under favorable,

nutrient-replete conditions (Yoon et al., 2012) but not under persistent nutrient-deficient stress

conditions (Liu et al., 2016). A possible explanation for lack of genetic compensation in the *pdat*

mutant is thus the relatively poor function of *PDAT* under stress conditions, in which *DGTTs* are

already highly expressed and responsible for TAG biosynthesis. Previous work showed P

metabolism, particularly P uptake, was impaired in a Δ*vtc1* mutant, in which the vacuolar

transport chaperon (VTC) complex was disrupted (Aksoy, Pootakham, & Grossman, 2014;

Plouviez et al., 2021; Sanz-Luque, Saroussi, Huang, Akkawi, & Grossman, 2020; Schmollinger

et al., 2021). As such, in this Δ*vtc1* mutant we reasoned that *PDAT* expression and function may

be especially critical to recycle phospholipids and balance P metabolism under stress. To test this

idea, we analyzed TAG biosynthesis in the Δ*vtc1* mutant and in a *VTC1*-rescued strain (*VTC1*)

under P replete and P-depleted conditions. In Δ*vtc1* cells TAG content was 90% higher (*t*-test, p

≤ 0.05) than the control under P-deprived conditions but not under P-replete conditions (Fig. 4A

and B), suggesting loss of *VTC1* had induced TAG biosynthesis under stress. When we analyzed

the TAG biosynthetic genes in P-depleted Δ*vtc1* cells, only the *PDAT* gene showed an increase

in expression (2.8-fold; Fig. 4C) while none of the *DGTT* or *PGD* genes were upregulated,

suggesting a role of *PDAT* in controlling TAG biosynthesis under stress in the Δ*vtc1* mutant.

We predicted that for the Δ*vtc1* mutant, in which *PDAT* function is critical, if *PDAT*

expression is interrupted, genetic compensation by other TAG biosynthetic genes may be

necessary for the resulting mutant to survive under stress. To test this idea, we generated a

Δ*vtc1*/*pdat* double mutant by crossing the Δ*vtc1* (CC5165) and the *pdat* (CC4502) single mutant

strains. Interestingly, the TAG level in the resulting strain increased compared with the control

and single mutant (Fig. 4B). After 48 h of P deprivation, the TAG content of Δ*vtc1*/*pdat* cells

was about 272.3% higher than the *VTC1* control and 95% higher than the Δ*vtc1* mutant (Fig. 4B).

We found that under stress, *DGTT1* transcripts were upregulated in Δ*vtc1*/*pdat* cells compared

with the control (by12.4-fold), but no other TAG biosynthetic genes were transcriptionally

upregulated (Fig. 4C). Moreover, Δ*vtc1*/*pdat* cells remained viable with moderate growth under

P deprviation. Δ*vtc1*/*pdat* cell density slightly increased from 0.69 x 10^6 cells/ml to 1.01 x 10^6

cells/ml between day 0 and day 4 (Fig. S4A), while the per cell dry weight of the Δ*vtc1*/*pdat*

mutant about doubled compared with the control (*VTC1*) (Fig. S4B). By contrast, the growth rate

and per cell dry weight of Δ*vtc1* cells were similar to that of the control (Fig. S4). Therefore, our

results suggest that upregulation of *PDAT* in the Δ*vtc1* mutant results in an increase in TAG

accumulation, while the effect of *PDAT* disruption in the Δ*vtc1* background is buffered by

upregulation of *DGTT1* in the *Δvtc1/pdat1* double mutant.

Discussion

Genetic compensation, and more specifically, transcriptional adaptation through

upregulation of related genes, is proposed as an underlying mechanism to explain discrepencies

between knockdown and knockout phenotypes in several model systems including zebrafish and

mouse (El-Brolosy et al., 2019; Z. Ma et al., 2019; Rossi et al., 2015), *Caenorhabditis elegans*

(Serobyan et al., 2020), Arabidopsis (Braun et al., 2008; Chen et al., 2014; Gao et al., 2015), and

tomato (Rodriguez-Leal et al., 2019). One mechanism for this phenomenon involves the

nonsense-mediated mRNA decay pathway that is activated by early nonsense mutations (El-

Brolosy et al., 2019; Z. Ma et al., 2019). However, it is not known whether this phenomenon

exists across eukaryotes, including microalgae (El-Brolosy & Stainier, 2017; Kontarakis &

Stainier, 2020). Here, we demonstrate that compensation by related genes buffers the effect of

deleterious mutations in Chlamydomonas lipid biosynthesis genes, in a process that resembles

transcriptional adaptation. Knockouts of *DGTT1* triggered upregulation of the functionally

related and homologous *PDAT* gene regardless of the strain background, mutagenesis approach,

or stress conditions (P or N deprivation), while knockout of *DGTT2* triggered upregulation of

PDAT, *DGTT1* and *PGD1* (Fig. 1 and Fig. 2). In the *dgtt1*, *CC-dgtt1* and *dgtt2* mutant, the TAG-

less phenotype found in respective knockdown lines (Liu et al., 2016) was not observed. Instead,

these mutants produced more TAG than the control, likely due to transcriptional adaptation of

homologous genes. This is unlikely the case for Arabidopsis mutants defective in TAG

biosynthestic genes, where knockout of *DGAT1* and *PDAT1* resulted in a reduced TAG

phenotype in oilseeds and vegetative tissues, respectively (Fan, Yan, & Xu, 2013; Zhang, Fan,

Taylor, & Ohlrogge, 2009). Arabidopsis *DGAT1* and *PDAT1* are suggested to be functionally

complementary genes essential for TAG biosynthesis and normal plant development (Fan et al.,

2013; Zhang et al., 2009). Whether genetic compensation contributes to the observed phenotype

in Arabidopsis mutants defective in TAG biosynthetic genes requires further study.

Recent advances in reverse genetic tools including CRISPR, TALEN, and ZFN genome

editing methods have greatly accelerated progress in generating knockout mutants in traditional

and new algal model systems, including Chlamydomonas (Greiner et al., 2017; Jiang et al., 2014;

Park, Asbury, & Miller, 2020), *Nannochloropsis* (Poliner, Takeuchi, Du, Benning, & Farré, 2018;

Wang et al., 2016), *Volvox carteri* (Ortega-Escalante, Jasper, & Miller, 2019), and *Ectocarpus*

(Badis et al., 2021). Moreover, there is a genome-wide insertional mutant pool, the CLiP library

(Xiaobo Li et al., 2019), for Chlamydomonas. Such tools and mutant libraries dramatically

facilitate analysis of gene function. However, as we demonstrated here, genetic compensation by

related genes might, in some cases, offset the effects of a null mutation, confounding genetic

analysis. Likewise, previous efforts to screen and characterize TAG-less Chlamydomonas

mutants from the CLiP library failed to identify any *DGTT* knockout mutants (XB Li, personal

communication), possibly due to genetic compensation, while knockdown lines of *DGTTs*

clearly showed a reduced-TAG phenotype (Liu et al., 2016). Therefore, when studying gene

function using null mutants, preferably multiple mutant strains targeting the same gene should be

obtained by a targeted genome editing method (e.g. CRISPR) or from the CLiP libarary (because

some might activate genetic compensation while others might not), and these mutants should be

carefully evaluated for their genotypic and phenotypic traits. If genetic compensation is observed

in the mutants, alternative methods such as knockdown should be used to confirm the respective

gene function. Ideally, deletion mutants should be analyzed as they will not trigger

transcriptional adaptation through the nonsense-mediated mRNA decay pathway (El-Brolosy et

al., 2019; Z. Ma et al., 2019), though to date a relatively small number of deletion mutants have

been generated in Chlamydomonas due to current technical limitations.

In Chlamydomonas, biosynthesis of TAG has been implicated in the protection of the

photosynthetic electron transport chain from over-reduction during stress conditions, and TAG

accumulation is suggested to be essential for cells to survive under stress (Du et al., 2018; X. Li

et al., 2012). Our initial experiments raised two interesting questions. First, does genetic

compensation of TAG biosynthesis, as appears to occur in the *dgtt1* and *dgtt2* mutants, play an

essential survival role under persistent stress conditions, and second, might genetic compensation

occur for other mutated TAG biosynthetic genes? To address these questions, we first examined

the *pdat* mutant. Unlike the *dgtt1* and *dgtt2* mutants, it has a similar TAG-less phenotype and

TAG biosynthetic gene expression profiles when stressed as the knockdown lines (Fig. 3 and

(Boyle et al., 2012; Yoon et al., 2012)), suggesting no genetic compensation in the *pdat* mutant.

This outcome may be due to the poor function of *PDAT* in stress conditions, under which *DGTTs*

are primarily responsible for TAG biosynthesis (Liu et al., 2016; Yoon et al., 2012). Consistent

with this notion is a recent *PDAT* analysis in the green lineage showing that *PDAT* plays a less

important role in algae than in plants (Falarz et al., 2020). Disruption of *AtPDAT1* in *A. thaliana*

led to an over 50% decrease in oil accumulation in growing tissues and significant gametophytic

and growth defects (Fan et al., 2013), whereas knockout of Chlamydomonas *PDAT* reduced

TAG content by only 25% - 30%, and *pdat* cells had a similar growth rate as the wild type (Fig

3A) (Boyle et al., 2012).

To determine whether there might be compensation for the *PDAT* knockout when PDAT

function is most essential, we evaluated the effect of the *PDAT* knockout in a Δ*vtc1* mutant, in

which *PDAT* expression and function are likely critical for Δ*vtc1* cells to store excess energy as

TAG under stress (Fig. 4). We expected that in the absence of compensation, the knockout of

PDAT in the Δ*vtc1* background should result in a TAG-less phenotype. Intriguingly, the

Δ*vtc1*/*pdat* mutant exhibited a 272.3% higher TAG content than the *VTC1* control, likely because

of a large (12.8-fold) upregulation of *DGTT1* (Fig. 4). The Δ*vtc1*/*pdat* mutant remained viable

under stress, as its cell density and dry weight increased in response to P deprivation (Fig. S4).

So while no genetic compensation was observed in the *pdat* mutant, in the Δ*vtc1*/*pdat* double

mutant, *DGTT1* was highly overexpressed, and more TAG was produced under P deprivation.

Under stress conditions, these cells continue to capture light energy through photosynthesis, and

if electron acceptors become overreduced due to the slow growth, cytotoxic reactive oxygen

species are produced that lead to cell death (Y. Li, Sommerfeld, Chen, & Hu, 2008). TAG

biosynthesis is likely essential for Δ*vtc1*/*pdat* cells to survive during stress because TAG is the

most reduced electron sink in algae (Hu et al., 2008). In line with our findings, it was previously

reported that a Chlamydomonas mutant defective for the MGDG-specific lipase PGD1

accumulated less TAG than the wild type while its cells lost viability under stress, suggesting

TAG plays an essential role as an electron and energy sink under stress conditions, thereby

preventing generation of reactive oxygen species (Du et al., 2018; X. Li et al., 2012). Together,

these data suggest that under stress conditions, loss of TAG biosynthetic gene function can be

buffered by upregulation of a related gene or genes through transcriptional adaptation in algae.

In summary, we have demonstrated that genetic compensation contributes to genetic

robustness in microalgal TAG biosynthesis, maintaining TAG biosynthesis and redox

homeostasis in knockout mutants under abiotic stress. This work demonstrates, to the best of our

knowledge, the first reported example of genetic compensation in microalgae, implies the

physiological relevance of genetic compensation in TAG biosynthesis under stress, and provides

guidance for future genetic engineering and mutant characterization efforts. The exact

mechanisms inducing genetic compensation in the Chlamydomonas mutants are not understood

and warrant further investigation. Since transcriptional adaptation in other systems is triggered

by mutant mRNA decay and involves Upf3a and COMPASS components (El-Brolosy et al.,

2019; Z. Ma et al., 2019), testing the involvement of these components in algal genetic

compensation is a logical next step.

Materials and Methods

Strains and growth conditions

Chlamydomonas reinhardtii LMJ.RY0402.223444, LMJ.RY0402.213587,

LMJ.RY0402.223565 and their parent strain CC5325 (*cw15*; mt$^-$) (Xiaobo Li et al., 2016),

CC4502 (*pdat1-1*; mt$^+$) and its parent strain CC4425 [D66] (*cw15*; mt$^+$) (Boyle et al., 2012),

CC3403 (*arg7*; *cw15*; mt$^-$), Δ*vtc1* mutant strain CC5165 (Δ*vtc1*; mt$^-$) and the *VTC1* rescue strain

CC5166 (*ars76::VTC1*) (Aksoy et al., 2014) were obtained from the Chlamydomonas Resource

Center (http://www.chlamycollection.org). CC3403 was used as the wild-type in CRISPR/Cas9

mutagenesis (Greiner et al., 2017). The Δ*vtc1/pdat* double mutant was derived from a cross

between CC5165 and CC4502 (methods described below). All strains were maintained in Tris-

acetate-phosphate (TAP) medium containing paramomycin as necessary. P deprivation was

imposed by transferring *C. reinhardtii* cells to P-free TAP, in which potassium phosphate was

replaced by 1.5 mM KCl. Ammonium chloride was omitted to generate N-free TAP. All batch

cultures were grown under continuous light illumination at 60 μmol photons m^{-2} s^{-1}, shaking at

150 rpm, room temperature.

Gene expression measurement by RT-qPCR

Total RNA was extracted and purified using Trizol reagent (Invitrogen) and RNeasy mini

kit (Qiagen). cDNA was synthesized using Protoscript II First Strand cDNA synthesis kit (NEB).

The expression of target genes was determined by RT-qPCR in Applied Biosystems 7500 using

the $2^{-\Delta\Delta CT}$ method normalized by the expression of 18s rRNA. Details are described in the

Supplemental Materials and Methods. The RT-qPCR analysis for each targeted gene was

performed twice, with three biological replicates and three technical replicates for each sample.

The Primer sequences used in RT-qPCR are listed in Table S2.

Generation and screening of *Chlamydomonas reinhardtii* mutants

The CC-*dgtt1* mutant was generated by the CRISPR/Cas9 method developed by Greiner

et al (Greiner et al., 2017). The parent strain of the *CC-dgtt1* mutant (CC3403), the SaCas9

expression vector (pHS_SaCas9) and the sgRNA cloning vector (pCrU6.4-SaCloning-aphVIII)

were obtained from the Chlamydomonas Resource Center. In brief, a customized targeting vector

that expressed sgRNA specifically targeting *DGTT1* was constructed in pCrU6.4-SaCloning-

aphVIII. The SaCas9 expression vector and the customized targeting vector that co-transformed

into CC3403 by electroporation (Park et al., 2020; Serobyan et al., 2020). The transformants

were selected for paromomycin resistance and screened by PCR using oligonucleotides

CrDGTT1F and CrDGTT1R (sequences in Table S2) for insertions at the *DGTT1* locus. The

candidates yielding PCR products larger than the expected size for the wild type *DGTT1* region

were sequenced. The detailed procedures are described in the Supplemental Materials and

Methods.

The Δ*vtc1*/*pdat1* double mutant was generated by mating as described in Goodenough,

1976 (Goodenough, Hwang, & Martin, 1976) with some modifications (procedures described in

the Supplemental Materials and Methods). The crossed candidates were selected for

paramomycin resistance and subsequently screened by colony PCR for a co-occurrence with the

deletion of the *VTC1* gene and the presence of the *pdat1-1* mutation (Boyle et al., 2012) in the

genome. Primers CrVTC1-pF1 and CrVTC1-stopR1 (sequences in Table S2) were used for

determination of the presence (yielding a 1148 bp PCR product) or the absence (yielding no PCR

products) of the *VTC1* gene. Primers CrPDAT1-1-F2 and RIM5-2 (sequences in Table S2)

(Boyle et al., 2012) were used to determine the presence (~ 1 kb PCR product) or the absence (no

PCR product) of the *AphIII* insertion in the *PDAT* gene.

TAG content measurement

Total lipid from *C. reinhardtii* cells was extracted using chloroform: methanol (2:1, v/v)

as previously described (Yoon et al., 2012). The lipid extracts were dried under a gentle stream

of nitrogen gas and re-dissolved in chloroform. Total lipid was resolved by thin-layer

chromatography (TLC) on a silica gel 60 F_{254} plates (EMD Millipore) using hexane: t-butyl

methyl ether: acetic acid (80:20:2, v/v/v) solvent mixture as a mobile phase to develop TAG. For

visualization, the developed TLC plates were sprayed with 8% (w/v) H_3PO_4 solution containing

10% (w/v) copper (II) sulfate pentahydrate, and then charred at 180°C for 3 min. The relative

intensity of signals corresponding to TAG can be measured using ImageJ software (NIH). The

absolute quantification of TAG can be calculated by the sum of fatty acid fractions composed of

TAG using gas chromatography-mass spectrometry (GC-MS) as previously described (Yantao Li

et al., 2010; Ohlrogge & Browse, 1995). Details are described in the Supplemental Materials and

Methods.

Statistical analysis

Statistical analysis was performed by Student's *t*-test with two-tailed distribution. *p* value ≤ 0.05

was the criteria to determine statistical significance.

Acknowledgements

This work was supported in part by grants from the U.S. National Science Foundation

(CBET-1511939) and DOE Office of Fossil Energy (FE-0031914) to YL, and by NSF 1332344

to SMM. We thank Drs. William Snell, Peeyush Ranjan, and Mayanka Awasthi at University of

Maryland College park for their help with generating double knockout mutants, Dr. Arthur

Grossman at Carnegie Institution for Science for providing the *Chlamydomonas vtc1* strains used

in this study, Dr. Xiaobo Li at Westlake University for discussion of unpublished mutant

screening data, and Drs. Fan Zhang and Russell Hill at the Institute of Marine and Environmental

Technology for their help with polyP analysis. We thank Mr. Mohamed Mahmoud-Aly at Cairo

University for his help with *Chlamydomonas* mutant characterization.

Competing financial interests

The authors declare no competing financial interests.

References

Aksoy, M., Pootakham, W., & Grossman, A. R. (2014). Critical Function of a *Chlamydomonas*
reinhardtii Putative Polyphosphate Polymerase Subunit during Nutrient Deprivation. *The*
Plant Cell, 26(10), 4214-4229. doi: 10.1105/tpc.114.129270

Badis, Y., Scornet, D., Harada, M., Caillard, C., Godfroy, O., Raphalen, M., . . . Cock, J. M.
(2021). Targeted CRISPR-Cas9-based gene knockouts in the model brown alga
Ectocarpus. *New Phytol*. doi: 10.1111/nph.17525
Bagnato, C., Prados, M. B., Franchini, G. R., Scaglia, N., Miranda, S. E., & Beligni, M. V.
(2017). Analysis of triglyceride synthesis unveils a green algal soluble diacylglycerol
acyltransferase and provides clues to potential enzymatic components of the chloroplast
pathway. *BMC Genomics, 18*(1), 223. doi: 10.1186/s12864-017-3602-0
Barabási, A. L., & Oltvai, Z. N. (2004). Network biology: understanding the cell's functional
organization. *Nat Rev Genet, 5*(2), 101-113. doi: 10.1038/nrg1272
Boyle, N. R., Page, M. D., Liu, B., Blaby, I. K., Casero, D., Kropat, J., . . . Merchant, S. S.
(2012). Three acyltransferases and nitrogen-responsive regulator are implicated in
nitrogen starvation-induced triacylglycerol accumulation in *Chlamydomonas*. *Journal of
Biological Chemistry, 287*(19), 15811-15825. doi: 10.1074/jbc.M111.334052
Braun, N., Wyrzykowska, J., Muller, P., David, K., Couch, D., Perrot-Rechenmann, C., &
Fleming, A. J. (2008). Conditional repression of AUXIN BINDING PROTEIN1 reveals
that it coordinates cell division and cell expansion during postembryonic shoot
development in Arabidopsis and tobacco. *Plant Cell, 20*(10), 2746-2762. doi:
10.1105/tpc.108.059048
Cases, S., Smith, S. J., Zheng, Y. W., Myers, H. M., Lear, S. R., Sande, E., . . . Farese, R. V., Jr.
(1998). Identification of a gene encoding an acyl CoA:diacylglycerol acyltransferase, a
key enzyme in triacylglycerol synthesis. *Proc Natl Acad Sci U S A, 95*(22), 13018-13023.
doi: 10.1073/pnas.95.22.13018
Chen, X., Grandont, L., Li, H., Hauschild, R., Paque, S., Abuzeineh, A., . . . Friml, J. (2014).
Inhibition of cell expansion by rapid ABP1-mediated auxin effect on microtubules.
Nature, 516(7529), 90-93. doi: 10.1038/nature13889
Dahlqvist, A., Stahl, U., Lenman, M., Banas, A., Lee, M., Sandager, L., . . . Stymne, S. (2000).
Phospholipid:diacylglycerol acyltransferase: an enzyme that catalyzes the acyl-CoA-
independent formation of triacylglycerol in yeast and plants. *Proc Natl Acad Sci U S A,
97*(12), 6487-6492. doi: 10.1073/pnas.120067297
Du, Z. Y., Lucker, B. F., Zienkiewicz, K., Miller, T. E., Zienkiewicz, A., Sears, B. B., . . .
Benning, C. (2018). Galactoglycerolipid Lipase PGD1 Is Involved in Thylakoid
Membrane Remodeling in Response to Adverse Environmental Conditions in
Chlamydomonas. *Plant Cell, 30*(2), 447-465. doi: 10.1105/tpc.17.00446
El-Brolosy, M. A., Kontarakis, Z., Rossi, A., Kuenne, C., Günther, S., Fukuda, N., . . . Stainier,
D. Y. R. (2019). Genetic compensation triggered by mutant mRNA degradation. *Nature,
568*(7751), 193-197. doi: 10.1038/s41586-019-1064-z
El-Brolosy, M. A., & Stainier, D. Y. R. (2017). Genetic compensation: A phenomenon in search
of mechanisms. *PLoS genetics, 13*(7), e1006780-e1006780. doi:
10.1371/journal.pgen.1006780
Falarz, L. J., Xu, Y., Caldo, K. M. P., Garroway, C. J., Singer, S. D., & Chen, G. (2020).
Characterization of the diversification of phospholipid:diacylglycerol acyltransferases in
the green lineage. *Plant J, 103*(6), 2025-2038. doi: 10.1111/tpj.14880
Fan, J., Yan, C., & Xu, C. (2013). Phospholipid:diacylglycerol acyltransferase-mediated
triacylglycerol biosynthesis is crucial for protection against fatty acid-induced cell death
in growing tissues of Arabidopsis. *Plant J, 76*(6), 930-942. doi: 10.1111/tpj.12343

Fedorov, Y., Anderson, E. M., Birmingham, A., Reynolds, A., Karpilow, J., Robinson, K., . . . Khvorova, A. (2006). Off-target effects by siRNA can induce toxic phenotype. *RNA (New York, N.Y.), 12*(7), 1188-1196. doi: 10.1261/rna.28106

Gao, Y., Zhang, Y., Zhang, D., Dai, X., Estelle, M., & Zhao, Y. (2015). Auxin binding protein 1 (ABP1) is not required for either auxin signaling or Arabidopsis development. *Proc Natl Acad Sci U S A, 112*(7), 2275-2280. doi: 10.1073/pnas.1500365112

Goodenough, U. W., Hwang, C., & Martin, H. (1976). Isolation and genetic analysis of mutant strains of Chlamydomonas reinhardi defective in gametic differentiation. *Genetics, 82*(2), 169-186.

Greiner, A., Kelterborn, S., Evers, H., Kreimer, G., Sizova, I., & Hegemann, P. (2017). Targeting of Photoreceptor Genes in Chlamydomonas reinhardtii via Zinc-Finger Nucleases and CRISPR/Cas9. *Plant Cell, 29*(10), 2498-2518. doi: 10.1105/tpc.17.00659

Hu, Q., Sommerfeld, M., Jarvis, E., Ghirardi, M., Posewitz, M., Seibert, M., & Darzins, A. (2008). Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances. *Plant J, 54*, 621-639.

Iwai, M., Ikeda, K., Shimojima, M., & Ohta, H. (2014). Enhancement of extraplastidic oil synthesis in Chlamydomonas reinhardtii using a type-2 diacylglycerol acyltransferase with a phosphorus starvation-inducible promoter. *Plant Biotechnology Journal, 12*(6), 808-819. doi: 10.1111/pbi.12210

Jiang, W., Brueggeman, A. J., Horken, K. M., Plucinak, T. M., & Weeks, D. P. (2014). Successful transient expression of Cas9/sgRNA genes in Chlamydomonas reinhardtii. *Eukaryotic Cell*. doi: 10.1128/ec.00213-14

Kontarakis, Z., & Stainier, D. Y. R. (2020). Genetics in Light of Transcriptional Adaptation. *Trends in Genetics, 36*(12), 926-935. doi: 10.1016/j.tig.2020.08.008

Lardizabal, K. D., Mai, J. T., Wagner, N. W., Wyrick, A., Voelker, T., & Hawkins, D. J. (2001). DGAT2 is a new diacylglycerol acyltransferase gene family: purification, cloning, and expression in insect cells of two polypeptides from Mortierella ramanniana with diacylglycerol acyltransferase activity. *J Biol Chem, 276*(42), 38862-38869. doi: 10.1074/jbc.M106168200

Lenka, S. K., Carbonaro, N., Park, R., Miller, S. K., Thorpe, I., & Li, Y. T. (2016). Current advances in molecular, biochemical, and computational modeling analysis of microalgal triacylglycerol biosynthesis. *Biotechnology Advances, 34*, 1046-1063. doi: 10.1016/j.biotechadv.2016.06.004

Li, X., Moellering, E. R., Liu, B., Johnny, C., Fedewa, M., Sears, B. B., . . . Benning, C. (2012). A galactoglycerolipid lipase is required for triacylglycerol accumulation and survival following nitrogen deprivation in Chlamydomonas reinhardtii. *Plant Cell., 24*. doi: 10.1105/tpc.112.105106

Li, X., Patena, W., Fauser, F., Jinkerson, R. E., Saroussi, S., Meyer, M. T., . . . Jonikas, M. C. (2019). A genome-wide algal mutant library and functional screen identifies genes required for eukaryotic photosynthesis. *Nature Genetics, 51*(4), 627-635. doi: 10.1038/s41588-019-0370-6

Li, X., Zhang, R., Patena, W., Gang, S. S., Blum, S. R., Ivanova, N., . . . Jonikas, M. C. (2016). An Indexed, Mapped Mutant Library Enables Reverse Genetics Studies of Biological Processes in *Chlamydomonas reinhardtii*. *The Plant Cell, 28*(2), 367-387. doi: 10.1105/tpc.15.00465

Li, Y., Han, D., Hu, G., Dauvillee, D., Sommerfeld, M., Ball, S., & Hu, Q. (2010).
Chlamydomonas starchless mutant defective in ADP-glucose pyrophosphorylase hyper-
accumulates triacylglycerol. *Metabolic Engineering, 12*, 387-391. doi: DOI:
10.1016/j.ymben.2010.02.002
Li, Y., Sommerfeld, M., Chen, F., & Hu, Q. (2008). Consumption of oxygen by astaxanthin
biosynthesis: A protective mechanism against oxidative stress in *Haematococcus*
pluvialis (Chlorophyceae). *J Plant Physiol*. doi: 10.1016/j.jplph.2007.12.007
Liu, J., Han, D., Yoon, K., Hu, Q., & Li, Y. (2016). Characterization of type 2 diacylglycerol
acyltransferases in Chlamydomonas reinhardtii reveals their distinct substrate
specificities and functions in triacylglycerol biosynthesis. *Plant Journal, 86*, 3-19. doi:
10.1111/tpj.13143
Ma, Y., Creanga, A., Lum, L., & Beachy, P. A. (2006). Prevalence of off-target effects in
Drosophila RNA interference screens. *Nature, 443*(7109), 359-363. doi:
10.1038/nature05179
Ma, Z., Zhu, P., Shi, H., Guo, L., Zhang, Q., Chen, Y., . . . Chen, J. (2019). PTC-bearing mRNA
elicits a genetic compensation response via Upf3a and COMPASS components. *Nature,*
568(7751), 259-263. doi: 10.1038/s41586-019-1057-y
Miller, R., Wu, G., Deshpande, R. R., Vieler, A., Gartner, K., Li, X., . . . Benning, C. (2010).
Changes in transcript abundance in Chlamydomonas reinhardtii following nitrogen
deprivation predict diversion of metabolism. *Plant Physiol, 154*(4), 1737-1752. doi:
10.1104/pp.110.165159
Molnar, A., Bassett, A., Thuenemann, E., Schwach, F., Karkare, S., Ossowski, S., . . .
Baulcombe, D. (2009). Highly specific gene silencing by artificial microRNAs in the
unicellular alga *Chlamydomonas reinhardtii*. *Plant Journal, 58*(1), 165-174.
Oelkers, P., Cromley, D., Padamsee, M., Billheimer, J. T., & Sturley, S. L. (2002). The DGA1
gene determines a second triglyceride synthetic pathway in yeast. *J Biol Chem, 277*(11),
8877-8881. doi: 10.1074/jbc.M111646200
Ohlrogge, J., & Browse, J. (1995). Lipid biosynthesis. *Plant Cell, 7*(7), 957-970.
Ortega-Escalante, J. A., Jasper, R., & Miller, S. M. (2019). CRISPR/Cas9 mutagenesis in Volvox
carteri. *The Plant Journal, 97*(4), 661-672. doi: https://doi.org/10.1111/tpj.14149
Park, R. V., Asbury, H., & Miller, S. M. (2020). Modification of a Chlamydomonas reinhardtii
CRISPR/Cas9 transformation protocol for use with widely available electroporation
equipment. *MethodsX, 7*, 100855. doi: https://doi.org/10.1016/j.mex.2020.100855
Plouviez, M., Fernández, E., Grossman, A. R., Sanz-Luque, E., Sells, M., Wheeler, D., &
Guieysse, B. (2021). Responses of Chlamydomonas reinhardtii during the transition from
P-deficient to P-sufficient growth (the P-overplus response): The roles of the vacuolar
transport chaperones and polyphosphate synthesis. *Journal of Phycology, 57*(3), 988-
1003. doi: https://doi.org/10.1111/jpy.13145
Poliner, E., Takeuchi, T., Du, Z. Y., Benning, C., & Farré, E. M. (2018). Nontransgenic Marker-
Free Gene Disruption by an Episomal CRISPR System in the Oleaginous Microalga,
Nannochloropsis oceanica CCMP1779. *ACS Synth Biol, 7*(4), 962-968. doi:
10.1021/acssynbio.7b00362
Rodriguez-Leal, D., Xu, C., Kwon, C.-T., Soyars, C., Demesa-Arevalo, E., Man, J., . . . Lippman,
Z. B. (2019). Evolution of buffering in a genetic circuit controlling plant stem cell
proliferation. *Nature Genetics, 51*(5), 786-792. doi: 10.1038/s41588-019-0389-8

Rohr, J., Sarkar, N., Balenger, S., Jeong, B. R., & Cerutti, H. (2004). Tandem inverted repeat
system for selection of effective transgenic RNAi strains in Chlamydomonas. *Plant*
Journal, 40(4), 611-621.
Rossi, A., Kontarakis, Z., Gerri, C., Nolte, H., Hölper, S., Krüger, M., & Stainier, D. Y. R.
(2015). Genetic compensation induced by deleterious mutations but not gene
knockdowns. *Nature, 524*(7564), 230-233. doi: 10.1038/nature14580
Sanz-Luque, E., Saroussi, S., Huang, W., Akkawi, N., & Grossman, A. R. (2020). Metabolic
control of acclimation to nutrient deprivation dependent on polyphosphate synthesis. *Sci*
Adv, 6(40). doi: 10.1126/sciadv.abb5351
Schmollinger, S., Chen, S., Strenkert, D., Hui, C., Ralle, M., & Merchant, S. S. (2021). Single-
cell visualization and quantification of trace metals in Chlamydomonas lysosome-related
organelles. *Proc Natl Acad Sci U S A, 118*(16). doi: 10.1073/pnas.2026811118
Schroda, M. (2006). RNA silencing in Chlamydomonas: mechanisms and tools. *Current*
Genetics, 49(2), 69-84. doi: 10.1007/s00294-005-0042-1
Serobyan, V., Kontarakis, Z., El-Brolosy, M. A., Welker, J. M., Tolstenkov, O., Saadeldein, A.
M., . . . Stainier, D. Y. R. (2020). Transcriptional adaptation in Caenorhabditis elegans.
eLife, 9, e50014. doi: 10.7554/eLife.50014
Tautz, D. (1992). Redundancies, development and the flow of information. *Bioessays, 14*(4),
263-266. doi: 10.1002/bies.950140410
Teng, X., Dayhoff-Brannigan, M., Cheng, W.-C., Gilbert, Catherine E., Sing, Cierra N., Diny,
Nicola L., . . . Hardwick, J. M. (2013). Genome-wide Consequences of Deleting Any
Single Gene. *Molecular Cell, 52*(4), 485-494. doi:
https://doi.org/10.1016/j.molcel.2013.09.026
Wang, Q., Lu, Y., Xin, Y., Wei, L., Huang, S., & Xu, J. (2016). Genome editing of model
oleaginous microalgae Nannochloropsis spp. by CRISPR/Cas9. *Plant J, 88*. doi:
10.1111/tpj.13307
Yoon, K., Han, D., Li, Y., Sommerfeld, M., & Hu, Q. (2012). Phospholipid:diacylglycerol
acyltransferase is a multifunctional enzyme involved in membrane lipid turnover and
degradation while synthesizing triacylglycerol in the unicellular green microalga
Chlamydomonas reinhardtii. *Plant Cell, 24*, 3708-3724. doi: 10.1105/tpc.112.100701
Zhang, M., Fan, J., Taylor, D. C., & Ohlrogge, J. B. (2009). DGAT1 and PDAT1
acyltransferases have overlapping functions in Arabidopsis triacylglycerol biosynthesis
and are essential for normal pollen and seed development. *The Plant Cell, 21*(12), 3885-
3901. doi: 10.1105/tpc.109.071795

Fig 1. Genetic structure, phenotype and gene expression of the *dgtt1*, *dgtt2*, *dgtt3* knockout mutants obtained from the CLiP library. (A) Locations of the insertion sites of the CIB1 cassette (brown triangles) and of the primer pairs (black arrows) used to detect gene expression in the *DGTT1*, *DGTT2* and *DGTT3* genes are presented. (B) TAG content of the *dgtt1* (blue), *dgtt2* (red), *dgtt3* (green) mutants and the parent strain CC5325 (black, WT) grown was measured at 72 h under P deprivation. (C) Relative transcriptional level of selected TAG biosynthesis genes (*DGTT1*, *DGTT2*, *DGTT3*, *PDAT* and *PGD1*) in the *dgtt1* (blue), *dgtt2* (red), *dgtt3* (green) mutants and the parent strain CC5325 (black) was measured at 24 h under P deprivation. Transcripts of *DGTT1* in the *dgtt1* mutant and *DGTT2* in the *dgtt2* mutant were nondetectable (N/D). Data represents mean ± standard deviation (SD) from three biological repeats. An asterisk indicates statistical significance by Student's *t*-test (p value ≤ 0.05) when compared to WT.

Fig 2. Genetic structure, phenotype and gene expression of the CRISPR-derived *dgtt1* knockout mutant. (A) The location of an *AphIII* cassette insertion (green triangle) generated by CRISPR in the *DGTT1* gene are presented. (B) TAG content of the *CC-dgtt1* mutant (blue) and the parent strain CC3403 (black, WT) grown was measured at 72 h under P deprivation. (C) Relative transcriptional level of selected TAG biosynthesis genes (*DGTT2*, *DGTT3*, *PDAT* and *PGD1*) in the *dgtt1* mutant (blue) and the parent strain CC3403 (black) was measured at 48 h under P deprivation. Data represents mean ± standard deviation (SD) from three biological repeats. An asterisk indicates statistical significance by Student's *t*-test (*p* value ≤ 0.05).

Fig 3. Phenotype and gene expression of the *pdat* mutant (CC4502). (A) Relative concentrations of TAG from the *pdat* knockout strain CC4502 (gray) and the wild-type CC4425 (black, WT) grown under 24 h-, 48 h-, 72 h- and 96 h-P deprivation was shown. (B) Relative transcriptional level of selected TAG biosynthesis genes (*DGTT1*, *DGTT2*, *DGTT3*) in the *pdat* mutant (gray) and the wild-type (black) was measured at 24 h under P deprivation. Data represents mean ± standard deviation (SD) from three biological repeats. An asterisk indicates statistical significance by Student's *t*-test (*p* value ≤ 0.05).

Fig 4. Phenotype and gene expression of the *VTC1* strain, Δ*vtc1* single mutant and Δ*vtc1*/*pdat* double mutant. (A) TAG content of the Δ*vtc1* strain CC5165 and the *VTC1* rescue strain CC5166 grown was measured at 48 h and 72 h under P replete. (B) TAG content of the *VTC1* strain, Δ*vtc1* strain and the Δ*vtc1*/*pdat* strain grown was measured at 48 h and 72 h under P deprivation. (C) Relative transcriptional level of selected genes (*DGTT1*, *DGTT2*, *DGTT3*, *PDAT* and *PGD1*) was measured at 24 h under P deprivation. Transcripts of *PDAT* in the Δ*vtc1*/*pdat* double mutant were nondetectable (N/D). Data represents mean ± standard deviation (SD) from three biological repeats. An asterisk indicates statistical significance by Student's *t*-test (*p* value ≤ 0.05).