Genetic compensation of triacylglycerol biosynthesis in the green microalga

Chlamydomonas reinhardtii

Yi-Ying Lee[a], Rudolph Park[b], Stephen M. Miller[b], and Yantao Li[a, c, 1]

[a] Institute of Marine and Environmental Technology, University of Maryland Center for

Environmental Science, Baltimore, MD 21202, USA

[b] Department of Biological Sciences, University of Maryland, Baltimore County, Baltimore, MD

21250, USA

[c] Department of Marine Biotechnology, University of Maryland Baltimore County, Baltimore,

MD 21202, USA

[1] Correspondence: Yantao Li

701 E Pratt Street, Baltimore, MD 21202

Phone: (410) 234-8833

Email: **yantao@umces.edu**

Key words: genetic compensation; triacylglycerol biosynthesis; *Chlamydomonas reinhardtii*;

CRISPR; gene knockout.

Abstract

Genetic compensation has been proposed to explain phenotypic differences between gene knockouts and knockdowns in several metazoan and plant model systems. With the rapid development of reverse genetic tools such as CRISPR/Cas9 and RNAi in microalgae, it is increasingly important to assess whether genetic compensation affects the phenotype of engineered algal mutants. While exploring triacylglycerol (TAG) biosynthesis pathways in the model alga *Chlamydomonas reinhardtii*, it was discovered that knockout of certain genes catalyzing rate-limiting steps of TAG biosynthesis, type-2 diacylglycerol acyltransferase genes (*DGTTs*), triggered genetic compensation under abiotic stress conditions. Genetic compensation of a *DGTT1* null mutation by a related *PDAT* gene was observed regardless of the strain background or mutagenesis approach, e.g., CRISPR/Cas 9 or insertional mutagenesis. However, no compensation was found in the *PDAT* knockout mutant. The effect of *PDAT* knockout was evaluated in a Δ*vtc1* mutant, in which *PDAT* was up-regulated under stress, resulting in a 90% increase in TAG content. Knockout of *PDAT* in the Δ*vtc1* background induced a 12.8-fold upregulation of *DGTT1* and a 272.3% increase in TAG content in Δ*vtc1*/*pdat1* cells, while remaining viable. These data suggest that genetic compensation contributes to the genetic robustness of microalgal TAG biosynthetic pathways, maintaining lipid and redox homeostasis in the knockout mutants under abiotic stress. This work demonstrates examples of genetic compensation in microalgae, implies the physiological relevance of genetic compensation in TAG biosynthesis under stress, and provides guidance for future genetic engineering and mutant characterization efforts.

Significance statement

Genetic compensation systems have been discovered in several metazoan and plant systems to

buffer deleterious mutations. With the rapid development of reverse genetic tools in microalgae,

it is increasingly important to assess whether genetic compensation affects the phenotype of

engineered algal mutants. We report that knockout of certain genes catalyzing rate-limiting steps

of triacylglyceroal (TAG) biosynthesis triggers up-regulation of homologous genes to buffer the

effect of a null mutation in *Chlamydomonas reinhardtii* under abiotic stress conditions. This

work provides evidence to show physiological relevance of genetic compensation in TAG

biosynthesis in the green lineage, and paves the way for future gene function analysis in

microalgae.

Introduction

Organisms have evolved various genetic buffering systems to maintain fitness in response to genetic perturbations, including functionally redundant genes (1), protein feedback loops (2), and acquisition of adaptive mutations (3). Such buffering or compensation systems ensure similar growth or developmental outcomes despite some genetic changes (4). Genetic compensation is defined as the phenomenon whereby the effect of a deleterious mutation is buffered by the genome (5). Recently, genetic compensation through transciptional adaptation, by which a homologous gene or genes are upregulated in response to an early nonsense mutation in a related gene, has been proposed as a mechanism for maintaining genetic robustness in a number of metazoans, including zebrafish, mice, and nematodes, and to explain profound differences between gene knockouts and knockdowns at certain loci (6-9). Likewise, comparison between gene knockouts and knockdowns in some model plant systems has revealed puzzling discrepencies that suggest transcriptional adaptation is acting to buffer the genome in these species as well (10-13).

Increasingly, gene knockdown and knockout tools have been deployed in the green microalga *Chlamydomonas reinhardtii* (Chlamydomonas; (14-19)), but little attention has been paid as to whether these tools might for some genes lead to very different outcomes, and more broadly, whether genetic compensation occurs in green algae (4, 5). Here we set out to address these questions as part of an ongoing study of lipid biosynthesis in Chlamydomonas, using triacylglycerol (TAG) biosynthesis pathways as examples.

Microalgae store TAG as a carbon and energy-rich compound in cells, whose synthesis can be stimulated under various stress conditions (20, 21). Nutrient deprivation such as nitrogen (N) deprivation and phosphorus (P) deprivation are most commonly used stressors to induce

TAG production. Under N deprivation, algal cell growth is almost completely arrested while P

deprivation promotes TAG production with moderate cell growth (22). Biosynthesis of TAG

involves sequential acylation to the precursor glycerol-3-phosphate. Two pathways are involved

in the last step of TAG biosynthesis. One pathway is a *de novo* acyl-CoA-dependent route

mediated by diacylglycerol acyltransferase (DGAT) (23). Three major types of DGATs

including two structurally distinctive, membrane bound type-1 and type-2 DGATs and a soluble

cytosolic type-3 DGAT have been identified in plants, algae, and other eukaryotes (24-26). The

other TAG biosynthesis pathway is an acyl-CoA-independent route that requires membrane lipid

breakdown and is mediated by a plastidic membrane bound enzyme, phospholipid:diacylglycerol

acyltransferase (PDAT) (27, 28).

The genome of *C. reinhardtii* harbors one type-1 *DGAT* gene (*DGAT1*), five type-2

DGAT genes (*DGTT1-5*), and one *PDAT* gene (28-31). RNA-Seq analysis previously revealed

that the abundance of *PDAT*, *DGAT1* and *DGTT1* transcripts was up-regulated, and that of

DGTT2 and *DGTT3* was moderate but constant, and the abundance of transcripts of *DGTT4* and

DGTT5 was low to undetectable in response to N deprivation (30). Subsequent RT-qPCR

analyses corroborated these findings for *PDAT*, *DGAT1* and *DGTT1*, while *DGTT2-3* were found

to be induced by N deprivation (28, 32). Similar results were found in response to P deprivation,

where *PDAT*, *DGAT1*, and *DGTT1-4* were up-regulated during TAG accumulation (22).

Functional studies using *in vitro* enzyme assays and gene knockdowns have largely yielded

results consistent with expectations based on these expression data. *DGTT1-3* and *PDAT* were

shown to catalyze TAG biosynthesis *in vitro* (28, 32). Knockdown of *DGTT1*, *DGTT2* or *DGTT3*

each resulted in a moderately reduced TAG (TAG-less) phenotype under persistent stress, while

knockdown of *PDAT* resulting in a moderate TAG-less phenotype during both favorable growth

and the early phase of stress induction (28, 32), suggesting *DGTT*s and *PDAT* are functionally

related genes involved in TAG biosynthesis in Chlamydomonas. However, in some cases,

analysis of gene knockdowns can be limited by off-target effects and low efficiency of gene

silencing (33, 34), so null mutants are needed to better understand gene function. Among

Chlamydomonas TAG biosynthesis genes, so far only *pdat* mutants have been characterized,

which had reduced TAG content under stress (30). However, whether the *PDAT* insertion

mutations analyzed were buffered by expression of related genes such as *DGTT*s is not known,

and up to now no *DGTT* mutants have been analyzed. So the precise contributions of these genes

to TAG accumulation remains obscure.

110   In this work we analyzed TAG accumulation in strains bearing null mutations in several

genes including *DGTT1-3* and *PDAT*, and found phenotypes that contradicted known

knockdown phenotypes in these genes. Our data showed genetic compensation through up-

regulation of related TAG biosynthetic genes for *dgtt1* and *dgtt2* mutants resulted in a higher

TAG content under stress, which is contrary to the TAG-less phenotype in the respective

knockdown lines. Interestingly, no genetic compensation was found in the *pdat* mutant. However,

when evaluated in the Δ*vtc1* background (defective for Vacuolar Transporter Chaperone 1, a

component of the polyphosphate polymerase complex), where *PDAT* expression and function is

likely critical for storing excess energy as TAG under stress, the *PDAT* knockout resulted in

compensation by up-regulation of *DGTT1* and a higher TAG content. Our data demonstrate

genetic compensation exists in microalgae and imply the physiological importance of genetic

compensation in TAG biosynthesis under stress.

Results

Genetic compensation is present in microalgae

Type-2 DGATs are believed to be major contributors for TAG biosynthesis in algae, but previous knockdown of Chlamydomonas type-2 *DGAT* genes *DGTT1*, *DGTT2* and *DGTT3* caused only relatively small (~10 - 20%) suppression of TAG accumulation (32). We set out to determine whether null mutations in these genes might have larger effects, so we obtained *dgtt1*, *dgtt2* and *dgtt3* CLiP library knockout mutants generated by the Jonikas group (14, 35) and analyzed their TAG accumulation. Each mutant strain carries a CIB1 cassette insertion that disrupts the respective *DGTT* gene. We further examined the genotype of these mutants and found the *dgtt1* mutant (LMJ.RY0402.223444) insertion is in the middle of intron 1, the *dgtt2* mutant (LMJ.RY0402.213587) insertion is at the junction between exon 4 and intron 4, and the *dgtt3* mutant (LMJ.RY0402.223565) insertion is at the beginning of exon 3 (Fig. 1A; Fig. S1). No detectable transcripts of *DGTT1*, *DGTT2* and *DGTT3* were found in the respective mutants (Fig. 1C), suggesting all the mutants are null.

TAG biosynthesis was comparatively studied in the *dgtt1*, *dgtt2*, and *dgtt3* mutants and their parent strain (CC5325; wild-type) under P deprivation, which was reported to induce TAG accumulation in *C. reinhardtii* while maintaining moderate cell growth (15). Surprisingly, the *dgtt1* and the *dgtt2* knockout mutants had significantly higher TAG content than the wild-type strain (t-test, p ≤ 0.05, Fig. 1B) either on a per cell basis or per dry weight (DW) basis. The TAG content of *dgtt1* and *dgtt2* mutant cells increased by about 90% and 70% per cell, respectively, and by 45% and 76% per unit dry weight, compared with wild-type cells at 72 h under P deprivation (Fig. 1B). These results are in stark contrast to those for the knockdown strains, which had reduced TAG content (32), and they suggest a compensation mechanism exists in Chlamydomonas that affects other TAG genes/enzymes in the *dgtt1* and *dgtt2* mutant. By

contrast, the *dgtt3* knockout mutant contained ~40% less TAG per cell and per dry weight

compared with the wild-type (Fig. 1B), and notably it had about twice the decrease in TAG as

that caused by the *dgtt3* knockdown (32). In Chlamydomonas, *DGTTs* and *PDAT* are homologs.

Sequence analysis revealed identities between *PDAT* and *DGTT1*, *DGTT2*, and *DGTT3* were

50.9%, 49.8%, and 49.2%, respectively (Table S1), all above 40% sequence identity needed to

induce genetic compensation by homologs in previous studies (8).

153   To explore possible genetic compensation in the knockout mutants, the expression of

TAG biosynthetic genes was examined, including that of *DGTT1*, *DGTT2*, *DGTT3*, *PDAT*, and

PGD1 (encodes an MGDG-specific lipase and presumably functions with PDAT in membrane

recycling) (28, 32, 36). In the *dgtt1* mutant, the expression of *DGTT2*, *DGTT3* and *PGD1*

remained unchanged, but the expression of *PDAT* was upregulated by 9-fold compared with the

wild-type at 24 h under P deprivation (Fig. 1C). In the *dgtt2* mutant, the expression of all tested

genes was upregulated; compared with the wild-type, the expression of *DGTT1*, *DGTT3*, *PDAT*

and *PGD1* in the *dgtt2* mutant was increased by 19-fold, 2.6-fold, 9-fold, and 7.6-fold,

respectively, at 24 h under P deprivation (Fig. 1C), correlating with higher TAG accumulation in

the *dgtt2* mutant. In the *dgtt3* knockout strain, the expression level of *PDAT* was significantly

higher than for the wild type (t-test, $p \leq 0.05$; Fig. 1C), but that increase did not lead to a

functional compensation for TAG accumulation (Fig. 1B).

165   Our findings suggest that a mutation leading to a premature stop codon in two different

type-2 *DGAT* genes likely activates a genetic compensatory network by upregulation of

homologous genes in Chlamydomonas. To determine whether this genetic compensation was

specific to insertional mutants in the CLiP library genetic background, or whether it might be a

more general phenomenon of null mutants with premature stop codons, we used CRISPR/Cas9

to generate a *dgtt1* mutant in a different strain background (*C. reinhardtii* CC3403). In the

resulting mutant (*CC-dgtt1*), the *DGTT1* gene carries a 1229-bp insertion corresponding to

pHS_SaCas9 sequence (Cas9 expression vector) near the beginning of exon 3 of the coding

region (Fig. 2A and Fig. S2A) that results in premature stop codons (Fig. S2B) and disrupts

expression of the *DGTT1* gene. The TAG content of *CC-dgtt1* was comparable to that of the

wild-type on a per cell basis, and 66% higher than the wild-type per unit dry weight under P

deprivation (Fig. 2B). Moreover, this phenotype was maintained under N deprivation (Fig. S3).

Similar to the situation for the *dgtt1* mutant from the CLiP library, RT-qPCR analyses revealed

that *PDAT* expression was upregulated by 6.3-fold in the *CC-dgtt1* mutant, while other TAG

biosynthesis genes (*DGTT2*, *DGTT3* and *PGD1*) remained the same as in the wild-type (Fig. 2C).

Taken together, genetic compensation of *DGTT* seems to occur in Chlamydomonas mutants

regardless of the strain background or early-frameshift mutant allele.

Compensation of *DGTT* knockouts is not reciprocal in the *pdat* mutant

Since *C. reinhardtii* cells are capable of compensating defects of *DGTT1* and *DGTT2*

mutations by upregulating related genes such as *PDAT*, we asked whether the mechanism is

reciprocal in the *pdat* mutant. To this end, we assessed TAG accumulation and the expression of

the *DGTT1-3* genes in a *pdat* knockout mutant (CC4502) previously described in (30). Unlike

the *dgtt1 and dgtt2* mutants, the *pdat* mutant had reduced TAG levels (Fig. 3), similar to the *pdat*

knockdown lines (28). The relative TAG content of the *pdat* knockout mutant was ~30% lower

than that of the wild-type strain at 48 h and 72 h under P deprivation (Fig. 3A), similar to the

decrease in TAG content previously reported for this strain in response to N deprivation (30). In

P-deprived *pdat* cells, the expression of *DGTT1-3* were the same as or slightly lower than in the

wild-type (Fig. 3B). These data suggest no genetic compensation of TAG biosynthesis occurs in

the *pdat* mutant.

Compensation by upregulation of *DGTT1* in a *Δvtc1/pdat1* double mutant

Our previous work suggests *PDAT* primarily regulates TAG biosynthesis under favorable,

nutrient-replete conditions (28) but not under persistent nutrient-deficient stress conditions (32).

A possible explanation for lack of genetic compensation in the *pdat* mutant is thus the relatively

poor function of *PDAT* under stress conditions, in which *DGTTs* are already highly expressed

and responsible for TAG biosynthesis. Previous work showed P metabolism, particularly P

uptake, was impaired in a $\Delta vtc1$ mutant, in which the vacuolar transport chaperon (VTC)

complex was disrupted (37-40). As such, in this $\Delta vtc1$ mutant we reasoned that *PDAT* expression

and function may be especially critical to recycle phospholipids and balance P metabolism under

stress. To test this idea, we analyzed TAG biosynthesis in the $\Delta vtc1$ mutant and in a *VTC1*-

rescued strain (*VTC1*) under P replete and P-depleted conditions. In $\Delta vtc1$ cells TAG content was

90% higher (*t*-test, $p \leq 0.05$) than the control under P-deprived conditions but not under P-replete

conditions (Fig. 4A and B), suggesting loss of *VTC1* had induced TAG biosynthesis under stress.

When we analyzed the TAG biosynthetic genes in P-depleted $\Delta vtc1$ cells, only the *PDAT* gene

showed an increase in expression (2.8-fold; Fig. 4C) while none of the *DGTT* or *PGD* genes

were upregulated, suggesting a role of *PDAT* in controlling TAG biosynthesis under stress in the

$\Delta vtc1$ mutant.

We predicted that for the $\Delta vtc1$ mutant, in which *PDAT* function is critical, if *PDAT*

expression is interrupted, genetic compensation by other TAG biosynthetic genes may be

necessary for the resulting mutant to survive under stress. To test this idea, we generated a

Δ*vtc1*/*pdat* double mutant by crossing the Δ*vtc1* (CC5165) and the *pdat* (CC4502) single mutant

strains. Interestingly, the TAG level in the resulting strain increased compared with the control

and single mutant (Fig. 4B). After 48 h of P deprivation, the TAG content of Δ*vtc1*/*pdat* cells

was about 272.3% higher than the *VTC1* control and 95% higher than the Δ*vtc1* mutant (Fig. 4B).

We found that under stress, *DGTT1* transcripts were upregulated in Δ*vtc1*/*pdat* cells compared

with the control (by12.4-fold), but no other TAG biosynthetic genes were transcriptionally

upregulated (Fig. 4C). Moreover, Δ*vtc1*/*pdat* cells remained viable with moderate growth under

P deprviation. Δ*vtc1*/*pdat* cell density slightly increased from 0.69 x 10^6 cells/ml to 1.01 x 10^6

cells/ml between day 0 and day 4 (Fig. S4A), while the per cell dry weight of the Δ*vtc1*/*pdat*

mutant about doubled compared with the control (*VTC1*) (Fig. S4B). By contrast, the growth rate

and per cell dry weight of Δ*vtc1* cells were similar to that of the control (Fig. S4). Therefore, our

results suggest that upregulation of *PDAT* in the Δ*vtc1* mutant results in an increase in TAG

accumulation, while the effect of *PDAT* disruption in the Δ*vtc1* background is buffered by

upregulation of *DGTT1* in the *Δvtc1/pdat1* double mutant.

Discussion

Genetic compensation, and more specifically, transcriptional adaptation through

upregulation of related genes, is proposed as an underlying mechanism to explain discrepencies

between knockdown and knockout phenotypes in several model systems including zebrafish and

mouse (6-8), *Caenorhabditis elegans* (9), Arabidopsis (10-12), and tomato (13). One mechanism

for this phenomenon involves the nonsense-mediated mRNA decay pathway that is activated by

early nonsense mutations (7, 8). However, it is not known whether this phenomenon exists

across eukaryotes, including microalgae (4, 5). Here, we demonstrate that compensation by

related genes buffers the effect of deleterious mutations in Chlamydomonas lipid biosynthesis

genes, in a process that resembles transcriptional adaptation. Knockouts of *DGTT1* triggered

upregulation of the functionally related and homologous *PDAT* gene regardless of the strain

background, mutagenesis approach, or stress conditions (P or N deprivation), while knockout of

DGTT2 triggered upregulation of *PDAT*, *DGTT1* and *PGD1* (Fig. 1 and Fig. 2). In the *dgtt1*, *CC-*

dgtt1 and *dgtt2* mutant, the TAG-less phenotype found in respective knockdown lines (32) was

not observed. Instead, these mutants produced more TAG than the control, likely due to

compensation by upregulation of homologous genes. In the *dgtt3* mutant, however, TAG content

was reduced despite compensation by upregulation of *PDAT* (Fig. 1). It is possible that DGTT3

and PDAT proteins function together, such that upregulation of PDAT without functional

DGTT3 protein could not restore TAG biosynthesis in *dgtt3* cells. Alternatively, knockout of

DGTT3 may trigger unknown mechanisms that couteract the effect of *PDAT* upregulation.

Recent advances in reverse genetic tools including CRISPR, TALEN, and ZFN genome

editing methods have greatly accelerated progress in generating knockout mutants in traditional

and new algal model systems, including Chlamydomonas (17, 19, 41), *Nannochloropsis* (42, 43),

Volvox carteri (44), and *Ectocarpus* (45). Moreover, there is a genome-wide insertional mutant

pool, the CLiP library (14), for Chlamydomonas. Such tools and mutant libraries dramatically

facilitate analysis of gene function. However, as we demonstrated here, genetic compensation by

related genes might, in some cases, offset the effects of a null mutation, confounding genetic

analysis. Likewise, previous efforts to screen and characterize TAG-less Chlamydomonas

mutants from the CLiP library failed to identify any *DGTT* knockout mutants (XB Li, personal

communication), possibly due to genetic compensation, while knockdown lines of *DGTTs*

clearly showed a reduced-TAG phenotype (32). Therefore, when studying gene function using

null mutants, preferably multiple mutant strains targeting the same gene should be obtained by a

targeted genome editing method (e.g. CRISPR) or from the CLiP libarary (because some might

activate genetic compensation while others might not), and these mutants should be carefully

evaluated for their genotypic and phenotypic traits. If genetic compensation is observed in the

mutants, alternative methods such as knockdown should be used to confirm the respective gene

function. Ideally, deletion mutants should be analyzed as they will not trigger transcriptional

adaptation through the nonsense-mediated mRNA decay pathway (7, 8), though to date a

relatively small number of deletion mutants have been generated in Chlamydomonas due to

current technical limitations.

In Chlamydomonas, biosynthesis of TAG has been implicated in the protection of the

photosynthetic electron transport chain from over-reduction during stress conditions, and TAG

accumulation is suggested to be essential for cells to survive under stress (36, 46). Our initial

experiments raised two interesting questions. First, does genetic compensation of TAG

biosynthesis, as appears to occur in the *dgtt1* and *dgtt2* mutants, play an essential survival role

under persistent stress conditions, and second, might genetic compensation occur for other

mutated TAG biosynthetic genes? To address these questions, we first examined the *pdat* mutant.

Unlike the *dgtt1* and *dgtt2* mutants, it has a similar TAG-less phenotype and TAG biosynthetic

gene expression profiles when stressed as the knockdown lines (Fig. 3 and (28, 30)), suggesting

no genetic compensation in the *pdat* mutant. This outcome may be due to the poor function of

PDAT in stress conditions, under which *DGTTs* are primarily responsible for TAG biosynthesis

(28, 32). Consistent with this notion is a recent *PDAT* analysis in the green lineage showing that

PDAT plays a less important role in algae than in plants (47). Disruption of *AtPDAT1* in *A.*

thaliana led to an over 50% decrease in oil accumulation in growing tissues and significant

gametophytic and growth defects (28), whereas knockout of Chlamydomonas *PDAT* reduced

TAG content by only 25% - 30%, and *pdat* cells had similar growth rate as the wild type (Fig 3A)

(30).

To determine whether there might be compensation for the *PDAT* knockout when PDAT

function is most essential, we evaluated the effect of the *PDAT* knockout in a Δ*vtc1* mutant, in

which *PDAT* expression and function are likely critical for Δ*vtc1* cells to store excess energy as

TAG under stress (Fig. 4). We expected that in the absence of compensation, the knockout of

PDAT in the Δ*vtc1* background should result in a TAG-less phenotype. Intriguingly, the

Δ*vtc1*/*pdat* mutant exhibited a 272.3% higher TAG content than the *VTC1* control, likely because

of a large (12.8-fold) upregulation of *DGTT1* (Fig. 4). The Δ*vtc1*/*pdat* mutant remained viable

under stress, as its cell density and dry weight increased in response to P deprivation (Fig. S4).

So while no genetic compensation was observed in the *pdat* mutant, in the Δ*vtc1*/*pdat* double

mutant, *DGTT1* was highly overexpressed, and more TAG was produced under P deprivation.

Under stress conditions, these cells continue to capture light energy through photosynthesis, and

if electron acceptors become overreduced due to the slow growth, cytotoxic reactive oxygen

species are produced that lead to cell death (48). TAG biosynthesis is likely essential for

Δ*vtc1*/*pdat* cells to survive during stress because TAG is the most reduced electron sink in algae

(20). In line with our findings, it was previously reported that a Chlamydomonas mutant

defective for the MGDG-specific lipase PGD1 accumulated less TAG than the wild type while

its cells lost viability under stress, suggesting TAG plays an essential role as an electron and

energy sink under stress conditions, thereby preventing generation of reactive oxygen species (36,

46). Together, these data suggest that under stress conditions, loss of TAG biosynthetic gene

function can be buffered by upregulation of a related gene or genes through transcriptional

adaptation in algae.

In summary, we have demonstrated that genetic compensation contributes to genetic

robustness in microalgal TAG biosynthesis, maintaining TAG biosynthesis and redox

homeostasis in knockout mutants under abiotic stress. This work demonstrates, to the best of our

knowledge, the first reported example of genetic compensation in microalgae, implies the

physiological relevance of genetic compensation in TAG biosynthesis under stress, and provides

guidance for future genetic engineering and mutant characterization efforts. The exact

mechanisms inducing genetic compensation in the Chlamydomonas mutants are not understood

and warrant further investigation. Since transcriptional adaptation in other systems is triggered

by mutant mRNA decay and involves Upf3a and COMPASS components (7, 8), testing the

involvement of these components in algal genetic compensation is a logical next step.

Materials and Methods

Strains and growth conditions

Chlamydomonas reinhardtii LMJ.RY0402.223444, LMJ.RY0402.213587,

LMJ.RY0402.223565 and their parent strain CC5325 (*cw15*; mt⁻) (35), CC4502 (*pdat1-1*; mt⁺)

and its parent strain CC4425 [D66] (*cw15*; mt⁺) (30), CC3403 (*arg7*; *cw15*; mt⁻), Δ*vtc1* mutant

strain CC5165 (Δ*vtc1*; mt⁻) and the *VTC1* rescue strain CC5166 (*ars76::VTC1*) (38) were

obtained from the Chlamydomonas Resource Center (http://www.chlamycollection.org).

CC3403 was used as the wild-type in CRISPR/Cas9 mutagenesis (17). The Δ*vtc1*/*pdat* double

mutant was derived from a cross between CC5165 and CC4502 (methods described below). All

strains were maintained in Tris-acetate-phosphate (TAP) medium containing paramomycin as

necessary. P deprivation was imposed by transferring *C. reinhardtii* cells to P-free TAP, in

which potassium phosphate was replaced by 1.5 mM KCl. Ammonium chloride was omitted to

generate N-free TAP. All batch cultures were grown under continuous light illumination at 60

μmol photons m^{-2} s^{-1}, shaking at 150 rpm, room temperature.

Gene expression measurement by RT-qPCR

Total RNA was extracted and purified using Trizol reagent (Invitrogen) and RNeasy mini

kit (Qiagen). cDNA was synthesized using Protoscript II First Strand cDNA synthesis kit (NEB).

The expression of target genes was determined by RT-qPCR in Applied Biosystems 7500 using

the $2^{-\Delta\Delta CT}$ method normalized by the expression of 18s rRNA. Details are described in the

Supplemental Materials and Methods. The RT-qPCR analysis for each targeted gene was

performed twice, with three biological replicates and three technical replicates for each sample.

The Primer sequences used in RT-qPCR are listed in Table S2.

Generation and screening of *Chlamydomonas reinhardtii* mutants

The CC-*dgtt1* mutant was generated by the CRISPR/Cas9 method developed by Greiner

et al (17). The parent strain of the *CC-dgtt1* mutant (CC3403), the SaCas9 expression vector

(pHS_SaCas9) and the sgRNA cloning vector (pCrU6.4-SaCloning-aphVIII) were obtained from

the Chlamydomonas Resource Center. In brief, a customized targeting vector that expressed

sgRNA specifically targeting *DGTT1* was constructed in pCrU6.4-SaCloning-aphVIII. The

SaCas9 expression vector and the customized targeting vector that co-transformed into CC3403

by electroporation (9, 41). The transformants were selected for paromomycin resistance and

screened by PCR using oligonucleotides CrDGTT1F and CrDGTT1R (sequences in Table S2)

for insertions at the *DGTT1* locus. The candidates yielding PCR products larger than the

expected size for the wild type *DGTT1* region were sequenced. The detailed procedures are

described in the Supplemental Materials and Methods.

The Δ*vtc1/pdat1* double mutant was generated by mating as described in Goodenough,

1976 (49) with some modifications (procedures described in the Supplemental Materials and

Methods). The crossed candidates were selected for paramomycin resistance and subsequently

screened by colony PCR for a co-occurrence with the deletion of the *VTC1* gene and the

presence of the *pdat1-1* mutation (30) in the genome. Primers CrVTC1-pF1 and CrVTC1-stopR1

(sequences in Table S2) were used for determination of the presence (yielding a 1148 bp PCR

product) or the absence (yielding no PCR products) of the *VTC1* gene. Primers CrPDAT1-1-F2

and RIM5-2 (sequences in Table S2) (30) were used to determine the presence (~ 1 kb PCR

product) or the absence (no PCR product) of the *AphIII* insertion in the *PDAT* gene.

TAG content measurement

Total lipid from *C. reinhardtii* cells was extracted using chloroform: methanol (2:1, v/v)

as previously described (28). The lipid extracts were dried under a gentle stream of nitrogen gas

and re-dissolved in chloroform. Total lipid was resolved by thin-layer chromatography (TLC) on

a silica gel 60 F_{254} plates (EMD Millipore) using hexane: t-butyl methyl ether: acetic acid

(80:20:2, v/v/v) solvent mixture as a mobile phase to develop TAG. For visualization, the

developed TLC plates were sprayed with 8% (w/v) H_3PO_4 solution containing 10% (w/v) copper

(II) sulfate pentahydrate, and then charred at 180°C for 3 min. The relative intensity of signals

corresponding to TAG can be measured using ImageJ software (NIH). The absolute

quantification of TAG can be calculated by the sum of fatty acid fractions composed of TAG

using gas chromatography-mass spectrometry (GC-MS) as previously described (23, 50). Details

are described in the Supplemental Materials and Methods.

Statistical analysis

Statistical analysis was performed by Student's *t*-test with two-tailed distribution. *p* value ≤ 0.05 was the criteria to determine statistical significance.

Acknowledgements

This work was supported in part by grants from the U.S. National Science Foundation (CBET-1511939) and DOE Office of Fossil Energy (FE-0031914) to YL, and by NSF 1332344 to SMM. We thank Drs. William Snell, Peeyush Ranjan, and Mayanka Awasthi at University of Maryland College park for their help with generating double knockout mutants, Dr. Arthur Grossman at Carnegie Institution for Science for providing the *Chlamydomonas vtc1* strains used in this study, Dr. Xiaobo Li at Westlake University for discussion of unpublished mutant screening data, and Drs. Fan Zhang and Russell Hill at the Institute of Marine and Environmental Technology for their help with polyP analysis. We thank Mr. Mohamed Mahmoud-Aly at Cairo University for his help with *Chlamydomonas* mutant characterization.

Competing financial interests

The authors declare no competing financial interests.

References

1. D. Tautz, Redundancies, development and the flow of information. *Bioessays* **14**, 263-266 (1992).
2. A. L. Barabási, Z. N. Oltvai, Network biology: understanding the cell's functional organization. *Nat Rev Genet* **5**, 101-113 (2004).
3. X. Teng *et al.*, Genome-wide Consequences of Deleting Any Single Gene. *Molecular Cell* **52**, 485-494 (2013).
4. M. A. El-Brolosy, D. Y. R. Stainier, Genetic compensation: A phenomenon in search of mechanisms. *PLoS genetics* **13**, e1006780-e1006780 (2017).

5.   Z. Kontarakis, D. Y. R. Stainier, Genetics in Light of Transcriptional Adaptation. *Trends*
in Genetics **36**, 926-935 (2020).
6.   A. Rossi *et al.*, Genetic compensation induced by deleterious mutations but not gene
knockdowns. *Nature* **524**, 230-233 (2015).
7.   M. A. El-Brolosy *et al.*, Genetic compensation triggered by mutant mRNA degradation.
Nature **568**, 193-197 (2019).
8.   Z. Ma *et al.*, PTC-bearing mRNA elicits a genetic compensation response via Upf3a and
COMPASS components. *Nature* **568**, 259-263 (2019).
9.   V. Serobyan *et al.*, Transcriptional adaptation in Caenorhabditis elegans. *eLife* **9**, e50014
(2020).
10.   Y. Gao *et al.*, Auxin binding protein 1 (ABP1) is not required for either auxin signaling
or Arabidopsis development. *Proceedings of the National Academy of Sciences of the*
United States of America **112**, 2275-2280 (2015).
11.   X. Chen *et al.*, Inhibition of cell expansion by rapid ABP1-mediated auxin effect on
microtubules. *Nature* **516**, 90-93 (2014).
12.   N. Braun *et al.*, Conditional repression of AUXIN BINDING PROTEIN1 reveals that it
coordinates cell division and cell expansion during postembryonic shoot development in
Arabidopsis and tobacco. *The Plant cell* **20**, 2746-2762 (2008).
13.   D. Rodriguez-Leal *et al.*, Evolution of buffering in a genetic circuit controlling plant stem
cell proliferation. *Nature Genetics* **51**, 786-792 (2019).
14.   X. Li *et al.*, A genome-wide algal mutant library and functional screen identifies genes
required for eukaryotic photosynthesis. *Nature Genetics* **51**, 627-635 (2019).
15.   A. Molnar *et al.*, Highly specific gene silencing by artificial microRNAs in the
unicellular alga *Chlamydomonas reinhardtii*. *Plant J.* **58**, 165-174 (2009).
16.   J. Rohr, N. Sarkar, S. Balenger, B. R. Jeong, H. Cerutti, Tandem inverted repeat system
for selection of effective transgenic RNAi strains in Chlamydomonas. *Plant J.* **40**, 611-
621 (2004).
17.   A. Greiner *et al.*, Targeting of Photoreceptor Genes in Chlamydomonas reinhardtii via
Zinc-Finger Nucleases and CRISPR/Cas9. *Plant Cell* **29**, 2498-2518 (2017).
18.   M. Schroda, RNA silencing in Chlamydomonas: mechanisms and tools. *Curr. Genet.* **49**,
69-84 (2006).
19.   W. Jiang, A. J. Brueggeman, K. M. Horken, T. M. Plucinak, D. P. Weeks, Successful
transient expression of Cas9/sgRNA genes in Chlamydomonas reinhardtii. *Eukaryot. Cell*
10.1128/ec.00213-14 (2014).
20.   Q. Hu *et al.*, Microalgal triacylglycerols as feedstocks for biofuel production:
perspectives and advances. *Plant J* **54**, 621-639 (2008).
21.   S. K. Lenka *et al.*, Current advances in molecular, biochemical, and computational
modeling analysis of microalgal triacylglycerol biosynthesis. *Biotech Adv* **34**, 1046-1063
(2016).
22.   M. Iwai, K. Ikeda, M. Shimojima, H. Ohta, Enhancement of extraplastidic oil synthesis in
Chlamydomonas reinhardtii using a type-2 diacylglycerol acyltransferase with a
phosphorus starvation-inducible promoter. *Plant Biotechnology Journal* **12**, 808-819
(2014).
23.   J. Ohlrogge, J. Browse, Lipid biosynthesis. *Plant Cell* **7**, 957-970 (1995).

24.   S. Cases *et al.*, Identification of a gene encoding an acyl CoA:diacylglycerol
acyltransferase, a key enzyme in triacylglycerol synthesis. *Proceedings of the National*
Academy of Sciences of the United States of America **95**, 13018-13023 (1998).
25.   K. D. Lardizabal *et al.*, DGAT2 is a new diacylglycerol acyltransferase gene family:
purification, cloning, and expression in insect cells of two polypeptides from Mortierella
ramanniana with diacylglycerol acyltransferase activity. *The Journal of biological*
chemistry **276**, 38862-38869 (2001).
26.   P. Oelkers, D. Cromley, M. Padamsee, J. T. Billheimer, S. L. Sturley, The DGA1 gene
determines a second triglyceride synthetic pathway in yeast. *The Journal of biological*
chemistry **277**, 8877-8881 (2002).
27.   A. Dahlqvist *et al.*, Phospholipid:diacylglycerol acyltransferase: an enzyme that catalyzes
the acyl-CoA-independent formation of triacylglycerol in yeast and plants. *Proceedings*
of the National Academy of Sciences of the United States of America **97**, 6487-6492
(2000).
28.   K. Yoon, D. Han, Y. Li, M. Sommerfeld, Q. Hu, Phospholipid:diacylglycerol
acyltransferase is a multifunctional enzyme involved in membrane lipid turnover and
degradation while synthesizing triacylglycerol in the unicellular green microalga
Chlamydomonas reinhardtii. Plant Cell **24**, 3708-3724 (2012).
29.   R. Miller *et al.*, Changes in transcript abundance in Chlamydomonas reinhardtii
following nitrogen deprivation predict diversion of metabolism. *Plant physiology* **154**,
1737-1752 (2010).
30.   N. R. Boyle *et al.*, Three acyltransferases and nitrogen-responsive regulator are
implicated in nitrogen starvation-induced triacylglycerol accumulation in
Chlamydomonas. J. Biol. Chem. **287**, 15811-15825 (2012).
31.   C. Bagnato *et al.*, Analysis of triglyceride synthesis unveils a green algal soluble
diacylglycerol acyltransferase and provides clues to potential enzymatic components of
the chloroplast pathway. *BMC genomics* **18**, 223 (2017).
32.   J. Liu, D. Han, K. Yoon, Q. Hu, Y. Li, Characterization of type 2 diacylglycerol
acyltransferases in Chlamydomonas reinhardtii reveals their distinct substrate
specificities and functions in triacylglycerol biosynthesis. *Plant J.* **86**, 3-19 (2016).
33.   Y. Fedorov *et al.*, Off-target effects by siRNA can induce toxic phenotype. *RNA* **12**,
1188-1196 (2006).
34.   Y. Ma, A. Creanga, L. Lum, P. A. Beachy, Prevalence of off-target effects in Drosophila
RNA interference screens. *Nature* **443**, 359-363 (2006).
35.   X. Li *et al.*, An Indexed, Mapped Mutant Library Enables Reverse Genetics Studies of
Biological Processes in *Chlamydomonas reinhardtii. The Plant Cell* **28**, 367-387 (2016).
36.   X. Li *et al.*, A galactoglycerolipid lipase is required for triacylglycerol accumulation and
survival following nitrogen deprivation in Chlamydomonas reinhardtii. *Plant Cell.* **24**
(2012).
37.   M. Plouviez *et al.*, Responses of Chlamydomonas reinhardtii during the transition from
P-deficient to P-sufficient growth (the P-overplus response): The roles of the vacuolar
transport chaperones and polyphosphate synthesis. *J. Phycol.* **57**, 988-1003 (2021).
38.   M. Aksoy, W. Pootakham, A. R. Grossman, Critical Function of a *Chlamydomonas*
reinhardtii Putative Polyphosphate Polymerase Subunit during Nutrient Deprivation. *The*
Plant Cell **26**, 4214-4229 (2014).

39.   E. Sanz-Luque, S. Saroussi, W. Huang, N. Akkawi, A. R. Grossman, Metabolic control
of acclimation to nutrient deprivation dependent on polyphosphate synthesis. *Sci Adv* **6**
(2020).
40.   S. Schmollinger *et al.*, Single-cell visualization and quantification of trace metals in
Chlamydomonas lysosome-related organelles. *Proc Natl Acad Sci U S A* **118** (2021).
41.   R. V. Park, H. Asbury, S. M. Miller, Modification of a Chlamydomonas reinhardtii
CRISPR/Cas9 transformation protocol for use with widely available electroporation
equipment. *MethodsX* **7**, 100855 (2020).
42.   Q. Wang *et al.*, Genome editing of model oleaginous microalgae Nannochloropsis spp.
by CRISPR/Cas9. *Plant J* **88** (2016).
43.   E. Poliner, T. Takeuchi, Z. Y. Du, C. Benning, E. M. Farré, Nontransgenic Marker-Free
Gene Disruption by an Episomal CRISPR System in the Oleaginous Microalga,
Nannochloropsis oceanica CCMP1779. *ACS Synth Biol* **7**, 962-968 (2018).
44.   J. A. Ortega-Escalante, R. Jasper, S. M. Miller, CRISPR/Cas9 mutagenesis in Volvox
carteri. *The Plant Journal* **97**, 661-672 (2019).
45.   Y. Badis *et al.*, Targeted CRISPR-Cas9-based gene knockouts in the model brown alga
Ectocarpus. *New Phytol* 10.1111/nph.17525 (2021).
46.   Z. Y. Du *et al.*, Galactoglycerolipid Lipase PGD1 Is Involved in Thylakoid Membrane
Remodeling in Response to Adverse Environmental Conditions in Chlamydomonas.
Plant Cell **30**, 447-465 (2018).
47.   L. J. Falarz *et al.*, Characterization of the diversification of phospholipid:diacylglycerol
acyltransferases in the green lineage. *The Plant journal : for cell and molecular biology*
103, 2025-2038 (2020).
48.   Y. Li, M. Sommerfeld, F. Chen, Q. Hu, Consumption of oxygen by astaxanthin
biosynthesis: A protective mechanism against oxidative stress in *Haematococcus*
pluvialis (Chlorophyceae). *J Plant Physiol* 10.1016/j.jplph.2007.12.007 (2008).
49.   U. W. Goodenough, C. Hwang, H. Martin, Isolation and genetic analysis of mutant
strains of Chlamydomonas reinhardi defective in gametic differentiation. *Genetics* **82**,
169-186 (1976).
50.   Y. Li *et al.*, *Chlamydomonas* starchless mutant defective in ADP-glucose
pyrophosphorylase hyper-accumulates triacylglycerol. *Metab. Eng.* **12**, 387-391 (2010).

Fig 1. Genetic structure, phenotype and gene expression of the *dgtt1*, *dgtt2*, *dgtt3* knockout mutants obtained from the CLiP library. (A) Locations of the insertion sites of the CIB1 cassette (brown triangles) and of the primer pairs (black arrows) used to detect gene expression in the *DGTT1*, *DGTT2* and *DGTT3* genes are presented. (B) TAG content of the *dgtt1* (blue), *dgtt2* (red), *dgtt3* (green) mutants and the parent strain CC5325 (black, WT) grown was measured at 72 h under P deprivation. (C) Relative transcriptional level of selected TAG biosynthesis genes (*DGTT1*, *DGTT2*, *DGTT3*, *PDAT* and *PGD1*) in the *dgtt1* (blue), *dgtt2* (red), *dgtt3* (green) mutants and the parent strain CC5325 (black) was measured at 24 h under P deprivation. Transcripts of *DGTT1* in the *dgtt1* mutant and *DGTT2* in the *dgtt2* mutant were nondetectable (N/D). Data represents mean ± standard deviation (SD) from three biological repeats. An asterisk indicates statistical significance by Student's *t*-test (p value ≤ 0.05) when compared to WT.

Fig 2. Genetic structure, phenotype and gene expression of the CRISPR-derived *dgtt1* knockout mutant. (A) The location of an *AphIII* cassette insertion (green triangle) generated by CRISPR in the *DGTT1* gene are presented. (B) TAG content of the *CC-dgtt1* mutant (blue) and the parent strain CC3403 (black, WT) grown was measured at 72 h under P deprivation. (C) Relative transcriptional level of selected TAG biosynthesis genes (*DGTT2*, *DGTT3*, *PDAT* and *PGD1*) in the *dgtt1* mutant (blue) and the parent strain CC3403 (black) was measured at 48 h under P deprivation. Data represents mean ± standard deviation (SD) from three biological repeats. An asterisk indicates statistical significance by Student's *t*-test (p value ≤ 0.05).

Fig 3. Phenotype and gene expression of the *pdat* mutant (CC4502). (A) Relative concentrations of TAG from the *pdat* knockout strain CC4502 (gray) and the wild-type CC4425 (black, WT) grown under 24 h-, 48 h-, 72 h- and 96 h-P deprivation was shown. (B) Relative transcriptional level of selected TAG biosynthesis genes (*DGTT1*, *DGTT2*, *DGTT3*) in the *pdat* mutant (gray) and the wild-type (black) was measured at 24 h under P deprivation. Data represents mean ± standard deviation (SD) from three biological repeats. An asterisk indicates statistical significance by Student's *t*-test (p value ≤ 0.05).

Fig 4. Phenotype and gene expression of the *VTC1* strain, Δ*vtc1* single mutant and Δ*vtc1*/*pdat* double mutant. (A) TAG content of the Δ*vtc1* strain CC5165 and the *VTC1* rescue strain CC5166 grown was measured at 48 h and 72 h under P replete. (B) TAG content of the *VTC1* strain, Δ*vtc1* strain and the Δ*vtc1*/*pdat* strain grown was measured at 48 h and 72 h under P deprivation. (C) Relative transcriptional level of selected genes (*DGTT1*, *DGTT2*, *DGTT3*, *PDAT* and *PGD1*) was measured at 24 h under P deprivation. Transcripts of *PDAT* in the Δ*vtc1*/*pdat* double mutant were nondetectable (N/D). Data represents mean ± standard deviation (SD) from three biological repeats. An asterisk indicates statistical significance by Student's *t*-test (*p* value ≤ 0.05).