Maximizing GWAS discovery and genomic prediction accuracy in Biobank data

Etienne J. Orliac[1], Daniel Trejo Banos[2], Sven E. Ojavee[3], Kristi Läll[4], Reedik Mägi[4], Peter M. Visscher[5,§]
Matthew R. Robinson[6,§,*]

1 Scientific Computing and Research Support Unit, University of Lausanne, Lausanne, Switzerland.
2 Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland.
3 Department of Computational Biology, University of Lausanne, Lausanne, Switzerland.
4 Estonian Genome Centre, Institute of Genomics, University of Tartu, Tartu, Estonia
5 Institute for Molecular Bioscience, University of Queensland, Brisbane, QLD, Australia.
6 Institute of Science and Technology Austria, Klosterneuburg, Austria.

§indicates joint last authorship
*corresponding author: matthew.robinson@ist.ac.at

Across 21 heritable traits in the UK and Estonian Biobank data, a Bayesian grouped mixture of regressions model (GMRM) obtains the highest genomic prediction accuracy reported to date, 15% (SD 10%) greater than a baseline model without MAF-LD-annotation groups, and 106% (SD 50%) greater than mixed-linear model association (MLMA) estimate polygenic scores. Prediction accuracy was up to 13% (mean 4%, SD 3%) higher than theoretical expectations, at 76% of the h^2_{SNP} for height (R^2 of 47%) and over 50% of the h^2_{SNP} for 12 traits. Using these predictors in MLMA, increased the independent GWAS loci detected from 16,899 using standard approaches to 18,837 using GMRM, an 11.5% increase. Modelling genetic associations accounting for MAF and LD differences among SNP markers, and incorporating prior knowledge of genomic function, is important for large-scale individual-level biobank-scale analyses and is facilitated by our scalable highly parallel open source GMRM software.

As biobank data sets increase in size, it is important to understand the factors limiting the prediction of phenotype from genotype. By combining recent work showing improved prediction accuracy from random effects models incorporating genomic annotations [1–4], with work on underlying heritability models [5–7], into a single flexible, scalable whole-genome regression modelling approach, we obtain vastly improved genomic prediction at Biobank scale, higher SNP heritability estimation (h^2_{SNP}) and often greater association power, as compared to common practices for analyzing large-scale biobank data. Our grouped Bayesian mixture of regressions model (GMRM) estimates genetic marker effects jointly, but with independent marker inclusion probabilities and independent h^2_{SNP} parameters across linkage disequilibrium (LD), minor allele frequency (MAF), and functional annotation groups (see Online Methods). Across 21 traits in the UK Biobank, we show that our GMRM improves h^2_{SNP} estimation by up to 35.6% (average of 9.09%, SD, 7.9%, Figure 1a) as compared to a REML model estimating a single h^2_{SNP} parameter. This improvement stems from the fact that genomic annotations differ greatly in the proportion of variance attributable to them, adjusted by that expected given the number of markers included within the model which map to each region (Figure S1). Our GMRM approach models the SNP-phenotype associations, capturing effect size differences across MAF-LD and genomic annotation groups and we obtain higher prediction accuracy than previously published estimates for all traits, in both a UK Biobank hold-out sample and within the Estonian Biobank data (Figure 1a, 1b, 1c). We improve prediction accuracy by an average of 14.6% (SD 10.2%) over the same model without MAF-LD-annotation groups, and by an average of 106% (SD 50%) over a genetic predictor taken from bolt-LMM summary statistics (Figure S2, S3). The h^2_{SNP} estimate sets the upper bound for prediction accuracy in an independent sample, and we achieve over 75% of the h^2_{SNP} for height, and over 50% for 12 of the traits in the UK Biobank hold-out sample (Figure 1b). The accuracy obtained is higher than that expected from theory by up to 12.5% (mean 4.1%, SD 3.4%, given ridge-regression assumptions and h^2_{SNP} estimates, Figure 1c), with prediction accuracy gains of GMRM increasing as power increases (Figure S4).

We meta-analysed the posterior mean effect sizes obtained from the UK Biobank, with those obtained from a GMRM analysis of 105,000 Estonian Genome Centre participants, and then predicted into an Estonian hold-out sample of 20,000 individuals, improving prediction of body mass index (BMI, prediction accuracy of 16.1%) and cardiovascular disease (CAD, prediction accuracy of 7%) over the accuracy obtained in the

UK Biobank hold-out sample (Figure 1d). Previous results have highlighted the lack of transfer of genomic predictors across populations and here we achieved reduced prediction accuracy for high blood pressure (BP) and type-2 diabetes (T2D) diagnoses in Estonia as compared to the UK Biobank. Thus, while these results highlight the potential of cross-biobank meta-analyses to improve genetic predictors, they also highlight the likely trait dependency of applicability of predictors across different health systems. Nevertheless, we show high stratification across both populations of early-onset risk groups with individuals in the top 1% of predicted genetic values having 7 times (95% CI 4-9) higher risk of cardiovascular disease (CAD), 8 times (95% CI 6-11) higher risk of high blood pressure (BP) in the UK Biobank hold-out sample, and 4 times (95% CI 3-8) higher risk of type-2 diabetes (T2D) prior to 60 years of age, as compared to the rest of the population (Figure 1e). Thus, we show how Bayesian posterior SNP effects size estimates can be meta-analysed across studies to improve identification of individuals at high risk of early common disease onset.

To facilitate comparison to other mixed-linear model approaches (MLMA), our GMRM approach provides leave-one-genomic-region-out SNP effect estimates (see Online Methods), and these yield greater association testing power than comparable MLMA methods [8–10] for 5 of the 21 traits (Figure 1f, Figure S5 to S7). At regions identified by all approaches, we find that the difference in the χ^2 values obtained by GMRM-MLMA as compared to other methods, scales with the percentage difference in prediction accuracy obtained in independent samples (Figure S5), a relationship expected by theory (see Online Methods). We validate this in simulations (Online Methods, Figure S8), showing that the GMRM-MLMA approach yields higher power at associated variants, whilst controlling for pervasive population stratification. However, in the UK Biobank we find that GMRM-MLMA yields more significantly associated regions only when prediction improvements are highest (Figure 1f, Figure S5, Figure S6). This is particularly true for height where we find twice the number of findings (6904 LD independent regions), and we also find a moderate increase for MCV (1510 LD-independent regions), MCH (1234 LD-independent regions), HDL (1105 regions), and BMD (927 regions, Figure 1f). Generally, the results of all approaches were concordant at a $p \leq 0.0005$ threshold (Figure S6). The GMRM approach also provides fine-mapping of the associations (WPPA aproach, see Online Methods) and we fine-map 170 associations to single markers with posterior inclusion probability $PIP \geq 0.95$, 307 associations to SNP sets of 2-5 markers with $PIP \geq 0.95$, 497 to groups of 6-20 markers with $PIP \geq 0.95$, 5674 to 20-100 markers with $PIP \geq 0.95$, and 9725 to 100 or more SNPs within a single set with $PIP \geq 0.95$ (Figure S7). Out of a total of 18,837 total GMRM-MLMA associations 87% (16,373) were significantly fine-mapped to regional associations, and 60% these fine-mapped regions contained ≥ 100 SNPs in LD, highlighting how LD in the genome creates difficulty for pinpointing the mechanistic basis of the associations found using Biobank data alone.

There remain important limitations. All of the above analyses were conducted using 2.17 million imputed genetic markers covering all rare high-quality imputed variants (minor allele frequency > 0.001) in a sample of UK Biobank European-ancestry individuals that included relatives (N=458,747, see Online Methods, Table S1) as a demonstration of the utility of our approach. The portability of polygenic scores across the spectrum of human ethnicity needs to be addressed to avoid polygenic risk stratification that is discriminatory towards groups little represented in currently available genomic data. This is an active research area, and our future work will involve accessing performance of analyses of diverse samples and how transfer learning can improve model estimation across world-wide biobank data. Second, summary statistic approaches have been developed, some of which also account for genomic annotations, MAF and LD when creating genetic predictors [4,11,12] and these methods should not be viewed as competing as they are essential for utilising currently available summary data and for situations when individual-level data are not accessible. However, as yet, summary statistics approaches have yet to yield the prediction accuracy obtained in this study [4,11,12], and here, we have shown using empirical data from two biobanks how gains in discovery and polygenic prediction can be obtained through a focus on creating powerful and efficient software applications to maximise individual-level data analysis. Thus, we hope that previous consortia analyses can be revisited with a range of improved methodology to facilitate further gains in discovery and polygenic prediction. Our approach is coded in highly optimized C++ code and can accommodate the analysis of multiple traits at once, obtaining trait specific estimates in sub-linear time as the data size increases (Online Methods, Figure S9). This work shows that SNP-heritability estimation, association discovery and genomic prediction can be improved simply by better utilizing current data and that as sample size increases, it may become more important to model genetic associations accounting for MAF and LD differences among SNP markers and to incorporate prior knowledge of genomic function.

Figure 1. SNP-heritability estimation, prediction accuracy and discovery of a GMRM approach. (a) SNP heritability estimates of the MAF-LD-annotation GMRM as a percentage difference from those obtained from a single component REML analysis across 20 traits. (b) Prediction accuracy of the GMRM effects sizes as compared to bolt-LMM effect sizes as a percentage of their upper bound (the SNP-heritability) for 20 traits. (c) Prediction accuracy obtained by GMRM for the 20 traits as compared to bolt-LMM plotted against that expected from ridge-regression theory. (d) Prediction accuracy obtained using GMRM UK Biobank estimates in UK Biobank hold-out data (UK->UK), GMRM UK Biobank estimates in Estonian data (UK->EE), and UK Biobank and Estonian meta-analysis GMRM estimates in Estonian hold-out data (UK+EE->EE) for five focal traits. (e) Odds ratio for top 1% of the GMRM genetic predictor as compared to all others, within UK->UK and UK+EE->EE for type-2 diabetes (T2D), cardiovascular disease (CAD), and high blood pressure (BP). (f) Number of LD independent genomic regions identified at $5x10^{-8}$ in bolt-LMM (bolt-MLMA) and GMRM (GMRM-MLMA), and the number of fine-mapped regions at ≥ 0.95 posterior inclusion probability (GMRM-WPPA). Error bars in give 95% confidence intervals. Full trait code descriptions are given in Table S1.

Author Contributions 82

MRR and PMV conceived and designed the study. MRR conducted the study with input from EJO, SEO, DTB, KL and RM, with RM providing Estonian Biobank study oversight. EJO and MRR developed the software. MRR wrote the paper. All authors approved the final manuscript prior to submission.

Author competing interests

The authors declare no competing interests.

Acknowledgements

This project was funded by an SNSF Eccellenza Grant to MRR (PCEGP3-181181), and by core funding from the Institute of Science and Technology Austria. PMV acknowledges funding from the Australian National Health and Medical Research Council (1113400) and the Australian Research Council (FL180100072). K.L. and R.M. were supported by the Estonian Research Council grant PRG687. EstBB computations were performed in the High Performance Computing Centre, University of Tartu.

Data availability

The individual-level genotype and phenotype data are available through formal application to the UK Biobank (http://www.ukbiobank.ac.uk). This project uses UK Biobank data under project 35520. The Estonian Biobank data are available upon request from the cohort authors according to data access protocols to researchers with approval from Estonian Committee on Bioethics and Human Research (Estonian Ministry of Social Affairs).

Summaries of all posterior distributions obtained are deposited on Dryad:

Robinson, Matthew (2021), Maximizing GWAS discovery and genomic prediction accuracy in Biobank data, Dryad, Dataset, `https://doi.org/10.5061/dryad.gtht76hmz`

Code availability

GMRM software is fully open source and available at:

`https://github.com/medical-genomics-group/gmrm`.

Online Methods

UK Biobank data

From the measurements, tests, and electronic health record data available in the UK Biobank data [13], we selected 12 blood based biomarkers, 3 of the most common heritable complex diseases, and 6 quantitative measures. The full list of traits, the UK Biobank coding of the data used, and the covariates adjusted for are given in Table S1. For the quantitative measures and blood-based biomarkers we adjusted the values by the covariates, removed any individuals with a phenotype greater or less than 7 SD from the mean (assuming these are measurement errors), and standardized the values to have zero mean and variance 1.

For the common complex diseases, we determined disease status using a combination of information available. For high blood pressure (BP), we used self-report information of whether high blood pressure was diagnosed by a doctor (UK Biobank code 6150-0.0), the age high blood pressure was diagnosed (2966-0.0), and whether the individual reported taking blood pressure medication (6153-0.0, 6177-0.0). For type-2 diabetes (T2D), we used self-report information of whether diabetes was diagnosed by a doctor (2443-0.0), the age diabetes was diagnosed (2976-0.0), and whether the individual reported taking diabetes medication (6153-0.0, 6177-0.0). For cardiovascular disease (CAD), we used self-report information of whether a heart attack was diagnosed by a doctor (3894-0.0), the age angina was diagnosed (3627-0.0), and whether the individual reported heart problem diagnosed by a doctor (6150-0.0) the date of myocardial infarction (42000-0.0). For each disease, we then combined this with primary death ICD10 codes (40001-0.0), causes of operative procedures (41201-0.0), and the main (41202-0.0), secondary (41204-0.0) and inpatient ICD10 codes (41270-0.0). For BP we selected ICD10 codes I10, for T2D we selected ICD10 codes E11 to E14 and excluded from the analysis individuals with E10 (type-1 diabetes), and for CAD we selected ICD10 code I20-I29. Thus, for the purposes of this analysis, we define these diseases broadly simply to maximise the number of cases available for analysis. For each disease, individuals with neither a self-report indication or a relevant ICD10 diagnosis, were then assigned a zero value as a control.

We restricted our discovery analysis of the UK Biobank to a sample of European-ancestry individuals. To infer ancestry, we used both self-reported ethnic background (21000-0) selecting coding 1 and genetic ethnicity (22006-0) selecting coding 1. We also took the 488,377 genotyped participants and projected them onto the first two genotypic principal components (PC) calculated from 2,504 individuals of the 1,000 Genomes project with known ancestries. Using the obtained PC loadings, we then assigned each participant to the closest population in the 1000 Genomes data: European, African, East-Asian, South-Asian or Admixed, selecting individuals with PC1 projection < absolute value 4 and PC 2 projection < absolute value 3. Samples were excluded if in the UK Biobank quality control procedures they (i) were identified as extreme heterozygosity or missing genotype outliers; (ii) had a genetically inferred gender that did not match the self-reported gender; (iii) were identified to have putative sex chromosome aneuploidy; (iv) were excluded from kinship inference; (v) had withdrawn their consent for their data to be used. We used the imputed autosomal genotype data of the UK Biobank provided as part of the data release. We used the genotype probabilities to hard-call the genotypes for variants with an imputation quality score above 0.3. The hard-call-threshold was 0.1, setting the genotypes with probability ≤ 0.9 as missing. From the good quality markers (with missingness less than 5% and p-value for Hardy-Weinberg test larger than 10-6, as determined in the set of unrelated Europeans) we selected those with minor allele frequency (MAF) > 0.0002 and rs identifier, in the set of European-ancestry participants, providing a data set 9,144,511 SNPs. From this we took the overlap with the Estonian Genome centre data described below to give a final set of 8,430,446 markers. For computational convenience we then removed markers in very high LD selecting one marker from any set of markers with LD $R^2 \geq 0.8$ within a 1MB window. These filters resulted in a data set with 458,747 individuals and 2,174,071 markers.

We split the sample into training and testing sets for each phenotype, selecting 30,000 individuals that were unrelated (SNP marker relatedness <0.05) to the training individuals to use as a testing set. This provides an independent sample of data with which to access prediction accuracy. For the complex diseases, we randomly select 1000 cases and match to 29,000 controls, again ensuring that these individuals were unrelated to those in the training sample.

Estonian Biobank data

The Estonian Biobank (EstBB) is a population-based cohort encompassing 20% of Estonia's adult population (200'000 individuals; 66% females; https://genomics.ut.ee/en/research/estonian-biobank). Individuals underwent microarray-based genotyping at the Core Genotyping Lab of the Institute of Genomics, University of Tartu. 136,421 individuals were genotyped on Illumina Global Screening (GSA) arrays and we imputed the data set to an Estonian reference, created from the whole genome sequence data of 2,244 participants [14]. From 11,130,313 markers with imputation quality score > 0.3, we selected SNPs that overlapped with those selected in the UK Biobank, resulting in a set of 2,174,071 markers.

General data, including basic body measurements, were collected at recruitment. Project-based questionnaires were sent later and filled on a voluntary basis. Health records are regularly updated through linkage with the national Health Insurance Fond and other relevant databases, providing sporadic access to blood biomarker measurements and medical diagnoses [15]. For the genotyped individuals, we had data available for height and body mass index and we removed individuals plus or minus 7SD from the mean and adjusted both phenotypes by the age at enrollment, sex, and the first 20 PCs of the SNP marker data. Prevalent cases of BP, CAD and T2D in the Estonian Biobank cohort were first identified on the basis of the baseline data collected at recruitment, where the information on prevalent diseases was either retrieved from medical records or self-reported by the participant. The cohort was subsequently linked to the Estonian Health Insurance database that provided additional information on prevalent cases (diagnoses confirmed before the date of recruitment) as well as on incident cases during the follow-up. For BP we selected ICD10 code I10, for CAD we selected codes of I20-I29, and for T2D we selected codes E11-E14 and excluded E10. We also split the sample into training and testing sets for each phenotype, selecting 20,000 individuals that were unrelated (SNP marker relatedness <0.05) to the training individuals to use as a testing set. This provides an independent sample of data with which to access prediction accuracy. For the complex diseases, we randomly select 1000 cases and match to 19,000 controls, again ensuring that these individuals were unrelated to those in the training sample.

All participants signed a broad informed consent form and analyses were carried out under ethical approval 1.1-12/2856 by Estonian Committee on Bioethics and Human Research and data release P03 from the EstBB.

Multiple outcome Bayesian grouped mixture of regressions model (GMRM)

We extend the software implementation of our recently developed Bayesian grouped mixture of regression model [3] to accommodate the analysis of multiple traits simultaneously and to provide standard mixed linear model association results. While this approach does not yet utilise estimates of the genetic or residual covariance among different outcomes when estimating the SNP effects, a number of coding developments were made to improve speed of the baseline calculations and facilitate the random number generation, vectorisation of the effect size estimation and missing data handling for multiple outcomes. We bench-marked the timing of our software increasing the sample size, marker number, number of traits analysed, and the number of MPI processes, to demonstrate the scalability of our approach as the dimensionality of the data increases. We present these results in Figure S9.

Following our previous work, we partition SNP markers into 7 location annotations using the knownGene table from the UCSC browser data, preferentially assigned SNPs to coding (exonic) regions first, then in the remaining SNPs we preferentially assigned them to intronic regions, then to 1kb upstream regions, then to 1-10kb regions, then to 10-500kb regions, then to 500-1Mb regions. Remaining SNPs were grouped in a category labelled "others" and also included in the model so that variance is partitioned relative to these also. Thus, we assigned SNPs to their closest upstream region, for example if a SNP is 1kb upstream of gene X, but also 10-500kb upstream of gene Y and 5kb downstream for gene Z, then it was assigned to be a 1kb region SNP. This means that SNPs 10-500kb and 500kb-1Mb upstream are distal from any known nearby genes. We further partition upstream regions to experimentally validated promoters, transcription factor binding sites (tfbs) and enhancers (enh) using the HACER, snp2tfbs databases (see Code Availability). All SNP markers assigned to 1kb regions map to promoters; 1-10kb SNPs, 10-500kb SNPs, 500kb-1Mb SNPs are split into enh, tfbs and others (un-mapped SNPs) extending the model to 13 annotation groups. Within each of these annotations, we have three minor allele frequency groups (MAF<0.01, 0.01>MAF>0.05, and MAF>0.05), and then each MAF group is further split into 2 based on median LD score. This gives 78 non-overlapping groups for which our model jointly estimates the phenotypic variation attributable to, and the SNP marker

effects within, each group. For each of the 78 groups, SNPs were modelled using five mixture groups with variance equal to the phenotypic variance attributable to the group multiplied by constants (mixture $0 = 0$, mixture $1 = 0.00001$, mixture $2 = 0.0001$, $3 = 0.001$, $4 = 0.01$, $5 = 0.1$). The probabilities that markers enter each of the five mixtures (the mixture components) and the variance attributable to the groups are all estimated independently, linked only by a residual updating scheme and thus a regression problem where the number of covariates greatly exceeds the number of measured individuals is broken down into a series of interdependent regressions where the number of covariates within a group is always far less than the total sample size.

We then extend the software to return the typical fixed effect SNP regression coefficients estimated by other mixed linear model approaches (MLMA). In the first step of running GMRM, the SNP marker effects that we obtain are jointly estimated and thus within each iteration estimation is made accounting for the effects of other markers in both short- and long-range LD. From these estimates, we can obtain partitioned predictors for each focal block k of the genome $\tilde{\mathbf{g}}_{\text{block}_k} = \sum_{j=1}^{j} \mathbf{x}_{j,-k} \beta_{j,-k}$, where $\mathbf{x}_{j,-k}$ are the values of SNP markers that are not part of block k and $\beta_{j,-k}$ their jointly estimated posterior mean effects. These can then be used when testing for association in a second step to yield standard frequentist mixed-model summary statistics, following other approaches. When testing for association of the phenotype with a marker $\mathbf{x}_j$ from focal block k we consider a simple linear model:

$$\tilde{\mathbf{y}}_{\text{block}_k} = \mathbf{x}_{j,k} \beta_{j,k} + \boldsymbol{\epsilon}_k \tag{1}$$

where $\tilde{\mathbf{y}}_{\text{block}_k} = \tilde{\mathbf{y}} - \tilde{\mathbf{g}}_{\text{block}_k}$ gives the phenotypic residuals where the polygenic effects estimated across the genome other than the focal testing block are adjusted for, $\mathbf{x}_j$ is the j^{th} marker in the focal block, β_j is the OLS estimate for the j^{th} marker in block k and $\boldsymbol{\epsilon}_k$ is the residual error, with $\boldsymbol{\epsilon} \sim N\left(0, \mathbf{I}_N \sigma_\epsilon^2\right)$ with M_k the number of markers within block k.

A t-test statistic is then straight forward to obtain as:

$$T_{j,k} = \frac{\mathbf{x}_{j,k}^T \tilde{\mathbf{y}}_{\text{block}_k}}{[\sigma_{\tilde{\mathbf{y}}_{\text{block}_k}}^2 \mathbf{x}_{j,k}^T \mathbf{x}_{j,k}]^{0.5}} \tag{2}$$

where $\sigma_{\tilde{\mathbf{y}}_{\text{block}_k}}^2$ is calculated as $\frac{1}{N}||\tilde{\mathbf{y}}_{\text{block}_k}^T||_2^2$ where N is the number of individuals. A normal approximation of $T_{j,k}^2 \approx \chi_1^2$ is used to give the p-value. A step-by-step algorithm is given in Algorithm 1.

The test statistic values obtained are an approximation of the mixed-model χ^2 statistic if one were to model the SNP as a fixed effect with a full mixed model equation. This approach follows recent studies (Eq. 23 and 24 of Supplementary Online Material of ref [8] and ref [10]), in particular as the χ^2 statistic obtained from bolt-LMM is equivalent to computing the squared correlations between SNPs being tested and a best linear unbiased predictor, which is the approach taken here. The power of mixed-model association is driven by the fact that focal test SNPs are tested against a 'denoised' residual phenotypes, from which other SNP effects estimated by the mixed model have been conditioned out. Here, we expect the SNP marker effect estimation to be improved as the genetic predictor used to residualise the phenotype should have a higher prediction accuracy within independent sample if the true underlying SNP marker effects differ across MAF-LD-annotation groups. While further work is required to optimise this approach for rare diseases using saddle point approximations and to account for the covariance of multiple outcomes, here we wished to simply demonstrate that modelling SNP effects in MAF-LD-annotation groups in the first step yields improved MLMA fixed effect marker estimates in the second step. We verify this also in a simulation study as described below.

Simulation study

We follow a similar simulation study design as presented in a number of recent studies [9]. From our quality controlled UK Biobank data, we first randomly selected 100,000 individuals with relatedness estimated from SNP markers of ≤ 0.05 and used marker data from chromosomes 1 through 10 with MAF ≥ 0.001 to give 1.36 million SNPs. For the odd chromosomes, we randomly selected 10,000 LD independent SNP markers as causal variants. We simulated effect sizes for the causal variants, $\mathbf{b}$, by drawing from a normal distribution with zero mean and variance $0.5/10000$. We then scaled the 10,000 causal variant SNP markers to have mean 0 and variance 1 and multiplied them by the simulated marker effect sizes to give genetic values with mean 0

variance 0.5. In previous work, we have extensively explored the ability of our GMRM approach to recover SNP effects sizes and accurately estimate SNP-heritability parameters independently of the relationship of effect sizes, MAF and LD, and so here we simply assume a simple underlying genetic model of randomly selected causal variants and no relationship between effect size and MAF or LD. We then simulate population stratification effects by scaling the loadings of the 100,000 individuals on the first principal component calculated from the genetic data and multiplying the values by $\sqrt{0.05}$, this gives a vector of values, ps with variance 0.05. Finally, individual environment values are drawing from a normal distribution with zero mean and variance $1 - \text{var}(\mathbf{Xb}) - \text{var}(ps)$, so that the sum of the genetic, ps, and environmental values give a phenotype with zero mean and variance 1. We replicate this simulation 10 times, referring to it in the main text and the figures as "random unrelated" causal variant allocation.

Using the same randomly selected individuals, we repeat the simulation but we separate the genetic markers into 4 groups randomly selecting 1000 LD independent exonic markers, 1000 LD independent intronic markers, 1000 LD independent transcription factor binding site markers, and 7000 LD independent markers from other annotation groups. We then sample the marker effects for each of the four groups, drawing from independent normal distributions with zero mean and variance 0.1/1000, 0.25/1000, 0.1/1000, and 0.05/7000 respectively. This gives three annotation groups with larger effect size variance but the same number of causal variants and a total variance explained by the SNP markers $\text{var}(\mathbf{Xb}) = 0.5$ as the previous simulation setting. We then simulated population stratification effects in the same way and individual environment values by drawing from a normal distribution with zero mean and variance $1 - \text{var}(\mathbf{Xb}) - \text{var}(ps)$, so that the sum of the genetic, ps, and environmental values give a phenotype with zero mean and variance 1. We also replicate this simulation setting 10 times, referring to it in the main text and the figures as "enrichment unrelated" causal variant allocation.

We then repeat the "random" and the "enrichment" simulation settings, but we change the selection of the individuals used for the simulation. We randomly select 10,000 unique sibling pairs from the UK Biobank data and combine this with 80,000 randomly selected unrelated individuals to give a mixture of relatedness similar to the proportions of related and unrelated individuals in the UK Biobank data. We simulate the same genetic and ps values, but we also add a common environmental variance by drawing a value for each sibling pair from a normal distribution with mean 0 and variance 1 and allocated a value of 0 to each of the unrelated individuals. We scale these values by $\sqrt{0.12}$ to give a vector of among-family differences, pe, with variance 0.12. Finally, individual environment values are drawing from a normal distribution with zero mean and variance $1 - \text{var}(\mathbf{Xb}) - \text{var}(ps) - \text{var}(pe)$, so that the sum of the genetic, ps, pe, and environmental values give a phenotype with zero mean and variance 1. We replicate this simulation 10 times, referring to it in the main text and the figures as "random related" and "enrichment related" causal variant allocation.

We then analyse this data using bolt-LMM [8], fastGWA [9], and GMRM-MLMA and calculate the average χ^2 values (using an approximation that $T^2 \approx \chi^2$) for the causal variants, which gives a comparative measure of the power of each approach. We also calculate the average χ^2 values (again using an approximation that $T^2 \approx \chi^2$) obtained from markers on the even chromosomes that contain no causal variants, comparing to the null expectation of 1. These results are presented in Figure S8. We find improved power of GMRM-MLMA over other approaches in all settings, with control of the pervasive population stratification. We find only moderate χ^2 inflation when relatives share strong common environment effects that is the same as that obtained by bolt-LMM. This inflation never increased the false discovery rate (FDR) above 2% in any of the simulations, but was best controlled by a fastGWA approach.

Utilizing the posterior distribution obtained

We apply our model to each UK Biobank and Estonian Genome Centre data trait, running two short chains for 5000 iterations and combining the last 2000 posterior samples together. We show in Figure S10 that the prediction accuracy obtained from our model and the hyperparameter estimates of h^2_{SNP} converge within the first 2000 iterations. While obtaining a full posterior distribution with many hundreds of independent samples would require running longer chains, we show that the posterior mean effect size for each SNP that we use for prediction (and thus also in the estimation of the MLMA effect sizes) is well approximated within this run time (Figure S10b), which is sufficient in this work to assess the prediction accuracy of our approach and to estimate the variance attributable to different genomic regions.

We estimate the proportion of total h^2_{SNP} attributable to each genomic annotation and we divide this

proportion, by the proportion of SNP markers in the model for this annotation given the total number of SNP markers in the model. This gives an estimate of the enrichment of the marker effects, whereby if the average effect sizes of markers within a given annotation are larger than expected given the number of markers entering the model for that annotation then the value obtained should be greater than 1. Conversely, smaller than expected marker effects will yield values less than 1. We present these results in Figure S1, where we find substantial enrichment of SNP-heritability in intronic regions across traits, and evidence that enrichment differs across traits, with exonic enrichment for height, forced vital capacity, mean corpuscular heamoglobin, low and high density lipoprotein, and blood cholesterol levels (Figure S1). Generally, blood based biomarkers show enrichment at proximal promotors, transcription factor binding sites and enhancers, with variation in complex diseases and quantitative traits attributable to distal transcription factor binding sites and enhancers in proportion to that expected given the number of markers in the model (Figure S1). SNP markers located greater than 500kb from a gene, explained a far smaller proportion of variance explained then expected given the number of markers which map to the region (Figure S1).

We previously presented a window posterior probability of association approach (WPPA) [3]. The WPPA, is estimated by counting the proportion of MCMC samples in which the regression coefficent β_j is greater than a given threshold for at least one SNP j in a given genomic window, which can be used as a proxy for the posterior probability that the genomic region contains a causal variant. Because WPPA for a given window is a partial association conditional on all other SNPs in the model, including those flanking the region, the influence of flanking markers on the WPPA signal for any given window will be inversely related to the distance k of the flanking markers. Thus, as the number of markers between a causal variant and the focal window increases, the influence of the causal variant on the WPPA signal will decrease and so WPPA computed for a given window can be used to locate associations for that given window, whilst also controlling the false discovery rate. Thus, it represents an approach to fine-mapping association results to groups of SNP markers. Here, we group markers by LD into 341,380 LD independent groups using plink's clumping procedure, which selects groups of markers (from single SNPs to groups of 100 or more) with LD $R^2 \geq 0.1$ within a 1Mb window. For each of these 341,380 SNP groups we calculate the WPPA, defined here as the posterior probability that a group explains at least 0.0001% of the phenotypic variance and we present the number of groups with WPPA ≥ 0.95.

Comparisons to theory

We note the important distinction between prediction SNP effect sizes that are obtained from mixed-models/penalized regression models, in which a single model is fit where all SNPs are included as random effects, and those obtained from an MLMA model where individual SNP association estimates are typically sought. Here, we sought to make this distinction clear by comparing the prediction accuracy obtained by GMRM and it's baseline BayesR model, to that obtained from using MLMA estimates to demonstrate that it is inappropriate to create polygenic risk scores from MLMA estimates as compared to random effect models. Thus, we also analysed the same data using existing approaches for large-scale analysis, which included BOLT-LMM (version 2.3; https://data.broadinstitute.org/alkesgroup/BOLT-LMM/) and fastGWA (GCTA version 1.93.0beta; https://cnsgenomics.com/software/gcta/#Overview). The expected prediction accuracy in an independent sample under ridge regression assumptions is given by ref [16] as:

$$R^2 = \frac{h_{SNP}^2}{1 + \frac{M}{Nh_{SNP}^2}(1 - R^2)} \tag{3}$$

and we use this equation, with the number of markers in the model as a proxy for M and the estimates of h_{SNP}^2 obtained from GMRM, to compare to the prediction accuracy we obtain.

For the common complex disease traits, we place the estimates of the proportion of variance explained by the SNP markers on the liability scale to facilitate comparison with the quantitative measures. The linear transformation of heritability from the observed 0-1 scale, h_o^2, to that of liability scale h_l^2 is:

$$h_l^2 \sim \frac{h_o^2 K(1 - K)}{z^2} \tag{4}$$

with K the population lifetime prevalence and z the height of the normal curve at the truncation point pertaining to K [17]. We did not observe the population lifetime prevalence of any disease within either

population and so we make the assumption that the prevalence in the UK Biobank and Estonian Biobank samples provides a very distant approximation. We scale the estimates of the prediction accuracy to also place these on the liability scale, R_l^2, as:

$$R_l^2 \sim \frac{R_o^2 K^2 (1 - K^2)}{z^2 P(1 - P)} \tag{5}$$

with P the proportion of cases in the testing sample. Note that this expression is also an approximation and can result in bias when ascertainment is extreme and heritability on the liability scale is high, though this is expected to be small in practice and negligible for the three disease trait considered here [18].

Previous work [19] has shown that the expectation of the mean χ^2 value at causal SNPs is given by:

$$E[\chi^2_{\text{causal SNPs}}] = 1 + \frac{N h_{SNP}^2}{M_{causal} \cdot \frac{1}{1 - R^2}} \tag{6}$$

with N the sample size. Thus, under the assumptions of no confounding, no case-control ascertainment-induced confounding, and no pervasive familial relatedness, increased prediction accuracy should yield increased power (χ^2 statistics) at SNPs that are associated with underlying causal variants. Here, the use of relatives in the analysis may complicate the relationship, but with sufficient power, theory predicts that if the model improves predictive performance than power in an MLMA approach should improve proportionally.

References

1. IM MacLeod, PJ Bowman, CJ Vander Jagt, M Haile-Mariam, KE Kemper, AJ Chamberlain, C Schrooten, BJ Hayes, and ME Goddard. Exploiting biological priors and sequence variants enhances qtl discovery and genomic prediction of complex traits. *BMC Genomics*, 17(1):144, 2016.

2. Yiming Hu, Qiongshi Lu, Ryan Powles, Xinwei Yao, Can Yang, Fang Fang, Xinran Xu, and Hongyu Zhao. Leveraging functional annotations in genetic risk prediction for human complex diseases. *PLOS Computational Biology*, 13(6):1–16, 06 2017.

3. Marion Patxot, Daniel Trejo Banos, Athanasios Kousathanas, Etienne J. Orliac, Sven E. Ojavee, Gerhard Moser, Alexander Holloway, Julia Sidorenko, Zoltan Kutalik, Reedik Mägi, Peter M. Visscher, Lars Rönnegård, and Matthew R. Robinson. Probabilistic inference of the genetic architecture underlying functional enrichment of complex traits. *medRxiv*, 2021.

4. Qianqian Zhang, Florian Privé, Bjarni Vilhjálmsson, and Doug Speed. Improved genetic prediction of complex traits from individual-level data or summary statistics. *Nature Communications*, 12(1):4192, 2021.

5. Doug Speed, John Holmes, and David J Balding. Evaluating and improving heritability models using summary statistics. *Nature Genetics*, 52(4):458–462, 2020.

6. Luke M Evans, Rasool Tahmasbi, Scott I Vrieze, Gonçalo R Abecasis, Sayantan Das, Steven Gazal, Douglas W Bjelland, Teresa R De Candia, Michael E Goddard, Benjamin M Neale, et al. Comparison of methods that use whole genome data to estimate the heritability and genetic architecture of complex traits. *Nature Genetics*, 50(5):737–745, 2018.

7. Kangcheng Hou, Kathryn S Burch, Arunabha Majumdar, Huwenbo Shi, Nicholas Mancuso, Yue Wu, Sriram Sankararaman, and Bogdan Pasaniuc. Accurate estimation of snp-heritability from biobank-scale data irrespective of genetic architecture. *Nature Genetics*, page 1, 2019.

8. Po-Ru Loh, George Tucker, Brendan K Bulik-Sullivan, Bjarni J Vilhjalmsson, Hilary K Finucane, Rany M Salem, Daniel I Chasman, Paul M Ridker, Benjamin M Neale, Bonnie Berger, et al. Efficient bayesian mixed-model analysis increases association power in large cohorts. *Nature Genetics*, 47(3):284, 2015.

9. Longda Jiang, Zhili Zheng, Ting Qi, Kathryn E. Kemper, Naomi R. Wray, Peter M. Visscher, and Jian Yang. A resource-efficient tool for mixed model association analysis of large-scale data. *Nature Genetics*, 51(12):1749–1755, 2019.

10. Joelle Mbatchou, Leland Barnard, Joshua Backman, Anthony Marcketta, Jack A. Kosmicki, Andrey Ziyatdinov, Christian Benner, Colm O'Dushlaine, Mathew Barber, Boris Boutkov, Lukas Habegger, Manuel Ferreira, Aris Baras, Jeffrey Reid, Goncalo Abecasis, Evan Maxwell, and Jonathan Marchini. Computationally efficient whole-genome regression for quantitative and binary traits. *Nature Genetics*, 53(7):1097–1103, 2021.

11. Gao Wang, Abhishek Sarkar, Peter Carbonetto, and Matthew Stephens. A simple new approach to variable selection in regression, with application to genetic fine mapping. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 82(5):1273–1300, 2020.

12. Luke R. Lloyd-Jones, Jian Zeng, Julia Sidorenko, Loïc Yengo, Gerhard Moser, Kathryn E. Kemper, Huanwei Wang, Zhili Zheng, Reedik Magi, Tõnu Esko, Andres Metspalu, Naomi R. Wray, Michael E. Goddard, Jian Yang, and Peter M. Visscher. Improved polygenic prediction by bayesian multiple regression on summary statistics. *Nature Communications*, 10(1):5086, 2019.

13. Clare Bycroft, Colin Freeman, Desislava Petkova, Gavin Band, Lloyd T Elliott, Kevin Sharp, Allan Motyer, Damjan Vukcevic, Olivier Delaneau, Jared O'Connell, et al. The uk biobank resource with deep phenotyping and genomic data. *Nature*, 562(7726):203–209, 2018.

14. Mario Mitt, Mart Kals, Kalle Pärn, Stacey B Gabriel, Eric S Lander, Aarno Palotie, Samuli Ripatti, Andrew P Morris, Andres Metspalu, Tõnu Esko, Reedik Mägi, and Priit Palta. Improved imputation accuracy of rare and low-frequency variants using population-specific high-coverage wgs-based imputation reference panel. *Eur J Hum Genet*, 25(7):869–876, Jun 2017.

15. Tõnis Tasa, Kristi Krebs, Mart Kals, Reedik Mägi, Volker M. Lauschke, Toomas Haller, Tarmo Puurand, Maido Remm, Tõnu Esko, Andres Metspalu, Jaak Vilo, and Lili Milani. Genetic variation in the estonian population: pharmacogenomics study of adverse drug effects using electronic health records. *European Journal of Human Genetics*, 27(3):442–454, 2019.

16. Hans D. Daetwyler, Beatriz Villanueva, and John A. Woolliams. Accuracy of predicting the genetic risk of disease using a genome-wide approach. *PLOS ONE*, 3(10):1–8, 10 2008.

17. E R Dempster and I M Lerner. Heritability of threshold characters. *Genetics*, 35(2):212–236, 03 1950.

18. Sang Hong Lee, Naomi R Wray, Michael E Goddard, and Peter M Visscher. Estimating missing heritability for disease from genome-wide association studies. *Am J Hum Genet*, 88(3):294–305, Mar 2011.

19. Jian Yang, Noah A Zaitlen, Michael E Goddard, Peter M Visscher, and Alkes L Price. Advantages and pitfalls in the application of mixed-model association methods. *Nature Genetics*, 46(2):100–106, 2014.

Algorithm 1: GMRM MLMA algorithm

Data: Matrix $\mathbf{X}$ of genotypes in plink binary format (.bed/.bim/.fam), matrix of posterior Bayesian marker estimates in binary format (.bet file), a file linking column number to rs SNP identifiers (.link), and file of phenotypic information (.phen file).

1 Assign markers present in SNP rs-identifier file to blocks across k MPI tasks

2 Assign the posterior Bayesian marker estimates to the markers of each task k based on the rs SNP identifiers

3 foreach *task k* **do**

4   Calculate means $\bar{x}_j$ and $\mathrm{SD}(x_j)$ of markers

5   Initialise $\mathbf{g}_{k,l}$ and $\bar{\mathbf{g}}_k$

6   **foreach** *iteration l in iter* **do**

7   **foreach** *marker j* **do**

8   **if** $\beta_j == 0$ **then**

9   next

10   **if** $\beta_j ! == 0$ **then**

11   calculate $\hat{\mathbf{g}}_j$ as $\frac{\mathbf{x}_j - \bar{x}_j}{\mathrm{SD}(x_j)} \beta_j$;

12   add $\hat{\mathbf{g}}_j$ to $\mathbf{g}_{k,l}$

13   Add $\mathbf{g}_{k,l}$ to $\mathbf{g}_k$

14   Communicate $\mathbf{g}_k$ to task 0 where task 0 calculates $\mathbf{g} = \sum_{l=1}^{l} \sum_{k=1}^{k} \mathbf{g}_{k,l}$

15   Receive $\mathbf{g}$ from node 0

16   Subtract $\mathbf{g} - \mathbf{g}_k$ then divide by iter to give $\bar{\mathbf{g}}_k$

17   Calculate $\tilde{\mathbf{y}}_{\mathrm{block}_k}$ as $\tilde{\mathbf{y}}_{\mathrm{block}_k} = \mathbf{y} - \bar{\mathbf{g}}_k$

18   Calculate $\sigma^2_{\tilde{\mathbf{y}}_{\mathrm{block}_k}}$ as $\frac{1}{N} \tilde{\mathbf{y}}^T_{\mathrm{block}_k} \tilde{\mathbf{y}}_{\mathrm{block}_k}$ where N is the number of individuals

19   **foreach** *marker j* **do**

20   calculate xtx as dot product $\mathbf{x}_j^T \mathbf{x}_j$

21   calculate xty as $\mathbf{x}_j^T \tilde{\mathbf{y}}_{\mathrm{block}_k}$

22   calculate $\beta_{j,k}$ as $[\mathrm{xtx}]^{-1}\mathrm{xty}$

23   calculate t_j as $\frac{\mathrm{xty}}{[\sigma^2_{\tilde{\mathbf{y}}_{\mathrm{block}_k}} \mathrm{xtx}]^{0.5}}$

24   calculate p-value j from χ_1^2 using t_j^2

25   output $\beta_{j,k}, t_j$ and corresponding χ_1^2 p-values.

Type	Phenotype	Code	Biobank codes	Adjusted covariate codes
Biomarker	Basophill count	BASO	30160-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Biomarker	Cholesterol	CHOL	30690-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20, Medication (6177-0.0)
Biomarker	Creatinine	CRET	30700-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Biomarker	Eosinophill count	EOSI	30150-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Biomarker	Glucose	GLU	30740-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20, Medication (6177-0.0)
Biomarker	Glycated haemoglobin	HbA1c	30750-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Biomarker	HDL cholesterol	HDL	30760-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20, Medication (6177-0.0)
Biomarker	LDL direct	LDL	30780-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20, Medication (6177-0.0)
Biomarker	Lymphocyte count	LYMPH	30120-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Biomarker	Mean corpuscular haemoglobin	MCH	30050-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Biomarker	Mean corpuscular volume	MCV	30040-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Biomarker	Red blood cell count	RBC	30010-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Disease	Cardiovascular disease	CAD	3894-0.0, 3627-0.0, 42000-0.0, 6150-0.0, 40001-0.0, 41202-0.0, 41204-0.0, 41270-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Disease	High blood pressure	BP	6150-0.0, 2966-0.0, 6153-0.0, 6177-0.0, 40001-0.0, 41201-0.0, 41202-0.0, 41204-0.0, 41270-0.0, 41262-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Disease	Type-2 diabetes	T2D	2443-0.0, 2976-0.0, 6153-0.0, 6177-0.0, 40001-0.0, 41201-0.0, 41202-0.0, 41204-0.0, 41270-0.0, 41262-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Measures	Heel bone mineral density T-score	BMD	3148-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Measures	Body mass index	BMI	21001-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Measures	Diastolic blood pressure	DBP	4079-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20, Medication (6177-0.0)
Measures	Forced vital capacity	FVC	3062-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Measures	Standing height	HT	50-0.0	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20
Measures	Systolic blood pressure	SBP	4080-0.1	Sex, Age, East-West coordinates, UK Biobank Centre, Genotype Batch, PCs 1-20, Medication (6177-0.0)

Table S1. UK Biobank phenotypes used within the study. Columns of the table give the type of trait, the name, the trait code used for the figures, the UK Biobank codes used to construct the trait values, and the covariates adjusted for within the analysis.

Figure S1. SNP-heritability enrichment estimated by GMRM. SNP heritability enrichment estimates calculated as the proportion of SNP-heritability attributable to each annotation group divided by the proportion of markers in the model given the total number of markers. If the average effect sizes of markers within a given annotation are larger than expected given the number of markers entering the model for that annotation then the value obtained should be greater than 1 (the red line shown). Conversely, smaller than expected marker effects will yield values less than 1.Error bars in give 95% credible intervals. Full trait codes are given in Table S1. exons = SNPs located in exonic regions, introns = SNPs located in intronic regions, prom = SNPs located in promotors, tfbs = SNPs located in transcription factor binding sites, enh = enhancers, oth = markers not mapping to a functional group but to a location from a nearest gene, others = markers not within 1Mb of a gene and that have no known regulatory function.

Figure S2. Prediction accuracy as percentage of SNP-heritability across approaches Prediction accuracy in a UK Biobank hold-out sample, showing improvements of random effect model implemented in GMRM against a baseline bayesR model (GMRM-bR) and predictors created from effects sizes obtained from mixed linear association model approaches. Error bars give the 95% confidence intervals of the estimates.

Figure S3. Prediction accuracy improvement of GMRM above other approaches. Error bars give the 95% confidence intervals for the estimates of the percentage difference in prediction accuracy of GMRM over other approaches for each trait.

Figure S4. Prediction accuracy obtained by GMRM differs to other methods in proportion to the SNP-heritability of the trait. As sample size in the UK Biobank is similar across traits, differences in power are approximately represented by the differences in SNP-heritability estimates. We find evidence that GMRM prediction improvements over MLMA and a baseline BayesR model (GMRM-bR) scale with the differences in SNP-heritability estimates across traits.

Figure S5. Difference in χ^2 values obtained by the GMRM mixed linear model association approach plotted against difference in prediction accuracy of GMRM from other approaches. As the prediction accuracy difference obtained by GMRM increases the difference in χ^2 value at the shared top associated loci for each trait increases.

Figure S6. Concordance of the findings of the mixed linear model association approaches. The top column labels gives the focal analysis which is compared to the analysis listed below in the second row of column labels. For example, the first column of barplots gives the proportion of regions containing a SNP with p-value less than $5\text{x}10^{-8}$ discovered by bolt-LMM software for each trait, that are replicated at different p-value thresholds using the GMRM-mlma software. GMRM-wppa is the Bayesian fine-mapping approach that localises the inclusion of markers in the model into SNP sets based on their LD.

Figure S7. GMRM Bayesian fine-mapping approach to localise the inclusion of markers in the model into SNP sets based on their LD. (a) The trait acronyms are given on the y-axis and the plot is faceted into five columns representing set of SNPs that are correlated at LD $R^2 \geq 0.1$, from some genomic regions having only a single SNP in LD with no others, to ≥ 100 markers in LD. Barplots give a count of the number of SNP sets for which the posterior probability of explaining greater than 0.001% of the phenotypic variance was greater than 95%. (b) For each trait, the percentage of SNP sets for which the posterior probability of explaining greater than 0.001% of the phenotypic variance was greater than 95% for different sizes of SNP set. (c) GMRM-MLMA significantly associated SNPs grouped into the same SNP sets based on LD. Across traits, a lower percentage of associations fine-map to either single SNPs, or to groups of five or less SNPs in LD as compared to the percentage of the discoveries from the GMRM-MLMA approach.

a

b

Figure S8. Simulation study for GMRM. (a) The y-axis gives the average χ^2 values for the approximate mixed linear model test statistics at the simulated causal variants as compared to models run using fastGWA and bolt-LMM. The plot is faceted into two columns representing phenotypes that were simulated using either a mixture of 10,000 randomly sampled UK Biobank sibling pairs and 80,000 randomly sampled unrelated UK Biobank individuals (related), or 100,000 randomly sampled UK Biobank individuals with SNP marker relatedness estimates <0.05 (unrelated). The plot is also faceted into two rows representing 10,000 causal variants from chromosomes 1, 3, 5, 7 and 9 that were either randomly sampled with effect sizes drawn from a normal distribution with zero mean variance 0.5/10000 (random) or were sampled differently across genomic annotations creating effects size differences acorss genomic groups (see Methods). Boxplots show the distribution of values obtained for ten simulation replicates. (b) shows the same simulation settings but the average χ^2 values for the approximate mixed linear model test statistics are given for the null chromosomes 2, 4, 6, 8, and 10 where no causal variants were simulated.

Figure S9. Scaling of GMRM software for an analysis of 2.2M markers. (a) For two resource settings of (1) 12 MPI processes each using 4 CPU cores on a single computer and 24 MPI processes each using 4 CPU cores across two computers, labelled as 1 and 2 on the x-axis, the factor speed-up per trait analysed is given on the y-axis. Speed-up factors are presented relative to the time taken to estimate 5,000 draws from the posterior distribution of a single trait in resource setting 1 (baseline time). We calculated the speed-up times for data from 10,000 individuals (10k) and 100,000 individuals (100k) separately. For example, for the analysis of 5 traits at the same time for 10k setting in resource setting 2 the value presented was calculated as: $t_{1,1}/(t_{5,2}/nt)$, where $t_{1,1}$ is the time at baseline for 1 trait in resource environment 1, $t_{5,1}$ is the time for analysing 5 traits simultaneously in resource environment 2, and nt is the number of traits analysed. Thus we obtain the proportional increase in time obtained relative to the baseline. (b) The number of hours required to obtain 5,000 posterior samples at two different sample sizes, decreases steeply but then begins to tail-off with a large number of simultaneously analysed traits. (c) In the configurations presented here, our algorithm displays sub-linear scaling of CPU core hours per trait per sample, with improved CPU use efficiency as sample size and number of traits analysed increases, essentially becoming more efficient as sample size and the number of traits grows. These examples were specifically designed to remain within memory and last level memory cache restrictions of the high-performance compute hardware used here in order to demonstrate the performance of our algorithm. Mostly, with the multi-trait version some MPI communication is shared among the traits, explaining the sub-linear scaling observed here. (d) In our hardware, at 400,000 individuals, 2.2M markers, and 72 MPI processes last-level cache size overflowed at greater than two traits analysed simultaneously, resulting in increased CPU hours per trait per sample (CPU hours), reduced speed-up, and an increase in the analysis time per trait. Thus, performance gains from our algorithm will become hardware limited. If the epsilon vectors stay in the cache we get continually improved performance as shown in (a) through (c). However, with more individuals, more traits, and more tasks per socket the hardware limits the sub-linear scaling to the point where degraded performance is observed as compared to single trait processing. The main considerations are the last-level cache size, the number of individuals and the number of traits to be analysed, and we advise users to set 1 MPI task per socket and estimate how many epsilon vectors can be held in the last level cache of the CPU.

Figure S10. Convergence of GMRM software. (a) Convergence of prediction accuracy in a hold out sample of 30,000 UK Biobank individuals for human height across three independent bulk synchronous Gibbs chains with synchronisation occurring after the update of 20 markers across 60 MPI process for 2.17 million SNP markers and 428,747 individuals. (b) The root mean square error of the marker effect size estimates of randomly assigned LD independent causal variants in simulation study of 100,000 unrelated individuals from the UK Biobank for three replicates.