Temperature and nutrients drive eco-phenotypic dynamics in a microbial food web

Ze-Yi Han[1], Daniel J. Wieczynski[1], Andrea Yammine[1], Jean P. Gibert[1]

[1]Department of Biology, Duke University, Durham, North Carolina, USA

Statement of authorship: ZH and JPG conceived the study; ZH and JPG designed the experimental work. ZH collected the data with support from AY. ZH analyzed the data with support from DJW and JPG; ZH wrote the first version of the manuscript; all authors contributed substantially to subsequent versions.

Data accessibility statement: All annotated code and data are available at our dedicated repository (https://github.com/ZeYiHan/Nut_Temp_Pheno_Eco).

Running title: Consequences of thermal asymmetries

Keywords: Global warming, Food-web temperature responses, Eutrophication, Anthropogenic impacts, Communities, Trait-based Ecology

To whom correspondence should be addressed: Ze-Yi Han, Department of Biology, Duke University, Durham, NC, USA, 402-405-3888, zeyihanw@gmail.com

ABSTRACT

Anthropogenic increases in temperature and nutrient loads will likely impact food web structure

and stability. Although their independent effects have been reasonably well studied, their joint

effects—particularly on coupled ecological and phenotypic dynamics—remain poorly

understood. Here we experimentally manipulated temperature and nutrient levels in microbial

food webs and used time-series analysis to quantify the strength of reciprocal effects between

ecological and phenotypic dynamics across trophic levels. We found that i) joint –often

interactive– effects of temperature and nutrients on ecological dynamics are more common at

higher trophic levels, ii) temperature and nutrients interact to shift the relative strength of top-

down vs. bottom-up control, and iii) rapid phenotypic change mediates observed ecological

responses to changes in temperature and nutrients. Our results expose how feedbacks between

ecological and phenotypic dynamics mediate food web responses to environmental change. This

suggests important but previously unknown ways that temperature and nutrients might jointly

control the rapid eco-phenotypic feedbacks that determine food web dynamics in a changing

world.

INTRODUCTION

Understanding how rapid global climate change (GCC) will affect the structure and dynamics of communities is a pressing goal of ecology (Karl 2003; Barbour & Gibert 2021). Increasing temperatures associated with GCC influence the metabolism of individuals (Gillooly *et al.* 2002; Brown *et al.* 2004; Clarke 2006), which strengthens species interactions (Barton *et al.* 2009; O'Connor 2009; Gounand *et al.* 2016), alters community structure (Bartley *et al.* 2019; Gibert 2019; Gauzens *et al.* 2020), and affects ecosystem function (Dossena *et al.* 2012). Additionally, increasing nutrient loads from agricultural runoffs can result in eutrophication and destabilize natural communities (Rosenzweig 1971; Fussmann 2000; Hautier *et al.* 2014), often leading to species loss (Hautier *et al.* 2009). Warming and eutrophication can independently and jointly impact food web structure and stability (Malcolm *et al.* 2006; Tabi *et al.* 2019). Counterintuitively, simultaneous increases in temperature and nutrient load can produce outcomes that are qualitatively different from the combined negative effects of each variable on its own (Binzer *et al.* 2012, 2016). These non-additive (interactive) outcomes are still poorly understood, but central to honing our understanding of GCC impacts on food web structure and dynamics in a highly anthropogenized world.

The mechanisms through which warming and increasing nutrient loads independently influence food webs are relatively well understood (McClelland & Valiela 1998; Carlier *et al.* 2008; Gibert 2019), specifically regarding their impacts on the relative strength of bottom-up and top-down effects (Binzer *et al.* 2012, 2016; Shurin *et al.* 2012). For example, warming can increase predation pressure (Kratina *et al.* 2012), thus decreasing resource biomass, while increasing the proportion of top predators (Vasseur & McCann 2005; O'Connor 2009; Shurin *et al.* 2012).

Alternatively, warming can increase metabolic demands while reducing conversion efficiency

(Barneche *et al.* 2021), leading to predator starvation at high temperatures, loss of top predators

(Gibert *et al.* 2022a), and reduced food chain length (Petchey *et al.* 1999). Unlike warming,

increasing nutrient loads tend to increase bottom-up effects, resulting in unstable dynamics and

species loss (i.e., paradox of enrichment, Rip and McCann, 2011; Rosenzweig, 1971), often

leading to top-heavy, unstable food webs (Rip & McCann 2011). Eutrophication resulting from

increasing nutrient loads can also change consumer trophic position, leading to changes in

species interactions and food web structure (Lee *et al.* 2021).

While warming and nutrients can independently influence food webs, they can also have non-

additive (interactive) effects when acting together (Binzer *et al.* 2012, 2016; Tabi *et al.* 2019),

but these are much less well understood. For example, although temperature increases are

typically considered to be destabilizing (Barton *et al.* 2009), at low temperatures, small

temperature increases can cause consumer starvation, stabilizing nutrient-induced instabilities

(Binzer *et al.* 2012, 2016). However, at high temperatures, increasing nutrient loads can counter

warming-induced consumer starvation by increasing carrying capacity and predator attack rates

(Binzer *et al.* 2012). Warming also weakens nutrient-induced increases in community biomass

(negative interactive effects, Tabi et al., 2019), in turn influencing food web structure (Sentis *et*

al. 2014), species richness, and community composition (Wang *et al.* 2016a).

In addition to their effects on entire food webs, temperature and nutrients can both determine the

physiology and morphology of organisms (Rosenblatt & Schmitz 2016; Tabi *et al.* 2019). For

example, higher temperatures often result in smaller sizes (Atkinson 1995; Atkinson *et al.* 2003),

while nutrient enrichment leads to larger organisms (Irwin *et al.* 2006; Marañón *et al.* 2013).

Additionally, temperature and nutrients can interactively affect body size, which increases with

increasing nutrients at low temperature but has the opposite effect at high temperature (Tabi *et*

al. 2019). Although body size is often considered a response variable, it also has well-known

effects on population growth (Fenchel 1974; Savage *et al.* 2004; Gibert *et al.* 2022b) and species

interactions (Ferenc & Sheppard 2020), so rapid body size responses to temperature, nutrients, or

both, may have consequences for food web structure and dynamics in warmer climates (Brose *et*

al. 2012; Bernhardt *et al.* 2018; Wieczynski *et al.* 2021). However, whether body size only

responds to environmental conditions, whether rapid changes in body size mediate how

environmental conditions affect food webs, or whether shifts in body size influence body size

responses to environmental conditions in another species, are not well understood.

Here, we study how temperature and nutrients influence feedbacks between species' ecological

dynamics and rapid changes in their body sizes (phenotypic dynamics) in a tractable microbial

food web. We describe observed food web and body size responses to temperature and nutrients

across trophic levels and also study the mechanisms of these responses. Specifically, we ask: 1)

do temperature and nutrients independently or interactively influence ecological dynamics in this

microbial food web? 2) Do these effects alter the relative importance of top-down vs. bottom-up

control? 3) Do these effects vary across trophic levels? And, 4) does body size only respond to

temperature and nutrients, as suggested elsewhere (e.g., Binzer et al 2016, Tabi et al 2019), or

does that response play a role in determining how the food web itself responds to environmental

change. To address these questions, we manipulate nutrient levels and temperature in a microbial

food web composed of a complex bacterial community, a bacterivorous protist (*Tetrahymena*

pyriformis), and an omnivorous top predator (*Euplotes sp.*), then track changes in the population

densities of all organisms and the body sizes of the protists over time. We use time-series

analysis to evaluate the relative strength of top-down vs. bottom-up processes across trophic

levels as well as whether and how the observed ecological and phenotypic dynamics influence

one another across temperature and nutrient treatments. Our results reveal complex temperature

and nutrient effects on food web dynamics that vary predictably across trophic levels by altering

the relative strength of trait-mediated bottom-up and top-down effects.

METHODS

Culture care

Euplotes sp. and *Tetrahymena pyriformis* stock cultures were acquired from Carolina Biological

Supply (Burlington, NC, USA) and grown in laboratory conditions for a year prior to this

experimental work. Both species were kept in autoclaved liquid protist medium (Carolina

Biological Supply), supplemented with one autoclaved wheat seed as a carbon source (Altermatt

et al 2014). Protists were fed a mixture of pond bacteria collected from an ephemeral pond at

Duke Forest (Wilbur/Gate 9 pond, Lat 36.013914, -Long 78.979720) and composed of thousands

of bacterial species, as described elsewhere (Rocca *et al.* 2022). We maintained all cultures on a

16:8 light:dark cycle at 22°C and 65% humidity in AL-22L2 Percival growth chambers (Perry,

IA, USA).

Experimental design

Microcosms were set up in autoclaved 250mL borosilicate jars filled with 200mL of protist

media. We manipulated temperature and nutrient loads by imposing two temperature levels

(22°C/25°C) and two nutrient levels (normal protist media concentration plus one wheat seed –

i.e., high nutrients–, or half concentration plus half a wheat seed, –i.e., low nutrients) in a

factorial design with 4 treatments and 6 replicates per treatment. Day/night cycle and humidity

levels mimicked rearing conditions and were kept constant. We inoculated the bacterial

communities from 2 ml of stock culture, the same used to rear the protists. The intermediate

consumer species, *T. pyriformis*, was introduced at a starting density of 37 ind/ml, and the larger

protists species, *Euplotes sp.*, at a starting density of 0.24 ind/ml in all microcosms. We recorded

the density of both protist species through fluid imaging (FlowCam; Yokogawa Fluid Imaging

Technologies, Portland, ME, USA), Monday through Friday, for 16 days. Fluid imaging

generates individual cell raster images that were used to quantify changes in protist size

(measured as cell area, in μm^2) over time. Bacteria density was quantified as optical density at a

wavelength of 600nm (OD600), using a BioTEK Epoch-2 microplate spectrophotometer

(Winooski, VT, USA).

Statistical analysis

To test for possible effects of temperature and nutrients on ecological dynamics, we fitted

Generalized Additive Mixed Models (GAMM) to time series of species density (OD600 for

bacteria) and protist body size, across all treatments, using the 'mgcv' package v1.8-31 (Wood

2011) in R v4.0.2 (R Core Team 2020). To control for temporal autocorrelation, we used an

Autoregressive Moving Average (ARMA) correlation structure in the GAMMs using the 'nlme'

R package (version 3.1-148). To account for repeated sampling within each replicate, we

included replicates as a random intercept in the model. We compared models with additive

and/or interactive temperature and nutrients effects, as well as different ARMA correlation

structures, using AICc (Appendix I Table S1-2). We discarded *T. pyriformis* phenotypic data

from days in which fewer than 10 individuals were measured (i.e., after populations collapsed).

Characterizing ecological dynamics

To better understand which aspects of the food web dynamics were most influenced by

temperature and nutrients, we characterized multiple aspects of the observed ecological

dynamics of the bacterial community and the protists across treatments. Specifically, we

quantified: 1) initial growth rate (day^{-1}) as $[\ln(Nf) - \ln(Ni)]/\text{time}$ for early-dynamics data, (up

to day 1 for *T. pyriformis* & bacteria and day 8 for *Euplotes sp.*, due to much slower growth), 2)

maximum density (individual/ml for protists and OD_{600} for bacteria), using the highest average

daily population density across replicates but within treatments for the protists, and the top three

observed OD600 values for each replicate from each treatment for the bacteria (because their

ecological dynamics remained relatively stable), 3) the coefficient of variation (CV = standard

deviation/mean) of the temporal population dynamics within treatments (typically used as a

measure of stability (McCann 2012)), 4) the time to population collapse in days (only *T.*

pyriformis), and 5) the time to population peak in days (only *Euplotes sp.*).

Quantifying top-down/bottom-up effects and eco-phenotypic feedbacks

To understand the mechanisms through which temperature and nutrients affected the observed

food web dynamics, we quantified the reciprocal effects of ecological dynamics and body size on

each other using Convergent Cross Mapping (CCM) (Sugihara *et al.* 2012). The CCM algorithm

has now been used multiple times across ecological systems and taxa reliably to estimate the

strength of causal effects between variables for which time-series are available (Matsuzaki *et al.*

2018; Wang *et al.* 2018; Rogers *et al.* 2020; Barraquand *et al.* 2021; Doi *et al.* 2021; Abidha *et*

al. 2022; Gibert *et al.* 2022b) and we followed this specialized literature to infer causation in our

data (see Appendix II Fig. S1-15). In a nutshell, CCM quantifies the strength of causation of one

dynamical variable onto another by measuring the extent to which the time series of one variable

can reliably estimate the state of another variable (Sugihara *et al.* 2012). The larger the causal

effect of X on Y, the better the ability of Y to predict X, as Y contains information about X (by

virtue of being 'forced' by X). Meanwhile, a variable X that does not influence a variable Y

cannot be predicted from the dynamics of Y, as no information regarding X is contained in Y

(Sugihara *et al.* 2012). The CCM algorithm yields a 'cross-mapping estimation skill' in the form

of a correlation coefficient (ρ) between observed and predicted points in the time-series

(Sugihara *et al.* 2012). The larger this number, the larger the dependence of one variable on the

other. Whether there is a cause-effect relationship between two dynamical variables, as opposed

to simple correlation, further depends on whether the cross-mapping estimation skill increases

with the length of the time series used for this estimation (called the 'library size'). Whenever

such an increase is observed, a causal effect of a variable on another one is likely (i.e.

convergence; Sugihara et al., 2012).

We used a modified CCM algorithm that allows for replication in the time series through the R

package 'multispatialCCM' (Version 1.0, Clark *et al.* 2015). The package can be used to detect

causality between short replicated time series, as in our case. Our time series lacked data on days

5, 6 and 12, 13 across all replicates. However, the CCM algorithm does not allow for missing

time points. To resolve this issue we interpolated the time series data for each replicate using

three methods: linear interpolation, spline interpolation, and smooth spline interpolation using

the 'approx', 'spline', and 'smooth.spline' functions in base R (see Appendix 2). To increase the

robustness of our inference, we then performed CCM in each of these time series and averaged

the estimation skill from our CCM analysis across these three separate interpolated time-series.

That said, CCM results based on each independent smoothing technique held qualitatively

(Appendix II Fig. 1-12), which corroborates the robustness of our inference. Additionally, we

only used CCM results that showed convergence in cross mapping skill (ρ) with increasing

library size (indicating causality) to focus only on likely causal effects between species

ecological and phenotypic dynamics (Appendix II Fig. 13-15).

Based on previous literature (Sugihara *et al.* 2012; Matsuzaki *et al.* 2018; Barraquand *et al.*

2021), we interpreted the cross-mapping skill (ρ) as the magnitude of the effect of one variable

on another, whenever convergence was present. We calculated this cross-mapping skill between

all predator and prey densities, between densities and trait dynamics (except the effect of protist

densities on bacterial traits as we lack phenotypic data for the bacteria community), and between

protist trait dynamics. The effect of prey density on predator density was thus considered as

representing bottom-up control, and the effect of predator density on prey density as top-down

control (dubbed "eco-eco" effects, for simplicity). The effects of change in body size on

ecological dynamics (density) where dubbed 'pheno-eco' effects, and the effects of density on

body size dynamics as 'eco-pheno' effects. Reciprocal effects of changes in predator and prey

protist body sizes were dubbed 'pheno-pheno' effects.

RESULTS

General Ecological and Phenotypic dynamics

Overall, bacterial density rapidly increased to carrying capacity (Fig. 1a, light grey), *T.*

pyriformis increased rapidly, then decreased (Fig. 1a, dark grey), while *Euplotes sp.* increased

almost monotonically to carrying capacity (Fig. 1a, black), ultimately resulting in a one protist +

bacterial community state. Temperature and nutrients affected all three species and led to

significantly different dynamics across treatments (Fig. 1b-d, Appendix I Table S2). The body

sizes of both protists changed rapidly over time (Fig. 1e, f) and responded to both temperature

and nutrients (Fig. 1e, f, Appendix I Table S2).

Temperature and nutrient effects changed across trophic levels in systematic ways

Temperature significantly affected a larger number of dynamical descriptors than nutrients (Fig.

2, 3). Individual temperature effects were more frequent in the upper trophic levels than in the

bacterial community (Fig. 3), as were interactions between temperature and nutrients (Fig. 3). In

particular, higher temperature alone i) had no significant effect on the intrinsic growth rate of the

bacteria (Fig a, Temperature effect (T from now on) = -0.003, $p = 0.28$), ii) strongly increased

initial growth rate in *T. pyriformis* (T = 0.50 $p = 1.16 \times 10^{-6}$, Fig. 2b), iii) decreased the maximum

density of bacteria and *T. pyriformis* (T_B = -0.005, $p = 0.004$; T_T = -975.3, $p < 0.02$; Fig. 2d-e)

but increased that of *Euplotes sp.* (T_E = 39.83, $p < 0.03$, Fig. 2f), iv) increased the variability in

T. pyriformis density (i.e. CV, T = 0.2, $p < 0.006$, Fig. 2h) but decreased that of *Euplotes sp.* (T =

-0.08, $p < 0.02$, Fig. 2i), v) accelerated the time to collapse in *T. pyriformis* (T = -2.8, $p = 1.6 \times 10^{-}$

^{5}; Fig. 2j), and vi) accelerated the time to peak density in *Euplotes sp.* (T = -2.2, $p < 0.02$; Fig.

2k).

Higher nutrients alone i) decreased the initial growth rate of *Euplotes sp.* (Nutrient effect (N

from now on) = -0.09, p = 5×10⁻⁴ Fig. 2c), ii) increased maximum density of all species (N_B=

0.01, p = 2×10⁻¹⁰; N_T = 1002.2, p = 0.01; N_E = 60.67, p < 0.002, Fig. 2d-f), and iii) increased the

CV of bacteria and *Euplotes sp.* (N_B = 0.2, p = 8.6×10⁻⁵, Fig. 2g; N_E = 0.16, p = 2.9×10⁻⁵, Fig. 2i)

while decreasing that of *T. pyriformis* (N_T = -0.19, p < 0.01, Fig. 2h). Interestingly, increasing

either temperature or nutrients had opposite effects on the CVs of *T. pyriformis* and *Euplotes sp.*,

indicating antagonistic additive effects of temperature and nutrients on food web stability.

Temperature and nutrients had complex interactive effects on ecological dynamics of both protist

consumers. Nutrients had a negative effect on the initial growth rate of *Euplotes sp.* at low

temperature, but a positive effect at high temperature (N×T = 0.13, p < 10^-3; Fig. 2c). The

maximum density of *T. pyriformis* increased with nutrients, but this effect was stronger at high

temperature (N×T = 1526.7, p < 0.01, Fig. 2e). Nutrients also had i) a negative effect on *T.*

pyriformis population variability (CV) at low temperature but no effect at high temperature (N×T

= 0.2, p = 0.045, Fig. 2h) and ii) a positive effect on *Euplotes sp.* CV that was stronger at low

temperature (N×T = -0.12, p < 0.01, Fig. 2i). Additional model stats can be found in Appendix I

Table S3. Taken together, these results imply that temperature and nutrients individually and

interactively influence species ecological dynamics at all trophic levels, with more of both types

of effects occurring at higher trophic levels (Fig. 3).

Temperature and nutrients influence top-down and bottom-up effects

Results from CCM analyses indicated that the relative strength of bottom-up and top-down

effects changed across treatments (Fig. 4a). Top-down and bottom-up effects between the two

protists were slightly stronger in the warmer temperature (Fig. 4a, yellow lines). Importantly,

temperature and nutrients showed interactive effects on both top-down and bottom-up effects

(Fig. 4a). For example, while the top-down effects of protists on bacteria (Fig.4a, green and

purple dashed lines) were generally stronger than the bottom-up effects of the bacteria on the

protists (Fig. 4a green and purple solid lines), temperature and nutrients interactively –but

differentially– shifted the magnitude of the top-down effects of the two protist predators on

bacteria (Fig.4a, open dots and dashed lines). Indeed, the top-down effect of *T. pyriformis* on

bacteria decreased in the high-nutrient treatment, but higher temperature strengthened this effect

(Fig. 4a purple dashed line). Meanwhile, high nutrients increased the top-down effect of *Euplotes*

sp. on bacteria in the lower temperature but decreased it at higher temperature (Fig. 4a green

dashed line).

The strength of species interactions in the food web, overall, also changed according to shifts in

the strength of top-down and bottom-up control between species pairs (Figure 5a). To visualize

these changes, we present CCM cross-mapping skill value at the largest library size as links

between species pairs (Fig. 5a). At low nutrient levels, higher temperature led to stronger species

interactions (Fig. 5a, bottom left and right) while at high nutrient levels, interaction strengths

were generally stronger and temperature only had a small effect (Fig. 5a, top left and right). Low

nutrients and high temperatures led to the strongest interactions among all three species (Fig. 5a,

bottom right).

Temperature and nutrients altered the reciprocal effects of ecological and phenotypic dynamics

Protist size not only responded to temperature and nutrients, but also played an important role in

determining the effects of temperature and nutrients on ecological dynamics (Fig. 4b-c). The

bidirectional effects between the body-size dynamics of *Euplotes sp.* and the ecological

dynamics of *T. pyriformis* were the strongest overall across treatments (Fig. 4b-c). The body size

dynamics of the omnivorous predator, *Euplotes sp.*, but not those of *T. pyriformis,* had relatively

strong causal effects on the ecological dynamics of all species, including its own (Fig. 4c).

Specifically, at low temperature, increasing nutrient level strengthened the effect of *Euplotes sp.*

phenotypic dynamics on its own ecological dynamics and those of *T. pyriformis*, but weakened

these effects on the ecological dynamics of bacteria (Fig. 4c). Yet, at the warmer temperature,

increasing nutrient levels weakened the effects of *Euplotes sp.* phenotypic dynamics on the

ecological dynamics of all species, including itself (Fig. 4c), indicating that plastic changes in

top predator body size can mediate how food web dynamics respond to temperature and

nutrients.

Trait on trait effects among the two predators (Fig. 4d, pheno-pheno) were weaker than their

pheno-eco or eco-pheno counterparts. Interestingly, pheno-pheno causal effects were only

observed at the higher temperature (Fig. 4d). Increasing nutrient levels increased the pheno-

pheno effects of *T. pyriformis* on *Euplotes sp.* but decreased those of *Euplotes sp.* on *T.*

pyriformis (Fig. 4d).

Overall, we also found stronger effects between ecological and phenotypic dynamics at higher

temperatures, especially in the low nutrient treatment (Fig. 5b, bottom right), consistent with

results from the top-down vs bottom-up effects (Fig 5a). Overall, changes in body size seem to

mediate the effect of temperature and nutrients on ecological dynamics (Fig. 5b), and these

effects seem stronger at higher temperatures but low nutrient levels.

DISCUSSION

Our results reveal complex but systematic effects of temperature and nutrients on the ecological

and phenotypic dynamics of a microbial food web. We show that temperature and nutrients can

each independently influence different aspects of food web dynamics (Fig. 1, 2), while their joint

effects get increasingly complex at higher trophic levels (Fig. 2, 3). We also found that changes

in the relative strength of top-down and bottom-up effects likely drive observed responses to

temperature and nutrients (Fig. 4, 5) and that rapid changes in body size mediate these effects

(Fig. 4, 5). Stronger species interactions at low nutrient levels and high temperature coupled with

phenotypic dynamics having more and stronger effects in both ecological and phenotypic change

at low nutrients and high temperature, suggest that phenotypic change may mediate the

temperature response of ecological dynamics, perhaps strengthening species interactions (Fig 4,

5). These results therefore suggest that body size not only responds to shifts in environmental

conditions, but also plays a role in determining ecological responses to such shifts (Fig. 5).

Evaluating feedbacks between ecological and phenotypic dynamics may therefore be integral to

understanding food web responses to environmental change.

Increasingly complex effects of temperature and nutrients at higher trophic levels

Our results reveal the pervasive effects of temperature and nutrients across all species, but also

how these effects are more numerous and increasingly complex (temperature-nutrient

interactions) at higher trophic levels (Fig. 2, 3). The effects of temperature and nutrients on

dietary preferences, species interactions, and foraging behavior may help explain why these

interactive effects may be more likely among consumer species (Fig. 1d; Fig. 2c).

Because energy enters at the bottom of a food web, basal species may be more strongly

influenced by direct effects of nutrients while species at higher trophic levels may be more

strongly affected by a combination of temperature and nutrients (Petchey *et al.* 1999; Voigt *et al.*

2003). 6/21/22 11:24:00 AM Some of these effects may be explained through the transitivity of

causality (Sugihara *et al.* 2012) as *Euplotes sp.* preys on *T. pyriformis* which in turn preys on the

bacterial community. So *Euplotes sp.* will be causally, albeit indirectly, linked to bacteria, and

also to the effects that temperature and nutrients have on bacteria. In ecological terms, as energy

flows from the basal species to the consumers, the temperature and nutrient effects that act on

basal species could be integrated by consumers across trophic levels, leading to additional

indirect temperature and nutrient effects on consumers at the top of the food web. Moreover,

these effects may be exaggerated in omnivores due to the complex nature of their trophic

interactions. As temperature and nutrients change species interactions and foraging behaviors

(Sentis *et al.* 2014), omnivorous consumers may shift diets between basal and intermediate

resources (Sentis *et al.* 2014), altering the strength/prevalence of competitive vs. predatory

interactions (Vandermeer 2006) and leading to climate-driven food web rewiring (Bartley *et al.*

2019; Barbour & Gibert 2021). Whether the joint effects of temperature and nutrients become

increasingly complex at higher trophic levels in food webs with a higher prevalence of

omnivory, however, is not known, but a promising avenue for future research.

Top-down control, bottom-up control, and food web stability

Consistent with our results (Fig. 2d), theory predicts that higher nutrient loads should increase

energy flux in food webs, through increases in basal species density (Rip & McCann 2011;

Shurin *et al.* 2012). Theory also predicts that increasing energy flux can be destabilizing, leading

to increasing oscillations in density (Rosenzweig 1971; Rip & McCann 2011), but increasing

temperatures should stabilize oscillations by weakening top-down effects (Vasseur & McCann

2005; Binzer *et al.* 2016; Tabi *et al.* 2019). We found that the interactive effects of temperature

and nutrients have divergent impacts on stability—with temperature stabilizing, and nutrients

destabilizing *Euplotes sp.* densities, but the opposite being true for *T. pyriformis* (Fig. 2h-i). We

also observed that top-down control on bacteria by both consumers was much stronger than

bottom-up effects across treatments (Fig. 4a). Temperature and nutrients had different but

interactive effects on top-down controls by the two predators even though bottom-up effects

remained unaffected by the treatments (Fig. 4a). These results thus indicate that changes in the

strength of top-down effects between basal resources and consumers—instead of bottom-up

effects—could be the dominant mechanism through which higher temperature may stabilize

instabilities caused by nutrients (Binzer *et al.* 2012, 2016).

Phenotypes mediate temperature and nutrient effects on food web dynamics

Our results indicate that phenotypic dynamics play a larger role in mediating environmental

impacts on food web dynamics than previously thought (Fig. 4, 5). Previous studies showed that

predator-prey body size ratios significantly influence temperature and nutrients effects (Binzer *et*

al. 2012; Gibert & DeLong 2014) and that body size responds to changes in nutrients,

temperature, and ecological dynamics in specific ways (Tabi *et al.* 2019). Our results support—

but also extend—those results by showing that 1) phenotypic effects on ecological dynamics can

be strong, especially among top predators, 2) changes in the prey ecological dynamics may

strongly drive changes in predator size, and 3) interactions involving phenotypic dynamics (eco-

pheno, pheno-eco, or pheno-pheno) vary across environmental conditions (Fig. 4b-d, Fig. 5).

Caveats

A caveat of this study is that, despite uniquely detailed, long, and well-replicated time-series, our

time-series have a few small gaps due to lack of sampling on weekends. To address this issue,

we interpolated these missing datapoints using three different methods, but, despite being

generally robust, some variation remained between these results and the CCM inference of

species interactions (Appendix II Figure 1-12). All three methods showed that bottom-up and

top-down effects between the protist predators were the strongest and remained strong across all

treatments. Moreover, top-down controls from both protists predators to bacteria were always

stronger than bottom-up effects (Appendix II Figure 13-15 Eco-Eco panels). Last, the effects of

Euplotes sp. body size on its own ecological dynamics and that of the *T. pyriformis* and bacteria

were consistent across interpolation methods (Appendix II Figure 13-15 Eco-Pheno, Pheno-Eco

and Pheno-Pheno panels). As CCM and related methods grow in use (Wang *et al.* 2016b;

Matsuzaki *et al.* 2018; Barraquand *et al.* 2021; Doi *et al.* 2021; Kitayama *et al.* 2021), there is a

real need for these tools to be robust even in the face of imperfect data, including missing

timepoints. Our results therefore underline the needs to better understand how missing data may

affect CCM and other time-series analyses and how to best predict missing data for analysis.

Concluding remarks

Rapid phenotypic change has been suggested as a main driver of food web rewiring in future

climates (Barbour & Gibert 2021), but little experimental evidence exists. Here, our results show

that strong feedbacks between ecological and phenotypic dynamics depend on environmental

conditions (temperature and nutrients), suggesting that rapid phenotypic change influences food

web responses to environmental change. Moreover, the joint effects of temperature and nutrients

do not equally affect all members of the community, as higher trophic levels are more likely to

experience both independent and joint effects. Together, our results emphasize the need to

incorporate phenotypic dynamics in future studies of food web responses to warming and

eutrophication in a changing world and show how shifts in distinct environmental stressors can

have complex but systematic effects on food web dynamics.

432 ACKNOWLEDGMENTS

433 This work was supported by a U.S. Department of Energy, Office of Science, Office of

434 Biological and Environmental Research, Genomic Science Program Grant under Award Number

435 DE-SC0020362 to JPG.

Figures

Figure 1. Ecological and phenotypic dynamics within experimental microcosms. (a) Average

population densities across treatments. (b-d) Density dynamics of bacteria, *T. pyriformis*, and

Euplotes sp. (e-f) Changes in the body size (measured as the area of the cell in microm^2) for

both protists over time. *T. pyriformis* were effectively extinct by day 9. In (b–f), dots are the

empirical measurements and lines are GAMM fits. The greyed area in (e) indicates days for

which data were insufficient to estimate trait distributions.

Figure 2. Additive and interactive effects of temperature and nutrients on descriptors of

population dynamics. Open circles and triangles represent data points from each replicate. Solid

shapes represent means across 6 replicates. Grey letters T, N, and (T×N) represent statistically

significant effects from temperature, nutrients, and interactive effects between temperature and

nutrients, respectively.

Fig. 3. Significant temperature and nutrient effects on ecological dynamics (i.e., initial growth,

maximum abundance, and density CV). T, N, and T×N denote effects of temperature, nutrients,

and interactive effects of temperature and nutrients, respectively.

Figure 4. Cross-mapping skill (ρ) of species ecological dynamics across treatments for each

species pair. (a) Shows CCM estimation skill of one species' effect on another. Solid circles and

solid lines represent bottom-up effects and open circles and dashed lines represent top-down

effects. (b) Shows all causal species interactions for all species considered. Line widths are

proportional to the magnitude of the CCM skill.

Figure 5. Interactions between phenotypic (Ph) and ecological (E) dynamics. (a-c) Show CCM

estimation skill of density on traits (eco-pheno), traits on density (pheno-eco), and trait on trait

(pheno-pheno). Circles represent CCM estimation skill of one species' effect on another species,

473 while triangles represent intraspecific effects. Dashed and solid lines represent top-down and

474 bottom-up effects respectively. Dotted lines represent intraspecific effects. In (d), line widths are

475 proportional to the CCM skill. Color code as in Fig. 4a.

References:

Abidha, C.A., Amoako, Y.A., Nyamekye, R.K., Bedu-Addo, G., Grziwotz, F., Mockenhaupt, F.P., *et al.* (2022). Fasting blood glucose in a Ghanaian adult is causally affected by malaria parasite load: a mechanistic case study using convergent cross mapping. *Malar. J.*, 21, 93.

Atkinson, D. (1995). Effects of temperature on the size of aquatic ectotherms: Exceptions to the general rule. *J. Therm. Biol.*, 20, 61–74.

Atkinson, D., Ciotti, B.J. & Montagnes, D.J.S. (2003). Protists decrease in size linearly with temperature: *ca* . 2.5% °C $^{-1}$. *Proc. R. Soc. Lond. B Biol. Sci.*, 270, 2605–2611.

Barbour, M.A. & Gibert, J.P. (2021). Genetic and plastic rewiring of food webs under climate change. *J. Anim. Ecol.*, 1365-2656.13541.

Barneche, D.R., Hulatt, C.J., Dossena, M., Padfield, D., Woodward, G., Trimmer, M., *et al.* (2021). Warming impairs trophic transfer efficiency in a long-term field experiment. *Nature*.

Barraquand, F., Picoche, C., Detto, M. & Hartig, F. (2021). Inferring species interactions using Granger causality and convergent cross mapping. *Theor. Ecol.*, 14, 87–105.

Bartley, T.J., McCann, K.S., Bieg, C., Cazelles, K., Granados, M., Guzzo, M.M., *et al.* (2019). Food web rewiring in a changing world. *Nat. Ecol. Evol.*, 3, 345–354.

Barton, B.T., Beckerman, A.P. & Schmitz, O.J. (2009). Climate warming strengthens indirect interactions in an old-field food web. *Ecology*, 90, 2346–2351.

Bernhardt, J.R., Sunday, J.M. & O'Connor, M.I. (2018). Metabolic Theory and the Temperature-Size Rule Explain the Temperature Dependence of Population Carrying Capacity. *Am. Nat.*, 192, 687–697.

Binzer, A., Guill, C., Brose, U. & Rall, B.C. (2012). The dynamics of food chains under climate change and nutrient enrichment. *Philos. Trans. R. Soc. B Biol. Sci.*, 367, 2935–2944.

Binzer, A., Guill, C., Rall, B.C. & Brose, U. (2016). Interactive effects of warming, eutrophication and size structure: impacts on biodiversity and food-web structure. *Glob. Change Biol.*, 22, 220–227.

Brose, U., Dunne, J.A., Montoya, J.M., Petchey, O.L., Schneider, F.D. & Jacob, U. (2012). Climate change in size-structured ecosystems. *Philos. Trans. R. Soc. B Biol. Sci.*, 367, 2903–2912.

Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004). Toward a metabolic theory of ecology. *Ecology*, 85, 1771–1789.

Carlier, A., Riera, P., Amouroux, J.-M., Bodiou, J.-Y., Desmalades, M. & Grémare, A. (2008). Food web structure of two Mediterranean lagoons under varying degree of eutrophication. *J. Sea Res.*, 60, 264–275.

Clark, A.T., Ye, H., Isbell, F., Deyle, E.R., Cowles, J., Tilman, G.D., *et al.* (2015). Spatial convergent cross mapping to detect causal relationships from short time series. *Ecology*, 96, 1174–1181.

Clarke, A. (2006). Temperature and the metabolic theory of ecology. *Funct. Ecol.*, 20, 405–412.

Doi, H., Yasuhara, M. & Ushio, M. (2021). Causal analysis of the temperature impact on deep-sea biodiversity. *Biol. Lett.*, 17, 20200666.

Dossena, M., Yvon-Durocher, G., Grey, J., Montoya, J.M., Perkins, D.M., Trimmer, M., *et al.* (2012). Warming alters community size structure and ecosystem functioning. *Proc. R. Soc. B Biol. Sci.*, 279, 3011–3019.

Fenchel, T. (1974). Intrinsic rate of natural increase: The relationship with body size. *Oecologia*,
14, 317–326.
Ferenc, V. & Sheppard, C.S. (2020). The stronger, the better – trait hierarchy is driving alien
species interaction. *Oikos*, 129, 1455–1467.
Fussmann, G.F. (2000). Crossing the Hopf Bifurcation in a Live Predator-Prey System. *Science*,
290, 1358–1360.
Gauzens, B., Rall, B.C., Mendonça, V., Vinagre, C. & Brose, U. (2020). Biodiversity of
intertidal food webs in response to warming across latitudes. *Nat. Clim. Change*, 10, 264–
269.
Gibert, J.P. (2019). Temperature directly and indirectly influences food web structure. *Sci. Rep.*,
9, 5312.
Gibert, J.P. & DeLong, J.P. (2014). Temperature alters food web body-size structure. *Biol. Lett.*,
10, 20140473.
Gibert, J.P., Grady, J.M. & Dell, A.I. (2022a). Food web consequences of thermal asymmetries.
Funct. Ecol., 1365-2435.14091.
Gibert, J.P., Han, Z., Wieczynski, D.J., Votzke, S. & Yammine, A. (2022b). Feedbacks between
size and density determine rapid eco-phenotypic dynamics. *Funct. Ecol.*, 1365-
2435.14070.
Gillooly, James.F., Charnov, E.L., West, G.B., Savage, V.M. & Brown, J.H. (2002). Effects of
size and temperature on developmental time. *Nature*, 417, 70–73.
Gounand, I., Kéfi, S., Mouquet, N. & Gravel, D. (2016). Trait selection during food web
assembly: the roles of interactions and temperature. *Theor. Ecol.*, 9, 417–429.
Hautier, Y., Niklaus, P.A. & Hector, A. (2009). Competition for Light Causes Plant Biodiversity
Loss After Eutrophication. *Science*, 324, 636–638.
Hautier, Y., Seabloom, E.W., Borer, E.T., Adler, P.B., Harpole, W.S., Hillebrand, H., *et al.*
(2014). Eutrophication weakens stabilizing effects of diversity in natural grasslands.
Nature, 508, 521–525.
Irwin, A.J., Finkel, Z.V., Schofield, O.M.E. & Falkowski, P.G. (2006). Scaling-up from nutrient
physiology to the size-structure of phytoplankton communities. *J. Plankton Res.*, 28,
459–471.
Karl, T.R. (2003). Modern Global Climate Change. *Science*, 302, 1719–1723.
Kitayama, K., Ushio, M. & Aiba, S. (2021). Temperature is a dominant driver of distinct annual
seasonality of leaf litter production of equatorial tropical rain forests. *J. Ecol.*, 109, 727–
736.
Kratina, P., Greig, H.S., Thompson, P.L., Carvalho-Pereira, T.S.A. & Shurin, J.B. (2012).
Warming modifies trophic cascades and eutrophication in experimental freshwater
communities. *Ecology*, 93, 1421–1430.
Lee, G.H., Vonk, J.A., Verdonschot, R.C.M., Kraak, M.H.S., Verdonschot, P.F.M. & Huisman,
J. (2021). Eutrophication induces shifts in the trophic position of invertebrates in aquatic
food webs. *Ecology*, 102.
Malcolm, J.R., Liu, C., Neilson, R.P., Hansen, L. & Hannah, L. (2006). Global Warming and
Extinctions of Endemic Species from Biodiversity Hotspots. *Conserv. Biol.*, 20, 538–548.
Marañón, E., Cermeño, P., López-Sandoval, D.C., Rodríguez-Ramos, T., Sobrino, C., Huete-
Ortega, M., *et al.* (2013). Unimodal size scaling of phytoplankton growth and the size
dependence of nutrient uptake and use. *Ecol. Lett.*, 16, 371–379.

Matsuzaki, S.S., Suzuki, K., Kadoya, T., Nakagawa, M. & Takamura, N. (2018). Bottom-up linkages between primary production, zooplankton, and fish in a shallow, hypereutrophic lake. *Ecology*, 99, 2025–2036.

McCann, K.S. (2012). *Food webs*. Monographs in population biology. Princeton University Press, Princeton, N.J.

McClelland, J. & Valiela, I. (1998). Changes in food web structure under the influence of increased anthropogenic nitrogen inputs to estuaries. *Mar. Ecol. Prog. Ser.*, 168, 259–271.

O'Connor, M.I. (2009). Warming strengthens an herbivore–plant interaction. *Ecology*, 90, 388–398.

Petchey, O.L., McPhearson, P.T., Casey, T.M. & Morin, P.J. (1999). Environmental warming alters food-web structure and ecosystem function. *Nature*, 402, 69–72.

Rip, J.M.K. & McCann, K.S. (2011). Cross-ecosystem differences in stability and the principle of energy flux: Cross-ecosystem differences in stability. *Ecol. Lett.*, 14, 733–740.

Rocca, J.D., Yammine, A., Simonin, M. & Gibert, J.P. (2022). Protist Predation Influences the Temperature Response of Bacterial Communities. *Front. Microbiol.*, 13, 847964.

Rogers, T.L., Munch, S.B., Stewart, S.D., Palkovacs, E.P., Giron-Nava, A., Matsuzaki, S.S., *et al.* (2020). Trophic control changes with season and nutrient loading in lakes. *Ecol. Lett.*, 23, 1287–1297.

Rosenblatt, A.E. & Schmitz, O.J. (2016). Climate Change, Nutrition, and Bottom-Up and Top-Down Food Web Processes. *Trends Ecol. Evol.*, 31, 965–975.

Rosenzweig, M.L. (1971). Paradox of Enrichment: Destabilization of Exploitation Ecosystems in Ecological Time. *Science*, 171, 385.

Savage, V.M., Gillooly, J.F., Brown, J.H., West, G.B. & Charnov, E.L. (2004). Effects of Body Size and Temperature on Population Growth. *Am. Nat.*, 163, 429–441.

Sentis, A., Hemptinne, J.-L. & Brodeur, J. (2014). Towards a mechanistic understanding of temperature and enrichment effects on species interaction strength, omnivory and food-web structure. *Ecol. Lett.*, 17, 785–793.

Shurin, J.B., Clasen, J.L., Greig, H.S., Kratina, P. & Thompson, P.L. (2012). Warming shifts top-down and bottom-up control of pond food web structure and function. *Philos. Trans. R. Soc. B Biol. Sci.*, 367, 3008–3017.

Sugihara, G., May, R., Ye, H., Hsieh, C. -h., Deyle, E., Fogarty, M., *et al.* (2012). Detecting Causality in Complex Ecosystems. *Science*, 338, 496–500.

Tabi, A., Petchey, O.L. & Pennekamp, F. (2019). Warming reduces the effects of enrichment on stability and functioning across levels of organisation in an aquatic microbial ecosystem. *Ecol. Lett.*, 22, 1061–1071.

Vandermeer, J. (2006). Omnivory and the stability of food webs. *J. Theor. Biol.*, 238, 497–504.

Vasseur, D.A. & McCann, K.S. (2005). A Mechanistic Approach for Modeling Temperature-Dependent Consumer-Resource Dynamics. *Am. Nat.*, 166, 184–198.

Voigt, W., Perner, J., Davis, A.J., Eggers, T., Schumacher, J., Bährmann, R., *et al.* (2003). Trophic levels are differentially sensitive to climate. *Ecology*, 84, 2444–2453.

Wang, J., Pan, F., Soininen, J., Heino, J. & Shen, J. (2016a). Nutrient enrichment modifies temperature-biodiversity relationships in large-scale field experiments. *Nat. Commun.*, 7, 13960.

Wang, J., Pan, F., Soininen, J., Heino, J. & Shen, J. (2016b). Nutrient enrichment modifies temperature-biodiversity relationships in large-scale field experiments. *Nat. Commun.*, 7, 13960.

Wang, Y., Yang, J., Chen, Y., De Maeyer, P., Li, Z. & Duan, W. (2018). Detecting the Causal Effect of Soil Moisture on Precipitation Using Convergent Cross Mapping. *Sci. Rep.*, 8, 12171.

Wieczynski, D., Singla, P., Doan, A., Singleton, A., Han, Z., Yammine, A., *et al.* (2021). *Simple traits predict complex temperature responses across scales* (preprint). In Review.

Wood, S.N. (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models: Estimation of Semiparametric Generalized Linear Models. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 73, 3–36.