Temperature and nutrients drive complex eco-phenotypic dynamics in a microbial
food web

Ze-Yi Han[1], Daniel J. Wieczynski[1], Andrea Yammine[1], Jean P. Gibert[1]
[1]Department of Biology, Duke University, Durham, North Carolina, USA

Statement of authorship: ZH and JPG conceived the study; ZH and JPG designed the
experimental work. ZH collected the data with support from AY. ZH analyzed the data with
support from DJW and JPG; ZH wrote the first version of the manuscript; all authors contributed
substantially to subsequent versions.

Data accessibility statement: All annotated code and data is available at our dedicated repository
(https://github.com/ZeYiHan/Nut_Temp_Pheno_Eco).

Running title: Consequences of thermal asymmetries

Keywords: Global warming, Food-web temperature responses, Eutrophication, Anthropogenic
impacts, Communities, Trait-based Ecology

Type: Letter

Number of words: abstract (150 words), main text (3983 words), boxes (0 words)

Number of figures: 5

Number of references: 48

Number of tables: 0

Number of boxes: 0

To whom correspondence should be addressed: Ze-Yi Han, Department of Biology, Duke
University, Durham, NC, USA, 402-405-3888, zeyihanw@gmail.com

ABSTRACT

Anthropogenic increases in temperature and nutrient loads will likely impact food web structure

and stability. Though their independent effects have been well studied, their joint effects—

particularly on coupled ecological and phenotypic dynamics—remain poorly understood. Here

we experimentally manipulated temperature and nutrient levels in microbial food webs and used

time-series analysis to quantify the strength and causality of reciprocal effects between

ecological and phenotypic dynamics across species. We found that i) temperature and nutrients

have more complex effects on ecological dynamics at higher trophic levels, ii) temperature and

nutrients interact antagonistically to shift the relative strength of top-down vs. bottom-up control,

iii) phenotypic dynamics have stronger impacts on ecological dynamics than vice versa

(especially at higher temperature), and iv) rapid phenotypic change mediates observed ecological

responses to changes in temperature and nutrients. Our results expose how feedbacks between

ecological and phenotypic dynamics mediate food web responses to environmental change.

INTRODUCTION

Understanding how rapid global climate change (GCC) will affect the structure and dynamics of

communities is a pressing goal of ecology (Karl 2003). Increasing temperatures associated with

GCC influence the metabolism of individuals (Clarke 2006), which strengthens species

interactions (Barton *et al.* 2009; O'Connor 2009; Gounand *et al.* 2016), alters community

structure (Bartley *et al.* 2019; Gibert 2019; Gauzens *et al.* 2020), and ecosystem function

(Dossena *et al.* 2012). Additionally, increasing nutrient loads from agricultural runoffs can result

in eutrophication and destabilize natural communities (Rosenzweig 1971; Fussmann 2000;

Hautier *et al.* 2014), often leading to species loss (Hautier *et al.* 2009). Warming and

eutrophication may therefore independently and jointly impact food web structure and stability

(Malcolm *et al.* 2006; Tabi *et al.* 2019). Counterintuitively, however, simultaneous increases in

temperature and nutrient load can produce outcomes that are qualitatively different from the

combined negative effects of each variable on its own (Binzer *et al.* 2012, 2016). These non-

additive (interactive) outcomes are still poorly understood, but central to honing our predictions

about GCC impacts on food web structure and dynamics in a highly anthropogenized world.

The mechanisms through which warming and increasing nutrient loads independently influence

food webs are relatively well understood (McClelland & Valiela 1998; Carlier *et al.* 2008; Gibert

2019), specifically regarding their impacts on the relative strength of bottom-up and top-down

effects (Binzer *et al.* 2012, 2016; Shurin *et al.* 2012). For example, warming can increase

predation pressure (Kratina *et al.* 2012), thus decreasing resource biomass, while increasing the

proportion of top predators (Vasseur & McCann 2005; O'Connor 2009; Shurin *et al.* 2012).

Alternatively, warming can increase metabolic demands while reducing conversion efficiency

(Barneche *et al.* 2021), leading to predator starvation at high temperatures, loss of top predators,

and reduced food chain length (Petchey *et al.* 1999)Unlike warming, increasing nutrient loads

tend to increase bottom-up effects, often resulting in unstable dynamics and species loss (i.e.,

paradox of enrichment, Rip and McCann, 2011; Rosenzweig, 1971). Increasing nutrient loads

can alter consumer-resources interactions, sometimes resulting in top-heavy, unstable food webs

(Rip & McCann 2011). Eutrophication resulting from increasing nutrient loads can also change

consumer trophic position, leading to changes in species interactions and food web structure (Lee

et al. 2021).

While warming and nutrients can independently influence food webs, they can also have non-

additive (interactive) effects when acting together (Binzer *et al.* 2012, 2016; Tabi *et al.* 2019).

For example, at low temperatures, small temperature increases can stabilize nutrient-induced

instabilities (Binzer *et al.* 2012, 2016), although temperature increases are themselves considered

destabilizing because they strengthen consumer-resource interactions (Barton *et al.* 2009). At

high temperatures, increasing nutrient loads can counteract warming-induced consumer

starvation by increasing carrying capacity and predator attack rates (Binzer *et al.* 2012).

Warming also weakens nutrient-induced increases in community biomass (negative interactive

effects, Tabi et al., 2019), in turn influencing food web structure (Sentis *et al.* 2014), species

richness, and community composition (Wang *et al.* 2016).

Temperature and nutrients can both determine the physiological and morphological traits of

organisms (Rosenblatt & Schmitz 2016; Tabi *et al.* 2019). For example, higher temperatures

often result in smaller sizes (Atkinson 1995; Atkinson *et al.* 2003), while nutrient enrichment

leads to larger organisms (Irwin *et al.* 2006; Marañón *et al.* 2013). Additionally, temperature and

nutrients can interactively affect body size, where body size may increase with nutrients at low

temperature but decrease at high temperature (Tabi *et al.* 2019). Although body size is often

considered a response variable, its strong influences on population growth (Fenchel 1974;

Savage *et al.* 2004) and species interactions (Ferenc & Sheppard 2020) mean that rapid

phenotypic responses to temperature, nutrients, or both, may have consequences for food web

structure and dynamics in warmer climates (Brose *et al.* 2012; Bernhardt *et al.* 2018; Wieczynski

et al. 2021). However, whether body size only responds to environmental conditions, whether

rapid changes in body size influences how environmental conditions affect food webs, or

whether shifts in body size in a species can influence body size responses to environmental

conditions in the species they interact with, are not well understood.

Here, we study how temperature and nutrients influence the feedbacks between ecological and

body size dynamics in a tractable microbial food web by first describing observed food web and

body size responses to temperature and nutrients across trophic levels, then studying the

mechanisms of response. Specifically, we ask: 1) do temperature and nutrients independently or

interactively influence ecological dynamics within microbial food webs?, 2) do these effects alter

the relative importance of top-down vs. bottom-up control?, 3) do these effects vary across

trophic levels?, and 4) is body size only responding to temperature and nutrients, as suggested

elsewhere (e.g., Binzer et al 2016, Tabi et al 2019), or does that response play a role in

determining how the food web itself responds to environmental change. To address these

questions, we manipulate nutrient inputs and temperature in a microbial food web composed of

bacteria, a bacterivorous protist (*Tetrahymena pyriformis*), and an omnivorous top predator

(*Euplotes sp.*), then track the changes in population densities and body sizes of over time. We

use state-of-the-art time-series analysis to evaluate the relative strength of top-down vs. bottom-

up processes across trophic levels, as well as whether and how the observed ecological and

phenotypic dynamics influence one another. Our results reveal complex temperature and nutrient

effects on food web dynamics that vary predictably across trophic levels by altering the relative

strength of trait-mediated bottom-up and top-down effects.

METHODS

Culture care

Euplotes sp. and *Tetrahymena pyriformis* stock cultures were acquired from Carolina Biological

Supply (Burlington, NC, USA). Both species were kept in autoclaved liquid protist medium

(Carolina Biological Supply), supplemented with one autoclaved wheat seed as a carbon source

(Altermatt et al 2014). Protists were fed a mixture of pond bacteria collected from an ephemeral

pond at Duke Forest (Wilbur/Gate 9 pond, Lat 36.013914, -Long 78.979720). We maintained all

cultures on a 16:8 light:dark cycle at 22°C and 65% humidity in AL-22L2 Percival growth

chambers (Perry, IA, USA).

Experimental design

Microcosms were set up in autoclaved 250mL borosilicate jars filled with 200mL of protist

media. We manipulated temperature and nutrient loads by imposing two temperature levels

(22°C/25°C) and two nutrient levels (normal protist media concentration plus one wheat seed –

i.e., high nutrients–, or half concentration plus half a wheat seed, –i.e., low nutrients) in a

factorial design with 4 treatments and 6 replicates per treatment. Day/night cycle and humidity

levels mimicked rearing conditions and were kept constant. We inoculated the bacterial

communities from 2 ml of stock culture, the same used to rear the protists. The intermediate

consumer species, *T. pyriformis*, was introduced at a starting density of 37 ind/ml, and the larger

protists species, *Euplotes sp.*, at a starting density of 0.24 ind/ml in all microcosms. We recorded

the density of both protist species through fluid imaging (FlowCam; Yokogawa Fluid Imaging

Technologies, Portland, ME, USA), Monday through Friday, for 16 days. Fluid imaging

generates individual cell raster images that were used to quantify changes in protist size

(measured as cell area, in μm^2) over time. Bacteria density was quantified as optical density at a

wavelength of 600nm (OD600), using a BioTEK Epoch-2 microplate spectrophotometer

(Winooski, VT, USA).

Data analysis

To test for possible effects of temperature and nutrients on ecological dynamics, we fitted

Generalized Additive Mixed Models (GAMM) to time series of species density (OD600 for

bacteria) and protist body size, across all treatments, using the 'mgcv' package (version 1.8-

31,(Wood 2011)) in R (version 4.0.2, R Core Team 2020). To control for temporal

autocorrelation, we used an Autoregressive Moving Average (ARMA) correlation structure in

the GAMM models using the 'nlme' R package (version 3.1-148). We compared models with

additive and/or interactive effects temperature and nutrients effects, as well as different ARMA

correlation structures, using AICc (Appendix I Table S1-2). We discarded *T. pyriformis*

phenotypic data from days on which less than 10 individuals were measured after populations

collapsed.

Characterization of ecological dynamics

To assess environmental effects on microcosm food web dynamics, we characterized multiple

aspects of the population dynamics of bacteria and protists and their responses to environmental

conditions. Specifically, we quantified: 1) initial growth rate (day^{-1}) as $[\ln(Nf) -$

$\ln(Ni)]$/time for early-dynamics data, (up to day 1 for *T. pyriformis* & bacteria and day 8 for

Euplotes sp., due to much slower growth), 2) maximum density (individual/ml for protists and

OD_{600} for bacteria), using the highest average daily population density across replicates but

within treatments for protists, and the top three observed OD600 values for each replicate from

each treatment for bacteria because their population dynamics remained relatively stable, 3) the

coefficient of variation (CV = standard deviation/mean) of the temporal population dynamics

within treatments (typically used as a measure of stability (McCann 2012)), 4) the time to

population collapse in days (only *T. pyriformis*), and 5) the time to population peak in days (only

Euplotes sp.).

Quantifying top-down/bottom-up effects and feedbacks between ecological and body size

dynamics

To understand the mechanisms through which temperature and nutrients affected the observed

food web dynamics, we quantified the reciprocal effects of ecological and body size dynamics

using Convergent Cross Mapping (CCM) from the R package 'rEDM' (version 1.5.0; Park et al

2020). CCM quantifies the strength of causation of one dynamical variable onto another by

measuring the extent to which the time series of one variable X can reliably estimate the state of

another variable Y in a future time step (Sugihara *et al.* 2012). The higher the ability of X to

predict Y, the higher the causal effect of Y on X. Meanwhile, a variable Y that does not

influence a variable X cannot be predicted from the dynamics of X (Sugihara *et al.* 2012). This

procedure yields a 'predictability' value that can be interpreted as a correlation coefficient

between past and future states of the two dynamical variables (Sugihara *et al.* 2012). Whether

there is a cause-effect relationship between two dynamical variables, as opposed to simple

correlation, further depends on whether the average predictability value increases with library

size (i.e., number of data points taken into account to run the analysis, Sugihara et al., 2012). We

took this into account to infer causation in our data (see Appendix II Fig. S1-14).

CCM is often applied to relatively long time series (Sugihara *et al.* 2012). However, the

dynamics in our system occurred much too fast (Fig. 1), precluding us from collecting longer

time series. To resolve this issue, we used CCM on the GAMM model predictions of changes in

density (for all species) and body size (for protists) with time, for each treatment. These GAMM

model predictions considered the effects of each treatment as a whole—i.e., each combination of

nutrient and temperature levels—because they more closely reproduced the observed dynamics

(Appendix I Table S1). We refer to these GAMM predictions as 'time-series' in the context of

CCM analyses. Using CCM on the density time series, we calculated the predictability of

changes in prey density from changes predator density (i.e., bottom-up control, as predator

density predicting prey density means changes in prey density causes changes in predator

densities, Sugihara et al., 2012) and the predictability of changes in predator density from

changes in prey (i.e., top-down control). We dubbed these "eco-eco" effects, for simplicity. To

assess feedbacks between body size and ecological dynamics, we calculated the predictability of

changes in protist body size from changes in species density (i.e., causal effects of changes in

body size on densities, or 'pheno-eco' effects), and vice versa ('eco-pheno' effects). Last, we

calculated the predictability of changes in predator and prey body sizes on each other ('pheno-

pheno' effects).

RESULTS

General Ecological and Phenotypic dynamics

Overall, bacterial density increased to carrying capacity early (Fig. 1a, light grey), *T. pyriformis*

increased rapidly, then decreased (Fig. 1a, dark grey), while *Euplotes sp.* increased almost

monotonically to carrying capacity (Fig. 1a, black), ultimately resulting in a two-species state.

Temperature and nutrients affected all three species and led to significantly different dynamics

across treatments (Fig. 1b-d, Appendix I Table S2). The body sizes of both protists changed

rapidly over time (Fig. 1e, f) and responded to both temperature and nutrients (Fig. 1e, f,

Appendix I Table S2).

Temperature and nutrient effects change across trophic levels in systematic ways

Temperature significantly affected a larger number of dynamical descriptors than nutrients (Fig.

2, 3). Individual temperature effects were more frequent in the upper trophic levels than in

bacteria (Fig. 3), as were interactions between temperature and nutrients (Fig. 3). In particular,

higher temperature alone i) strongly increased initial growth rate in Tetrahymena ($T = 0.50025$, p

$= 1.16 \times 10^{-6}$, Fig. 2b), ii) decreased the maximum density of bacteria and *T. pyriformis* ($T_B = $ -

0.004917, $p = 0.00419$; $T_T = -975.3$, $p < 0.0155$; Fig. 2d-e) but increased that of *Euplotes sp.* (T_E

$= 39.83$, $p = 0.02668$, Fig. 2f), iii) increased the variability in *T. pyriformis* density (i.e. CV, $T = $

0.20033, p = 0.00554, Fig. 3h) but decreased that of *Euplotes sp.* (T = -0.07689, p = 0.01692,

Fig. 3i), iv) accelerated the time to collapse in *T. pyriformis* (T = -2.7576, p = 1.59×10^{-5}; Fig. 2j),

and v) accelerated the time to peak density in *Euplotes sp.* (T = -2.1667, p = 0.0196; Fig. 2k).

Higher nutrients alone i) decreased the initial growth rate of *Euplotes sp.* (N = -0.087, p =

5.02×10^{-4} Fig. 2c), ii) increased maximum density of all species (N_B= 0.012717, p = 2.03×10^{-10};

N_T = 1002.2, p = 0.0132; N_E = 60.67, p = 0.00157, Fig. 2d-f), and iii) increased the CV of

bacteria and *Euplotes sp.* (N_B = 0.17521, p = 8.54×10^{-5}, Fig. 2g; N_E = 0.15853, p = 2.94×10^{-5},

Fig. 2i) while decreasing that of *T. pyriformis* (N_T = -0.18727, p = 0.00875, Fig. 2h).

Interestingly, increased temperature and nutrients had opposite effects on the CV of *T. pyriformis*

and *Euplotes sp.*, indicating antagonistic additive effects of temperature and nutrients on food

web stability.

Temperature and nutrients had complex interactive effects on ecological dynamics of both protist

consumers. Nutrients had a negative effect on the initial growth rate of *Euplotes sp.* at low

temperature, but a positive effect at high temperature (N×T = 0.12621, p = 0.000391; Fig. 2c).

The maximum density of *T. pyriformis* increased with nutrients, but this effect was stronger at

high temperature (N×T = 1526.7, p = 0.0083, Fig. 2e). Nutrients also had i) a negative effect on

T. pyriformis population variability (CV) at low temperature but no effect at high temperature

(N×T = 0.19492, p = 0.04500, Fig. 2h), and ii) a positive effect on *Euplotes sp.* CV that was

stronger at low temperature (N×T = -0.12045, p = 0.00913, Fig. 2i). Additional model stats can

be found in Appendix I Table S3. Taken together, these results imply that temperature and

nutrients individually and interactively influence population dynamics at all trophic levels, with

more of both types of effects occurring at higher trophic levels (Fig. 3).

Temperature and nutrients influence top-down and bottom-up effects

Using CCM analysis, we estimated changes in the relative strength of bottom-up and top-down

ecological effects across treatments (Fig. 4a). In general, top-down and bottom-up effects

between the top predators were the strongest (Fig. 4a, yellow). Top-down effects involving

bacteria were the most strongly influenced by temperature and nutrients (Fig.4a, green & purple

dashed lines). Importantly, temperature and nutrients showed interactive effects on top-down and

bottom-up control. Most notably, top-down control by both protists on bacteria decreased with

nutrients at low temperature but increased with nutrients at high temperatures (Fig.4a, open dots

and dashed lines).

The most dominant species interactions in the overall food web also changed according to shifts

in the strength of top-down and bottom-up control between species pairs (Figure 5a). To be

conservative and increase the chance of picking up biologically relevant effects, we only present

effects whose CCM predictability value was above 0.75 in Figure 5 (Appendix III Table S1).

Specifically, increasing nutrients weakened top-down control from *T. pyriformis* to bacteria at

low temperatures (Fig. 5a left panel, dashed purple line), but strengthened top-down control from

both protists to bacteria at high temperatures (Fig. 5a right panel, dashed lines). In other words,

top-down effects were strongest when temperature and nutrients were both low, or both high,

suggesting that shifts in the relative importance of top-down and bottom-up effects across trophic

levels explain some effects of temperature and nutrients on the overall food web.

Temperature and nutrients alter the reciprocal effects of ecological and phenotypic dynamics

Protist size not only responded to temperature and nutrients, but also played an important role in

determining the effects of temperature and nutrients on ecological dynamics (Fig. 4b-c). On

average, higher temperature generally led to stronger eco-pheno effects (mean CCM at 22°C =

0.55, mean CCM at 25°C = 0.66, Fig. 4b, Appendix II Table S1) and pheno-eco interactions (size

driving ecological dynamics, mean CCM at 22°C = 0.71, mean CCM at 25°C = 0.79, Fig. 4c,

Appendix II Table S1), indicating stronger coupling between ecological and phenotypic

dynamics at higher temperatures. Pheno-eco effects were generally stronger (14 strong effects,

i.e., CCM predictability > 0.75, Fig. 4b) than eco-pheno effects (11 strong effects, Fig. 4c),

suggesting stronger effects of body size on ecological dynamics than the other way around. For

example, *Euplotes sp.* size had stronger effects on its own population dynamics than the opposite

(Fig. 4b-c, black dotted lines).

Both eco-pheno and pheno-eco effects generally responded interactively to changes in

temperature and nutrients. For instance, *T. pyriformis* eco-pheno effects on *Euplotes. sp* (yellow

solid lines in Fig.4b) decreased with nutrients at low temperature but remained relatively

unchanged at high temperature. Also, nutrients increased the effect of *T. pyriformis* size on

bacterial dynamics and this pattern was more exaggerated at higher temperature (Fig. 4c, dashed

purple lines). Meanwhile, nutrients decreased the effects of *Euplotes sp.* size on bacteria

dynamics at low temperature and this pattern was reversed at high temperature (Fig. 4c, dashed

green lines). In 22 out of the 25 eco-pheno or pheno-eco interactions, we found strong support

for causation (Appendix II Fig. S1-14).

Pheno-pheno effects between protists (Fig. 4d) were relatively weaker than pheno-eco or eco-

pheno effects. In general *T. pyriformis* size more strongly affected *Euplotes sp*. size than the

other way around (Fig. 4d). Additionally, temperature generally increased body size effects

between protists and we observed the strongest phenotypic coupling between protists when

temperature and nutrients were both high (Fig. 4d), indicating that higher temperatures may

increase the role of phenotypic dynamics in consumer-resource interactions.

Overall, these results indicate that the effect of temperature and nutrients on ecological dynamics

may be mediated by changes in body size (Fig. 5). Temperature and nutrients showed negative

interactions on strong eco-eco effects (CCM predictability > 0.75), leading to stronger eco-eco

effects when temperature and nutrients were both low or both high (Fig. 5a). However, the

number of strong causal effects between ecological and phenotypic dynamics increased with

temperature but were weakened by nutrients (Fig. 5b).

DISCUSSION

Our results reveal complex but systematic effects of temperature and nutrients on the ecological

and phenotypic dynamics of a microbial food web. We show that temperature and nutrients can

each independently influence different aspects of food web dynamics (Fig. 1, 2), while their joint

effects get increasingly complex at higher trophic levels (Fig. 2, 3). We also found that changes

in the relative strength of top-down and bottom-up effects likely drive observed responses to

temperature and nutrients (Fig. 4, 5), and that rapid changes in body size mediate these effects

(Fig. 4, 5). These results suggest that body size not only responds to shifts in environmental

conditions, but also plays a role in determining ecological responses to such shifts. Evaluating

feedbacks between ecological and phenotypic dynamics is therefore integral to understanding

food web responses to environmental change.

Increasingly complex effects of temperature and nutrients at higher trophic levels

Our results reveal the pervasive effects of temperature and nutrients across all species, but these

effects are more numerous and increasingly complex (temperature-nutrient interactions) at higher

trophic levels (Fig. 2, 3). The effects of temperature and nutrients on dietary preference, species

interactions, and foraging behavior may help explain why these interactive effects may be more

likely among consumer species (Fig. 1d; Fig. 2c).

Because energy enters at the bottom of a food web, basal species are more likely to be influenced

by direct effects of nutrients while species at higher trophic levels may be more strongly affected

by temperature (Petchey *et al.* 1999; Voigt *et al.* 2003)8/12/21 11:03:00 AM. As energy flows

from the basal species to the consumers, the temperature and nutrient effects that act on basal

species could be integrated by consumers, leading to additional indirect temperature and nutrient

effects on consumers. Additionally, these effects may be exaggerated in omnivores due to the

complex nature of their trophic interactions. As temperature and nutrients change species

interactions and foraging behaviors (Sentis *et al.* 2014), omnivorous consumers may shift diets

between basal and intermediate resources (Sentis *et al.* 2014), altering the strength/prevalence of

competitive vs. predatory interactions (Vandermeer 2006), thus rewiring food webs (Bartley *et*

al. 2019). Taken together, this may indicate why the joint effects of temperature and nutrients

may be more complex at higher trophic levels in food webs with higher prevalence of omnivory.

Top-down control, bottom-up control, and food web stability

Consistent with our results (Fig. 2d), theory predicts that higher nutrient loads should increase

energy flux in food webs, increasing basal species density (Rip & McCann 2011; Shurin *et al.*

2012). Theory also predicts that increasing energy flux can destabilize population

oscillations(Rosenzweig 1971; Rip & McCann 2011), but increasing temperatures should

stabilize oscillations by weakening top-down effects (Vasseur & McCann 2005; Binzer *et al.*

2016; Tabi *et al.* 2019). We found that temperature and nutrients interact to have divergent

impacts on population stability—with temperature stabilizing, and nutrients destabilizing

Euplotes sp. densities, but the opposite being true for *T. pyriformis* (Fig. 2h-i). We also observed

that top-down control of bacteria by both consumers was strongest when temperature and

nutrients were both high (and both low), while bottom-up effects were relatively invariant across

treatments (Fig. 4a, Fig. 5). Interestingly, these shifts in top-down control across environments

(Fig. 4a) coincided with significant changes in population stability that varied across trophic

levels (Fig. 2i). These results thus indicate that changes in the strength of top-down effects

between basal resources and consumers—instead of bottom-up effects—could be the dominant

mechanism through which higher temperature stabilize unstable fluctuations caused by nutrients.

Phenotypes mediate temperature and nutrients effects on food web dynamics

Our results indicate that phenotypic dynamics play a larger role in mediating environmental

impacts on food web dynamics than previously thought (Fig. 5, Appendix III Fig. S1). Previous

studies showed that predator-prey body size ratios significantly influence temperature and

nutrients effects (Binzer *et al.* 2012) and that body size responds to changes in nutrients,

temperature, and ecological dynamics in specific ways (Tabi *et al.* 2019). Our results support—

but also extend—those results by showing that 1) phenotypic effects on ecological dynamics

may be stronger than the other way around, as also recently reported elsewhere (Gibert *et al.*

2021), 2) changes in prey size may more strongly drive changes in predator size than the other

way around, and 3) interactions involving phenotypic dynamics (eco-pheno, pheno-eco, or

pheno-pheno) vary in complex ways across environmental conditions, but seem to be generally

stronger at higher temperature (Fig. 4b-d, Fig. 5). Rapid phenotypic change has been suggested

as a main driver of food web rewiring in future climates (Barbour & Gibert 2021), but little

experimental evidence exists. Here, we found that strong feedbacks between ecological and

phenotypic dynamics vary with temperature and nutrients, suggesting that rapid phenotypic

change influences food web responses to changing environmental conditions. Together, our

results emphasize the need to incorporate phenotypic dynamics in future studies of food web

responses to warming and eutrophication in a changing world.

ACKNOWLEDGMENTS

This work was supported by a U.S. Department of Energy, Office of Science, Office of

Biological and Environmental Research, Genomic Science Program Grant under Award Number

DE-SC0020362 to JPG.

Figures

Figure 1. Population and phenotypic dynamics within experimental microcosms. (a) The average population density across treatments shows the general population dynamics of each species. (b-d) show GAMM smoothed population dynamics of bacteria, *T. pyriformis* and *Euplotes sp.*. (e-f) show the changes in the area (body size) of both protists body over time. *T. pyriformis* were effectively extinct by day 9. Greyed area indicates days for which data are insufficient to estimate trait distributions.

Figure 2. Interactive plot of effects of temperature and nutrients on descriptors of population

dynamics. Open circles and triangles represent data points from each replicate. Solid shapes are

the mean across 6 replicates. Grey letters T, N, and (T×N) represent statistically significant

effects from temperature, nutrients, and interactive effects between temperature and nutrients,

respectively.

Fig. 3. Significant temperature and nutrients effects on ecological dynamics. T, N, and T×N

denote effects of temperature, nutrients, and interactive effects of temperature and nutrients,

respectively.

Figure 4. CCM predictabilities ecological and phenotypic dynamics across treatments for each

species pair. Circles represents CCM predictability of one species' effect on another species,

while the triangles representing intraspecific. Solid circles and solid lines represent CCM

predictability of bottom-up effects and open circles and dashed lines represent top-down effects.

Interactions with CCM predictability higher than 0.75 are considered to be strong interactions.

Figure 5. Strong effects (i.e., CCM predictability higher than 0.75) within food webs across

treatments. Panel (a) shows strong eco-eco effects (Fig. 4a). Panel (b) shows all strong

interactions between phenotypic (Ph) and ecological (E) dynamics (pheno-eco, eco-pheno and

pheno-pheno effects; Fig. 4b-d). Dashed and solid lines represent top-down and bottom-up

effects respectively. Dotted lines represent intraspecific effects. Color code is the same as in Fig.

4a.

453 References:
454
Atkinson, D. (1995). Effects of temperature on the size of aquatic ectotherms: Exceptions to the
general rule. *J. Therm. Biol.*, 20, 61–74.
Atkinson, D., Ciotti, B.J. & Montagnes, D.J.S. (2003). Protists decrease in size linearly with
temperature: *ca* . 2.5% °C $^{-1}$. *Proc. R. Soc. Lond. B Biol. Sci.*, 270, 2605–2611.
Barbour, M.A. & Gibert, J.P. (2021). Genetic and plastic rewiring of food webs under climate
change. *J. Anim. Ecol.*, 1365-2656.13541.
Barneche, D.R., Hulatt, C.J., Dossena, M., Padfield, D., Woodward, G., Trimmer, M., *et al.*
(2021). Warming impairs trophic transfer efficiency in a long-term field experiment.
Nature.
Bartley, T.J., McCann, K.S., Bieg, C., Cazelles, K., Granados, M., Guzzo, M.M., *et al.* (2019).
Food web rewiring in a changing world. *Nat. Ecol. Evol.*, 3, 345–354.
Barton, B.T., Beckerman, A.P. & Schmitz, O.J. (2009). Climate warming strengthens indirect
interactions in an old-field food web. *Ecology*, 90, 2346–2351.
Bernhardt, J.R., Sunday, J.M. & O'Connor, M.I. (2018). Metabolic Theory and the Temperature-
Size Rule Explain the Temperature Dependence of Population Carrying Capacity. *Am.
Nat.*, 192, 687–697.
Binzer, A., Guill, C., Brose, U. & Rall, B.C. (2012). The dynamics of food chains under climate
change and nutrient enrichment. *Philos. Trans. R. Soc. B Biol. Sci.*, 367, 2935–2944.
Binzer, A., Guill, C., Rall, B.C. & Brose, U. (2016). Interactive effects of warming,
eutrophication and size structure: impacts on biodiversity and food-web structure. *Glob.
Change Biol.*, 22, 220–227.
Brose, U., Dunne, J.A., Montoya, J.M., Petchey, O.L., Schneider, F.D. & Jacob, U. (2012).
Climate change in size-structured ecosystems. *Philos. Trans. R. Soc. B Biol. Sci.*, 367,
2903–2912.
Carlier, A., Riera, P., Amouroux, J.-M., Bodiou, J.-Y., Desmalades, M. & Grémare, A. (2008).
Food web structure of two Mediterranean lagoons under varying degree of
eutrophication. *J. Sea Res.*, 60, 264–275.
Clarke, A. (2006). Temperature and the metabolic theory of ecology. *Funct. Ecol.*, 20, 405–412.
Dossena, M., Yvon-Durocher, G., Grey, J., Montoya, J.M., Perkins, D.M., Trimmer, M., *et al.*
(2012). Warming alters community size structure and ecosystem functioning. *Proc. R.
Soc. B Biol. Sci.*, 279, 3011–3019.
Fenchel, T. (1974). Intrinsic rate of natural increase: The relationship with body size. *Oecologia*,
14, 317–326.
Ferenc, V. & Sheppard, C.S. (2020). The stronger, the better – trait hierarchy is driving alien
species interaction. *Oikos*, 129, 1455–1467.
Fussmann, G.F. (2000). Crossing the Hopf Bifurcation in a Live Predator-Prey System. *Science*,
290, 1358–1360.
Gauzens, B., Rall, B.C., Mendonça, V., Vinagre, C. & Brose, U. (2020). Biodiversity of
intertidal food webs in response to warming across latitudes. *Nat. Clim. Change*, 10, 264–
269.
Gibert, J.P. (2019). Temperature directly and indirectly influences food web structure. *Sci. Rep.*,
9, 5312.
Gibert, J.P., Han, Z.-Y., Wieczynski, D.J. & Yammine, A. (2021). Rapid plastic shifts in body
size precede and determine population growth. BioRxiv. doi: 10.1101/2021.07.15.452551

Gounand, I., Kéfi, S., Mouquet, N. & Gravel, D. (2016). Trait selection during food web assembly: the roles of interactions and temperature. *Theor. Ecol.*, 9, 417–429.

Hautier, Y., Niklaus, P.A. & Hector, A. (2009). Competition for Light Causes Plant Biodiversity Loss After Eutrophication. *Science*, 324, 636–638.

Hautier, Y., Seabloom, E.W., Borer, E.T., Adler, P.B., Harpole, W.S., Hillebrand, H., *et al.* (2014). Eutrophication weakens stabilizing effects of diversity in natural grasslands. *Nature*, 508, 521–525.

Irwin, A.J., Finkel, Z.V., Schofield, O.M.E. & Falkowski, P.G. (2006). Scaling-up from nutrient physiology to the size-structure of phytoplankton communities. *J. Plankton Res.*, 28, 459–471.

Joseph Park, Cameron Smith, George Sugihara and Ethan Deyle (2020). rEDM: Empirical Dynamic Modeling ('EDM'). R package version 1.5.0. https://CRAN.R-project.org/package=rEDM

Karl, T.R. (2003). Modern Global Climate Change. *Science*, 302, 1719–1723.

Kratina, P., Greig, H.S., Thompson, P.L., Carvalho-Pereira, T.S.A. & Shurin, J.B. (2012). Warming modifies trophic cascades and eutrophication in experimental freshwater communities. *Ecology*, 93, 1421–1430.

Lee, G.H., Vonk, J.A., Verdonschot, R.C.M., Kraak, M.H.S., Verdonschot, P.F.M. & Huisman, J. (2021). Eutrophication induces shifts in the trophic position of invertebrates in aquatic food webs. *Ecology*, 102.

Malcolm, J.R., Liu, C., Neilson, R.P., Hansen, L. & Hannah, L. (2006). Global Warming and Extinctions of Endemic Species from Biodiversity Hotspots. *Conserv. Biol.*, 20, 538–548.

Marañón, E., Cermeño, P., López-Sandoval, D.C., Rodríguez-Ramos, T., Sobrino, C., Huete-Ortega, M., *et al.* (2013). Unimodal size scaling of phytoplankton growth and the size dependence of nutrient uptake and use. *Ecol. Lett.*, 16, 371–379.

McCann, K.S. (2012). *Food webs*. Monographs in population biology. Princeton University Press, Princeton, N.J.

McClelland, J. & Valiela, I. (1998). Changes in food web structure under the influence of increased anthropogenic nitrogen inputs to estuaries. *Mar. Ecol. Prog. Ser.*, 168, 259–271.

O'Connor, M.I. (2009). Warming strengthens an herbivore–plant interaction. *Ecology*, 90, 388–398.

Petchey, O.L., McPhearson, P.T., Casey, T.M. & Morin, P.J. (1999). Environmental warming alters food-web structure and ecosystem function. *Nature*, 402, 69–72.

R Core Team (2020). R: A language and for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rip, J.M.K. & McCann, K.S. (2011). Cross-ecosystem differences in stability and the principle of energy flux: Cross-ecosystem differences in stability. *Ecol. Lett.*, 14, 733–740.

Rosenblatt, A.E. & Schmitz, O.J. (2016). Climate Change, Nutrition, and Bottom-Up and Top-Down Food Web Processes. *Trends Ecol. Evol.*, 31, 965–975.

Rosenzweig, M.L. (1971). Paradox of Enrichment: Destabilization of Exploitation Ecosystems in Ecological Time. *Science*, 171, 385.

Savage, V.M., Gillooly, J.F., Brown, J.H., West, G.B. & Charnov, E.L. (2004). Effects of Body Size and Temperature on Population Growth. *Am. Nat.*, 163, 429–441.

Sentis, A., Hemptinne, J.-L. & Brodeur, J. (2014). Towards a mechanistic understanding of
temperature and enrichment effects on species interaction strength, omnivory and food-
web structure. *Ecol. Lett.*, 17, 785–793.
Shurin, J.B., Clasen, J.L., Greig, H.S., Kratina, P. & Thompson, P.L. (2012). Warming shifts top-
down and bottom-up control of pond food web structure and function. *Philos. Trans. R.*
Soc. B Biol. Sci., 367, 3008–3017.
Sugihara, G., May, R., Ye, H., Hsieh, C. -h., Deyle, E., Fogarty, M., *et al.* (2012). Detecting
Causality in Complex Ecosystems. *Science*, 338, 496–500.
Tabi, A., Petchey, O.L. & Pennekamp, F. (2019). Warming reduces the effects of enrichment on
stability and functioning across levels of organisation in an aquatic microbial ecosystem.
Ecol. Lett., 22, 1061–1071.
Vandermeer, J. (2006). Omnivory and the stability of food webs. *J. Theor. Biol.*, 238, 497–504.
Vasseur, D.A. & McCann, K.S. (2005). A Mechanistic Approach for Modeling Temperature-
Dependent Consumer-Resource Dynamics. *Am. Nat.*, 166, 184–198.
Voigt, W., Perner, J., Davis, A.J., Eggers, T., Schumacher, J., Bährmann, R., *et al.* (2003).
TROPHIC LEVELS ARE DIFFERENTIALLY SENSITIVE TO CLIMATE. *Ecology*,
84, 2444–2453.
Wang, J., Pan, F., Soininen, J., Heino, J. & Shen, J. (2016). Nutrient enrichment modifies
temperature-biodiversity relationships in large-scale field experiments. *Nat. Commun.*, 7,
13960.
Wieczynski, D., Singla, P., Doan, A., Singleton, A., Han, Z., Yammine, A., Gibert, J.P. (2021).
Simple traits predict complex temperature responses across scales. Research Square.
doi:10.21203/rs.3.rs-116110/v1
Wood, S.N. (2011). Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models: Estimation of Semiparametric
Generalized Linear Models. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 73, 3–36.
569