Microbial colonization of tannin-rich tropical plants: interplay between

degradability, methane production and tannin disappearance in the rumen

M. Rira[1,2†], D.P. Morgavi[1†], M. Popova[1], G. Maxin[1], and M. Doreau[1]

[1]INRAE, VetAgro Sup, UMR1213 Herbivores, F-63122 Saint-Genès-Champanelle,

France

[2]Ecole Nationale Supérieure de Biotechnologie, Ali Mendjli, BP E66, 25100

Constantine, Algeria

11
12 [†] E-mail: moufida_r@yahoo.fr;
13 [†]diego.morgavi@inrae.fr
14
15

Abstract

Condensed tannins in plants are found free and attached to protein and fibre but it is

not known whether these fractions influence rumen degradation and microbial

colonization. This study explored the rumen degradation of tropical tannins-rich

plants and the relationship between condensed tannins fractions' disappearance and

microbial communities colonising plant particles. Leaves from *Calliandra calothyrsus*,

Gliricidia sepium, and *Leucaena leucocephala*, pods from *Acacia nilotica* and the

leaves of two agricultural by-products: *Manihot esculenta* and *Musa* spp. were

incubated in situ in the rumen of dairy cows. An In vitro approach was also used to

assess the effects of these plants on rumen fermentation parameters. All plants

contained more than 100 g/kg of condensed tannins with a large proportion (32 to

61%) bound to proteins. *Calliandra calothyrsus* had the highest concentration of

condensed tannins at 361 g/kg, whereas *Acacia nilotica* was particularly rich in

hydrolysable tannins (350 g/kg). Hydrolysable and free condensed tannins from all

plants completely disappeared after 24 h incubation in the rumen. Disappearance of

protein-bound condensed tannins was variable with values ranging from 93% for

Gliricidia sepium to 21% for *Acacia nilolitica*. In contrast, fibre-bound condensed

tannins disappearance averaged ~82% and did not vary between plants.

Disappearance of bound fractions of condensed tannins was not associated with

degradability of plant fractions. The presence of tannins interfered with the microbial

colonisation of plants. Each plant had distinct bacterial and archaeal communities

after 3 and 12 h of incubation in the rumen and distinct protozoal communities at 3 h.

Adherent communities in tannin-rich plants had a lower relative abundance of

fibrolytic microbes, notably *Fibrobacter* spp. Whereas, archaea diversity was reduced

in high tannin-containing *Calliandra calothyrsus* and *Acacia nilotica* at 12 h of

incubation. Concurrently, in vitro methane production was lower for *Calliandra*

calothyrsus, *Acacia nilotica* and *Leucaena leucocephala* although for the latter total

volatile fatty acids production was not affected and was similar to control. Here we

show that the total amount of hydrolysable and condensed tannins contained in a

plant govern the interaction with rumen microbes affecting degradability and

fermentation. The effect of protein- and fibre-bound condensed tannins on

degradability is less important.

Keywords: tropical plants; hydrolysable tannins; condensed tannins; methane, *in situ*

degradability

Introduction

Feed availability is a major limitation in many tropical ruminant production systems. One way for farmers to increase forage availability in these grazing systems is to use leguminous trees for supplementing diets. Leaves from leguminous trees are a high-protein feed resource that balances the nutrition of ruminants grazing tropical grasses poor in protein (Roothaert and Paterson, 1997). However, they remain underutilised because of the presence of tannins that are often found at high concentrations. High concentrations of tannins can reduce voluntary feed intake and nutrient digestibility (Frutos et al., 2004). The decrease in nutritive value is associated with tannins' property to bind to proteins, both from the diet and digestive enzymes. Tannins also bind to structural carbohydrates present in plant cell walls (Mueller-Harvey et al., 2019). Despite these adverse effects, a low concentration of tannins in the diet can improve nitrogen (N) utilisation efficiency and have a positive enteric methane-reduction effect (Goel and Makkar, 2012). This latter effect is because tannins may reduce organic matter digestibility in the rumen even when total-tract digestibility is unchanged, or because they inhibit microbial populations, or both (Frutos et al., 2004).

Tannins are conventionally classified into two major groups: hydrolysable (HT) and condensed tannins (CT). Hydrolysable tannins consist of polyphenols (gallic acid and/or hexahydroxydiphenic acid) ester-linked to a hexose moiety. They are categorised according to their structural characteristics into two subgroups: gallotannins and ellagitannins. In contrast, CT are polymers of varying molecular weight composed of flavan-3-ol (e.g., catechin) or flavan-3,4-diol (proanthocyanidins) linked by C–C or C–O–C bonds. Condensed tannins are found in different fractions in plants: free, protein-bound, and fibre-bound (Mueller-Harvey and McAllan, 1992,

Terrill et al., 1992, Schofield et al., 2001). A better understanding of the effects of

HT-rich and CT-rich forages on nutrient digestibility and methane mitigation

properties would improve the management of such resources. To this end, the

relationship between feed degradation and the disappearance of CT fractions must

be established for developing feeding strategies overcoming undesirable effects

when using tannin-rich forages. We hypothesised that rumen microbial degradation

of tannin-rich forages is influenced not only by the abundance of tannins but also by

their chemical form and binding to plants' structural components. This knowledge

would be of considerable importance for the efficient utilisation of these forages in the

tropics.

Our objective was to study the degradation of tannin-rich forages in the rumen and

establish the relationship with the amount and disappearance of their different tannin

fractions. We connected these effects on forage degradation and tannin

disappearance with the microbial communities colonising feed particles in the rumen

and with fermentation parameters. We used six tropical forages from leguminous

shrubs and agricultural by-products with differing amounts and nature of tannins. We

carried out 1) an in situ experiment, in order to determine the ruminal degradation of

forage components, including different tannin fractions, and the colonisation of feed

particles by microbes, and 2) an in vitro experiment, in order to measure methane

production and feed fermentation.

Materials and methods

The use of experimental animals followed the guidelines for animal research of the

French Ministry of Agriculture and other applicable guidelines and regulations for

animal experimentation in the European Union. Animals were housed at the INRAE

UE1414 Herbipôle Unit (Saint Genès Champanelle, France;

https://doi.org/10.15454/1.5572318050509348E12). Procedures were approved by

French Ministry of Education and Research (APAFIS #8218-20161151782412).

Plant material

Four browse species and two crops by-products rich in tannins and available in the

tropics were selected. The browse species were leguminous shrubs: *Acacia nilotica*

pods, *Calliandra calothyrsus, Gliricidia sepium* and *Leucaena leucocephala* leaves.

The by-products were cassava (*Manihot esculenta*) leaves and banana (*Musa* spp.)

leaves. These species are consumed by ruminants and have a large range of CT

and HT content.

Acacia nilotica samples were collected from the Ferlo region of Senegal (15°N,

15°W) having a mean annual rainfall of 200-400 mm and a mean temperature of 28-

30°C. *Gliricidia sepium*, *Leucaena leucocephala*, *Manihot esculenta* and *Musa* spp.

were collected in Guadeloupe, Basse-Terre Island, France (16°N, 61°W), having a

mean annual rainfall of 1500-2000 mm and a mean temperature of 24-28°C.

Calliandra calothyrsus was collected from native shrubs in the south of the Réunion

Island, France (21°S, 55°E) having a mean annual rainfall of 1000-1500 mm and a

mean temperature of 20-24°C; leaves were harvested at a late vegetative stage.

Following collection, fresh material was dried at 40 °C to avoid degradation or

modification of tannins, ground to pass through a 1-mm grid and placed in air-tight

plastic bags. Upon reception at the laboratory in Metropolitan France, samples were

stored at room temperature until use for *in situ* and *in vitro* measurements. Forages

poor in tannins were used as a control when necessary for a better understanding of

processes. For studying microbial colonisation of feed particles, the control forage

was the natural grassland hay used for feeding animals. For studying changes in

methane production and feed fermentation, the control forage was hay from a 75-d

regrowth of natural grassland based on *Dichanthium* spp. harvested in Guadeloupe,

Grande-Terre Island, France.

Experiment 1: In situ rumen degradation

Three rumen-cannulated Holstein non-lactating cows were used for the study with an

average body weight of 737 ± 40 kg. Cows were housed in individual pens and fed

natural grassland hay, first cycle harvested in Auvergne, France. A fixed amount of

7.6 kg DM of hay was offered to each cow twice a day: 2/3 at 0900 h, 1/3 at 1600 h.

Water was on free access. In situ ruminal incubations started after 15 days of

adaptation to the diet.

Feeds were incubated in the rumen for 3, 6, 12, 24, 48 and 96 h. Three grams of

ground samples were put into 5.5 × 12 cm polyester bags (pore size ca. 50 μm,

model R1020, Ankom, Fairport, NY). Bags were hooked to a stainless-steel weight

and inserted in the ventral sac of the rumen at 0800 h. Two successive series of

incubations were performed for each cow. Each series had one bag per feed and

incubation time for measurements of DM, N and NDF degradation, two bags per feed

for measurements of CT disappearance at 24 h of incubation to get enough residues

for further tannin analyses and two additional bags per feed for measurements of

microbial colonisation at 3 h and at 12 h. At each designated incubation time, bags

for measurements of DM, N and NDF degradation and tannin disappearance were

removed from the rumen, immersed in cold water and then washed under running tap

water until the water became clear. Zero-time disappearance was obtained by

washing two non-incubated bags per feed as described above. Bags were kept at 4

°C for 48 h then washed in a washing machine without detergent until clean water

was obtained (4 cycles of 10 min), dried at 40 °C for 96 h and weighed to determine

rumen residual DM content. Bags for microbial analysis, taken out after 3 and 12 h of

incubation, were gently squeezed, put on ice and immediately transferred to the lab

where they were washed three times in phosphate-buffered saline at 4°C for 5 min

on a rocking shaker at ~40 cycles per min. Bags were then snap-frozen in liquid N

and stored at -80°C until gDNA extraction.

Experiment 2: In vitro fermentation

Fermentations were performed using a batch technique (Rira et al., 2019). The

donor animals were four Texel wethers fitted with a ruminal cannula and weighed on

average 80.7 ± 6.9 kg. Wethers were fed daily 900 g hay (natural grassland based

on *Dichanthium* spp.) divided into equal amounts at 0700 and 1900 h. Wethers were

adapted to the diet for 2 weeks before being used as donors. Four series of 24-h

incubations were performed, one per wether. In each series, control *Dichanthium*

and the six tannin-rich forages were anaerobically incubated in duplicate as

described (Rira et al., 2019).

Gas production was measured at 24 h using a pressure transducer. After recording

pressure, a gas sample (5 mL) was taken for methane analysis. For VFA

determination, 0.8 mL of filtrate was mixed with 0.5 mL of 4 mg/mL crotonic acid and

20 mg/mL metaphosphoric acid in 0.5 M HCl and frozen at -20°C until analysis.

Chemical analyses

Feeds and feed residues in bags were analysed according to the Association of

Official Analytical Chemists (AOAC, 2005). Organic matter in feeds was determined

by ashing at 550 °C for 6 h (AOAC method number 923.03). Nitrogen (N) in feeds

and feed residues was determined by the Dumas method (crude protein (CP) = N ×

6.25, AOAC method number 992.15). For residues at 24, 48 and 96 h, residues of

the 3 cows were pooled to get enough residue for analysis. Cell wall components in

feeds (NDF, ADF, and ADL) were determined using sodium sulphite, without heat-

stable amylase and including residual ash (AOAC methods number 200.04 and

973.18). Enzymatic dry matter digestibility was estimated in feeds by hydrolysis with

pepsin in 0.1 N HCl then with fungal cellulose (Aufrere and Michalet-Doreau, 1988).

Soluble CT, protein-bound CT and fibre-bound CT fractions in tannins-rich plants

samples were extracted and analysed according to Terrill et al. (1992) as previously

detailed (Rira et al., 2019). The concentration of CT in soluble, protein-bound and

fibre-bound fractions was calculated with a standard calibration curve of purified

quebracho tannins.

For HT, the rhodanine method was used for determination of gallotannins in feeds

(Inoue and Hagerman, 1988). The potassium iodate (KIO3) method was used to

estimate total HT (gallotannins and ellagitannins) in feeds (Hartzfeld et al., 2002).

Details of these two methods are mentioned in (Rira et al., 2019). Ellagitannins were

calculated using the difference between total HT and gallotannins.

After in vitro fermentation, gas composition was determined by gas chromatography

(Micro GC 3000A; Agilent Technologies, Les Ulis, France) within 2 h after sampling.

Gas molar concentration was calibrated using a certified standard. Volatile fatty

acids were analysed by gas chromatography using crotonic acid as internal standard

on a Perkin Elmer Clarus 580 GC (Perkin Elmer, Courtaboeuf, France) equipped with

a Stabilwax-DA column (30 m by 0.53 mm i.d.) (Morgavi et al., 2013).

Microbial analysis: DNA extraction and sequencing strategy

DNA was extracted following the protocol described by Yu and Morrison (2004). Total

genomic DNA was sent to Roy J. Carver Biotechnology Center (Illinois, USA) for

fluidigm amplification and Illumina sequencing using primers targeting bacterial 16S

rRNA gene (V3-V5 region), archaeal 16S rRNA gene, fungal ITS2 and 18S rRNA

gene for protozoa, as described (Saro et al., 2018, Popova et al., 2019).

Microbial bioinformatics analyses.

Raw sequencing data were trimmed for quality (Phred score > 25), expected

amplicon length (570 nt for bacteria 16S rRNA gene, 457 nt for archaea 16S rRNA

gene, 660 nt for 18S rRNA gene and 356 nt for ITS) and maximum 5 primer

mismatches. Sequences were analysed using computational pipelines as described

(Saro et al., 2018). On average per sample, we obtained 28 045 (±6 457) reads of

bacterial 16S rRNA gene, 23 718 (±4 726) for archaeal 16S rRNA gene and 18 645

(±2 696) for eukaryotic 18S rRNA gene. For fungi, most samples had a low number

of reads, precluding further comparative analysis between samples. As forward and

reverse reads for bacterial 16S rRNA gene amplicons reads and eukaryotic 18S

rRNA gene amplicons reads were not overlapping, sequence data were analysed

using IM Tornado pipeline (Jeraldo et al., 2014) and taxonomy assigned according to

Silva v128. Archaeal sequences were analysed following standard QIIME pipeline

(Caporaso et al., 2010) and taxonomy assigned with RIM DB (Seedorf et al., 2014).

Sequencing data are available in the Sequence Read Archive (SRA) under

accession ID PRJNA554299.

OTU tables were analysed in R using the package "vegan" (Oksanen et al., 2016).

Diversity indices (Shannon, Simpson, Richness and Evenness) were computed using

implemented functions and statistical differences were tested using the non-

parametric Kruskal-Wallis test to evaluate the effect of plant at each incubation time.

For β-diversity analysis, OTU tables were rarefied to an even depth: 3460 reads for

bacteria, 4103 for Archaea and 1000 for protozoa. Bray-Curtis method was used for

computing dissimilarity indices with the vegdist function. Principal component

analysis (PCoA) was performed on dissimilarity matrices with prcomp function.

Permutational multivariate analysis of variance was performed using Adonis function.

Correlations were computed using vegan's function "corr" and "Hmisc" and "corrplot"

packages were used for plotting correlation matrices. Differential abundance

analysis to evaluate the effect of plant at each incubation time were done with

MicrobiomeAnalyst (Chong et al., 2020) using default data filtering (4 minimum count,

20% prevalence, low variance filter: 10% inter-quantile range), relative log expression

(RLE) for normalisation and the metagenomeSeq package (Paulson et al., 2013).

Degradation and fermentation data: calculations and statistical analyses

Rumen degradability (D) of DM and N were calculated using an exponential model

with lag time (Denham et al., 1989):

$D(t) = a + b(1 - e^{-c(t-L)})$

where D is the degradation after t hours of rumen incubation; t = hours of rumen

incubation (0, 3, 6, 12, 24, 48 and 96 h); a = rapidly degradable fraction (%); b =

slowly degradable fraction (%); c = degradation rate constant of the b fraction (h-1);

L= lag time before the beginning of degradation of the b fraction (h).

The nonlinear procedure (PROC NLIN) of SAS v9.4 (SAS Inst. Inc., Cary, NC, USA)

was used to fit degradation data to the model. This model was chosen after a

preliminary comparison of models with or without lag time, because lag time was

higher than 3 h for 4 of the 6 feeds, and because the model with lag time globally led to lower sum of squares than the model without lag time. The theoretical degradability in the rumen derived from the model (TDm) was calculated from the equation:

$$TDm = a + [b \times c /(c + kp)]$$

where kp is the passage rate of solid contents out of the rumen. A unique value of 0.04 h^{-1} was used for all feeds. Rumen degradability of NDF did not fit to the exponential model due to a linear degradation rate for *Calliandra calothyrsus* and to erratic variations for other two forages. For NDF calculations of degradability, we used instead a stepwise method (Kristensen et al., 1982):

$$TDs = \Sigma_{(i = 0 \text{ to } n)} (Dt_{(i+1)} - Dt_i) \times p (t_i , t_{i+1})$$

where (Dt(i+1)-Dti) is the amount of feed degraded between times ti+1 and ti, and p (ti , ti+1) is the proportion of feed remaining in the rumen between times ti and ti+1 with pti = e-kpti.

From VFA production in the in vitro experiment, fermented organic matter (FOM) was calculated by the stoichiometric equation of Demeyer and Van Nevel (1975):

FOM = 162 (0.5 acetate + 0.5 propionate + butyrate + valerate) where FOM is expressed in mg and VFA in mmol.

Data of both experiments were submitted to the same mixed model using the MIXED procedure of SAS including feed (n = 6 for in situ experiment and n = 7 for in vitro experiment) as fixed effect and animal (n = 3 for in situ experiment and n = 4 for in vitro experiment) as random effect. Differences between feeds were analysed using the Tukey t-test. Effects were declared significant when $P < 0.05$. Principal component analyses were performed using Minitab® version 17 software (Minitab Inc., State College, PA). Average values for each of the 6 tannin-rich plants were

included for 19 variables: 7 from chemical composition, 5 from experiment 1 and 7

from experiment 2.

Results

Forage characteristics

The chemical composition of forages is shown in Table 1. Tannin-containing plants

have higher concentration of CP and lower concentration of fibre than the control

forage, except *Musa* spp. The enzymatic DM digestibility test showed high values

(~70%) for *Acacia nilotica* pods, *Leucaena leucocephala, Gliricidia sepium* and

Manihot esculenta leaves, whereas digestibility of *Musa* spp. and *Calliandra*

calothyrsus leaves was much lower at 40% or less. There were marked differences

among forages in both the total amount of CT and the fraction that tannins were

associated with (Table 1). For all forages, bound CT were predominantly (≥ 60%)

linked to protein compared to fibre. The proportion of free CT was particularly large

in *Calliandra calothyrsus* leaves (54% of total CT) and *Acacia nilotica* pods (58% of

total CT). In addition, *Acacia nilotica* pods were particularly rich in HT (350 g/kg DM

versus less than 33 g/kg DM for the other plants). For the others forages consisting

in plant leaves, the HT content was minor. For all forages, ellagitannins were

predominant representing ≥ 80% of total HT.

In situ dry matter, N and NDF degradability

Dry matter degradability was similar for *Acacia nilotica* pods, *Gliricidia sepium,*

Manihot esculenta, and *Leucaena leucocephala* with values around 65%, whereas

only a third of *Musa* spp. and *Calliandra calothyrsus* leaves were degraded (Table 2).

However, for forages presenting similar degradability, there were differences in the

proportion of degraded fractions and rate of degradation. *Acacia nilotica* pods and

Gliricidia sepium had a higher proportion of soluble fraction (a), whereas *Leucaena*

leucocephala and *Manihot esculenta* leaves had a higher proportion of potentially

degradable fraction (b). *Musa* spp. and *Calliandra calothyrsus*, had lower values for

the (a) and (b) fractions than the others forages. Although they have similar

degradability, *Musa* spp. leaves had a low proportion of slowly degraded fraction (b)

at 14%, whereas *Calliandra calothyrsus* leaves were the most slowly degraded

(0.022% h^{-1}).

The N degradability showed similar values and ranking than DM degradability.

Notwithstanding, there was a higher variation among forages with *Manihot esculenta*

leaves presenting the highest overall degradability due to its high proportion of

soluble fraction (a) and high rate of degradation (c). For both DM and N, the

theoretical degradability calculated according to the model was similar to values

obtained by a stepwise calculation. The rumen degradability of NDF was calculated

with the stepwise method only because the exponential method with lag time failed to

produce suitable models for most forages. Compared to N, NDF was generally less

degraded. In most plants around one third of NDF was degraded in the rumen;

notable exceptions were *Acacia nilotica* pods with 14% disappearance and

Calliandra calothyrsus with only 6%.

Condensed tannins, N and NDF disappearance

Results of 24-h disappearance of tannins, are presented in Table 3, together with N

and NDF disappearance to assess if there is a relationship between these

parameters. The disappearance of N and NDF at 24 h was calculated from

degradability values above to have the same point in time as tannins. After 24 h in

the rumen, free CT disappeared completely or almost completely (~98%) in all

forages. Among those plants exhibiting a high proportion of protein-bound CT, the

disappearance of this fraction was variable with up to 93% loss in *Gliricidia sepium*

and no disappearance in *Calliandra calothyrsus*. For this latter forage, it is noted that

N disappeared at 24 h was ~32% but it had also the highest amount of CT linked to

protein (Table 1). For fibre-bound CT, the average disappearance at 24 h was ~80%

and, despite numerical differences, did not differ significantly between forages (P>

0.05). The disappearance of total CT reflected the differences observed in the

disappearance of the various fractions and ranged from 58% for *Calliandra*

calothyrsus leaves to 95% for *Gliricidia sepium.*

Microbial community attached to tannin-rich plants

We studied ruminal microbial communities attached to plants after 3 and 12 h of

incubation in the rumen. Incubation times were chosen in order to catch the biphasic

primary and secondary colonisation process described in temperate plants (Elliott et

al., 2018). Based on the known differences in communities between these two

phases (Mayorga et al., 2016, Elliott et al., 2018), most results are presented by

incubation time to better identify the effect of plants.

For anaerobic fungi, most samples had a low number of reads and no downstream

analysis was made. However, it is noted that the only plant that was consistently

colonised by anaerobic fungi was *Musa* with a threefold increase from 3 to 12 h

reaching more than 4000 reads on average (Supplementary Figure 1). For protozoa,

Leucaena leucocephala had low numbers of reads but it was considered a

characteristic of the plant and these samples were included in downstream analysis.

Changes on alpha diversity indices were more marked for bacteria at 3 h

(Supplementary Table 1). *Calliandra calothyrsus* had high alpha diversity values that

remained numerically higher than for other plants at 12 h. Whereas, archaeal indices

differed more at 12 h than at 3 h; *Acacia nilotica* and, to a lesser degree, *Calliandra*

calothyrsus were the plants with the lowest values.

Principal coordinated analysis plots showed differences in community structure that

were influenced by the type of plant (Figure 1). Permanova analyses highlighted

significant differences between plants for all microbial communities at 3 h (Adonis P <

0.001; R^2= 0.53, 0.43 and 0.59 for bacteria, archaea and protozoa, respectively).

These differences remained at 12 h for bacteria (Adonis R^2= 0.69, P < 0.001) and

archaea (Adonis R^2= 0.48, P < 0.001). At 3 h, the first component separated

bacterial communities attached to *Acacia nilotica, Calliandra calothyrsus* and

Leucaena leucocephala from *Musa*, *Manihot esculenta*, and the control hay (same

hay fed to cows). However, at 12 h, only *Musa* and control hay grouped together

clearly separated from the other plants. For archaea, *Acacia nilotica* and *Calliandra*

calothyrsus were separated from other plants both at 3 and 12 h of incubation.

Whereas, for protozoa, *Leucaena leucocephala* was clearly separated from all other

plants at 3 h, undoubtedly due to the low number of reads recovered from this plant;

and, at 12 h not clear grouping of plants was observed in agreement with permanova

results. The bacterial and archaeal communities correlated with some chemical

features of plants; NDF and ADF contents influenced the bacterial community

structure and total concentration of tannins, both CT and HT, had a stronger

influence on the archaeal community structure (Supplementary Figure 2).

Differential abundance analyses showed a numerically higher proportion of

Proteobacteria for plants richer in tannins at 3 h of incubation (Table 4). At lower

taxonomical levels, the family Rhodospirillaceae from the Proteobacteria was

particularly abundant in *Calliandra calothyrsus* (Supplementary Table 4), whereas the

γ-proteobacterium *Pantoea* sp. was more abundant in *Calliandra calothyrsus* and

Leucaena leucocephala (Supplementary Table 2). In contrast, the phylum

Fibrobacteres was more abundant in *Musa* and control hay, which were the plants

containing a higher amount of fibre and less tannins. The differences were

particularly striking at 12 h of incubation with ~25% of sequences belonging to

Fibrobacteres in these two plants compared to values as low as 3% for *Acacia*

nilotica (Table 4). Results for families and genera levels are shown in Supplementary

Tables 2 to 5. *Lachnospiraceae* were proportionally more abundant in *Calliandra*

calothyrsus, Leucaena leucocephala and *Glyricidia sepium* at 3 h of incubation and

generally more abundant in tannin-rich plants at 12 h of incubation. Representative

genera of this family, such as *Butyrivibrio* and *Oribacterium* showed the same trend.

There were no marked differences in relative abundance of archaea (Supplementary

Tables 6 and 7) and, for protozoa, the main change was observed in the relative

abundance of *Isotricha* spp. that was higher at 3 h in *Acacia nilotica*; differences were

less marked at 12 h. In accord with these results, there was a positive correlation

between *Isotricha* spp. and the levels of HT and TCT (Supplementary Tables 8 and

9, and Supplementary Figure 2).

Correlation analyses considering the chemical composition of plants, including the

various fractions of tannins, and the most abundant (≥ 1%) OTUs only show a few

highlighted negative and positive associations. At 3 h, a *Succiniclasticum*_uncl. OTU

was negatively correlated to TCT and HT and a *Chistensenellaceae* R7 OTU was

positively associated to the fibre-linked fraction of CT. However, at 12 h, the same

Chistensenellaceae R7 OTU and a *Prevotella* OTU were negatively correlated to the

protein linked fraction of CT. A *Fibrobacter* OTU was negatively correlated to HT,

whereas *Ruminococcaceae*_NK4A214_group, *Rikenellaceae*_RC9_gut_group_uncl.

and a *Butyrivibrio*_2_uncl. were positively correlated to CT (total, free and protein

linked) (Figure 2).

In vitro rumen fermentation of forages

The in vitro incubation of each tannin-rich plant separately allowed to obtain

additional information on their effects on rumen function and methane production. All

tannin-rich plants had lower production of gas than control (natural grassland hay

based on Dichanthium spp.; Table 5). *Musa* spp. and *Calliandra calothyrsus* were

those that produced less gas. Only *Musa* spp., *Calliandra calothyrsus* and *Acacia*

nilotica had lower VFA production and FOM than control ($P < 0.05$). All tannin-rich

plants produced more acetate and less butyrate than control, resulting in the absence

of difference in the ratio of acetate or acetate+butyrate to propionate between control

and tannin-rich plants. Compared to control, methane production when expressed as

mL/24 h was reduced for *Leucaena leucocephala*, and to a greater extent for *Musa*

spp., *Calliandra calothyrsus* and *Acacia nilotica.* When methane production was

expressed per 100 mM of VFA produced, only *Acacia nilotica* and *Musa* spp.

decreased production ($P < 0.05$) compared to control.

The relationships between the content and type of tannins in plants and the in vitro

and in situ parameters were further explored through a principal component analysis

(Supplementary Figure 3) that showed that indicators of the extent of ruminal

degradation and fermentation, including methane, were opposed on the first axis to

total CT content and CT fractions, but not to HT.

Discussion

All plant species used in this experiment were rich in CT. The amount of CT was within those reported in the literature although mostly on the higher end. The higher CT values in this work can be explained by the analytical method, involving a double-extraction procedure and different solvents for different fractions, potentially yielding a higher amount of extracted tannins than faster methods. Four plants had a higher amount of protein-bound tannins than free tannins whereas two plants (*Acacia* and *Calliandra*) had a higher amount of free tannins. It is generally reported, even for plants used in this study such as *Leucaena leucocephala* and *Calliandra calothyrsus* that free CT are higher than protein-bound CT that in turn is higher than fibre-bound CT (Terrill et al., 1992, Jackson et al., 1996, Dentinho and Bessa, 2016). However, similar to total CT, there are important divergences in the literature (Perez-Maldonado and Norton, 1996, Rubanza et al., 2005).

In animal nutrition studies, the HT content of plant species is seldom measured and when done it is generally by using non-specific methods, e.g. they are often calculated by the difference between total tannins and CT. We used methods that specifically measured the content of total HT and gallotannins. Among the plants used in our study only *Acacia nilotica* was rich in HT. Goel et al. (2015) also reported high values of HT for this plant (186 g/kg DM, estimated by the difference between total tannins and CT). The other plants used in our study are recognised sources of CT but their HT content is seldom reported. Our results show that these plants rich in CT are also a source of HT, which albeit minor can also have a biological effect.

Degradation of tannin-rich plants in the rumen - relationship with tannin content

Calliandra calothyrsus and *Musa* spp. had a lower DM and N degradability (~33%) than the other plants (~65%). However, there is no a single reason that may explain these differences. *Calliandra calothyrsus* had a markedly low NDF degradability at 6% but it was the disappearance of condensed tannins, representing one third of plant weight, that affected the calculation of DM degradation. Whereas, the low DM degradability of *Musa* spp. is mainly explained by the high proportion of NDF (65% on a DM base). In both cases the rate constant c was low. In contrast, the high degradability of *Leucaena leucocephala*, *Glyricidia sepium*, *Acacia nilotica*, and *Manihot esculenta* was due to the high N degradability, to the extensive disappearance of CT, and to the total disappearance of HT from bags. Disproving our hypothesis, we did not observe any relationship either between N degradability and total or protein-bound tannin content, or between NDF degradability and total or fibre-bound tannin content. Similarly, the absence of relationship between total extractible tannins of plants and DM disappearance in situ of seven temperate browses was reported by Khazaal et al. (1993). This contrasts with in vitro degradation results obtained with 72 African browses where protein and NDF degradability were negatively correlated with soluble CT but not with insoluble CT (Rittner and Reed, 1992). The difference with our results may be explained either by the low number of forages in our experiment or by the methodology (in vitro *vs* in situ).

Rumen disappearance of tannins from plants

Information on tannin disappearance in the digestive tract of ruminants is scarce.

Hydrolysable tannins can be degraded by rumen microbes (Brooker et al., 1994,

Goel et al., 2005). However, as HT would be washed out of the bags, we did not

attempt to quantify them after incubation.

Most authors agree that there is no evidence for free CT degradation by microbes in

the rumen (McSweeney et al., 2001b, Patra et al., 2012). Notwithstanding, the

complete rumen disappearance of free CT observed is logical because these

compounds are water-soluble and are washed out of the bags. A nearly complete

(99%) disappearance of free CT of *Calliandra calothyrsus* between mouth and faeces

was reported by Perez-Maldonado and Norton (1996).

The disappearance of protein-bound CT varied largely between plants from 21 to

98%. This variation may be due to differences in strength of binding between

proteins and tannins (Le Bourvellec and Renard, 2019). As stated above, no

relationship was observed between disappearance of proteins and protein-bound CT

except for *Calliandra calothyrsus* that had a low N degradation and no disappearance

of protein-bound tannins. For this plant, a negative value for disappearance was

even obtained. This may be due to a technical problem as interactions with the

insoluble matrix, proteins, polysaccharides, and other plant polymers can decrease

the solubility of tannins in the extractant, resulting in an underestimation of tannin

content in feeds (Dentinho and Bessa, 2016). Another possible reason is a linkage

of dietary free CT with proteins. According to Hagerman (1989), if CT are present in

excess, all proteins available are bound to tannins, leading to insoluble complexes.

This is the case for *Calliandra calothyrsus* which contains more tannins than proteins

(361 vs 217 g/kg, respectively).

The fibre-bound CT represented a small proportion of the total CT content of the

forages used in our study. Their disappearance in the rumen varied between 61 and

98% but differences between forages were not significant.

Tannin-rich plants and methane production

Tannin concentration is often considered a critical factor affecting ruminal

fermentation (Patra and Saxena, 2011). Correspondingly, we generally observed

that the CT content of plants was negatively associated to in situ rumen degradability

and to in vitro fermentation parameters including methane production. However, the

effect on methane of some plants could also be due to parameters other than CT. As

protein concentration affects the volume of gas produced by the bicarbonate buffer in

the in vitro fermentation system, we will discuss methane production normalised by

VFA for assessing the impact of tannin-rich plants independently of the extent of gas

production. Compared to control, _Acacia nilotica_ and _Musa_ spp. reduced methane

production but probably not for the same reasons. For _Musa_ spp. the effect cannot

be ascribed to the amount of tannins as this plant has a low concentration compared

to others. The effect could be due to other secondary compounds such as

polyphenols that are present in _Musa_ spp. leaves in large amounts (Marie-

Magdeleine et al., 2010). For _Acacia nilotica_, tannins are the most plausible cause of

methane inhibition as this plant contains CT but is especially rich in HT. We

previously showed that HT could be more efficient than CT for reducing methane

(Rira et al., 2019). Several authors have also reported that HT can reduce methane

production without compromising overall rumen fermentation, but to date only

extracts were studied (Bhatta et al., 2009, Hassanat and Benchaar, 2013,

Jayanegara et al., 2015). It is likely that HT do not interfere with rumen fermentation

because they do not bind to protein or fibre and do not have an inhibitory effect on

microbes that can also degrade HT in the rumen (Patra et al., 2012).

In this study, *Gliricidia sepium* and *Manihot esculenta* did not decrease methane

production. In previous studies, *Gliricidia sepium* was less effective than *Leucaena*

leucocephala and *Manihot esculenta* for decreasing methane *in vitro* (Rira et al.,

2015) and *in vivo* (Archimede et al., 2016). The absence of effect of *Manihot*

esculenta on methane production in this study is unexpected, but is consistent with

the large variability of response of methane production to CT for a same plant

(Piluzza et al., 2014).

Tannin-rich plants modulate adherent rumen microbes

The colonising microbial community differed between plants and was influenced by

tannins as well as other chemical features such as fibre content. As plants were

incubated in the same rumen environment and exposed to the same microbiota there

is no doubt that the plant itself is selecting for their adherent microbiota, at least in the

initial colonisation stages monitored in this work. The process is probably a

combination of microbial tolerance to tannins' toxicity (Frutos et al., 2004) and

substrate preferences. Our results brings new insight on how methane production in

the rumen is affected by tannin-rich plants and concurs with the reported absence of

a relationship between chemical structure and biological effect of tannins including

methane production (McAllister et al., 2005, Naumann et al., 2018).

There is no equivalent published information on the attachment of rumen microbes to

these plants for a straight comparison. In contrast, the effect of tannin-rich plants or

extracts added to the diet have been reported in several studies.

The fungal community was targeted in our study but it was not further analysed as a

low number of reads was recovered. Nevertheless, *Musa* spp. with one of the lowest

amounts of tannins among the tested plants was the only one colonised by fungi.

This result could be interpreted as an inhibitory effect of tannins on fungi as reported

previously (Muhammed et al., 1995). For protozoa, it is noted that *Isotricha* spp.

were more abundant on *Acacia nilotica* samples at 3 h of incubation suggesting that

this genus was attracted to HT particularly abundant in this plant. This result is only

comparable within this study as the pore size of the in sacco incubation bags may

obstruct the free passage of protozoa but it can help explain the variable effect of

different tannin sources on protozoa reported in the literature (Patra and Saxena,

2011).

The plants' characteristics clearly influenced the structure of the attached bacterial

community. Initially, each plant seems to harbour a distinct community of primary

colonisers. Then, at 12 h when most of soluble tannins are no longer interfering, the

communities in tannin-rich plants became more similar but distinctly separated from

communities found in the control forage without tannin and the low-tannin containing

Musa. For archaea the pattern is different with only *Acacia nilotica* and *Calliandra*

calothyrsus clearly separating from other plants. This indicate that the plant effect is

more important on colonising bacteria than on archaea, which is logical as the latter

rely on metabolic end products from other microbes. The different archaeal structure

for tannin-rich *Acacia nilotica* and *Calliandra calothyrsus* could reflect a toxicity

threshold attained at the biofilm microenvironment level that needs to be proved.

Our results bring new evidence indicating that some of the differences reported on

rumen microbes when tannins are supplemented to the diet are due to changes in

the colonisation and development of feed-attached communities. The low proportion

of *Fibrobacter* attached to tannin-rich plants may explain why this fibrolytic bacterium

is often affected in supplemented animals (McSweeney et al., 2001a, Diaz Carrasco

et al., 2017, Harun et al., 2017, Salami et al., 2018). On the other hand, some

bacteria that were positively associated plants rich in tannins could provide new

avenues of exploration for improving the nutritional value of these forages through

modulation of the microbiota.

Conclusion

We used an integrated approach for assessing the effect of tannin-rich plants on

rumen processes. Tannin-rich plants have contrasting proportions of tannins bound

to protein or fibre. However, there was no relationship between the amount and

disappearance of bound tannins and the rumen degradability of protein or fibre of the

plants studied in this work, except for *Calliandra calothyrsus* that was extremely rich

in CT. Tannins present in plants restrained the colonization of certain rumen

microbial populations known to play a role in the degradation of recalcitrant feeds.

These results expand our understanding of the effects of tannins in the rumen

opening the way to improve their use in the diet of ruminants.

Acknowledgements

The authors thank the staff of UE 1414 Herbipôle for their technical collaboration,

animal care and management, A. Torrent and L. Genestoux for their contribution in

the experimental setup and sample analysis, and A. Roche (intern student at INRAE)

for her contribution in the analysis of condensed tannins. We are grateful to the

INRAE MIGALE bioinformatics platform (http://migale.jouy.inra.fr) for providing

computational resources and A. Bernard for initial bioinformatic analysis.

References

AOAC. 2005. Official methods of analysis of AOAC International. 16th ed. AOAC International, Washington, DC.

Archimede, H., M. Rira, D. J. Barde, F. Labirin, C. Marie-Magdeleine, B. Calif, F. Periacarpin, J. Fleury, Y. Rochette, D. P. Morgavi, and M. Doreau. 2016. Potential of tannin-rich plants, *Leucaena leucocephala*, *Glyricidia sepium* and *Manihot esculenta*, to reduce enteric methane emissions in sheep. J. Anim. Physiol. Anim. Nutr. (Berl) 100:1149-1158. http://doi.org/10.1111/jpn.12423

Aufrere, J. and B. Michalet-Doreau. 1988. Comparison of methods for predicting digestibility of feeds. Anim. Feed Sci. Technol. 20:203-218. http://doi.org/https://doi.org/10.1016/0377-8401(88)90044-2

Bhatta, R., Y. Uyeno, K. Tajima, A. Takenaka, Y. Yabumoto, I. Nonaka, O. Enishi, and M. Kurihara. 2009. Difference in the nature of tannins on in vitro ruminal methane and volatile fatty acid production and on methanogenic archaea and protozoal populations. J. Dairy Sci. 92:5512-5522. http://doi.org/10.3168/jds.2008-1441

Brooker, J. D., L. A. Odonovan, I. Skene, K. Clarke, L. Blackall, and P. Muslera. 1994. *Streptococcus caprinus* sp nov, a tannin-resistant ruminal bacterium from feral goats. Letters in Applied Microbiology 18:313-318. http://doi.org/10.1111/j.1472-765X.1994.tb00877.x

Caporaso, J. G., J. Kuczynski, J. Stombaugh, K. Bittinger, F. D. Bushman, E. K. Costello, N. Fierer, A. G. Pena, J. K. Goodrich, J. I. Gordon, G. A. Huttley, S. T. Kelley, D. Knights, J. E. Koenig, R. E. Ley, C. A. Lozupone, D. McDonald, B. D. Muegge, M. Pirrung, J. Reeder, J. R. Sevinsky, P. J. Turnbaugh, W. A. Walters, J. Widmann, T. Yatsunenko, J. Zaneveld, and R. Knight. 2010. QIIME allows analysis of high-throughput community sequencing data. Nat Methods 7:335-336. http://doi.org/10.1038/nmeth.f.303

Chong, J., P. Liu, G. Zhou, and J. Xia. 2020. Using MicrobiomeAnalyst for comprehensive statistical, functional, and meta-analysis of microbiome data. Nat. Protocols 15:799-821. http://doi.org/10.1038/s41596-019-0264-1

Demeyer, D. and C. Van Nevel. 1975. Methanogenesis, an integrated part of carbohydrate fermentation and its control. Pages 366–382 in Digestion and metabolism in the ruminant. I. McDonald and A. Warner, ed. The University of New England Publishing Unit, Armidale, Australia.

Denham, S. C., G. A. Morantes, D. B. Bates, and J. E. Moore. 1989. Comparison of two models used to estimate in situ nitrogen disappearance. J. Dairy Sci. 72:708-714. http://doi.org/10.3168/jds.S0022-0302(89)79163-3

Dentinho, M. T. P. and R. J. B. Bessa. 2016. Effect of tannin source and pH on stability of tannin-protein and fibre complexes. Rev. Ciênc. Agrárias 39:114-121. http://doi.org/10.19084/RCA15062

Diaz Carrasco, J. M., C. Cabral, L. M. Redondo, N. D. Pin Viso, D. Colombatto, M. D. Farber, and M. E. Fernandez Miyakawa. 2017. Impact of chestnut and quebracho tannins on rumen microbiota of bovines. Biomed Res. Int. 2017:9610810. http://doi.org/10.1155/2017/9610810

Elliott, C. L., J. E. Edwards, T. J. Wilkinson, G. G. Allison, K. McCaffrey, M. B. Scott, P. Rees-Stevens, A. H. Kingston-Smith, and S. A. Huws. 2018. Using 'omic approaches to compare temporal bacterial colonization of Lolium perenne, Lotus corniculatus, and Trifolium pratense in the rumen. Front. Microbiol. 9:2184. http://doi.org/10.3389/fmicb.2018.02184

Frutos, P., G. Hervás, F. J. Giráldez, and A. R. Mantecón. 2004. Review. Tannins and ruminant nutrition. Span. J. Agric. Res. 2:191-202.

Goel, G. and H. P. Makkar. 2012. Methane mitigation from ruminants using tannins and saponins. Trop. Anim. Health Prod. 44:729-739. http://doi.org/10.1007/s11250-011-9966-2

Goel, G., A. K. Puniya, C. N. Aguilar, and K. Singh. 2005. Interaction of gut microflora with tannins in feeds. Naturwissenschaften 92:497-503. http://doi.org/10.1007/s00114-005-0040-7

Goel, G., M. Raghav, V. Beniwal, and A. K. Puniya. 2015. Anaerobic degradation of tannins in *Acacia nilotica*
pods by *Enterococcus faecalis* in co-culture with ruminal microbiota. J. Gen. Appl. Microbiol. 61:31-33.
http://doi.org/10.2323/jgam.61.31

Hagerman, A. E. 1989. Chemistry of Tannin-Protein Complexation. Pages 323-333 in Chemistry and Significance
of Condensed Tannins. R. W. Hemingway, J. J. Karchesy, and S. J. Branham, ed. Springer US, Boston,
MA.

Hartzfeld, P. W., R. Forkner, M. D. Hunter, and A. E. Hagerman. 2002. Determination of hydrolyzable tannins
(gallotannins and ellagitannins) after reaction with potassium iodate. J. Agric. Food. Chem. 50:1785-
1790. http://doi.org/10.1021/jf0111155

Harun, N. L. A., A. R. Alimon, M. F. Jahromi, and A. A. Samsudin. 2017. Effects of feeding goats with *Leucaena*
leucocephala and *Manihot esculenta* leaves supplemented diets on rumen fermentation profiles,
urinary purine derivatives and rumen microbial population. J. Appl. Anim. Res. 45:409-416.
http://doi.org/10.1080/09712119.2016.1205499

Hassanat, F. and C. Benchaar. 2013. Assessment of the effect of condensed (acacia and quebracho) and
hydrolysable (chestnut and valonea) tannins on rumen fermentation and methane production in vitro.
J. Sci. Food Agric. 93:332-339. http://doi.org/10.1002/jsfa.5763

Jackson, F. S., T. N. Barry, C. Lascano, and B. Palmer. 1996. The extractable and bound condensed tannin
content of leaves from tropical tree, shrub and forage legumes. J. Sci. Food Agric. 71:103-110.
http://doi.org/10.1002/(Sici)1097-0010(199605)71:1<103::Aid-Jsfa554>3.0.Co;2-8

Jayanegara, A., G. Goel, H. P. S. Makkar, and K. Becker. 2015. Divergence between purified hydrolysable and
condensed tannin effects on methane emission, rumen fermentation and microbial population in
vitro. Anim. Feed Sci. Technol. 209:60-68. http://doi.org/10.1016/j.anifeedsci.2015.08.002

Jeraldo, P., K. Kalari, X. Chen, J. Bhavsar, A. Mangalam, B. White, H. Nelson, J. P. Kocher, and N. Chia. 2014. IM-
TORNADO: a tool for comparison of 16S reads from paired-end libraries. PLoS One 9:e114804.
http://doi.org/10.1371/journal.pone.0114804

Khazaal, K., X. Markantonatos, A. Nastis, and E. R. Ørskov. 1993. Changes with maturity in fibre composition
and levels of extractable polyphenols in Greek browse: Effects on in vitro gas production and in sacco
dry matter degradation. J. Sci. Food Agric. 63:237-244. http://doi.org/10.1002/jsfa.2740630210

Kristensen, E. S., P. Moller, and T. Hvelplund. 1982. Estimation of the effective protein degradability in the
rumen of cows using the nylon bag technique combined with the outflow rate. Acta Agric. Scand. A
Anim. Sci. 32:123-127. http://doi.org/10.1080/00015128209435738

Le Bourvellec, C. and C. M. G. C. Renard. 2019. Interactions between polyphenols and macromolecules: effect
of tannin structure. Pages 515-521 in Encyclopedia of Food Chemistry. L. Melton, F. Shahidi, and P.
Varelis, ed. Academic Press, Oxford.

Marie-Magdeleine, C., M. Boval, L. Philibert, A. Borde, and H. Archimède. 2010. Effect of banana foliage (*Musa*
x paradisiaca) on nutrition, parasite infection and growth of lambs. Livest. Sci. 131:234-239.
http://doi.org/10.1016/j.livsci.2010.04.006

Mayorga, O. L., A. H. Kingston-Smith, E. J. Kim, G. G. Allison, T. J. Wilkinson, M. J. Hegarty, M. K. Theodorou, C. J.
Newbold, and S. A. Huws. 2016. Temporal metagenomic and metabolomic characterization of fresh
perennial ryegrass degradation by rumen bacteria. Front. Microbiol. 7:23.
http://doi.org/10.3389/fmicb.2016.01854

McAllister, T. A., T. Martinez, H. D. Bae, A. D. Muir, L. J. Yanke, and G. A. Jones. 2005. Characterization of
condensed tannins purified from legume forages: chromophore production, protein precipitation, and
inhibitory effects on cellulose digestion. J. Chem. Ecol. 31:2049-2068. http://doi.org/10.1007/s10886-
005-6077-4

McSweeney, C. S., B. Palmer, R. Bunch, and D. O. Krause. 2001a. Effect of the tropical forage calliandra on
microbial protein synthesis and ecology in the rumen. J. Appl. Microbiol. 90:78-88.
http://doi.org/10.1046/j.1365-2672.2001.01220.x

McSweeney, C. S., B. Palmer, D. M. McNeill, and D. O. Krause. 2001b. Microbial interactions with tannins:
nutritional consequences for ruminants. Anim. Feed Sci. Technol. 91:83-93.
http://doi.org/10.1016/S0377-8401(01)00232-2

Morgavi, D. P., C. Martin, and H. Boudra. 2013. Fungal secondary metabolites from Monascus spp. reduce
rumen methane production in vitro and in vivo. J. Anim. Sci. 91:848-860.
http://doi.org/10.2527/jas.2012-5665

Mueller-Harvey, I., G. Bee, F. Dohme-Meier, H. Hoste, M. Karonen, R. Kolliker, A. Luscher, V. Niderkorn, W. F.
Pellikaan, J. P. Salminen, L. Skot, L. M. J. Smith, S. M. Thamsborg, P. Totterdell, I. Wilkinson, A. R.
Williams, B. N. Azuhnwi, N. Baert, A. G. Brinkhaus, G. Copani, O. Desrues, C. Drake, M. Engstrom, C.
Fryganas, M. Girard, N. T. Huyen, K. Kempf, C. Malisch, M. Mora-Ortiz, J. Quijada, A. Ramsay, H. M.
Ropiak, and G. C. Waghorn. 2019. Benefits of condensed tannins in forage legumes fed to ruminants:
importance of structure, concentration, and diet composition. Crop Science 59:861-885.
http://doi.org/10.2135/cropsci2017.06.0369

Mueller-Harvey, I. and A. McAllan. 1992. Tannins: their biochemistry and nutritional properties. Pages 151-217
in Advances in plant cell biochemistry and biotechnology. Vol. 1. I. M. Morrison, ed. JAI Press, London.

Muhammed, S., C. S. Stewart, and T. Acamovic. 1995. Effects of tannic acid, ellagic acid, gallic acid and catechin
on cellulose degradation by the rumen fungus *Neocallimastix frontalis* strain rel. Anim. Sci. 1995:147-
147. http://doi.org/10.1017/S0308229600029135

Naumann, H., R. Sepela, A. Rezaire, S. E. Masih, W. E. Zeller, L. A. Reinhardt, J. T. Robe, M. L. Sullivan, and A. E.
Hagerman. 2018. Relationships between structures of condensed tannins from texas legumes and
methane production during in vitro rumen digestion. Molecules 23:2123. http://doi.org/ARTN 2123

10.3390/molecules23092123

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R. B. O'Hara, G. L.
Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner. 2016. vegan: Community Ecology
Package. R package version 2.4-1. https://CRAN.R-project.org/package=vegan.

Patra, A. K., B.-R. Min, and J. Saxena. 2012. Dietary tannins on microbial ecology of the gastrointestinal tract in
ruminants. Pages 237-262 in Dietary phytochemicals and microbes. A. K. Patra, ed. Springer.

Patra, A. K. and J. Saxena. 2011. Exploitation of dietary tannins to improve rumen metabolism and ruminant
nutrition. J. Sci. Food Agric. 91:24-37. http://doi.org/10.1002/jsfa.4152

Paulson, J. N., O. C. Stine, H. C. Bravo, and M. Pop. 2013. Differential abundance analysis for microbial marker-
gene surveys. Nat. Methods 10:1200-1202. http://doi.org/10.1038/nmeth.2658

Perez-Maldonado, R. and B. Norton. 1996. Digestion of 14 C-labelled condensed tannins from *Desmodium
intortum* in sheep and goats. Br. J. Nutr. 76:501-513. http://doi.org/10.1079/BJN19960059

Piluzza, G., L. Sulas, and S. Bullitta. 2014. Tannins in forage plants and their role in animal husbandry and
environmental sustainability: a review. Grass Forage Sci. 69:32-48. http://doi.org/10.1111/gfs.12053

Popova, M., J. Guyader, M. Silberberg, A. R. Seradj, C. Saro, A. Bernard, C. Gérard, C. Martin, and D. P. Morgavi.
2019. Changes in the rumen microbiota of cows in response to dietary supplementation with nitrate,
linseed, and saponin alone or in combination. Appl. Environ. Microbiol. 85:e02657-02618.
http://doi.org/10.1128/aem.02657-18

Rira, M., D. P. Morgavi, H. Archimède, C. Marie-Magdeleine, M. Popova, H. Bousseboua, and M. Doreau. 2015.
Potential of tannin-rich plants for modulating ruminal microbes and ruminal fermentation in sheep. J.
Anim. Sci. 93:334-347. http://doi.org/10.2527/jas.2014-7961

Rira, M., D. P. Morgavi, L. Genestoux, S. Djibiri, I. Sekhri, and M. Doreau. 2019. Methanogenic potential of
tropical feeds rich in hydrolysable tannins. J. Anim. Sci. 97:2700-2710.
http://doi.org/10.1093/jas/skz199

Rittner, U. and J. D. Reed. 1992. Phenolics and in-vitro degradability of protein and fibre in West African
Browse. J. Sci. Food Agric. 58:21-28. http://doi.org/https://doi.org/10.1002/jsfa.2740580105

Roothaert, R. L. and R. T. Paterson. 1997. Recent work on the production and utilization of tree fodder in East
Africa. Anim. Feed Sci. Technol. 69:39-51. http://doi.org/https://doi.org/10.1016/S0377-
8401(97)81621-5

Rubanza, C., M. Shem, R. Otsyina, S. Bakengesa, T. Ichinohe, and T. Fujihara. 2005. Polyphenolics and tannins
effect on in vitro digestibility of selected *Acacia* species leaves. Anim. Feed Sci. Technol. 119:129-142.
http://doi.org/10.1016/j.anifeedsci.2004.12.004

Salami, S. A., B. Valenti, M. Bella, M. N. O'Grady, G. Luciano, J. P. Kerry, E. Jones, A. Priolo, and C. J. Newbold.
2018. Characterisation of the ruminal fermentation and microbiome in lambs supplemented with
hydrolysable and condensed tannins. FEMS Microbiol. Ecol. 94. http://doi.org/10.1093/femsec/fiy061

Saro, C., U. M. Hohenester, M. Bernard, M. Lagrée, C. Martin, M. Doreau, H. Boudra, M. Popova, and D. P.
Morgavi. 2018. Effectiveness of interventions to modulate the rumen microbiota composition and
function in pre-ruminant and ruminant lambs. Front. Microbiol. 9:1273.
http://doi.org/10.3389/fmicb.2018.01273

Schofield, P., D. M. Mbugua, and A. N. Pell. 2001. Analysis of condensed tannins: a review. Anim. Feed Sci.
Technol. 91:21-40. http://doi.org/10.1016/S0377-8401(01)00228-0

Seedorf, H., S. Kittelmann, G. Henderson, and P. H. Janssen. 2014. RIM-DB: a taxonomic framework for
community structure analysis of methanogenic archaea from the rumen and other intestinal
environments. PeerJ 2:e494. http://doi.org/10.7717/peerj.494

Terrill, T. H., A. M. Rowan, G. B. Douglas, and T. N. Barry. 1992. Determination of extractable and bound
condensed tannin concentrations in forage plants, protein concentrate meals and cereal grains. J. Sci.
Food Agric. 58:321-329. http://doi.org/10.1002/jsfa.2740580306

Yu, Z. and M. Morrison. 2004. Improved extraction of PCR-quality community DNA from digesta and fecal
samples. BioTechniques 36:808-812. http://doi.org/10.2144/04365ST04

 Table 1. Chemical composition and enzymatic digestibility, in g/kg DM, of plants used in experiments

	Control hay Exp. 1	Control hay Exp. 2	*Acacia nilotica* pods	*Calliandra calothyrsus* leaves	*Gliricidia sepium* leaves	*Leucaena leucocephala* leaves	*Manihot esculenta* leaves	*Musa spp* leaves
Organic matter	915	927	956	956	891	897	914	895
Crude protein	83	84	140	217	250	336	294	69
NDF	625	701	227	333	240	223	285	653
ADF	351	326	152	228	147	125	202	349
ADL	44	20	38	63	57	46	69	63
Total condensed tannins	ND[1]	ND	157	361	112	180	166	128
Free	ND	ND	91	194	5	54	59	38
Linked to protein	ND	ND	51	125	69	83	95	57
Linked to fibre	ND	ND	15	42	38	43	12	33
Hydrolysable tannins	ND	ND	350	33	9	13	12	4
Gallotannins	ND	ND	84	6	Traces	Traces	Traces	Traces
Ellagitannins	ND	ND	266	27	9	13	12	4
Enzymatic DM digestibility	-[2]	529	720	406	714	745	668	359

740   [1]ND: not detected
741   [2]Not measured
742   Control hay Exp. 1: Natural grassland hay harvested in Auvergne, France
743   Control hay Exp. 2: Natural grassland hay based on *Dichanthium spp.* harvested in Guadeloupe, French West Indies

744 **Table 2.** Dry matter, N and NDF degradability[1] of tropical tannin-rich plants in the rumen

	Acacia nilotica	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	SEM	P Value
Dry matter degradability								
a (%)	53.5[a]	22.1[e]	49.5[b]	37.3[c]	39.2[c]	27.7[d]	0.88	< 0.001
b (%)	30.4[b]	25.2[b2]	27.6[b]	45.7[a]	41.4[a]	14.0[c]	1.86	< 0.001
c (h^{-1})	0.038[ab]	0.022[b]	0.170[ab]	0.075[ab]	0.189[a]	0.027[b]	0.0337	0.013
L (h)	4.24[ab]	0.00[b]	5.84[a]	2.68[ab]	5.95[a]	4.90[ab]	1.595	0.026
TDm[3] (%)	65.2[a]	30.9[c]	65.6[a]	63.3[b]	66.0[a]	32.2[c]	0.46	< 0.001
TDs[2] (%)	65.7[a]	32.0[b]	65.1[a]	63.8[a]	65.5[a]	33.2[b]	0.70	< 0.001
N degradability								
a (%)	40.1[b]	24.7[e]	37.8[bc]	35.6[c]	47.5[a]	29.7[d]	0.74	< 0.001
b (%)	58.6[a]	45.1[ab]	47.3[a]	54.6[a]	45.0[ab]	13.4[b]	6.29	0.004
c (h^{-1})	0.031[bc]	0.013[c]	0.076[b]	0.054[bc]	0.155[a]	0.025[bc]	0.0115	< 0.001
L (h)	1.88[bc]	0.00[c]	5.95[a]	0.52[c]	5.40[ab]	8.54[a]	0.912	< 0.001
TDm[2] (%)	63.7[c]	33.2[d]	61.8[c]	65.7[b]	76.2[a]	33.3[d]	0.57	< 0.001
TDs[2] (%)	64.1[c]	34.7[e]	61.1[d]	66.2[b]	75.7[a]	34.1[e]	0.64	< 0.001
NDF degradability								
TDs[2] (%)	14.5[c]	6.3[d]	28.4[b]	30.0[b]	35.8[a]	30.7[ab]	1.27	< 0.001

[a-d] Values within a row with different superscripts differ significantly at P<0.05, n= 3.

[1] Degradation D was modelled according to the equation $D(t) = a + b\,(1-e^{-c(t+L)})$ where a is the rapidly degraded fraction, b the

slowly degraded fraction, c the rate of degradation of fraction b, t the time of incubation in the rumen and L the lag time.

[2] n= 2

[3] TDm : theoretical degradability calculated according to the model ; TDs : theoretical degradability calculated according to a

stepwise calculation.

Table 3. Rumen disappearance (%) of condensed tannins, N and NDF from tropical tannin-rich plants after 24 h of incubation

	Acacia nilotica	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	SEM	P Value
Condensed tannins	73.0[d]	58.2[e]	94.9[a]	77.8[cd]	88.6[ab]	84.3[bc]	2.10	<0.001
Free	98.3	97.4	100.0	100.0	100.0	100.0	-	-
Linked to protein	21.3[d]	-7.8[e]	92.9[a]	58.0[c]	83.0[ab]	72.7[bc]	3.24	<0.001
Linked to fibre	83.0	60.8	98.0	87.9	76.4	86.4	8.23	0.10
N	65.3[b]	31.7[c]	68.8[b]	73.9[b]	89.1[a]	33.8[c]	2.93	0.001
NDF	11.5[ab]	5.1[b]	31.6[ab]	33.8[ab]	43.4[a]	27.8[ab]	8.21	0.012

754 [a-d]Values within a row with different superscripts differ significantly at P<0.05.

Table 4. Relative abundance of rumen bacterial phyla colonizing tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 3 and 12 h

	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena Leucocephala*	*Manihot esculenta*	*Musa spp*	P value	FRD[2]
After 3 h of incubation									
Firmicutes	47.61	41.16	43.47	57.90	60.08	43.58	50.26	0.554	0.727
Bacteroidetes	36.59	37.89	42.87	32.39	29.41	42.71	32.71	0.375	0.687
Fibrobacteres	11.53	2.12	3.87	5.19	3.50	8.13	12.88	0.089	0.327
Spirochaetae	1.53	1.68	3.24	1.47	2.30	3.39	1.57	0.464	0.727
Actinobacteria	1.52	6.46	2.52	1.75	1.25	0.92	1.84	0.221	0.607
Proteobacteria	0.52	9.77	3.15	0.70	2.95	0.43	0.33	0.015	0.083
Tenericutes	0.46	0.84	0.74	0.40	0.34	0.63	0.22	0.868	0.868
Chloroflexi	0.25	0.08	0.14	0.19	0.15	0.21	0.19	0.595	0.727
After 12 h of incubation									
Firmicutes	38.54	55.98	45.32	54.81	43.48	35.15	40.42	0.592	0.609
Bacteroidetes	30.78	29.22	32.76	27.63	34.39	33.63	28.41	0.093	0.249
Fibrobacteres	26.76a	2.72d	5.88c	11.09b	13.93b	11.02b	23.15a	0.006	0.045
Spirochaetae	1.94	5.17	7.53	3.20	5.37	13.89	7.13	0.609	0.609
Tenericutes	0.93	5.31	7.50	2.57	2.45	5.93	0.38	0.076	0.249
Actinobacteria	0.65	0.32	0.37	0.36	0.10	0.19	0.31	0.550	0.609
Proteobacteria	0.25	1.24	0.46	0.22	0.20	0.11	0.11	0.155	0.309
Chloroflexi	0.16	0.05	0.18	0.10	0.07	0.09	0.09	0.484	0.609

[1]Control: Natural grassland hay harvested in Auvergne, France

[2]FDR= false discovery rate post-hoc adjustment

762 **Table 5.** *In vitro* ruminal fermentation of tropical tannin-rich plants

	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	SEM	P value
Total gas production (mL / 24 h)	34.41[a]	23.26[c]	16.25[d]	28.01[b]	24.28[bc]	27.95[b]	11.90[d]	1.630	< 0.001
Total VFA[2] production (mM / 24 h)	45.61[a]	31.98[bc]	27.14[c]	52.49[a]	41.53[ab]	48.11[a]	31.96[bc]	3.251	< 0.001
VFA composition (%)									
Acetate	54.15[b]	68.05[a]	65.01[a]	63.56[a]	61.21[ab]	64.92[a]	62.79[ab]	2.079	0.003
Propionate	19.08[ab]	18.07[b]	19.43[ab]	18.34[b]	23.67[a]	17.20[b]	18.20[b]	1.084	0.010
Butyrate	19.92[a]	6.28[bc]	5.32[c]	9.21[ab]	6.50[bc]	8.28[bc]	6.77[bc]	0.861	< 0.001
Isobutyrate	1.40[bc]	0.92[c]	2.24[ab]	2.19[ab]	1.78[bc]	2.25[ab]	2.97[a]	0.319	< 0.001
Isovalerate	2.30[b]	2.26[b]	4.24[a]	3.19[ab]	3.13[ab]	3.52[ab]	4.28[a]	0.761	0.005
Valerate	2.91[b]	4.28[ab]	3.46[ab]	3.30[ab]	3.52[ab]	3.67[ab]	4.64[a]	0.469	0.010
Caproate	0.25[abc]	0.13[c]	0.29[ab]	0.21[abc]	0.20[abc]	0.16[bc]	0.35[a]	0.045	0.003
Acetate:propionate	2.88[a]	3.90[a]	3.41[a]	3.49[a]	2.59[a]	3.78[a]	3.47[a]	0.281	0.038
(Acetate + butyrate):propionate	3.94[ab]	4.25[ab]	3.69[ab]	3.99[ab]	2.86[b]	4.27[a]	3.84[ab]	0.301	0.060
Fermented organic matter (mg / 24 h)	177[a]	112[bc]	90[c]	180[a]	142[ab]	165[a]	105bc	11.3	< 0.001
Fermented organic matter (% OM)	47.3[ab]	28.8[cd]	23.4[d]	51.0[a]	39.0[bc]	44.9[ab]	29.9[cd]	2.95	< 0.001
Methane production (mL / 24 h)	3.92[a]	1.75[c]	1.66[c]	3.54[a]	2.59[b]	4.00[a]	1.41[c]	0.201	< 0.001
Methane production (mL/100 mM VFA)	8.69[a]	5.52[b]	6.49[ab]	6.75[ab]	6.35[ab]	8.56[a]	4.45[b]	0.700	0.003

763   [a-d]Values within a row with different superscripts differ significantly at P<0.05.
764   [1]Control: Natural grassland hay based on *Dichanthium spp*. harvested in Guadeloupe, French West Indies.
765   [2] VFA: volatile fatty acids.
766

Figure 1. Distribution of bacterial, archaeal and protozoal communities associated to tropical tannin-rich plants after 3 and 12 h of in situ incubation in the rumen (PCoA).

● = control hay (no tannin); ● = *Acacia nilotica*; ● = *Calliandra calothyrsus*; ● = *Gliricidia sepium*; ● = *Leucaena leucocephala*; ● = *Manihot esculenta*; ● = *Musa spp*.

Figure 2. Correlations between microbial species and characteristics of tropical tannin-rich plants after 3 and 12 h of incubation of feeds in the rumen. A: bacteria at 3 h of incubation; B: bacteria, 12 h of incubation; C: protozoa, 3 h of incubation; D: protozoa, 12 h of incubation. For bacteria, operational taxonomic units with abundance > 1% are shown. Green and red dots indicate positive and negative significant correlations (P<0.05), respectively.

Dry matter degradability (Msdeg), Nitrogen degradability (Ndeg), Theoretical degradability of dry matter calculated according to a stepwise calculation (TDmspap), Theoretical degradability of nitrogen calculated according to a stepwise calculation (TDNpap), Fibre linked condensed tannins disappearance (FCTdisp), Theoretical degradability of NDF calculated according to a stepwise calculation (TDNDFpap), Free condensed tannins disappearance (freedisp), Total condensed tannins disappearance (TCTdisp), Proteins linked condensed tannins disappearance (PCTdisp), Fibre linked condensed tannins (FCT), Proteins linked condensed tannins (PCT), Total condensed tannins (TCT), Free condensed tannins (free), NDF degradability (NDFdeg), Hydrolysable tannins (HT).

Supplementary information

Supplementary Table 1. Diversity indices for bacterial, archaeal and protozoal populations colonising tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 3 and 12 h, statistical analysis was performed using the non parametric Kruskal-Wallis test

Microbial type Index	Incubation time (h)	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	SEM	P
Bacteria										
Shannon	3	4.57abc	4.55ab	4.78a	4.39bc	4.44bc	4.41bc	4.31c	0.166	0.036
	12	4.50	4.35	4.72	4.28	4.48	4.49	4.33	0.193	0.072
Simpson	3	0.97abc	0.97ab	0.98a	0.96bc	0.96abc	0.97bc	0.96c	0.009	0.049
	12	0.96ab	0.96ab	0.97a	0.95b	0.97a	0.97a	0.96ab	0.010	0.046
Evenness	3	0.76ab	0.80a	0.81a	0.73b	0.76ab	0.73b	0.72b	0.036	0.014
	12	0.75	0.75	0.79	0.71	0.74	0.74	0.73	0.028	0.099
Richness	3	266ab	269ab	287a	244bc	251bc	250bc	236c	17.4	0.014
	12	376	331	381	339	338	346	322	25.1	0.085
Archaea										
Shannon	3	5.82	5.86	5.67	5.79	5.80	5.45	5.75	0.231	0.74
	12	6.10ab	5.65b	5.84b	6.10a	6.17a	5.94ab	6.08ab	0.202	0.029
Simpson	3	0.99	0.99	0.99	0.98	0.98	0.98	0.99	0.005	0.51
	12	0.99abc	0.97c	0.98bc	0.99ab	0.99a	0.99ab	0.99ab	0.007	0.034
Evenness	3	0.74	0.75	0.76	0.72	0.77	0.75	0.73	0.022	0.52
	12	0.78abc	0.74c	0.76bc	0.80a	0.79ab	0.78abc	0.77abc	0.018	0.022
Richness	3	1242	1210	1242	1089	983	1147	1192	110.4	0.20
	12	1958a	1690b	1778ab	1944a	1980a	1807ab	1958a	130.9	0.042
Protozoa										
Shannon	3	2.08	1.87	2.04	2.00	1.96	1.61	1.53	0.284	0.12
	12	1.67	1.83	2.19	2.07	1.92	2.05	1.94	0.230	0.26
Simpson	3	0.82ab	0.76ab	0.84a	0.83a	0.84a	0.65b	0.72b	0.087	0.040
	12	0.67	0.78	0.85	0.83	0.80	0.81	0.77	0.066	0.13
Evenness	3	0.66ab	0.57b	0.70ab	0.69ab	0.81a	0.53b	0.51b	0.116	0.025
	12	0.53	0.58	0.73	0.72	0.69	0.68	0.60	0.098	0.11
Richness	3	25ab	20a	23ab	22ab	14b	22a	23a	3.520	0.031
	12	21	20	20	17	16	21	23	2.739	0.16

[1]Control: Natural grassland hay based on *Dichanthium spp.* harvested in Guadeloupe, French West Indies

Supplementary Table 2. Proportional abundance of bacterial taxa colonising tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 3 h

	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	P value[2]	FRD
Uncultured_rumen_bacterium	62.8	60.6	60.7	67.3	57.7	67.6	67.0	0.498	0.755
*Fibrobacter*_unclassified	9.8	1.7	3.0	4.8	3.0	7.2	10.8	0.030	0.248
*Streptococcus*_unclassified	4.2	0.2	0.1	0.8	0.3	0.1	0.5	0.451	0.747
*Prevotella*_1_unclassified	4.0	3.2	4.3	3.8	2.6	4.2	3.3	0.318	0.722
Uncultured_rumen_bacterium_unclassified	3.0	6.6	5.4	3.3	2.6	2.5	2.2	0.878	0.934
Uncultured_bacterium	2.9	3.3	4.3	3.1	2.9	3.7	2.7	0.193	0.653
*Butyrivibrio*_2_unclassified	1.8	1.3	4.5	4.4	7.0	3.3	1.4	0.003	0.041
*Christensenellaceae*_R_7_group_unclassified	1.4	1.7	1.5	1.5	0.9	1.6	1.5	0.622	0.864
Ruminococcus_flavefaciens	0.8	0.8	0.5	0.5	0.4	0.9	1.6	0.705	0.904
Prevotella_ruminicola	0.8	1.2	1.7	1.0	1.4	1.5	0.9	0.306	0.722
*Oribacterium*_unclassified	0.6	0.3	1.2	1.6	3.9	0.8	0.4	0.005	0.047
Uncultured_rumen_bacterium_5C0d_11	0.5	0.2	0.3	0.2	0.3	0.3	0.6	0.694	0.904
Uncultured_bacterium_unclassified	0.5	1.9	1.3	0.4	0.7	0.3	0.5	0.821	0.933
Unidentified_rumen_bacterium_RC2	0.5	0.3	0.5	0.2	0.4	0.2	0.3	0.108	0.542
Uncultured_rumen_bacterium_4C28d_15_unclassified	0.4	0.2	0.1	0.0	0.1	0.1	0.1	0.389	0.747
*Selenomonas*_1_unclassified	0.4	0.5	0.8	0.4	2.0	0.2	0.1	0.921	0.940
*Ruminococcus_sp*_HUN007	0.4	0.3	0.3	0.1	0.0	0.1	0.8	0.663	0.895
uncultured_*Lachnospiraceae*_bacterium	0.4	0.2	0.3	0.5	0.4	0.6	0.7	0.139	0.612
Bacterium_AC2043	0.3	0.4	0.3	0.3	0.2	0.3	0.5	0.814	0.933
*Bacteroidales*_BS11_gut_group_unclassified	0.3	0.4	0.4	0.1	0.3	0.1	0.3	0.595	0.864
Uncultured_rumen_bacterium_3C0d_9	0.3	0.1	0.3	0.1	0.1	0.1	0.2	0.227	0.709
Uncultured_*Prevotellaceae*_bacterium	0.3	0.1	0.4	0.4	0.3	0.4	0.3	0.095	0.530
Uncultured_*Clostridium_sp*	0.3	0.1	0.1	0.1	0.1	0.2	0.5	0.294	0.722
*Selenomonas_ruminantium*_AB3002	0.2	1.6	1.1	0.2	1.3	0.2	0.2	0.278	0.722
Butyrivibrio_fibrisolvens	0.2	0.3	0.5	0.9	1.7	0.4	0.2	0.155	0.612
*Lachnospiraceae*_bacterium_AC2012	0.2	0.2	0.1	0.4	0.4	0.1	0.1	0.361	0.747
*Prevotella_sp*_CA17	0.2	0.2	0.3	0.3	0.2	0.3	0.1	0.943	0.943
X_*Eubacterium_ventriosum*_group_unclassified	0.2	0.3	0.2	0.0	0.1	0.1	0.1	0.772	0.929
*Lachnospiraceae*_unclassified	0.2	0.1	0.1	0.1	0.2	0.2	0.1	0.919	0.940
*Succinivibrio*_unclassified	0.2	0.6	0.4	0.3	0.9	0.2	0.1	0.404	0.747
*Lachnospiraceae*_bacterium_NK4A144	0.2	0.2	0.4	0.3	0.8	0.2	0.1	0.159	0.612
Unidentified_rumen_bacterium_RC17	0.2	0.2	0.1	0.1	0.1	0.1	0.2	0.383	0.747
*Rikenellaceae*_RC9_gut_group_unclassified	0.2	0.5	0.2	0.1	0.1	0.1	0.1	0.463	0.747
*Lachnospiraceae*_NK3A20_group_unclassified	0.2	0.1	0.2	0.2	0.0	0.1	0.1	0.196	0.653
Bacterium_XPD3003	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.444	0.747
Uncultured_rumen_bacterium_3C0d_20	0.1	0.1	0.0	0.2	0.1	0.2	0.2	0.460	0.747
Unidentified_rumen_bacterium_RFN89	0.1	0.1	0.0	0.3	0.2	0.2	0.1	0.242	0.712
*Roseburia*_unclassified	0.1	0.1	0.1	0.3	0.3	0.2	0.1	0.463	0.747
*Bacteroidales*_RF16_group_unclassified	0.1	0.2	0.3	0.1	0.1	0.1	0.1	0.723	0.904
*Bacteroidetes*_unclassified	0.1	0.1	0.1	0.2	0.2	0.3	0.1	0.872	0.934
Bacterium_XPB1013	0.1	0.0	0.2	0.2	0.2	0.1	0.1	0.072	0.516

X_*Eubacterium_hallii*_group_unclassified	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.610	0.864
Ruminococcus_albus	0.1	0.0	0.0	0.1	0.0	0.1	0.2	0.781	0.929
Bacterium_XPB4001	0.1	0.3	0.3	0.1	0.1	0.1	0.0	0.306	0.722
Unidentified_rumen_bacterium_RF31	0.1	0.2	0.2	0.3	0.2	0.2	0.1	0.857	0.934
Unidentified_rumen_bacterium_RFN13	0.1	0.0	0.0	0.2	0.2	0.2	0.1	0.497	0.755
Bacteria_unclassified	0.0	0.0	0.8	0.1	0.1	0.1	0.0	<0.001	0.001
*Clostridiales_va*dinBB60_group_unclassified	0.0	0.5	0.1	0.0	0.2	0.0	0.0	0.090	0.530
*Pantoea*_unclassified	0.0	0.1	1.1	0.0	3.0	0.0	0.0	0.001	0.018

[1]Control: Natural grassland hay based on *Dichanthium spp*. harvested in Guadeloupe, French West Indies

[2]Differential abundance was estimated with the metagenomeSeq package (Paulson et al., 2013)

Supplementary Table 3. Proportional abundance of bacterial taxa colonising tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 12 h

	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	P value[2]	FRD
Uncultured_rumen_bacterium	53.1	52.5	54.2	56.7	47.6	51.3	55.4	0.938	0.979
*Fibrobacter*_unclassified	23.7	2.0	4.6	9.9	12.2	9.6	19.9	0.030	0.142
Uncultured_bacterium	2.5	6.4	3.8	2.2	3.8	5.8	2.4	0.157	0.430
Uncultured_rumen_bacterium_unclassified	2.0	1.9	3.3	3.1	2.7	2.2	1.1	0.202	0.482
*Prevotella*_1_unclassified	2.0	3.8	4.5	3.2	6.3	3.0	2.9	0.545	0.799
Uncultured_rumen_bacterium_4C28d_15_unclassified	1.4	0.1	0.8	0.2	0.4	0.1	0.2	<0.001	<0.001
Prevotella_ruminicola	1.4	0.7	1.0	2.9	1.1	2.3	1.1	0.517	0.795
Unidentified_rumen_bacterium_RC2	1.3	0.5	1.0	0.3	0.3	0.3	0.5	0.131	0.394
Uncultured_rumen_bacterium_5C0d_11	1.1	0.1	0.1	0.1	0.2	0.5	2.6	0.977	0.979
Ruminococcus_flavefaciens	1.0	0.1	0.2	0.3	0.3	0.5	1.7	0.206	0.482
Uncultured_*Prevotellaceae*_bacterium	0.9	2.9	3.1	2.6	5.2	4.5	2.4	0.130	0.394
*Christensenellaceae*_R_7_group_unclassified	0.8	1.3	0.7	0.6	0.6	0.6	0.7	0.077	0.271
*Ruminococcus_sp*_HUN007	0.8	0.1	0.1	0.0	0.0	0.1	1.8	0.961	0.979
*Butyrivibrio*_2_unclassified	0.8	4.8	7.0	4.0	5.9	6.9	1.4	0.147	0.422
Uncultured_*Lachnospiraceae*_bacterium	0.6	0.4	0.5	2.5	0.9	0.5	0.7	0.757	0.924
Uncultured_*Clostridium_sp*	0.6	0.1	0.1	0.1	0.0	0.1	0.4	0.481	0.757
*Oribacterium*_unclassified	0.5	0.4	0.4	2.0	2.4	0.4	0.3	0.778	0.924
*Treponema*_2_unclassified	0.4	1.6	3.2	0.6	1.2	1.4	0.5	0.024	0.137
Uncultured_rumen_bacterium_3C0d_9	0.3	0.3	0.3	0.1	0.1	0.0	0.2	0.979	0.979
*Prevotella_sp*_CA17	0.3	0.1	0.1	0.2	0.1	0.2	0.2	0.269	0.527
*Streptococcus*_unclassified	0.3	1.1	0.0	0.2	0.0	0.0	0.1	<0.001	<0.001
*Selenomonas_ruminantium*_AB3002	0.2	0.5	0.1	0.2	0.2	0.0	0.1	0.010	0.073
bacterium_AC2043	0.2	0.3	0.2	0.2	0.2	0.1	0.2	0.270	0.527
Butyrivibrio_fibrisolvens	0.2	4.4	1.2	1.5	1.1	0.8	0.2	0.005	0.042
Bacterium_XPD3003	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.955	0.979
*Lachnospiraceae*_unclassified	0.2	0.2	0.4	0.3	0.3	0.3	0.1	0.071	0.263
*Lachnospiraceae*_bacterium_AC2012	0.2	1.4	0.3	0.8	0.2	0.1	0.2	<0.001	0.004
Uncultured_bacterium_unclassified	0.2	0.0	0.3	0.1	0.1	0.1	0.1	0.032	0.142
Unidentified_rumen_bacterium_RFN66	0.2	0.1	0.0	0.3	0.2	0.0	0.1	0.004	0.037
*Selenomonas*_1_unclassified	0.2	0.7	0.1	0.1	0.2	0.1	0.1	0.046	0.181
Unidentified_rumen_bacterium_RC11	0.2	0.2	0.2	0.1	0.1	0.1	0.2	0.735	0.924
*X_Eubacterium_coprostanoligenes*_group_unclassified	0.2	0.1	0.2	0.1	0.0	0.1	0.1	0.848	0.953
Ruminococcus_albus	0.1	0.1	0.1	0.0	0.1	0.1	0.3	0.694	0.911
Unidentified_rumen_bacterium_RFN89	0.1	0.1	0.1	0.2	0.1	0.0	0.0	0.617	0.845
Unidentified_rumen_bacterium_RFN13	0.1	0.1	0.1	0.1	0.2	0.2	0.0	0.042	0.176
*X_Eubacterium_hallii*_group_unclassified	0.1	0.2	0.1	0.1	0.1	0.0	0.1	0.085	0.282
*Bacteroidales*_BS11_gut_group_unclassified	0.1	0.1	0.1	0.0	0.0	0.0	0.1	0.820	0.939
Human_gut_metagenome	0.1	0.1	0.2	0.0	0.1	0.1	0.1	0.643	0.861
Uncultured_prokaryote	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.440	0.710
*Succinivibrio*_unclassified	0.1	0.8	0.1	0.1	0.1	0.1	0.0	0.000	0.001
*Roseburia*_unclassified	0.1	0.8	0.2	1.1	0.4	0.5	0.1	0.002	0.019

Bacterium_XPB1013	0.1	0.3	0.4	0.2	0.8	0.5	0.3	0.293	0.527
*Lachnospiraceae*_bacterium_NK4A144	0.1	0.7	0.3	0.3	0.5	0.1	0.1	0.023	0.137
*Ruminococcus*_1_unclassified	0.1	0.0	0.0	0.0	0.1	0.1	0.0	0.164	0.430
*Bacteroidales*_S24_7_group_unclassified	0.1	0.2	0.4	0.3	0.3	0.2	0.0	0.792	0.924
Uncultured_rumen_bacterium_3C0d_20	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.615	0.845
Unidentified_rumen_bacterium_RF31	0.1	0.5	0.2	0.4	0.2	0.2	0.1	0.002	0.019
*X_Eubacterium_ventriosum*_group_unclassified	0.1	0.1	0.1	0.0	0.0	0.1	0.1	0.900	0.979
*Lachnospiraceae*_NK3A20_group_unclassified	0.1	0.1	0.2	0.1	0.0	0.1	0.1	0.349	0.611
X_Clostridium_aminophilum	0.1	0.0	0.0	0.1	0.2	0.1	0.0	0.272	0.527
*Bacteroidetes*_unclassified	0.1	0.2	0.4	0.1	0.7	0.7	0.1	0.176	0.444
*Mollicutes*_unclassified	0.1	1.8	2.4	0.2	0.3	1.4	0.0	0.289	0.527
*Saccharofermentans*_unclassified	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.361	0.615
*Anaeroplasma*_unclassified	0.0	1.5	1.7	0.3	0.2	1.4	0.0	0.279	0.527
*Bacteroidales*_unclassified	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.289	0.527
Unidentified_rumen_bacterium_RC17	0.0	0.1	0.1	0.1	0.1	0.1	0.2	0.951	0.979
*X_Eubacterium_rectale*_group_unclassified	0.0	0.0	0.1	0.1	0.0	0.2	0.1	0.760	0.924
*Treponema_sp*_S	0.0	0.2	0.2	0.1	0.2	0.6	0.1	0.541	0.799
Unidentified	0.0	3.0	0.3	0.1	0.7	0.3	0.0	0.249	0.527
*X_Eubacterium_cellulosolvens*_6	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.605	0.845
Bacterium_ND2006	0.0	0.2	0.1	0.1	0.2	0.0	0.0	0.026	0.138
*Porphyromonadaceae*_unclassified	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.429	0.710
*Prevotella_bryantii*_B14	0.0	0.1	0.1	0.3	0.1	0.3	0.0	0.779	0.924

[1]Control: Natural grassland hay based on *Dichanthium spp*. harvested in Guadeloupe, French West Indies

[2]Differential abundance was estimated with the metagenomeSeq package (Paulson et al., 2013)

Supplementary Table 4. Proportional abundance of bacterial families colonising tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 3 h

	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	P value[2]	FRD
Prevotellaceae	28.9	24.2	30.6	26.2	23.0	37.7	27.5	0.042	0.147
Lachnospiraceae	15.6	12.0	21.2	30.2	41.8	16.9	14.0	0.010	0.045
Christensenellaceae	15.3	15.9	10.9	14.1	7.5	15.6	20.2	0.635	0.846
Fibrobacteraceae	11.5	2.1	3.9	5.2	3.5	8.1	12.9	0.005	0.030
Ruminococcaceae	7.3	8.0	6.0	5.2	3.8	5.9	9.4	0.911	0.966
Streptococcaceae	4.1	0.2	0.1	0.8	0.3	0.1	0.5	0.916	0.966
Rikenellaceae	4.0	4.9	4.7	2.2	2.5	2.0	2.3	0.626	0.846
Acidaminococcaceae	3.5	1.5	2.5	6.4	2.3	3.7	5.3	0.042	0.147
*Bacteroidales*_BS11_gut_group	1.7	3.1	2.1	1.4	1.5	1.0	1.4	0.762	0.927
Spirochaetaceae	1.6	1.8	3.3	1.5	2.4	3.4	1.6	0.017	0.069
Coriobacteriaceae	1.5	6.5	2.5	1.7	1.2	0.9	1.8	0.165	0.421
*Bacteroidales*_RF16_group	1.3	4.5	4.4	2.0	1.7	1.1	1.0	0.444	0.654
Veillonellaceae	0.7	2.3	2.1	0.7	3.7	0.4	0.4	0.309	0.633
*Bacteroidales*_S24_7_group	0.6	0.5	0.4	0.4	0.3	0.5	0.3	0.369	0.633
*Clostridiales*_vadinBB60_group	0.5	0.2	0.1	0.0	0.1	0.1	0.1	0.074	0.207
Family_XIII	0.4	0.5	0.5	0.4	0.3	0.2	0.2	0.758	0.927
Rhodospirillaceae	0.3	0.8	2.3	0.4	0.3	0.2	0.2	0.004	0.030
Anaeroplasmataceae	0.3	0.6	0.6	0.3	0.3	0.5	0.2	0.407	0.633
Anaerolineaceae	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.917	0.966
*Bacteroidales*_UCG_001	0.2	0.9	0.3	0.1	0.3	0.1	0.2	0.287	0.633
Succinivibrionaceae	0.2	0.6	0.4	0.3	0.9	0.2	0.1	0.392	0.633
Erysipelotrichaceae	0.1	0.3	0.2	0.0	0.1	0.0	0.1	0.370	0.633
*Bacteroidetes*_unclassified	0.1	0.1	0.2	0.2	0.2	0.3	0.1	0.966	0.966
*Bacteroidales*_unclassified	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.916	0.966
*Mollicutes*_unclassified	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.392	0.633

[1]Control: Natural grassland hay based on *Dichanthium spp*. harvested in Guadeloupe, French West Indies

[2]Differential abundance was estimated with the metagenomeSeq package (Paulson et al., 2013)

Supplementary Table 5. Proportional abundance of bacterial families colonising tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 12 h

	Control[1]	Acacia nilotica	Calliandra calothyrsus	Gliricidia sepium	Leucaena leucocephala	Manihot esculenta	Musa spp	P value[2]	FRD
Fibrobacteraceae	26.9	2.7	5.9	11.1	13.9	11.0	23.1	0.022	0.099
Prevotellaceae	20.8	24.5	21.5	21.8	28.0	27.8	24.7	0.571	0.727
Christensenellaceae	12.7	9.9	7.3	7.7	7.3	6.3	12.9	0.112	0.253
Lachnospiraceae	12.5	39.6	30.7	39.9	29.8	22.3	14.5	0.309	0.504
Ruminococcaceae	8.2	2.7	4.9	3.3	3.1	4.6	9.7	0.004	0.021
Rikenellaceae	7.7	2.4	7.0	2.5	2.7	2.8	2.6	0.002	0.015
Acidaminococcaceae	2.4	0.9	1.1	3.2	2.1	1.5	2.5	0.670	0.787
Spirochaetaceae	1.9	5.1	7.0	3.2	5.3	13.9	7.2	0.940	0.976
*Clostridiales*_vadinBB60_group	1.4	0.1	0.9	0.2	0.4	0.1	0.2	<0.001	<0.001
*Bacteroidales*_S24_7_group	1.2	1.7	3.1	2.2	2.5	2.0	0.6	0.044	0.133
Anaeroplasmataceae	0.7	3.2	4.8	2.0	1.9	4.2	0.2	0.069	0.169
Coriobacteriaceae	0.6	0.3	0.4	0.4	0.1	0.2	0.3	0.420	0.629
*Bacteroidales*_BS11_gut_group	0.6	0.3	0.7	0.5	0.2	0.3	0.3	0.027	0.102
Veillonellaceae	0.5	1.3	0.3	0.3	0.5	0.2	0.2	0.030	0.102
Family_XIII	0.4	0.2	0.3	0.3	0.1	0.1	0.2	0.708	0.796
*Bacteroidales*_RF16_group	0.3	0.3	0.4	0.6	0.5	0.2	0.2	0.521	0.727
Streptococcaceae	0.3	1.2	0.0	0.2	0.0	0.0	0.1	0.000	0.000
*Bacteroidales*_UCG_001	0.2	0.1	0.1	0.0	0.0	0.0	0.2	0.989	0.989
Rhodospirillaceae	0.2	0.1	0.3	0.1	0.1	0.1	0.1	0.317	0.504
Anaerolineaceae	0.2	0.0	0.2	0.1	0.1	0.1	0.1	0.845	0.912
Succinivibrionaceae	0.1	0.8	0.1	0.1	0.1	0.1	0.0	<0.001	<0.001
*Bacteroidetes*_unclassified	0.1	0.2	0.4	0.1	0.7	0.7	0.1	0.153	0.317
Erysipelotrichaceae	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.547	0.727
*Mollicutes*_unclassified	0.1	2.0	2.5	0.2	0.3	1.4	0.0	0.592	0.727
*Bacteroidales*_unclassified	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.237	0.456

[1]Control hay: Natural grassland hay based on *Dichanthium spp*. harvested in Guadeloupe, French West Indies

[2]Differential abundance was estimated with the metagenomeSeq package (Paulson et al., 2013)

Supplementary Table 6. Proportional abundance of archaea colonising tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 3 h

	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	P value[2]	FRD
*Methanobrevibacter_gottschalkii_*clade	61.4	65.1	63.0	65.3	66.6	65.5	69.6	0.929	0.929
Not_Assigned	17.5	14.3	13.9	17.9	15.1	12.9	11.2	0.772	0.836
Group9_sp	9.2	8.7	10.8	8.8	5.3	7.7	8.6	0.406	0.627
*Methanobrevibacter_ruminantium_*clade	6.2	4.6	6.0	4.1	4.8	5.3	5.9	0.434	0.627
Group10_sp	2.8	1.4	2.7	1.4	7.1	5.5	1.9	0.054	0.213
*Methanosphaera_sp_*ISO3_F5	2.0	4.1	1.9	1.4	1.0	1.8	1.6	0.065	0.213
Methanosphaera_stadtmanae	0.3	0.3	0.4	0.2	0.0	0.2	0.7	0.554	0.655
Methanobrevibacter_oralis	0.2	0.4	0.3	0.1	0.1	0.2	0.2	0.334	0.621
Methanosphaera_cuniculi	0.2	0.4	0.1	0.1	0.0	0.1	0.2	0.008	0.098
Methanobrevibacter_smithii	0.1	0.0	0.1	0.1	0.0	0.1	0.1	0.331	0.621
Methanomicrobium_mobile	0.1	0.5	0.7	0.3	0.0	0.1	0.0	0.232	0.604
Methanobacterium_alkaliphilum	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.038	0.213

[1]Control: Natural grassland hay based on *Dichanthium spp*. harvested in Guadeloupe, French West Indies

[2]Differential abundance was estimated with the metagenomeSeq package (Paulson et al., 2013)

Supplementary Table 7. Proportional abundance of archaea colonising tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 12 h

	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	P value[2]	FRD
*Methanobrevibacter_gottschalkii_*clade	40.8	53.8	40.8	53.9	50.4	52.6	58.0	0.673	0.828
Not_Assigned	24.0	17.5	21.9	15.8	16.8	13.3	16.6	0.776	0.887
Group9_sp	11.5	7.2	11.5	10.1	11.6	12.9	9.9	0.246	0.636
*Methanobrevibacter_ruminantium_*clade	11.3	9.8	11.8	8.2	9.4	8.6	8.0	0.585	0.813
Group10_sp	9.1	6.0	9.4	9.1	8.8	9.3	4.7	0.077	0.636
Methanosphaera_sp_ISO3_F5	1.3	3.6	2.2	2.1	1.4	2.1	1.5	0.429	0.765
Group11_sp	0.7	0.2	0.5	0.1	0.3	0.6	0.0	0.928	0.928
Methanosphaera_stadtmanae	0.5	0.7	1.0	0.2	0.8	0.1	0.3	0.461	0.765
Methanobrevibacter_oralis	0.3	0.4	0.5	0.3	0.2	0.3	0.3	0.478	0.765
Methanomicrobium_mobile	0.2	0.5	0.2	0.1	0.0	0.0	0.3	0.874	0.928
Methanobrevibacter_smithii	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.190	0.636
Group8_sp	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.166	0.636
Methanosphaera_cuniculi	0.1	0.2	0.1	0.0	0.0	0.1	0.1	0.115	0.636
Methanobacterium_alkaliphilum	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.610	0.813

[1]Control: Natural grassland hay based on *Dichanthium spp*. harvested in Guadeloupe, French West Indies

[2]Differential abundance was estimated with the metagenomeSeq package (Paulson et al., 2013)

Supplementary Table 8. Proportional abundance of protozoa associated to tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 3 h

	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	P value	FRD
Uncultured_rumen_protozoa_unclassified	63.8	28.7	51.2	51.9	60.2	79.8	79.6	0.029	0.113
*Isotricha_prostoma*_unclassified	20.8	63.1	30.2	32.6	18.3	7.5	13.6	<0.001	0.001
*Diplodinium*_unclassified	3.1	0.2	1.1	1.5	3.0	2.7	0.3	0.388	0.657
*Trichostomatia*_unclassified	3.1	0.7	2.9	2.1	2.7	2.4	0.8	0.802	0.935
Uncultured_protist_unclassified	2.4	0.5	1.6	1.2	1.2	1.9	2.8	0.813	0.935
*Eudiplodinium_maggii*_unclassified	2.4	1.0	4.0	3.2	6.0	1.9	0.2	0.870	0.953
*Ophryoscolex*_unclassified	1.5	0.2	1.2	1.4	1.7	1.4	0.3	0.376	0.657
*Isotricha_intestinalis*_unclassified	1.3	3.3	1.0	1.1	1.3	1.0	2.0	<0.001	<0.001
*Dasytricha_ruminantium*_unclassified	0.9	0.5	1.2	1.5	0.2	0.6	0.3	0.206	0.475
*Isotricha*_unclassified	0.7	2.0	5.6	3.3	5.5	0.7	0.2	0.095	0.273

[1]Control: Natural grassland hay based on *Dichanthium spp*. harvested in Guadeloupe, French West Indies

[2]Differential abundance was estimated with the metagenomeSeq package (Paulson et al., 2013)

Supplementary Table 9. Proportional abundance of protozoa associated to tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 12 h

	Control[1]	*Acacia nilotica*	*Calliandra calothyrsus*	*Gliricidia sepium*	*Leucaena leucocephala*	*Manihot esculenta*	*Musa spp*	P value	FRD
Uncultured_rumen_protozoa_unclassified	62.9	53.4	36.2	36.2	55.8	65.7	66.6	0.005	0.046
*Eudiplodinium_maggii*_unclassified	17.2	10.9	14.6	14.6	16.9	8.2	3.8	0.787	0.871
*Isotricha_prostoma*_unclassified	10.6	29.3	33.1	33.1	19.3	10.2	21.9	0.046	0.208
*Diplodinium*_unclassified	3.3	0.3	1.2	1.2	0.4	4.2	0.4	0.663	0.871
*Trichostomatia*_unclassified	3.0	1.3	2.4	2.4	2.2	4.4	1.0	0.401	0.871
*Isotricha_intestinalis*_unclassified	1.1	2.2	3.5	3.5	1.0	1.2	2.5	0.588	0.871
Uncultured_protist_unclassified	0.8	0.8	1.2	1.2	0.0	3.3	3.5	0.848	0.871
*Ophryoscolex*_unclassified	0.6	0.2	0.2	0.2	0.2	0.9	0.1	0.871	0.871
*Isotricha*_unclassified	0.5	1.5	7.3	7.3	3.7	1.3	0.3	0.273	0.819

[1]Control: Natural grassland hay based on *Dichanthium spp*. harvested in Guadeloupe, French West Indies

[2]Differential abundance was estimated with the metagenomeSeq package (Paulson et al., 2013)

Supplementary Fig. 1. Abundance of anaerobic fungal taxa (ITS amplicon) colonising tropical tannin-rich plants incubated in the rumen of cows (n= 3) for 3 and 12 h

Supplementary Figure 2. Relationship between chemical composition and degradability in the rumen of tropical tannin-rich plants and their associated microbial communities after 3 and 12 h of incubation.

● = control; ● = *Acacia nilotica*; ● = *Calliandra calothyrsus*; ● = *Gliricidia sepium*; ● = *Leucaena leucocephala*; ● = *Manihot esculenta*; ● = *Musa spp*. Variables included in the analysis were NDF, ADF and N content; DM, NDF and N degradability (Deg) after 3 or 12 h of incubation; DM, N and NDF theoretical degradability (DT); contents in hydrolysable (HT) and total condensed tannins (TCT) and their fractions (free for free CT, FCT for fibre-bound CT, PCT for protein-bound CT); disappearance (dis) of total condensed tannins and their fractions. Variables significantly correlated are depicted in the graphs.

Supplementary Fig. 3. Contribution of variables of tropical tannin-rich plants and variables obtained in vitro and in situ.

Abscissa and intercept represent the two main components accounting for 49 and 21% of total variability, respectively. Values in the axes are eigenvalues of the correlation matrix. Observations are average values for each tannin-rich plant (n = 6).

Variables of feed characterisation: NDF: neutral detergent fiber; FbCT: fibre-bound condensed tannins; FrCT: free condensed tannins; PCT: protein-bound condensed tannins; TCT: total condensed tannins; HT: hydrolysable tannins; IVD: *in vitro* digestibility. Variables from Exp. 1: DMD: dry matter degradability; CPD: crude protein degradability; FbCTD: fibre-bound condensed tannins disappearance; PCTD: protein-bound condensed tannins disappearance; TCTD: total condensed tannins disappearance. Variables from Exp. 2: VFA: total volatile fatty acid production; A:P: acetate:propionate ratio; AB:P: (acetate + butyrate):propionate ratio; B: butyrate, % total volatile fatty acids; CH4: methane production; Gas: total gas production; FOM: fermented organic matter.