BubR1 recruitment to the kinetochore via Bub1 enhances Spindle Assembly Checkpoint

signaling

Anand Banerjee[1,*,#], Chu Chen[2,*], Lauren Humphrey[2], John J. Tyson[1], Ajit P. Joglekar[2,#]

1 – Academy of Integrated Science, Virginia Polytechnic Institute & State University, Blacksburg,

VA 24601, USA

2 – Cell & Developmental Biology, University of Michigan Medical School, Ann Arbor, MI 48109,

USA

* - These authors contributed equally.

[#] – corresponding authors

Abstract

During mitosis, unattached kinetochores in a dividing cell generate the anaphase-inhibitory Mitotic

Checkpoint Complex (MCC) to activate the Spindle Assembly Checkpoint (SAC) and delay

anaphase onset. To generate MCC, these kinetochores recruit MCC constituent proteins

including the protein BubR1. The increased local concentration of BubR1 resulting from this

recruitment should enhance MCC generation, but prior studies found this not to be the case. We

analyzed the contribution of two BubR1 recruitment pathways to MCC generation in human

kinetochores. For these analyses, we isolated a subset of the MCC generation reactions to the

cytosol using ectopic SAC activation systems. These analyses and mathematical modeling show

that BubR1 binding to the SAC protein Bub1, but not to the 'KI' motifs in the kinetochore protein

Knl1, significantly enhances the rate of Bub1-mediated MCC generation in the kinetochore. Our

work also suggests that Bub1-BubR1 stoichiometry will strongly influence the dose-response

characteristics of SAC signaling.

Keywords: mitosis, Spindle Assembly Checkpoint, eSAC, kinetochores, mathematical modeling

Introduction

The "Spindle Assembly Checkpoint" (SAC) is a cell cycle control that minimizes chromosome missegregation during cell division (Musacchio, 2015). It is activated by kinetochores that are not stably attached to the plus-ends of spindle microtubules. These kinetochores generate a diffusible anaphase-inhibitory signal in the form of the Mitotic Checkpoint Complex (MCC), which in turn delays anaphase onset to avert cell division. The amount of MCC in the dividing cell controls the delay in anaphase onset. This amount depends on the number of unattached kinetochores and the rate at which each kinetochore generates MCC (Collin et al., 2013; Dick and Gerlich, 2013). Therefore, the latter parameter is a key determinant of the delay in anaphase onset and chromosome segregation accuracy (Roy et al., 2020).

The rate at which an unattached kinetochore generates MCC depends on its ability to recruit MCC components Mad2, and Cdc20 (Vleugel et al., 2015; Vleugel et al., 2013) (Figure 1A, dashed square). Upon their recruitment to unattached kinetochores, these proteins interact to generate either the MCC itself (or the C-Mad2-Cdc20 subcomplex). Recent discoveries show that Bub1 plays a major role in coordinating the formation of the C-Mad2-Cdc20 subcomplex, which is a crucial, but rate-limiting step in MCC formation (Faesen et al., 2017; Lara-Gonzalez et al., 2021; Piano et al., 2021). Bub1 also heterodimerizes with BubR1 (Overlack et al., 2015). Given this knowledge, it is reasonable to expect that BubR1 recruitment to the kinetochore will increase the rate of MCC generation. Interestingly, however, this expectation does not appear to hold. Disruption of BubR1 recruitment to unattached kinetochores does not reduce the duration of SAC-induced mitotic arrest in nocodazole-treated cells (Overlack et al., 2015; Zhang et al., 2016). This finding is counter-intuitive, because decreased local BubR1 concentration in the kinetochore should reduce the rate of MCC formation by the kinetochore, and therefore lessen the delay in anaphase onset. A possible explanation for this finding is that BubR1 recruits Protein Phosphatase 2A (PP2A) to the kinetochore, which indirectly promotes SAC silencing (Foley et al., 2011; Nijenhuis et al., 2014).

In this study, we re-examine the contribution of kinetochore recruitment of BubR1 to SAC signaling in HeLa cells. To overcome the challenges associated with studying SAC signaling within kinetochores, we use the ectopic SAC activation system or the 'eSAC' to isolate specific signaling interactions in the cytosol (Chen et al., 2019). We find that Bub1-BubR1 heterodimerization elevates Bub1-mediated MCC generation. On the other hand, the recruitment of Bub1 and BubR1 via the 'KI' motifs in the kinetochore protein Knl1 does not contribute to MCC generation mediated by the MELT motifs within the Knl1 phosphodomain (Bolanos-Garcia et al.,

2011a; Bolanos-Garcia et al., 2011b; Kiyomitsu et al., 2011; Krenn et al., 2012; Primorac et al.,

2013; Vleugel et al., 2013; Zhang et al., 2014). We also establish a mathematical model to

elucidate the mechanistic details of the SAC signaling cascade that generates the MCC. We

confirm these findings in the context of kinetochore-based SAC signaling.

Results

The BubR1 binding domain of Bub1 promotes Bub1-mediated MCC assembly

BubR1 is recruited to the human kinetochore by: (1) its heterodimerization with Bub1, which itself

binds to the kinetochore protein Knl1 as part of the Bub1-Bub3 complex, and (2) its binding to the

'KI motif' located within the N-terminal domain of Knl1 (Figure 1A) (Bolanos-Garcia et al., 2011b;

Kiyomitsu et al., 2011; Krenn et al., 2014; Leontiou et al., 2019; Nijenhuis et al., 2013). We first

examined whether Bub1-BubR1 heterodimerization contributes to Bub1-mediated MCC

assembly. Conventional assays are ill-suited to directly address this question, because both Bub1

and BubR1 recruit regulatory enzymes to the kinetochore. Consequently, any disruption of the

recruitment of the SAC proteins to the kinetochore is bound to perturb the intricate regulation of

signaling activities within the kinetochore. Therefore, we first decided to employ the ectopic SAC

activation system (the eSAC) to isolate Bub1-mediated MCC generation in the cytoplasm and

then quantitatively dissect the role of Bub1-BubR1 heterodimerization in MCC assembly.

We and others have previously found that the induced dimerization of a fragment of the central

domain of Bub1 with Mps1 delays anaphase onset in HeLa cells, budding yeast, and fission yeast

(Chen et al., 2019; Leontiou et al., 2019; Yuan et al., 2017). In HeLa cells, induced dimerization

of the central domain of Bub1 (Bub1$^{231-620}$-mNG-2xFkbp12, diagram in Figure 1B) with the Mps1

kinase domain (Frb-mCherry-Mps1$^{500-817}$) produced a dose-dependent delay in anaphase onset

with a maximal duration of ~ 145 minutes (Figure 1B replotted for comparison from (Chen et al.,

2019)). To test whether Bub1-BubR1 heterodimerization contributes to Bub1-mediated MCC

assembly, we performed dose-response analysis on a truncated Bub1 phosphodomain that lacks

the BubR1 heterodimerization domain (Bub1$^{444-620}$-mNG-2xFkbp12) (Overlack et al., 2015).

Ectopic SAC activation using this truncated Bub1 phosphodomain elicited a significantly weaker

response, with a maximal mitotic duration of ~ 105 minutes (Figure 1C, plateau of the 4-parameter

sigmoid predicted as 105 ± 6 minutes, the range indicates 95% confidence intervals, see Materials

and Methods for details). It should be noted that the removal of the Bub3-binding GLEBS domain

in the truncated phosphodomain is not responsible for the observed decrease in the maximal

mitotic delay induced by the eSAC activity (Roy et al., 2021). Thus, Bub1-BubR1
heterodimerization contributes to the ability of the Bub1 phosphodomain to generate MCC. A
similar observation was reported by a recent *in vitro* study (Piano et al., 2021).

To understand whether a mechanism based on local BubR1 enrichment can explain the dose-
response data, we constructed a numerical model that simulates key steps in MCC generation by
the eSAC system and the effect of this MCC on the metaphase-to-anaphase transition (see
Materials and Methods). This model consists of two stages. In the first stage, we calculate the
equilibrium concentrations of two types of signaling complexes that the Bub1 phosphodomain
(Bub1$^{231\text{-}620}$) can form: one that contains both BubR1 and Mad1 and one containing only Mad1
(Figure 1D, left; phosphodomains that do not bind Mad1 cannot generate MCC). BubR1 interacts
with the Bub1 phosphodomain regardless of its phosphorylation state, whereas Mad1 interacts
only with the phosphorylated phosphodomain (Faesen et al., 2017; Ji et al., 2017). Therefore, the
latter interaction depends on the amount of Frb-mCherry-Mps1$^{500\text{-}817}$, i.e., the eSAC dosage. We
assume that the cellular concentration of the eSAC phosphodomain is higher than both BubR1
and Mad1 concentration (this is experimentally confirmed later). Consequently, the number of
signaling complexes saturates at high eSAC dosage (Figure S1A). Cdc20 recruitment to either
signaling complex is assumed to be independent of phosphorylation as well as BubR1 or Mad1
binding t*o* the phosphodomain (not depicted).

The signaling complex containing Bub1, BubR1, Mad1, and Cdc20 facilitates the formation of
either the Mad2-Cdc20 complex or the MCC. Since the rate-limiting step in MCC assembly is the
formation of Mad2-Cdc20, we choose the same rate constant for the formation of MCC and Mad2-
Cdc20 (Faesen et al., 2017). We assume that a newly formed Mad2-Cdc20 at the phosphodomain
can combine with BubR1 to form the MCC with probability *p*, or escape to the cytoplasm with
probability 1−*p*. Thus, the rates of MCC and Mad2-Cdc20 formation at the Bub1 phosphodomain
are given by:

$$\text{MCC formation rate at Bub1}^{231\text{-}620} = p \cdot k_{\text{asmcc2}} \cdot [\text{Mad2}]_{\text{I}} \cdot [\text{Cdc20-Bub1-BubR1}]$$

$$\text{Mad2-Cdc20 formation rate at Bub1}^{231\text{-}620} \text{or Bub1}^{444\text{-}620}$$

$$= (1-p) \cdot k_{\text{asmcc2}} \cdot [\text{Mad2}]_{\text{I}} \cdot [\text{Cdc20-Bub1-BubR1}],$$

where [Mad2]$_\text{I}$ is the concentration of inactive (open) form of Mad2 in the cytoplasm and [Cdc20
-Bub1-BubR1] is the concentration of the complex between Bub1, BubR1, and Cdc20. This
simplified reaction scheme assumes MCC formation to be a one-step reaction.

The important question is whether Bub1-BubR1 heterodimerization promotes MCC formation at
the Bub1 phosphodomain. If we assume that it does not contribute to MCC formation (or in the
case of Bub1[444-620], which lacks the BubR1 binding site), then only Mad2-Cdc20 will be produced
at the phosphodomain. This sub-complex must interact with BubR1 in the cytosol to complete
MCC formation. We calculate this rate as:

$$\text{MCC formation rate in the cytoplsam} = k_{\text{asmcc1}} \cdot [\text{BubR1}]_{\text{f}} \cdot [\text{Mad2-Cdc20}],$$

where $[\text{BubR1}]_{\text{f}}$ is the concentration of free BubR1 in the cytoplasm. To simulate the effect of
MCC generation by these two mechanisms on mitotic progression, we modified a previously
described model of the metaphase-to-anaphase transition (Chen et al., 2019; He et al., 2011)
(Methods, also see Figure S1B).

We first simulated the dose-response curve for Bub1[444-620] by assuming a suitable value for the
k_{asmcc1} parameter (assumed protein concentrations and rate constants are listed in Table S1).
Next, we simulated dose-response data for Bub1[231-620]. When we assumed that BubR1 binding to
the kinetochore does not affect the rate of MCC generation, i.e., $p = 0$ in the above equations,
the simulation generated a weaker dose-response curve with a lower maximal response (dotted
blue line in Figure 1E). The response is weaker because BubR1 bound to Bub1 cannot participate
in cytosolic MCC assembly, and consequently, the rate of cytoplasmic MCC generation is
reduced. This result is inconsistent with the observed dose-response curves. The nominal choice
of $p = 1$ and $k_{\text{asmcc2}} = 0.5 \, \text{nM}^{-1}\text{min}^{-1}$ resulted in longer delays in anaphase onset - matching
the observed maximal response (Cyclin B dynamics shown in Figure S1C).

A comparison of the total MCC formation rates according to Eq. 1 and 2 shows that the cumulative
MCC formation rate is only modestly higher (~ 12%) when the BubR1 heterodimerization domain
is present in the phosphodomain (Figure S1D). However, this cumulative rate includes
contributions from both types of signaling complexes present in the cell. We wanted to assess the
magnitude of enhancement in MCC generation for each signaling complex. To roughly estimate
this local speed-up, we first observe that, in the case of a full Bub1 phosphodomain, the rate of
MCC formation is proportional to the concentration of Bub1-Cdc20-BubR1 (Figure S1E left).
Similarly, in the case of Bub1[444-620], the rate of MCC assembly in the cytoplasm is roughly
proportional to the Bub1-Cdc20 complex concentration (this complex must bind BubR1 in the
cytoplasm to complete MCC assembly, Figure S1E right). Therefore, upon taking the ratio of the
concentrations of these two complexes, we find that the rate of MCC assembly per
phosphodomain due to Bub1-Cdc20-BubR1 complex is ~ 10 times faster than the effective rate

of MCC assembly due to Bub1-Cdc20 complex (note the difference in the scaling factors used to

display the concentrations of the two complexes in Figure S1E).

156   KI motifs suppress the signaling strength of the eSAC phosphodomain

The KI motifs, so named because of the critical role that Lysine and iso-Leucine residues play in

their activities, also recruit Bub1 and BubR1 to the human kinetochore (Figure 1A). The first KI

motif (KI1) is thought to exclusively bind Bub1, whereas the second motif (KI2) is thought to bind

BubR1 (Krenn et al., 2012). A prior study concluded that the KI motifs cooperate with the first

MELT motif in KNL1 to strengthen SAC signaling (Krenn et al., 2014). However, it is unclear

whether this is because Bub1 and BubR1 recruited by the KI motifs participate in MCC assembly

or because they recruit regulatory enzymes (e.g., PLK1, PP2A) that promote SAC signaling

(Nijenhuis et al., 2014). Therefore, to test whether Bub1 and BubR1 bound to the KI motifs directly

promote MCC assembly and to delineate the roles of the two KI motifs, we developed another

eSAC system. In this system, a fragment of Knl1 spanning the first MELT motif and the two KI

motifs serves as the phosphodomain (Knl1$^{160-256}$, Figure 2A). Additionally, we created a variant

phosphodomain wherein the two KI motifs are inactivated by mutating the residues identified as

critical for their activity (Krenn et al., 2012). If Bub1 and BubR1 recruitment via the KI motifs

enhances MELT motif activity, this will be apparent as increased signaling strength of the eSAC

phosphodomain.

To establish the basal activity of the first MELT motif, we first studied the dependence of mitotic

duration on the dosage of the eSAC signaling complex consisting of the eSAC phosphodomain

with mutant, inactive KI motifs (M1-KI1*-KI2*-mNG-2xFkbp12, Figure 2A left). As before, the data

was fit with a sigmoidal curve (Figure 2A, middle panel) (Chen et al., 2019). The maximal

response for this phosphodomain was higher than the previously defined maximal response for

the eSAC phosphodomain containing the 12th MELT motif (211 versus 157 minutes with ± 95%

confidence intervals of 169-357 and 152-163 respectively predicted by a 4-parameter sigmoidal

fit to the binned data). This difference likely reflects the difference in the Bub1-Bub3 binding

affinities of the first and twelfth MELT motif (Vleugel et al., 2015). Interestingly, the inclusion of

active KI motifs in the phosphodomain significantly depressed its signaling strength: the eSAC

concentration required for half-maximal response nearly doubled (EC50 = 9.7 and 19.4 a. u.

respectively, Figure 2A middle). Decrease in the apparent signaling strength of the

phosphodomain was eventually compensated by high eSAC dosage, and therefore, the maximal

response was similar in value (Figure 2A middle and right).

Thus, the inclusion of the KI motifs in the eSAC phosphodomain suppressed MCC production
mediated by the phosphorylated MELT motif within the same phosphodomain. Our eSAC-based
analyses of the Bub1 phosphodomain in the prior section provides a possible explanation for this
unexpected observation: if one assumes that the KI motifs interact with Bub1 and BubR1
independently of the MELT motif, and if the Bub1 and BubR1 molecules recruited by the KI motifs
do not participate in MCC formation mediated by the MELT motif, then the KI motifs will effectively
sequester Bub1 and BubR1 from MELT-motif mediated MCC assembly. This will suppress the
signaling activity of the eSAC.

Bub1 and BubR1 interactions with the KI motifs do not require Mps1-mediated
phosphorylation of KNL1; Bub1 interacts with both KI motifs

To delineate the activities of the KI motifs in the eSAC system, we constructed a new
phosphodomain by fusing an unstructured region of the Knl1 phosphodomain spanning three
previously characterized MELT motifs (Knl1$^{881-1014}$) to the C-terminus of the phosphodomain used
above (Knl1$^{160-256}$, see Figure 2B) (Chen et al., 2019; Vleugel et al., 2015). We also created mutant
versions of this phosphodomain wherein the KI motifs were rendered inactive either individually
or together using point mutations (KI1*-KI2, KI1-KI2*, and KI1*-KI2* with the asterisk denoting
loss of activity, Figure 2B) (Krenn et al., 2012). By incorporating multiple MELT motifs in the
phosphodomain, we wanted to test whether the KI motifs influence the ability of the four MELT
motifs in the phosphodomain to engage in synergistic signaling (Chen et al., 2019).

We first determined whether the two KI motifs interact with Bub1 and BubR1 exclusively, and
whether MELT motif phosphorylation is necessary for these interactions. We immunoprecipitated
the four, mNeonGreen-tagged eSAC phosphodomains from whole cell extracts of mitotic HeLa
cells in the absence of the Mps1 kinase domain using mNeonTrap beads and probed the
precipitates for Bub1 and BubR1 (Methods). When both KI motifs were active (KI1-KI2), Bub1
and BubR1 co-precipitated with the eSAC phosphodomain. When both KI motifs were inactive
(KI1*-KI2*), Bub1 and BubR1 did not co-precipitate with the eSAC phosphodomain as expected
(Figure 2B). Surprisingly, when the first KI motif, which is expected to interact with Bub1 alone,
was inactive (KI1*-KI2), Bub1 still co-precipitated with the phosphodomain albeit at a lower level.
BubR1 coprecipitation was also reduced. This observation suggests that Bub1 competes with
BubR1 to bind to the second KI motif. Consistent with this inference, inactivation of second KI
motif (KI1-KI2*) made BubR1 undetectable in the precipitate and reduced the amount of Bub1.

We obtained similar results from immunoprecipitation experiments involving eSAC
phosphodomains with only one MELT motif and the two KI motifs (Supplementary Figure S2A).
These results show that the first KI motif binds only Bub1, but the second motif can bind Bub1 or
BubR1. Furthermore, these interactions are not dependent on MELT motif phosphorylation.

Bub1 and BubR1 recruited by the KI motifs do not contribute to MCC assembly mediated
by the MELT motifs

Having established the biochemical activities of the KI motifs, we performed dose-response
analysis for the eSAC phosphodomain containing inactive KI motifs to define the baseline activity
of the four MELT motifs (KI1*-KI2* in Figure 2B left, purple circles). The response elicited by the
eSAC phosphodomain with four MELT motifs to increasing eSAC concentration was non-
monotonic and characterized by an initial steep rise that is followed by a gradual decay in the
mitotic duration (residuals from a sigmoidal fit to the dose-response data points is shown in Figure
S2B, left). It should be noted that such non-monotonic response was not observed for a previously
characterized eSAC phosphodomain also containing four MELT motifs (numbers 11-14) (Chen et
al., 2019). The different behaviors of the two phosphodomains may be due to a higher affinity of
the 1^{st} MELT motif compared to the 11^{th} MELT motif for binding Bub3-Bub1 (Vleugel et al., 2015).
Interestingly, when both KI motifs were active, the maximal duration of mitosis was significantly
reduced (~ 87 minutes estimated by 4-parameter sigmoidal fit to the binned averages of the data,
see Figure 2C). Qualitatively similar results were obtained for an extended phosphodomain
containing 7 MELT motifs and the two KI motifs (Figure S2C).

We next determined the dose-response dependence for two eSAC phosphodomains containing
four MELT motifs and only one active KI motif. When only the first KI motif was active (KI1-KI2*,
only Bub1 depleted, see Figure 2B), the dose-response data were monotonic with a slightly lower
maximal response (Figure 2C green circles and curve). When only the second KI motif was active
(KI1*-KI2, Bub1 and BubR1 depleted, Figure 2C), the maximal response was significantly
attenuated (blue circles and line in Figure 2C). Interestingly, the response to this eSAC was non-
monotonic: an initial overshoot was followed by decay to a lower maximal response (residuals
from the sigmoidal fit in Figure S2B, right). These results further show that Bub1 and BubR1
recruited by the two KI motifs do not directly contribute to MCC generation by the MELT motifs
within the same phosphodomain in the eSAC system (Figure 2C). The likely explanation for the
suppression of eSAC signaling activity is that Bub1 and BubR1 recruitment by the KI motifs
sequesters these proteins preventing them from participating in eSAC signaling.

One simple explanation for the strong suppression of eSAC signaling activity by the second KI

motif is that this motif sequesters BubR1 along with Bub1. Since BubR1 is an MCC component,

its sequestration will limit the maximal amount of MCC (MCC_{max}) that the eSAC can generate.

Although we could not quantify the amount of MCC produced by the eSAC, this amount and

MCC_{max} can both be compared qualitatively by studying the duration of mitotic delays. The eSAC

systems discussed here delay mitosis by at most 300 minutes. In contrast, RPE-1 cells treated

with high doses of nocodazole arrest for ~ 1500 minutes on average (Collin et al., 2013; Dick and

Gerlich, 2013). The significantly smaller delay caused by eSAC signaling, when compared to the

delay induced by nocodazole treatment, suggests that the eSAC produces MCC levels smaller

than MCC_{max} (Dick and Gerlich, 2013). Therefore, a reduced MCC_{max} due to BubR1 sequestration

by the second KI motif alone cannot explain the weaker signaling activity.

Numerical simulation of the dose-response data

We returned to numerical simulations to understand the mechanistic underpinnings of the dose-

response data. The suppression of eSAC signaling activity by the KI motifs is in part because the

eSAC phosphodomains are significantly more abundant than Bub1 and BubR1 (shown later).

Nonetheless, the quantitative dose-response data provided us with an excellent opportunity to

delineate the contribution of BubR1 recruitment at the MELT motifs to MCC formation. Bub1 and

BubR1 sequestration will differentially affect the composition of the two signaling complexes:

MELpT-Bub1 and MELpT-Bub1-BubR1 that the eSAC phosphodomains can assemble. If these

signaling complexes assemble the MCC at different rates, their effects on mitotic progression will

also be different (Dick and Gerlich, 2013). Using simulations, we wanted to understand the

mechanisms that give rise to the ~ 1.75-fold difference in the maximal mitotic delay generated by

KI1-KI2* and KI1*-KI2 (Figure 2C).

To understand underlying mechanisms, it is necessary to consider: (a) the effects of the

differential sequestration of Bub1 and BubR1 by the two KI motifs on the concentrations of the

two signaling complexes, and (b) the rates at which these signaling complexes generate MCC.

Both types of signaling complexes will form on the eSAC phosphodomain, because we have

previously shown the eSAC phosphodomain recruits Bub1, BubR1, and Mad1 only in cells treated

with rapamycin (Chen et al., 2019). The stoichiometry of these two signaling complexes depends

on the interplay of two sets of parameters: (1) the stoichiometry of Bub1, BubR1, and the eSAC

phosphodomain (quantification presented after the simulations), and (2) the relative affinities of

Bub1 and BubR1 for the two KI motifs, of Bub1 for phosphorylated MELT motifs, and of BubR1

for Bub1, which are known (Krenn et al., 2014; Vleugel et al., 2015; Zhang et al., 2016).

To simulate the activities of the KI motifs, we added two new elements to our previously described

model of the eSAC (Chen et al., 2019): the two KI motifs, and the factors Bub1 and BubR1

representing Bub1-Bub3 and BubR1-Bub3 respectively (Figure 3A). As before, all interactions are

described by mass action kinetics. In accord with our IP results (Figure 2), the model assumes

that the first KI motif (KI1) interacts only with Bub1, whereas the second KI motif (KI2) can interact

with either Bub1 or BubR1. Based on prior observations, we also assume that only BubR1-Bub3

is recruited by the Bub1 bound to a phosphorylated MELT motif (Overlack et al., 2017; Overlack

et al., 2015). Based on *in vitro* data, the dissociation constant for Bub1 and BubR1 bound to the

KI motifs was assumed to be 6 to 50-fold higher relative to the dissociation constant for Bub1

bound to a phosphorylated MELT motif (Chen et al., 2019; Krenn et al., 2012; Overlack et al.,

2015; Primorac et al., 2013) (rate constants tabulated in Table S2). We assume that the KI motifs

and phosphorylated MELT motifs recruit Bub1 and BubR1 independently. Finally, we also assume

that Bub1 and BubR1 bound to the KI motifs do not participate in the signaling events mediated

by the phosphorylated MELT motifs (Figure 3A). Three of the four MELT motifs used in the eSAC

phosphodomain were used in our previous study. Therefore, their affinities for Bub1 were similar

to the values used in the original simulation (Chen et al., 2019). The rate constants for the first,

12^{th} and 14^{th} MELT motifs were set to reflect that they have a higher affinity for Bub1 compared

to the weaker 13^{th} MELT motif (Vleugel et al., 2015). The protein concentrations used are

tabulated in Table S2.

Effects of the differential activity of the KI motifs on the distribution of the signaling
complexes formed by the eSAC phosphodomains

We first calculated the steady-state concentrations of MELpT-Bub1 and MELpT-Bub1-BubR1

assembled by KI1-KI2* and KI1*-KI2. Bub1-BubR1 stoichiometry plays a critical role in

determining these concentrations. If Bub1-BubR1 stoichiometry is high, e.g., 1:4, nearly every

MELpT-Bub1 complex assembled by KI1-KI2* and KI1*-KI2 will recruit BubR1 to form MELpT-

Bub1-BubR1 despite the differential depletion of Bub1 and BubR1 by the two phosphodomains

(converging dashed lines at high eSAC dosage in Figure 3B, right). Consequently, the

concentration of MELpT-Bub1 is negligible (dashed green and blue lines respectively near the X

axis in Figure 3B, left). At high eSAC dosage, both KI1-KI2* and KI1*-KI2 form the same signaling

complex (MELpT-Bub1-BubR1). Because both phosphodomains form similar amounts of the

same signaling complex, they will generate similar amounts of MCC, and therefore produce
mitotic delays of similar magnitude.

This picture changes when Bub1-BubR1 stoichiometry is around 1:1. In this case, KI1-KI2* and
KI1*-KI2 phosphodomains assemble different equilibrium concentrations of MELpT-Bub1 and
MELpT-Bub1-BubR1 especially at high eSAC dosage. BubR1 depletion by KI1*-KI2 lowers the
amount of MELpT-Bub1-BubR1 that it forms compared to KI1-KI2* (Figure 3B, right, solid lines).
Conversely, the concentration of MELpT-Bub1 formed by KI1*-KI2 is now higher, because BubR1
availability is reduced (solid lines in Figure 3B, left). Figure 3C shows how Bub1-BubR1
stoichiometry affects MELpT-Bub1-BubR1 concentration (Figure 3C left, for the high and constant
eSAC dosage = 20). These calculations show that if the KI1-KI2* and KI1*-KI2 phosphodomains
are to exhibit different dose-response characteristics, BubR1 (i.e., BubR1-Bub3) cannot be in too
much excess of Bub1 (i.e., Bub1-Bub3). Equal Bub1 and BubR1 abundance will produce the
maximal difference in the levels of signaling complexes assembled by KI1-KI2* and KI1*-KI2 at
high eSAC dosage. This difference in the composition of the signaling complexes can provide the
basis for the differential activity of the KI1-KI2* and KI1*-KI2 phosphodomains.

A higher rate of MCC generation by MELpT-Bub1-BubR1 compared to MELpT-Bub1 can
explain the differential behavior of the two phosphodomains

Differences in the maximal concentrations of the MELpT-Bub1 and MELpT-Bub1-BubR1 will
translate into different levels of MCC only if they generate MCC at different rates (the ratio of
maximal mitotic duration achieved by KI1-KI2* and KI1*-KI2 is indicated by the red horizontal line
in Figure 3C). Specifically, MELpT-Bub1-BubR1 must generate MCC at a higher rate than
MELpT-Bub1. To simulate MCC generation by the two signaling complexes, we used a simplified
process of MCC generation wherein we do not explicitly include the dynamics of Mad2, Bub1 and
BubR1. Instead, a single reaction between the signaling complex and free cytoplasmic Cdc20
generates MCC, with the forward rate constants k_{MCC} for MELpT-Bub1 and k'_{MCC} for MELpT-Bub1-
BubR1 as the only parameters. This simplified, coarse-grained scheme is justified because, the
rate of MCC generation at the phosphodomain and in the cytoplasm will be proportional to the
concentration of the corresponding signaling complex (see Figure S1E). To simulate the effect of
the MCC generated in this manner on mitotic progression, we suitably modified the model of
metaphase-to-anaphase transition described by He et al. to reflect the simplified MCC generation
scheme (Figure S3B) (Chen et al., 2019; He et al., 2011).

To study how k_{MCC} and k'_{MCC} affect the maximal mitotic duration, we fixed the eSAC dosage at a

high and constant value, and then calculated the ratio of maximal mitotic duration produced by

KI1-KI2* over that produced by KI1*-KI2 as a function of Bub1-BubR1 stoichiometry and

$k'_{MCC}:k_{MCC}$ ratio (contour plot in Figure 3D). From this plot we see that the simulations of the ratio

of maximal mitotic durations for KI1-KI2* and KI1*-KI2 is 1.4 (close to the observed ratio of ~

1.75), when (1) BubR1-Bub1 stoichiometry is ~ 1, and (2) the signaling strength of the MELpT-

Bub1-BubR1 signaling complex is at least 6-fold higher than the signaling strength of the MELpT-

Bub1 signaling complex.

We used this framework to simulate the dose-response curves for all four phosphodomains. In

this simulation, we assumed [Bub1]:[BubR1] = 1 and $k_{MCC}:k'_{MCC}$ = 0.1 (corresponding to a 10-fold

increase in MCC generation at the eSAC phosphodomain relative to cytoplasmic MCC

generation). With these nominal parameter values, dose-response curves generated by the model

capture key characteristics of the dose-response data (Figure 3E). As before, the assumption of

synergistic signaling is necessary to reproduce the non-monotonicity of the dose-response data

for the phosphodomain containing four MELT motifs (Chen et al., 2019). Without it, the responses

elicited by all phosphodomains become monotonic (Figure S3D). An interesting feature of the

observed dose-response data for the KI1*-KI2 phosphodomain is their non-monotonic nature

(Figure 2E). Our simulations suggest that this difference arises due to Bub1 sequestration by the

KI motif, which in turn suppresses synergistic signaling. Synergistic signaling by MELT motifs

requires high occupancy of MELpT motifs by Bub1-BubR1. For this, the binding between MELpT

and Bub1 is the necessary first step. In the KI1*-KI2 eSAC, Bub1 is not sequestered as strongly.

This allows the assembly of multiple MELpT-Bub1-BubR1 complexes on the same

phosphodomain only at low eSAC concentration. Resultant synergistic signaling becomes

observable as a peak in the dose-response curve. In the case of the KI1-KI2* eSAC, such

synergistic signaling is modulated due to Bub1 sequestration, and cannot be detected. We found

that simply increasing the dissociation constant of KI1-Bub1 binding by a factor of two (thereby

decreasing Bub1 sequestration) made the dose-response curve for KI1-KI2* eSAC visibly non-

monotonic (not shown).

Stoichiometry of Bub1, BubR1, Mad1, and the eSAC phosphodomain in HeLa cells

The simulations show that the stoichiometric ratios among Bub1, BubR1, and the eSAC

phosphodomain are critical determinants of the behavior of the eSAC systems. To experimentally

measure these ratios, we constructed genome-edited HeLa cell lines wherein mNeonGreen

(abbreviated as mNG) was fused to the N-terminus of Bub1 and BubR1, and to the C-terminus of
Mad1 (see Materials and Methods). In all three cases, we obtained clonal cell lines with partially
edited genomes: approximately half of the protein was labeled; the remaining protein fraction was
unlabeled (Figure 4A, also see Methods).

Using mNG fluorescence quantitation, we found that the abundance of the eSAC phosphodomain
containing two active KI motifs and four MELT motifs was 2-4-fold higher than the three SAC
proteins in mitotic cells (Figure 4B). The over-abundance of the eSAC phosphodomain relative to
Bub1, BubR1, and Mad1, and the constitutive activity of the two KI motifs together support our
assumption that the highly expressed eSAC phosphodomains sequester Bub1, BubR1 or both.
Moreover, BubR1 is ~ 2-fold more abundant than both Bub1 and Mad1 (Figure 4C). It should be
noted that this abundance corresponds to the total BubR1 in a cell including free BubR1-Bub3 as
well as BubR1-Bub3 incorporated into MCC (Overlack et al., 2017). The partitioning of BubR1
between the BubR1-Bub3 complex and the MCC remains unknown. Therefore, we conclude that
the concentration of BubR1-Bub3 available for SAC signaling is less than 2-fold higher than that
of Bub1-Bub3. This is consistent with the prediction of our model that the BubR1:Bub1
stoichiometry must be near 1:1 for the observed dose-response data for the KI1-KI2* and KI1*-
KI2 phosphodomains.

Recruitment of BubR1 by BuB1 per se contributes to the signaling strength of the SAC

Observations from the eSAC systems and our numerical simulations forced us to re-examine the
role of Bub1-BubR1 heterodimerization in kinetochore-based SAC signaling. Previous studies
using the knock-down and rescue approach found that the disruption of Bub1-BubR1
heterodimerization increases the SAC strength in nocodazole-treated cells, increasing the
duration of mitotic arrest rather than decreasing it (Krenn et al., 2014; Zhang et al., 2016).
However, we noted three factors that may have influenced this result. First, the deletion of different
segments of BubR1 in the two studies may have resulted in different levels of residual BubR1 at
signaling kinetochores. The second factor arises from the loss of the SAC-antagonizing activity
of PP2A due to the disruption of BubR1 recruitment to the kinetochore. The kinetochore
attachment and regulatory domain (KARD) of BubR1 was either left intact or mutated to eliminate
the binding of the PP2A-recruiting subunit B56α, but not necessarily its other isoforms. Most
importantly, the mutant BubR1 may have been overexpressed in these experiments. Higher levels
of BubR1 in the cytosol will proportionally increase the rate of cytosolic MCC formation, and the
increased rate may mask impaired MCC assembly within kinetochores.

To account for these three factors, we designed and performed new BubR1 knock-down/rescue
experiments. We knocked down endogenous BubR1 in HeLa-A12 using RNA interference and
rescued these cells with mNG-BubR1$^{\Delta665\text{-}682}$ or mNG-BubR1$^{\Delta432\text{-}484,\ \Delta665\text{-}682}$. Both lack the KARD to
interact with PP2A-B56 (ΔKARD), while mNG-BubR1$^{\Delta432\text{-}484,\ \Delta665\text{-}682}$ cannot heterodimerize with
Bub1 (ΔHD, ΔKARD). Therefore, mNG-BubR1$^{\Delta432\text{-}484,\ \Delta665\text{-}682}$ has nearly undetectable localization
at unattached kinetochores (Figure 5A). In these cells, we induced mitotic arrest using nocodazole
treatment following the previous studies. Importantly, we used the genome-edited *mNG-BUBR1*
HeLa-A12 cells treated with control siRNA and imaged under identical conditions to establish the
physiological BubR1expression level. In our analysis of cells expressing the mutant, mNG-tagged
BubR1, we only considered those cells exhibiting mNG intensity that is 0.5–2-times the average
mNG intensity of mitotic mNG-BubR1 HeLa-A12 cells (because only around half of total BubR1
is tagged in the CRISPR-Cas9-edited cell line, Figure 5B, left).

We found that in BubR1 knock-down cells rescued with BubR1 lacking the KARD domain arrested
for a significantly longer duration as compared to control cells. This is likely due to the loss of
PP2A activity from the kinetochores (Saurin et al., 2011). Importantly, cells rescued with BubR1
lacking both the KARD domain and the Bub1 heterodimerization domain arrested for a
significantly shorter amount of time in comparison (Figure 5B, right). Thus, the recruitment of
BubR1 *per se* contributes to the efficient assembly of MCC.

Discussion

Unattached kinetochores are poised to assemble the MCC by recruiting its component subunits:
BubR1-Bub3, Mad2, and Cdc20. Therefore, it was surprising that the disruption of BubR1
recruitment to the kinetochore was found to be unimportant for SAC signaling in nocodazole-
treated cells (Overlack et al., 2015; Zhang et al., 2016). We used the eSAC system to isolate a
subset of the biochemical reactions in the MCC generation cascade from the kinetochore and
analyzed them in the physiological context. These analyses show that Bub1-BubR1
heterodimerization significantly enhances the rate of Bub1-mediated MCC assembly. This
enhancement may be simply due to enrichment of BubR1 at the site of formation of Mad2-Cdc20
(Lara-Gonzalez et al., 2021; Piano et al., 2021), although more complicated mechanisms may
also be at work.

We also find that Bub1 and BubR1 recruitment via the KI motifs does not contribute to Bub1-
mediated MCC assembly. It is important to note that the eSAC signaling complexes contain only
one Knl1 phosphodomain in contrast to the spatially clustered organization of many Knl1
phosphodomains within the kinetochore. Therefore, it remains possible that KI motifs directly or
indirectly prime SAC signaling by constitutively recruiting Bub1 and BubR1 (Jia et al., 2016; Krenn
et al., 2014). Nonetheless, because of the higher affinity of phosphorylated MELT motifs for Bub1-
Bub3 and their superior numbers (~ 19 MELT motifs versus 1 KI motif each per Knl1), the MELT
motifs will take over Bub1 and BubR1 recruitment once Mps1 becomes fully active within
unattached kinetochores. KI motif activity is unlikely to contribute significantly under this condition.

We also constructed a numerical model to simulate the dose-response data using experimental
data wherever possible and assuming plausible parameter values when data were unavailable.
This simple model based on differential sequestration of Bub1 and BubR1 can simulate key
features of the dose-response data. The simulations support the following main conclusions. First,
Bub1-BubR1 heterodimerization significantly elevates the rate of Bub1-mediated MCC
generation. To simulate the observed dose-response data, the rate of MCC assembly due to
Bub1-bound BubR1 was assumed to be 10-fold higher when compared to cytoplasmic MCC
assembly in our model. This enhanced MCC generation rate is reasonable given that the local
concentration of BubR1 will increase by orders of magnitude upon its binding to Bub1, the site
where Mad2-Cdc20 is formed. Second, our model further supports the observation that Bub1 and
BubR1 recruited by the KI motifs do not contribute to MCC generation. Third, the stoichiometry of
Bub1 and BubR1 emerges as a key parameter influencing how the MCC generation output of
each kinetochore will change as a function of the number of signaling kinetochores in the cell.
These simulations thus establish a useful platform for future studies of SAC signaling.

Human kinetochores also use the RZZ pathway to promote MCC assembly, and this pathway can
compensate for a partial loss of Bub1 from the kinetochore (Qian et al., 2017; Silio et al., 2015;
Zhang et al., 2015). Nevertheless, MELT motif and Bub1-mediated MCC assembly is likely to be
critical over the course of a normal cell division. Bub1-mediated recruitment of BubR1 to the
kinetochore is especially important because this will significantly enhance the rate of MCC
generation within the kinetochore, unlike the RZZ pathway, which likely generates only Mad2-
Cdc20 and relies on cytosolic assembly of the MCC. Our study does not clarify the role of the KI
motifs. The possibilities include the 'priming' of SAC signaling in the early stages of mitosis by
enhancing local Bub1 and BubR1 concentration and the recruitment of regulatory factors via Bub1
and BubR1 for kinetochore functions unrelated to SAC signaling.

477
Acknowledgements

We would like to thank Dr. Stephen S. Taylor for the generous gift of antibodies, Dr. Eugene

Makeyev for the HeLa acceptor cell line, and Dr. Michael Lampson for plasmids used in this study.

We also thank members of the Joglekar lab for stimulating discussions. Use of the Incucyte

system was made possible by a generous gift from the Richard Tam Foundation to Prof. Sue

O'Shea (Cell & Developmental Biology, University of Michigan Medical School).

The authors declare that no competing financial interests exist.

Materials and Methods

Plasmid construction

The plasmids used for the stable cell lines were based on plasmids that have been described previously (Chen et al., 2019). Briefly, the phosphodomain was integrated either *Not*I or *Asc*I and *Xho*I restriction sites to create constitutively expressed phosphodomain-mNeonGreen-2xFkbp12. A Mps1$^{500-857}$ fragment corresponding to the Mps1 kinase domain was integrated into the *Fse*I and *Bgl*II restriction sites to create conditionally expressed Frb-mCherry-Mps1$^{500-857}$.

eSAC phosphodomains spanning the 1st MELT motif (M1) and the two KI motifs were created by fusing Knl1$^{160-256}$ to mNeonGreen-2xFkbp12. eSAC phosphodomains containing four MELT motifs and the two KI motifs were created by fusing Knl1$^{160-256}$ to Knl1 fragment Knl1$^{881-1014}$ spanning MELT motifs 12-14, which has been characterized previously (Vleugel et al., 2015). The activity of the first KI motif was disrupted by changing its amino acid sequence from 'K**I**D**TT**S**F**LANLK' to 'K**A**D**AA**S**A**LANLK' (KI1*). Similarly, the activity of the second KI motif was disrupted by mutating its amino acid sequence from 'KIDFND**F**IK**R**LK' to 'KIDFND**A**IK**A**LK' (KI2*) following (Krenn et al., 2012).

DNA repair templates used for CRISPR/Cas9 mediated genome editing were constructed via DNA assembly using the NEB HiFi DNA assembly kit as per manufacturer's instructions. Successfully edited alleles encode mNeonGreen-tagged SAC proteins that separate the corresponding wildtype protein and the fluorescent protein mNeonGreen by a short flexible linker (mNG-BUBR1 and mNG-BUB1: GSGGSG; MAD1-mNG: GGAGGSGG). The sequences of all homology-directed repair template plasmids and Cre-Lox recombination-mediated cassette exchange plasmids are available upon request.

Tissue culture and cell line construction for eSAC analyses

Henrietta Lacks (HeLa) cells were grown in DMEM media supplemented with 10% FBS, 1% Pen/Strep, 1x-Glutamax, and 25 mM HEPES under standard tissue culture conditions (37 °C and 5% CO2). Stable cell lines expressing the two eSAC components were generated by integrating a bi-cistronic eSAC plasmid at engineered *lox* sites in the HeLa genome according to the protocol described in (Khandelia et al., 2011). Upon transfection, DMEM media supplemented with 1 µg/ml Puromycin was used to select transformed cells, and all the colonies were pooled to culture the transformed cells used in the experiments.

To conduct dose-response analysis, each eSAC cell line was plated ~ 40-48 hours prior to the start of the experiment in DMEM media without Puromycin. Doxycycline was added at the time of plating to induce the expression of Frb-mCherry-Mps1. Prior to imaging, the cells were washed

with PBS. Fluorobrite media with 10% FBS, 1% Pen/Strep with or without Rapamycin were added

to each well.

Genome editing HeLa cells using CRISPR/Cas9

The guide RNAs (gRNAs) for in situ BUBR1 and BUB1 N-terminal mNeonGreen-tagging were 5'-

CAGGAUGGCGGCGGUGAAGA-3' and 5'-GGUUCAGGUUUGGCCGCUGC-3', respectively.

The gRNA for in situ MAD1 C-terminal mNeonGreen-tagging was 5'-

CAGACCGUGGCGUAGCCUGC-3'. Single guide RNAs (sgRNAs) were synthesized using the

EnGen sgRNA Synthesis Kit (for the *Streptococcus pyogenes*-originated Cas9, New England

Biolabs). The SpCas9-sgRNA ribonucleoprotein (RNP) complex was assembled at room

temperature in a buffer consisting of 20 mM of HEPES-KOH (pH 7.5), 150 mM of KCl, 1 mM of

$MgCl_2$, ten percent (by volume) of glycerol, and 1 mM of DTT using 100 pmol of SpCas9- 2×NLS

(the QB3 MacroLab) and 120 pmol of sgRNA. The RNP complex and 1.5 µg of a linearized

homology-directed repair template plasmid were transfected into $2×10^5 – 5×10^5$ nocodazole-

arrested mitotic HeLa A12 cells using a Nucleofector and the associated Cell Line Kit R (Lonza)

following manufacturer's instructions. After five weeks, green fluorescence-positive mitotic cells

(arrested by 330 nM of nocodazole for 16 hours) were sorted directly into 96-well plates at 1

cell/well. Healthy colonies were subject to further validation by genotyping and sequencing, as

well as immunoblotting.

For genotyping, HeLa-A12 genomic DNAs were purified using the Wizard SV Genomic DNA

Purification System (Promega). Genotyping primers (BUBR1 forward primer 5'-

CCTGGTCACATCTGAGCTAT-3', BUBR1 reverse primer 5'-CTCAGTGAGACTCCAGTGTT-3',

BUB1 forward primer 5'-CCCTCTACATGAAGGCGCTA-3', BUB1 reverse primer 5'-

GCTCGCCCAAGGTAAACATT-3', MAD1 forward primer 5'-GGACTTTTCAGGGACGTGGT-3',

and MAD1 reverse primer 5'-GAGTTGGGAGGAGGGGACTC-3') were designed to bind outside

of homology arms to avoid false-positive colonies from integration of the homology-directed repair

template plasmid to an off-target genomic locus.

Drug and RNAi treatments

To induce the expression of the mCherry-Frb-Mps1 kinase domain doxycycline was added to a

final concentration of 2 ug/ml (stock concentration 2 mg/ml in DMSO) ~ 48 hours prior to the start

of the experiment. Prior to prior to the start of each experiment, Rapamycin was added ~ 1 hour

of the experiment to a final concentration of 500 nM (stock concentration 500 µM in DMSO) to

induce the dimerization of the eSAC kinase domain with the eSAC phosphodomain. Nocodazole

was added to the final concentration of 330 nM (stock concentration 330 µM in DMSO). The cocktail of siRNA against five different B56 isoforms was added to a final concentration of 40 nM (stock concentration 10 µM). The siRNA sequences were obtained from ref. (Nijenhuis et al., 2014). Cell cycle synchronization was in G1/S was achieved by treating cells with 2.5 mM thymidine (from a 100 mM stock in PBS) for 16-18 hours. Cells were washed with DMEM for release from the G1/S arrest.

Immunoprecipitation

HeLa A12 cells constitutively expressing either MELT1-KI or MELT1-KI-M3 were synchronized at G1/S by 2.5 mM of thymidine. 9 hours after being released from the (double) thymidine block, cells were synchronized in metaphase using 10 µM MG132. After another 1.5 hours, cells were scraped off the plate, washed once with PBS, pelleted, snap-frozen, and stored at −80 °C. Cells were thawed, resuspended in the complete lysis buffer (75 mM HEPES-HCl (pH 7.5), 150 mM KCl, 1.5 mM EGTA, 1.5 mM MgCl2, 10% (by volume) glycerol, and 0.075% (by volume) Nonidet P-40 (AmericanBio); immediately before use one cOmplete protease inhibitor cocktail tablet (EDTA-free, Roche Diagnostics) and a phosphatase inhibitor cocktail (1 mM Na4P2O7, 0.1 mM Na3VO4, 5 mM NaF, and 2 mM sodium β-glycerophosphate) were added), and lysed for 1 hour at 4 °C while rotating. Cell lysates were then centrifuged at 18,000 g for 25 min at 4 °C. The supernatant was subsequently cleared by control agarose beads for 1.5 hours at 4 °C to reduce non-specific binding. Cleared supernatant was then mixed with mNeonGreen-Trap agarose beads (ChromoTek) and rotated for 1.5 hours at 4 °C. Beads were washed 4 times using the lysis buffer. Finally, 2×Laemmli buffer (Bio-Rad Laboratories) supplemented with β-mercaptoethanol was added to the beads. Samples were boiled in a water bath for 10 min before subjecting to SDS-PAGE.

The following antibodies and working dilutions were used in immunoblotting:

SB1.3 antibody against BUB1 (Taylor et al., 2001), 1:500 or 1:1000, sheep polyclonal; BUB3 antibody (Sigma B7811), 1:500, rabbit polyclonal; FKBP12 antibody (Abcam ab2918), 1:2000, rabbit polyclonal; BUBR1 antibody (Bethyl A300-995A), 1:1000, rabbit polyclonal.

Long term live-cell imaging of HeLa cells

Imaging was conducted over a period of 24 hours as described in detail previously (Chen et al.,
2019). We used either the Incucyte Zoom live cell imaging system (Sartorus Inc.) or the
ImageXpress Nano live cell imaging system (Molecular Devices Inc), both equipped with a 20x
Phase objective. To image cells on the Incucyte system, cells were plated in 12-well plastic tissue
culture plates, whereas they were plated in 24-well plate glass-bottom dishes (Corning) for the
ImageExpress Nano system. At each position, one phase, GFP, and mCherry image was
acquired every 10 minutes. The exposure times for mCherry and GFP images were adjusted to
minimize photobleaching while enabling accurate determination of intensity values. It should be
noted that the excitation sources, optics, and detector on the ImageXpress Nano and the Incucyte
microscope are entirely different. Therefore, the mCherry intensity values across different
experiments are not directly comparable.

Image analysis

Prior to intensity quantification, acquired images were pre-processed using functions from the
'Image Processing Toolbox' provided with Matlab as follows. First, the phase image sequence
was registered to remove any movement of the field of view between adjacent time point, and at
each time point, the same transform was applied to the GFP and mCherry images to register
them. Additionally, image intensity from a blank, unseeded well was used to for background
correction of the fluorescence channels. Next, GFP and mCherry fluorescence signals were
quantified using a custom graphical user interface (GUI) written in Matlab as described previously.
Briefly, this interface uses cross-correlation of each phase image with a circular kernel to identify
cells with circular shapes close to the diameter of the circular kernel. The centroids of these
shapes were then linked along the time axis. These images were presented to the user via the
GUI to: (a) discard false positive, non-mitotic cells or debris and (b) to visually correct the time of
entry into or exit from mitosis. The GUI then calculated the GFP and mCherry signals per cell as
the average fluorescence intensity
In all the dose-response assays discussed in this study, the phosphodomain is highly, and
constitutively, expressed in all cells, whereas the kinase domain is expressed conditionally by an
inducible promoter. Consequently, the amount of the kinase domain expressed varies from cell
to cell, and it is lower than the amount of the phosphodomain in most cells. Because of this design
of the eSAC system, the dosage of the dimerized signaling complex in a cell can be inferred from
the amount of Frb-mCherry-Mps1[500-817] in that cell. Therefore, we defined the dosage of the eSAC
signaling complex by quantifying the average mCherry fluorescence within a cell, and the

response as the duration of mitosis (the amount of time that the cell spends with a spherical
morphology that is characteristic of mitotic HeLa cells).

Mathematical modeling

Modeling of dose-response curve for eSAC systems with Bub1 phosphodomains

We constructed this model to understand the mechanisms underlying delayed mitotic exit
produced by the eSAC consisting of Bub1$^{221\text{-}620}$ and Bub1$^{441\text{-}620}$. In the equations below, we refer
to these phosphodomains simply as 'Bub1'. The eSAC activator complex is formed by the
dimerization of Bub1 with the kinase domain of Mps1. Upon phosphorylation by Mps1, Bub1 can
bind Mad1/2. The concentrations of different species of the Bub1 phosphodomain are related by
the equation:

$$[\text{Bub1}]_T = [\text{Bub1-Mps1-Mad1/2}] + [\text{Bub1-Mps1}] + [\text{Bub1}]. \qquad (1)$$

Assuming reversible binding between phosphorylated Bub1 phosphodomain and Mad1/2
complex, we determined the concentration of eSAC activator complex, [Bub1-Mps1-Mad1/2] - it
saturates due to the finite concentration of Mad1-Mad2 complex (set at 100 nM, (Fig. S1A). Note,
that the phosphodomain can produce MCC only if it recruits Mad1/2. Bub1-Mps1 and Bub1 can
both bind BubR1 and Cdc20, but they do not participate in SAC signaling.

We modified a model of metaphase-to-anaphase transition by He et al. (He et al., 2011) as shown
in Figure S1B to capture key proteins and steps in the process of Bub1-mediated MCC formation
and to simulate the effect of MCC on mitotic exit. In the He model, Cyclin B ('CycB') is synthesized
at a constant rate and degraded by APC-Cdc20-dependent ubiquitination (denoted simply as
Cdc20). The abundance of CycB determines the activity of CDK1-CycB complexes. The activity
of the eSAC complex depends positively on CDK1-CycB activity and negatively on PP2-B55, a
counter-acting protein phosphatase (CAPP) (Bouchoux and Uhlmann, 2011; Sullivan et al.,
2004). This is a reasonable assumption, because recent data reveals that Bub1 must be
phosphorylated by both CDK1-CycB and Mps1 before it can bind Mad1 (Ji et al., 2017). The MCC
undergoes spontaneous dissociation, releasing BubR1, Cdc20 and inactive/open Mad2.
Furthermore, active APC-Cdc20 promotes the inactivation of closed/active Mad2 in MCC; this
positive feedback of active Cdc20 on its own release from the MCC accelerates the activation of
APC-Cdc20 during the transition into anaphase (Chen et al., 2019; He et al., 2011).

The molecular species, chemical reactions, parameters and constraints involved in the model are
presented in an Excel file and Table S1 (Supporting Information). A MATLAB script was used to

read this file and produce a file with Ordinary Differential Equations (ODEs) describing the rate of
change of all molecular species. We numerically integrated the ODEs to calculate the time
evolution of [CycB] and the other molecular species. For the initial conditions of species involving
BubR1 and Cdc20 bound to Bub1 phosphodomain (both signaling and non-signaling) we used
the equilibrium concentrations. These concentrations depended on the concentration of eSAC
activator complex and were calculated at the start of each simulation. The initial concentrations
of the following species were kept constant: [CycB] = 45 nM, [Mad2-Cdc20] = 0 nM, [MCC] = 25
nM, [CAPP] = 5 nM. In our model the system temporally evolves towards a steady state
corresponding to anaphase (low [CycB] and high [Cdc20]). We assume that a cell exits mitosis
when [CycB] drops below 1 nM (Fig. S1E). The choice of initial conditions does not qualitatively
affect the outcome of the model. Furthermore, the main result of this analysis – Bub1-BubR1
produces MCC at a higher rate than Bub1 – is robust, even though many different combinations
of parameters produce similar looking dose-response curves. Experimental measurements of
some of these parameters will be needed to constrain the model further.
667
Modeling of dose-response curve for eSAC with MELT and KI motifs
We constructed this model to simulate the observed dose-response data for the different eSAC
systems involving MELT and KI motifs. In these simulations, the phosphodomain consists of four
MELT motifs and either active or inactive KI motifs. All four MELT motifs in contact with the Mps1
kinase domain are assumed to be phosphorylated, whereupon they can recruit Bub1
(representing Bub1-Bub3 in this model. Upon binding to the MELT motif, Bub1 recruits BubR1
(representing BubR1-Bub3). The KI1 motif can bind to Bub1, whereas the KI2 motif can bind to
either Bub1 or BubR1 with characteristic binding affinities. The KI-bound Bub1 and BubR1 do not
participate in SAC signaling. Therefore, the KI motifs act like sinks that reduce Bub1 and BubR1
concentration. Interaction of the MELT motifs with Bub1 and BubR1 produces two types of
signaling complexes: MELpT-Bub1 or MELpT-Bub1-BubR1. For simplicity, we did not explicitly
model in detail how these signaling complexes catalyze MCC formation. Instead, we assume that
the MELpT-Bub1 and MELpT-Bub1-BubR1 complexes catalyze the assembly of MCC at the
apparent rates constants k_{MCC} and k'_{MCC}. A schematic diagram of the molecular mechanism
underlying this model is displayed in Figure S3A. The model simulates dose-response data in two
stages.
684
1. <u>eSAC signaling cascade</u>. We first calculate the steady-state concentrations of SAC signaling
proteins (Bub1 and BubR1) bound to the phosphorylated MELT motifs and KI motifs of an

eSAC phosphodomain. Using the steady-state concentrations of MELpT-Bub1 and MELpT-
Bub1-BubR1, we calculate the rates of generation of MCC by the two phosphodomains,
assuming that these rates is proportional their steady-state concentrations.

2. <u>Mitotic delay</u>. In the second stage of the model, we simulate the duration of mitosis, according
to the overall reaction scheme developed by He et al. and discussed in the previous section.

Stage 1: Simulation of SAC protein recruitment by the eSAC phosphodomains

We assume that MELT and KI motifs in the eSAC phosphodomain interact with SAC proteins
independently. We assigned the same rate of binding of the 'Bub1' protein to each MELT motif
(*kf* in Table S2), but assigned a low unbinding rate (*kr*, Table S2) for the strong MELT motifs
(MELT 1, 12 and 14), and a higher unbinding rate for the weak MELT motif (MELT 13) following
previous studies (Chen et al., 2019; Vleugel et al., 2015). We also chose the dissociation constant
for KI1-Bub1 binding to be equal to the dissociation constant for the KI2-BubR1 binding.

Each phosphorylated MELT motif can be in one of three possible states: MELpT (unbound MELT),
MELpT-Bub1 (MELT bound by Bub1), and MELpT-Bub1-BubR1 (MELT bound by Bub1 and
BubR1). Since there are four MELT motifs in each eSAC activator complex, the number of
possible states for the phosphomimic become $3^4 = 81$. Similarly, each KI motif of the
phosphodomain can be in two states: bound or unbound. The time evolution of concentrations of
different Bub1 and BubR1 bound states of eSAC activator complex is given by:

$$\dot{\mathbf{X}} = \mathbf{AX} \tag{1}$$

where $\mathbf{X} = \{x_1, x_1, \ldots, x_N\}$ is a vector of concentrations of the $N = 81$ different Bub1 and BubR1
binding states of the eSAC, and **A** is the rate matrix. The binding of Bub1 and BubR1 to KI motifs
is described by the set of equations:

711
$$\dot{y}_1 = \text{kfbub} \cdot [\text{Bub1}]_f \cdot y_0 - \text{krbub} \cdot y_1 - \text{kfbub} \cdot [\text{Bub1}]_f \cdot y_1 - \text{kfbubr} \cdot [\text{BubR1}]_f \cdot y_1$$
$$+ \text{krbubr} \cdot y_4 + \text{krbub}' \cdot y_5 \tag{2}$$

712
$$\dot{y}_2 = \text{kfbub} \cdot [\text{Bub1}]_f \cdot y_0 - \text{krbub}' \cdot y_2 - \text{kfbub} \cdot [\text{Bub1}]_f \cdot y_2 + \text{krbub}' \cdot y_5 \tag{3}$$

713
$$\dot{y}_3 = \text{kfbubr} \cdot [\text{BubR1}]_f \cdot y_0 - \text{krbubr} \cdot y_3 - \text{kfbub} \cdot [\text{Bub1}]_f \cdot y_3 + \text{krbubr} \cdot y_4 \tag{4}$$

714
$$\dot{y}_4 = \text{kfbubr} \cdot [\text{BubR1}]_f \cdot y_1 - \text{krbubr} \cdot y_3 - \text{krbub} \cdot y_3 - \text{krbubr} \cdot y_4 \tag{5}$$

715
$$\dot{y}_5 = \text{kfbub}' \cdot [\text{Bub1}]_f \cdot y_1 + \text{kfbub} \cdot [\text{Bub1}]_f \cdot y_2 - \text{krbub} \cdot y_5 - \text{krbub}' \cdot y_5, \tag{6}$$

716   where, y_0 is the concentration of free KI motifs (both KI motifs unbound),

$$y_1 = [\text{KI1-Bub1}],$$

$$y_2 = [\text{KI2-Bub1}],$$

$$y_3 = [\text{KI2-BubR1}],$$

$$y_4 = [\text{KI1-Bub1, KI2-BubR1}],$$

$$y_5 = [\text{KI1-Bub1, KI2-Bub1}].$$

$[\text{Bub1}]_\text{f}$ and $[\text{BubR1}]_\text{f}$ are the concentrations of free Bub1 and BubR1 in the cytoplasm, respectively. The parameters kfbub and kfbub′ are the rate constants for Bub1 binding to the first and the second KI motif respectively (krbub and krbub′ are the corresponding unbinding rates), and kfbubr and krbubr are the binding (unbinding) rates between BubR1 and KI2.

The concentrations satisfy the constraints:

$$[\text{eSAC}] = \sum_{i=1}^{N} x_i \tag{7}$$

$$[\text{Bub1}]_\text{T} = [\text{Bub1}]_\text{f} + y_1 + y_2 + y_4 + 2y_5 + \sum_{i=1}^{N} n_i \cdot x_i \tag{8}$$

$$[\text{BubR1}]_\text{T} = [\text{BubR1}]_\text{f} + y_3 + y_4 + \sum_{i=1}^{N} m_i \cdot x_i \tag{9}$$

Here, x_i is the concentration of the i-th species, n_i = number of Bub1 units bound to i-th species, and m_i = number of BubR1 units bound to the i-th species. We assume $[\text{Bub1}]_\text{T} = 100$ nM and $[\text{BubR1}]_\text{T} = 100$ nM. The equilibrium concentration of each state was obtained by numerically solving $\dot{\mathbf{X}} = \dot{y}_i = 0, i = 1,2 \ldots ,5$.

In experiments, the concentration of eSAC activator complex is measured in arbitrary units of mCherry fluorescence (a. u.), whereas in our model the unit of concentration is nano moles (nM). In our simulations, we chose the maximum value of concentration of eSAC activator complex (the value corresponding to 20 a. u. in experiments) to be 200 nM. For easier comparison to experimental figures, in our simulation results the eSAC activator complex concentration is expressed in arbitrary units, with 1 a. u. of fluorescence corresponding to 10 nM.

Figure S3A shows the abundance of different Bub1 and BubR1-bound states as functions of the total concentration of eSAC activator complex for KI1*-KI2*. At low eSAC concentrations, the eSAC tends to be highly loaded, with Bub1 and BubR1 on every MELT motif. However, for cells

with a high eSAC concentration, [eSAC] >> [Bub1]$_T$, the most abundant eSAC species is one that

does not bind any Bub1 at all (not shown), followed by species that bind either Bub1 or Bub1-

BubR1 at only one of the MELT motifs. We define the sum of concentration of MELpT-Bub1 and

MELpT-Bub1-BubR1 as [eSAC]$_T$:

$$[eSAC]_T = [MELpT\text{-}Bub1] + [MELpT\text{-}Bub1\text{-}BubR1] \qquad (10)$$

$$[MELpT\text{-}Bub1] = \sum_{i=1}^{N} (n_i - m_i) \cdot x_i \qquad (11)$$

$$[MELpT\text{-}Bub1\text{-}BubR1] = \sum_{i=1}^{N} m_i \cdot x_i \qquad (12)$$

<u>Formation of MCC by the eSAC signaling complexes</u>

We assume that the recruitment of SAC proteins (Bub1 and BubR1) to MELT motifs of the eSAC

activator complexes enables the incorporation of Cd20 into MCC. Since different species catalyze

this reaction at different rates, we define the effective rate of conversion, k_{asmcc}, as the

concentration-weighted sum of the conversion rates of each eSAC complex:

$$k_{asmcc} = \frac{1}{[eSAC]_T} \cdot \left[k_{MCC} \cdot \sum_{i=1}^{N} (n_i - m_i) \cdot x_i + k'_{MCC} \cdot \sum_{i=1}^{N} m_i \cdot x_i \right], \qquad (13)$$

where k_{MCC} and k'_{MCC} are the MCC formation rates due to MELpT-Bub1 and MELpT-Bub1-

BubR1, respectively. Note, for most species, the rate of MCC generation is additive. For example,

$k_{0012} = k_{MCC} + k'_{MCC}$ for a phosphodomain that binds only Bub1 at the 13th MELT motif and

Bub1-BubR1 at 14th MELT motif (Table S3). Using [eSAC]$_T$ and k_{asmcc} as inputs for the He model

(discussed below), we calculated the time evolution of Cyclin B concentration and from it the time

in mitosis.

Stage 2: The effect of MCC produced on exit from mitosis

To calculate the effect of MCC generated by the eSAC on mitotic exit, we used a simplified version

of the model of the mitotic checkpoint proposed by the He model (He et al., 2011). Active eSAC

signaling complexes (eSAC$_A$) generate MCC, as described in the previous section. The temporal

dynamics of our mitotic checkpoint model are determined by the ordinary differential equations:

$$\frac{d[CycB]}{dt} = k_{scyc} - \left(k_{dcyc} + k_{dcyc,c20} \cdot [Cdc20] \right) \cdot [CycB] \qquad (14a)$$

$$\frac{d[eSAC_A]}{dt} = k_{ancyc} \cdot [CycB] \cdot ([eSAC]_T - [eSAC_A]) - k_{in,capp} \cdot [CAPP] \cdot [eSAC_A] \qquad (14b)$$

$$\frac{d[MCC]}{dt} = k_{asmcc} \cdot [Cdc20] \cdot [eSAC]_A - (k_{dimcc} + k_{imad} + k_{imad,c20} \cdot [Cdc20]) \cdot [MCC] \qquad (14c)$$

$$\frac{d[CAPP]}{dt} = k_{acapp} \cdot ([CAPP]_T - [CAPP]) - k_{icapp,cyc} \cdot [CycB] \cdot [CAPP] \qquad (14d)$$

In these equations, $[CycB] = [CDK\text{-}CycB]$, $[Cdc20] = [APC\text{-}Cdc20]$, $[eSAC]_T$ is the total

concentration of Bub1 and BubR1-bound eSAC signaling complexes (Eq. 10), which is either in

the active, signaling-competent state eSAC$_A$, or the inactive state, $[eSAC_I] = [eSAC]_T - [eSAC_A]$.

[MCC] and [CAPP] refer to the concentrations of the mitotic checkpoint complex and the CDK-

counteracting protein phosphatase, respectively. In addition, the total concentration of Cdc20 is:

$$[Cdc20]_T = [Cdc20] + [MCC]$$

Here, the subscript T indicates total concentration. The values of the parameters in the model and

of the fixed concentrations of some components, as listed in Table S2, are taken from (He et al.,

2011).

<u>Simulation of time in mitosis</u>

To simulate 'time in mitosis', we assume that a cell exits mitosis when [CycB] drops below 1 nM.

We numerically integrated Ordinary Differential Equations (ODEs) to calculate the time evolution

of [CycB] and the other components of the mitotic checkpoint. As before, the initial conditions for

the simulation are chosen to be [CycB] = 45 nM, [eSAC$_A$] = 0, [MCC] = 25 nM, and [CAPP] = 5

nM. The qualitative aspects of our results do not change if the initial conditions are changed a

little bit. Figure S3C displays typical time courses for [CycB], for different eSACs, for [eSAC

activator complex] = 10 a. u. The system always comes out of mitosis (as seen by the drop in

[CycB]), albeit after different time delays.

Statistical analysis

To determine the overall trend in the dose-response data, the data were first binned (in MATLAB)

and then the mean values of each bin were overlayed on the data. The number of observations,

technical replicates are noted in the figure legends. These mean values were fit with a four-

parameter sigmoidal curve using the GraphPad Prism 9 software. Statistical significance of the

difference between the mean values in Figure 5B was assessed using the unpaired t-test with

Welch's correction.

Figure 1

Figure 1 The contribution of Bub1-BubR1 heterodimerization to MCC assembly during eSAC signaling. A Schematic at the top displays the domain organization of human Knl1. Below is a simplified schematic displaying the pathways that recruit SAC signaling proteins to unattached kinetochores. **B** Schematic at the top displays the domain organization of human Bub1. The

cartoon represents the ectopic SAC activation or 'eSAC' system. **C** Dose-response data for an
eSAC system that uses Bub1$^{444-620}$ as the phosphodomains. Each gray circle displays the average
cellular mCherry fluorescence (X axis) and the time in mitosis (Y axis) for a single cell (n = 1920
from 2 experiments; 139 and 155 are the lower and upper 95% confidence intervals). Blue
squares represent mean values of the dose-response data binned according to mCherry
fluorescence. Vertical and horizontal lines represent S.E.M. The solid blue curve displays the
least squares 4-parameter sigmoidal fit to the binned mean values. Sigmoidal fits for dose-
response data for one MELT motif (MELT12, dashed red curve) and an extended Bub1$^{231-620}$
phosphodomain containing the Bub3- and BubR1-binding sites (dashed blue curve) from our
previous study have been super-imposed for comparison (Chen et al., 2019). Bar graph on the
right displays the maximal mitotic duration estimated by the sigmoidal fit. Vertical lines represent
95% confidence interval on the fit parameter. **D** Cartoon displays the simplified scheme used to
simulate the generation of MCC by the eSAC system. **E** Numerical simulation of the dose-
response curves by assuming the assembly of signaling complexes and ultimately the MCC as
show in **D** followed by a bi-stable switch based on negative feedback described by He et al (He
et al., 2011).
826
827
828

Figure 2

Figure 2 Recruitment of Bub1 and BubR1 by the KI motifs in KNL1 and their contribution to eSAC signaling. A Left: Schematic of the two phosphodomains used to test whether the KI motifs contribute to MCC assembly mediated by the MELT motif in the eSAC phosphodomain. KI1-KI2 indicates phosphodomain with intact KI motifs, KI1*-KI2* indicates phosphodomains wherein the KI motifs are inactivated using suitable point mutations (see Methods for details). Scatter plot in the middle displays the dose-response data for the two phosphodomains (n = 1888 in green and n = 836 in magenta from ≥ 2 experiments respectively for KI1-KI2 and KI1*-KI2*; symbol usage follows the scheme established in 1**C**). The bar graph on the right displays the

maximal mitotic duration predicted by 4-parameter sigmoidal fits to the binned mean values as in

Figure 1C. Vertical lines display 95% confidence intervals on the fit parameter. **B** Left: Schematic

of the phosphodomains consisting of 4 MELT motifs and either active or inactive (indicated by *)

KI motifs. Right: Immunoprecipitation of the eSAC phosphodomains using mNeonGreenTrap

beads followed by Western blot analysis to probe for the co-IP of the indicated proteins. **C** Dose-

response data for the indicated phosphodomains. Only the binned mean values are shown for

clarity. Statistical analysis performed as in 1D (n = 1019, 1024, 666, and 3219 for KI1-KI2, KI1-

KI2*, KI1*-K2, and KI1*-KI2* respectively from ≥ 2 experiments). Bar graph on the right displays

the maximal time in mitosis predicted by the sigmoidal fits.

Figure 3

850
Figure 3 Numerical simulation of the dose-response data for the eSAC phosphodomains
containing KI motifs. A Schematic of the two-stage model used to simulate the dose-response
curves. **B** The influence of Bub1-BubR1 stoichiometry and the differential depletion of Bub1 and
BubR1 by the KI motifs in the simulations on the equilibrium concentrations of the two signaling
complexes formed on phosphorylated MELT motifs. **C** Comparison of the equilibrium
concentration of the MELpT-Bub1-BubR1 complex assembled by KI1*-KI2 and KI1-KI2* (left) and
their ratio (right) as a function of the Bub1-BubR1 stoichiometry. For the two phosphodomains to
generate different responses, the Bub1-BubR1 stoichiometry must be around 1:1 (indicated by

the blue shaded area). **D** Contour plot shows the ratio of the maximal responses produced by

KI1*-KI2 and KI1-KI2* as a function of the Bub1-BubR1 stoichiometry and the ratio of the rates at

which the MELpT-Bub1-BubR1 and the MELpT-Bub1 signaling complexes produce MCC (i.e.,

k'_{MCC}/k_{MCC}). **E** Simulation of the dose-response curves for the four phosphodomains using the

same set of parameter values.

Figure 4

Figure 4 Generation of CRISPR/Cas9 genome-edited HeLa cell lines expressing mNeonGreen-Bub1, mNeonGreen-BubR1, and Mad1-mNeonGreen. A Western blots show that showing that roughly half the BubR1 and Mad1 (left) and Bub1 (middle) is tagged with mNeonGreen in the three partially genome-edited cell lines (asterisks on the right of each displayed blot mark the mNG fusion protein). Western blot on the right shows whole cell extracts of the three cell lines probed with anti-mNeonGreen antibodies. **B** Average mNG-Bub1, mNG-BubR1, Mad1-mNG, and M4-KI1-KI2-mNG-2xFkbp12 signals from mitotic HeLa cells (left) and estimation of the relative protein abundance, assuming that the total protein is twice as high as the abundance of the mNG-labeled species (mean ± std. dev.).

Figure 5

Figure 5 Recruitment of BubR1 by BuB1 per se contributes to the signaling strength of the

SAC. A Micrographs of representative cells displaying the indicated antigens at the top. The top

row displays micrographs of the partially genome-edited cell line expressing mNG-BubR1. The

bottom two rows display representative cells after endogenous BubR1 was knocked down and

the indicated mutant, mNG-tagged version was ectopically expressed (scale bar ~ 1.22 microns).

B Scatterplot on the left displays the duration of mitosis of either genome-edited cells expressing

WT BubR1 or cells that ectopically expressed the indicated mutant after BubR1 RNAi in media

containing 50 nM nocodazole. Scatterplot on the right displays the average intracellular

mNeonGreen signal from the same cells (mean ± 95% confidence intervals).

888 **Supplementary Figure 1**

889
890
Figure S1 Simulation of the dose-response curves for the Bub1 eSAC system. A The
amount of the eSAC signaling complex (Bub1-Mad1) saturates due to the limited availability of
Mad1 (100 nM). **B** Schematic of the model used to simulate the dose-response data for the "full"
and "truncated" Bub1 phosphodomain used in the eSAC system. **C** Temporal evolution of Cyclin
B concentration for the two eSAC systems. **D** Assessment of MCC formation rates for the full and
truncated phosphodomain. **E** Superimposition of the dynamics of the MCC formation rate and the
concentration of signaling complexes that produce MCC.

898
899

Supplementary Figure 2

Supplementary Figure 2 A Western blot probing for the indicated protein for assessing the co-immunoprecipitation of Bub1 and BubR1 with the indicated eSAC phosphodomain (at the top). The asterisk marks the position of the band corresponding to the full-length protein. The cross marks bands that likely degradation products of the full-length protein. **B** Residual plots from a 4-

parameter sigmoidal fit to the dose-response data for the KI1*-KI2* (left) and KI1-KI2* (right)

phosphodomains. **C** Dose-response data for the indicated phosphodomains. Circles represent

average values of binned data (vertical and horizontal lines display standard error on each mean

value. n= 498, 1313, and 394 respectively for M1-KI1-KI2-M3-M3, M1-KI1*-KI2*-M3-M3, and M1-

KI1-KI2-M3-M3 without rapamycin respectively pooled from two or more experiments).

Supplementary Figure 3

916
917
Figure S3 Simulation of the dose-response data for eSAC systems containing the KI motifs

and MELT motifs. A Abundance of the indicated eSAC signaling complex (of the form n*(MELpT-

Bub1-BubR1)) as a function of the total concentration of the eSAC activator complex. The number

n indicates the number of MELT motifs bound to Bub1-BubR1. Also see Table S3. **B.** Schematic

of the model used to simulate the dose-response data for eSAC with MELT and KI motifs. **C**

Temporal evolution of Cyclin B concentration for the MELT and KI motif eSAC system. **D**

Simulation of dose-response curves for the four phosphodomains without synergistic interactions

between the MELT motifs.

References

Bolanos-Garcia, V.M., Lischetti, T., Matak-Vinkovic, D., Cota, E., Simpson, P.J., Chirgadze, D.Y., Spring, D.R., Robinson, C.V., Nilsson, J., and Blundell, T.L. (2011a). Structure of a Blinkin-BUBR1 complex reveals an interaction crucial for kinetochore-mitotic checkpoint regulation via an unanticipated binding Site. Structure *19*, 1691-1700.

Bolanos-Garcia, Victor M., Lischetti, T., Matak-Vinković, D., Cota, E., Simpson, Pete J., Chirgadze, Dimitri Y., Spring, David R., Robinson, Carol V., Nilsson, J., and Blundell, Tom L. (2011b). Structure of a Blinkin-BUBR1 Complex Reveals an Interaction Crucial for Kinetochore-Mitotic Checkpoint Regulation via an Unanticipated Binding Site. Structure *19*, 1691-1700.

Bouchoux, C., and Uhlmann, F. (2011). A quantitative model for ordered Cdk substrate dephosphorylation during mitotic exit. Cell *147*, 803-814.

Chen, C., Whitney, I.P., Banerjee, A., Sacristan, C., Sekhri, P., Kern, D.M., Fontan, A., Kops, G., Tyson, J.J., Cheeseman, I.M.*, et al.* (2019). Ectopic Activation of the Spindle Assembly Checkpoint Signaling Cascade Reveals Its Biochemical Design. Curr Biol *29*, 104-119 e110.

Collin, P., Nashchekina, O., Walker, R., and Pines, J. (2013). The spindle assembly checkpoint works like a rheostat rather than a toggle switch. Nat Cell Biol *15*, 1378-1385.

Dick, A.E., and Gerlich, D.W. (2013). Kinetic framework of spindle assembly checkpoint signalling. Nat Cell Biol *15*, 1370-1377.

Faesen, A.C., Thanasoula, M., Maffini, S., Breit, C., Muller, F., van Gerwen, S., Bange, T., and Musacchio, A. (2017). Basis of catalytic assembly of the mitotic checkpoint complex. Nature.

Foley, E.A., Maldonado, M., and Kapoor, T.M. (2011). Formation of stable attachments between kinetochores and microtubules depends on the B56-PP2A phosphatase. Nat Cell Biol *13*, 1265-1271.

He, E., Kapuy, O., Oliveira, R.A., Uhlmann, F., Tyson, J.J., and Novák, B. (2011). System-level feedbacks make the anaphase switch irreversible. Proceedings of the National Academy of Sciences.

Ji, Z., Gao, H., Jia, L., Li, B., and Yu, H. (2017). A sequential multi-target Mps1 phosphorylation cascade promotes spindle checkpoint signaling. Elife *6*.

Jia, L., Li, B., and Yu, H. (2016). The Bub1-Plk1 kinase complex promotes spindle checkpoint signalling through Cdc20 phosphorylation. Nat Commun *7*, 10818.

Khandelia, P., Yap, K., and Makeyev, E.V. (2011). Streamlined platform for short hairpin RNA interference and transgenesis in cultured mammalian cells. Proc Natl Acad Sci USA *108*, 12799-12804.

Kiyomitsu, T., Murakami, H., and Yanagida, M. (2011). Protein interaction domain mapping of human kinetochore protein Blinkin reveals a consensus motif for binding of spindle assembly checkpoint proteins Bub1 and BubR1. Mol Cell Biol *31*, 998-1011.

Krenn, V., Overlack, K., Primorac, I., van Gerwen, S., and Musacchio, A. (2014). KI motifs of human Knl1 enhance assembly of comprehensive spindle checkpoint complexes around MELT repeats. Curr Biol *24*, 29-39.

Krenn, V., Wehenkel, A., Li, X., Santaguida, S., and Musacchio, A. (2012). Structural analysis reveals features of the spindle checkpoint kinase Bub1-kinetochore subunit Knl1 interaction. J Cell Biol *196*, 451-467.

Lara-Gonzalez, P., Kim, T., Oegema, K., Corbett, K., and Desai, A. (2021). A tripartite mechanism
catalyzes Mad2-Cdc20 assembly at unattached kinetochores. Science *371*, 64-67.
Leontiou, I., London, N., May, K.M., Ma, Y., Grzesiak, L., Medina-Pritchard, B., Amin, P.,
Jeyaprakash, A.A., Biggins, S., and Hardwick, K.G. (2019). The Bub1-TPR Domain Interacts Directly
with Mad3 to Generate Robust Spindle Checkpoint Arrest. Curr Biol *29*, 2407-2414 e2407.
Musacchio, A. (2015). The Molecular Biology of Spindle Assembly Checkpoint Signaling Dynamics.
Curr Biol *25*, R1002-1018.
Nijenhuis, W., Vallardi, G., Teixeira, A., Kops, G.J., and Saurin, A.T. (2014). Negative feedback at
kinetochores underlies a responsive spindle checkpoint signal. Nat Cell Biol *16*, 1257-1264.
Nijenhuis, W., von Castelmur, E., Littler, D., De Marco, V., Tromer, E., Vleugel, M., van Osch, M.H.,
Snel, B., Perrakis, A., and Kops, G.J. (2013). A TPR domain-containing N-terminal module of MPS1
is required for its kinetochore localization by Aurora B. J Cell Biol *201*, 217-231.
Overlack, K., Bange, T., Weissmann, F., Faesen, A.C., Maffini, S., Primorac, I., Muller, F., Peters,
J.M., and Musacchio, A. (2017). BubR1 Promotes Bub3-Dependent APC/C Inhibition during
Spindle Assembly Checkpoint Signaling. Curr Biol *27*, 2915-2927 e2917.
Overlack, K., Primorac, I., Vleugel, M., Krenn, V., Maffini, S., Hoffmann, I., Kops, G.J., and
Musacchio, A. (2015). A molecular basis for the differential roles of Bub1 and BubR1 in the spindle
assembly checkpoint. Elife *4*, e05269.
Piano, V., Alex, A., Stege, P., Maffini, S., Stoppiello, G.A., Huis In 't Veld, P.J., Vetter, I.R., and
Musacchio, A. (2021). CDC20 assists its catalytic incorporation in the mitotic checkpoint complex.
Science *371*, 67-71.
Primorac, I., Weir, J.R., Chiroli, E., Gross, F., Hoffmann, I., van Gerwen, S., Ciliberto, A., and
Musacchio, A. (2013). Bub3 reads phosphorylated MELT repeats to promote spindle assembly
checkpoint signaling. Elife *2*, e01030.
Qian, J., Garcia-Gimeno, M.A., Beullens, M., Manzione, M.G., Van der Hoeven, G., Igual, J.C.,
Heredia, M., Sanz, P., Gelens, L., and Bollen, M. (2017). An Attachment-Independent Biochemical
Timer of the Spindle Assembly Checkpoint. Mol Cell *68*, 715-730 e715.
Roy, B., Han, S.J.Y., Fontan, A.N., and Joglekar, A.P. (2020). The copy-number and varied strength
of MELT motifs in Spc105 balance the strength and responsiveness of the Spindle Assembly
Checkpoint. Elife *9*.
Roy, B., Han, S.J.Y., Fontan, A.N., and Joglekar, A.P. (2021). Aurora B phosphorylates Bub1 to
promote spindle assembly checkpoint signaling. bioRxiv, 2021.2001.2005.425459.
Saurin, A.T., van der Waal, M.S., Medema, R.H., Lens, S.M., and Kops, G.J. (2011). Aurora B
potentiates Mps1 activation to ensure rapid checkpoint establishment at the onset of mitosis.
Nat Commun *2*, 316.
Silio, V., McAinsh, A.D., and Millar, J.B. (2015). KNL1-Bubs and RZZ Provide Two Separable
Pathways for Checkpoint Activation at Human Kinetochores. Dev Cell *35*, 600-613.
Sullivan, M., Higuchi, T., Katis, V.L., and Uhlmann, F. (2004). Cdc14 phosphatase induces rDNA
condensation and resolves cohesin-independent cohesion during budding yeast anaphase. Cell
117, 471-482.
Taylor, S.S., Hussein, D., Wang, Y., Elderkin, S., and Morrow, C.J. (2001). Kinetochore localisation
and phosphorylation of the mitotic checkpoint components Bub1 and BubR1 are differentially
regulated by spindle events in human cells. J Cell Sci *114*, 4385-4395.

Vleugel, M., Omerzu, M., Groenewold, V., Hadders, M.A., Lens, S.M., and Kops, G.J. (2015).
Sequential multisite phospho-regulation of KNL1-BUB3 interfaces at mitotic kinetochores. Mol
Cell *57*, 824-835.
Vleugel, M., Tromer, E., Omerzu, M., Groenewold, V., Nijenhuis, W., Snel, B., and Kops, G.J.
(2013). Arrayed BUB recruitment modules in the kinetochore scaffold KNL1 promote accurate
chromosome segregation. J Cell Biol *203*, 943-955.
Yuan, I., Leontiou, I., Amin, P., May, K.M., Soper Ni Chafraidh, S., Zlamalova, E., and Hardwick,
K.G. (2017). Generation of a Spindle Checkpoint Arrest from Synthetic Signaling Assemblies. Curr
Biol *27*, 137-143.
Zhang, G., Lischetti, T., Hayward, D.G., and Nilsson, J. (2015). Distinct domains in Bub1 localize
RZZ and BubR1 to kinetochores to regulate the checkpoint. Nat Commun *6*, 7162.
Zhang, G., Lischetti, T., and Nilsson, J. (2014). A minimal number of MELT repeats supports all the
functions of KNL1 in chromosome segregation. J Cell Sci *127*, 871-884.
Zhang, G., Mendez, B.L., Sedgwick, G.G., and Nilsson, J. (2016). Two functionally distinct
kinetochore pools of BubR1 ensure accurate chromosome segregation. Nat Commun *7*, 12256.
1027