A B cell actomyosin arc network couples integrin co-stimulation to mechanical force-dependent immune synapse formation

Jia C. Wang[1], Yang-In Yim[1], Xufeng Wu[2], Valentin Jaumouillé[1], Clare M. Waterman[1], John H. Kehrl[3], and John A. Hammer[1,4]

[1] Cell and Developmental Biology Center, National Heart, Lung and Blood Institute, National Institutes of Health, Bethesda MD 20892

[2] Light Microscopy Core, National Heart, Lung and Blood Institute, National Institutes of Health, Bethesda MD 20892

[3] B Cell Molecular Immunology Section, National Institute of Allergy and Infectious Disease, National Institutes of Health, Bethesda MD 20892

[4] Author for correspondence: hammerj@nhlbi.nih.gov

Abstract

B-cell activation and immune synapse (IS) formation with membrane-bound
antigens are actin-dependent processes that scale positively with the strength of
antigen-induced signals. Importantly, ligating the B-cell integrin, LFA-1, with ICAM-1
promotes IS formation when antigen is limiting. Whether the actin cytoskeleton plays a
specific role in integrin-dependent IS formation is unknown. Here we show using super-
resolution imaging of primary B cells that LFA-1: ICAM-1 interactions promote the
formation of an actomyosin network that dominates the B-cell IS. This network is
created by the formin mDia1, organized into concentric, contractile arcs by myosin 2A,
and flows inward at the same rate as B-cell receptor (BCR): antigen clusters.
Consistently, individual BCR microclusters are swept inward by individual actomyosin
arcs. Under conditions where integrin is required for synapse formation, inhibiting
myosin impairs synapse formation, as evidenced by reduced antigen centralization,
diminished BCR signaling, and defective signaling protein distribution at the synapse.
Together, these results argue that a contractile actomyosin arc network plays a key role
in the mechanism by which LFA-1 co-stimulation promotes B-cell activation and IS
formation.

Introduction

32   BCR engagement with cognate antigen triggers striking changes in B cell physiology that promote B cell activation, immune synapse (IS) formation, and B cell effector functions (1-3). These changes include dramatic increases in actin filament assembly and dynamics that are thought to drive IS formation in B cells engaged with membrane-bound antigen (1-4). For B cells *in vivo*, this usually involves interactions with antigen bound to the surface of an antigen-presenting cell (APC) (4-6), although activating surfaces such as antigen-coated glass and planar lipid bilayers containing freely-diffusing antigen are used to mimic these *in vivo* interactions. IS formation in these contexts is initiated by the formation of a radially-symmetric, Arp2/3 complex-dependent branched actin network at the outer edge of the IS (i.e. in the distal supramolecular activation cluster or dSMAC) (7, 8). This lamellipodia-like actin network drives the spreading of the B cell across the antigen-coated surface, thereby promoting BCR: antigen interactions (3, 9). Once the B cell is fully spread, the continued polymerization of branched actin at the outer edge of the dSMAC generates a centripetal or retrograde flow of actin that drives the movement of BCR: antigen clusters (10-12) towards the center of the synapse (i.e. to the central SMAC or cSMAC) (8, 13). This centripetal actin flow, combined with an overall contraction of the B cell, is thought to be responsible for the transport of BCR: antigen clusters to the center of the maturing synapse (13). Importantly, this process of antigen centralization is required for robust BCR signaling (3, 11, 14), and is thought to be a prerequisite for antigen internalization by follicular B cells (15-17).

54   Antigen-induced IS formation scales with the strength of antigen-induced signals such that IS formation and B cell activation are attenuated when membrane-bound antigen binds the BCR weakly or is presented at low density. Importantly, co-stimulatory signals can promote IS formation and B cell activation under both of these conditions (18, 19). Seminal work from Carrasco and colleagues showed that the B-cell integrin LFA-1, which binds the adhesion molecule ICAM-1 present on the surface of APCs (20, 21), serves as one such co-stimulatory signal (18). This conclusion was based on four key observations. First, B cells responded robustly to higher affinity membrane-bound antigens presented at high density whether or not ICAM-1 was present on the membrane. Second, the robust activation of B cells in response to antigens of all affinities increasingly required ICAM-1 in the membrane as the density of the antigen was lowered. Third, this co-stimulatory effect was most dramatic for weaker antigens. Finally, this latter effect was not observed in B cells lacking LFA-1. With regard to the underlying mechanism, IRM imaging suggested that LFA-1: ICAM-1 interactions, which were shown to concentrate in the medial portion of the synapse (i.e. the peripheral SMAC or pSMAC), lower the threshold for B cell activation by enhancing cell adhesion.

While the actin cytoskeleton clearly plays a central role in driving IS formation, whether it plays a specific role in integrin-dependent IS formation is unknown. This is an important question, as most B cell interactions with professional APCs presenting cognate antigen involve integrin ligation. Relevant to this question, the dendritic actin network occupying the outer dSMAC ring, which is thought to be the main driver of IS formation, has been observed primarily in cells that received antigen stimulation alone, and almost exclusively in immortalized B cell lines (13, 22-25). It is not known, therefore, whether integrin-co-stimulation alters the organization and/or dynamics of actin at the B-cell IS. Moreover, we are only just beginning to elucidate the organization and dynamics of synaptic actin networks formed by primary B cells. Here we show that LFA-1: ICAM-1 interactions in primary B cells stimulate the formation of a contractile actomyosin arc network that occupies the pSMAC portion of the synapse. This actomyosin network represents the major actin structure at the IS of primary B cells receiving integrin co-stimulation, and its dynamics drive antigen centralization by sweeping antigen centripetally. Importantly, under conditions of limiting antigen, where integrin co-stimulation is required for IS formation, blocking the contractility of this pSMAC network inhibits IS formation and BCR signaling. Finally, we show that germinal center (GC) B cells can also create this actomyosin structure, suggesting that it contributes to the process of antibody affinity maturation. Together, our data demonstrate that a contractile actomyosin arc network created downstream of integrin ligation plays a major role in the mechanism by which integrin co-stimulation promotes B cell activation and IS formation when antigen is limiting. Importantly, these findings highlight the need for including integrin co-stimulation when examining the role of actin during B cell activation, especially under physiologically relevant conditions.

                                              Results

Integrin co-stimulation promotes the formation of an actin arc network in the
pSMAC

To investigate the possibility that LFA-1 ligation might promote B cell activation
by triggering a significant change in synaptic actin organization, we imaged F-actin at
ISs formed by primary mouse B cells on glass surfaces coated with either anti-IgM or
anti-IgM plus ICAM-1. F-actin was visualized using GFP-F-Tractin, a dynamic reporter
for F-actin (26, 27), and two super-resolution imaging modalities: Airyscan (*xy* resolution
~140 nm) and total internal reflection-structured illumination (TIRF-SIM; *xy* resolution
~100 nm). Individual video frames of anti-IgM-engaged B cells using both imaging
modalities (Fig. 1A, B), together with the corresponding movies (Movies 1A and 1B),
revealed a thin, bright, highly-dynamic outer rim of F-actin (white arrows in Fig. 1A, B)
that likely corresponds to the branched actin network comprising the dSMAC (13, 22,
25). Both modalities (but especially TIRF-SIM) showed that the F-actin present inside
this outer dSMAC rim is composed of a highly disorganized mixture of short actin
filaments/fibers and actin foci (blue brackets in Fig. 1A, B). In sharp contrast, individual
video frames of anti-IgM+ICAM-1-engaged B cells using both modalities (Fig. 1C, D),
together with the corresponding movies (Movies 2A and 2B), showed a highly organized
network inside the outer dSMAC rim (i.e. in the pSMAC) that is comprised of concentric
actin arcs (blue brackets and red arrows in Fig. 1C, D). The difference in synaptic actin
organization between anti-IgM-engaged B cells and anti-IgM+ICAM-1-engaged B cells
is very evident in enlarged TIRF-SIM images. While it is challenging to define SMAC
boundaries and any pattern of F-actin organization in the pSMAC of B cells engaged
with anti-IgM alone (Fig. 1E1, E2), SMAC boundaries and pSMAC F-actin organization
are both very distinct in B cells engaged using anti-IgM+ICAM-1 (Fig. 1F1, F2).
Consistently, scoring B cells for the presence of a discernable actin arc network showed
that the addition of ICAM-1 increases the percentage of such cells from ~30% to ~70%
(Fig. 1G). Moreover, dynamic imaging showed that the arcs in cells engaged with anti-
IgM alone are typically sparse and transient (Movies 1A and 1B), while those in cells
engaged with both anti-IgM and ICAM-1 are dense and persistent (Movies 2A and 2B).
Finally, measuring the percentage of total synaptic F-actin content within each SMAC
(Fig. 1H), and the percentage of total IS footprint occupied by each SMAC (Fig. 1I),
showed that the actin arc-containing pSMAC comprises the major actin network at the
IS of primary B cells engaged using both anti-IgM and ICAM-1. Together, these results
demonstrate that LFA-1 co-stimulation promotes the formation of a pSMAC actin arc
network that dominates the B cell IS.

Linear actin filaments generated by the formin mDia1 at the outer edge of the
synapse give rise to the pSMAC actin arc network

We next sought to define the origin of the actin arcs that comprise the pSMAC of
B cells stimulated using both anti-IgM and ICAM-1. Primary B cells stimulated in this
way exhibit small, actin-rich surface spikes at the outer synapse edge (Fig. 2A).
Importantly, magnified images revealed that the actin within these spikes continues into
the cytoplasm in the form of linear actin filaments (Fig. 2B1, B2). Moreover, tracing
these linear actin filaments showed that they are contiguous with the pSMAC actin arcs
(Fig. 2C1, C2; Movies 3A and 3B). These results argue that linear actin filaments
nucleated at the plasma membrane at the outer edge of the synapse give rise to the
actin arcs populating the pSMAC. While these results do not identify the specific
nucleator involved, they do point to it being a member of the formin family based on the
fact that the actin being made is linear and nucleated at the plasma membrane (28, 29).
Consistent with this conjecture, and with the fact that formins incorporate fluorescent
protein-labelled actin monomer into filaments poorly (26, 27, 30), we did not see
fluorescent actin arcs in B cells expressing mEOS-labeled G-actin (Movie 4).

To test if a formin is indeed responsible for creating the pSMAC actin arc
network, we used the pan-formin inhibitor SMIFH2 (31). Fig. 2D1 and 2D2, together with
the line scan in Fig. 2E, show that the pSMAC actin arcs present in a representative
primary B cell immediately before SMIFH1 addition (blue trace) had largely disappeared
6 minutes after adding SMIFH2 (red trace). Given recent concerns about the specificity
of SMIFH2 (32), we used three different miRNAs to knock down the formin mDia1 in the
lymphoma B cell line A20 (Fig. S1), which also forms pSMAC actin arcs when
stimulated using anti-IgG+ICAM-1 (Fig. 1J1, J2; Movie 5A). mDia1 was chosen as the
miRNA target as it is highly expressed in B cells (Immgen Database) and is largely
responsible for making linear actin filaments in T cells (27). Compared to control A20 B
cells (Fig. 2F1), representative B cells expressing each of the three miRNAs (Fig. 2F2-
F4) were largely devoid of actin arcs. This difference was supported by quantitating the
ratio of pSMAC to dSMAC F-actin (Fig. 2G), as well as the amount of F-actin in the
pSMAC (Fig. 2H). Together, these results argue that the pSMAC actin arcs are indeed
created by a formin, and that the formin mDia1 plays a major role.

To provide one final piece of evidence that the arcs are created by a formin, we
imaged A20 B cells following the addition of the Arp2/3 inhibitor CK-666. The rationale
for this experiment lies in the recent revelation that the two major consumers of actin
monomer in cells, the Arp2/3 complex and formins, are always competing for a limiting
pool of actin monomer (33-36). One consequence of this competition is that when one
of these nucleators is inhibited, the actin structures created by the other nucleator get
more robust because that nucleator now gets more monomer. For example, inhibiting
the Arp2/3 complex promotes the formation of formin-dependent actin networks in both
yeast and vertebrate cells (27, 33-36). Given this, and assuming that the arcs in B cells
are formin-generated, then inhibiting the Arp2/3 complex in B cells should lead not only

to a diminution of the branched actin network in the dSMAC, but also to an amplification
of the arc network in the pSMAC. Consistently, Figure S2A1/A2 (before CK-666
addition) and Figure S2A3/A4 (after CK-666 addition) together show that CK-666
addition leads not only to a reduction in the size of the dSMAC (red brackets), but also
to an increase in arc content in the pSMAC (blue brackets). These changes were
supported by measuring the percentage of total synaptic F-actin content residing within
each SMAC (Fig. S2B), which revealed a significant shift away from dSMAC F-actin and
toward pSMAC F-actin following CK-666 treatment. This shift was also reflected in
measurements of total pSMAC F-actin content (Fig. S2C), the ratio of pSMAC to
cSMAC F-actin content (Fig. S2D), and the ratio of pSMAC to cSMAC area (Fig. S2E).
Taken together, these data argue strongly that linear actin filaments generated by the
formin mDia1 at the outer edge of the synapse give rise to the pSMAC actin arc
network.

Myosin 2A co-localizes with the actin arcs

Having established that ICAM-1 co-stimulation promotes the formin-dependent
formation of actin arcs in the pSMAC, we asked how these arcs are organized into
concentric structures. Formin-derived linear actin filaments are commonly organized
into well-defined structures such as stress fibers, transverse arcs, and the contractile
ring in dividing cells by bipolar filaments of the actin-based motor protein myosin 2 (37-
39). We decided, therefore, to test whether myosin 2 co-localizes with the actin arcs and
is required for their concentric organization.

To define the localization and dynamics of myosin 2 at the B cell IS, we used
primary B cells isolated from a mouse in which GFP had been knocked into the N-
terminus of the myosin 2A (M2A) heavy chain gene (40), as M2A is the only myosin 2
isoform expressed in B cells (Immgen Database). Individual video frames of these cells
following transfection with Td-Tomato-F-Tractin and attachment to coverslips coated
with anti-IgM and ICAM-1 revealed a dramatic co-localization between M2A and the
actin arcs in the pSMAC (Fig. 3A1-A3; Movie 6). Magnified TIRF-SIM images show that
the myosin signals align with actin arcs in a periodic fashion (Fig. 3A4) that resembles
other myosin 2-rich, linear actin structures like stress fibers and the contractile ring (41).
Moreover, these myosin signals exhibit the SIM signature for M2A bipolar filaments
when M2A is GFP-labeled at its N-terminus (42), which is a pair of GFP puncta spaced
~300 nm apart (Fig. 3A5; 304 ± 32 nm; n = 230 filaments from 12 cells). The presence
of M2A filaments in the medial portion of the synapse was also evident in primary B
cells isolated from a mouse in which mCherry had been knocked into the N-terminus of
M2A (Fig. S3A), in primary B cells that we genome edited using CRISPR to place GFP
at the N-terminus of M2A (Fig. S3B), and in A20 B cells that we genome edited using
CRISPR to place mScarleti at the N-terminus of M2A and then transfected with GFP-F-
Tractin (Fig. S3C1-C3; Movie 5B). Finally, 3D-SIM images of A20 B cells that were fixed

and stained for M2A and actin showed that endogenous M2A also co-localizes with the actin arcs (Fig. S3D1-D3; note that the signature for M2A filaments using this antibody, which recognizes the C-terminus of M2A, is a single fluorescent punctum that corresponds to the center of an individual M2A filament (41, 43)). The extent of this colocalization was even clearer in enlarged images of immunostained cells (Fig. S3E1-E3), where line scans showed endogenous M2A coinciding with actin arcs (Fig. S3F). Together, these results show that the actin arc network in primary B cells receiving ICAM-1 co-stimulation is in fact an actomyosin arc network.

To gain insight into how the arcs become decorated with M2A filaments, we examined time lapse TIRF-SIM images of GFP-M2A knockin primary B cells expressing Td-Tomato F-Tractin. Individual video frames (Fig. 3B1-B6), as well as the corresponding movie (Movie 7), show that bipolar filaments of M2A begin to appear near the dSMAC: pSMAC boundary in association with the linear actin filaments/bundles exiting the dSMAC (white, yellow and fuchsia arrowheads mark such myosin filaments at time 0s in Fig. 3B1). As time progresses, these filaments move centripetally and undergo expansion into filament clusters (Fig. 3B1-B6; see also Movie 7). This expansion, in which individual myosin filaments expand into a small cluster of filaments, is presumably driven by the same sequential amplification pathway described previously for M2A filament assembly in Hela cells (42). Finally, the myosin filaments in these clusters begin to align with the arcs forming at the outer edge of the pSMAC, which then merge with the larger actomyosin arc network in the pSMAC (Fig. 3B1-B6). As all this is happening, new myosin filaments keep appearing near the dSMAC: pSMAC boundary to repeat the process (Fig. 3B2-B6; follow the blue, green and purple arrowheads).

Given that ICAM-1 co-stimulation promotes the formation of actin arcs, and that the arcs recruit M2A, ICAM-1 co-stimulation should also result in an increase in the amount of M2A at the IS. Consistently, primary GFP-M2A knockin B cells receiving both anti-IgM and ICAM-1 stimulation exhibited a greater amount of synaptic M2A than B cells receiving only anti-IgM stimulation (Fig. 3C-3E). Of note, this difference remained significant even after normalizing the M2A fluorescence for a small difference in the average cell-spread area under these two conditions (Fig. S3G1, G2).

Myosin 2A contractility is required for the concentric organization of the actin arcs and for integrin-dependent traction force

The organization of formin-generated linear actin filaments into well-defined structures is typically driven by the contractility of myosin 2 filaments (37, 38). Therefore, we asked if M2A contractility is required for the concentric organization of the pSMAC actin arcs by treating cells with para-nitroblebbistatin (pnBB), a blue light-insensitive version of the cell-permeable, small molecule myosin 2 inhibitor blebbistatin

(BB) that blocks myosin 2-based contractility by locking the myosin in its weak actin
binding state (44). While control, DMSO-treated cells exhibited concentric actin arcs in
their pSMAC as expected (Fig. 3F), cells treated with 25 µM pnBB displayed highly-
disorganized, mesh-like actin arrays in their pSMAC (Fig. 3G). To quantify this defect in
pSMAC actin organization, we used FibrilTool (45) to measure the anisotropy of actin
filaments in the pSMAC. This tool measures how well structures of interest (here actin
filaments) within a given region of interest (ROI; here the pSMAC) are arranged in
parallel (anisotropy values range from 0 when the orientation of the structures is
completely random to 1.0 when all of the structures are perfectly aligned/parallel to each
other). To measure actin anisotropy in the radially-symmetric IS, we "linearized"
pSMACs by dividing them into 10-12 trapezoid-shaped ROIs of similar size (Fig. S3H).
As anticipated, the anisotropy values revealed a dramatic shift towards more
disorganized pSMAC actin when B cells are treated with pnBB (Fig. 3H). Together,
these results demonstrate that M2A contractility is indeed required for the concentric
organization of the pSMAC actin arcs.

We used traction force microscopy in combination with pnBB to ask if integrin-
dependent traction forces that B cells exert on a deformable substate require M2A
contractility. As expected (46, 47), B cells engaged with substrate coated with anti-IgM
and ICAM-1 generated significantly more traction force than B cells engaged with
substrate coated with anti-IgM alone (Fig. S4A1, A2, B1, B2, D). Importantly, ICAM-1-
dependent traction forces were completely abrogated by pre-treating the cells with
pnBB (Fig. S4C1, C2, D), indicating that the generation of integrin-dependent traction
forces requires M2A contractility. This requirement likely reflects pulling forces exerted
by M2A on the substrate through LFA-1: ICAM-1 pairs, combined with the increase in
M2A content at the synapse caused by ligating LFA-1 with ICAM-1, and the contribution
that M2A-dependent pulling forces make in keeping LFA-1 in its open, active
conformation (48-50). These results, together with the fact that integrin clusters are
known to accumulate in the pSMAC portion of the B cell IS (18, 19), suggest a feed-
forward relationship where integrin ligation promotes the formation of pSMAC
actomyosin arcs, and the contractile forces exerted by these actomyosin arcs promote
further integrin activation and robust adhesion in the pSMAC.

The actomyosin arc network in the pSMAC exhibits centripetal flow

Inward flows of cortical actin networks are thought to drive the transport of
antigen receptor clusters to the center of maturing synapses in both T cells and B cells
((7, 36, 50); although see (51, 52)). For B cells, the clearest example of this to date is
the demonstration that the centripetal flow of the branched actin network comprising the
dSMAC propels BCR: antigen clusters towards the cSMAC (13). As a prelude to asking
whether the actomyosin arcs comprising the pSMAC also contribute to antigen
centralization, we asked if this contractile network exhibits centripetal flow. Kymograph

analyses of actin flow across synapses made by primary B cells expressing GFP-F-
Tractin showed that their pSMAC actomyosin arc network indeed flows centripetally at
1.07 +/-0.07 μm/min, or about one third the rate of centripetal actin flow in the dSMAC
(2.89 +/- 0.18 μm/min) (Fig. S5A1-A3). Similar results were obtained for A20 B cells
(pSMAC rate: 0.97 +/- 0.13 μm/min; dSMAC rate: 3.16 +/- 0.35 μm/min) (Fig. S5B1-B3).
Together, these results indicate that the actomyosin arcs could contribute along with the
branched actin network in the dSMAC to the inward transport of BCR: antigen clusters.

Actomyosin arcs contribute to antigen centralization by sweeping BCR: antigen clusters inward

We used planar lipid bilayers (PLBs) to determine if the actomyosin arcs do in
fact contribute to antigen centralization. As expected, primary B cells expressing GFP-
F-Tractin readily formed actin arcs when PLBs contained both anti-IgM and ICAM-1
(Movie 8A), but not when they contained anti-IgM alone (Movie 8B). Also as expected,
primary B cells engaged with PLBs containing fluorescent anti-IgM (red) and unlabeled
ICAM-1 yielded mature synapses in which concentric actin arcs surrounded antigen
accumulated in the cSMAC (Fig. 4A1-A3, white arrows). To obtain a holistic view of
antigen centralization, we imaged antigen clusters in the dSMAC and pSMAC of primary
B cells over time with the aim of correlating their rates of centripetal transport with the
distinct rates of centripetal actin flow exhibited by these two IS zones (Movie 9).
Tracking of single antigen microclusters showed that they moved inward at 2.36 +/- 1.1
μm/min and 1.03 +/- 0.3 μm/min across the dSMAC (red tracks) and pSMAC (green
tracks), respectively (Fig. 4B and 4C). Importantly, these rates are very similar to the
rates of centripetal actin flow across the dSMAC and pSMAC, respectively (Fig. S5A1-
A3). Together, these observations argue that the pSMAC actomyosin arc network works
together with the dSMAC branched actin network to drive antigen centralization.

To identify the mechanism by which the actomyosin arcs drive antigen
centralization, we imaged F-actin and anti-IgM in the medial portion of forming synapses
at high magnification using TIRF-SIM. Anti-IgM microclusters were seen to move across
the pSMAC towards the cSMAC (which in the following images was in the down
direction) while embedded in an arc network moving in the same direction (Movie 10).
White lines in Movie 10 and in the corresponding still images in Fig. 4D1-D6 mark actin
arcs that were sweeping an individual anti-IgM microcluster inward (Fig. 4E1-E6).
Figure 4F shows the trajectory of this microcluster (temporally color-coded) as it moved
towards the cSMAC. Finally, a kymograph of this trajectory (Fig. 4G) shows that several
actin arcs contributed to the inward movement of this microcluster (areas bracketed in
white), and that pauses in movement (areas bracketed in pink) occurred where no actin
signal was immediately adjacent to the microcluster. Together, these results argue that
individual actin arcs move individual BCR: antigen microclusters inward via a sweeping
mechanism that likely depends on frictional coupling between the actin arc and the

microcluster (53-55). While arcs can slip past microclusters, the overall incidence of such slippage must be fairly small as the rate of inward antigen transport across the pSMAC (Fig. 4C) is not significantly slower than the rate of inward actin arc flow across the pSMAC (Fig. S5A3).

Integrin ligation-dependent IS formation requires myosin 2A contractility

B cells engaged with membrane-bound antigen at low density fail to centralize antigen unless their integrin LFA-1 is also engaged with ICAM-1 in the target membrane (18). As a prelude to investigating the myosin dependence of this integrin co-stimulatory effect, we sought to recapitulate these findings using primary B cells and PLBs containing varying amounts of mobile, fluorophore-labeled anti-IgM antibody in the presence or absence of unlabeled ICAM-1. Using this approach, we determined an amount of anti-IgM antibody that would not elicit robust antigen centralization in the absence of ICAM-1, but would in its presence. B cells exhibited robust antigen centralization/cSMAC formation over 10 minutes without the need for ICAM-1 when the PLB was loaded using a solution containing anti-IgM at a concentration of 2 µg/ml (hereafter referred to as "high density antigen") (Fig. S6A1-A3). By contrast, B cells formed antigen microclusters across their synaptic interface but failed to centralize them over 10 minutes when the PLB was loaded using a solution containing anti-IgM at a concentration 0.15 µg/ml (hereafter referred to as "low or limiting density antigen") (Fig. S6B1-B3). Importantly, when unlabeled ICAM-1 was included in these low-density antigen bilayers, B cells now exhibited robust antigen centralization/cSMAC formation (Fig. S6C1-C3). This co-stimulatory effect was supported by scoring antigen distribution as centralized, partially centralized or non-centralized (Fig. S6D1-D3 and E). It was also supported by scoring the percent of total synaptic antigen present within the cSMAC, which was defined by a circular area encompassing 20% of the entire synaptic interface and centered around the center of mass of the fluorescent antigen-containing pixels within the interface (Fig. S6F). Finally, it was supported by measuring the size of antigen clusters as a function of their distance from the center of the cSMAC (defined as above) (Fig. S6G). Specifically, B cells engaged with PLBs containing antigen at the limiting density and no ICAM-1 exhibited small antigen clusters (~0.3 µm^2) located roughly evenly across the synaptic interface (Fig. S6G, black trace), while B cells engaged with PLBs containing ICAM-1 in addition to antigen at the limiting density exhibited large antigen clusters (up to 3 µm^2), the largest of which were located at the center of the cSMAC (Fig. S6G, green trace). Of note, the total amount of antigen present at the synaptic interface was also greater for cells engaged with low density anti-IgM+ICAM-1 than for cells engaged with low density anti-IgM alone (Fig. S6H). Together, these results recapitulated a central aspect of the integrin co-stimulatory effect described by Carrasco *et al.* (18), and they established the specific conditions we used next to test the myosin dependence of this co-stimulatory effect.

To score the myosin dependence of the integrin co-stimulatory effect, we
measured the ability of primary B cells treated with either vehicle control (DMSO) or
pnBB to centralize antigen and form a cSMAC when engaged for 10 minutes with PLBs
containing ICAM-1 and anti-IgM at the limiting density. While DMSO-treated cells
exhibited robust antigen centralization/cSMAC formation (Fig. 5A1-A3), pnBB-treated
cells failed to centralize antigen/create a cSMAC (Fig. 5B1-B3). Consistently, the actin
arcs that surround centralized antigen in DMSO-treated cells (Fig. 5C1-C3; white
arrows) were absent in pnBB-treated cells (Fig. 5D1-D3). The fact that myosin inhibition
abrogates the integrin co-stimulatory effect was further supported by scoring antigen
distribution in control and pnBB-treated cells as centralized, partially centralized or non-
centralized (Fig. 5E), by scoring the percent of total synaptic antigen present within the
cSMAC (Fig. 5F), and by measuring the size of antigen clusters as a function of their
distance from the center of the cSMAC (Fig. 5G). Of note, the total amount of antigen
present at the synaptic interface was also greater for cells treated with DMSO than for
cells treated with pnBB (Fig. 5H). Together, these results show that the ability of integrin
ligation to promote antigen centralization and cSMAC formation when antigen is limiting
requires myosin contractility. This in turn argues that the contractile actomyosin arc
network created downstream of integrin ligation plays an important role in the
mechanism by which LFA-1 co-stimulation promotes B cell activation.

Finally, we were curious if the robust centralization of antigen that occurs in the
absence of LFA-1 ligation when the density of antigen is high is also dependent on
myosin contractility, at least to some extent. Indeed, we found that treatment with para-
amino BB (paBB), a newer, slightly more water soluble version of BB (56), attenuated
antigen centralization significantly even when the density of antigen was high (Fig. S6;
see legend for details), although the magnitude of the inhibition was smaller than for B
cells engaged with limiting antigen plus ICAM-1 (compare the results in Fig. S6 to the
results in Fig. 5). We conclude, therefore, that M2A contractility potentiates antigen
centralization when antigen density is high as well as when antigen density is low
enough that LFA-1 co-stimulation becomes important for IS formation.

Myosin 2A contractility promotes BCR-dependent signaling

To measure the contribution that actomyosin arcs might make to BCR-dependent
signaling, we determined the effect that pnBB has on the distribution and synaptic
content of phosphorylated CD79a (P-CD79a), an early signaling molecule responsible
for signal transduction downstream of BCR-antigen interaction (14, 57). Consistent with
results above and with the known properties of CD79a, DMSO-treated primary B cells
engaged for 10 minutes with PLBs containing ICAM-1 and limiting antigen and then
fixed/stained for P-CD79a exhibited robust cSMAC formation, with P-CD79a and anti-
IgM concentrated in the cSMAC (Fig. 6A1-A4). Also as expected, pnBB-treated B cells
failed to form a clear cSMAC, resulting in CD79a and anti-IgM spread across the

synapse (Fig. 6B1-B4). Importantly, quantitation showed that pnBB-treated cells also
exhibited a significant reduction relative to control cells in synaptic P-CD79a content
(Fig. 6C). This defect was also seen after only 5 minutes on PLBs (Fig. S8A), and the
defects at both time points were not due to differences between BB-treated cells and
control cells in synaptic CD79a content (Fig. S8B).

To extend these results, we determined the effect that pnBB has on the
distribution and synaptic content of phosphorylated CD19, an important co-receptor for
the BCR that is responsible for PI3K activation (3, 58-60). DMSO-treated primary B cells
engaged with PLBs as above exhibited robust cSMAC formation, with P-CD19 enriched
at the outer edge of the IgM concentrated in the cSMAC (Fig. 6D1-D4). This enrichment
of P-CD19 at the pSMAC/cSMAC boundary was confirmed by line scans of the
fluorescence intensities for F-actin, anti-IgM and P-CD19 (Fig. 6G, see boxed pSMAC
regions). In contrast to control cells, pnBB-treated B cells failed to concentrate anti-IgM
at the center of the synapse, and P-CD19 staining was now spread across the synaptic
interface (Fig. 6E1-E4 and H). Moreover, quantitation showed that pnBB-treated cells
also exhibited a significant reduction relative to control cells in synaptic P-CD19 content
(Fig. 6F) that was not due to a difference in synaptic CD19 content (Fig. S8C).
Together, these results indicate that the actomyosin arcs promote BCR-dependent
signaling.

Germinal center B cells can make actomyosin arcs and centralize antigen

Recent studies have presented evidence that germinal center (GC) B cells differ
markedly from naive B cells with regard to the distribution and fate of antigen at mature
synapses. Rather than concentrating antigen at the center of the synapse and using
actomyosin force to extract it there, GC B cells accumulate antigen in clusters at the
periphery of the synapse and use actomyosin force to extract it there (36, 61, 62).
These and other results argue that GC B cells differ dramatically from naïve B cells with
regard to the organization of actomyosin at their synapse. We wondered, however, if
actomyosin arcs could be detected in mouse GC B cells using our imaging approaches.
Consistently, TIRF-SIM imaging of mouse GC B cells isolated from the GFP-M2A
knockin mouse that were stained with Cell Mask Deep Red and plated on coverslips
coated with anti-IgM, anti-IgG and ICAM-1 revealed a subset of cells exhibiting
enrichment of M2A filaments in the medial, pSMAC portion of the synapse (Movie 11),
just as in naïve B cells. Moreover, these myosin filaments move centripetally (Movie 11)
and co-localize with pSMAC actin arcs in phalloidin-stained samples (Fig. 7A1-A3, white
arrows), just as in naive B cells. Importantly, scoring showed that about one third of GC
B cells exhibit robust accumulation of M2A filaments in the pSMAC (Fig. 7B). Similar
results were obtained when GFP-M2A knockin GC B cells were engaged for 10 minutes
with PLBs containing fluorophore-labeled anti-IgM/IgG and unlabeled ICAM-1, and then

fixed and stained with phalloidin, where about one third of the cells exhibited
actomyosin arcs in their pSMAC (Fig. 7C1-C4, white arrows, and D).

Given these results, we asked if our PLB-engaged mouse GC B cells can
centralize antigen. In partial agreement with previous findings (61, 62), ~45% of
synapses exhibited small to medium sized antigen clusters distributed to varying
degrees in the synapse periphery (Fig. 7E1, E2 and F). In addition, ~20% of synapses
exhibited antigen microclusters spread throughout the synaptic interface (Fig. 7E3 and
F). Importantly, the remaining ~35% of synapses exhibited highly centralized antigen
(Fig. 7E4 and F). We conclude, therefore, that GC B cells can make actomyosin arcs
and centralize antigen like naïve B cells, although the degree to which they do these
two things is considerably less than for naïve B cells.

Discussion

Integrin co-stimulation promotes B cell activation and IS formation when antigen is limiting by promoting B cell adhesion (18, 19). Here we identified an actomyosin-dependent component of this integrin co-stimulatory effect. By combining super-resolution imaging with specific cytoskeletal perturbations, we showed that integrin ligation induces the formation of a pSMAC actomyosin arc network that comprises the major actin network at the primary B cell IS. This network is created by the formin mDia1, organized into a concentric, contractile structure by the molecular motor M2A, and promotes synapse formation by mechanically sweeping antigen clusters centripetally into the cSMAC. Most importantly, we showed that integrin-dependent synapse formation under conditions of limiting antigen requires M2A, as inhibiting its contractility significantly impairs antigen centralization. Consistently, myosin inhibition also diminishes the synaptic content of the key BCR signaling proteins P-CD79a and P-CD19 and disrupts their synaptic distribution. Finally, we showed that a significant fraction of GC B cells also make this contractile pSMAC actomyosin arc network. Together, our results argue that integrin co-stimulation promotes B cell activation and synapse formation not only by enhancing B cell adhesion (18), but also by eliciting the formation of a contractile actomyosin arc network that drives mechanical force-dependent IS formation. These findings invite a critical "reset" for the way in which future B cell studies should be approached by highlighting the need for integrin co-stimulation when examining the role of actin during B cell activation. This reset is especially important given that most *in vitro* studies of B cell IS formation and activation have been performed under conditions of excess antigen, while antigen is rarely available in excess *in vivo*.

A central player in the link between integrin co-stimulation and the formation of the actomyosin arc network is almost certainly active RhoA. First, active RhoA would drive arc formation by simultaneously targeting, unfolding and activating mDia1 at the plasma membrane (63, 64). Second, active RhoA would drive arc organization and contractility by activating the ROCK-dependent phosphorylation of the regulatory light chains on M2A (43), thereby promoting the assembly of the M2A bipolar filaments that decorate, organize and contract the arcs. Finally, it is likely that active RhoA would promote actomyosin arc formation by activating the ROCK-dependent phosphorylation of mDia1's autoinhibitory domain, thereby blocking its refolding and subsequent inactivation (65-67). Given all this, it seems very likely that integrin ligation promotes actomyosin arc formation at least in part by promoting the loading of RhoA with GTP. Consistent with this idea, adhesion signaling has been linked in a variety of systems to the activation of guanine nucleotide exchange factors (GEFs) for RhoA (e.g. p190RhoGEF, GEF H1) (68, 69). Future work should seek, therefore, to clarify the outside-in signaling pathway in B cells that links integrin ligation to the activation of one

or more GEFs for RhoA. Such efforts should also take into account parallel activation
pathways, such as the PI3K-dependent activation of RhoA downstream of BCR
signaling (70), the myosin-dependent activation of B cell adhesion downstream of
CXCR5 signaling (71), and the diacylglycerol kinase-dependent regulation of adhesion
and actomyosin force generation at the B cell synapse (72). Given our results here, the
ability of the B-cell integrin VLA-4, which binds VCAM-1 on APCs, to promote IS
formation under limiting antigen conditions (19) may also involve an actomyosin-
dependent mechanism. Indeed, actomyosin-dependent B cell IS formation may be a
mechanism harnessed by multiple co-stimulatory pathways to promote B cell activation.

Consistent with our findings, a recent study by Bolger-Munro *et al.* reported that
GFP-tagged M2A localizes to the medial portion of synapses formed by A20 B cells
(13). In their hands, however, BB treatment did not inhibit antigen centralization, arguing
that synapse formation does not require M2A. The disparity between their results and
ours as regards the functional significance of M2A may be due to numerous differences
in experimental design, including the cell type used (primary B cells versus the A20 B
cell line), the mode of antigen presentation (anti-IgM-containing PLBs versus
transmembrane antigen expressed by APCs), and the density of antigen (known in
PLBs versus unknown and variable on APCs). Our pSMAC actomyosin arcs may also
be related to the myosin-rich regions that form in primary HEL-specific naïve B cells
bound to acrylamide gels coated with HEL antigen (47).

The contractile actomyosin structure identified here occupies the portion of the B-
cell synapse defined by the presence of an integrin ring, i.e. the pSMAC (3, 18, 19, 73).
This co-localization supports a feed-forward relationship where integrin co-stimulation
promotes the formation of the actomyosin arcs, and the contractile forces that these
arcs then exert promote further integrin activation and robust adhesion. Indeed, the B
cell pSMAC can be viewed as roughly analogous to the lamellar region of mesenchymal
cells, where integrins present within ECM-anchored focal adhesions are kept in their
open, extended, high-affinity conformation by the forces that myosin-rich stress fibers
exert on them (74, 75). By analogy, the contribution that the centering forces exerted by
the actomyosin arcs make to integrin activation in the pSMAC may be enhanced in the
context of an APC by the fact that the APC restricts ICAM-1 mobility (76).

Having established that naive B cells use actomyosin arcs to promote IS
formation, an obvious next question is whether they also use this contractile structure to
drive the extraction of membrane-bound antigens, a process known to require M2A (61,
77-79). Imaging synaptic actin and myosin in naïve B cells during the process of antigen
extraction should shed considerable light on this question. In that vein, Roper *et al.*
reported recently that the synapse of naïve B cells undergoing antigen extraction from
antigen-bearing plasma membrane sheets (PMSs) is composed of a dynamic mixture of

Arp2/3 complex-dependent actin foci and formin-dependent linear filaments/fibers (80).
While static images showed little co-localization between the actin foci and antigen
clusters, dynamic imaging suggested that the foci promote antigen extraction (although
the Arp2/3 complex and formins were both required for antigen uptake). Based on these
and other observations, Roper *et al.* concluded that naive B cells use a foci-filament
network to drive force-dependent antigen extraction (80). How M2A provides the force
in this context was unclear, however, as M2A (visualized using an antibody to the
phosphorylated form of M2A's RLC) did not co-localize with either actin structure.
Moreover, neither actin structure was affected by BB treatment. Of note, these two
findings are at odds with our findings here that M2A (visualized by endogenous tagging
of the M2A heavy chain) co-localizes extensively with actin arcs, and that BB treatment
profoundly disrupts the organization of the pSMAC actin arc network. Finally, we note
that the images of synaptic actin presented by Roper *et al.* look similar to our images of
naïve B cells stimulated with anti-IgM alone (i.e. without ICAM-1), where the synapse
was also composed of a disorganized and dynamic mixture of actin foci and short actin
filaments/fibers. The fact that the PMSs used by Roper *et al.* do not present any integrin
ligands may explain, therefore, why they did not see a more organized synapse
containing actomyosin arcs.

While the force-dependent extraction of antigens by naïve B cells commonly
initiates the process of antibody production, the force-dependent extraction of antigens
by GC B cells is thought to drive the subsequent process of antibody affinity maturation
by selecting for BCR: antigen interactions of higher affinity (2, 81). Consistently, GC B
cells possess higher levels of active M2A and exert stronger pulling forces than naive B
cells, allowing them to extract antigens with higher affinities for the BCR (61, 62).
Recent studies have shown that GC B cells accumulate antigen at the periphery of their
synapse in association with actin-rich surface projections that mediate antigen
extraction (61, 62). The striking difference between this behavior and that of naïve B
cells, which centralize antigen prior to extraction, argues that GC B cells must differ
dramatically from naïve B cells with regard to the organization of actomyosin at their
synapse. We found, however, that about one third of GC B cells formed actomyosin
arcs that were indistinguishable from those formed by naïve B cells. Moreover, about
one third of GC B cells centralized antigen (although about half exhibited peripheral
antigen clusters like those previously reported). While additional work is required to
prove that the subset of GC B cells with actomyosin arcs are the ones that centralize
antigen, this seems likely given our evidence here that actomyosin arcs drive antigen
centralization in naïve B cells. Future work will also be required to understand why GC
B cells vary with regard to actomyosin organization and the ability to centralize antigen
(e.g. dark zone versus light zone GCs). That said, our results argue that the manner in
which GC B cells harness actomyosin forces to extract antigen during the process of
antibody affinity maturation may not be restricted to the previously reported pathway

(61, 62). Finally, we note that the PLBs and PMSs employed in those studies to image
synaptic actin and antigen extraction did not contain integrin ligands. Given our results
here, this might explain why the GC B cells in those studies did not contain actomyosin
arcs.

Finally, future efforts should seek to define the role(s) played by the actomyosin
arcs described here in the process of extracting membrane-bound antigen, as this
process is impaired when M2A is inhibited or deleted (61, 77, 79). At a minimum, the
ability of the actomyosin arcs to centralize and corral antigen at the cSMAC should
promote antigen extraction no matter the exact extraction mechanism (15-17). As for a
more direct role, the ability of the actomyosin arcs to create a ring of strong adhesion
within the pSMAC that can grip the APC tightly may be required to extract antigens with
high affinity for the BCR, much as a tight grip on the wine bottle is required to pull out its
cork. This contractile structure also promotes both proximal signaling and the formation
of larger antigen clusters, two processes that are likely linked mechanistically through
the phase separation-dependent enhancement of signaling pathways (82). Whether the
pSMAC actomyosin arcs contribute directly to the upward pulling forces inside the B cell
that are thought to drive antigen extraction is, at the moment, unknown. Relevant to this
question, M2A was not detected in the dispersed actin foci thought to mediate antigen
extraction by naïve B cells engaged with antigen-bearing PMSs (80), or in the actin-rich,
invadopodia-like structures thought to mediate antigen extraction by naïve B cells
engaged with antigen-bearing acrylamide gels (47). Moreover, an association between
M2A and the peripheral actin-rich pods thought to mediate antigen extraction by GC B
cells engaged with antigen-bearing PMSs (62) has yet to be reported. We also note that
none of the activating surfaces used in these studies contained integrin ligands, and
that none of these studies examined antigen extraction in the context of a
physiologically relevant form of antigen presentation (e.g. via Fc or complement
receptors on the surface of a living cell). Given all this, further efforts to define the
mechanism by which M2A powers the extraction of membrane-bound antigens should
follow the myosin as the B cell extracts antigen from an APC, where the B cell's
integrins will be engaged, and where the antigen can be presented in a physiologically
relevant way. Perhaps only then will the role(s) played by M2A in antigen extraction be
fully revealed.
607

 Materials and Methods

Mice and cell culture
Primary B cells were isolated from the spleens of 6 to 12 week-old C57BL/6 mice
(Jackson Laboratories #002595) and M2A-GFP KI mice (gift of R. Adelstein,
NHLBI/NIH) of either sex using negative selection B cell isolation (StemCell
Technologies). Euthanasia was performed in accordance with protocols approved by
the National Human Genome Research Institute Animal Use and Care Committee at the
National Institutes of Health. The A20 murine IgG$^+$ B cell line was from ATCC (ATCC®
TIB-208™). B cells were cultured in complete medium (RPMI-1640, 10% heat-
inactivated fetal calf serum (FCS), 2 mM L-glutamine, 1 mM sodium pyruvate, 50 µM 2-
mercaptoethanol and 1X Antibiotic-Antimycotic) at 37 ^{o}C with 5% CO_2. Primary B cell
complete media also contains 5 ng/ml of BAFF (R&D Systems).

Plasmids and Reagents
GFP- and tdTomato-tagged F-Tractin were gifts from Michael Schell (Uniformed
Services University, Maryland). Alexa Fluor-conjugated phalloidins were purchased
from Thermo Fisher. Anti-mDia1 antibody was purchased from Thermo Fisher (PA5-
27607). HRP-conjugated mouse anti-β-actin antibody was purchased from Santa Cruz
(SC-47778 HRP). Rabbit anti-CD79a (#3351), anti-PCD79a (#5173), anti-CD19 (#3574)
and anti-PCD19 (#3571) were purchased from Cell Signaling Technologies. Anti-M2A
was purchased from Millipore Sigma (#M8064) CK-666 and SMIFH2 were purchased
from Millipore Sigma and used at final concentrations of 100 µM and 25 µM,
respectively. pnBB and paBB were purchased from Cayman Chemicals and used at a
final concentration of 25 µM. DMSO vehicle control was purchased from Millipore
Sigma. CellMask™ Deep Red Plasma Membrane Stain was purchased from Thermo
Fisher. Alexa Fluor 488- (#111-545-003), 594- (#111-585-003) and 647- (#111-605-003)
conjugated goat, anti-rabbit secondary antibodies were purchased from Jackson
ImmunoResearch. Goat anti-mouse IgG Fcγ fragment specific antibody (#115-005-008)
and goat anti-mouse IgM, µ-chain specific antibodies (#115-005-020) were purchased
from Jackson ImmunoResearch. Anti-rabbit-HRP (#32260) was purchased from Thermo
Fisher.

GC B cell generation and sorting
GC B cells were generated and sorted using a previously described protocol (83).
Briefly, 6 to 12 week-old M2A-GFP KI mice were immunized with sheep's red blood
cells. After 8 to 10 days, total B cells from the spleens and lymph nodes were isolated
using the Negative Selection B cell isolation kit (Stemcell Technologies) according to
the manufacturer's instructions. Dead cells were stained using Zombie Yellow viability
stain (Biolegend) and Fc receptors were blocked with the mouse TruStain FcX™
antibody (#156604). Cells were immunostained with anti-mouse CD38 (#102719), B220

648 (#103235) and GL-7 (#144617) purchased from Biolegend. GC B cells were sorted on a
649 BD Aria III FACs sorter (Beckton Dickinson) for GFP+, Zombie Yellow⁻, B220⁺, CD38^{low}
650 and GL-7⁺ cells, and were used immediately.
651

B cell transfection

A20 B cells and primary B cells were transfected as previously described (25). Briefly,
e*x vivo* primary B cells were first cultured for 12 h in complete media supplemented with
5 ng/ml BAFF (R&D Systems) and 2.5 µg/ml *Escherichia coli* O111:B4 LPS (Millipore
Sigma) (LPS was included to promote cell survival during nucleofection). $2x10^6$ B cells
were then nucleofected with 2 µg of plasmid DNA using Nucleofector Kit V (Lonza) and
rested for 16-24 hours using complete media containing 5 ng/ml BAFF and lacking LPS.
We refer to both rested, transfected cells and *ex vivo* non-manipulated cells as naïve B
cells because neither had been activated by antigen.
661

CRISPR

Mouse GFP-M2A and Scarleti-M2A template plasmids were gifts from Jordan Beach
(Loyola University, Chicago). Mouse M2A sgRNAs were synthesized by Synthego and
used according to the manufacturer's instructions. Briefly, sgRNAs were mixed with
Cas9 (New England Biolabs) to form ribonucleoproteins and then added together with
0.5 µg of template plasmid to $2x10^6$ cells suspended in the solution for Nucleofector kit
V. Following nucleofection, the cells were cultured in complete media for 24 hours
before Fluorescence-activated cell sorting (FACS) for GFP or Scarleti expression using
the Aria III (Becton Dickinson).
671

miRNA-mediated knockdown of mDia1

miRNAs targeting the 3' UTR of mouse mDia1 were designed as previously described
(84) using BLOCK-iT RNAi Designer (Thermo Fisher), synthesized (Gene Universal),
and fused to the C-terminus of mNeonGreen-F-Tractin using In-Fusion cloning (Takara).
A20 B cells were transfected with 2 µg of F-Tractin-mNeonGreen vector control or F-
Tractin-mNeonGreen-mDia1-miRNAs and cultured in complete media for 16 hrs. Cells
were then lysed and immunoblotted using an antibody to DIAPH1 (1:250; Thermo
Fisher, PA5-27607) and an HRP-conjugated antibody to β-actin (1:5000; Santa Cruz,
SC-47778 HRP) to confirm knockdown. Cells that had received the miRNA were
identified based on the expression of F-Tractin-mNeonGreen and then quantified. F-
Tractin-mNeonGreen-positive cells were also used in a cell spreading assay as
described above. See Supplementary information for additional details.
684

Cell spreading on functionalized glass

8-well Labtek chambers (Nunc) were coated with 15 µg/ml of anti-IgM and/or anti-IgG
with or without 0.5 µg/ml of mouse histidine-tagged ICAM-1 (Sino Biological) for 1 hour

at room temperature. B cells were resuspended in modified HEPES-buffered saline
(mHBS) (25) and adhered to functionalized glass for 15 min at 37°C before live-imaging
or fixing with 4% paraformaldehyde for staining (see SI Materials and Methods). Where
inhibitors were used, cells were pretreated for 30 min with 100 µM CK-666, 25 µM
SMIFH2, 25 µM pnBB or paBB, or dH_2O/DMSO vehicle control in mHBS at 37 ^{o}C. Cells
were then added to functionalized Labtek chambers in mHBS containing the same
concentrations of inhibitors or vehicle control as the pretreatment.
695
Supported planar lipid bilayers
Liposomes were prepared as described previously (26, 27, 85). Briefly, 0.4mM 1,2-
dioleoyl-*sn*-glycero-3-phosphocholine, biotin–CAP-PE, 1,2-dioleoyl-*sn*-glycero-3-[(*N*-(5-
amino-1-carboxypentyl)iminodiacetic acid)succinyl] (DGS)–NTA and 1,2-dioleoyl-sn-
glycero-3-phosphocholine (Avanti Polar Lipids, Inc.) were mixed at 1:3:96 molar % ratio.
Lipids were dried under a stream of argon and then desiccated in a vacuum chamber.
Unilamellar liposomes were generated from lyophilized lipids hydrated in Tris-buffered
saline via extrusion through a 50-nm pore membrane using a mini-extruder kit (Avanti
Polar Lipids, Inc.). PLBs were assembled in Sticky-Slide VI$^{0.4}$ Luer closed chambers
(Ibidi) as previously described (86). 25 × 75-mm glass coverslips (Ibidi) were cleaned
using Piranha solution (1:3 ratio of sulfuric acid and 30% hydrogen peroxide). After
depositing liposomes onto the flow channels, the channels were washed with HBS
buffer containing 1% BSA. A solution containing mono-biotinylated, Alexa Fluor 647-
labeled anti-IgM antibody (0.15 µg/ml for the limiting antigen condition and 2 µg/ml for
the high antigen condition) and streptavidin (Sigma-Aldrich) were added to the flow
chambers with or without 0.5 µg/ml unlabeled histidine-tagged ICAM-1. Anti-IgM
antibody (µ-chain specific) was monobiotinylated and labeled with Alexa Fluor 647
(Thermo Fisher) as described previously (18). The uniformity and lateral mobility of
PLBs were assessed using FRAP as described previously (26). Photo bleached spots
typically recovered within 60 seconds). B cells were resuspended in modified HEPES-
buffered saline and allowed to engage PLBs at 37°C and imaged immediately, or fixed
with 4% paraformaldehyde after 5 and 10 mins for immunostaining.
718
Traction force microscopy
Polyacrylamide gels (PA, 0.23 kPa shear modulus, 40 µm thickness) were prepared on
glass coverslips with embedded 40 nm fluorescent beads (TransFluoSpheres (633/720),
Thermo Scientific), as described previously (87). B cells were resuspended in mHBS
with 2% FCS and added to PA gels. Images of B cells that had engaged PA gels for 20
mins were captured. A no-stress reference image of the PA gels with beads was
captured after lifting cells from the PA gel by adding 1% sodium dodecyl sulfate in 1X
PBS to the imaging chamber at a final concentration of 0.04%. Particle image
velocimetry was used to calculate bead displacements relative to the reference position,

and the corresponding contractile energy was quantified using ImageJ plugins as
previously described (87, 88). Traction forces were reported as the mean magnitude of
traction stress within the cell relative to the cell surface area.
731

Immunostaining
733 Fixed cells were permeabilized with 0.2% Triton-X-100 and blocked for 30 min at room
734 temperature using PBS containing 2% BSA. Cells were incubated with primary
735 antibodies (1:200) overnight at 4°C and then secondary antibodies (1:250) with Alexa
736 Fluor-conjugated phalloidins for 1h at room temperature. Antibodies and phalloidins
737 were diluted in blocking buffer. All washes were performed with 1X PBS.
738

Microscopy
All live cell imaging was performed at 37 °C in mHBS supplemented with 2% FCS.
TIRF-SIM and 3D-SIM imaging were performed on a GE DeltaVision OMX SR
microscope (Cytiva) equipped with a 60X 1.42 NA oil objective (Olympus). For 3D-SIM,
z-stacks were acquired at 0.125 μm increments. Raw data were reconstructed using
Softworx software (Cytiva) with a Wiener filter constant of 0.002-0.003. Airyscan
imaging was performed using an LSM 880 Zeiss confocal microscope equipped with
Airyscan and using a Plan-Apochromat 63X 1.4 NA oil objective. Airyscan image
reconstruction was performed using Zeiss ZEN imaging software. TFM was imaged
using a Nikon Eclipse Ti2 microscope equipped with a 60X 1.2 NA water objective.
Linear adjustments to images were made using ImageJ 1.53 (NIH).
750

Image analyses
All image analysis was performed using ImageJ (NIH). To compare the relative fraction
of total IS footprint occupied by the cSMAC, pSMAC and dSMAC, the SMAC areas
were manually drawn based on actin morphology (dendritic for dSMAC, actin arc for
pSMAC, and hypodense for cSMAC) and then compared to the total IS area, which was
determined based on thresholds for F-actin intensity at the synaptic interface.
Fluorescence intensity within the SMAC regions was quantified using the ROIs from
above and reported as the total background-corrected fluorescence within the ROI,
which was quantified as described (89) using the following equation: Integrated density–
[(area of ROI)×(mean background fluorescence per unit area)], where the integrated
density is equal to [(area of ROI)×(mean fluorescence per unit area within the ROI)].
Mean background fluorescence was determined using the same ROI size at 3 separate
positions less than 3 μm away from the cell. The myosin fluorescence intensity in 3D
SIM images was quantified using a maximum projection image of the image stacks
where the cell ROI was determined based on the F-actin threshold and the background-
corrected myosin fluorescence within the cell ROI was reported. The FibrilTool plugin
for ImageJ was used to measure actin arc morphology as described previously (27, 45).

Briefly, the pSMAC regions in TIRF-SIM images were divided into 10-12 trapezoid-
shaped ROIs of similar size to measure the anisotropy of arcs in the radially symmetric
pSMAC. The values obtained range from 0, when the orientation of the structures is
random, to 1, when the structures show higher orientation in the same direction. The
velocity of centripetal actin flow was assessed by assembling kymographs from TIRF-
SIM videos using the Kymograph Builder plugin from ImageJ, as previously described
(27). Briefly, the dSMAC and pSMAC regions were identified by the relatively abrupt
slope change for F-actin flow, and slope angles were used to quantify the rates of actin
movement. The size of each antigen cluster and their relative distance from the cSMAC
center were quantified using an ImageJ macro. First, the perimeter of the synaptic
interface was determined based on thresholds for F-actin, and an ROI that
encompassed the interface area was drawn (the synaptic ROI). The coordinates for
each pixel contained in the ROI was determined and the linear distance of each pixel
from the center of mass of the total synaptic antigen (defined as cSMAC center) was
determined. The longest distance was defined as the furthest distance to travel from the
outermost edge of the cell. A binary image of the antigen channel combined with the
ImageJ watershed algorithm was used to segment individual antigen clusters within the
synaptic ROI. The area of each antigen cluster was quantified using the Analyze
Particles function in ImageJ. The relative distance of each antigen cluster was reported
as the distance between the center of mass of the antigen cluster and the cSMAC
center after normalizing to the furthest distance from the cell edge to the cSMAC center.
To quantify the antigen fluorescence in the cSMAC, a circular ROI corresponding to
20% of the total synaptic area (based on the average area of the cSMAC at the synaptic
interface) was drawn such that the center of the circle lies at the same coordinates as
the center of mass of the total antigen signal Antigen fluorescence within this circle was
quantified and presented as a percent of the total synaptic antigen fluorescence. The
fluorescence intensity of the signaling molecules CD79a, P-CD79a, CD19 and P-CD19
were all reported as the total fluorescence intensity within the synaptic ROI. All
fluorescence intensities were corrected for background as described above.
Fluorescence intensity profiles were obtained by drawing a 10 μm line across the center
of the synaptic interface and using the ImageJ function "Plot Profiles" to obtain
fluorescence intensity values across the line. The intensity profiles of several cells were
combined and the average fluorescence intensity ± standard deviation was reported.
The speeds of antigen cluster movement were quantified using the ImageJ plugin
TrackMate as previously described (90) where a combination of automated and manual
tracking were performed. Prior to quantification, the perimeter of the cell was identified
by over-saturating the signal for GFP-M2A, and the anti-IgM fluorescence signal outside
of the cell was removed so that only antigen clusters formed by that cell were quantified.
Antigen clusters were determined using a blob diameter of 0.2 $μm^2$ and tracks were
obtained using a threshold of 2000 units with sub-pixel localization. Mean antigen

cluster movement speeds were reported as distance traveled over time. Kymographs of
moving antigen clusters were created using the ImageJ plugin Kymograph Builder.
810

Statistical analyses
All statistical analyses were performed using Prism 9 (GraphPad). Statistical
comparisons of dot plots were performed using unpaired, two-tailed t-tests, and data are
represented as mean ± standard deviation. Statistical comparisons of bar charts were
performed using paired, two-tailed t-tests, and data are represented as mean ±
standard error of the mean. The following annotations are used to indicate significance:
* = $P<0.05$, ** = $P<0.01$, *** = $P<0.001$, and **** = $P<0.0001$.

Acknowledgments

This work was supported by the Intramural Research Program of the National Heart, Lung, and Blood Institute (NHLBI) (1ZIAHL006121-04 to JAH). The authors thank the NHLBI Flow Cytometry Core, Dr. Xuefei Ma for providing M2A-GFP mice, Dr. Il-Young Hwang for immunizing mice to prepare GC B cells, and Dr. Christopher J. Alexander for advice on miRNA design.

Competing interests

We have no competing interests to disclose.

 Figures

Figure 1. ICAM-1 co-stimulation promotes the formation of actin arcs at the B cell IS.

(A-F) GFP-F-Tractin-expressing primary B cells on glass coated with anti-IgM alone (A, B, E1, E2) or with anti-IgM+ICAM-1 (C, D, F1, F2) and imaged using Airyscan (A, C) or TIRF-SIM (B, D, E1, E2, F1, F2). The white arrows in A and B indicate the thin outer rim of dendritic actin in the dSMAC. The blue bars in A-D indicate the pSMAC. E2 and F2 correspond to the boxed regions in E1 and F1, respectively. (G) Percent of cells with pSMAC actin arcs (N>67 cells/condition from 3 experiments). (H, I) Percent of total synaptic F-actin (H) and percent of total IS footprint (I) contained within the dSMAC, pSMAC and cSMAC portions of the synapse for primary B cells on anti-IgG/ICAM-1-coated glass (N=44 cells/condition from 6 experiments). (J1, J2) GFP-F-Tractin-expressing A20 B cell on anti-IgG/ICAM-1-coated glass. J2 corresponds to the boxed region in J1. The red arrows in C, D and J1 indicate actin arcs. Scale bars: 10 μm.

Figure 2. The actin arcs are created by the formin mDia1 acting at the outer edge of the IS.

(A) GFP-F-Tractin-expressing primary B cell on anti-IgG/ICAM-1-coated glass. (B1, B2) Boxed regions in (A). (C1, C2) B1 and B2 with red lines applied to highlight linear actin filaments/bundles arising from surface spikes at the IS edge that are contiguous with actin arcs in the pSMAC. (D1, D2) GFP-F-Tractin-expressing primary B cell on anti-IgG/ICAM-1-coated glass before (D1) and 6 minutes after SMIFH2 addition (D2). (E) F-actin intensity profiles corresponding to the line scans in D1 (blue, before SMIFH2 addition) and D2 (red, after SMIFH2 addition). (F1-F4) GFP-F-Tractin-expressing A20 B cells transfected with vector-only or the indicated mDia1 miRNA constructs and activated on anti-IgG/ICAM-1-coated glass. (G) Ratio of pSMAC to dSMAC F-actin (N>20 cells/condition from 2 experiments). (H) pSMAC F-actin content (N=20-26 cells/condition from 2 experiments). A-C, and F: TIRF-SIM images; D: Airyscan images. Scale bars: 5 μm in A, D2, and F1; 2 μm in B1.

Figure 3. Myosin 2A decorates the actin arcs and is required for their concentric organization.

(A1-A5) Td-Tomato-F-Tractin expressing primary B cell from the M2A-GFP knockin mouse on anti-IgM/ICAM-1-coated glass. A4 and A5 correspond to the boxed regions in A1 and A2, respectively. (B1-B6) Still images at the indicated time points taken from a region within Movie 7 of a Td-Tomato-F-Tractin expressing primary B cell from the M2A-GFP knockin mouse. Different color arrowheads mark the formation and centripetal movement of individual M2A bipolar filaments (see text for details). (C, D) Phalloidin-

stained primary B cell from the M2A-GFP knockin mouse on glass coated with anti-IgM
alone (C) or with anti-IgM+ICAM-1 (D). (E) Total synaptic M2A content (N=91-115
cells/condition from 3 experiments). (F, G) GFP-F-Tractin-expressing primary B cells
that had been pretreated with DMSO (F) or pnBB (G) for 30 minutes and activated on
anti-IgM/ICAM-1-coated glass. (H) Anisotropy of the actin filaments/bundles present
within the pSMAC (see also Fig. S2H) (N=369-423 ROIs from 30-37 cells from 3
experiments). All panels: TIRF-SIM images. Scale bars: 3 µm in A3 and B6; 250 nm in
A5; 5 µm in D and G.

Figure 4. Actin arcs sweep antigen clusters centripetally.

(A1-A3) Phalloidin-stained (green) primary B cell 15 min after engagement with a PLB
containing unlabeled ICAM-1 and limiting anti-IgM (red). The white arrows in A1 and A3
mark the actin arcs. (B) Tracks of single anti-IgM microclusters traveling centripetally
across the dSMAC (red tracks) and pSMAC (green tracks) acquired from Movie 9. The
white line indicates the outer edge of this cell. (C) Mean speed of single anti-IgM
microclusters moving centripetally across the dSMAC and pSMAC (N=180-273 tracks
from 3 well-spread cells). (D1-D6) Still images at the indicated time points from Movie
10 showing the centripetal movement of actin arcs and a representative anti-IgM
microcluster (white arrows) (the center of the synapse is directly below the images).
Transparent white lines highlight the actin arcs that moved the microcluster
centripetally. (E1-E6) Same as D1-D6 except showing only the anti-IgM microcluster,
and indicating its centripetal path in blue. (F) Temporally pseudo-colored, projected
image of the anti-IgM microcluster in (D) and (E). (G) Kymograph of the 3 µm-long paths
taken by the microcluster and the actin arcs in (D) and (E) over a period of 400s. The
white brackets on the right indicate where actin arcs overlapped with and moved the
microcluster, while the red brackets indicate where the movement of the microcluster
stalled. A: Airyscan images; D-G: TIRF-SIM images. Scale bars: 5 µm in A3 and B; 300
nm in D6 and F.

Figure 5. Integrin ligation-dependent IS formation requires myosin 2A
contractility.

(A1-A3) DMSO-treated, phalloidin-stained primary B cells 15 min after engagement with
a PLB containing ICAM-1 and limiting anti-IgM. (B1-B3) Same as A1-A3 except the B
cells were treated with pnBB. (C1-C3) Images of a representative, DMSO-treated
primary B cell (white arrows mark actin arcs). (D1-D3) Images of a representative,
pnBB-treated primary B cell. (E) Percent of cells exhibiting centralized, partially
centralized and non-centralized antigen (see Figure S5D1-D3 for representative
examples of these three types of antigen distribution) (N=126-144 cells/condition from 3
experiments). (F) Percent of total synaptic antigen in the cSMAC (N=81-86
cells/condition from 3 experiments). (G) Antigen cluster size as a function of normalized

distance from the cSMAC center (N=113-144 cells/condition from 3 experiments). (H) Total synaptic antigen content (N=56-62 cells/condition from 3 experiments). All panels: Airyscan images. Scale bars: 10 µm in A1, B1, A3, and B3; 5 µm in D3.

Figure 6. Myosin 2A contractility promotes BCR signaling.

(A1-A4) DMSO-treated primary B cell 10 min after engagement with a PLB containing ICAM-1 and limiting anti-IgM, and stained for F-actin and P-CD79a. (B1-B4) Same as A1-A4 except the B cell was treated with pnBB. (C) Synaptic P-CD79a content (N=55-81 cells/condition from 3 experiments). (D1-D4) DMSO-treated primary B cell 10 min after engagement with a PLB containing ICAM-1 and limiting anti-IgM, and stained for F-actin and P-CD19. (E1-E4) Same as D1-D4 except the cell was treated with pnBB. (F) Synaptic P-CD19 content (N=115-140 cells/condition from 3 experiments). (G) Fluorescence intensities across synapses for P-CD19 (red), antigen (grey), and F-actin (green) in B cells treated with DMSO (N=22 cells from 2 experiments). The position of the pSMAC is highlighted in blue. (H) Same as G except the cells were treated with pnBB (N=16 cells from 2 experiments). All panels: Airyscan images. Scale bars: 5 µm in B4; 3 µm in E4.

Figure 7. GC B cells make actomyosin arcs.

(A1-A3) Phalloidin-stained primary GC B cell from the M2A-GFP knockin mouse on anti-IgM/anti-IgG/ICAM-1-coated glass. White arrows mark the actomyosin arcs. (B) Percent of cells on glass that did or did not show M2A enrichment in the pSMAC (N=140 cells from 4 experiments). (C) Phalloidin-stained primary GC B cell from the M2A-GFP knockin mouse 15 min after engagement with a PLB containing anti-IgM, anti-IgG, and ICAM-1. (D) Percent of cells on PLBs that did or did not show M2A enrichment in the pSMAC (N=89 cells from 4 experiments). (E1-E4) Representative images of the three types of anti-Ig distribution exhibited by GC B cells 15 min after engagement with a PLB containing anti-IgG and ICAM-1 (cell outlines are shown in blue). (F) Percent of GC cells displaying the three types of anti-Ig distribution shown in E1-E4 (N=157 cells from 6 experiments). All panels: TIRF-SIM images. Scale bars: 5 µm in A3; 3 µm in C4 and E4.

References

1. Forthal DN, Functions of Antibodies. *Microbiol. Spectr.* **2,** 1-17 (2014).
2. Heesters BA, van der Poel CE, Das A, & Carroll MC, Antigen Presentation to B Cells. *Trends in Immunology* **37,** 844-854 (2016). 10.1016/j.it.2016.10.003.
3. Harwood NE & Batista FD, The cytoskeleton coordinates the early events of B-cell activation. *Cold Spring Harb. Perspect. Biol.* **3,** a002360 (2011). 10.1101/cshperspect.a002360.
4. Gonzalez SF*, et al.*, Trafficking of B cell antigen in lymph nodes. *Annu. Rev. Immunol.* **29,** 215-233 (2011). 10.1146/annurev-immunol-031210-101255.
5. Carrasco YR & Batista FD, B cell recognition of membrane-bound antigen: an exquisite way of sensing ligands. *Curr. Opin. Immunol.* **18,** 286-291 (2006). 10.1016/j.coi.2006.03.013.
6. Cyster JG, B cell follicles and antigen encounters of the third kind. *Nat. Immunol.* **11,** 989-996 (2010). 10.1038/ni.1946.
7. Wang JC & Hammer JA, The role of actin and myosin in antigen extraction by B lymphocytes. *Semin. Cell Dev. Biol.* **102,** 90-104 (2020). 10.1016/j.semcdb.2019.10.017.
8. Song W, Liu C, & Upadhyaya A, The pivotal position of the actin cytoskeleton in the initiation and regulation of B cell receptor activation. *Biochim. Biophys. Acta* **1838,** 569-578 (2014). 10.1016/j.bbamem.2013.07.016.
9. Fleire SJ*, et al.*, B cell ligand discrimination through a spreading and contraction response. *Science* **312,** 738-741 (2006). 10.1126/science.1123940.
10. Tolar P, Sohn HW, Liu W, & Pierce SK, The molecular assembly and organization of signaling active B-cell receptor oligomers. *Immunol. Rev.* **232,** 34-41 (2009). 10.1111/j.1600-065X.2009.00833.x.
11. Mattila PK, Batista FD, & Treanor B, Dynamics of the actin cytoskeleton mediates receptor cross talk: An emerging concept in tuning receptor signaling. *J. Cell Biol.* **212,** 267-280 (2016). 10.1083/jcb.201504137.
12. Treanor B, Harwood NE, & Batista FD, Microsignalosomes: spatially resolved receptor signalling. *Biochem. Soc. Trans.* **37,** 1014-1018 (2009). 10.1042/BST0371014.
13. Bolger-Munro M*, et al.*, Arp2/3 complex-driven spatial patterning of the BCR enhances immune synapse formation, BCR signaling and cell activation. *Elife* **8,** (2019). 10.7554/eLife.44574.
14. Batista FD, Treanor B, & Harwood NE, Visualizing a role for the actin cytoskeleton in the regulation of B-cell activation. *Immunol. Rev.* **237,** 191-204 (2010). 10.1111/j.1600-065X.2010.00943.x.
15. Batista FD, Iber D, & Neuberger MS, B cells acquire antigen from target cells after synapse formation. *Nature* **411,** 489-494 (2001). 10.1038/35078099.
16. Yuseff MI, Pierobon P, Reversat A, & Lennon-Dumenil AM, How B cells capture, process and present antigens: a crucial role for cell polarity. *Nat. Rev. Immunol.* **13,** 475-486 (2013). 10.1038/nri3469.
17. Yuseff MI & Lennon-Dumenil AM, B cells use conserved polarity cues to regulate their antigen processing and presentation functions. *Front. Immunol.* **6,** 251 (2015). 10.3389/fimmu.2015.00251.
18. Carrasco YR, Fleire SJ, Cameron T, Dustin ML, & Batista FD, LFA-1/ICAM-1 interaction lowers the threshold of B cell activation by facilitating B cell adhesion and synapse formation. *Immunity* **20,** 589-599 (2004).
19. Carrasco YR & Batista FD, B-cell activation by membrane-bound antigens is facilitated by the interaction of VLA-4 with VCAM-1. *EMBO J.* **25,** 889-899 (2006). 10.1038/sj.emboj.7600944.

983   20.   Springer TA, Adhesion receptors of the immune system. *Nature* **346,** 425-434 (1990).
984   10.1038/346425a0.
985   21.   Springer TA, Dustin ML, Kishimoto TK, & Marlin SD, The lymphocyte function-associated LFA-1,
986   CD2, and LFA-3 molecules: cell adhesion receptors of the immune system. *Annu. Rev. Immunol.*
987   **5,** 223-252 (1987). 10.1146/annurev.iy.05.040187.001255.
988   22.   Wang JC, Bolger-Munro M, & Gold MR, Visualizing the Actin and Microtubule Cytoskeletons at
989   the B-cell Immune Synapse Using Stimulated Emission Depletion (STED) Microscopy. *J. Vis. Exp.*
990   (2018). 10.3791/57028.
991   23.   Freeman SA*, et al.*, Cofilin-mediated F-actin severing is regulated by the Rap GTPase and
992   controls the cytoskeletal dynamics that drive lymphocyte spreading and BCR microcluster
993   formation. *J. Immunol.* **187,** 5887-5900 (2011). 10.4049/jimmunol.1102233.
994   24.   Liu C*, et al.*, Actin reorganization is required for the formation of polarized B cell receptor
995   signalosomes in response to both soluble and membrane-associated antigens. *J. Immunol.* **188,**
996   3237-3246 (2012). 10.4049/jimmunol.1103065.
997   25.   Wang JC*, et al.*, The Rap1-cofilin-1 pathway coordinates actin reorganization and MTOC
998   polarization at the B cell immune synapse. *J. Cell Sci.* **130,** 1094-1109 (2017).
999   10.1242/jcs.191858.
26.   Yi J, Wu XS, Crites T, & Hammer JA, 3rd, Actin retrograde flow and actomyosin II arc contraction
drive receptor cluster dynamics at the immunological synapse in Jurkat T cells. *Mol. Biol. Cell* **23,**
834-852 (2012). 10.1091/mbc.E11-08-0731.
27.   Murugesan S*, et al.*, Formin-generated actomyosin arcs propel T cell receptor microcluster
movement at the immune synapse. *J. Cell Biol.* **215,** 383-399 (2016). 10.1083/jcb.201603080.
28.   Goode BL & Eck MJ, Mechanism and function of formins in the control of actin assembly. *Annu.*
Rev. Biochem. **76,** 593-627 (2007). 10.1146/annurev.biochem.75.103004.142647.
29.   Breitsprecher D & Goode BL, Formins at a glance. *J. Cell Sci.* **126,** 1-7 (2013). 10.1242/jcs.107250.
30.   Chen Q, Nag S, & Pollard TD, Formins filter modified actin subunits during processive elongation.
J. Struct. Biol. **177,** 32-39 (2012). 10.1016/j.jsb.2011.10.005.
31.   Rizvi SA*, et al.*, Identification and characterization of a small molecule inhibitor of formin-
mediated actin assembly. *Chem. Biol.* **16,** 1158-1168 (2009). 10.1016/j.chembiol.2009.10.006.
32.   Nishimura Y*, et al.*, The formin inhibitor SMIFH2 inhibits members of the myosin superfamily. *J.*
Cell Sci. **134,** (2021). 10.1242/jcs.253708.
33.   Burke TA*, et al.*, Homeostatic actin cytoskeleton networks are regulated by assembly factor
competition for monomers. *Curr. Biol.* **24,** 579-585 (2014). 10.1016/j.cub.2014.01.072.
34.   Lomakin AJ*, et al.*, Competition for actin between two distinct F-actin networks defines a
bistable switch for cell polarization. *Nat. Cell Biol.* **17,** 1435-1445 (2015). 10.1038/ncb3246.
35.   Fritzsche M, Erlenkamper C, Moeendarbary E, Charras G, & Kruse K, Actin kinetics shapes
cortical network structure and mechanics. *Sci. Adv.* **2,** e1501337 (2016).
10.1126/sciadv.1501337.
36.   Hammer JA, Wang JC, Saeed M, & Pedrosa AT, Origin, Organization, Dynamics, and Function of
Actin and Actomyosin Networks at the T Cell Immunological Synapse. *Annu. Rev. Immunol.* **37,**
201-224 (2019). 10.1146/annurev-immunol-042718-041341.
37.   Vicente-Manzanares M, Ma X, Adelstein RS, & Horwitz AR, Non-muscle myosin II takes centre
stage in cell adhesion and migration. *Nat. Rev. Mol. Cell Biol.* **10,** 778-790 (2009).
10.1038/nrm2786.
38.   Sellers JR, Myosins: a diverse superfamily. *Biochim. Biophys. Acta* **1496,** 3-22 (2000).
10.1016/s0167-4889(00)00005-7.
39.   Shutova MS & Svitkina TM, Common and Specific Functions of Nonmuscle Myosin II Paralogs in
Cells. *Biochemistry (Mosc)* **83,** 1459-1468 (2018). 10.1134/S0006297918120040.

40. Zhang Y*, et al.*, Mouse models of MYH9-related disease: mutations in nonmuscle myosin II-A.
Blood **119,** 238-250 (2012). 10.1182/blood-2011-06-358853.
41. Beach JR*, et al.*, Nonmuscle myosin II isoforms coassemble in living cells. *Curr. Biol.* **24,** 1160-
1166 (2014). 10.1016/j.cub.2014.03.071.
42. Beach JR*, et al.*, Actin dynamics and competition for myosin monomer govern the sequential
amplification of myosin filaments. *Nat. Cell Biol.* **19,** 85-93 (2017). 10.1038/ncb3463.
43. Beach JR & Hammer JA, 3rd, Myosin II isoform co-assembly and differential regulation in
mammalian systems. *Exp. Cell Res.* **334,** 2-9 (2015). 10.1016/j.yexcr.2015.01.012.
44. Kepiro M*, et al.*, para-Nitroblebbistatin, the non-cytotoxic and photostable myosin II inhibitor.
Angew. Chem. Int. Ed. Engl. **53,** 8211-8215 (2014). 10.1002/anie.201403540.
45. Boudaoud A*, et al.*, FibrilTool, an ImageJ plug-in to quantify fibrillar structures in raw microscopy
images. *Nat. Protoc.* **9,** 457-463 (2014). 10.1038/nprot.2014.024.
46. Wang J*, et al.*, Profiling the origin, dynamics, and function of traction force in B cell activation.
Sci. Signal. **11,** (2018). 10.1126/scisignal.aai9192.
47. Kumari A*, et al.*, Actomyosin-driven force patterning controls endocytosis at the immune
synapse. *Nat. Commun.* **10,** 2870 (2019). 10.1038/s41467-019-10751-7.
48. Gardel ML, Schneider IC, Aratyn-Schaus Y, & Waterman CM, Mechanical integration of actin and
adhesion dynamics in cell migration. *Annu. Rev. Cell. Dev. Biol.* **26,** 315-333 (2010).
10.1146/annurev.cellbio.011209.122036.
49. Case LB & Waterman CM, Integration of actin dynamics and cell adhesion by a three-
dimensional, mechanosensitive molecular clutch. *Nat. Cell Biol.* **17,** 955-963 (2015).
10.1038/ncb3191.
50. Comrie WA & Burkhardt JK, Action and Traction: Cytoskeletal Control of Receptor Triggering at
the Immunological Synapse. *Front. Immunol.* **7,** 68 (2016). 10.3389/fimmu.2016.00068.
51. Schnyder T*, et al.*, B cell receptor-mediated antigen gathering requires ubiquitin ligase Cbl and
adaptors Grb2 and Dok-3 to recruit dynein to the signaling microcluster. *Immunity* **34,** 905-918
(2011). 10.1016/j.immuni.2011.06.001.
52. Hashimoto-Tane A*, et al.*, Dynein-driven transport of T cell receptor microclusters regulates
immune synapse formation and T cell activation. *Immunity* **34,** 919-931 (2011).
10.1016/j.immuni.2011.05.012.
53. Yu CH, Wu HJ, Kaizuka Y, Vale RD, & Groves JT, Altered actin centripetal retrograde flow in
physically restricted immunological synapses. *PLoS One* **5,** e11878 (2010).
10.1371/journal.pone.0011878.
54. Ditlev JA*, et al.*, A composition-dependent molecular clutch between T cell signaling
condensates and actin. *Elife* **8,** (2019). 10.7554/eLife.42695.
55. Smoligovets AA, Smith AW, Wu HJ, Petit RS, & Groves JT, Characterization of dynamic actin
associations with T-cell receptor microclusters in primary T cells. *J. Cell Sci.* **125,** 735-742 (2012).
10.1242/jcs.092825.
56. Varkuti BH*, et al.*, A highly soluble, non-phototoxic, non-fluorescent blebbistatin derivative. *Sci.
Rep.* **6,** 26141 (2016). 10.1038/srep26141.
57. Tanaka S & Baba Y, B Cell Receptor Signaling. *Adv. Exp. Med. Biol.* **1254,** 23-36 (2020).
10.1007/978-981-15-3532-1_2.
58. Tuveson DA, Carter RH, Soltoff SP, & Fearon DT, CD19 of B cells as a surrogate kinase insert
region to bind phosphatidylinositol 3-kinase. *Science* **260,** 986-989 (1993).
10.1126/science.7684160.
59. Keppler SJ*, et al.*, Wiskott-Aldrich Syndrome Interacting Protein Deficiency Uncovers the Role of
the Co-receptor CD19 as a Generic Hub for PI3 Kinase Signaling in B Cells. *Immunity* **43,** 660-673
(2015). 10.1016/j.immuni.2015.09.004.

60. Depoil D*, et al.*, CD19 is essential for B cell activation by promoting B cell receptor-antigen
microcluster formation in response to membrane-bound ligand. *Nat. Immunol.* **9,** 63-72 (2008).
10.1038/ni1547.
61. Nowosad CR, Spillane KM, & Tolar P, Germinal center B cells recognize antigen through a
specialized immune synapse architecture. *Nat. Immunol.* **17,** 870-877 (2016). 10.1038/ni.3458.
62. Kwak K*, et al.*, Intrinsic properties of human germinal center B cells set antigen affinity
thresholds. *Sci. Immunol.* **3,** (2018). 10.1126/sciimmunol.aau6598.
63. Kuhn S & Geyer M, Formins as effector proteins of Rho GTPases. *Small GTPases* **5,** e29513
(2014). 10.4161/sgtp.29513.
64. Rose R*, et al.*, Structural and mechanistic insights into the interaction between Rho and
mammalian Dia. *Nature* **435,** 513-518 (2005). 10.1038/nature03604.
65. Nezami AG, Poy F, & Eck MJ, Structure of the autoinhibitory switch in formin mDia1. *Structure*
14, 257-263 (2006). 10.1016/j.str.2005.12.003.
66. Maiti S*, et al.*, Structure and activity of full-length formin mDia1. *Cytoskeleton (Hoboken, N.J.)*
69, 393-405 (2012). 10.1002/cm.21033.
67. Staus DP, Taylor JM, & Mack CP, Enhancement of mDia2 activity by Rho-kinase-dependent
phosphorylation of the diaphanous autoregulatory domain. *Biochem. J.* **439,** 57-65 (2011).
10.1042/BJ20101700.
68. Guilluy C*, et al.*, The Rho GEFs LARG and GEF-H1 regulate the mechanical response to force on
integrins. *Nat. Cell Biol.* **13,** 722-727 (2011). 10.1038/ncb2254.
69. Lawson CD & Burridge K, The on-off relationship of Rho and Rac during integrin-mediated
adhesion and cell migration. *Small GTPases* **5,** e27958 (2014). 10.4161/sgtp.27958.
70. Saci A & Carpenter CL, RhoA GTPase regulates B cell receptor signaling. *Mol. Cell* **17,** 205-214
(2005). 10.1016/j.molcel.2004.12.012.
71. Saez de Guinoa J, Barrio L, Mellado M, & Carrasco YR, CXCL13/CXCR5 signaling enhances BCR-
triggered B-cell activation by shaping cell dynamics. *Blood* **118,** 1560-1569 (2011).
10.1182/blood-2011-01-332106.
72. Merino-Cortes SV*, et al.*, Diacylglycerol kinase zeta promotes actin cytoskeleton remodeling and
mechanical forces at the B cell immune synapse. *Sci. Signal.* **13,** (2020).
10.1126/scisignal.aaw8214.
73. Dustin ML, Chakraborty AK, & Shaw AS, Understanding the structure and function of the
immunological synapse. *Cold Spring Harb. Perspect. Biol.* **2,** a002311 (2010).
10.1101/cshperspect.a002311.
74. Nordenfelt P, Elliott HL, & Springer TA, Coordinated integrin activation by actin-dependent force
during T-cell migration. *Nat. Commun.* **7,** 13119 (2016). 10.1038/ncomms13119.
75. Parsons JT, Horwitz AR, & Schwartz MA, Cell adhesion: integrating cytoskeletal dynamics and
cellular tension. *Nat. Rev. Mol. Cell Biol.* **11,** 633-643 (2010). 10.1038/nrm2957.
76. Comrie WA, Li S, Boyle S, & Burkhardt JK, The dendritic cell cytoskeleton promotes T cell
adhesion and activation by constraining ICAM-1 mobility. *J. Cell Biol.* **208,** 457-473 (2015).
10.1083/jcb.201406120.
77. Natkanski E*, et al.*, B cells use mechanical energy to discriminate antigen affinities. *Science* **340,**
1587-1590 (2013). 10.1126/science.1237572.
78. Spillane KM & Tolar P, Mechanics of antigen extraction in the B cell synapse. *Mol. Immunol.* **101,**
319-328 (2018). 10.1016/j.molimm.2018.07.018.
79. Hoogeboom R*, et al.*, Myosin IIa Promotes Antibody Responses by Regulating B Cell Activation,
Acquisition of Antigen, and Proliferation. *Cell Rep.* **23,** 2342-2353 (2018).
10.1016/j.celrep.2018.04.087.

80. Roper SI*, et al.*, B cells extract antigens at Arp2/3-generated actin foci interspersed with linear
filaments. *Elife* **8,** (2019). 10.7554/eLife.48093.
81. Spillane KM & Tolar P, B cell antigen extraction is regulated by physical properties of antigen-
presenting cells. *J. Cell Biol.* **216,** 217-230 (2017). 10.1083/jcb.201607064.
82. Huang WYC*, et al.*, A molecular assembly phase transition and kinetic proofreading modulate
Ras activation by SOS. *Science* **363,** 1098-1103 (2019). 10.1126/science.aau5721.
83. Hwang IY*, et al.*, An essential role for RGS protein/Galphai2 interactions in B lymphocyte-
directed cell migration and trafficking. *J. Immunol.* **194,** 2128-2139 (2015).
10.4049/jimmunol.1401952.
84. Alexander CJ & Hammer JA, 3rd, Optimization of cerebellar purkinje neuron cultures and
development of a plasmid-based method for purkinje neuron-specific, miRNA-mediated protein
knockdown. *Methods Cell Biol.* **131,** 177-197 (2016). 10.1016/bs.mcb.2015.06.004.
85. Hong J, Murugesan S, Betzig E, & Hammer JA, Contractile actomyosin arcs promote the
activation of primary mouse T cells in a ligand-dependent manner. *PLoS One* **12,** e0183174
(2017). 10.1371/journal.pone.0183174.
86. Comrie WA, Babich A, & Burkhardt JK, F-actin flow drives affinity maturation and spatial
organization of LFA-1 at the immunological synapse. *J. Cell Biol.* **208,** 475-491 (2015).
10.1083/jcb.201406121.
87. Jaumouille V, Cartagena-Rivera AX, & Waterman CM, Coupling of beta2 integrins to actin by a
mechanosensitive molecular clutch drives complement receptor-mediated phagocytosis. *Nat.*
Cell Biol. **21,** 1357-1369 (2019). 10.1038/s41556-019-0414-2.
88. Martiel JL*, et al.*, Measurement of cell traction forces with ImageJ. *Methods Cell Biol.* **125,** 269-
287 (2015). 10.1016/bs.mcb.2014.10.008.
89. Burgess A*, et al.*, Loss of human Greatwall results in G2 arrest and multiple mitotic defects due
to deregulation of the cyclin B-Cdc2/PP2A balance. *Proc. Natl. Acad. Sci.* **107,** 12564-12569
(2010). 10.1073/pnas.0914191107.
90. Tinevez JY*, et al.*, TrackMate: An open and extensible platform for single-particle tracking.
Methods **115,** 80-90 (2017). 10.1016/j.ymeth.2016.09.016.

1154

1155

1156

Figure 1

Anti-IgM

Airyscan
A Primary B cell
F-Tractin

TIRF-SIM
B

Anti-IgM + ICAM-1

Airyscan
C

TIRF-SIM
D

Anti-IgM
E1 Primary B cell
F-Tractin

E2
cSMAC? pSMAC? dSMAC

Anti-IgM+ICAM-1
F1

F2
cSMAC pSMAC dSMAC

G

Percent cells with actin arcs (%)

Anti-IgM   Anti-IgM +ICAM-1

H

Percent of total synaptic F-actin

dSMAC   pSMAC   cSMAC

I

Percent of total IS footprint

dSMAC   pSMAC   cSMAC

J1 A20 B cell
F-Tractin

J2
dSMAC   pSMAC   cSMAC

Figure 2

A Primary B cell   B1   B2   Before SMIFH2   +SMIFH2 6'
F-Tractin   C1   C2   D1 Primary B cell   D2
                                    F-Tractin

E dSMAC pSMAC cSMAC

F-actin (AUx10³)

Before
+SMIFH2

Distance (µm)

F1 A20 B cell   F2 miRNA#1   F3 miRNA#2   F4 miRNA#3
F-Tractin
Vector

G

Ratio of pSMAC to
dSMAC F-actin

****
****
****

Vector miRNA#1 miRNA#2 miRNA#3

H

pSMAC F-actin
content (AUx10⁵)

****
****
****

Vector miRNA#1 miRNA#2 miRNA#3

Figure 3

A3 F-Tractin

A2 GFP-M2A

A1 F-Tractin GFP-M2A

A4 GFP-M2A F-Tractin

A5 300 nm GFP-M2A

B1 0s B2 33s B3 75s B4 123s B5 177s B6 255s

C Anti-IgM

D Anti-IgM+ICAM-1

Phalloidin GFP-M2A

E ****

Total synaptic M2A content (AU×10³)

Anti-IgM   Anti-IgM +ICAM-1

F DMSO F-Tractin

G pnBB

H DMSO pnBB

Number of ROIs

Anisotropy

Figure 4

Figure 5

Limiting Anti-IgM + ICAM-1 + DMSO

Limiting Anti-IgM + ICAM-1 + pnBB

Figure 6

Figure 7