Learning from the path not taken: Sensory prediction error computation and implicit motor adaptation proceed without movement

Olivia A. Kim[1], Alexander D. Forrence[2], and Samuel D. McDougle[2]

1. Department of Psychology, Princeton University, Princeton, NJ 08544
2. Department of Psychology, Yale University, New Haven, CT 06511

Correspondence to:
    Olivia Kim
    oakim@princeton.edu
    kim.olivia.a@gmail.com

Number of Pages: 29
Number of Figures in Main Text: 3
Number of Supplemental Figures: 4
Number of Supplemental Tables: 3
References: 60 (45 main text, 15 methods-only)
Number of Characters (including spaces):
    Abstract (1,293)
    Highlights (306)
    Main text, including main figure legends (27,858)
    Methods (15,013)

AUTHOR CONTRIBUTIONS
Conceptualization, OK, AF, & SM; Methodology, OK, AF, & SM; Software, OK & AF; Validation, OK & AF; Formal Analysis, OK & AF; Investigation, OK & AF; Resources, OK & SM; Data Curation, OK & AF; Supervision, SM; Project Administration, SM; Funding Acquisition, OK & SM; Writing – original draft, OK; Writing – review & editing, OK, AF, & SM

CONFLICT OF INTEREST
The authors declare no competing financial interests.

ACKNOWLEDGEMENTS
This work was supported by a grant from the National Institutes of Health to OK (F32-NS122921).

ABSTRACT

Prediction errors guide many forms of learning, providing teaching signals that help us improve our

performance. Implicit motor adaptation, for instance, is driven by sensory prediction errors (SPEs), which

occur when the expected and observed consequences of a movement differ. Traditionally, SPE

computation is thought to require movement execution. However, recent work suggesting that the brain

generates and accounts for sensory predictions based on motor imagery or planning alone calls this

assumption into question. Here, by measuring implicit adaptation during a visuomotor task, we tested

whether motor planning and well-timed sensory feedback are sufficient for SPE computation. Human

participants were cued to reach to a target and were, on a subset of trials, rapidly cued to withhold these

movements. Errors displayed both on trials with and without movements induced single-trial implicit

learning. Learning following trials without movements persisted even when movement trials had never been

paired with errors, and when the direction of movement and sensory feedback trajectories were decoupled.

These observations demonstrate that the brain can compute SPEs without generating overt movements,

leading to the adaptation of planned movements even when they are not performed.

HIGHLIGHTS

• The brain can learn to update movements that are not performed, representing a mechanism for

learning based only on movement planning and sensory expectation.

• Supports a fundamental role for prediction in adaptation.

• Provides further support for the role of forward models in predictive motor control.

KEYWORDS

predictive coding, forward model, mental imagery, supervised learning

INTRODUCTION

Prediction errors help to optimize behavior by driving learning processes that correct for our

mistakes. Accordingly, they are thought to be a fundamental computation of the nervous system.[1] Specific

types of prediction errors are associated with specific learning processes, and sensory prediction errors

(SPEs) are the teachers of the implicit motor system, triggering adaptation and refinement of movements.[2–

6] Traditionally, SPEs are thought to be computed by comparing the expected and observed consequences

of a movement, with movement execution being a critical element in this process.[4,7] However, the

predictions factoring into SPE computation are thought to be generated by an internal forward model, which

estimates the consequences of movements before relevant sensory feedback can reach the brain.[7] This

predictive capacity suggests that the forward model predicts the sensory consequences of an intended

movement regardless of whether that movement occurs. That is, because movements are synchronous

with the sensory outcomes that they produce, motor execution cannot itself be necessary for the generation

of predictions by the forward model.

Recent work supports the claim that the brain generates predictions related to the sensory

consequences of movements before they can be performed, and even when there is never any intention to

move.[9–12] Considering that these studies imply that movements are not needed for the generation of

sensorimotor predictions, it may be that these predictions can be combined with sensory observations to

support SPE computation without actual motor execution. In other words, SPEs should be effectively

computed based upon only two events – generation of a sensory prediction and observation of sensory

feedback (**Fig. 1a**). Here, we used implicit motor adaptation as a readout for SPE occurrence, as SPEs are

both necessary and sufficient for this form of learning, and asked if motor planning and subsequent sensory

events are sufficient to drive SPE computation without movement.[5,13–18] While prior work has illustrated that

cognitive processes support visuomotor learning without movement by observers who witnesses others'

motor errors, these processes may be recruited by performance errors beyond SPE, or by a combination

of multiple sources of error and multiple learning processes.[19–21] Here, we isolate implicit motor learning to

specifically test whether SPE computation requires movement execution.

Figure 1. Schematics showing the proposed learning framework and task design. (**a**) Schematic
showing how the forward model may support implicit motor adaptation in the presence of sensory feedback
not causally related to self-generated movement. (**b**) Schematic of events on trials with visual feedback.
The robotic apparatus brought the participant's hand to the starting location to initiate a trial. On Movement
trials (top), the target turned green ('GO'), cueing participants to reach through the target. On trials with
visual feedback, participants observed a white feedback cursor move along a rotated trajectory ('Rotation').
On No-Movement trials (bottom), the target turned magenta 100 ms after turning green, cueing participants
to withhold movement ('STOP'). After a delay, an animation showing the feedback cursor moving 15° off-
target played ('Animation'). The hand is shown in the figure for illustrative purposes but was not visible
during the experiment. (**c**) Schematic showing the trials that constituted a triplet and how single-trial learning
(STL) was computed. Triplets were composed of 2 Movement trials without visual feedback flanking either
a Movement or a No-Movement trial with visual feedback. STL was measured as the difference between
reach angles on the flanking trials. (**d**) Schematic showing the pseudorandomized order in which trials were
presented for an example participant, with color reflecting the movement condition (Movement: green, No-
Movement: magenta).

We measured trial-by-trial implicit adaptation during a visuomotor task in which human participants

saw visual feedback while performing – or withholding – hand and arm movements, using a modified stop-

signal paradigm. At the onset of each trial, a "Go" cue was presented, instructing participants to make a

movement through a visual target. On some trials, a "Stop" cue was presented shortly after presenting the

"Go" cue, instructing participants to withhold the movement ("No-Movement" trials, **Fig. 1b**). We reasoned

that presenting the "Go" cue on "Stop" trials would be a sufficient stimulus to induce prediction generation

and support sensory prediction error computation, given that prior work has shown that cued – but ultimately

withheld – movements can still serve as an input to a forward model and support the generation of

differential sensory predictions.[12] To isolate implicit adaptation, we employed a recently-developed

approach that requires participants to aim their movements directly for presented targets and disregard

visual feedback.[17,22–26] We predicted that single-trial motor adaptation would occur following both typical

movement (Go) trials that generated sensory error, as well as trials where movements were withheld (Stop)

but "simulated" sensory errors were observed. If confirmed, this result would demonstrate that SPEs can

be computed in the absence of movement and can cause adaptation of planned but not performed

movements. Furthermore, such a result would provide strong support for predictive models of motor

adaptation.

RESULTS

Simulated and typical visuomotor rotations cause motor adaptation

In our first study, we measured implicit motor adaptation in humans (n = 20) performing straight

reaches during a visuomotor adaptation task (**Fig. 1b**). Vision of the hand and arm was occluded by a mirror

that reflected visual feedback from a horizontally mounted monitor. A white cursor provided feedback about

participants' hand positions as they reached from a starting location to a displayed target. After a brief

acclimation period, trials were organized into triplets, such that each trial with cursor feedback was flanked

by trials without cursor feedback. This allowed for a reliable measurement of single-trial learning (STL) in

response to feedback, quantified as the difference between hand angles on the first and third trials of each

triplet (**Fig. 1c**). Trials with cursor feedback were either "Movement" trials on which a "Go" signal prompted

movement or No-Movement trials on which a "Stop" signal indicated that movements should be withheld.

On Movement trials, feedback involved a visuomotor error (±15° rotation added to the visual cursor path; +

= clockwise; **Fig. 1b right**). On No-Movement trials, sensory feedback involved a simulation of the cursor's

path, using timing variables based on ongoing measurements of participant behavior (see *Methods*). All

flanking trials of each triplet were "Go" trials and thus required movements. The direction of the error

(clockwise [CW] or counterclockwise [CCW]) was pseudorandomly varied across triplets to maintain overall

adaptation near 0 throughout the session (**Fig. 1d**). This straightforward design allowed us to test the

hypothesis that SPE computation and motor adaptation do not require movement and sensory feedback to

be causally linked (**Fig. 1a**).

Consistent with our predictions, rotated cursor paths on Movement and No-Movement trials both

caused subsequent hand trajectories to shift opposite the direction of the rotation (**Fig. 2b**-**c**), with a two-

way repeated measures ANOVA revealing statistically significant main effects of the direction (CW vs CCW)

of the perturbation applied ($F(1, 19) = 98.62$, $p = 5.89$ x 10^{-9}, $\eta_G^2 = 0.76$). While there was no significant main

effect of withholding movement ($F(1, 19) = 1.79$, $p = 0.20$), we observed a significant interaction between

the perturbation applied and withholding movement ($F(1, 19) = 137.32$, $p = 3.87$ x 10^{-10}, $\eta_G^2 = 0.49$). Post-

hoc pairwise comparisons confirmed that STL was sensitive to perturbation direction during both Movement

(paired t-test: $t(19) = 12.92$, $p_{adj} = 2.96$ x 10^{-10}, *Cohen's d* = 5.12) and No-Movement triplets ($t(19) = 4.39$,

$p_{adj} = 3.13$ x 10^{-4}, *Cohen's d* = 1.63), and indicated that STL magnitude was greater across Movement than

No-Movement triplets (paired-samples signed-rank test, CW rotations: $V = 210$, $p_{adj} = 2.55$ x 10^{-6}, $r = 0.88$;

CCW rotations: $t(19) = 9.43$, $p_{adj} = 2.70$ x 10^{-8}, *Cohen's d* = 2.02) rotations. However, we note that the

amplitude of adaptation observed both with and without movement was within the range of implicit learning

rates measured in previous visuomotor rotation studies (**Supplemental Fig. 1**).

Figure 2. Effects of typical and simulated errors during a visuomotor reach adaptation task. (**a**) Schematic illustrating how STL and remembered STL measurements were computed. (**b**) An example subject's (top) and the group's (bottom) mean ± SEM changes in reach paths across triplets with a rotation applied (green: triplets with perturbations on Movement trials, magenta: triplets with perturbations on No-Movement trials, solid lines: perturbation was a CW rotation, dashed lines: perturbation was a CCW rotation). (**c**) STL across Movement (green) and No-Movement (magenta) triplets for all participants (*n* = 20). In the reference frame used here, positive changes in hand angle are CCW. Please refer to Supplemental Table 1 for details on all statistical tests. (**d**) Group mean ± SEM Δhand angle values after exposure to Movement (green) and No-Movement (magenta) trial perturbations. Positive Δ values indicate that the change in hand angle proceeded opposite the direction of the perturbation (*i.e.,* the direction that would counter the error, "Right-Way"). (**e**) Group mean of participants' ratios of remembered STL to initial STL during Movement and No-Movement trials. (**f**) The relationship between Right-Way STL observed during Movement and No-Movement triplets. (**g**) As in (**f**), but for STL observed on trials where adaptation proceeded in the direction that would exacerbate the error (*i.e.,* the same direction as the perturbation applied, "Wrong Way"). Statistical significance (* = $p_{adj} < 0.05$; n.s. = $p_{adj} \geq 0.05$) is indicated. Abbreviations: STL – single-trial learning, CW – clockwise, CCW – counterclockwise, Δ – change in.

Supplemental Figure 1. Learning rates reported in the literature and observed in the current study. Learning rates for motor adaptation observed in previous studies are shown at left in black, and learning rates observed in each experiment in the current report are shown at the right, with data from Movement triplets shown in green and data from No-Movement triplets shown in magenta. Data are shown as mean ± SEM, and are shown for rotational/error clamp perturbations of 15°, with the exception of Wei & Kording, 2009 E2, where an 11° perturbation was applied. Papers referred to and their corresponding reference numbers: Kim et al., 2018[23]; Morehead et al., 2017[17]; Wei & Körding, 2009[27]. "Rotation, Go" and "Rotation, Stop" show data from the in-lab experiment where participants saw 15° rotated feedback on Movement trials (i.e., data from Fig. 2), "Rotation online, Go" and "Rotation online, Stop" show data from the online experiment where participants saw 0-15° rotated feedback on Movement trials (i.e., data from Supplemental Fig. 2). "Clamp online, Go" and "Clamp online, Stop" show data from the online experiment where participants saw 0-15° error-clamed feedback (i.e., data from Supplemental Fig., 4). "Rotation 0 go" and "Clamp 0 go" show data from the online experiments where participants saw 0° perturbed feedback on Movement trials. Abbreviations: E, experiment.

To address whether the STL measured represented genuine implicit learning, we checked whether

adaptation persisted beyond the trial after an error was experienced. We examined participants' hand

angles on the second trial after a perturbation relative to the pre-perturbation baseline trial (i.e., hand angle

on trial 1 of triplet $t + 1$ relative to hand angle on trial 1 of triplet t, subsequently referred to as remembered

STL, **Fig. 2a**). Because visual feedback was withheld on both trial types, this approach provided a pure

measure of persistent memory in the absence of error-driven changes in performance. Hand angle

remained adapted in the direction opposite the rotation on trials with nonzero perturbations regardless of

movement condition (**Fig. 2d**), suggesting that genuine implicit learning was observed in response to errors

under both movement conditions. Closer examination of the relative ratio of remembered STL to initial STL

revealed that retention of adaptation differed significantly from zero after both Movement (t(19) = 26.20, p_{adj}

= 6.71 x 10^{-16}, *Cohen's d* = 5.86) and No-Movement triplets (t(19) = 9.20, p_{adj} = 2.98 x 10^{-8}, *Cohen's d* =

2.06). Moreover, the amount of retention observed was not statistically significantly different between the

movement conditions ($t(19) = 2.07$, $p_{adj} = 0.053$, **Fig. 2e**).

To assess the potential similarity of mechanisms underlying adaptation after errors on Movement

and No-Movement trials, we compared STL amplitude under each condition, reasoning that there should

be a reliable relationship between the two measures if STL is supported by the same mechanism for both

Movement and No-Movement trials. When we considered instances of STL in the direction that would

compensate for the observed error (the direction opposite the rotation, i.e., the "Right-Way"), within-subject

changes in hand angle were correlated between Movement and No-Movement trials (*Pearson's r* = 0.55, *p*

= 0.01; **Fig. 2f**). Conversely, changes in hand angle in the direction that would exacerbate the observed

error (the direction of the rotation, i.e., the "Wrong-Way") were uncorrelated between Movement and No-

Movement trials (*Pearson's r* = 0.17, *p* = 0.48, **Fig. 2g**). Together, these observations suggest that the

same learning process may underlie adaptive single-trial learning events in response to errors on both kinds

of trials, while maladaptive changes in hand-angle may be attributable to noise.

Implicit motor adaptation proceeds after simulated errors in an online visuomotor task

Illustrating that the above observations are reproducible and generalize across experimental

contexts, we again observed that simulated errors in No-Movement trials also induced motor adaptation in

an online, crowd-sourced version of the task. Participants (*n* = 40) made hand movements using their

computer mouse or trackpad to move a cursor towards a target. As in the experiment described above,

trials were presented in triplets, allowing us to measure STL that occurred in response to cursor feedback

presented during Movement and No-Movement trials at the center of each triplet (**Fig. 1b**-**d**). For this online

study, triplets with 0° perturbations/simulated errors were also included to provide an estimate of baseline

changes in hand angle, in the event that participants exhibited strong movement biases in the online

platform.

Participants showed STL that was directionally appropriate for the perturbation applied during both

Movement and No-Movement trials (**Supplemental Fig. 2a**-**c**, please refer to the supplemental material

details of the statistical analysis). Further echoing the results of the in-person study, STL on both Movement

and No-Movement trials was retained beyond the triplet in which the relevant error occurred (**Supplemental**

Fig. 2d-e), and the magnitude of STL in the direction that would counter the perturbation was again
correlated across the two movement conditions (**Supplemental Fig. 2e**). These data provide further
support for the claim that movements that are not performed can undergo implicit motor adaptation and
extend our findings to a task with different movement demands.

Supplemental Figure 2. Single-trial learning in response to errors on Movement and No-Movement
trials during an online visuomotor adaptation task. (**a**) An example participant's mean ± SEM changes
in reach paths across triplets (green: triplets with perturbations on Movement trials, magenta: triplets with
perturbations on No-Movement trials, solid lines: perturbation was a CW rotation, dashed lines: perturbation
was a CCW rotation). (**b**) Boxplot showing STL across Movement (green) and No-Movement (magenta)
triplets for participants in an online version of the task described in Figure 1 (n = 40). (**c**) Estimated marginal
means (EMMs) ± 95% confidence intervals from the linear mixed model (LMM) fit to participants' STL
performance (summarized in **b**). The LMM (fixed effects: rotation [15° counterclockwise {CCW}, 0°, and 15°
clockwise {CW}], movement condition [Movement, No-Movement], rotation x movement condition
interaction; random effects: participant) revealed significant main effects of rotated cursor feedback (F(2,
2223) = 136.46, p = 2.2 x 10^{-16}, *partial R^2* = 0.11) and movement condition (F(1, 2248) = 4.74, p = 0.03,
partial R^2 = 0.002), as well as a significant interaction (F(2, 2229) = 12.40, p = 4.41 x 10^{-6}, *partial R^2* = 0.01).
Post-hoc pairwise comparisons of the EMMs from the model support the claim that rotated feedback
induced a statistically significant degree of STL on both Movement (0° vs 15° CW: t(2227) = 9.14, p_{adj} =
6.39 x 10^{-19}, *Cohen's d* = 0.61; 0° vs 15° CCW: t(2220) = 7.81, p_{adj} = 2.61 x 10^{-14}, *Cohen's d* = 0.52) and
No-Movement trials (0° vs 15° CW: t(2225) = 3.92, p_{adj} = 1.39 x 10^{-4}, *Cohen's d* = 0.31; 0° vs 15° CCW:
t(2229) = 3.56, p_{adj} = 4.84 x 10^{-4}, *Cohen's d* = 0.29). Adaptation in the presence of a rotation was significantly
greater in Movement trials than No-Movement trials for CW (t(2238) = 4.98, p_{adj} = 1.26 x 10^{-6}, *Cohen's d* =
0.37) and CCW rotations (t(2239) = 2.06, p_{adj} = 0.04, *Cohen's d* = 0.15). (**d**) Group mean ± SEM change in

(Δ) hand angle after exposure to Movement (green) and No-Movement (magenta) triplets' perturbations.
Positive Δ values indicate that the change in hand angle proceeded opposite the direction of the
perturbation (*i.e.,* in the direction that would counter the error). (**e**) Group mean ± SEM ratio of remembered
STL to STL. Remembered STL was statistically significantly greater than 0 for both Movement (one-sample
signed-rank test: $V = 819$, $p_{adj} = 1.09 \times 10^{-11}$, $r = 0.87$) and No-Movement triplets ($V = 769$, $p_{adj} = 9.69 \times 10^{-8}$, $r = 0.76$), but remembered STL did not significantly differ between movement conditions (paired-samples
signed-rank test: $V = 441$, $p_{adj} = 0.68$). (**f**) Scatter plot showing the relationship between individual subjects'
signed-rank test: $V = 441$, $p_{adj} = 0.68$). (**f**) Scatter plot showing the relationship between individual subjects'
STL amplitude in the direction opposite the rotation on Movement and No-Movement trials. When we
considered instances of STL in the direction that would compensate for the observed error (update opposite
rotation, "Right-Way"), within-subject changes in hand angle were correlated between Movement and No-
Movement trials (*Pearson's r* = 0.49, $p_{adj} = 0.002$). (**g**) As in **f**, but showing data from trials with changes in
hand angle in the direction that would exacerbate the observed error (update in direction of rotation,
"Wrong-Way"). These ΔHand Angle values were uncorrelated between Movement and No-Movement trials
(*Pearson's r* = 0.10, $p_{adj} = 0.52$). These observations support the idea that the same learning process may
underlie adaptive single-trial learning events in response to errors on both kinds of trials, while maladaptive
changes in hand-angle may be attributable to potential sources of random noise. Boxplot centers: median,
notch: 95% confidence interval of the median, box edges: 1st and 3rd quartiles, whiskers: most extreme
value within 1.5*interquartile range of the median. Statistical significance (* = $p < 0.05$; n.s. = $p \geq 0.05$) is.
Abbreviations: STL – single-trial learning, Δ – change, CW – clockwise, CCW – counterclockwise.

Motor adaptation during No-Movement triplets does not depend on participants' control over cursor

trajectory during Movement trials

We note that rotated visual feedback on Movement trials was sensitive to people's actual reaching

directions. That is, the rotation was simply added to the measured reach direction, as is typical in visuomotor

rotation tasks. It is possible that these directional contingencies affected participants' responses to error,

potentially encouraging them to attempt to deliberately control the cursor's position via an explicit re-aiming

process.[18] To rule this out, we recruited a new group of participants ($n = 37$) to perform a variant of the task

where the visual cursor moved in a fixed path ("error-clamped" feedback[17]; **Supplemental Fig. 3**) in one of

three directions (0° or 15° CW/CCW) on trials with feedback.

Supplemental Figure 3. Schematics illustrating the difference between rotation and error-clamp
perturbations. Diagrams showing the relationship between hand and cursor feedback movement
directions under rotational error regimes (left) and error-clamp regimes (right). When rotational

perturbations are applied to cursor feedback, the cursor's movement direction is contingent upon the
participant's movement direction, although there is an experimenter-induced offset. If a rotation is
consistently applied, participants explicit aiming strategies are a viable supplement for imperfect implicit
motor performance. When error-clamp perturbations are applied to cursor feedback, the cursor travels in a
fixed direction, regardless of the direction that the hand travels, rendering explicit aiming strategies useless.
Thus, error-clamp perturbations are often used in studies attempting to isolate implicit motor adaptation
processes.

Replicating and extending the findings from the two studies employing rotated cursor feedback,

participants assigned to the error-clamp condition exhibited STL after Movement and No-Movement trials

(**Supplemental Fig. 4a-c**; please refer to supplemental material for details). We also observed significant

retention of STL (**Supplemental Fig. 4d-e**) and correlation between STL amplitude on Movement and No-

Movement trials where adaptation proceeded opposite the direction of the perturbation (**Supplemental Fig.**

4f-g). These data further strengthen the claim that motor adaptation does not require movement.

Supplemental Figure 4. Single-trial learning in response to errors on Movement and No-Movement
trials with error-clamped feedback or simulated errors. (**a**) An example participant's mean ± SEM
changes in reach paths across triplets (green: triplets with perturbations on Movement trials, magenta:
triplets with perturbations on No-Movement trials, solid lines: perturbation was a CW error-clamp, dashed
lines: perturbation was a CCW error-clamp). (**b**) Boxplot showing STL across Movement (green) and No-
Movement (magenta) triplets for participants (*n* = 37) in an online experiment where cursor feedback was
error-clamped on Movement trials. (**c**) Estimated marginal means (EMMs) ± 95% confidence intervals from

the linear mixed model (LMM) fit to participants' STL performance (summarized in **b**). The LMM (fixed
effects: rotation [15° counterclockwise {CCW}, 0°, and 15° clockwise {CW}], movement condition
[Movement, No-Movement], error-clamp x movement condition interaction; random effects: participant)
revealed significant main effects of error-clamped cursor feedback (F(2, 1829) = 79.46, p = 2.2 x 10^{-16},
partial R^2 = 0.08) and an interaction between error-clamp and movement condition (F(2, 1832) = 8.45, p =
0.0002, *partial R^2* = 0.0003), although there was no main effect of movement condition (F(1, 1844) = 0.60,
p = 0.44). Post-hoc comparisons of the EMMs from the model revealed significant STL in response to non-
zero error-clamped feedback on both Movement (0° vs 15° CW: t(1827) = 7.55, p_{adj} = 3.08 x 10^{-13}, *Cohen's*
d = 0.56; 0° vs 15° CCW: t(1828) = 5.57, p_{adj} = 8.84 x 10^{-8}, *Cohen's d* = 0.41) and No-Movement trials (0°
vs 15° CW: t(1830) = 3.21, p_{adj} = 0.002, *Cohen's d* = 0.29; 0° vs 15° CCW: t(1832) = 2.25, p_{adj} = 0.03,
Cohen's d = 0.22). Adaptation in the presence of a 15° error-clamp was significantly greater on Movement
trials than No-Movement trials for CW (t(1846) = 3.49, p_{adj} = 0.0009, *Cohen's d* = 0.29) and CCW clamps
(t(1846) = 2.29, p_{adj} = 0.03, *Cohen's d* = 0.19). Please refer to Supplementary Table 2 for further details on
post-hoc comparisons in this panel. (**d**) Group mean ± SEM change in (Δ) hand angle one and two trials
after exposure to Movement (green) and No-Movement (magenta) triplets' perturbations. Positive Δ values
indicate that the change in hand angle proceeded opposite the direction of the perturbation (*i.e.,* in the
direction that would counter the error). (**e**) Remembered STL shown as the ratio of relative hand angle 2
trials after experiencing a perturbation to the relative hand angle 1 trial after the perturbation (STL).
Remembered STL was significantly greater than 0 after both Movement (green; one-sample t-test: t(36) =
11.31, p_{adj} = 6.23 x 10^{-13}, *Cohen's d* = 1.86) and No-Movement triplets (magenta, one-sample signed-rank
test: V = 579, p_{adj} = 5.95 x 10^{-5}, r = 0.64), but did not exhibit statistically significant differences between
movement conditions (paired t-test: t(35) = 1.71, p_{adj} = 0.09). Remembered STL on No-Movement trials
could not be computed for one participant, so n = 36 instead of 37 in this panel. (**f**) Scatter plot showing the
relationship between individual subjects' STL amplitude in the direction opposite the error-clamp on
Movement and No-Movement trials (i.e., the "Right-Way"). Right-way changes in hand angle were
correlated between Movement and No-Movement trials (*Pearson's r* = 0.40, p_{adj} = 0.03). (**g**) As in (**f**), but
showing data from trials on which STL proceeded in the same direction as the error-clamp (*i.e.*, the "Wrong-
Way"). Wrong-Way changes in hand angle were not statistically significantly correlated between Movement
and No-Movement trials (r = 0.33, p_{adj} = 0.06). These observations support the idea that the same learning
process may underlie adaptive single-trial learning events in response to errors on both kinds of trials, while
maladaptive changes in hand-angle may be attributable to potential sources of random noise. Boxplot
centers: median, notch: 95% confidence interval of the median, box edges: 1st and 3rd quartiles, whiskers:
most extreme value within 1.5*interquartile range of the median. Statistical significance (* = p < 0.05; n.s.
= p ≥ 0.05) is indicated for selected comparisons. Abbreviations: STL – single-trial learning, Δ – change,
CW – clockwise, CCW – counterclockwise.

Adaptation on No-Movement trials does not depend on within-session adaptation on Movement trials

In two further experiments, we asked if adaptation to errors in the No-Movement condition was

contingent on sharing a context with the Movement condition. In other words, if learning in the No-

Movement condition only occurs when there are neighboring trials in the Movement condition producing

typical SPEs, it is possible that adaptive responses observed in the No-Movement condition reflect a

"cueing" effect, whereby an adapted sensorimotor map is cued by observation of the visual error and then

retrieved on the subsequent trial(s). While our previous retention (**Fig. 2d**-**e**, **Supplemental Fig. 2d-e**,

Supplemental Fig. 4d-**e**) and correlation (**Fig. 2f-g**, **Supplemental Fig. 2f-g**, **Supplemental Fig. 4f**-**g**)

results argue against this interpretation as they suggest a shared learning mechanism across movement

conditions, we opted to directly test this alternative explanation in another pair of experiments. Here, we

only included 0° rotated (**Fig. 3a**, left, n = 24 participants) or clamped (**Fig. 3a**, right, n = 37 participants)

error feedback on Movement trials, but maintained 0° or 15° CW/CCW errors on the No-Movement trials.

The key results were again replicated – learning was preserved in the No-Movement condition even when

error feedback had never been associated with executed movements (**Fig. 3c**, rotation: LMM: $F(557)$ 23.01,

p = 2.07 x 10^{-6}, *partial R^2* = 0.04; error-clamp: $F(802)$ = 9.41, p = 9.14 x 10^{-5}, *partial R^2* = 0.02). Post-hoc

pairwise comparisons showed that adaptation was significantly different between triplets with clockwise and

counterclockwise errors for both the rotation ($t(557)$ = 4.80, p = 2.07 x 10^{-6}, *Cohen's d* = 0.4) and error-

clamp experiments ($t(1453)$ = 4.32, p = 5.34 x 10^{-5}, *Cohen's d* = 0.37) – a hallmark of implicit motor

adaptation (please refer to **Supplemental Table 3** for all post-hoc test results). Overall levels of STL

observed on No-Movement trials were comparable during these two experiments to those discussed above,

and within the range of learning rates previously observed in the literature (**Supplemental Fig. 1**).

Furthermore, both groups of participants showed retention of STL that differed significantly from 0 (rotation,

mean ± SEM: 0.53 ± 0.06 retention ratio, one-sample t-test: $t(22)$ = 8.28, p = 3.34 x 10^{-8}, *Cohen's d* = 1.73;

error-clamp, median: 0.45, interquartile-range: 0.58, one-sample signed-rank test: V = 507, p = 0.001, r =

0.53). Overall, these experiments support our hypothesis that motor adaptation can proceed without

movement execution.

Figure 3. Effects of simulated errors when perturbations were never applied during Movement trials.
(**a**) Schematic illustrating the relationship between movement and visual feedback on Movement trials
during an experiment where visuomotor rotations (left) or error-clamps (right) were never applied during
Movement trials. (**b**) An example participant's mean ± SEM changes in reach paths across No-Movement
triplets from studies in which non-zero rotations (left) and error-clamps (right) were never applied (solid
lines: perturbation was CW, dashed lines: perturbation was CCW). (**c**) Boxplots showing STL in response
to different directions of simulated errors (No-Movement triplets indicated in magenta) from rotation (left, *n*
= 24) and error-clamp (right, *n* = 37) studies. (**d**) Estimated marginal means ± 95% confidence intervals
from the linear mixed models fit to each participant's STL performance summarized in (**c**). Asterisks indicate
statistically significant differences. (**e**) Mean ± SEM relative hand angles on the two trials after a perturbation
was presented on a No-Movement trial. Please refer to Supplemental Table 3 for detailed statistical results.
Boxplot centers: median, notches: 95% confidence interval of the median, box edges: 1st and 3rd quartiles,
whiskers: most extreme values within 1.5*IQR of the median. Statistical significance (* = p_{adj} < 0.05; n.s. =
p_{adj} ≥ 0.05) is indicated for selected comparisons. Abbreviations: STL – single-trial learning, CW –
clockwise, CCW – counterclockwise, Δ – change in.

DISCUSSION

The results presented above demonstrate that movements can be implicitly refined even when they

are not performed. Participants who were cued to perform a movement towards a target but suppressed

that movement after observation of a Stop cue showed consistent, robust STL in response to simulated

errors (**Figs. 2-3**, **Supplemental Fig. 2**, **Supplemental Fig. 4**). As implicit learning necessarily proceeds

following SPEs, our data also provide evidence that SPEs are computed even when movements are not

performed. These findings strongly support the fundamental assumptions of predictive processing

frameworks of motor adaptation, where precise sensory predictions are generated from a movement intent

(or "plan", "goal") and compared against sensory observations to induce error-based learning.[12,25]

We argue that we have measured learning via an implicit process, and, by extension, that the STL

observed in our study provides evidence that SPEs are computed regardless of whether a movement is

performed. Although it is important to note that visuomotor learning tasks do sometimes recruit cognitive

learning strategies (e.g., the deliberate "re-aiming" of movements), multiple factors indicate that our studies

successfully measured implicit adaptation.[18,28,29] First, in line with previous work isolating the implicit

adaptation process, participants were instructed to ignore the visually displayed cursor and try to contact

the target on every trial, a straightforward technique which has been consistently shown to eliminate the

explicit re-aiming of movements.[15,17,22,23,26] Second, the randomization of both the presence and direction

of visual errors also discourages explicit learning, reducing motivation to apply ineffective re-aiming

strategies (see [30]). Third, we excluded data from participants who did not understand or recall the instruction

to always aim directly at the target (see *Methods*), decreasing the likelihood that strategic re-aiming

contaminated the analysis. Fourth, adaptation persisted into subsequent no-feedback trials (**Fig. 2d-e**, **Fig.**

3e, **Supplemental Fig. 2d-e**, **Supplemental Fig. 4d-e**), a finding that is consistent with lingering implicit

motor learning; it is unlikely that strategies would be maintained through trials where no feedback is present.

Fifth, the magnitude of STL observed was consistent with multiple previous studies that measured implicit

motor adaptation rates (**Supplemental Fig. 1**).[17,23,31] Lastly, the adaptation effects observed in the No-

Movement conditions were not attributable to recall of learning that occurred on Movement trials (**Fig. 3**.

Our data thus provide striking evidence that movement is not required for implicit adaptation, and, by

extension, SPE computation.

While motor planning and concurrent sensory observations are sufficient to drive SPE computation

and motor adaptation, our data also indicate that errors paired with movements support a greater degree

of learning than errors without preceding movements, as participants showed significantly stronger STL

over triplets with Movement trials versus No-Movement trials. This suggests that movement provides

additional training input beyond what erroneous visual feedback provides alone. Alternately (though not

mutually exclusive), it may be that the absence of movement-related proprioceptive feedback dampens the

SPE signals that drive adaptation, perhaps by failing to counterbalance signals predicting proprioceptive

feedback.[8,32] Simpler explanations are that participants simply did not recognize the Go cue to set a

movement goal and plan a movement on a subset of the No-Movement trials, and/or that the timing of the

simulated error was inherently too variable relative to the behaviorally unobservable internal prediction.

Further experimentation will be required to address these possibilities. Nonetheless, our data demonstrate

the significant influence of the brain's prediction signals on learning – even without the ability to directly

attribute sensory feedback to an actual movement, prediction of a planned movement's sensory

consequences supports the error computations that drive adaptation of future behavior.

While our results demonstrate that overt movement is not necessary for motor adaptation, we are

not able to discern whether the generation of a descending motor command is a necessary component for

updating the forward model. To reliably elicit prediction generation from the forward model, we employed a

motor inhibition paradigm, briefly cueing participants to reach before presenting them with an abrupt stop

signal. This strategy is likely to have recruited mechanisms of motor inhibition beyond the primary motor

cortex, leaving open the possibility that a descending motor command and not an upstream motor plan *per*

se provide the necessary input for forward model adaptation.[33–35] Future experimentation quantifying neural

(EEG) or physiological (EMG) signals during this task and measuring their relationship with adaptation on

No-Movement trials will be crucial for addressing this point. Inspired by work in non-human primates, we

speculate that similar motor cortical preparatory states in the Movement and No-Movement conditions may

produce comparable inputs to downstream internal models.[36]

The current report provides evidence that implicit motor adaptation can proceed without movement,

adding to a substantial body of work indicating that many forms of motor learning do not strictly require

movement-based practice. For instance, in [35], after human participants observed others adapting to a force

field applied during reaching movements, the observers were able to partially compensate for that same

force field when they encountered it. Interestingly, this observational learning did not proceed if participants

were executing other movements during the observation period. This finding combined with subsequent

neuroimaging data showing that observational learning recruits brain areas associated with motor planning

were taken to suggest that engagement in a covert motor planning process may be the critical element

allowing for force-field adaptation following error observation without movement.[37–39] Together, this related

prior work and the evidence we have provided that motor planning is sufficient for the generation of sensory

predictions and can support SPE computations suggest that there are multiple routes to inducing motor

planning and ultimately driving motor adaptation processes.

Other reports in the motor learning literature have provided evidence for cognitive compensation

for observed motor errors during reaching, improved visual tracking following observation of target

movement without engagement in visual pursuit, and improvement in movement speeds as a result of

mental imagery training, highlighting the breadth of motor performance-related processes that can be

trained without engagement in physical movements.[19,20,40,41] Together with the findings related to implicit

motor adaptation discussed above, this work suggests that many features of motor performance can be

improved by training regiments that do not involve movement. This work points to a potential opportunity

for the development of motor training or rehabilitation protocols that can be used when people are unable

to physically perform target motor behaviors, perhaps improving performance beyond what physical

practice can do alone.

Finally, our results echo the fact that other types of learning also appear to occur without overt task

execution. Outside of motor learning, fear associations can be extinguished by instructing participants to

imagine a fear-predicting stimulus even when they are not presented with the stimulus, with this

"imagination" protocol generating neural signatures of the negative prediction errors observed during

naturalistic fear extinction.[42,43] Considering this prior work and the findings presented in this study, it may

be that the generation of predictions for comparison with subsequent sensory observations is sufficient for

error-based learning across motor and non-motor domains alike. In other words, task execution may not

always be required for learning, so long as the predictions and observations needed to compute errors are

both present.

REFERENCES

1. Friston, K. The free-energy principle: a unified brain theory? *Nat Rev Neurosci* **11**, 127–138 (2010). doi: 10.1038/nrn2787

2. Jordan, M. I. & Rumelhart, D. E. Forward models: Supervised learning with a distal teacher. *Cognitive Science* **16**, 307–354 (1992). doi: 10.1016/0364-0213(92)90036-T

3. Wallman, J. & Fuchs, A. F. Saccadic gain modification: Visual error drives motor adaptation. *Journal of Neurophysiology* **80**, 2405–2416 (1998). doi: 10.1152/jn.1998.80.5.2405

4. Shadmehr, R., Smith, M. A. & Krakauer, J. W. Error correction, sensory prediction, and adaptation in motor control. *Annu. Rev. Neurosci.* **33**, 89–108 (2010). doi:10.1146/annurev-neuro-060909-153135

5. Tseng, Y., Diedrichsen, J., Krakauer, J. W., Shadmehr, R. & Bastian, A. J. Sensory prediction errors drive cerebellum-dependent adaptation of reaching. *Journal of Neurophysiology* **98**, 54–62 (2007). doi: 10.1152/jn.00266.2007

6. Wong, A. L. & Shelhamer, M. Sensorimotor adaptation error signals are derived from realistic predictions of movement outcomes. *Journal of Neurophysiology* **105**, 1130-1140 (2011). doi: 10.1152/jn.00394.2010

7. Held, R. & Freedman, S. J. Plasticity in human sensorimotor control. *Science* **142**, 455–462 (1963).

8. Miall, R. C. & Wolpert, D. M. Forward Models for Physiological Motor Control. *Neural Networks* **9**, 1265–1279 (1996). doi: 10.1016/S0893-6080(96)00035-4

9. Kilteni, K., Andersson, B. J., Houborg, C. & Ehrsson, H. H. Motor imagery involves predicting the sensory consequences of the imagined movement. *Nat Commun* **9**, 1617 (2018). doi: 10.1038/s41467-018-03989-0

10. Voss, M., Ingram, J. N., Haggard, P. & Wolpert, D. M. Sensorimotor attenuation by central motor command signals in the absence of movement. *Nat Neurosci* **9**, 26–27 (2006). doi: 10.1038/nn1592

11. Sheahan, H. R., Franklin, D. W. & Wolpert, D. M. Motor Planning, Not Execution, Separates Motor Memories. *Neuron* **92**, 773–779 (2016). doi: 10.1016/j.neuron.2016.10.017

12. Sheahan, H. R., Ingram, J. N., Žalalytė, G. M. & Wolpert, D. M. Imagery of movements immediately following performance allows learning of motor skills that interfere. *Sci Rep* **8**, 14330 (2018). doi: 10.1038/s41598-018-32606-9

13. Butcher, P. A. & Taylor, J. A. Decomposition of a sensory prediction error signal for visuomotor adaptation. *Journal of Experimental Psychology: Human Perception and Performance* **44**, 176–194 (2018). doi: 10.1037/xhp0000440

14. Tsay, J. S., Haith, A. M., Ivry, R. B. & Kim, H. E. *Interactions between sensory prediction error and task error during implicit motor learning*. 2021.06.20.449180 https://www.biorxiv.org/content/10.1101/2021.06.20.449180v2 (2021) doi:10.1101/2021.06.20.449180.

15. McDougle, S. D., Ivry, R. B. & Taylor, J. A. Taking aim at the cognitive side of learning in sensorimotor adaptation tasks. *Trends in Cognitive Sciences* **20**, 535–544 (2016). doi: 10.1016/j.tics.2016.05.002

16. Mazzoni, P. & Krakauer, J. W. An implicit plan overrides an explicit strategy during visuomotor adaptation. *J. Neurosci.* **26**, 3642–3645 (2006). doi: 10.1523/JNEUROSCI.5317-05.2006

17. Morehead, J. R., Taylor, J. A., Parvin, D. E. & Ivry, R. B. Characteristics of implicit sensorimotor adaptation revealed by task-irrelevant clamped feedback. *J Cogn Neurosci* **29**, 1061–1074 (2017). doi: 10.1162/jocn_a_01108

18. Taylor, J. A., Krakauer, J. W. & Ivry, R. B. Explicit and implicit contributions to learning in a sensorimotor adaptation task. *The Journal of Neuroscience* **34**, (2014).

19. Ong, N. T. & Hodges, N. J. Absence of after-effects for observers after watching a visuomotor adaptation. *Exp Brain Res* **205**, 325–334 (2010). doi: 10.1007/s00221-010-2366-4

20. Lim, S. B., Larssen, B. C. & Hodges, N. J. Manipulating visual–motor experience to probe for
observation-induced after-effects in adaptation learning. *Exp Brain Res* **232**, 789–802 (2014). doi:
10.1007/s00221-013-3788-6

21. Taylor, J. A. & Ivry, R. B. Flexible cognitive strategies during motor learning. *PLOS Computational*
Biology **7**, e1001096 (2011). doi: 10.1371/journal.pcbi.1001096

22. Tsay, J. S., Parvin, D. E. & Ivry, R. B. Continuous reports of sensed hand position during
sensorimotor adaptation. *Journal of Neurophysiology* **124**, 1122–1130 (2020). doi:
10.1152/jn.00242.2020

23. Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R. & Ivry, R. B. Invariant errors reveal
limitations in motor correction rather than constraints on error sensitivity. *Communications Biology* **1**,
1–7 (2018). doi: 10.1038/s42003-018-0021-y

24. Avraham, G., Morehead, J. R., Kim, H. E. & Ivry, R. B. Reexposure to a sensorimotor perturbation
produces opposite effects on explicit and implicit learning processes. *PLOS Biology* **19**, e3001147
(2021). doi: 10.1371/journal.pbio.3001147

25. Poh, E., Al-Fawakari, N., Tam, R., Taylor, J. A. & McDougle, S. D. Generalization of motor learning in
psychological space. *bioRxiv* 2021.02.09.430542 (2021) doi:10.1101/2021.02.09.430542. doi:
10.1101/2021.02.09.430542

26. Hadjiosif, A. M., Krakauer, J. W. & Haith, A. M. Did we get sensorimotor adaptation wrong? Implicit
adaptation as direct policy updating rather than forward-model-based learning. *J. Neurosci.* **41**,
2747–2761 (2021). doi: 10.1523/JNEUROSCI.2125-20.2021

27. Wei, K. & Körding, K. Relevance of error: What drives motor adaptation? *Journal of Neurophysiology*
101, 655–664 (2009). doi: 10.1152/jn.90545.2008

28. Krakauer, J., W., Hadjiosif, A., Xu, J. & Wong, A., L. Motor Learning. in *Comprehensive Physiology*
(ed. Terjung, R.) (Wiley, 2019). doi:10.1002/cphy.

29. McDougle, S. D. & Taylor, J. A. Dissociable cognitive strategies for sensorimotor learning. *Nat*
Commun **10**, 40 (2019). doi: 10.1038/s41467-018-07941-0

30. Hutter, S. A. & Taylor, J. A. Relative sensitivity of explicit reaiming and implicit motor adaptation.
Journal of Neurophysiology **120**, 2640–2648 (2018). doi: 10.1152/jn.00283.2018

31. Wei, K. & Körding, K. Uncertainty of feedback and state estimation determines the speed of motor
adaptation. *Front. Comput. Neurosci.* **4**, (2010). doi: 10.3389/fncom.2010.00011

32. Requarth, T., Kaifosh, P. & Sawtell, N. B. A role for mixed corollary discharge and proprioceptive
signals in predicting the sensory consequences of movements. *J. Neurosci.* **34**, 16103–16116 (2014).
doi: 10.1523/JNEUROSCI.2751-14.2014

33. Xu, K. Z., Abderson, B. A., Emeric, E .E., Sali, A. W., Stuphorn, V., Yantis, S., Courtney, S. M. Neural
basis of cognitive control over movement inhibition: Human fMRI and primate electrophysiology
evidence. *Neuron* **96**, 1447-1458.e6 (2017). doi: 10.1016/j.neuron.2017.11.010

34. Osada, T., Ohta, S., Ogawa, A., Tanaka, M., Suda, A., Kamagata, K., Hori, M., Shigeki, A., Shimo,
Y., Hattori, N., Shimizu, T., Enomoto, H., Hanajima, R., Ugawa, Y., Konishi, S. An essential role of
the intraparietal sulcus in response inhibition predicted by parcellation-based network. *J. Neurosci.*
39, 2509–2521 (2019). doi: 10.1523/JNEUROSCI.2244-18.2019

35. Duque, J., Greenhouse, I., Labruna, L. & Ivry, R. B. Physiological Markers of Motor Inhibition during
Human Behavior. *Trends in Neurosciences* **40**, 219–236 (2017). doi: 10.1016/j.tins.2017.02.006

36. Vyas, S, Even-Chen, N., Stavitsky, S. D., Ryu, S. I., Nuyujukian, P., Shenoy, K. V. Neural population
dynamics underlying motor learning transfer. *Neuron* **97**, 1177-1186.e3 (2018). doi:
10.1016/j.neuron.2018.01.040

37. Mattar, A. A. G. & Gribble, P. L. Motor Learning by Observing. *Neuron* **46**, 153–160 (2005). doi:
10.1016/j.neuron.2005.02.009

38. Malfait, N., Valyear, K. F., Culham, J. C., Anton, J., Brown, L. E., Gribble, P. L. fMRI activation during
observation of others' reach errors. *Journal of Cognitive Neuroscience* **22**, 1493–1503 (2010). doi:
10.1162/jocn.2009.21281
39. McGregor, H. R. & Gribble, P. L. Functional connectivity between somatosensory and motor brain
areas predicts individual differences in motor learning by observing. *Journal of Neurophysiology* **118**,
1235–1243 (2017). doi: 10.1016/j.cub.2016.01.064
40. Gentili, R., Han, C. E., Schweighofer, N. & Papaxanthis, C. Motor Learning Without Doing: Trial-by-
Trial Improvement in Motor Performance During Mental Training. *Journal of Neurophysiology* **104**,
(2010).
41. Cain, M., Botschko, Y. & Joshua, M. Passive motor learning: Oculomotor adaptation in the absence
of behavioral errors. *eNeuro* **8**, (2021). doi: 10.1523/ENEURO.0232-20.2020
42. Mendelsohn, A., Pine, A. & Schiller, D. Between thoughts and actions: Motivationally salient cues
invigorate mental action in the human brain. *Neuron* **81**, 207–217 (2014). doi:
10.1016/j.neuron.2013.10.019
43. Reddan, M. C., Wager, T. D. & Schiller, D. Attenuating neural threat expression with imagination.
Neuron **100**, 994-1005.e4 (2018). doi: 10.1016/j.neuron.2018.10.047
44. Herzfeld, D. J., Vaswani, P. A., Marko, M. K. & Shadmehr, R. A memory of errors in sensorimotor
learning. *Science* **345**, 1349–1353 (2014). doi: 10.1126/science.1253138
45. van Beers, R. J. Motor learning Is optimally tuned to the properties of motor noise. *Neuron* **63**, 406–
417 (2009). doi: 10.1016/j.neuron.2009.06.025
46. Wu, H. G., Miyamoto, Y. R., Castro, L. N. G., Ölveczky, B. P. & Smith, M. A. Temporal structure of
motor variability is dynamically regulated and predicts motor learning ability. *Nat Neurosci* **17**, 312–
321 (2014). doi: 10.1038/nn.3616
47. Photon Storm. Phaser (V. 3.24.1). https://phaser.io (2020).
48. Tsay, J. S., Ivry, R. B., Lee, A. & Avraham, G. Moving outside the lab: The viability of conducting
sensorimotor learning studies online. *NBDT* 1–22 (2021) doi:10.51628/001c.26985.
49. Kassambara, A. *rstatix: Pipe-Friendly Framework for Basic Statistical Tests*. (2021).
50. Hothorn, T., Hornik, K., van de Wiel, M. A. & Zeileis, A. Implementing a class of permutation tests:
The coin package. *Journal of Statistical Software* **28**, 1–23 (2008). doi: 10.18637/jss.v028.i08
51. Bartoń, K. *MuMIn: Multi-Model Inference*. (2020).
52. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest package: Tests in linear mixed
effects models. *Journal of Statistical Software* **82**, 1–26 (2017). doi: 10.18637/jss.v082.i13
53. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using lme4.
arXiv:1406.5823 [stat] (2014).
54. Jaeger, B. *r2glmm: Computes R Squared for Mixed (Multilevel) Models*. (2017).
55. Lenth, R. V. *emmeans: Estimated Marginal Means, aka Least-Squares Means*. (2021).
56. Torchiano, M. *effsize: Efficient Effect Size Computation*. (2020).
57. Ben-Shachar, M., Lüdecke, D. & Makowski, D. effectsize: Estimation of effect size indices and
standardized parameters. *Journal of Open Source Software* **5**, 7 (2020). doi: 10.21105/joss.02815
58. Bache, S. M., Wickham, H. & RStudio. *magrittr: A Forward-Pipe Operator for R*. (2020).
59. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, 2016).
60. Kassambara, A. *ggpubr: 'ggplot2' Based Publication Ready Plots*. (2020).
61. Lüdecke, D. ggeffects: Tidy Data Frames of Marginal Effects from Regression Models. *JOSS* **3**, 772
(2018). doi: 10.21105/joss.00772

METHODS

Participants. Participants (n = 233, aged 18-35, 126 female) recruited from the research participation pools

at Princeton/Yale University and on Prolific provided informed consent, approved by each University's IRB.

Seventy-five participants were excluded (10 from the dataset collected for Supplemental Fig. 2, 13 from the

dataset collected for Supplemental Fig. 4, 26 from the dataset collected for the rotation perturbation

experiments described in Fig. 3, and 26 from the dataset collected for the error-clamp perturbation

experiments described in Fig. 3) for failure to sufficiently recall task instructions, as ascertained by a

questionnaire at the end of the experiment, leaving 158 participants for our analyses. See the *Questionnaire*

section below the *Test phase* sections for more details. We note that all the key results described here (*i.e.,*

statistically significant learning after No-Movement trials) held with or without these exclusions; we opted to

be conservative in our exclusion criteria to limit potential effects of explicit learning. We note that the

samples used for the online experiments described in the text are around twice the size of similar studies

in the literature, providing additional statistical power to compensate for the experiment being conducted

remotely.[44–46]

Task Setup: In Lab. Participants were seated in a chair and made ballistic reaching movements while

grasping the handle of a robotic manipulandum with their dominant hand (Kinarm End-Point). The

manipulandum restricted movements to the horizontal plane. All visual stimuli were projected to the

participant via a horizontal display screen (60 Hz) reflected onto a semi-silvered mirror mounted above the

robotic handle. The mirror occluded vision of the arm, hand, and robotic handle, preventing direct visual

feedback of hand position. Tasks were programmed in Matlab 2019a's Simulink for deployment in Kinarm's

Dexterit-E software (version 3.9). Movement kinematics were recorded at 1 kHz. Each participant viewed

a single target located at either 45°, 135°, 225°, or 315° (with target position counterbalanced across

participants), 8 cm from a central starting location. The target was visible throughout the experiment.

Task Setup: Online. Experiments were conducted remotely using a custom JavaScript web application

based on Phaser 3.24 (download available at [47]), similar to an approach previously described.[48] Each

participant viewed a single target located at either 45°, 135°, 225°, or 315° (with target position

counterbalanced across participants), 250 pixels from a central starting location. The target was visible

throughout the experiment.

Participants used an input device of their choice to control their computer cursor during center-out

movements. One participant reported using a touchscreen device and was excluded from all analyses. The

remaining participants reported using either a trackpad (n = 112), an optical mouse (n = 86), or a trackball

(n = 14). A linear mixed model (LMM) did not show effects of Mouse Type on single-trial learning (STL),

although we observed that participants using a trackpad exhibited longer reaction times than others,

consistent with a previous report.[48]

Mouse position sampling rates depended on the exact hardware that each participant used to

complete the task. Sampling rates were likely affected by features of the specific mouse used, along with

features of the specific computer used, as computers may limit the rate at which the browser samples data

in order to cope with limited processing power. In general, sampling rates were around 60 Hz (median ±

interquartile range across all 213 online participants recruited: 62.46 ± 2.17 Hz) but ranged from 19.23 Hz

to 249.69 Hz. Note that the vast majority of sampling rates were near 60 Hz: Only 5% of sampling rates

were < 41.79 Hz, and only 5% of sampling rates were > 126.65 Hz.

Baseline training phase. For in-lab participants, the robot moved the participant's hand to a central starting

location (depicted by a grey circle) at the middle of the display while hand and cursor feedback were hidden.

They were instructed to hold their hand still in the starting location util the target turned green, at which

point they should make a straight slicing movement through the target. After a 100 ms delay, the robot

moved the hand back to the starting location. Participants completed 5 of these trials with online and

endpoint cursor feedback, followed by 5 trials without visual feedback of the cursor location. Endpoint

feedback was constituted by the cursor remaining at the position where it had passed the target radius for

50 ms. Participants then completed 10 alternating trials on which the target turned green and stayed green

(Execution, 'Go' trials) and on which the target turned magenta 100 ms after turning green, signaling that

participants should withhold their movement (No-Movement, 'Stop' trials). After this baseline phase,

participants were instructed to continue following these instructions for the remainder of the experiment.

Online participants experienced an identical baseline phase, with the exception that they were

instructed to move their mouse into a central starting location on the first trial and subsequently saw their

mouse cursor reappear near the starting location 100 ms after the completion of the reaching movement,

so that participants could quickly return to the start location to initiate the next trial.

Test phase: Rotation and Error-Clamp Experiments (Fig. 2, Supplemental Fig. 2, Supplemental Fig. 4).

During the test phase, 480 (in-lab) or 270 (online) total trials were divided into 3-trial triplets (**Fig. 1c**). The

first and last trials of all triplets were Go trials, and participants received neither online nor endpoint feedback

about cursor location on these trials. The second trial of each triplet was either a Movement or a No-

Movement trial. On Movement trials, participants either received rotated/error-clamped[17] visual feedback

(15° clockwise [-, CW] or counterclockwise [+, CCW], with sign randomized across trials) or veridical/0°

error-clamped visual feedback of their cursor location. On No-Movement (Stop) perturbation trials,

participants viewed a brief animation of the cursor moving straight to the center of the target following a

trajectory deflected by ±15° from the target center. Animation onset latency was set as a running median

of the participant's reaction times on the previous 5 trials. Animation duration was set as a running median

of the participant's movement times on the previous 5 trials. If a participant took longer than 400 ms to

execute a movement, 800 ms to initiate the movement, their reach trajectory changed by >10° during the

movement, or the reach terminated ≥ 60° away from the target, they received a warning and a 4s time-out.

If a participant moved their hand (>5 mm in-lab [radius of the starting location]; anything >0 pixels online)

on a No-Movement trial, the trial was immediately aborted, and they received a warning and a 4s time-out.

The Stop manipulation was successful: Across the experiments, participants erroneously moved on 34.39

± 20.63% (mean ± standard deviation) of Stop trials, suggesting that, for the most part, they were

consistently planning movements on Stop trials.

For in-lab studies, we used 4 possible triplet perturbation trial types (Movement/No-Movement:

±15°), each of which occurred 40 times throughout each session. For online studies, we used 6 possible

triplet perturbation trial types (Movement/No-Movement: ±15° or 0°), each of which occurred 15 times

throughout each session. Triplets were pseudorandomly presented within each block, with the constraints

that a single rotation (±15° or 0°) could not occur on more than 2 consecutive triplets and that the same

movement condition (i.e., Movement or No-Movement) could not occur on more than 3 consecutive triplets.

Three repetitions of each triplet type occurred in blocks of 18 triplets, and participants received a break after

each of these blocks.

Test phase: Rotation and Error-Clamp Experiments with 0° Perturbations on Movement Trials (Fig. 3).

Experiments were conducted as described above for the other online experiments, with the exception of

the details described in this section. For the experiments described in **Fig. 3a-d**, we used a reduced set of

3 possible triplet Perturbation trial types (No-Movement, 15° clockwise error; No-Movement, 15°

counterclockwise error; Movement, 0° rotation). We maintained an equal number of Movement and No-

Movement triplets throughout the session in order to ensure that participants would reliably respond to the

"Go" cue presented at the start of each trial. So, each No-Movement triplet type occurred 22 times, while

the Movement triplet type occurred 44 times. Triplets were pseudorandomly presented within each block,

with the constraints that a single non-zero rotation (15° clockwise, 15° counterclockwise) could not occur

on more than 2 consecutive triplets.

For the experiments described in **Fig. 3e-h**, we used a set of 4 possible triplet Perturbation trial

types (No-Movement, 15° clockwise error; No-Movement, 15° counterclockwise error; No-Movement, 0°

error; Movement, 0° error-clamp). To maintain an equal number of Movement and No-Movement triplets

throughout the session, each No-Movement triplet type occurred 15 times and the Movement triplet type

occurred 45 times.

Questionnaire: As we could not receive verbal confirmation that participants understood the task

instructions in the online version of the task, we asked subjects to fill out a brief questionnaire to query their

understanding of the task. The questionnaire asked participants to attest whether or not 1) their goal was

to move the real mouse and not the cursor straight through the green targets and whether or not 2) their

goal was to move the white cursor (not the real mouse) straight through the green targets. Participants

could select the options, "True," "False," or "Not Sure." Participants were considered to have understood

the instructions if they answered both questions correctly (i.e., answered "True" to question 1 and "False"

to question 2). The majority of participants answered both questions correctly (138 of 213 participants

[65%]), suggesting that most participants understood the task instructions. Nonetheless, these participants

made up the dataset for all reported analyses for the online experiments, and all other online participants

were excluded from analyses to exclude potential effects of explicit re-aiming.

Data analysis. Data were processed in Python 3.8.5 and Matlab 2018a. Trials with movement were

excluded from analysis 1) if any of the reaches in the triplet were not straight (aspect ratio > participant-

wise mean + 3 * participant-wise standard deviation), 2) if the participant received any warning for failure

to follow task instructions (see *Feedback for failure to follow task instructions*, above), or 3) if the triplet

included a No-Movement No-Go perturbation trial with any detectable mouse movement (>0 pixels online,

>5 mm in lab).

Reach endpoint angle was computed as the angular distance between the center of the target and

the point at which the mouse passed the target's radial distance. Because mouse sampling rates did not

always allow us to measure mouse position at the exact target radius during the online study, we used the

last sample before and the first sample after the mouse passed the target radius to compute an interpolated

mouse position at the target radius, as described in a previous report.[48] We note that analyses comparing

these measures to measurements at the last sample of the reach (even when it was beyond the target) or

the hand angle at peak velocity did not result in substantially different hand angle measurements or

statistical outcomes.

Single-trial learning (STL) was measured as the difference between reach endpoint angle on the

third and first trial of each triplet. For our initial analyses, the sign of STL corresponded to the direction of

the relative change in hand angle, with clockwise changes in hand angle taking a negative sign and

counterclockwise changes in hand angle taking a positive sign. When we collapsed STL data across

rotation directions, we normalized the sign of STL so that changes in hand angle opposite the direction of

the imposed rotation took a positive sign and changes in the direction of the rotation took a negative sign.

Remembered STL was quantified as the difference between reach endpoint angle on the first trial

of one triplet and reach endpoint angle on the first trial of the previous triplet. When remembered STL is

reported as a ratio, this value was computed by dividing remembered STL by the STL attributable to a given

triplet.

Statistics. Statistical tests were conducted in R (v. 4.0.3; packages rstatix[49], coin[50], MuMIn[51],

lmerTest[52], lme4[53], r2glmm[54], emmeans[55], effsize[56], effectsize[57], magrittr[58], ggplot2[59], ggpubr[60],

ggeffects[61]). Data from in-lab experiments were analyzed using a two-way repeated measures ANOVA. If

an ANOVA showed a significant main effect or interaction, post-hoc pairwise tests were performed. When

samples failed to satisfy the normality assumption of the pairwise t-test (assessed via a Shapiro-Wilk test),

we used the more robust paired-samples Wilcoxon signed-rank test. Otherwise, we used the more powerful

paired t-test. Effect sizes for ANOVA main effects/interactions were quantified via generalized η^2 (η_G^2), we

quantified the effect sizes for t-tests using Cohen's d, and we used the Wilcoxon effect size (r) to quantify

effect sizes for signed-rank tests. For these and all subsequent analyses, we corrected for multiple

comparisons using the false-discovery rate approach to maintain family-wise alpha at 0.05.

Data from the experiments conducted online did not satisfy multiple assumptions of the two-way

repeated measures ANOVA (non-existence of extreme outliers, sphericity), so we employed a linear mixed

modeling (LMM; R package lmerTest and lme4) approach for analysis of these data. All LMM's included

fixed effects of perturbation size and movement condition, as well as random effects of subject. Degrees of

freedom were estimated using the Kenward-Rogers approach, and LMM outcomes were reported using

ANOVA-style statistics. Partial R^2 was computed to report effect sizes for the LMM factors (R package

r2glmm). Post-hoc pairwise comparisons were performed between estimated marginal means computed

from the LMM (R package emmeans).

For one-off comparisons between samples or to distributions with 0-mean, we checked samples

for normality. When samples were normally distributed, we ran t-tests and computed Cohen's d to report

effect sizes for statistically significant results. Otherwise, we ran Wilcoxon-signed rank tests and measured

effect sizes using the Wilcoxon effect size (r).

FIGURE LEGENDS

Figure 1. Schematics showing the proposed learning framework and task design. (**a**) Schematic

showing how the forward model may support implicit motor adaptation in the presence of sensory feedback

not causally related to self-generated movement. (**b**) Schematic of events on trials with visual feedback.

The robotic apparatus brought the participant's hand to the starting location to initiate a trial. On Movement

trials (top), the target turned green ('GO'), cueing participants to reach through the target. On trials with

visual feedback, participants observed a white feedback cursor move along a rotated trajectory ('Rotation').

On No-Movement trials (bottom), the target turned magenta 100 ms after turning green, cueing participants

to withhold movement ('STOP'). After a delay, an animation showing the feedback cursor moving 15° off-

target played ('Animation'). The hand is shown in the figure for illustrative purposes but was not visible

during the experiment. (**c**) Schematic showing the trials that constituted a triplet and how single-trial learning

(STL) was computed. Triplets were composed of 2 Movement trials without visual feedback flanking either

a Movement or a No-Movement trial with visual feedback. STL was measured as the difference between

reach angles on the flanking trials. (**d**) Schematic showing the pseudorandomized order in which trials were

presented for an example participant, with color reflecting the movement condition (Movement: green, No-

Movement: magenta).

Figure 2. Effects of typical and simulated errors during a visuomotor reach adaptation task. (**a**)

Schematic illustrating how STL and remembered STL measurements were computed. (**b**) An example

subject's (top) and the group's (bottom) mean ± SEM changes in reach paths across triplets with a rotation

applied (green: triplets with perturbations on Movement trials, magenta: triplets with perturbations on No-

Movement trials, solid lines: perturbation was a CW rotation, dashed lines: perturbation was a CCW

rotation). (**c**) STL across Movement (green) and No-Movement (magenta) triplets for all participants ($n =$

20). In the reference frame used here, positive changes in hand angle are CCW. Please refer to

Supplemental Table 1 for details on all statistical tests. (**d**) Group mean ± SEM Δhand angle values after

exposure to Movement (green) and No-Movement (magenta) trial perturbations. Positive Δ values indicate

that the change in hand angle proceeded opposite the direction of the perturbation (*i.e.,* the direction that

would counter the error, "Right-Way"). (**e**) Group mean of participants' ratios of remembered STL to initial

STL during Movement and No-Movement trials. (**f**) The relationship between Right-Way STL observed

during Movement and No-Movement triplets. (**g**) As in (**f**), but for STL observed on trials where adaptation

proceeded in the direction that would exacerbate the error (*i.e.*, the same direction as the perturbation

applied, "Wrong Way"). Statistical significance (* = p_{adj} < 0.05; n.s. = p_{adj} ≥ 0.05) is indicated. Abbreviations:

STL – single-trial learning, CW – clockwise, CCW – counterclockwise, Δ – change in.

Figure 3. Effects of simulated errors when perturbations were never applied during Movement trials.

(**a**) Schematic illustrating the relationship between movement and visual feedback on Movement trials

during an experiment where visuomotor rotations (left) or error-clamps (right) were never applied during

Movement trials. (**b**) An example participant's mean ± SEM changes in reach paths across No-Movement

triplets from studies in which non-zero rotations (left) and error-clamps (right) were never applied (solid

lines: perturbation was CW, dashed lines: perturbation was CCW). (**c**) Boxplots showing STL in response

to different directions of simulated errors (No-Movement triplets indicated in magenta) from rotation (left, *n*

= 24) and error-clamp (right, *n* = 37) studies. (**d**) Estimated marginal means ± 95% confidence intervals

from the linear mixed models fit to each participant's STL performance summarized in (**c**). Asterisks indicate

statistically significant differences. (**e**) Mean ± SEM relative hand angles on the two trials after a perturbation

was presented on a No-Movement trial. Please refer to Supplemental Table 3 for detailed statistical results.

Boxplot centers: median, notches: 95% confidence interval of the median, box edges: 1st and 3rd quartiles,

whiskers: most extreme values within 1.5*IQR of the median. Statistical significance (* = p_{adj} < 0.05; n.s. =

p_{adj} ≥ 0.05) is indicated for selected comparisons. Abbreviations: STL – single-trial learning, CW –

clockwise, CCW – counterclockwise, Δ – change in.

Supplemental information accompanying the submission of "Learning from the path not taken:
Sensory prediction error computation and implicit motor adaptation proceed without movement"
3
4
5   Olivia A. Kim, Alexander D. Forrence, and Samuel D. McDougle
12   **Contents:**

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4

SUPPLEMENTAL FIGURE CAPTIONS

Supplemental Figure 1. Learning rates reported in the literature and observed in the current study.

Learning rates for motor adaptation observed in previous studies are shown at left in black, and learning

rates observed in each experiment in the current report are shown at the right, with data from Movement

triplets shown in green and data from No-Movement triplets shown in magenta. Data are shown as mean

± SEM, and are shown for rotational/error clamp perturbations of 15°, with the exception of Wei & Kording,

2009 E2, where an 11° perturbation was applied. Papers referred to and their corresponding reference

numbers: Kim et al., 2018[23]; Morehead et al., 2017[17]; Wei & Körding, 2009[27]. "Rotation, Go" and "Rotation,

Stop" show data from the in-lab experiment where participants saw 15° rotated feedback on Movement

trials (i.e., data from Fig. 2), "Rotation online, Go" and "Rotation online, Stop" show data from the online

experiment where participants saw 0-15° rotated feedback on Movement trials (i.e., data from Supplemental

Fig. 2). "Clamp online, Go" and "Clamp online, Stop" show data from the online experiment where

participants saw 0-15° error-clamed feedback (i.e., data from Supplemental Fig., 4). "Rotation 0 go" and

"Clamp 0 go" show data from the online experiments where participants saw 0° perturbed feedback on

Movement trials. Abbreviations: E, experiment.

Supplemental Figure 2. Single-trial learning in response to errors on Movement and No-Movement

trials during an online visuomotor adaptation task. (**a**) An example participant's mean ± SEM changes

in reach paths across triplets (green: triplets with perturbations on Movement trials, magenta: triplets with

perturbations on No-Movement trials, solid lines: perturbation was a CW rotation, dashed lines: perturbation

was a CCW rotation). (**b**) Boxplot showing STL across Movement (green) and No-Movement (magenta)

triplets for participants in an online version of the task described in Figure 1 ($n = 40$). (**c**) Estimated marginal

means (EMMs) ± 95% confidence intervals from the linear mixed model (LMM) fit to participants' STL

performance (summarized in **b**). The LMM (fixed effects: rotation [15° counterclockwise {CCW}, 0°, and 15°

clockwise {CW}], movement condition [Movement, No-Movement], rotation x movement condition

interaction; random effects: participant) revealed significant main effects of rotated cursor feedback ($F(2,$

$2223) = 136.46$, $p = 2.2 \times 10^{-16}$, *partial R^2* = 0.11) and movement condition ($F(1, 2248) = 4.74$, $p = 0.03$,

partial R^2 = 0.002), as well as a significant interaction ($F(2, 2229) = 12.40$, $p = 4.41 \times 10^{-6}$, *partial R^2* = 0.01).

Post-hoc pairwise comparisons of the EMMs from the model support the claim that rotated feedback

induced a statistically significant degree of STL on both Movement (0° vs 15° CW: $t(2227) = 9.14$, $p_{adj} =$

6.39×10^{-19}, *Cohen's d* $= 0.61$; 0° vs 15° CCW: $t(2220) = 7.81$, $p_{adj} = 2.61 \times 10^{-14}$, *Cohen's d* $= 0.52$) and

No-Movement trials (0° vs 15° CW: $t(2225) = 3.92$, $p_{adj} = 1.39 \times 10^{-4}$, *Cohen's d* $= 0.31$; 0° vs 15° CCW:

$t(2229) = 3.56$, $p_{adj} = 4.84 \times 10^{-4}$, *Cohen's d* $= 0.29$). Adaptation in the presence of a rotation was significantly

greater in Movement trials than No-Movement trials for CW ($t(2238) = 4.98$, $p_{adj} = 1.26 \times 10^{-6}$, *Cohen's d* $=$

0.37) and CCW rotations ($t(2239) = 2.06$, $p_{adj} = 0.04$, *Cohen's d* $= 0.15$). (**d**) Group mean ± SEM change in

(Δ) hand angle after exposure to Movement (green) and No-Movement (magenta) triplets' perturbations.

Positive Δ values indicate that the change in hand angle proceeded opposite the direction of the

perturbation (*i.e.,* in the direction that would counter the error). (**e**) Group mean ± SEM ratio of remembered

STL to STL. Remembered STL was statistically significantly greater than 0 for both Movement (one-sample

signed-rank test: $V = 819$, $p_{adj} = 1.09 \times 10^{-11}$, $r = 0.87$) and No-Movement triplets ($V = 769$, $p_{adj} = 9.69 \times 10^{-8}$

$r = 0.76$), but remembered STL did not significantly differ between movement conditions (paired-samples

signed-rank test: $V = 441$, $p_{adj} = 0.68$). (**f**) Scatter plot showing the relationship between individual subjects'

STL amplitude in the direction opposite the rotation on Movement and No-Movement trials. When we

considered instances of STL in the direction that would compensate for the observed error (update opposite

rotation, "Right-Way"), within-subject changes in hand angle were correlated between Movement and No-

Movement trials (*Pearson's r* $= 0.49$, $p_{adj} = 0.002$). (**g**) As in **f**, but showing data from trials with changes in

hand angle in the direction that would exacerbate the observed error (update in direction of rotation,

"Wrong-Way"). These ΔHand Angle values were uncorrelated between Movement and No-Movement trials

(*Pearson's r* $= 0.10$, $p_{adj} = 0.52$). These observations support the idea that the same learning process may

underlie adaptive single-trial learning events in response to errors on both kinds of trials, while maladaptive

changes in hand-angle may be attributable to potential sources of random noise. Boxplot centers: median,

notch: 95% confidence interval of the median, box edges: 1st and 3rd quartiles, whiskers: most extreme

value within 1.5*interquartile range of the median. Statistical significance (* = p < 0.05; n.s. = p ≥ 0.05) is.

Abbreviations: STL – single-trial learning, Δ – change, CW – clockwise, CCW – counterclockwise.

Supplemental Figure 3. Schematics illustrating the difference between rotation and error-clamp

perturbations. Diagrams showing the relationship between hand and cursor feedback movement

directions under rotational error regimes (left) and error-clamp regimes (right). When rotational

perturbations are applied to cursor feedback, the cursor's movement direction is contingent upon the

participant's movement direction, although there is an experimenter-induced offset. If a rotation is

consistently applied, participants explicit aiming strategies are a viable supplement for imperfect implicit

motor performance. When error-clamp perturbations are applied to cursor feedback, the cursor travels in a

fixed direction, regardless of the direction that the hand travels, rendering explicit aiming strategies useless.

Thus, error-clamp perturbations are often used in studies attempting to isolate implicit motor adaptation

processes.

Supplemental Figure 4. Single-trial learning in response to errors on Movement and No-Movement

trials with error-clamped feedback or simulated errors. (**a**) An example participant's mean ± SEM

changes in reach paths across triplets (green: triplets with perturbations on Movement trials, magenta:

triplets with perturbations on No-Movement trials, solid lines: perturbation was a CW error-clamp, dashed

lines: perturbation was a CCW error-clamp). (**b**) Boxplot showing STL across Movement (green) and No-

Movement (magenta) triplets for participants (n = 37) in an online experiment where cursor feedback was

error-clamped on Movement trials. (**c**) Estimated marginal means (EMMs) ± 95% confidence intervals from

the linear mixed model (LMM) fit to participants' STL performance (summarized in **b**). The LMM (fixed

effects: rotation [15° counterclockwise {CCW}, 0°, and 15° clockwise {CW}], movement condition

[Movement, No-Movement], error-clamp x movement condition interaction; random effects: participant)

revealed significant main effects of error-clamped cursor feedback (F(2, 1829) = 79.46, p = 2.2 x 10^{-16},

partial R^2 = 0.08) and an interaction between error-clamp and movement condition (F(2, 1832) = 8.45, p =

0.0002, *partial R^2* = 0.0003), although there was no main effect of movement condition (F(1, 1844) = 0.60,

p = 0.44). Post-hoc comparisons of the EMMs from the model revealed significant STL in response to non-

zero error-clamped feedback on both Movement (0° vs 15° CW: t(1827) = 7.55, p_{adj} = 3.08 x 10^{-13}, *Cohen's*

d = 0.56; 0° vs 15° CCW: t(1828) = 5.57, p_{adj} = 8.84 x 10^{-8}, *Cohen's d* = 0.41) and No-Movement trials (0°

vs 15° CW: t(1830) = 3.21, p_{adj} = 0.002, *Cohen's d* = 0.29; 0° vs 15° CCW: t(1832) = 2.25, p_{adj} = 0.03,

Cohen's d = 0.22). Adaptation in the presence of a 15° error-clamp was significantly greater on Movement

trials than No-Movement trials for CW ($t(1846)$ = 3.49, p_{adj} = 0.0009, *Cohen's d* = 0.29) and CCW clamps

($t(1846)$ = 2.29, p_{adj} = 0.03, *Cohen's d* = 0.19). Please refer to Supplemental Table 2 for further details on

post-hoc comparisons in this panel. (**d**) Group mean ± SEM change in (Δ) hand angle one and two trials

after exposure to Movement (green) and No-Movement (magenta) triplets' perturbations. Positive Δ values

indicate that the change in hand angle proceeded opposite the direction of the perturbation (*i.e.,* in the

direction that would counter the error). (**e**) Remembered STL shown as the ratio of relative hand angle 2

trials after experiencing a perturbation to the relative hand angle 1 trial after the perturbation (STL).

Remembered STL was significantly greater than 0 after both Movement (green; one-sample t-test: $t(36)$ =

11.31, p_{adj} = 6.23 x 10^{-13}, *Cohen's d* = 1.86) and No-Movement triplets (magenta, one-sample signed-rank

test: V = 579, p_{adj} = 5.95 x 10^{-5}, r = 0.64), but did not exhibit statistically significant differences between

movement conditions (paired t-test: $t(35)$ = 1.71, p_{adj} = 0.09). Remembered STL on No-Movement trials

could not be computed for one participant, so n = 36 instead of 37 in this panel. (**f**) Scatter plot showing the

relationship between individual subjects' STL amplitude in the direction opposite the error-clamp on

Movement and No-Movement trials (i.e., the "Right-Way"). Right-way changes in hand angle were

correlated between Movement and No-Movement trials (*Pearson's r* = 0.40, p_{adj} = 0.03). (**g**) As in (**f**), but

showing data from trials on which STL proceeded in the same direction as the error-clamp (*i.e.,* the "Wrong-

Way"). Wrong-Way changes in hand angle were not statistically significantly correlated between Movement

and No-Movement trials (r = 0.33, p_{adj} = 0.06). These observations support the idea that the same learning

process may underlie adaptive single-trial learning events in response to errors on both kinds of trials, while

maladaptive changes in hand-angle may be attributable to potential sources of random noise. Boxplot

centers: median, notch: 95% confidence interval of the median, box edges: 1st and 3rd quartiles, whiskers:

most extreme value within 1.5*interquartile range of the median. Statistical significance (* = p < 0.05; n.s.

= p ≥ 0.05) is indicated for selected comparisons. Abbreviations: STL – single-trial learning, Δ – change,

CW – clockwise, CCW – counterclockwise.

SUPPLEMENTAL TABLES

Supplemental Table 1.

Pairwise post-hoc comparisons between estimated marginal means in Supplemental Fig. 2c

Group 1		Group 2		Est. Diff.	t	df	FDR-adjusted p	Cohen's d
Trial Type	Rotation	Trial Type	Rotation					
M	0°	M	15° CW	-3.67°	-9.14	2227	* 6.39 x 10^{-19}	-0.61
M	0°	M	15° CCW	3.09°	7.81	2220	* 2.60 x 10^{-14}	0.52
M	15° CW	M	15° CCW	6.76°	17.08	2223	* 1.31 x 10^{-60}	1.13
No-M	0°	No-M	15° CW	-1.88°	-3.92	2225	* 0.0001	-0.31
No-M	0°	No-M	15° CCW	1.75°	3.56	2229	* 0.0005	0.29
No-M	15° CW	No-M	15° CCW	3.64°	7.41	2231	* 3.86 x 10^{-13}	0.61
M	0°	No-M	0°	0.41°	0.93	2240	0.35	--
M	15° CW	No-M	15° CW	2.20°	4.98	2238	* 1.25 x 10^{-6}	0.37
M	15° CCW	No-M	15° CCW	-0.93°	-2.06	2239	* 0.04	-0.15

Note. Abbreviations: M – Movement; No-M – No-Movement; Est. Diff. – Estimated Differences in degrees. Note that degrees of freedom pertain to the inputs to the LMM and are estimated using the Kenward-Rogers approach. Statistically significant p-values are indicated with asterisks. * p < 0.05.

132   **Supplemental Table 2.**

Pairwise post-hoc comparisons between estimated marginal means in Supplemental Fig. 4c

Group 1		Group 2		Est. Diff.	t	df	FDR-adjusted p	Cohen's d
Trial Type	Rotation	Trial Type	Rotation					
M	0°	M	15° CW	3.54°	-7.55	1827	* 3.08×10^{-13}	-0.56
M	0°	M	15° CCW	2.57°	5.57	1828	* 8.84×10^{-8}	0.41
M	15° CW	M	15° CCW	6.11°	13.14	1828	* 8.51×10^{-37}	0.97
No-M	0°	No-M	15° CW	-1.81°	-3.21	1830	* 0.002	-0.29
No-M	0°	No-M	15° CCW	1.29°	2.25	1832	* 0.03	0.20
No-M	15° CW	No-M	15° CCW	3.10°	5.46	1833	* 1.24×10^{-7}	0.49
M	0°	No-M	0°	0.08°	0.16	1846	0.87	--
M	15° CW	No-M	15° CW	1.81°	3.49	1846	* 0.0009	0.29
M	15° CCW	No-M	15° CCW	-1.19°	-2.28	1846	* 0.03	-0.19

Note. Abbreviations: M – Movement; No-M – No-Movement; Est. Diff. – Estimated Differences in degrees. Note that degrees of freedom pertain to the inputs to the model and are estimated using the Kenward-Rogers approach. Statistically significant p-values are indicated with asterisks. * p < 0.05.

134 **Supplemental Table 3.**

Pairwise post-hoc comparisons between estimated marginal means in Fig. 3

Group 1		Group 2		Est. Diff.	t	df	FDR-adjusted p	Cohen's d
Trial Type	Rotation	Trial Type	Rotation					
0° Rotation Applied on Movement Trials								
No-M	15° CCW	No-M	15° CW	2.45°	4.80	557	* 2.07 x 10^{-6}	0.40
0° Error-Clamped Feedback on Movement Trials								
No-M	15° CCW	No-M	15° CW	2.37°	4.32	805	* 0.0001	0.37
No-M	15° CW	No-M	0°	1.36°	2.48	801	* 0.020	0.21
No-M	15° CCW	No-M	0°	1.01°	1.83	800	0.067	--

Note. Abbreviations: No-M – No-Movement; Est. Diff. – Estimated Differences in degrees. Note that degrees of freedom pertain to the inputs to the model and are estimated using the Kenward-Rogers approach. Statistically significant p-values are indicated with asterisks. * $p < 0.05$.

135