1 **Movement planning and feedback are sufficient for sensory prediction error computation and**
2 **motor adaptation**
3

4 Olivia A. Kim[1], Alexander D. Forrence[2], and Samuel D. McDougle[2]
5
6 *1. Department of Psychology, Princeton University, Princeton, NJ 08544*
7 *2. Department of Psychology, Yale University, New Haven, CT 06511*
8
9
10
11
12 Correspondence to:
13   Olivia Kim
14   oakim@princeton.edu
15   kim.olivia.a@gmail.com

16
17
18
Number of Pages: 29
Number of Figures in Main Text: 3
Number of Supplemental Figures: 3
Number of Supplemental Tables: 3
References: 61 main text, 18 methods-only
Number of Words:
Abstract: 186
Introduction: 465
Results: 1,736
Main figure captions: 663
Discussion: 1,030
30
31
32
33
34 **AUTHOR CONTRIBUTIONS**
35 Conceptualization, OK, AF, & SM; Methodology, OK, AF, & SM; Software, AF & OK; Validation, OK & AF;
36 Formal Analysis, OK & AF; Investigation, OK & AF; Resources, OK & SM; Data Curation, OK & AF;
37 Supervision, SM; Project Administration, SM; Funding Acquisition, OK & SM; Writing – original draft, OK;
38 Writing – review & editing, OK, AF, & SM
39
40
41
42
43 **CONFLICT OF INTEREST**
44 The authors declare no competing financial interests.
45
46
47
48
ACKNOWLEDGEMENTS

This work was supported by a grant from the National Institutes of Health to OK (F32-NS122921).

ABSTRACT

Prediction errors guide many forms of learning, providing teaching signals that help us improve our

performance. Implicit motor adaptation, for instance, is driven by sensory prediction errors (SPEs), which

occur when the expected and observed consequences of a movement differ. Traditionally, SPE

computation is thought to require movement execution. However, recent work suggesting that the brain

generates and accounts for sensory predictions based on motor imagery or planning alone calls this

assumption into question. Here, by measuring implicit adaptation during a visuomotor task, we tested

whether motor planning and well-timed sensory feedback are sufficient for SPE computation. Human

participants were cued to reach to a target and were, on a subset of trials, rapidly cued to withhold these

movements. Errors displayed both on trials with and without movements induced single-trial implicit

learning. Learning following trials without movements persisted even when movement trials had never been

paired with errors, and when the direction of movement and sensory feedback trajectories were decoupled.

These observations demonstrate that the brain can compute SPEs without generating overt movements,

leading to the adaptation of planned movements even when they are not performed.

KEYWORDS

INTRODUCTION

Prediction errors help to optimize behavior by driving learning processes that correct for our

mistakes. Accordingly, they are thought to be a fundamental computation of the nervous system.[1] Specific

types of prediction errors are associated with specific learning processes, and sensory prediction errors

(SPEs) are the teachers of the implicit motor system, triggering adaptation and refinement of movements.[2−

5] Traditional formulations suggest that movement execution is critical for SPE computation.[6,7] However,

current thinking posits that the forward model estimates the consequences of movements before relevant

sensory feedback can reach the brain.[8] Consistent with this proposal, neurophysiological evidence

indicates that the cerebellum predictively encodes expected kinematics ahead of actual movement, and

that preparatory activity in the primary motor cortex reflects learned predictive changes during visuomotor

adaptation.[9−12] Thus, the literature indicates that the sensory consequences of intended movements are

predicted before those movements occur. This suggests that movement itself is not necessary for the

computation of sensory predictions or prediction errors.

Recent work supports the claim that the brain predicts the sensory consequences of movements

before they can be performed, even when the agent does not have a clear intention to move.[13−17]

Considering that sensorimotor prediction does not in theory require movement, it may be that a prediction

can be combined with an observation to support SPE computation without any actual motor execution. In

other words, SPEs should be effectively computed based upon only two events − the generation of a

sensory prediction and the observation of sensory feedback (**Fig. 1a**).

Some prior work has illustrated that higher-level cognitive processes support visuomotor learning

without movement, for instance when observers witness others' motor errors: motor learning in this case

might be driven by SPEs, or by other types of performance errors beyond SPE (e.g., reward prediction

errors), or perhaps by a combination of multiple sources of error.[18−20] Here, we isolated implicit motor

adaptation to specifically test whether SPE computation requires movement execution, as SPEs are both

necessary and sufficient for this form of learning.[21−27] Having isolated implicit motor adaptation, we then

asked whether withheld movements could undergo adaptation.

Figure 1. Schematics showing the proposed learning framework and task design. (**a**) Schematic
showing how the forward model may support implicit motor adaptation in the presence of sensory feedback
not causally related to self-generated movement. (**b**) Events on trials with visual feedback. The robotic
apparatus brought the participant's hand to the starting location to initiate a trial. On Movement trials (top),
the target turned green (GO), cueing participants to reach through the target. On trials with visual feedback,
participants observed a white feedback cursor move along a rotated trajectory (Rotation). On No-Movement
trials (bottom), the target turned magenta 100 ms after turning green, cueing participants to withhold
movement (STOP). After a delay, an animation showing the feedback cursor moving 15° off-target played
(Animation). The hand is shown in the figure for illustrative purposes but was not visible during the
experiment. (**c**) How single-trial learning (STL) was computed using a triplet paradigm. Triplets were
composed of 2 Movement trials without visual feedback flanking either a Movement or a No-Movement trial
with visual feedback. STL was measured as the difference between reach angles on the flanking trials. (**d**)
The pseudorandomized order in which trials were presented for an example participant. Color indicates
movement condition (Movement: green, No-Movement: magenta).

To that end, we measured trial-by-trial implicit adaptation during a visuomotor task in which human

participants saw visual feedback while performing – or withholding – hand and arm movements, using a

modified stop-signal paradigm. To isolate implicit adaptation, we employed a recently-developed approach

that requires participants to aim their movements directly for presented targets and disregard visual

feedback.[26,28–32] We predicted that single-trial motor adaptation would occur following both typical

movement trials that generated sensory error, as well as trials where movements were withheld but

"simulated" sensory errors were observed. If confirmed, this result would demonstrate that the brain can

compute SPEs in the absence of movement and can drive the adaptation of planned movements that were

not performed.

RESULTS

Simulated and typical visuomotor rotations cause motor adaptation

In our first experiment, we measured implicit motor adaptation in humans (n = 20) performing or

withholding straight reaches during a visuomotor adaptation task (**Fig. 1b**). Vision of the hand and arm was

occluded by a mirror that reflected visual feedback from a horizontally mounted monitor. A white cursor

provided feedback about participants' hand positions as they reached from a starting location to a displayed

target. After a brief acclimation period, trials were organized into triplets, such that each trial with cursor

feedback was flanked by trials without cursor feedback. This allowed for a reliable measurement of single-

trial learning (STL) in response to feedback, quantified as the difference between the direction of hand

movement (hand angle) on the first and third trials of each triplet (**Fig. 1c**). Trials with cursor feedback were

either Movement trials on which a Go signal prompted movement or No-Movement trials on which a Stop

signal immediately followed the Go signal, indicating that movements should be withheld. On Movement

trials, feedback involved a visuomotor error (±15° rotation added to the visual cursor path; + = clockwise;

Fig. 1b right). On No-Movement trials, sensory feedback involved a simulation of the cursor's path, using

timing variables based on ongoing measurements of participant behavior (see *Methods*). All flanking trials

of each triplet were Go trials and required movements. The direction of the error (clockwise [CW] or

counterclockwise [CCW]) was pseudorandomly varied across triplets to maintain overall adaptation near 0

throughout the session (**Fig. 1d**). This straightforward design allowed us to test the hypothesis that SPE

computation and motor adaptation do not require that movement and sensory feedback to be causally

linked (**Fig. 1a**).

Consistent with our predictions, rotated cursor paths on Movement and No-Movement trials both

caused subsequent hand trajectories to shift opposite the direction of the rotation (**Fig. 2b-c**), with a 2-way

repeated measures ANOVA revealing statistically significant main effects of the direction (CW vs CCW) of

the perturbation applied (F(1, 19) = 98.62, p = 5.89 x 10^{-9}, η_G^2 = 0.76). While there was no significant main

effect of withholding movement (F(1, 19) = 1.79, p = 0.20), we observed a significant interaction between

the perturbation applied and withholding movement ($F(1, 19) = 137.32$, $p = 3.87 \times 10^{-10}$, $\eta_G^2 = 0.49$). Post-

hoc pairwise comparisons confirmed that STL was sensitive to perturbation direction during both Movement

(paired t-test: $t(19) = 12.92$, $p_{adj} = 2.96 \times 10^{-10}$, *Cohen's d* = 5.12) and No-Movement triplets ($t(19) = 4.39$,

$p_{adj} = 3.13 \times 10^{-4}$, *Cohen's d* = 1.63), and also indicated that STL magnitude was greater across Movement

than No-Movement triplets (paired-samples signed-rank test, CW rotations: $V = 210$, $p_{adj} = 2.55 \times 10^{-6}$, $r =$

0.88; CCW rotations: $t(19) = 9.43$, $p_{adj} = 2.70 \times 10^{-8}$, *Cohen's d* = 2.02) rotations. The overall amplitude of

adaptation observed both with and without movement was within the range of implicit learning rates

measured in previous studies (**Supplemental Fig. 1**).

Figure 2. Effects of typical and simulated errors during a visuomotor reach adaptation task. (**A**)
Schematic illustrating how STL and remembered STL measurements were computed. (**B**) An example
subject's (top) and the group's (bottom) mean ± SEM changes in reach paths across triplets with a rotation
applied (green: triplets with perturbations on Movement trials, magenta: triplets with perturbations on No-
Movement trials, solid lines: perturbation was a CW rotation, dashed lines: perturbation was a CCW

rotation). (**C**) STL across Movement (green) and No-Movement (magenta) triplets for all participants (*n* =
20). Positive changes in hand angle are CCW. Refer to Supplemental Table 1 for details on all statistical
tests. (**D**) Group mean ± SEM Δhand angle values after exposure to Movement (green) and No-Movement
(magenta) trial perturbations. Positive Δ values indicate that the change in hand angle proceeded opposite
the direction of the perturbation (*i.e.,* the direction that would counter the error, "Right-Way"). (**E**) Group
mean of participants' ratios of remembered STL to initial STL during Movement and No-Movement trials.
(**F**) The relationship between Right-Way STL observed during Movement and No-Movement triplets. (**G**)
As in (**F**), but for STL observed on trials where adaptation proceeded in the direction that would exacerbate
the error (*i.e.,* the same direction as the perturbation applied, "Wrong Way"). Statistical significance (* = p_{adj}
< 0.05; n.s. = p_{adj} ≥ 0.05) is indicated. Abbreviations: STL – single-trial learning, CW – clockwise, CCW –
counterclockwise, Δ – change in.

Supplemental Figure 1. Learning rates reported in the literature and observed in the current study.
Learning rates for motor adaptation observed in previous studies are shown at left in black, and learning
rates observed in each experiment in the current report are shown at the right, with data from Movement
triplets shown in green and data from No-Movement triplets shown in magenta. Data are shown as mean
± SEM, and are shown for rotational/error clamp perturbations of 15°, with the exception of Wei & Kording,
2009 E2, where an 11° perturbation was applied. Papers referred to and their corresponding reference
numbers: Kim et al., 2018[29]; Morehead et al., 2017[26]; Wei & Körding, 2009[33]. "Rotation, Go" and "Rotation,
Stop" show data from the in-lab experiment where participants saw 15° rotated feedback on Movement
trials (i.e., data from Fig. 2), "Rotation online, Go" and "Rotation online, Stop" show data from the online
experiment where participants saw 0-15° rotated feedback on Movement trials (i.e., data from Supplemental
Fig. 2). "Clamp online, Go" and "Clamp online, Stop" show data from the online experiment where
participants saw 0-15° error-clamed feedback (i.e., data from Supplemental Fig., 4). "Rotation 0 go" and
"Clamp 0 go" show data from the online experiments where participants saw 0° perturbed feedback on
Movement trials. Abbreviations: E, experiment.

To address whether observed STL measured genuine implicit learning, we checked whether

adaptation persisted beyond the trial after an error was experienced. We examined participants' hand

angles on the second trial after a perturbation relative to the pre-perturbation baseline trial (i.e., hand angle

on trial 1 of triplet *t + 1* relative to hand angle on trial 1 of triplet *t,* subsequently referred to as remembered

STL, **Fig. 2a**). As visual feedback was withheld on both trial types, this approach provided a pure measure

of persistent memory in the absence of error-driven changes in performance. Hand angle remained adapted

in the direction opposite the rotation on trials with nonzero perturbations regardless of movement condition

(**Fig. 2d**), suggesting that genuine implicit learning was observed in response to errors under both

movement conditions. Closer examination of the relative ratio of remembered STL to initial STL revealed

that retention of adaptation differed significantly from zero after both Movement (t(19) = 26.20, p_{adj} = 6.71 x

10^{-16}, *Cohen's d* = 5.86) and No-Movement triplets (t(19) = 9.20, p_{adj} = 2.98 x 10^{-8}, *Cohen's d* = 2.06), and

the amount of retention observed was not statistically significantly different between the movement

conditions (t(19) = 2.07, p_{adj} = 0.053, **Fig. 2e**).

To assess the potential similarity of mechanisms underlying adaptation after errors on Movement

and No-Movement trials, we compared STL amplitude under each condition, reasoning that there should

be a reliable relationship between the two measures if STL is supported by the same mechanism following

both Movement and No-Movement trials. When we considered instances of STL in the direction that would

compensate for the observed error (the direction opposite the rotation, i.e., the "Right-Way"), within-subject

changes in hand angle were correlated between Movement and No-Movement trials (*Pearson's r* = 0.55, *p*

= 0.01; **Fig. 2f**). Conversely, changes in hand angle in the direction that would exacerbate the observed

error (the direction of the rotation, i.e., the "Wrong-Way") were uncorrelated between Movement and No-

Movement trials (*Pearson's r* = 0.17, *p* = 0.48, **Fig. 2g**). Together, these observations suggest that the

same learning process may underlie adaptive STL events in response to errors during both kinds of trials,

while maladaptive changes in hand angle may be attributable to noise.

Implicit motor adaptation proceeds after simulated errors in an online visuomotor task

Illustrating that the above observations are reproducible and generalize across experimental

contexts, we again observed that simulated errors in No-Movement trials also induced motor adaptation in

an online, crowd-sourced version of the task. Participants (*n* = 40) made hand movements using their

computer mouse or trackpad to move a cursor towards a target. As in the experiment described above,

trials were presented in triplets, allowing us to measure STL in response to cursor feedback presented

during Movement and No-Movement trials at the center of each triplet (**Fig. 1b-d**). For this online study,

triplets with 0° perturbations/simulated errors were also included to provide an estimate of baseline changes

in hand angle, in the event that participants exhibited strong movement biases in the online platform.

STL was directionally appropriate for the perturbation applied during both Movement and No-

Movement trials (**Supplemental Fig. 2a-c**, please refer to the supplemental material details of the statistical

analysis). Further echoing the results of the in-person study, STL on both Movement and No-Movement

trials was retained beyond the triplet in which the relevant error occurred (**Supplemental Fig. 2d-e**), and

the magnitude of STL in the direction that would counter the perturbation was again correlated across the

two movement conditions (**Supplemental Fig. 2e**). These data provide further support for the claim that

movements that are not performed can undergo implicit motor adaptation, and they extend our findings to

a task with different movement demands (e.g., finger or wrist movements versus full, center-out reaches).

Supplemental Figure 2. Single-trial learning in response to errors on Movement and No-Movement
trials during an online visuomotor adaptation task. (**a**) An example participant's mean ± SEM changes
in reach paths across triplets (green: triplets with perturbations on Movement trials, magenta: triplets with
perturbations on No-Movement trials, solid lines: perturbation was a CW rotation, dashed lines: perturbation
was a CCW rotation). (**b**) Boxplot showing STL across Movement (green) and No-Movement (magenta)
triplets for participants in an online version of the task described in Figure 1 (*n* = 40). (**c**) Estimated marginal
means (EMMs) ± 95% confidence intervals from the linear mixed model (LMM) fit to participants' STL
performance (summarized in **b**). The LMM (fixed effects: rotation [15° counterclockwise {CCW}, 0°, and 15°
clockwise {CW}], movement condition [Movement, No-Movement], rotation x movement condition

interaction; random effects: participant) revealed significant main effects of rotated cursor feedback (F(2,
2223) = 136.46, p = 2.2 x 10^{-16}, *partial R^2* = 0.11) and movement condition (F(1, 2248) = 4.74, p = 0.03,
partial R^2 = 0.002), as well as a significant interaction (F(2, 2229) = 12.40, p = 4.41 x 10^{-6}, *partial R^2* = 0.01).
Post-hoc pairwise comparisons of the EMMs from the model support the claim that rotated feedback
induced a statistically significant degree of STL on both Movement (0° vs 15° CW: t(2227) = 9.14, p_{adj} =
6.39 x 10^{-19}, *Cohen's d* = 0.61; 0° vs 15° CCW: t(2220) = 7.81, p_{adj} = 2.61 x 10^{-14}, *Cohen's d* = 0.52) and
No-Movement trials (0° vs 15° CW: t(2225) = 3.92, p_{adj} = 1.39 x 10^{-4}, *Cohen's d* = 0.31; 0° vs 15° CCW:
t(2229) = 3.56, p_{adj} = 4.84 x 10^{-4}, *Cohen's d* = 0.29). Adaptation in the presence of a rotation was significantly
greater in Movement trials than No-Movement trials for CW (t(2238) = 4.98, p_{adj} = 1.26 x 10^{-6}, *Cohen's d* =
0.37) and CCW rotations (t(2239) = 2.06, p_{adj} = 0.04, *Cohen's d* = 0.15). Please refer to Supplemental Table
1 for full details on post-hoc pairwise comparisons. (**d**) Group mean ± SEM change in (Δ) hand angle after
exposure to Movement (green) and No-Movement (magenta) triplets' perturbations. Positive Δ values
indicate that the change in hand angle proceeded opposite the direction of the perturbation (*i.e.,* in the
direction that would counter the error). (**e**) Group mean ± SEM ratio of remembered STL to STL.
Remembered STL was statistically significantly greater than 0 for both Movement (one-sample signed-rank
test: V = 819, p_{adj} = 1.09 x 10^{-11}, r = 0.87) and No-Movement triplets (V = 769, p_{adj} = 9.69 x 10^{-8}, r = 0.76),
but remembered STL did not significantly differ between movement conditions (paired-samples signed-rank
test: V = 441, p_{adj} = 0.68). (**f**) Scatter plot showing the relationship between individual subjects' STL
amplitude in the direction opposite the rotation on Movement and No-Movement trials. When we considered
instances of STL in the direction that would compensate for the observed error (update opposite rotation,
"Right-Way"), within-subject changes in hand angle were correlated between Movement and No-Movement
trials (*Pearson's r* = 0.49, p_{adj} = 0.002). (**g**) As in **f**, but showing data from trials with changes in hand angle
in the direction that would exacerbate the observed error (update in direction of rotation, "Wrong-Way").
These ΔHand Angle values were uncorrelated between Movement and No-Movement trials (*Pearson's r* =
0.10, p_{adj} = 0.52). These observations support the idea that the same learning process may underlie
adaptive single-trial learning events in response to errors on both kinds of trials, while maladaptive changes
in hand angle may be attributable to potential sources of random noise. Boxplot centers: median, notch:
95% confidence interval of the median, box edges: 1st and 3rd quartiles, whiskers: most extreme value within
1.5*interquartile range of the median. Statistical significance (* = p < 0.05; n.s. = p ≥ 0.05) is. Abbreviations:
STL – single-trial learning, Δ – change, CW – clockwise, CCW – counterclockwise.

Supplemental Table 1.

Pairwise post-hoc comparisons between estimated marginal means in Supplemental Fig. 2c

Group 1		Group 2		Est. Diff.	t	df	FDR-adjusted p	Cohen's d
Trial Type	Rotation	Trial Type	Rotation					
M	0°	M	15° CW	-3.67°	-9.14	2227	* 6.39 x 10⁻¹⁹	-0.61
M	0°	M	15° CCW	3.09°	7.81	2220	* 2.60 x 10⁻¹⁴	0.52
M	15° CW	M	15° CCW	6.76°	17.08	2223	* 1.31 x 10⁻⁶⁰	1.13
No-M	0°	No-M	15° CW	-1.88°	-3.92	2225	* 0.0001	-0.31
No-M	0°	No-M	15° CCW	1.75°	3.56	2229	* 0.0005	0.29
No-M	15° CW	No-M	15° CCW	3.64°	7.41	2231	* 3.86 x 10⁻¹³	0.61
M	0°	No-M	0°	0.41°	0.93	2240	0.35	--
M	15° CW	No-M	15° CW	2.20°	4.98	2238	* 1.25 x 10⁻⁶	0.37
M	15° CCW	No-M	15° CCW	-0.93°	-2.06	2239	* 0.04	-0.15

Note. Abbreviations: M – Movement; No-M – No-Movement; Est. Diff. – Estimated Differences in degrees. Note that degrees of freedom pertain to the inputs to the LMM and are estimated using the Kenward-Rogers approach. Statistically significant p-values are indicated with asterisks. * p < 0.05.

Motor adaptation during No-Movement triplets does not depend on participants' control over cursor

trajectory during Movement trials

We note that rotated visual feedback on Movement trials was sensitive to people's actual reaching

directions because the rotation was simply added to the measured reach direction, as is typical in

visuomotor rotation tasks. It is possible that these directional contingencies affected participants' responses

to error, potentially encouraging them to attempt to deliberately control the cursor's position via an explicit

re-aiming process.[27] To rule this out, we recruited a new group of participants ($n = 37$) to perform a variant

of the task where the visual cursor moved in a fixed path ("error-clamped" feedback[26]; **Supplemental Fig.**

3a) in one of three directions (0° or 15° CW/CCW) on the trials with feedback.

Replicating and extending the findings reported above, participants assigned to the error-clamp

condition exhibited STL after Movement and No-Movement trials (**Supplemental Fig. 3b-d**; please refer to

supplemental material for details). We also observed significant retention of STL (**Supplemental Fig. 3e-f**)

and a significant correlation between STL amplitude on Movement and No-Movement trials where

adaptation proceeded opposite the direction of the perturbation (**Supplemental Fig. 3g-h**). These data

further strengthen the claim that motor adaptation does not require movement.

Supplemental Figure 3. Single-trial learning in response to errors on Movement and No-Movement
trials with error-clamped feedback or simulated errors. (**a**) Diagrams showing the relationship between
hand and cursor feedback movement directions under rotational (left) and error-clamp regimes (right).
When rotations are applied, the cursor's movement direction is contingent upon the participant's movement
direction. When error-clamp perturbations are applied, the cursor travels in a fixed direction, regardless of
the direction that the hand travels. As error-clamp perturbations render deliberate changes in movement
direction useless, they are often used in studies attempting to isolate implicit motor adaptation processes.
(**b**) An example participant's mean ± SEM changes in reach paths across triplets (green: triplets with
perturbations on Movement trials, magenta: triplets with perturbations on No-Movement trials, solid lines:
perturbation was CW error-clamp, dashed lines: perturbation was CCW error-clamp). (**c**) Boxplot showing

STL across Movement (green) and No-Movement (magenta) triplets for participants (n = 37) in an online
experiment where cursor feedback was error-clamped on Movement trials. (**d**) Estimated marginal means
(EMMs) ± 95% confidence intervals from the linear mixed model (LMM) fit to participants' STL performance
(summarized in **c**). The LMM (fixed effects: rotation [15° counterclockwise {CCW}, 0°, and 15° clockwise
{CW}], movement condition [Movement, No-Movement], error-clamp x movement condition interaction;
random effects: participant) revealed significant main effects of error-clamped cursor feedback (F(2, 1829)
= 79.46, p = 2.2 x 10^{-16}, *partial R^2* = 0.08) and an interaction between error-clamp and movement condition
(F(2, 1832) = 8.45, p = 0.0002, *partial R^2* = 0.0003), although there was no main effect of movement
condition (F(1, 1844) = 0.60, p = 0.44). Post-hoc comparisons of the EMMs from the model revealed
significant STL in response to non-zero error-clamped feedback on both Movement (0° vs 15° CW: t(1827)
= 7.55, p_{adj} = 3.08 x 10^{-13}, *Cohen's d* = 0.56; 0° vs 15° CCW: t(1828) = 5.57, p_{adj} = 8.84 x 10^{-8}, *Cohen's d*
= 0.41) and No-Movement trials (0° vs 15° CW: t(1830) = 3.21, p_{adj} = 0.002, *Cohen's d* = 0.29; 0° vs 15°
CCW: t(1832) = 2.25, p_{adj} = 0.03, *Cohen's d* = 0.22). Adaptation in the presence of a 15° error-clamp was
significantly greater on Movement trials than No-Movement trials for CW (t(1846) = 3.49, p_{adj} = 0.0009,
Cohen's d = 0.29) and CCW clamps (t(1846) = 2.29, p_{adj} = 0.03, *Cohen's d* = 0.19). Please refer to
Supplementary Table 2 for further details on post-hoc comparisons in this panel. (**e**) Group mean ± SEM
change in (Δ) hand angle one and two trials after exposure to Movement (green) and No-Movement
(magenta) triplets' perturbations. Positive Δ values indicate that the change in hand angle proceeded
opposite the direction of the perturbation (*i.e.,* in the direction that would counter the error). (**f**) Remembered
STL shown as the ratio of relative hand angle 2 trials after experiencing a perturbation to the relative hand
angle 1 trial after the perturbation (STL). Remembered STL was significantly greater than 0 after both
Movement (green; one-sample t-test: t(36) = 11.31, p_{adj} = 6.23 x 10^{-13}, *Cohen's d* = 1.86) and No-Movement
triplets (magenta, one-sample signed-rank test: V = 579, p_{adj} = 5.95 x 10^{-5}, r = 0.64), but did not exhibit
statistically significant differences between movement conditions (paired t-test: t(35) = 1.71, p_{adj} = 0.09).
Remembered STL on No-Movement trials could not be computed for one participant, so n = 36 instead of
37 in this panel. (**g**) Scatter plot showing the relationship between individual subjects' STL amplitude in the
direction opposite the error-clamp on Movement and No-Movement trials (i.e., the "Right-Way"). Right-way
changes in hand angle were correlated between Movement and No-Movement trials (*Pearson's r* = 0.40,
p_{adj} = 0.03). (**h**) As in (**g**), but showing data from trials on which STL proceeded in the same direction as the
error-clamp (*i.e.*, the "Wrong-Way"). Wrong-Way changes in hand angle were not statistically significantly
correlated between Movement and No-Movement trials (r = 0.33, p_{adj} = 0.06). Boxplot centers: median,
notch: 95% confidence interval of the median, box edges: 1st and 3rd quartiles, whiskers: most extreme
value within 1.5*interquartile range of the median. Statistical significance (* = p < 0.05; n.s. = p ≥ 0.05) is
indicated for selected comparisons. Abbreviations: STL – single-trial learning, Δ – change, CW – clockwise,
CCW – counterclockwise.

Supplemental Table 2.

Pairwise post-hoc comparisons between estimated marginal means in Supplemental Fig. 3d

Group 1		Group 2		Est. Diff.	t	df	FDR-adjusted p	Cohen's d
Trial Type	Rotation	Trial Type	Rotation					
M	0°	M	15° CW	3.54°	-7.55	1827	* 3.08 x 10⁻¹³	-0.56
M	0°	M	15° CCW	2.57°	5.57	1828	* 8.84 x 10⁻⁸	0.41
M	15° CW	M	15° CCW	6.11°	13.14	1828	* 8.51 x 10⁻³⁷	0.97
No-M	0°	No-M	15° CW	-1.81°	-3.21	1830	* 0.002	-0.29
No-M	0°	No-M	15° CCW	1.29°	2.25	1832	* 0.03	0.20
No-M	15° CW	No-M	15° CCW	3.10°	5.46	1833	* 1.24 x 10⁻⁷	0.49
M	0°	No-M	0°	0.08°	0.16	1846	0.87	--
M	15° CW	No-M	15° CW	1.81°	3.49	1846	* 0.0009	0.29
M	15° CCW	No-M	15° CCW	-1.19°	-2.28	1846	* 0.03	-0.19

Note. Abbreviations: M – Movement; No-M – No-Movement; Est. Diff. – Estimated Differences in degrees. Note that degrees of freedom pertain to the inputs to the model and are estimated using the Kenward-Rogers approach. Statistically significant p-values are indicated with asterisks. * p < 0.05.

Adaptation on No-Movement trials does not depend on within-session adaptation on Movement trials

In two further experiments, we asked if adaptation to errors in the No-Movement condition was

contingent on sharing a context with the Movement condition. In other words, if learning in the No-

Movement condition only occurs when there are neighboring trials in the Movement condition producing

typical SPEs, it is possible that adaptive responses observed in the No-Movement condition reflect a

"cueing" effect, whereby an adapted sensorimotor map is cued by observation of the visual error and then

retrieved on the subsequent trial(s).[34,35] While our previous retention (**Fig. 2d-e**, **Supplemental Fig. 2d-e**,

Supplemental Fig. 3e-f) and correlation (**Fig. 2f-g**, **Supplemental Fig. 2f-g**, **Supplemental Fig. 3g-h**)

results argue against this interpretation as they suggest a shared learning mechanism across movement

conditions, we opted to directly test this alternative explanation in another pair of experiments. Here, we

only included 0° rotated (**Fig. 3a**, left, *n* = 24 participants) or clamped (**Fig. 3a**, right, *n* = 37 participants)

error feedback on Movement trials, but maintained 0° or 15° CW/CCW errors on the No-Movement trials.

Thus, visual perturbations were never paired with movement. The key results were again replicated –

learning was preserved in the No-Movement condition even when error feedback had never been

associated with executed movements (**Fig. 3c**, rotation: LMM: $F_{(557)}$ 23.01, $p = 2.07 \times 10^{-6}$, *partial R^2* =

0.04; error-clamp: $F_{(802)} = 9.41$, $p = 9.14 \times 10^{-5}$, *partial R^2 = 0.02*). Post-hoc pairwise comparisons showed

that adaptation was significantly different between triplets with clockwise and counterclockwise errors for

both the rotation ($t_{(557)} = 4.80$, $p = 2.07 \times 10^{-6}$, *Cohen's d* = 0.4) and error-clamp experiments ($t_{(1453)} =$

4.32, $p = 5.34 \times 10^{-5}$, *Cohen's d* = 0.37) – a hallmark of implicit motor adaptation (please refer to

Supplemental Table 3 for all post-hoc test results). Overall levels of STL observed on No-Movement trials

were comparable during these two experiments to those discussed above, and within the range of learning

rates previously observed in the literature (**Supplemental Fig. 1**). Furthermore, both groups of participants

showed retention of STL that differed significantly from 0 (rotation, mean ± SEM: 0.53 ± 0.06 retention ratio,

one-sample t-test: $t_{(22)} = 8.28$, $p = 3.34 \times 10^{-8}$, *Cohen's d* = 1.73; error-clamp, median: 0.45, interquartile-

range: 0.58, one-sample signed-rank test: $V = 507$, $p = 0.001$, $r = 0.53$). Overall, these experiments support

the hypothesis that motor adaptation can proceed without movement execution.

Figure 3. Effects of simulated errors when perturbations were never applied during Movement trials.
(**a**) Schematic illustrating the relationship between movement and visual feedback on Movement trials
during an experiment where visuomotor rotations (left) or error-clamps (right) were never applied during
Movement trials. (**b**) An example participant's mean ± SEM changes in reach paths across No-Movement
triplets from studies in which non-zero rotations (left) and error-clamps (right) were never applied (solid

lines: perturbation was CW, dashed lines: perturbation was CCW). (**c**) Boxplots showing STL in response
to different directions of simulated errors (No-Movement triplets indicated in magenta) from rotation (left, *n*
= 24) and error-clamp (right, *n* = 37) studies. (**d**) Estimated marginal means ± 95% confidence intervals
from the linear mixed models fit to each participant's STL performance summarized in (**c**). Asterisks indicate
statistically significant differences. (**e**) Mean ± SEM relative hand angles on the two trials after a perturbation
was presented on a No-Movement trial. Please refer to Supplemental Table 3 for detailed statistical results.
Boxplot centers: median, notches: 95% confidence interval of the median, box edges: 1st and 3rd quartiles,
whiskers: most extreme values within 1.5*IQR of the median. Statistical significance (* = $p_{adj} < 0.05$; n.s. =
$p_{adj} \geq 0.05$) is indicated for selected comparisons. Abbreviations: STL – single-trial learning, CW –
clockwise, CCW – counterclockwise, Δ – change in.

Supplemental Table 3.

Pairwise post-hoc comparisons between estimated marginal means in Fig. 3

Group 1		Group 2		Est. Diff.	t	df	FDR-adjusted p	Cohen's d
Trial Type	Rotation	Trial Type	Rotation					
0° Rotation Applied on Movement Trials								
No-M	15° CCW	No-M	15° CW	2.45°	4.80	557	* 2.07 x 10^{-6}	0.40
0° Error-Clamped Feedback on Movement Trials								
No-M	15° CCW	No-M	15° CW	2.37°	4.32	805	* 0.0001	0.37
No-M	15° CW	No-M	0°	1.36°	2.48	801	* 0.020	0.21
No-M	15° CCW	No-M	0°	1.01°	1.83	800	0.067	--

Note. Abbreviations: No-M – No-Movement; Est. Diff. – Estimated Differences in degrees. Note that

degrees of freedom pertain to the inputs to the model and are estimated using the Kenward-Rogers

approach. Statistically significant p-values are indicated with asterisks. * p < 0.05.

DISCUSSION

Our results demonstrate that movements can be implicitly refined even when they are not

performed. Participants who were cued to reach towards a target but suppressed that movement after

observation of a Stop cue showed consistent, robust STL in response to simulated errors (**Figs. 2-3**,

Supplemental Fig. 2, **Supplemental Fig. 3**). As implicit learning necessarily proceeds following SPEs, our

data also provide evidence that SPEs are computed even when movements are not performed. These

findings strongly support the fundamental assumptions of predictive processing frameworks of motor

adaptation, where precise sensory predictions are generated from a movement intent (or "plan", "goal") and

compared against sensory observations to induce error-based learning.[15,17,36,37]

We argue that we have measured learning via an implicit process, and, by extension, that the STL

observed in our study provides evidence that SPEs are computed regardless of whether a movement is

performed. Although visuomotor tasks sometimes recruit cognitive strategies (e.g., deliberate "re-aiming"

of movements), multiple factors indicate that our studies successfully measured implicit adaptation.[27,38,39]

First, participants were instructed to ignore the displayed cursor and try to contact the target on every trial,

a straightforward technique which has been consistently shown to eliminate the explicit re-aiming of

movements.[24,26,28–30,32] Second, randomization of the presence and direction of errors discouraged explicit

learning, reducing motivation to apply ineffective re-aiming strategies (see [40]). Third, data from participants

who did not understand or recall the instruction to always aim directly at the target were excluded (see

Methods), decreasing the likelihood that strategic re-aiming contaminated the analysis. Fourth, adaptation

persisted into subsequent no-feedback trials (**Fig. 2d-e**, **Fig. 3e**, **Supplemental Fig. 2d-e**, **Supplemental**

Fig. 3e-f), consistent with lingering implicit motor learning; it is unlikely that strategies would be maintained

through trials where no feedback is present. Fifth, the magnitude of STL observed was generally consistent

with multiple previous studies that measured implicit motor adaptation rates (**Supplemental Fig. 1**).[26,29,41]

Lastly, the adaptation effects observed in the No-Movement conditions were not attributable to recall of

learning that occurred on Movement trials (**Fig. 3**). Our data thus provide striking evidence that movement

is not required for implicit adaptation, and, by extension, SPE computation.

While motor planning and concurrent sensory observations are sufficient to drive SPE computation

and motor adaptation, our data also indicate that participants showed significantly stronger STL over triplets

with Movement trials versus No-Movement trials. This suggests that movement provides additional training

input to the brain. Interestingly, this is consistent with patterns of cerebellar activity during motor behaviors

and current thinking about mechanisms for learning in cerebellar-dependent tasks like implicit reach

adaptation.[26] Purkinje cell complex spikes are a powerful teaching signal in the cerebellum, and these

complex spikes exhibit firing patterns that may be movement-dependent.[42–45] During target-directed

reaching, complex spikes related to reach goal locations are generated *after* reach onset.[46] If these complex

spikes are tied to motor performance and not motor planning, then the absence of these error signals on

No-Movement trials may account for reduced levels of STL without movement.[47–50] Another non-mutually

exclusive possibility is that the precise timing of SPEs is less effective in our No-Movement condition than

under normal, movement conditions: in the former case, the timing of simulated feedback is controlled by

the experimenter and not triggered by the subject's actual movement, potentially adding a novel source of

noise into the adaptation process.[51,52] Irrespective of the fact that STL was of lesser amplitude across No-

Movement than Movement triplets, our data demonstrate the significant influence of the brain's prediction

signals on learning – even without the ability to directly attribute sensory feedback to an actual movement,

prediction of a planned movement's sensory consequences supports the error computations that drive

adaptation of future behavior.

Our findings add to a body of work indicating that many forms of motor learning do not strictly

require movement-based practice. For instance, in [53], after human participants observed others adapting

to a force field applied during reaching movements, the observers were able to partially compensate for

that same force field when they encountered it themselves. Interestingly, this observational learning did not

proceed if participants were executing other task-irrelevant movements during the observation period. This

finding combined with subsequent neuroimaging data showing that observational learning recruits brain

areas associated with motor planning were taken to suggest that engagement in a covert motor planning

process may be the critical element allowing for force-field adaptation following error observation without

movement.[53–55] Together, this related prior work and the evidence we have provided that motor planning is

sufficient for the generation of sensory predictions and can support SPE computations suggest that there

are multiple routes to inducing motor planning and ultimately driving motor adaptation.

Other reports in the motor learning literature have provided evidence for cognitive compensation

for observed motor errors during reaching, improved visual tracking following observation of target

movement without engagement in visual pursuit, and improvement in movement speeds as a result of

mental imagery training; this work highlights the breadth of motor performance-related processes that can

be trained without engagement in physical movements.[18,19,56,57] Together with the findings related to motor

adaptation via observation discussed above, the findings of the present report suggest that many features

of motor performance can be improved by training regiments that do not involve movement. This points to

a potential opportunity for the development of motor training or rehabilitation protocols that can be used

when people are unable to physically perform target motor behaviors, perhaps improving performance

beyond what physical practice can do alone.

Finally, our results echo the fact that other types of learning have occur without overt task execution.

As an example, fear associations can be extinguished by instructing participants to imagine a fear-

predicting stimulus even when they are not presented with the stimulus, with this "imagination" protocol

generating neural signatures of the negative prediction errors observed during naturalistic fear

extinction.[60,61] Considering this prior work and the findings presented in this study, it may be that the

generation of predictions for comparison with subsequent sensory observations is sufficient for error-based

learning across motor and non-motor domains alike. In other words, task execution may not always be

required for learning, so long as the predictions and observations needed to compute errors are both

present.

REFERENCES

1. Friston, K. The free-energy principle: a unified brain theory? *Nat Rev Neurosci* **11**, 127–138 (2010).

2. Wolpert, D. M., Miall, R. C. & Kawato, M. Internal models in the cerebellum. *Trends in Cognitive Sciences* **2**, 338–347 (1998).

3. Miall, R. C., Weir, D. J., Wolpert, D. M. & Stein, J. F. Is the Cerebellum a Smith Predictor? *Journal of Motor Behavior* **25**, 203–216 (1993).

4. Popa, L. S. & Ebner, T. J. Cerebellum, Predictions and Errors. *Frontiers in Cellular Neuroscience* **12**, (2019).

5. Wolpert, D. M. & Flanagan, J. R. Motor prediction. *Current Biology* **11**, R729–R732 (2001).

6. Shadmehr, R., Smith, M. A. & Krakauer, J. W. Error Correction, Sensory Prediction, and Adaptation in Motor Control. *Annu. Rev. Neurosci.* **33**, 89–108 (2010).

7. Held, R. & Freedman, S. J. Plasticity in Human Sensorimotor Control. *Science* **142**, 455–462 (1963).

8. Miall, R. C. & Wolpert, D. M. Forward Models for Physiological Motor Control. *Neural Networks* **9**, 1265–1279 (1996).

9. Hewitt, A. L., Popa, L. S., Pasalar, S., Hendrix, C. M. & Ebner, T. J. Representation of limb kinematics in Purkinje cell simple spike discharge is conserved across multiple tasks. *Journal of Neurophysiology* **106**, 2232–2247 (2011).

10. Popa, L. S., Hewitt, A. L. & Ebner, T. J. Predictive and Feedback Performance Errors Are Signaled in the Simple Spike Discharge of Individual Purkinje Cells. *J. Neurosci.* **32**, 15345–15358 (2012).

11. Paz, R., Boraud, T., Natan, C., Bergman, H. & Vaadia, E. Preparatory activity in motor cortex reflects learning of local visuomotor skills. *Nat Neurosci* **6**, 882–890 (2003).

12. Sun, X. *et al.* Cortical preparatory activity indexes learned motor memories. *Nature* 1–6 (2022) doi:10.1038/s41586-021-04329-x.

13. Kilteni, K., Andersson, B. J., Houborg, C. & Ehrsson, H. H. Motor imagery involves predicting the sensory consequences of the imagined movement. *Nat Commun* **9**, 1617 (2018).

14. Voss, M., Ingram, J. N., Haggard, P. & Wolpert, D. M. Sensorimotor attenuation by central motor command signals in the absence of movement. *Nat Neurosci* **9**, 26–27 (2006).

15. Sheahan, H. R., Franklin, D. W. & Wolpert, D. M. Motor Planning, Not Execution, Separates Motor Memories. *Neuron* **92**, 773–779 (2016).

16. Sheahan, H. R., Ingram, J. N., Žalalytė, G. M. & Wolpert, D. M. Imagery of movements immediately following performance allows learning of motor skills that interfere. *Sci Rep* **8**, 14330 (2018).

17. Howard, I. S., Wolpert, D. M. & Franklin, D. W. The Value of the Follow-Through Derives from Motor Learning Depending on Future Actions. *Current Biology* **25**, 397–401 (2015).

18. Ong, N. T. & Hodges, N. J. Absence of after-effects for observers after watching a visuomotor adaptation. *Exp Brain Res* **205**, 325–334 (2010).

19. Lim, S. B., Larssen, B. C. & Hodges, N. J. Manipulating visual–motor experience to probe for observation-induced after-effects in adaptation learning. *Exp Brain Res* **232**, 789–802 (2014).

20. Taylor, J. A. & Ivry, R. B. Flexible Cognitive Strategies during Motor Learning. *PLOS Computational Biology* **7**, e1001096 (2011).

21. Tseng, Y., Diedrichsen, J., Krakauer, J. W., Shadmehr, R. & Bastian, A. J. Sensory Prediction Errors Drive Cerebellum-Dependent Adaptation of Reaching. *Journal of Neurophysiology* **98**, 54–62 (2007).

22. Butcher, P. A. & Taylor, J. A. Decomposition of a sensory prediction error signal for visuomotor adaptation. *Journal of Experimental Psychology: Human Perception and Performance* **44**, 176–194 (2018).

23. Tsay, J. S., Haith, A. M., Ivry, R. B. & Kim, H. E. *Interactions between sensory prediction error and task error during implicit motor learning.* 2021.06.20.449180 https://www.biorxiv.org/content/10.1101/2021.06.20.449180v2 (2021) doi:10.1101/2021.06.20.449180.

24. McDougle, S. D., Ivry, R. B. & Taylor, J. A. Taking Aim at the Cognitive Side of Learning in
Sensorimotor Adaptation Tasks. *Trends in Cognitive Sciences* **20**, 535–544 (2016).
25. Mazzoni, P. & Krakauer, J. W. An Implicit Plan Overrides an Explicit Strategy during Visuomotor
Adaptation. *J. Neurosci.* **26**, 3642–3645 (2006).
26. Morehead, J. R., Taylor, J. A., Parvin, D. E. & Ivry, R. B. Characteristics of Implicit Sensorimotor
Adaptation Revealed by Task-irrelevant Clamped Feedback. *J Cogn Neurosci* **29**, 1061–1074 (2017).
27. Taylor, J. A., Krakauer, J. W. & Ivry, R. B. Explicit and Implicit Contributions to Learning in a
Sensorimotor Adaptation Task | Journal of Neuroscience. *The Journal of Neuroscience* **34**, (2014).
28. Tsay, J. S., Parvin, D. E. & Ivry, R. B. Continuous reports of sensed hand position during
sensorimotor adaptation. *Journal of Neurophysiology* **124**, 1122–1130 (2020).
29. Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R. & Ivry, R. B. Invariant errors reveal
limitations in motor correction rather than constraints on error sensitivity. *Communications Biology* **1**,
1–7 (2018).
30. Avraham, G., Morehead, J. R., Kim, H. E. & Ivry, R. B. Reexposure to a sensorimotor perturbation
produces opposite effects on explicit and implicit learning processes. *PLOS Biology* **19**, e3001147
(2021).
31. Poh, E., Al-Fawakari, N., Tam, R., Taylor, J. A. & McDougle, S. D. Generalization of motor learning in
psychological space. *bioRxiv* 2021.02.09.430542 (2021) doi:10.1101/2021.02.09.430542.
32. Hadjiosif, A. M., Krakauer, J. W. & Haith, A. M. Did We Get Sensorimotor Adaptation Wrong? Implicit
Adaptation as Direct Policy Updating Rather than Forward-Model-Based Learning. *J. Neurosci.* **41**,
2747–2761 (2021).
33. Wei, K. & Körding, K. Relevance of Error: What Drives Motor Adaptation? *Journal of Neurophysiology*
101, 655–664 (2009).
34. Pollmann, S. & Maertens, M. Shift of activity from attention to motor-related brain areas during visual
learning. *Nat Neurosci* **8**, 1494–1496 (2005).
35. Hommel, B. Event Files: Evidence for Automatic Integration of Stimulus-Response Episodes. *Visual*
Cognition **5**, 183–216 (1998).
36. Wong, A. L. & Shelhamer, M. Sensorimotor adaptation error signals are derived from realistic
predictions of movement outcomes. *Journal of Neurophysiology* **105**, 1130–1140 (2011).
37. Franklin, D. W. & Wolpert, D. M. Computational Mechanisms of Sensorimotor Control. *Neuron* **72**,
425–442 (2011).
38. Krakauer, J., W., Hadjiosif, A., Xu, J. & Wong, A., L. Motor Learning. in *Comprehensive Physiology*
(ed. Terjung, R.) (Wiley, 2019). doi:10.1002/cphy.
39. McDougle, S. D. & Taylor, J. A. Dissociable cognitive strategies for sensorimotor learning. *Nat*
Commun **10**, 40 (2019).
40. Hutter, S. A. & Taylor, J. A. Relative sensitivity of explicit reaiming and implicit motor adaptation.
Journal of Neurophysiology **120**, 2640–2648 (2018).
41. Wei, K. & Koerding, K. Uncertainty of feedback and state estimation determines the speed of motor
adaptation. *Front. Comput. Neurosci.* (2010) doi:10.3389/fncom.2010.00011.
42. Lisberger, S. G. The Rules of Cerebellar Learning: Around the Ito Hypothesis. *Neuroscience* **462**,
175–190 (2021).
43. Ito, M. Mechanisms of motor learning in the cerebellum. *Brain Research* **886**, 237–245 (2000).
44. De Zeeuw, C. I., Lisberger, S. G. & Raymond, J. L. Diversity and dynamism in the cerebellum. *Nat*
Neurosci **24**, 160–167 (2021).
45. Mauk, M. D., Medina, J. F., Nores, W. L. & Ohyama, T. Cerebellar function: Coordination, learning or
timing? *Current Biology* **10**, R522–R525 (2000).
46. Kitazawa, S., Kimura, T. & Yin, P.-B. Cerebellar complex spikes encode both destinations and errors
in arm movements. *Nature* **392**, 494–497 (1998).

47. ten Brinke, M. M. *et al.* Evolving Models of Pavlovian Conditioning: Cerebellar Cortical Dynamics in
Awake Behaving Mice. *Cell Reports* **13**, 1977–1988 (2015).
48. Ohmae, S. & Medina, J. F. Climbing fibers encode a temporal-difference prediction error during
cerebellar learning in mice. *Nat Neurosci* **18**, 1798–1803 (2015).
49. Llinás, R. R. Inferior olive oscillation as the temporal basis for motricity and oscillatory reset as the
basis for motor error correction. *Neuroscience* **162**, 797–804 (2009).
50. Romano, V. *et al.* Olivocerebellar control of movement symmetry. *Current Biology* **0**, (2022).
51. Brudner, S. N., Kethidi, N., Graeupner, D., Ivry, R. B. & Taylor, J. A. Delayed feedback during
sensorimotor learning selectively disrupts adaptation but not strategy use. *Journal of
Neurophysiology* **115**, 1499–1511 (2016).
52. Suvrathan, A., Payne, H. L. & Raymond, J. L. Timing Rules for Synaptic Plasticity Matched to
Behavioral Function. *Neuron* **92**, 959–967 (2016).
53. Mattar, A. A. G. & Gribble, P. L. Motor Learning by Observing. *Neuron* **46**, 153–160 (2005).
54. Malfait, N. *et al.* fMRI Activation during Observation of Others' Reach Errors. *Journal of Cognitive
Neuroscience* **22**, 1493–1503 (2010).
55. McGregor, H. R. & Gribble, P. L. Functional connectivity between somatosensory and motor brain
areas predicts individual differences in motor learning by observing. *Journal of Neurophysiology* **118**,
1235–1243 (2017).
56. Gentili, R., Han, C. E., Schweighofer, N. & Papaxanthis, C. Motor Learning Without Doing: Trial-by-
Trial Improvement in Motor Performance During Mental Training. *Journal of Neurophysiology* **104**,
(2010).
57. Cain, M., Botschko, Y. & Joshua, M. Passive Motor Learning: Oculomotor Adaptation in the Absence
of Behavioral Errors. *eNeuro* **8**, (2021).
58. Roberts, T. F., Gobes, S. M. H., Murugan, M., Ölveczky, B. P. & Mooney, R. Motor circuits are
required to encode a sensory model for imitative learning. *Nat Neurosci* **15**, 1454–1459 (2012).
59. Rizzolatti, G., Fogassi, L. & Gallese, V. Neurophysiological mechanisms underlying the
understanding and imitation of action. *Nat Rev Neurosci* **2**, 661–670 (2001).
60. Mendelsohn, A., Pine, A. & Schiller, D. Between Thoughts and Actions: Motivationally Salient Cues
Invigorate Mental Action in the Human Brain. *Neuron* **81**, 207–217 (2014).
61. Reddan, M. C., Wager, T. D. & Schiller, D. Attenuating Neural Threat Expression with Imagination.
Neuron **100**, 994-1005.e4 (2018).
62. Herzfeld, D. J., Vaswani, P. A., Marko, M. K. & Shadmehr, R. A memory of errors in sensorimotor
learning. *Science* **345**, 1349–1353 (2014).
63. Beers, R. J. van. Motor Learning Is Optimally Tuned to the Properties of Motor Noise. *Neuron* **63**,
406–417 (2009).
64. Wu, H. G., Miyamoto, Y. R., Castro, L. N. G., Ölveczky, B. P. & Smith, M. A. Temporal structure of
motor variability is dynamically regulated and predicts motor learning ability. *Nat Neurosci* **17**, 312–
321 (2014).
65. Photon Storm. Phaser (V. 3.24.1). https://phaser.io (2020).
66. Tsay, J. S., Ivry, R. B., Lee, A. & Avraham, G. Moving outside the lab: The viability of conducting
sensorimotor learning studies online. *NBDT* 1–22 (2021) doi:10.51628/001c.26985.
67. Kassambara, A. *rstatix: Pipe-Friendly Framework for Basic Statistical Tests.* (2021).
68. Hothorn, T., Hornik, K., van de Wiel, M. A. & Zeileis, A. Implementing a Class of Permutation Tests:
The coin Package. *Journal of Statistical Software* **28**, 1–23 (2008).
69. Bartoń, K. *MuMIn: Multi-Model Inference.* (2020).
70. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear Mixed
Effects Models. *Journal of Statistical Software* **82**, 1–26 (2017).
71. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models using lme4.
arXiv:1406.5823 [stat] (2014).

72. Jaeger, B. *r2glmm: Computes R Squared for Mixed (Multilevel) Models*. (2017).
73. Lenth, R. V. *emmeans: Estimated Marginal Means, aka Least-Squares Means*. (2021).
74. Torchiano, M. *effsize: Efficient Effect Size Computation*. (2020).
75. Ben-Shachar, M., Lüdecke, D. & Makowski, D. effectsize: Estimation of Effect Size Indices and
Standardized Parameters. *Journal of Open Source Software* **5**, 7 (2020).
76. Bache, S. M., Wickham, H. & RStudio. *magrittr: A Forward-Pipe Operator for R*. (2020).
77. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, 2016).
78. Kassambara, A. *ggpubr: 'ggplot2' Based Publication Ready Plots*. (2020).
79. Lüdecke, D. ggeffects: Tidy Data Frames of Marginal Effects from Regression Models. *JOSS* **3**, 772
(2018).
609

METHODS

Participants. Participants (*n* = 233, aged 18-35, 126 female) recruited from the research participation pools

at Princeton/Yale University and on Prolific provided informed consent, approved by each University's IRB.

Seventy-five participants were excluded (10 from the dataset collected for Supplemental Fig. 2, 13 from the

dataset collected for Supplemental Fig. 3, 26 from the dataset collected for the rotation perturbation

experiments described in Fig. 3, and 26 from the dataset collected for the error-clamp perturbation

experiments described in Fig. 3) for failure to sufficiently recall task instructions, as ascertained by a

questionnaire at the end of the experiment, leaving 158 participants for our analyses. See the *Questionnaire*

section below the *Test phase* sections for more details. We note that all the key results described here (*i.e.,*

statistically significant learning after No-Movement trials) held with or without these exclusions; we opted to

be conservative in our exclusion criteria to limit potential effects of explicit learning. We note that the

samples used for the online experiments described in the text are around twice the size of similar studies

in the literature, providing additional statistical power to compensate for the experiment being conducted

remotely.[62–64]

Task Setup: In Lab. Participants were seated in a chair and made ballistic reaching movements while

grasping the handle of a robotic manipulandum with their dominant hand (Kinarm End-Point). The

manipulandum restricted movements to the horizontal plane. All visual stimuli were projected to the

participant via a horizontal display screen (60 Hz) reflected onto a semi-silvered mirror mounted above the

robotic handle. The mirror occluded vision of the arm, hand, and robotic handle, preventing direct visual

feedback of hand position. Tasks were programmed in Matlab 2019a's Simulink for deployment in Kinarm's

Dexterit-E software (version 3.9). Movement kinematics were recorded at 1 kHz. Each participant viewed

a single target located at either 45°, 135°, 225°, or 315° (with target position counterbalanced across

participants), 8 cm from a central starting location. The target was visible throughout the experiment.

Task Setup: Online. Experiments were conducted remotely using a custom JavaScript web application

based on Phaser 3.24 (download available at [65]), similar to an approach previously described.[66] Each

participant viewed a single target located at either 45°, 135°, 225°, or 315° (with target position

counterbalanced across participants), 250 pixels from a central starting location. The target was visible

throughout the experiment.

Participants used an input device of their choice to control their computer cursor during center-out

movements. One participant reported using a touchscreen device and was excluded from all analyses. The

remaining participants reported using either a trackpad (n = 112), an optical mouse (n = 86), or a trackball

(n = 14). A linear mixed model (LMM) did not show effects of Mouse Type on single-trial learning (STL),

although we observed that participants using a trackpad exhibited longer reaction times than others,

consistent with a previous report.[66]

Mouse position sampling rates depended on the exact hardware that each participant used to

complete the task. Sampling rates were likely affected by features of the specific mouse used, along with

features of the specific computer used, as computers may limit the rate at which the browser samples data

in order to cope with limited processing power. In general, sampling rates were around 60 Hz (median ±

interquartile range across all 213 online participants recruited: 62.46 ± 2.17 Hz) but ranged from 19.23 Hz

to 249.69 Hz. Note that the vast majority of sampling rates were near 60 Hz: Only 5% of sampling rates

were < 41.79 Hz, and only 5% of sampling rates were > 126.65 Hz.

Baseline training phase. For in-lab participants, the robot moved the participant's hand to a central starting

location (depicted by a grey circle) at the middle of the display while hand and cursor feedback were hidden.

They were instructed to hold their hand still in the starting location util the target turned green, at which

point they should make a straight slicing movement through the target. After a 100 ms delay, the robot

moved the hand back to the starting location. Participants completed 5 of these trials with online and

endpoint cursor feedback, followed by 5 trials without visual feedback of the cursor location. Endpoint

feedback was constituted by the cursor remaining at the position where it had passed the target radius for

50 ms. Participants then completed 10 alternating trials on which the target turned green and stayed green

(Execution, 'Go' trials) and on which the target turned magenta 100 ms after turning green, signaling that

participants should withhold their movement (No-Movement, 'Stop' trials). After this baseline phase,

participants were instructed to continue following these instructions for the remainder of the experiment.

Online participants experienced an identical baseline phase, with the exception that they were

instructed to move their mouse into a central starting location on the first trial and subsequently saw their

mouse cursor reappear near the starting location 100 ms after the completion of the reaching movement,

so that participants could quickly return to the start location to initiate the next trial.

Test phase: Rotation and Error-Clamp Experiments (Fig. 2, Supplemental Fig. 2, Supplemental Fig. 3).

During the test phase, 480 (in-lab) or 270 (online) total trials were divided into 3-trial triplets (**Fig. 1C**). The

first and last trials of all triplets were Go trials, and participants received neither online nor endpoint feedback

about cursor location on these trials. The second trial of each triplet was either a Movement or a No-

Movement trial. On Movement trials, participants either received rotated/error-clamped[26] visual feedback

(15° clockwise [-, CW] or counterclockwise [+, CCW], with sign randomized across trials) or veridical/0°

error-clamped visual feedback of their cursor location. On No-Movement (Stop) perturbation trials,

participants viewed a brief animation of the cursor moving straight to the center of the target following a

trajectory deflected by ±15° from the target center. Animation onset latency was set as a running median

of the participant's reaction times on the previous 5 trials, and animation duration was set as a running

median of the participant's movement times on the previous 5 trials. If a participant took longer than 400

ms to execute a movement, 800 ms to initiate the movement, their reach trajectory changed by >10° during

the movement, or the reach terminated ≥ 60° away from the target, they received a warning and a 4s time-

out. If a participant moved their hand (>5 mm in-lab [radius of the starting location]; anything >0 pixels

online) on a No-Movement trial, the trial was immediately aborted, and they received a warning and a 4s

time-out. The Stop manipulation was successful: Across the experiments, participants erroneously moved

on 34.39 ± 20.63% (mean ± standard deviation) of Stop trials, suggesting that, for the most part, they were

consistently planning movements on Stop trials.

For in-lab studies, we used 4 possible triplet perturbation trial types (Movement/No-Movement:

±15°), each of which occurred 40 times throughout each session. For online studies, we used 6 possible

triplet perturbation trial types (Movement/No-Movement: ±15° or 0°), each of which occurred 15 times

throughout each session. Triplets were pseudorandomly presented within each block, with the constraints

that a single rotation (±15° or 0°) could not occur on more than 2 consecutive triplets and that the same

movement condition (i.e., Movement or No-Movement) could not occur on more than 3 consecutive triplets.

Three repetitions of each triplet type occurred in blocks of 18 triplets, and participants received a break after

each of these blocks.

Test phase: Rotation and Error-Clamp Experiments with 0° Perturbations on Movement Trials (Fig. 3).

Experiments were conducted as described above for the other online experiments, with the exception of

the details described in this section. For the experiments described in **Fig. 3A-D**, we used a reduced set of

3 possible triplet Perturbation trial types (No-Movement, 15° clockwise error; No-Movement, 15°

counterclockwise error; Movement, 0° rotation). We maintained an equal number of Movement and No-

Movement triplets throughout the session in order to ensure that participants would reliably respond to the

"Go" cue presented at the start of each trial. So, each No-Movement triplet type occurred 22 times, while

the Movement triplet type occurred 44 times. Triplets were pseudorandomly presented within each block,

with the constraints that a single non-zero rotation (15° clockwise, 15° counterclockwise) could not occur

on more than 2 consecutive triplets.

For the experiments described in **Fig. 3E-H**, we used a set of 4 possible triplet Perturbation trial

types (No-Movement, 15° clockwise error; No-Movement, 15° counterclockwise error; No-Movement, 0°

error; Movement, 0° error-clamp). To maintain an equal number of Movement and No-Movement triplets

throughout the session, each No-Movement triplet type occurred 15 times and the Movement triplet type

occurred 45 times.

Questionnaire: As we could not receive verbal confirmation that participants understood the task

instructions in the online version of the task, we asked subjects to fill out a brief questionnaire to query their

understanding of the task. The questionnaire asked participants to attest whether or not 1) their goal was

to move the real mouse and not the cursor straight through the green targets and whether or not 2) their

goal was to move the white cursor (not the real mouse) straight through the green targets. Participants

could select the options, "True," "False," or "Not Sure." Participants were considered to have understood

the instructions if they answered both questions correctly (i.e., answered "True" to question 1 and "False"

to question 2). The majority of participants answered both questions correctly (138 of 213 participants

[65%]), suggesting that most participants understood the task instructions. Nonetheless, these participants

made up the dataset for all reported analyses for the online experiments, and all other online participants

were excluded from analyses to exclude potential effects of explicit re-aiming.

Data analysis. Data were processed in Python 3.8.5 and Matlab 2018a. Trials with movement were

excluded from analysis 1) if any of the reaches in the triplet were not straight (aspect ratio > participant-

wise mean + 3 * participant-wise standard deviation), 2) if the participant received any warning for failure

to follow task instructions (see *Feedback for failure to follow task instructions*, above), or 3) if the triplet

included a No-Movement No-Go perturbation trial with any detectable mouse movement (>0 pixels online,

>5 mm in lab).

Reach endpoint angle was computed as the angular distance between the center of the target and

the point at which the mouse passed the target's radial distance. Because mouse sampling rates did not

always allow us to measure mouse position at the exact target radius during the online study, we used the

last sample before and the first sample after the mouse passed the target radius to compute an interpolated

mouse position at the target radius, as described in a previous report.[48] We note that analyses comparing

these measures to measurements at the last sample of the reach (even when it was beyond the target) or

the hand angle at peak velocity did not result in substantially different hand angle measurements or

statistical outcomes.

Single-trial learning (STL) was measured as the difference between reach endpoint angle on the

third and first trial of each triplet. For our initial analyses, the sign of STL corresponded to the direction of

the relative change in hand angle, with clockwise changes in hand angle taking a negative sign and

counterclockwise changes in hand angle taking a positive sign. When we collapsed STL data across

rotation directions, we normalized the sign of STL so that changes in hand angle opposite the direction of

the imposed rotation took a positive sign and changes in the direction of the rotation took a negative sign.

Remembered STL was quantified as the difference between reach endpoint angle on the first trial

of one triplet and reach endpoint angle on the first trial of the previous triplet. When remembered STL is

reported as a ratio, this value was computed by dividing remembered STL by the STL attributable to a given

triplet.

Statistics. Statistical tests were conducted in R (v. 4.0.3; packages rstatix[67], coin[68], MuMIn[69],

lmerTest[70], lme4[71], r2glmm[72], emmeans[73], effsize[74], effectsize[75], magrittr[76], ggplot2[77], ggpubr[78],

ggeffects[79]). Data from in-lab experiments were analyzed using a two-way repeated measures ANOVA. If

an ANOVA showed a significant main effect or interaction, post-hoc pairwise tests were performed. When

samples failed to satisfy the normality assumption of the pairwise t-test (assessed via a Shapiro-Wilk test),

we used the more robust paired-samples Wilcoxon signed-rank test. Otherwise, we used the more powerful

paired t-test. Effect sizes for ANOVA main effects/interactions were quantified via generalized η^2 (η^2_G), we

quantified the effect sizes for t-tests using Cohen's d, and we used the Wilcoxon effect size (r) to quantify

effect sizes for signed-rank tests. For these and all subsequent analyses, we corrected for multiple

comparisons using the false-discovery rate approach to maintain family-wise alpha at 0.05.

Data from the experiments conducted online did not satisfy multiple assumptions of the two-way

repeated measures ANOVA (non-existence of extreme outliers, sphericity), so we employed a linear mixed

modeling (LMM; R package lmerTest and lme4) approach for analysis of these data. All LMM's included

fixed effects of perturbation size and movement condition, as well as random effects of subject. Degrees of

freedom were estimated using the Kenward-Rogers approach, and LMM outcomes were reported using

ANOVA-style statistics. Partial R^2 was computed to report effect sizes for the LMM factors (R package

r2glmm). Post-hoc pairwise comparisons were performed between estimated marginal means computed

from the LMM (R package emmeans).

For one-off comparisons between samples or to distributions with 0-mean, we checked samples

for normality. When samples were normally distributed, we ran t-tests and computed Cohen's d to report

effect sizes for statistically significant results. Otherwise, we ran Wilcoxon-signed rank tests and measured

effect sizes using the Wilcoxon effect size (r).