Learning from the path not taken: Sensory prediction errors are sufficient for implicit adaptation of withheld movements

Olivia A. Kim[1], Alexander D. Forrence[2], and Samuel D. McDougle[2]

1. Department of Psychology, Princeton University, Princeton, NJ 08544
2. Department of Psychology, Yale University, New Haven, CT 06511

Correspondence to:
  Olivia Kim
  oakim@princeton.edu

Number of Pages: 31
Number of Figures in Main Text: 3
Number of Supplementary Figures: 5
Number of Supplementary Tables: 3
References: 43
Number of Characters:
  Abstract (1263)
  Highlights (306)
  Introduction (3383), Results (16,126), & Discussion (8,736)
  Online Methods (12,136)
  Main Figure Captions (4,921)
  Supplementary Figure Captions (10,579)

AUTHOR CONTRIBUTIONS
Conceptualization, OK, AF, & SM; Methodology, OK, AF, & SM; Software, OK & AF; Validation, OK & AF; Formal Analysis, OK & AF; Investigation, OK & AF; Resources, OK & SM; Data Curation, OK & AF; Supervision, SM; Project Administration, SM; Funding Acquisition, SM; Writing – original draft, OK; Writing – review & editing, OK, AF, & SM

CONFLICT OF INTEREST
The authors declare no competing financial interests.

ABSTRACT

Current theories of motor control emphasize forward models as a critical component of the brain's motor

execution and learning networks. These internal models are thought to predict the consequences of

movement before sensory feedback from these movements can reach the brain, allowing for smooth,

continuous online motor performance and for the computation of prediction errors that drive implicit motor

learning. Taking this framework to its logical extreme, we tested the hypothesis that movements are not

necessary for the generation of predictions, the computation of prediction errors, and implicit motor

adaptation. Human participants were cued to move a computer mouse to a visually displayed target and

were subsequently cued to withhold those movements on a subset of trials. Visual errors displayed on both

trials with and without movements to the target induced single-trial learning. Furthermore, learning on trials

without movements persisted when accompanying movement trials were never paired with errors and when

movement and sensory feedback trajectories were decoupled. These data provide compelling evidence

supporting an internal model framework in which forward models generate sensory predictions independent

of the generation of movements.

HIGHLIGHTS

• The brain can learn to update movements that are not performed, representing a mechanism for

learning based only on movement planning and sensory expectation.

• Supports a fundamental role for prediction in adaptation.

• Provides further support for the role of forward models in predictive motor control.

KEYWORDS

predictive coding, mental imagery, supervised learning, cerebellum

INTRODUCTION

Forward models are thought to guide our motor behaviors by providing an internal representation

of body-environment interactions. A key feature of these models is their ability to predict the consequences

of movements before relevant sensory feedback can reach the motor controller. As delays between

movements and sensory feedback processing are an inherent byproduct of the time needed for signal

detection and neural propagation, forward models' predictive capacity is essential to motor control.[1] When

sensory feedback indicates that a prediction of a forward model is incorrect, the resulting sensory prediction

error is used to update future predictions and optimize motor commands.[2–5] Such motor adaptation

proceeds implicitly regardless of task goals, reflecting the powerful influence of prediction errors on the

motor system's internal models.[6–8]

The predictive capacity of forward models led us to a strong logical inference: the forward model

should predict and learn from the sensory consequences of an intended movement regardless of whether

that movement occurs. That is, because movements are synchronous with the sensory outcomes that they

produce, motor execution cannot itself be necessary for the generation of predictions by the forward model

or the ensuing computation of sensory prediction error. Indirect evidence for this claim comes from

observations that the brain predicts and accounts for the sensory consequences of movement during motor

imagery in the absence of overt motor behavior.[9–11] This suggests that prediction alone should be sufficient

for the computation of the sensory prediction errors that drive motor adaptation. An internal model for motor

control should thus be effectively trained with only two constituent events – generation of a sensory

prediction and observation of sensory feedback (**Fig. 1a**). Here, we asked if performing a particular

movement is necessary for its adaptation.

To do so, we measured trial-by-trial implicit adaptation during a visuomotor task in which human

participants saw visual feedback while performing – or withholding – hand movements, using a so-called

"Go/No-Go" paradigm. At the onset of each trial, a "Go" cue was presented, instructing participants to make

a movement through a visual target. On some trials, a "No-Go" cue was presented shortly after presenting

the "Go" cue, instructing participants to withhold the movement (**Fig. 1b**). We reasoned that presenting the

"Go" cue on "No-Go" trials would be a sufficient stimulus to induce prediction generation and support

sensory prediction error computation, given that prior work has shown that cued – but ultimately withheld –

movements can still serve as an input to the forward model and support the generation of differential

sensory predictions.[12] To isolate implicit adaptation, we employed a recently-developed approach that

requires participants to aim their movements directly for presented targets and disregard visual

feedback.[8,13–17] We predicted that single-trial motor adaptation would occur following both typical movement

(Go) trials that generated sensory error, as well as trials where movements were withheld (No-Go) but

"simulated" sensory errors were observed. If confirmed, this result would demonstrate that sensory

prediction errors can cause adaptation of selected movements, regardless of whether those movements

are executed.

Figure 1. Schematics showing the proposed learning framework and task design. (**a**) Schematic
showing how the forward model may support motor adaptation in the presence of sensory feedback not
causally related to self-generated movement. (**b**) Schematic of events on trials with visual feedback.
Participants brought their mouse cursor (black pointer) to the starting location to initiate a trial. On
Movement trials (top), the target turned green ('Go Cue'), cueing participants to move the mouse through
the target. Depending on the trial type, participants either observed a white feedback cursor move along
their mouse trajectory, or along a rotated trajectory ('Rotation'). On No-Movement trials (bottom), the target
turned magenta 100 ms after turning green, cueing participants to withhold movement on this trial. After a
delay, an animation showing the feedback cursor moving straight to the target or 15° off-target played
('Animation'). The black mouse pointer is shown in the figure for illustrative purposes but was not visible
during the experiment. (**c**) Schematic showing the trials that constituted a triplet and how single-trial learning
(STL) was computed. Triplets were composed of 2 Movement trials without visual feedback flanking either
a Movement or a No-Movement trial with visual feedback. STL was measured as the difference between
reach angles on the flanking trials.

RESULTS

Simulated and genuine visuomotor rotations cause motor adaptation

We measured implicit motor adaptation in humans (n = 35) performing straight hand movements

during a computerized version of a visuomotor adaptation task (**Fig. 1b**). Trials were organized into triplets,

such that each trial with feedback was flanked by trials without feedback, allowing for a reliable

measurement of single-trial learning (STL) as the difference between the third and first trial of each triplet

(**Fig. 1c**). Trials in which feedback was presented were designated as either "Movement" trials (a "Go"

signal prompted movement) or No-Movement trials (a "No-Go" signal prompted the withholding of

movement). On Movement trials, feedback either involved a visuomotor error (±15° rotation added to the

visual cursor path; + = clockwise) or was veridical (no rotation added). On No-Movement trials, sensory

feedback involved a simulation of the cursor's path, using timing variables based on ongoing participant

performance (**Fig. 2a**, see *Methods*). All flanking trials were "Go" trials and thus required movements. This

straightforward design allowed us to test the hypothesis that forward-model-based motor adaptation does

not require a causal link between movement and sensory feedback (**Fig. 1a**).

Consistent with our predictions, rotated cursor paths on Movement and No-Movement trials both

induced significant adaptation: Movement trajectories on the following trial tended to shift opposite the

direction of the cursor path. This was reflected by STL observed when participants experienced non-zero

rotations of cursor feedback (**Fig. 2b-c**). A linear mixed model (LMM; fixed effects: rotation [15°

counterclockwise {CCW}, 0°, and 15° clockwise {CW}], movement condition [Movement, No-Movement],

rotation x movement condition interaction; random effects: participant) was fit to participants' triplet-wise

STL performance. The LMM revealed significant main effects of rotated cursor feedback (F(2, 1959) =

131.13, p = 2.2 x 10^{-16}, *partial R^2* = 0.12) and movement condition (F(1, 1982) = 4.23, p = 0.04, *partial R^2* =

0.01), as well as a significant interaction (F(2, 1965) = 9.71, p = 6.36 x 10^{-5}, *partial R^2* = 0.002). Post-hoc

pairwise comparisons of the estimated marginal means from the model (**Supplementary Table 1**) support

the claim that rotated feedback induced a statistically significant degree of STL on both Movement (0° vs

15° CW: t(1964) = 8.94, p_{adj} = 3.88 x 10^{-18}, *Cohen's d* = 0.64; 0° vs 15° CCW: t(1971) = 7.143, p_{adj} = 2.89

x 10^{-12}, *Cohen's d* = 0.51) and No-Movement trials (0° vs 15° CW: t(1964) = 4.24, p_{adj} = 3.49 x 10^{-5}, *Cohen's*

d = 0.36; 0° vs 15° CCW: t(1966) = 3.44, p_{adj} = 0.0008, *Cohen's d* = 0.30). Adaptation in the presence of a

rotation was significantly greater in Movement trials than No-Movement trials for CW rotations ($t(1974) =$

4.44, $p_{adj} = 1.67$, *Cohen's d* = 0.35) but not CCW rotations ($t(1971) = 1.76$, $p_{adj} = 0.09$).

Figure 2. Single-trial learning in response to errors on Movement and No-Movement trials in the context of visuomotor rotations and error-clamps. (**a**) Schematic of the relationship between mouse movement and cursor feedback on Movement trials where a rotation was applied to visual feedback. Corresponding mouse (black pointer) and cursor (white circle) movement vectors are color coded. (**b**) An example participant's mean ± SEM changes in reach paths across triplets with a rotation applied. Green shading indicates that data are from triplets in which a perturbation was applied on a Movement trial, while magenta shading indicates that the perturbation was applied on a No-Movement trial. Solid lines indicate that a clockwise (CW) rotation was applied on the relevant perturbation trial, while dashed lines indicate that a counterclockwise (CCW) rotation was applied. (**c**) Boxplot showing change in hand angle across Movement (green) and No-Movement (magenta) triplets in an experiment where feedback on Movement trials was rotated relative to mouse position. In the reference frame used in these experiments, positive changes in hand angle are CW. (**d**) Estimated marginal means ± 95% confidence intervals from the linear mixed models (LMM) fit to each participant's single-trial learning performance summarized in (**c**). Please refer to Supplementary Table 1 for details on all post-hoc comparisons. (**e** - **h**) As in (**a** - **d**), but for an experiment where cursor feedback on Movement trials was error-clamped relative to target location. Please refer to Supplementary Table 2 for details on post-hoc comparisons. Boxplot center: median, notch: 95% confidence interval of the median, box edges: 1st and 3rd quartiles, whiskers: most extreme value within 1.5*interquartile range of the median. Statistical significance (* = p < 0.05; n.s. = p ≥ 0.05) is indicated for selected comparisons. Abbreviations: Δ – change, CW – clockwise, CCW – counterclockwise.

176   **Supplementary Table 1.**

Pairwise post-hoc comparisons between estimated marginal means in Fig. 2d

Group 1		Group 2		Est. Diff.	t	df	FDR-adjusted p	Cohen's d
Trial Type	Rotation	Trial Type	Rotation					
M	0°	M	15° CW	3.77°	8.942	1964	* 3.88 x 10^{-18}	0.65
M	0°	M	15° CCW	2.96°	7.14	1958	* 2.89 x 10^{-12}	0.51
M	15° CW	M	15° CCW	6.73°	16.20	1960	* 1.45 x 10^{-54}	1.15
No-M	0°	No-M	15° CW	2.12°	4.24	1962	* 3.49 x 10^{-5}	0.36
No-M	0°	No-M	15° CCW	1.74°	3.44	1966	* 0.0008	0.30
No-M	15° CW	No-M	15° CCW	3.86°	7.61	1964	* 1.27 x 10^{-13}	0.66
M	0°	No-M	0°	0.40°	0.88	1975	0.38	--
M	15° CW	No-M	15° CW	2.06°	4.44	1974	* 1.67 x 10^{-5}	0.35
M	15° CCW	No-M	15° CCW	0.82°	1.76	1971	0.09	--

Note. Abbreviations: M – Movement; No-M – No-Movement; Est. Diff. – Estimated Differences in degrees. Note that degrees of freedom pertain to the inputs to the LMM and are estimated using the Kenward-Rogers approach. Statistically significant p-values are indicated with asterisks. * $p < 0.05$.

To address whether the STL measured represented genuine implicit learning, we checked whether
adaptation persisted beyond the trial after an error was experienced. We examined participants' hand
angles on the second trial after a perturbation relative to the pre-perturbation baseline trial (i.e., hand angle
on trial 1 of triplet *t + 1* relative to hand angle on trial 1 of triplet *t,* henceforth referred to as remembered
STL, **Supplementary Fig. 1a**). Because visual feedback was withheld on both trial types, this approach
provides a relatively pure measure of persistent learning in the absence of error-driven changes in
performance. Hand angle appeared to remain adapted in the direction opposite the rotation on trials with
nonzero perturbations regardless of movement condition (**Supplementary Fig. 1b-c**), suggesting that
genuine implicit learning was observed in response to errors under both movement conditions. Participant-
wise GLMs fit to individual subjects' remembered STL measures (rotation, movement condition, and their
interaction included as predictors) support this interpretation: the distribution of estimated coefficients for

rotation as a predictor was significantly different from zero (Wilcoxon's signed-rank test: $V = 25$, $p_{adj} = 1.58$

x 10^{-7}, $r = -0.80$), whereas estimated movement coefficients ($V = 228$, $p_{adj} = 0.24$) and movement/rotation

interaction coefficients ($V = 365$, $p_{adj} = 0.42$) did not differ significantly from 0 (**Supplementary Fig. 1d**).

Importantly, additional participant-wise GLMs fit to the adaptation observed the trial after error presentation

(i.e., the STL measurements shown in **Fig. 2**) with the perturbations applied on that and the preceding

triplet used as predictors showed that STL was not attributable to errors observed on preceding Movement

triplets (Wilcoxon's signed-rank test: $V = 283$, $p_{adj} = 0.70$; **Supplementary Fig. 2a**). Thus, while STL

persisted at least one trial beyond the triplet in which an error was observed, STL observed on No-

Movement trials was not attributable to residual adaptation from preceding triplets with errors on Movement

trials.

Supplementary Figure 1. Evidence that STL memories are maintained. (**a**) Schematic of the trial order
in two consecutive triplets, illustrating that the 2 trials after each Movement/No-Movement trial are
performed without visual feedback. (**b**) Change in hand angle relative to the first trial of the triplet in which
the perturbation being considered occurs for Movement (green) and No-Movement (magenta) trials
(remembered STL), for participants in an experiment where feedback was rotated relative to actual mouse
position on Movement trials. Data are normalized to the change in hand angle observed on this trial on
triplets with 0° perturbations, and data are shown as mean ± SEM. (**c**) Boxplots showing the distribution of
participants' median persistent STL (change in hand angle relative to the pre-perturbation trial) two trials
after the perturbation was observed. (**d**) Estimated coefficients from GLMs fitted to each participants'
remembered STL performance with rotation applied, movement condition, and rotation:movement condition
interaction provided as predictors. Wilcoxon's signed-rank tests showed statistically significant differences
between rotation coefficients and 0 ($V = 25$, $p_{adj} = 1.58 \times 10^{-7}$, $r = -0.80$), but not between movement
coefficients ($V = 228$, $p_{adj} = 0.24$) or rotation:movement interaction coefficients ($V = 365$, $p_{adj} = 0.42$) and 0.
(**e-f**) As in (**b-d**), but showing data collected in an experiment where participants saw error-clamped
feedback on Movement trials. (**g**) Wilcoxon's signed-rank tests showed statistically significant differences
between rotation coefficients and 0 ($V = 31$, $p_{adj} = 1.04 \times 10^{-7}$, $r = -0.80$) and between rotation:movement

217 interaction coefficients and 0 ($V = 490$, $p_{adj} = 0.02$, $r = 0.41$), but not between movement coefficients ($V =$
218 428, $p_{adj} = 0.25$) and 0. Boxplot center: median, notch: 95% confidence interval of the median, box edges:
219 1st and 3rd quartiles, whiskers: most extreme value within 1.5 * interquartile range of the median. Notches
220 have been omitted for panels **d** and **g** because 95% confidence intervals exceeded the interquartile range.
221 * indicates p < 0.05.

Supplementary Figure 2. Coefficients from participant-wise GLMs with the perturbations applied on current and preceding Movement and No-Movement trials used as predictors. (**a**) Estimated coefficients from GLMs fit to each participant's single-trial learning (STL) measurements, using the perturbation applied on the current and preceding triplets as predictors. Participants were exposed to rotated cursor feedback on Movement trials. In the x-axis labels, t indicates that the perturbation was applied on the current triplet, while t-1 indicates that the perturbation was applied on the preceding triplet. The distribution of coefficient estimates for Movement perturbations applied on the current triplet (Movement, t) was significantly different from 0 (Wilcoxon's signed-rank test, $V = 16$, $p_{adj} = 7.87 \times 10^{-8}$, $r =$ -0.83). The same was true for the coefficient estimates for No-Movement perturbations applied on the current triplet (No-Movement, t; $V = 74$, $p_{adj} = 9.50 \times 10^{-5}$, $r = -0.67$). No-Movement perturbations applied on the preceding triplet (No-Movement, $t - 1$) also differed from 0 ($V = 463$, $p_{adj} = 0.04$, $r = 0.40$). Notably, No-Movement, t-1 coefficients tended to be positive (median: 0.05), suggesting that any effects of perturbations on previous No-Movement trials are not examples of residual adaptation from those trials, as any such effects would take on a negative coefficient. No other coefficient estimates differed significantly from 0 (Movement, t-1: $V = 283$, $p_{adj} = 0.70$; No-Movement, t : Movement, t-1: $V = 265$, $p_{adj} = 0.80$; No-Movement, t : No-Movement, t-1: $V = 246$, $p_{adj} = 0.70$; Movement, t : Movement, $t - 1$: $V = 254$, $p_{adj} = 0.70$; Movement, t : No-Movement, $t - 1$: $V = 235$, $p_{adj} = 0.70$). (**b**) As in (**a**), except that participants were exposed to error-clamped cursor feedback on Movement trials. The distribution of coefficient estimates for Movement perturbations applied on the current triplet (Movement, t) was significantly different from 0 (Wilcoxon signed-rank test, $V = 92$, $p_{adj} = 0.0002$, $r = -0.64$). The same was true for the coefficient estimates for No-Movement perturbations applied on the current triplet (No-Movement, t; $V = 149$, $p_{adj} = 0.008$, $r = -0.47$). No-Movement perturbations applied on the preceding triplet (No-Movement, $t - 1$) also differed from 0 ($V = 576$, $p_{adj} =$ 0.002, $r = 0.56$). No other coefficient estimates differed significantly from 0 (Movement, t-1: $V = 300$, $p_{adj} =$ 0.71; No-Movement, t : Movement, t-1: $V = 248$, $p_{adj} = 0.87$; No-Movement, t : No-Movement, t-1: $V = 178$, $p_{adj} = 0.96$; Movement, t : Movement, $t - 1$: $V = 240$, $p_{adj} = 0.87$; Movement, t : No-Movement, $t - 1$: $V =$

250   243, $p_{adj} = 0.32$). Boxplot center: median, notches: 95% confidence interval of the median, edges: 1st and
251   3rd quartile, whiskers: most extreme datapoints within 1.5 IQR of the median. * indicates $p_{adj} < 0.05$
252

To assess the potential similarity of mechanisms underlying adaptation after errors on Movement

and No-Movement trials, we compared STL amplitude under each condition, reasoning that there should

be a relationship if STL is supported by the same mechanisms on Movement and No-Movement trials.

When we considered instances of STL in the direction that would compensate for the observed error

(update opposite rotation, **Supplementary Fig. 3a**), within-subject changes in hand angle were correlated

between Movement and No-Movement trials (*Pearson's r* = 0.45, *p* = 0.007; **Supplementary Fig. 3b**).

Conversely, changes in hand angle in the direction that would exacerbate the observed error (update in

direction of rotation, **Supplementary Fig. 3a**) were uncorrelated between Movement and No-Movement

trials (*Pearson's r* = 0.03, *p* = 0.86, **Supplementary Fig. 3c**). Together, these observations suggest that

the same learning process may underlie adaptive single-trial learning events in response to errors on both

kinds of trials, while maladaptive changes in hand-angle may be attributable to potential sources of random

noise. In spite of the correlation between the amplitudes of adaptive changes in hand angle on Movement

and No-Movement trials, the overall amplitude of the changes in hand angle opposite the direction of the

rotation were nonetheless smaller on No-Movement trials than on Movement trials (Wilcoxon's signed-rank

test: *W* = 88, *p* = 0.0002, *rank-biserial correlation* = 0.14), indicating that errors on No-Movement trials may

not drive adaptation as strongly as errors on Movement trials. (We return to this issue in the *Discussion*.)

Supplementary Figure 3. Correlations between STL magnitude after Movement and No-Movement trials. (**a**) Schematic illustrating the implications of changes in hand angle (updates) in the same direction as or opposite direction to the perturbation applied. Under "normal" circumstances when we encounter a consistent perturbation, adaptation that causes the hand to move in the direction opposite the perturbation will counter that perturbation and tend to minimize future errors. (**b**) Scatter plot showing the relationship between individual subjects' STL amplitude in the direction opposite the rotation on Movement and No-Movement trials. Data were collected from an experiment where participants saw rotated feedback on Movement trials. (**c**) As in (**b**), but for STL amplitude in the same direction as the rotation applied. (**d-e**) As in (**b** – **c**), but for participants who received error-clamped feedback on Movement trials.

Motor adaptation in response to errors during error clamp trials

We note that rotated visual feedback on Movement trials was sensitive to people's actual reaching directions (i.e., the rotation was simply added to their natural reach direction, as is typical in visuomotor rotation tasks). Thus, it is possible that introducing these directional contingencies affected how people behaved in response to perceived errors. To rule this out, we recruited participants (n = 37) for a second experiment where the visual cursor moved in a fixed path ("error-clamped" feedback[8]) in one of three directions (0° or 15° CW/CCW) on trials with feedback (**Fig. 2e**). In line with the effects of rotated cursor feedback, a LMM revealed main effects of error-clamped feedback (F(2, 1827.8) = 79.46, p = 2.2 x 10^{-16}, *partial R^2* = 0.08) and an interaction between rotation and movement condition (F(2, 1831.8) = 8.45, p =

0.0002, *partial* R^2 = 0.0003), although there was no main effect of movement condition (F(1, 1844.1) = 0.60,

p = 0.44; **Fig. 2h**). Post-hoc comparisons of the model's estimated marginal means (**Supplementary Table**

2) revealed significant STL in response to non-zero error-clamped feedback on both Movement (0° vs 15°

CW: t(1827) = 7.55, p_{adj} = 3.08 x 10^{-13}, *Cohen's d* = 0.56; 0° vs 15° CCW: t(1828) = 5.57, p_{adj} = 8.84 x 10^{-}

^{8}, *Cohen's d* = 0.41) and No-Movement trials (0° vs 15° CW: t(1830) = 3.21, p_{adj} = 0.002, *Cohen's d* = 0.29;

0° vs 15° CCW: t(1832) = 2.25, p_{adj} = 0.03, *Cohen's d* = 0.20). Adaptation in the presence of a 15° error-

clamp was significantly greater on Movement trials than No-Movement trials for CW (t(1846) = 3.50, p_{adj} =

0.0009, *Cohen's d* = 0.29) and CCW clamps (t(1846) = 2.29, p_{adj} = 0.03, *Cohen's d* = 0.19).

Supplementary Table 2.

Pairwise post-hoc comparisons between estimated marginal means in Fig. 2h

Group 1		Group 2		Est. Diff.	t	df	FDR-adjusted p	Cohen's d
Trial Type	Rotation	Trial Type	Rotation					
M	0°	M	15° CW	3.54°	7.55	1827	* 3.08 x 10^{-13}	0.56
M	0°	M	15° CCW	2.57°	5.57	1828	* 8.84 x 10^{-8}	0.41
M	15° CW	M	15° CCW	6.11°	13.14	1828	* 8.51 x 10^{-37}	0.97
No-M	0°	No-M	15° CW	1.81°	3.21	1830	* 0.002	0.29
No-M	0°	No-M	15° CCW	1.29°	2.25	1832	* 0.03	0.20
No-M	15° CW	No-M	15° CCW	3.10°	5.46	1833	* 1.24 x 10^{-7}	0.49
M	0°	No-M	0°	0.08°	0.16	1846	0.87	--
M	15° CW	No-M	15° CW	1.81°	3.49	1846	* 0.0009	0.29
M	15° CCW	No-M	15° CCW	1.19°	2.29	1846	* 0.03	0.19

Note. Abbreviations: M – Movement; No-M – No-Movement; Est. Diff. – Estimated Differences in degrees. Note that degrees of freedom pertain to the inputs to the model and are estimated using the Kenward-Rogers approach. Statistically significant p-values are indicated with asterisks. * p < 0.05.

Like participants who observed rotated cursor feedback on Movement trials, participants who

observed error-clamped cursor feedback on Movement trials showed persistent STL beyond the triplet in

which the error was observed (remembered STL, **Supplementary Fig. 1e-f**), and participant-wise GLMs'

fitted rotation coefficients differed significantly from 0 (**Supplementary Fig. 1g**; Wilcoxon's signed-rank

test; $V = 31$, $p_{adj} = 1.04 \times 10^{-7}$, $r = -0.80$). Additionally, STL for error-clamp participants was not attributable

to errors observed on preceding Movement trials (**Supplementary Fig. 2b**). Unlike the rotation study

participants, error-clamp study participants' rotation:movement interaction coefficients also differed from 0

($V = 490$, $p_{adj} = 0.02$, $r = 0.41$), suggesting that they showed less remembered STL after No-Movement

trials than Movement trials. Nonetheless, these observations indicate that changing the nature of feedback

on Movement trials did not eliminate the capacity of errors on No-Movement trials to induce genuine

learning.

Also echoing the rotation study, the amplitude of error-clamp study participants' STL in the direction

counter to the error observed was correlated between Movement and No-Movement conditions (*Pearson's*

$r = 0.34$, $p_{adj} = 0.04$), while the amplitude of STL events that proceeded in the same direction as the

presented error was not correlated between Movement and No-Movement conditions ($r = 0.28$, $p_{adj} = 0.11$).

Thus, these data generally replicate the rotation study, strengthening the claim that the same learning

process may underlie adaptive single-trial learning events in response to errors on both Movement and No-

Movement trials.

Adaptation on No-Movement trials does not depend on within-session adaptation on Movement trials

In two further experiments, we asked if adaptation to errors in the No-Movement condition was

contingent on sharing a context with the Movement condition. That is, does learning in the No-Movement

condition only occur if it happens alongside similar learning in the Movement condition? If so, this would

suggest that adaptive responses in the No-Movement condition may reflect a "cueing" effect, whereby an

adapted sensorimotor map is cued by the visual error and then retrieved on the subsequent trial(s). While

our retention (**Supplementary Figs. 1-2**) and correlation (**Supplementary Fig. 3**) results argue against

this interpretation, we directly tested this in another pair of experiments. Here, we only included 0° rotated

(**Fig. 3a**) or clamped (**Fig. 3d**) error feedback on Movement trials, and the normal mix of 0°, 15° CW/CCW

errors on the No-Movement trials. The key results were replicated – learning was preserved in the No-

Movement condition even when error feedback had never been associated with executed movements (**Fig.**

 333   **3b – c**, LMM: $F(557)$ 23.01, p = 2.07 x 10⁻⁶, *partial R^2* = 0.04, **e – f**, $F(1456)$ = 11.63, p = 9.76 x 10⁻⁶, *partial*

R^2 = 0.02). Post-hoc pairwise comparisons showed that adaptation was significantly different between

triplets with clockwise and counterclockwise errors for both the rotation ($t(557)$ = 4.80, p = 2.07 x 10⁻⁶,

Cohen's d = 0.4) and error-clamp experiments ($t(1453)$ = 4.03, p = 1.16 x 10⁻⁵, *Cohen's d* = 0.4) – a hallmark

of implicit motor adaptation (see **Supplementary Table 3** for all post-hoc tests and results).

Figure 3. Single-trial learning on No-Movement trials when feedback on Movement trials underwent
0° rotation or clamp. (**a**) Schematic of visual feedback on Movement trials when 0° rotation was applied.
(**b**) An example participant's mean ± SEM changes in reach paths across triplets with a rotation applied.
Green shading indicates that data are from triplets in which a perturbation was applied on a Movement trial,
while magenta shading indicates that the perturbation was applied on a No-Movement trial. Solid lines
indicate that a clockwise (CW) rotation was applied on the relevant perturbation trial, while dashed lines
indicate that a counterclockwise (CCW) rotation was applied. (**c**) Single-trial learning in response to different
directions of simulated errors on No-Movement trials (magenta) and no rotation on Movement trials (green).
(**d**) Estimated marginal means ± 95% confidence intervals from the linear mixed models (LMM) fit to each
participant's single-trial learning performance summarized in (**c**). (**e-h**) As (**a-f**), but for participants in an
experiment with 0° error-clamped feedback on Movement trials. Note that data in (**g**) from the 0° rotation
applied condition are collapsed across Movement and No-Movement trials. Asterisks indicate statistically
significant differences between post-hoc comparisons of estimated marginal means. Please refer to
Supplementary Table 3 for detailed statistical results. Boxplot centers: median, notches: 95% confidence

interval of the median, box edges: 1st and 3rd quartiles, whiskers: most extreme values within 1.5*IQR of
the median.

Supplementary Table 3.

Pairwise post-hoc comparisons between estimated marginal means in Fig. 3

Group 1		Group 2		Est. Diff.	t	df	FDR-adjusted p	Cohen's d
Trial Type	Rotation	Trial Type	Rotation					
0° Rotation Applied on Movement Trials								
No-M	15° CCW	No-M	15° CW	2.45°	4.80	557	* 2.07 x 10^{-6}	0.40
0° Error-Clamped Feedback on Movement Trials								
No-M	15° CCW	No-M	15° CW	2.47°	4.03	1453	* 0.001	0.40
No-M	15° CW	M + No-M	0°	2.21°	4.64	1457	* 1.16 x 10^{-5}	0.36
No-M	15° CCW	M + No-M	0°	0.26°	0.56	1456	0.57	--

Note. Abbreviations: M – Movement; No-M – No-Movement; Est. Diff. – Estimated Differences in

degrees. Note that degrees of freedom pertain to the inputs to the model and are estimated using the

Kenward-Rogers approach. Statistically significant p-values are indicated with asterisks. * p < 0.05.

Although these results demonstrated that sharing a context with movement-accompanied

adaptation was not a necessary condition for No-Movement adaptation, we did observe attenuated learning

on No-Movement Trials: Adaptation in response to errors on both Movement and No-Movement trials was

greater in experiments in which non-zero rotations and clamps were applied on Movement trials

(**Supplementary Fig. 4a**, two-sample t-test: *t*(85) = 5.94, *p* = 6.15 x 10^{-8}, *Cohen's d* = 1.15). Crucially, STL

across all four of our experiments was within the range of learning rates observed in previous studies with

identical error sizes (**Supplementary Fig. 5**).

Prior work has indicated that subjects' within-session histories of error affect the amplitude of motor

adaptation, offering a potential explanation of the observed differences.[18–20] Here, differences in the

experimental designs mandated by the need to counterbalance the number of Movement and No-

Movement trials within a session may have affected the prevalence of different kinds of errors and error

transitions in the experiments described in **Figs. 2**-**3**, potentially influencing the amplitude of STL observed

within each experiment.

Following the analyses in (ref. [20]), we first investigated whether observation of two successive

errors in opposite directions decreased sensitivity to error and may have accounted for the differences in

overall STL amplitude between the experiments contributing to **Figs. 2**-**3**. To our surprise, in each

experiment, we observed an increase in STL after a change in the direction of the perturbation relative to

STL after maintaining the direction of the perturbation (**Supplementary Fig. 4b**): Across all experiments,

when the direction of the observed error was maintained between triplets, STL was significantly smaller

than STL when an error abruptly emerged (i.e., when a non-zero error occurred after a triplet with a 0°

perturbation trial; $t(113) = 2.18$, $p_{adj} = 0.047$, *Cohen's d* = 0.29), or when the error's direction changed ($t(109)$

$= 2.38$, $p_{adj} = 0.047$, *Cohen's d* = 0.32). These data suggest that, when participants are exclusively exposed

to triplets with randomly varying error, inconsistent errors may increase the rate of implicit adaptation, in

contrast to what was previously reported in a blocked learning design.[20] We note that participants in

experiments where only 0° perturbations were applied on Movement trials were exposed to significantly

fewer instances of inconsistent errors than participants who experienced 0° and 15° perturbations on

Movement trials (**Supplementary Fig. 4d**, two-sample Kolmogorov-Smirnov tests; Rotation 0°/15° vs

Rotation 0° only on Movement Trials: $D = 1$, $p_{adj} = 1.29 \times 10^{-12}$; Error-clamp 0°, 15° vs Error-clamp 0° only

on Movement Trials: $D = 1$, $p_{adj} = 1.66 \times 10^{-13}$). Taken together, these observations suggest that overall

STL was smaller in experiments where participants only saw 0° perturbations on Movement trials because

of decreases in error sensitivity arising from a paucity of inconsistent error transitions.

In addition to effects of error direction transitions, prior studies have also reported putative effects

of reward on implicit adaptation.[22–24] Specifically, when unexpected feedback deviations still result in task

success (i.e., the cursor movement terminates within the target), implicit adaptation is attenuated. When

we retrospectively considered the proportion of trials where the visually displayed cursor landed completely

on target and may have satisfied the implicit task goal of "hitting" the target, we found that participants who

experienced exclusively 0° error-clamped feedback on Movement trials saw the cursor hit the target >25%

more than participants in the other experiments (**Supplementary Fig. 4e**). This greater rate of apparent

task success may have established a context that suppressed adaptation in a manner consistent with prior

work.[22,24]

Together, these data suggest that the order and content of trials and the nature of the stimuli

employed in the rotation and Error Clamp experiments without non-zero perturbations on Movement trials

effectively reduced learning rates in these experiments and may account for the lower observed levels of

adaptation between the two sets of studies. Overall, these experiments support our hypothesis that, in the

right circumstances, motor adaptation can proceed without movement execution.

Supplementary Figure 4. Effects of changes in error sign on learning rate and graphics showing
details of experiments' trial composition. (**a**) Boxplots showing STL observed across Movement and
No-Movement trials in the experiments shown in **Fig. 2** (0°, ±15° on Go) and the experiments shown in **Fig.**
3 (0° on Go). A two-sample t-test revealed a significant difference between STL amplitude in each kind of
experiment (t(85) = 5.94, p = 6.15 x 10^{-8}, *Cohen's d* = 1.15). (**b**) Boxplots showing STL observed on triplets
where the perturbation direction changed relative to the previous triplet (open boxes), triplets where a non-
zero perturbation occurred after a 0° perturbation (filled boxes), and triplets where the perturbation direction
was the same as the previous triplet (hashed boxes). (**c**) STL across all experiments in the present study,
divided by the type of error transition observed. Data are shown as mean ± SEM. Paired t-tests between
STL amplitudes between error transition conditions showed significant differences between triplets where
error direction remained the same and triplets where errors appeared following a 0°-perturbation trial (t(113)
= 2.18, p_{adj} =0.047 , *Cohen's d* = 0.29) and triplets where the direction of the perturbation changed (t(109)
= 2.38, p_{adj} = 0.047, *Cohen's d* = 0.32). There was no significant difference between triplets where error

appeared following a 0°-perturbation triplet and triplets where the direction of the perturbation changed
(t(117) = 0.43, p_{adj} = 0.67). (**d**) Cumulative distribution plots showing the proportion of participants with a
certain proportion of trials or lower on which an error appeared or the sign of the perturbation changed
between triplets. Two-sample Kolmogorov-Smirnov tests revealed that the proportion of trials where errors
appeared or the perturbation sign changed was significantly lower for participants who received 0°-error-
clamp perturbations on all Movement trials than for participants who received 0°-rotation perturbations on
all Movement trials (D = 1, p_{adj} = 2.21 x 10^{-11}), who received 0°/±15° rotation perturbations on all Movement
trials (D = 1, p_{adj} = 2.31 x 10^{-13}), and who received 0°/±15° error-clamp perturbations on all Movement trials
(D = 1, p_{adj} = 1.66 x 10^{-13}). The proportion of trials where errors appeared/changed for participants who
received 0°-rotation perturbations on all Movement trials was also significantly lower that that observed for
participants who received 0°/±15° rotation perturbations on all Movement trials (D = 1, p_{adj} = 1.29 x 10^{-12}),
and who received 0°/±15° error-clamp perturbations on all Movement trials (D = 1, p_{adj} = 2.72 x 10^{-14}). The
proportion of trials where errors appeared/changed directions did not differ between participants who
received 0°/±15° rotation or error-clamp perturbations on all Movement trials (D = 0.21, p_{adj} = 0.42). (**e**) Bar
plots showing the proportion of feedback trials on which the cursor landed completely on the target. The
number of trials on which the cursor hit the target was inflated in the Clamp: 0° study, because 1) all
Movement trials were 0° error-clamped trials, 2) there were as many Movement trials as No-Movement
trials, and 3) there were also No-Movement trials showing 0° simulated errors. The number of trials on
which the cursor hit the target was deflated in the Rotation: 0° study because there was no No-Movement
trials showing 0° simulated errors (i.e., there was no subset of trials on which the cursor was guaranteed to
hit the target for these participants). Data are shown as mean ± SEM when values varied between
participants (rotation experiments) and as mean only when values did not vary (clamp experiments).
Boxplot centers: median, notches: 95% confidence interval of the median, box edges: 1st and 3rd quartiles,
whiskers: most extreme values within 1.5*IQR of the median. * indicates p_{adj} < 0.05.
446
447

Supplementary Figure 5. Learning rates observed in the literature and the current study. Learning
rates observed in previous studies are shown at left, and learning rates observed in each experiment in the
current report are shown at the right. Data are shown as mean ± SEM, and are shown for rotations/error
clamp manipulations of 15°, with the exception of Wei & Kording E2, where an 11° perturbation was applied.
Papers referred to and their corresponding reference numbers: Kim et al., 2018[14]; Morehead et al., 2017[8];
Wei & Kording, 2009[21]. "E1" in the current report refers to conditions where participants saw 0-15°
perturbations on Movement trials, and "E2" in the current report refers to conditions where participants saw
only 0° perturbations on Movement trials. Abbreviations: E, experiment.
457

DISCUSSION

The results presented above demonstrate that movements can be implicitly refined even when they

are not performed. Participants who were cued to perform a movement towards a target but suppressed

that movement after observation of a No-Go cue showed consistent, robust single-trial learning in response

to an animation of erroneous cursor movement (**Fig. 2b-d**, **f-h; Fig. 3b-d, f-h**). This learning carried the key

signatures of implicit adaptation of an internal model. Our findings support the fundamental assumptions of

predictive processing frameworks of motor adaptation, where precise sensory predictions are generated

from a movement intent (or "plan", "goal") and compared against sensory observations to induce error-

based learning.[12,25]

We argue that we have measured learning via an implicit process, and, by extension, that implicit

adaptation of specific movements can proceed even when those movements are not performed. While

visuomotor learning tasks do sometimes recruit cognitive learning strategies (e.g., the deliberate "re-aiming"

of movements), multiple factors indicate that our studies measured implicit adaptation.[7,26,27] First, in line

with previous work isolating the implicit process, participants were instructed to ignore the visually displayed

cursor and try to contact the target on every trial, a simple technique which has proven consistently effective

at eliminating the explicit re-aiming of movements.[8,13,14,28] Second, in addition to instructing participants in

a manner that should have suppressed recruitment of explicit processes, the randomization of both the

presence and direction of visual errors discourages explicit learning, reducing any motivation that

participants may have had to apply ineffective re-aiming strategies (see [29]). Third, we excluded data from

any participants who did not appear to understand or recall our instruction to always aim at the target (see

Methods), decreasing the likelihood that strategic re-aiming contaminated the analyzed data sets (however,

we note that our key results did not change if these participants were included). Fourth, adaptation persisted

into subsequent no-feedback trials (**Supplementary Fig. 1b-g**), a finding that is consistent with lingering

implicit motor learning; it is unlikely that strategies would be maintained across trials where no feedback is

present. Lastly, the magnitude of single-trial learning we observed was in line with multiple previous studies

designed to measure implicit trial-by-trial motor adaptation (**Supplementary Fig. 5**).[8,14,21] Our data thus

provide striking evidence that internal forward models predict the state of the world following action

selection, instead of predicting the sensory consequences of movement *per se*, and that this prediction is

then compared with observed sensory events to compute a sensory-prediction error to drive learning.

We also note that the adaptation effects observed in the No-Movement conditions were not

attributable to a recall process driven by movement-based learning. That is, No-Movement adaptation was

not related to errors on preceding Movement triplets (**Supplementary Fig. 2**) and proceeded independently

of visual errors being present on Movement trials (**Fig. 3**). While overall levels of adaptation were smaller

during the two experiments with no perturbations on Movement trials, this reduced level of adaptation was

likely attributable to features of the experimental design: The decreased likelihood of errors and changes

in trial-by-trial error variance in these experiments may have reduced learning rates (**Supplementary Fig.**

4). Additionally, an increased likelihood of task success (hitting the target) during the error-clamp study with

no perturbations on Movement trials may have further suppressed learning rates.[22,24]

Our observation that changes in the direction of the perturbation enhanced the degree of single-

trial adaptation (**Supplementary Fig. 4b**) appears to conflict with a recently published report suggesting

that changes in the direction of error decrease error sensitivity and accompanying learning rate.[20] However,

several differences between this study and the present study may account for these differences. For

instance, the experiments described in [20] varied perturbation amplitude, but held perturbation sign

consistent, whereas we varied perturbation sign. Additionally, we observed effects of changes in the sign

of the perturbation applied on STL, while [20] reported effects of changes in the sign of the visual error

observed on cumulative learning across sessions. Finally, in our study, changes in errors took place across

triplets (sets of 3 trials); in contrast, in the experiments in [20] errors may have changed between single trials.

It may be that a different mechanism for tracking and adjusting error sensitivity in response to environmental

variability is recruited when visual feedback is not present on all trials, or when the sign in addition to the

magnitude of the perturbation varies.

While our results demonstrate that overt movement is not necessary for motor adaptation, we are

not able to discern whether the generation of a descending motor command is a necessary component for

updating the forward model. To reliably elicit prediction generation from the forward model, we employed a

motor inhibition paradigm, briefly cueing participants to reach before presenting them with an abrupt stop

signal. This strategy is likely to have recruited mechanisms of motor inhibition beyond the primary motor

cortex, leaving open the possibility that a descending motor command and not an upstream motor plan *per*

se provide the necessary input for forward internal model adaptation.[30–32] Future experimentation

quantifying neural (EEG) or physiological (EMG) signals during this task and measuring their relationship

with adaptation on No-Movement trials will be crucial for addressing this point. Inspired by work in non-

human primates, we speculate that shared motor cortical preparatory states in the Movement and No-

Movement conditions may produce comparable inputs to downstream internal models.[33]

In addition to suggesting that motor planning is sufficient for motor adaptation, our data also

demonstrate that errors paired with movements support a greater degree of learning than errors without

movement, as participants showed significantly stronger STL over triplets with Movement trials than No-

Movement trials. This may suggest that movement provides additional training input to the forward model

beyond what erroneous visual feedback provides alone. Alternately (though not mutually exclusive), it may

be that the absence of movement-related proprioceptive feedback dampens the sensory prediction error

signals that drive adaptation, perhaps by failing to counterbalance inhibitory signals predicting

proprioceptive feedback.[1,34] A simpler set of explanations are that participants simply did not recognize the

Go cue to set a movement goal and engage the forward model on a subset of No-Movement trials, or that

the timing of the simulated error was too variable, relative to the internal prediction, to elicit maximal

adaptation. Further experimentation will be required to address these possibilities. Nonetheless, our data

demonstrate the significant influence of the brain's prediction signals on learning – even without the ability

to directly attribute sensory feedback to an actual movement, the brain's prediction of a planned

movement's sensory consequences support the error computations that adapt future behavior.

We note that other forms of learning also appear to occur without overt task execution. First, motor

learning can occur through observation of another individual learning the same task or observation of target

movement without active tracking – the degree to which such learning reflects action selection processes

or implicit motor adaptation (or both) is unclear, though our current results may support observation-based

implicit motor adaptation.[35,36] Outside of motor learning, fear associations can be extinguished by instructing

participants to imagine a fear-predicting stimulus even when they are not presented with the stimulus, with

this "imagination" protocol generating neural signatures of the negative prediction errors observed during

naturalistic fear extinction.[37,38] Considering this prior work and the findings presented in this study, it may

be that the generation of predictions for comparison with subsequent sensory feedback is sufficient for

error-based learning across motor and non-motor domains alike. In other words, task execution may not

always be required for learning, so long as the prediction and sensory feedback needed to compute error

are present.

References

1.   Miall, R. C. & Wolpert, D. M. Forward Models for Physiological Motor Control. *Neural Networks* **9**,
1265–1279 (1996).

2.   Jordan, M. I. & Rumelhart, D. E. Forward Models: Supervised Learning with a Distal Teacher.
Cognitive Science **16**, 307–354 (1992).

3.   Wallman, J. & Fuchs, A. F. Saccadic Gain Modification: Visual Error Drives Motor Adaptation. *Journal*
of Neurophysiology **80**, 2405–2416 (1998).

4.   Shadmehr, R., Smith, M. A. & Krakauer, J. W. Error Correction, Sensory Prediction, and Adaptation
in Motor Control. *Annu. Rev. Neurosci.* **33**, 89–108 (2010).

5.   Tseng, Y., Diedrichsen, J., Krakauer, J. W., Shadmehr, R. & Bastian, A. J. Sensory Prediction Errors
Drive Cerebellum-Dependent Adaptation of Reaching. *Journal of Neurophysiology* **98**, 54–62 (2007).

6.   Mazzoni, P. & Krakauer, J. W. An Implicit Plan Overrides an Explicit Strategy during Visuomotor
Adaptation. *J. Neurosci.* **26**, 3642–3645 (2006).

7.   Taylor, J. A., Krakauer, J. W. & Ivry, R. B. Explicit and Implicit Contributions to Learning in a
Sensorimotor Adaptation Task | Journal of Neuroscience. *The Journal of Neuroscience* **34**, (2014).

8.   Morehead, J. R., Taylor, J. A., Parvin, D. E. & Ivry, R. B. Characteristics of Implicit Sensorimotor
Adaptation Revealed by Task-irrelevant Clamped Feedback. *J Cogn Neurosci* **29**, 1061–1074 (2017).

9.   Kilteni, K., Andersson, B. J., Houborg, C. & Ehrsson, H. H. Motor imagery involves predicting the
sensory consequences of the imagined movement. *Nat Commun* **9**, 1617 (2018).

10.   Voss, M., Ingram, J. N., Haggard, P. & Wolpert, D. M. Sensorimotor attenuation by central motor
command signals in the absence of movement. *Nat Neurosci* **9**, 26–27 (2006).

11.   Sheahan, H. R., Ingram, J. N., Žalalytė, G. M. & Wolpert, D. M. Imagery of movements immediately
following performance allows learning of motor skills that interfere. *Sci Rep* **8**, 14330 (2018).

12.   Sheahan, H. R., Franklin, D. W. & Wolpert, D. M. Motor Planning, Not Execution, Separates Motor
Memories. *Neuron* **92**, 773–779 (2016).

13.   Tsay, J. S., Parvin, D. E. & Ivry, R. B. Continuous reports of sensed hand position during
sensorimotor adaptation. *Journal of Neurophysiology* **124**, 1122–1130 (2020).

14.   Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R. & Ivry, R. B. Invariant errors reveal
limitations in motor correction rather than constraints on error sensitivity. *Communications Biology* **1**,
1–7 (2018).

15.   Avraham, G., Morehead, J. R., Kim, H. E. & Ivry, R. B. Reexposure to a sensorimotor perturbation
produces opposite effects on explicit and implicit learning processes. *PLOS Biology* **19**, e3001147
(2021).

16.   Poh, E., Al-Fawakari, N., Tam, R., Taylor, J. A. & McDougle, S. D. Generalization of motor learning in
psychological space. *bioRxiv* 2021.02.09.430542 (2021) doi:10.1101/2021.02.09.430542.

17.   Hadjiosif, A. M., Krakauer, J. W. & Haith, A. M. Did We Get Sensorimotor Adaptation Wrong? Implicit
Adaptation as Direct Policy Updating Rather than Forward-Model-Based Learning. *J. Neurosci.* **41**,
2747–2761 (2021).

18.   Wei, K. & Koerding, K. Uncertainty of feedback and state estimation determines the speed of motor
adaptation. *Front. Comput. Neurosci.* **0**, (2010).

19. Semrau, J. A., Daitch, A. L. & Thoroughman, K. A. Environmental experience within and across
testing days determines the strength of human visuomotor adaptation. *Exp Brain Res* **216**, 409–418
(2012).

20. Albert, S. T. *et al.* An implicit memory of errors limits human sensorimotor adaptation. *Nat Hum
Behav* **5**, 920–934 (2021).

21. Wei, K. & Körding, K. Relevance of Error: What Drives Motor Adaptation? *Journal of Neurophysiology*
101, 655–664 (2009).

22. Kim, H. E., Parvin, D. E. & Ivry, R. B. The influence of task outcome on implicit motor learning. *eLife*
8, e39882 (2019).

23. Galea, J. M., Mallia, E., Rothwell, J. & Diedrichsen, J. The dissociable effects of punishment and
reward on motor learning. *Nat Neurosci* **18**, 597–602 (2015).

24. Leow, L.-A., Marinovic, W., Rugy, A. de & Carroll, T. J. Task errors contribute to implicit aftereffects in
sensorimotor adaptation. *European Journal of Neuroscience* **48**, 3397–3409 (2018).

25. Franklin, D. W. & Wolpert, D. M. Computational Mechanisms of Sensorimotor Control. *Neuron* **72**,
425–442 (2011).

26. Krakauer, J., W., Hadjiosif, A., Xu, J. & Wong, A., L. Motor Learning. in *Comprehensive Physiology*
(ed. Terjung, R.) (Wiley, 2019). doi:10.1002/cphy.

27. McDougle, S. D. & Taylor, J. A. Dissociable cognitive strategies for sensorimotor learning. *Nat
Commun* **10**, 40 (2019).

28. McDougle, S. D., Ivry, R. B. & Taylor, J. A. Taking Aim at the Cognitive Side of Learning in
Sensorimotor Adaptation Tasks. *Trends in Cognitive Sciences* **20**, 535–544 (2016).

29. Hutter, S. A. & Taylor, J. A. Relative sensitivity of explicit reaiming and implicit motor adaptation.
Journal of Neurophysiology **120**, 2640–2648 (2018).

30. Xu, K. Z. *et al.* Neural Basis of Cognitive Control over Movement Inhibition: Human fMRI and Primate
Electrophysiology Evidence. *Neuron* **96**, 1447-1458.e6 (2017).

31. Osada, T. *et al.* An Essential Role of the Intraparietal Sulcus in Response Inhibition Predicted by
Parcellation-Based Network. *J. Neurosci.* **39**, 2509–2521 (2019).

32. Duque, J., Greenhouse, I., Labruna, L. & Ivry, R. B. Physiological Markers of Motor Inhibition during
Human Behavior. *Trends in Neurosciences* **40**, 219–236 (2017).

33. Vyas, S. *et al.* Neural Population Dynamics Underlying Motor Learning Transfer. *Neuron* **97**, 1177-
1186.e3 (2018).

34. Requarth, T., Kaifosh, P. & Sawtell, N. B. A Role for Mixed Corollary Discharge and Proprioceptive
Signals in Predicting the Sensory Consequences of Movements. *J. Neurosci.* **34**, 16103–16116
(2014).

35. Mattar, A. A. G. & Gribble, P. L. Motor Learning by Observing. *Neuron* **46**, 153–160 (2005).

36. Cain, M., Botschko, Y. & Joshua, M. Passive Motor Learning: Oculomotor Adaptation in the Absence
of Behavioral Errors. *eNeuro* **8**, (2021).

37. Mendelsohn, A., Pine, A. & Schiller, D. Between Thoughts and Actions: Motivationally Salient Cues
Invigorate Mental Action in the Human Brain. *Neuron* **81**, 207–217 (2014).

38. Reddan, M. C., Wager, T. D. & Schiller, D. Attenuating Neural Threat Expression with Imagination.
Neuron **100**, 994-1005.e4 (2018).

39. Herzfeld, D. J., Vaswani, P. A., Marko, M. K. & Shadmehr, R. A memory of errors in sensorimotor
learning. *Science* **345**, 1349–1353 (2014).

40. Beers, R. J. van. Motor Learning Is Optimally Tuned to the Properties of Motor Noise. *Neuron* **63**,
406–417 (2009).

41. Wu, H. G., Miyamoto, Y. R., Castro, L. N. G., Ölveczky, B. P. & Smith, M. A. Temporal structure of
motor variability is dynamically regulated and predicts motor learning ability. *Nat Neurosci* **17**, 312–
321 (2014).

42. photonstorm.com, P. S.-. Phaser - Download - Phaser 3.24.1. https://phaser.io.

43. Tsay, J. S., Ivry, R. B., Lee, A. & Avraham, G. Moving outside the lab: The viability of conducting
sensorimotor learning studies online. *NBDT* 1–22 (2021) doi:10.51628/001c.26985.

ONLINE METHODS

Participants. Participants (n = 213, aged 18-35, 105 female) recruited from the research participation pools

at Princeton/Yale University and Prolific provided informed consent, approved by each University's IRB.

Ninety participants were excluded for failure to recall task instructions, as ascertained by a questionnaire

at the end of the experiment, leaving 123 participants for our analysis. See the *Questionnaire* section below

the *Test phase* sections for more details. We note that all the key results described here (significant learning

after No-Movement trials) held with or without these exclusions. We note that the samples used for each

experiment described in the text are around twice the size of similar studies in the literature, providing

additional statistical power to compensate for the experiment being conducted remotely.[39–41]

Task Setup. Experiments were conducted remotely using a custom JavaScript web application based on

Phaser3.24 (download available at [42]), similar to an approach previously described.[43] Each participant

viewed a single target located at either 45°, 135°, 225°, or 315° (with target position counterbalanced across

participants), 250 pixels from a central starting location. The target was visible throughout the experiment.

Participants used an input device of their choice to control their computer cursor during center-out

movements. One participant reported using a touchscreen device and was excluded from all analyses. The

remaining participants reported using either a trackpad (n = 112), an optical mouse (n = 86), or a trackball

(n = 14). A linear mixed model (LMM) did not show effects of Mouse Type on single-trial learning (STL),

although we observed that participants using a trackpad exhibited longer reaction times than others,

consistent with a previous report.[43]

Mouse position sampling rates depended on the exact hardware that each participant used to

complete the task. Sampling rates were likely affected by features of the specific mouse used, along with

features of the specific computer used, as computers may limit the rate at which the browser samples data

in order to cope with limited processing power. In general, sampling rates were around 60 Hz (median ±

interquartile range across all 213 participants recruited: 62.46 ± 2.17 Hz), but ranged from 19.23 Hz to

249.69 Hz. Note that the vast majority of sampling rates were near 60 Hz: Only 5% of mouse sampling

rates were < 41.79 Hz, and only 5% of mouse sampling rates were > 126.65 Hz.

Baseline training phase. During a baseline phase, participants were instructed to move their mouse into a

central starting location and then quickly make a straight, slicing movement through the target when it

turned green. After a 100 ms delay following the completion of the movement, the cursor reappeared near

the central starting location so that participants could return to that location to initiate the next trial.

Participants completed 5 of these trials with online and endpoint cursor feedback, followed by 5 trials without

visual feedback of the cursor location. Endpoint feedback was constituted by the cursor remaining at the

position where it had passed the target radius for 50 ms. Participants then completed 10 alternating trials

on which the target turned green and stayed green (Execution, 'Go' trials) and on which the target turned

magenta 100 ms after turning green, signaling that participants should withhold their movement (No-

Movement, 'No-Go' trials). After this baseline phase, participants were instructed to continue following these

instructions for the remainder of the experiment.

Test phase: Rotation and Error-Clamp Experiments (Fig. 2). During the test phase, 270 total trials were

divided into 3-trial triplets (**Fig. 1c**). The first and last trials of all triplets were Go trials, and participants

received neither online nor endpoint feedback about cursor location on these trials. The second trial of each

triplet was either a Movement or a No-Movement trial. On Movement trials, participants either received

rotated/error-clamped[8] visual feedback (15° clockwise [+] or counterclockwise [-], with sign randomized

across trials) or veridical/0° error-clamped visual feedback of their cursor location. On No-Movement (No-

Go) perturbation trials, participants viewed a brief animation of the cursor moving straight to the center of

the target following a trajectory deflected by 15° (clockwise or counterclockwise) from the target center.

Animation onset latency was set as a running median of the participant's reaction times on the previous 5

trials. Animation duration was set as a running median of the participant's movement times on the previous

5 trials. If a participant took longer than 400 ms to execute a movement, 800 ms to initiate the movement,

their reach trajectory changed by >10° during the movement, or the reach terminated ≥ 60° away from the

target, they received a warning and a 4s time-out. If a participant moved the cursor at all (>0 pixels) on a

No-Movement trial, the trial was immediately aborted, and they received a warning and a 4s time-out. The

No-Go manipulation was successful: Across the experiments, participants erroneously moved on 29.55 ±

0.18% (mean ± standard deviation) of No-Go trials, suggesting that, for the most part, they were consistently

planning movements on No-Go trials.

We used 6 possible triplet perturbation trial types (Movement/No-Movement: ±15° or 0°), each of

which occurred 15 times throughout each session. Triplets were pseudorandomly presented within each

block, with the constraints that a single rotation (±15° or 0°) could not occur on more than 2 consecutive

triplets and that the same movement condition (i.e., Movement or No-Movement) could not occur on more

than 3 consecutive triplets. Three repetitions of each triplet type occurred in blocks of 18 triplets, and

participants received a break after each of these blocks.

Test phase: Rotation and Error-Clamp Experiments with 0° Perturbations on Movement Trials (Fig. 3).

Experiments were conducted as described above, with the exception of the details described in this section.

For the experiments described in **Fig. 3a-d**, we used a reduced set of 3 possible triplet Perturbation trial

types (No-Movement, 15° clockwise error; No-Movement, 15° counterclockwise error; Movement, 0°

rotation). We maintained an equal number of Movement and No-Movement triplets throughout the session

in order to ensure that participants would reliably respond to the "Go" cue presented at the start of each

trial. So, each No-Movement triplet type occurred 22 times, while the Movement triplet type occurred 44

times. Triplets were pseudorandomly presented within each block, with the constraints that a single non-

zero rotation (15° clockwise, 15° counterclockwise) could not occur on more than 2 consecutive triplets.

For the experiments described in **Fig. 3e-h**, we used a set of 4 possible triplet Perturbation trial

types (No-Movement, 15° clockwise error; No-Movement, 15° counterclockwise error; No-Movement, 0°

error; Movement, 0° error-clamp). To maintain an equal number of Movement and No-Movement triplets

throughout the session, each No-Movement triplet type occurred 15 times and the Movement triplet type

occurred 45 times.

Questionnaire: The questionnaire asked participants to attest whether or not 1) they changed how they

moved the mouse after seeing the cursor miss the target, 2) their goal was to move the real mouse and not

the cursor straight through the green targets, and 3) their goal was to move the white cursor (not the real

mouse) straight through the green targets. Participants could select the options, "True," "False," or "Not

Sure."

Data analysis. Data were processed in Python 3.8.5. Trials with movement were excluded from analysis 1)

if any of the reaches in the triplet were not straight (aspect ratio > participant-wise mean + 3 * participant-

wise standard deviation), 2) if the participant received any warning for failure to follow task instructions (see

Feedback for failure to follow task instructions, above), or 3) if the triplet included a No-Movement No-Go

perturbation trial with any detectable mouse movement (>0 pixels).

Reach endpoint angle was computed as the angular distance between the center of the target and

the point at which the mouse passed the target's radial distance. Because mouse sampling rates did not

always allow us to measure mouse position at the exact target radius, we used the last sample before and

the first sample after the mouse passed the target radius to compute an interpolated mouse position at the

target radius, as described in a previous report.[43] We note that analyses comparing these measures to

measurements at the last sample of the reach (even when it was beyond the target) or the hand angle at

peak velocity did not result in substantially different hand angle measurements or statistical outcomes.

Single-trial learning (STL) was measured as the difference between reach endpoint angle on the

third and first trial of each triplet. For our initial analyses, the sign of STL corresponded to the direction of

the relative change in hand angle, with clockwise changes in hand angle taking a positive sign and

counterclockwise changes in hand angle taking a negative sign. When we collapsed STL data across

rotation directions, we normalized the sign of STL so that changes in hand angle opposite the direction of

the imposed rotation took a positive sign and changes in the direction of the rotation took a negative sign.

Statistics. Statistical tests were conducted in R (v. 4.0.3; packages rcompanion, rstatix, magrittr, ggplot2,

ggeffects, MuMIn, lmerTest, lme4, r2glmm, emmeans, stats). Data from 2 of the 4 experiments conducted

did not satisfy assumptions of two-way repeated measures ANOVA (non-existence of extreme outliers,

sphericity), so we employed a linear mixed modeling (LMM; R package lmerTest and lme4) approach for

our analyses. All LMM's included fixed effects of perturbation size and movement condition, as well as

random effects of subject. Degrees of freedom were estimated using the Kenward-Rogers approach, and

LMM outcomes were reported using ANOVA-style statistics. Partial R^2 was computed to report effect sizes

for the LMM factors (R package r2glmm). Post-hoc pairwise comparisons were performed between

estimated marginal means computed from the LMM (R package emmeans). For these and all subsequent

analyses, we corrected for multiple comparisons using the false-discovery rate approach to maintain

experiment-wise alpha at 0.05.

For one-off comparisons between samples or to distributions with 0-mean, we checked samples

for normality. When samples were normally distributed, we ran t-tests and computed Cohen's d to report

effect sizes for statistically significant results. Otherwise, we ran Wilcoxon-signed rank tests and measured

effect sizes using the rank-biserial correlation.

Initial efforts to model the remembered STL and the effects of errors experienced on preceding

trials on STL using LMM approaches suffered from singularity. Thus, we employed generalized linear

models (GLM; R package stats) assuming underlying Gaussian distributions.

To investigate whether STL was remembered following Movement and No-Movement trials, GLMs

with terms for the perturbation applied (15° CCW, 0°, or 15° CW), movement condition (Movement or No-

Movement), and their interaction were fit to triplet-wise remembered STL performance (hand angle on the

first trial of triplet *t + 1* minus hand angle on the first trial of triplet *t*) for each participant. We then compared

the distributions of coefficients for each term using Wilcoxon's signed-rank tests.

To investigate whether errors affected STL measurements beyond the triplet in which they

occurred, GLMs with terms for errors on Movement trials and No-Movement trials in the current triplet or

one triplet back, plus pairwise interactions for these 4 terms, were fit to triplet-wise STL performance for

each participant. We then compared the distributions of coefficients for each term using Wilcoxon's signed-

rank tests.