**Cross-species incompatibility between a DNA satellite and
the Drosophila Spartan homolog poisons germline genome integrity**

Cara L. Brand and Mia T. Levine*

Department of Biology and Epigenetics Institute, University of Pennsylvania, Philadelphia, PA

Running title: adaptive evolution to preserve genome integrity

Keywords: DNA satellite, coevolution, maternal haploid, 359, Spartan, Topoisomerase II

*Corresponding Author:
Dr. Mia Levine, University of Pennsylvania, 433 S. University Avenue, Philadelphia, PA 19104.
p: 215-573-9709 e: m.levine@sas.upenn.edu.

ABSTRACT

Satellite DNA spans megabases of eukaryotic sequence and evolves rapidly. Paradoxically,

satellite-rich genomic regions mediate strictly conserved, essential processes like chromosome

segregation and nuclear structure. A leading resolution to this paradox posits that satellite DNA

and satellite-associated chromosomal proteins coevolve to preserve these essential functions.

We experimentally test this model of intra-genomic coevolution by conducting the first evolution-

guided manipulation of both chromosomal protein and DNA satellite. The 359bp satellite spans

an 11Mb array in *D. melanogaster* that is absent from its sister species, *D. simulans*. This

species-specific DNA satellite colocalizes with the adaptively evolving, ovary-enriched protein,

Maternal Haploid (MH)–the Drosophila homolog of Spartan. To determine if MH and 359

coevolve, we swapped the *D. simulans* version of MH ("MH[sim]") into *D. melanogaster*.

MH[sim] triggers ovarian cell death, reduced ovary size, and loss of mature eggs. Surprisingly,

the *D. melanogaster mh* null mutant has no such ovary phenotypes, suggesting that MH[sim] is

toxic in a *D. melanogaster* background. Using both cell biology and genetics, we discovered that

MH[sim] poisons oogenesis through a DNA damage pathway. Remarkably, deleting the

D. melanogaster-specific 359 satellite array completely restores *mh[sim]* germline genome

integrity and fertility, consistent with a history of coevolution between these two fast-evolving

loci. Germline genome integrity and fertility are also restored by overexpressing Topoisomerase

II (Top2), suggesting that MH[sim] interferes with Top2-mediated processing of 359. The

observed 359-MH[sim] cross-species incompatibility supports a model under which ostensibly

inert repetitive DNA and essential chromosomal proteins must coevolve to preserve germline

genome integrity.

RESULTS AND DISCUSSION

DNA satellite-enriched genomic regions evolve rapidly and yet support strictly conserved

nuclear functions.[1-9] A classic resolution to this paradox posits that DNA satellite-associated

proteins evolve adaptively to mitigate deleterious proliferation of DNA satellite sequence

variants.[10] Repeated bouts of DNA satellite evolution and chromosomal protein adaptation result

in exquisitely coevolved satellites and satellite-associated proteins. This model of coevolution

predicts pervasive incompatibilities between satellite DNA and chromosomal proteins from

closely related species: adaptively evolving chromosomal proteins from one species should fail

to package or process DNA satellites from another.[10-12]

Evidence for this coevolution model has emerged from engineering "evolutionary mismatches"

between the adaptively evolving chromosomal protein(s) of one species and the DNA satellite

landscape of a close relative. Under one approach, a diverged chromosomal protein is

introduced into a closely related species, generating an evolutionary mismatch between the

manipulated protein and one or more DNA satellites.[12-15] Consistent with disrupted DNA

satellite:chromosomal protein coevolution, the naïve protein typically perturbs a satellite-

mediated function, such as chromosome segregation or nuclear organization.[12,14,15] In these

cases, however, the incompatible DNA satellites are unknown. A second approach crosses

sister species to generate evolutionary mismatches between chromosomal proteins and DNA

satellites in hybrid progeny. Consistent with disrupted DNA satellite:chromosomal protein

coevolution, interspecies hybrid inviability has been linked to satellite-rich genomic loci.[16,17] In

these systems, however, the incompatible chromosomal proteins are unknown. To date, there

are no cases of experimental identification of both chromosomal protein and satellite engaged in

coevolution.

To experimentally probe *both sides* of the coevolution model, we searched for a rapidly evolving

DNA satellite associated with an adaptively evolving chromosomal protein. In *Drosophila*

melanogaster, the 359bp satellite spans an 11Mb array at the base of the *X* chromosome.[18,19]

Close relatives of *D. melanogaster*, including *D. simulans* and *D. erecta*, lack this *X*-linked

satellite array.[20] Instead, these species have shorter arrays of "359-like" sequence dispersed

throughout heterochromatin and euchromatin.[9,21,22] Such extreme lineage-restriction to

D. melanogaster makes this DNA satellite array an ideal locus for testing the coevolution model.

On the protein side, we identified from the literature Maternal Haploid (MH), an ovary-enriched

protein that is maternally provisioned to the embryo and colocalizes with the 359 satellite.[23,24]

Embryos of *mh* null mothers suffer paternal chromosome mis-segregation at the very first

mitosis, suggesting that the maternally-provisioned MH prepares the otherwise inert, sperm-

deposited paternal chromosomes for participation in embryonic mitosis.[23,24] Most of these

embryos arrest around the first mitotic division. A smaller fraction develop beyond the first

division, cycling only the "maternal haploid" complement of chromosomes until arrest prior to

hatching.[23-25]

If the *D. melanogaster*-specific 359 proliferation triggered *mh* to innovate, we should detect

evidence of positive selection at *mh* between *D. melanogaster* and *D. simulans*. To determine if

mh evolves adaptively, we conducted a McDonald-Kreitman test[26] using polymorphism within

D. melanogaster and *D. simulans* populations and divergence between *D. melanogaster* and

D. simulans (2.5 million years diverged[27]). This comparison revealed an excess of

nonsynonymous fixations, consistent with a history of adaptive evolution (Figure 1A). The

dynamic evolution of the 359 satellite and adaptive evolution of a 359-associated protein, MH,

raises the possibility that *mh* recurrently evolves to preserve a biological function compromised

by 359 satellite proliferation.

To test the possibility of MH:359 coevolution, we first conducted an evolution-guided

manipulation of *mh* to generate an "evolutionary mismatch" between protein and satellite. We

used CRISPR/Cas9 to integrate into the native *mh* locus of *D. melanogaster* either a 3xFLAG-

tagged *mh* coding sequence from *D. melanogaster* (our control fly, "*mh[mel]*") or a 3xFLAG-

tagged *mh* coding sequence from *D. simulans* (our experimental fly, "*mh[sim]*", Figure 1B). Both

the *D. melanogaster* and the *D. simulans* coding sequences were codon-optimized for

D. melanogaster. We observed equivalent expression of the two transgenes (Figure S1A).

The *mh* null mutant phenotype in the early embryo motivated our prediction that an evolutionary

mismatch between the *D. simulans mh* and the *D. melanogaster* 359 *X*-linked array would

compromise paternal chromosome segregation during the first mitotic division. Specifically, we

predicted that MH[sim] would fail to recognize and process 359, triggering mis-segregation of

the paternal *X*-chromosome. This defect would result in reduced female fertility and a dearth of

female progeny. We discovered that *mh[sim]* females produced significantly fewer progeny than

control *mh[mel]* females (Figure 1C); however, contrary to our prediction, the progeny sex ratio

did not deviate from 50/50 (Figure S1B). These data suggest that paternal 359 is *not* uniquely

Brand and Levine Figure 1

Figure 1. MH evolves adaptively to preserve female fertility. (A) Counts of synonymous and nonsynonymous polymorphic and fixed sites within and between *D. melanogaster* and *D. simulans;* χ^2 test, p = 0.04. (B) Swap strategy: the *D. melanogaster* ("*D. mel*", blue) or *D. simulans* ("*D. sim*," yellow) *mh* coding sequence, codon-optimized for *D. melanogaster* and 3xFLAG-tagged, replaced the native *mh* gene on the *X* chromosome. (C) Total offspring from *mh[mel]* or *mh[sim]* females crossed to wildtype (w^{1118}) males. (D) Frequency distribution of embryos at increasing mitotic cycle numbers collected for 70 minutes from *mh[mel]*, *mh[sim]*, and *mh^1* females. Dashed bars from *mh^1* females correspond to embryos undergoing mitotic catastrophe likely triggered at the first mitosis. Solid gray bars representing *mh^1*-derived embryos are presumed to be maternal haploid. (E) Representative images and ovary size estimates from *mh[mel]* and *mh[sim]* females. (F) Number of mature eggs per ovary pair from *mh[mel]* and *mh[sim]* females. (*t*-test: "***" = p < 0.001, scale bar = 100μm)

vulnerable to the presence of MH[sim] during the first mitotic division. Consistent with this

inference, we observed that *mh[sim]* completely rescues the first mitotic division: embryos from

mh[mel] and *mh[sim]* mothers show equivalent, normal distributions of embryonic stages from a

70-minute collection (Figure 1D). In contrast, embryos produced by *mh* null mothers typically

arrest during the first division (Figure 1D). Moreover, we observed no evidence of elevated

maternal haploid embryos from *mh[sim]* mothers (Figure S1C). These data suggest that *mh[sim]*

does not phenocopy the *mh* null early embryonic phenotype.

To uncover an alternative source of the *mh[sim]* fertility defect, we looked at the developmental

stage just before the first embryonic mitosis: oogenesis. Although *mh* is highly expressed during

oogenesis, previous reports suggested that *mh* null alone yields no ovary phenotype[23,24]. We

similarly detected no difference in ovary size or mature egg number of *mh* null mothers

compared to heterozygous controls (Figure S1D,E). In contrast, *mh[sim]* ovaries are significantly

smaller than *mh[mel]* ovaries and are depleted of the most mature egg stages (Figure 1E,F).

This unexpected *mh[sim]* ovary phenotype, combined with the complete rescue of the first

embryonic division by *mh[sim]*, suggests that *mh[sim]* does not behave as a loss-of-function

allele. Instead, MH[sim] might be toxic.

To explore the possibility that MH[sim] is toxic, we first asked if MH[sim] localizes aberrantly in

the ovary. We visualized MH[mel] and MH[sim] by staining ovaries with anti-FLAG. We

discovered that MH[mel] localized primarily in the earliest stages of oogenesis (the germarium,

Figure 2A,B). MH[sim] localized not only in these early stages of oogenesis but also on the

"nurse cell" nuclei of later stage egg chambers and weakly on the somatic follicle cell nuclei

(Figure 2A,B). The aberrant persistence of MH[sim] during oogenesis, combined with

compromised *mh[sim]* ovary development, raised the possibility that MH mislocalization alone

might be toxic. To test this hypothesis, we used the UAS/GAL4 system to overexpress MH[mel]

in the female germline (driver nos-Gal4-VP16). In ovaries overexpressing MH[mel], we indeed

observed elevated levels and aberrant persistence of the protein in later stage egg chambers

(Figure S2A). Nevertheless, these females gave rise to abundant progeny (Figure S2B),

suggesting that mislocalization alone cannot explain the compromised ovary development of

mh[sim] females. In contrast, overexpression of MH[sim] in otherwise wildtype ovaries resulted

in a complete absence of mature eggs (nosGAL4 > UASp-*mh[sim]*: $\mu = 0$ eggs/ovary pair versus

native > *mh[sim]*: $\mu = 15$ eggs/ovary pair, n = 63 ovary pairs/genotype). Consequently, these

females were completely sterile (Figure. S2B). These data suggest that MH[sim] – which

Brand and Levine Figure 2

Figure 2. MH[sim] poisons oogenesis through a DNA damage pathway. (A) Diagram of a Drosophila ovary (above) and a single ovariole (below) with the germline stem cells in the germarium at the anterior (left) position and the mature eggs at the posterior position (right). The dashed box shows the developmental stages shown in images 2B-2D (Created using BioRender.com). (B) *mh[mel]* and *mh[sim]* ovaries stained with anti-FLAG to visualize MH localization (left). Merged images of single nuclei (*) show no MH foci on the DNA (dashed inset boxes). Merged images of enlarged egg chambers imaged under higher laser power reveal that MH[sim] but not MH[mel] localizes to the peripheral follicle cells (right). (C) Incidence of cell death captured by the fraction of ovarioles with condensed nuclei (arrowheads) in *mh[mel]* and *mh[sim]* ovaries. (D) γH2Av signal in *mh[mel]* and *mh[sim]* ovaries and the quantification of normalized fluorescent signal intensity. Note that the expected γH2Av-positive cells in the germarium in *mh[mel]* are absent under the imaging parameters used but are indeed present, see Figure S2D. (E) Ovary size estimates from *mh[mel];Chk2$^{-/-}$* and *mh[sim];Chk2$^{-/-}$* females. (F) Number of mature eggs per ovary pair from *mh[mel];Chk2$^{-/-}$* and *mh[sim];Chk2$^{-/-}$* females. In panels E and F, dotted lines correspond to *mh[mel]* and *mh[sim]* averages reported in Figure 1E and F, respectively. (*t*-test: "***" = $p < 0.001$, "n.s." $p > 0.05$, scale bar = 25µm)

functions normally in its native *D. simulans* genome – is toxic to oogenesis in *D. melanogaster*.

This toxicity appears to be dose-dependent. Heterozygous *mh[mel]/mh[sim]* females give rise to

progeny counts similar to *mh[mel]* homozygotes (Figure S2C).

To study the cell biological basis of this block to oogenesis, we turned back to ovaries of

females expressing the CRISPR-introduced *mh[mel]* or *mh[sim]* transgene under the native

promoter (Figure 1B). We observed an excess of hyper-condensed nuclei in *mh[sim]* ovaries,

consistent with elevated cell death (Figure 2C[28,29]). A classic trigger of cell death is the

accumulation of DNA damage.[30] To visualize DNA damage, we stained *mh[mel]* and *mh[sim]*

ovaries for the double-stranded break marker, γH2Av.[31,32] We observed elevated DNA damage

signaling in *mh[sim]* ovaries (Figure 2D, S2D). This phenotype further distinguishes *mh[sim]*

from *mh* null ovaries – the latter show no evidence of elevated DNA damage (Figure S2E). To

address the hypothesis that MH[sim] compromises oogenesis through a DNA repair pathway,

we combined *mh[sim]* with a null mutation in a DNA damage checkpoint gene. The gene, *Chk2*

(also known as *mnk*), normally blocks egg production in the presence of DNA damage.[33,34]

Chk2$^{-/-}$ ovaries bypass this checkpoint, allowing a female to make mature but damaged eggs in

the presence of elevated DNA damage. We discovered that *Chk2$^{-/-}$* restores *mh[sim]* ovaries to

mh[mel]-like ovary size and *mh[mel]*-like egg production (Figure 2E,F). However, the

mh[sim];*Chk2$^{-/-}$* females are sterile while *mh[mel]]*;*Chk$^{-/-}$* females retain fertility (Figure S2F).

These data suggest that MH[sim] compromises oogenesis by triggering DNA damage.

Applying these phenotypic data to the coevolution model, we hypothesized that MH[sim]-

induced DNA damage depends on the 11Mb array of 359 satellite in *D. melanogaster*. Under

this model, MH[sim]-specific residues are incompatible with 359. Removing 359 should restore

germline genome integrity and fertility of *mh[sim]* females. To directly test this prediction, we

took advantage of a fly strain that lacks the 11Mb array of *X*-linked 359 satellite (Figure 3A[35]).

We recombined this 359-deletion, called *Zygotic hybrid rescue*, or "*Zhr*," onto both the *mh[mel]*

and the *mh[sim] X* chromosomes (Figure 3A). If MH[sim]-induced toxicity depends on the

presence of the 359 expansion, *mh[sim],Zhr* females should have minimal DNA damage and

recover fertility. Remarkably, the 359-deletion completely restores the DNA damage marker,

γH2Av, to wildtype (low) levels (Figure 3B). Consistent with restored germline genome integrity

of *mh[sim]* females, we observed no difference in ovary size and no difference in egg production

between *mh[mel]* and *mh[sim]* females that lack 359 (Figure 3C,D). Finally, the 359-deletion

completely restores *mh[sim]* fertility to *mh[mel]* levels (Figure 3E). These data reveal that

Brand and Levine Figure 3

Figure 3. The 359 satellite deletion rescues *mh[sim]* genome integrity and fertility. (A) The *Zhr X* chromosome lacks the 11Mb pericentromeric 359 satellite array (left). A 10-cycle PCR distinguishes between wildtype 359 copy number and the 359-deletion (*Zhr*) and validates the recombined *mh[mel],Zhr* and recombined *mh[sim],Zhr X* chromosomes (right). (B) γH2Av signal in *mh[mel],Zhr* and *mh[sim],Zhr* ovaries and the quantification of normalized fluorescent signal intensity. (C) Ovary size of *mh[mel],Zhr* and *mh[sim],Zhr* females. (D) Number of mature eggs per ovary pair from *mh[mel],Zhr* and *mh[sim],Zhr* females. (E) Progeny counts from *mh[mel],Zhr* and *mh[sim],Zhr* females crossed to wildtype (*w[1118]*) males. In panel B, dotted lines correspond to *mh[mel]* and *mh[sim]* averages reported in Figure 2D. In panels C, D, and E, dotted lines correspond to *mh[mel]* and *mh[sim]* averages reported in Figure 1E, F, and C, respectively. (*t*-test: "n.s." p > 0.05, scale bar = 25μm)

MH[sim] toxicity depends on 359, consistent with a history of coevolution between these two

fast-evolving components of the Drosophila genome.

The observed 359-dependent toxicity rather than loss of function suggests that MH[sim] may

interfere with the preservation of 359 integrity. To define a molecular basis for this interference,

we used two, well-characterized MH homologs as guides. The MH homologs in worm (DVC-1)

and human (Spartan) use the conserved Spartan metalloprotease domain to cleave DNA-

protein crosslinks that form between DNA and Topoisomerase II (Top2[36-38]). These crosslinks

dock Top2 at DNA entanglements, which Top2 resolves by inducing double strand breaks

followed by re-ligation.[39-41] In *D. melanogaster*, Top2 specifically cleaves 359[42] and resolves

DNA entanglements involving 359 during female meiosis.[43] We hypothesized that MH[sim]

interferes with Top2 resolution of 359 entanglements. This interference model predicts that Top2

is limiting in the presence of MH[sim]. To test this prediction, we reduced Top2 using a

heterozygous loss of function mutant and overexpressed Top2 in the ovary using the UAS/

GAL4 system in an *mh[mel]* or *mh[sim]* background. Reduction of Top2 exacerbates *mh[sim]*-

dependent subfertility (Figure 4A) while Top2 overexpression in the ovary completely rescues

mh[sim] fertility (Figure 4B). The rescued *mh[sim]* ovaries also showed restored genome

integrity (Figure 4C). Excess Top2 appears to overcome MH[sim] interference. Combined with

the literature on DNA-Top2 crosslink cleavage by MH homologs DVC-1 and Spartan, these data

raise the possibility that MH[sim] over-actively clears the 359-Top2 crosslinks necessary to

resolve 359 entanglements in the female germline. Persistent DNA entanglements would trigger

the observed DNA damage that blocks oogenesis progression (Figure 4D).

Our model is motivated in part by the observation that repeat-rich heterochromatin, but

especially the 11Mb array of 359, is uniquely vulnerable to DNA entanglements.[16,43,44] If 359 is

so deleterious, how could it have proliferated? DNA satellites can behave selfishly, gaining a

transmission advantage from one generation to the next.[45,46] We suspect that such non-

Mendelian segregation led to 359 proliferation, triggering MH to evolve adaptively (Figure 4E).

However, we cannot formally rule out the possibility that a selection pressure distinct from 359

proliferation triggered MH adaptive evolution. Under this alternative model, the *D. melanogaster*

version of MH evolved first, releasing constraint on 359 copy number. Most likely, both selection

and loss of constraint operate cyclically.

Brand and Levine Figure 4

Figure 4. MH[sim] interferes with Top2 processing of 359 entanglements. (A) Progeny counts from *mh[mel];Top2⁻/+* and *mh[sim];Top2⁻/+* females crossed to wildtype (w^{1118}) males. (B) Progeny counts from nos-Gal4-VP16 (female germline GAL4) driven *mh[mel];*UAS-*Top2* or *mh[sim];*UAS-*Top2* females crossed to wildtype (w^{1118}) males. (C) γH2Av signal from ovaries of nos-Gal4-VP16 driven *mh[mel];*UAS-*Top2* or *mh[sim];*UAS-*Top2* females and quantification of normalized fluorescent signal intensity. (D) Model of MH[sim] interference with Top2 processing of 359 entanglements. These entanglements threaten genome integrity and ultimately, fertility. MH[mel], in contrast, has no measurable function in the ovaries, suggesting that it avoids interfering with 359 processing by Top2. (E) Model of MH evolution tracking 359 satellite proliferation. In panel A, dotted lines correspond to *mh[mel]* and *mh[sim]* averages reported in Figure 1C. In panel C, dotted lines correspond to *mh[mel]* and *mh[sim]* averages reported in Figure 2D. (*t*-test: "***" = p < 0.001, "n.s." p > 0.05, scale bar = 25µm)

Regardless of the force(s) that promoted 359 proliferation, the 359:MH system offers an

important elaboration of the canonical model of intra-genomic coevolution.[10,47-49] This canonical

model posits that chromosomal proteins evolve adaptively to recognize and process novel

satellite repeat variants. Under this model, the mismatched *mh[sim]* allele should fail to perform

an *mh* function; that is, act as a loss of function allele. Instead, we demonstrate that *mh[sim]* is

toxic, suggesting that *mh[mel]* instead evolved adaptively to *avoid interfering* with 359

processing, and likely 359 processing by Top2. The observed colocalization of MH[mel] foci with

Top2 and 359 in the embryo ([24], many cycles after the first mitosis) motivates future work

dissecting the functional consequences of MH adaptive evolution during this distinct

developmental stage.

359-mediated toxicity to oogenesis highlights the catastrophic functional consequences of DNA

satellite evolution. Importantly, 359-mediated toxicity is also apparent in *D. melanogaster-*

D. simulans hybrid embryos: a distinct, unmapped gene on *D. simulans* chromosome 2[50-53]

interacts deleteriously with 359 to cause embryonic chromosome mis-segregation, genome

instability, and lethality.[16,35,54] This interspecies hybrid dysfunction in the embryo, together with

the 359:*mh[sim]* toxicity in the ovary reported here, suggests that recurrent bouts of coevolution

not only shape essential genome functions within species but also can trigger hybrid

incompatibilities between species.

ACKNOWLEDGEMENTS

We thank Isabella Farkas and Courtney Christopher for technical assistance. We also thank the

Levine Lab, M. Patel, N. Phadnis, A. Das, and D. Dudka for feedback on the manuscript and the

Levine Lab, P. Geyer, H. Malik, R.S. Hawley, and M. Buszczak for discussions about the

project. This work was supported by a Life Sciences Research Foundation fellowship to C.L.B.

and National Institutes of Health (NIH) NIGMS grant R35GM124684 to M.T.L.

AUTHOR CONTRIBUTIONS

Conceptualization, M.T.L and C.L.B.; Methodology, M.T.L and C.L.B.; Investigation, M.T.L and

C.L.B; Writing M.T.L and C.L.B; Funding Acquisition, M.T.L and C.L.B.

DECLARATION OF INTERESTS

The authors declare no competing interests.

STAR METHODS

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse Monoclonal anti-FLAG M2	Sigma-Aldrich	Cat#F3165
Mouse Monoclonal 12G10 anti-αTubulin	Developmental Studies Hybridoma Bank	Cat#12G10 anti-alpha-tubulin, RRID:AB_1157911
Mouse Monocolonal anti-Histone 2A Gamma Variant, Phosphorylated	Developmental Studies Hybridoma Bank	Cat#UNC93-5.2.1, RRID:AB_261807
Chicken Polyconal anti-Green Fluorescent Protein	Aves Labs	Cat# GFP-1010, RRID:AB_230731
Chemicals, peptides, and recombinant proteins		
Phusion High-Fidelity DNA Polymerase	NEB	Cat#M0530
RIPA buffer	Cell Signaling Technology	Cat#9806
Protease Inhibitor Cocktail	Roche	Cat#11873580001
PMSF	Cell Signaling Technology	Cat#8553
Benzonase	Sigma-Aldrich	Cat#E1014
ProLong Gold Antifade Mountant with DAPI	ThermoFisher Scientific	Cat#P36931
Critical commercial assays		
Kwikquant Western Blot Detection Kit	Kindle Biosciences	Cat#R1004
Deposited data		
mh alleles from *D. simulans*	This paper	OL546458..64
Experimental models: Organisms/strains		
yw; nos-Cas9(II-attP40)	Best Gene, Inc	N/A
yw; PBac[y⁺-attP-9A]VK00018	Bloomington Drosophila Stock Center	BDSC:9736
Gal4::VP16-nos	Bloomington Drosophila Stock Center	BDSC:64277
Zhr¹	Bloomington Drosophila Stock Center	BDSC:25140
mnkˡ⁶	N. Phadnis	N/A
Top2¹⁷⁶	P. Geyer	N/A
w¹¹¹⁸	MTL laboratory	N/A
mh¹	Bloomington Drosophila Stock Center	BDSC:7130
P{gcid.EGFP.cidIII}.2	K. McKim,[55]	N/A
Oligonucleotides		

Primers to amplify mh alleles from *D. simulans*, see Table S1	This paper	N/A
Primers to screen *mh* CRISPR transformants, see Table S1	This paper	N/A
Primers to detect native *mh* locus, see Table S1	This paper	N/A
Primers to screen for PCR-introduced mutations in the cloned *mh* UAS constructs, see Table S1	This paper	N/A
Primers to screen for 359 deletion (*Zhr*), see Table S1	[6]	N/A
Recombinant DNA		
Plasmid: pBFv-U6.2	Addgene	Addgene#138400
Plasmid: pBFv-U6.2B	Addgene	Addgene#138401
Plasmid: 3xFLAG:*mh[mel]* and 3xFLAG:*mh[sim]* HDR plasmids	This paper	N/A
Plasmid: pUASp-attB	Drosophila Genomics Resource Center	DGRC#1358
Software and algorithms		
Geneious v 11.1.5	Biomatters	https://www.geneious.com/
FIJI	ImageJ2	https://imagej.net/software/fiji/

Contact for reagent and resource sharing

240 Further information and requests for resources and reagents should be directed to and will be

241 fulfilled by the Lead Contact, Mia Levine (m.levine@sas.upenn.edu).

Population genetic and molecular evolution analyses

We conducted population genetic analysis of *mh* using multiple alleles from both

D. melanogaster and *D. simulans*. We obtained eight *D. melanogaster mh* alleles (coordinates

X:15472804-15475400, dmel r6.4) from lines collected in Lyon, France.[56] We amplified seven *D.*

simulans mh alleles from lines collected in Nairobi, Kenya (Accession OL546458-OL546464)[57].

Importantly, a duplication event occurred along the *D. simulans* lineage, resulting in a full-length

copy of the *mh* ortholog, and a tandem partial duplicate.[58] To specifically amplify the full length

mh ortholog, we designed primers that anneal to unique genomic sequence (Table S1). We

then prepared genomic DNA and conducted PCR amplification followed by Sanger sequencing

using standard protocols. We aligned the sequences in Geneious using the Geneious Alignment

algorithm with default settings (Geneious v11.1.5, Biomatters, Auckland, New Zealand) and

confirmed alignment quality by eye. We performed a McDonald-Kreitman test[26] with the

D. melanogaster and *D. simulans mh* coding sequences.

Fly stock construction and husbandry

Constructing gene swaps

We used CRISPR/Cas9 to generate *D. melanogaster* flies that encode a transgenic

D. melanogaster allele or a *D. simulans* allele of *mh,* integrated into the native location. We first

generated a U6 promoter-driven guide RNA construct by cloning sgRNAs flanking the coding

sequence of *mh* (5': GGATTGGCCCAGGATCAACA, 3': CGTGGAGAGCTTCTGCCGCG) into

pBFv-U6.2 and pBFv-U6.2B backbones. We shuttled the 3' sgRNA into pBFv-U6.2 to create a

dual sgRNA vector (University of Utah Mutagenesis Core). In parallel, we constructed homology

directed repair (HDR) plasmids encoding one kilobase homology arms 5' and 3' of their

respective guide RNAs. Between the homology arms we synthesized a codon-optimized (for

D. melanogaster) *mh* coding sequence from either *D. melanogaster* or *D. simulans* (GenScript,

Piscataway, NJ). We N-terminally tagged each sequence with 3xFLAG along with a linker

sequence (GGTGGTTCATCA). We injected the dual sgRNA vector and a single HDR plasmid

into the Cas9-expressing line, *yw; nos-Cas9(II-attP40)* (BestGene Inc, Chino Hills, CA).

We crossed single males, injected as embryos, to FM7 (*X*-chromosome balancer) females. We

screened F_1 females to identify positive transformants using forward primer

5'-AAGTGTCGCGCTATTTCACC-3' and reverse primer

5'-TCACCGTCATGGTCTTTGTAGTCCAT-3'. We then backcrossed the positive F_1 females to

FM7 males and self-crossed the balanced F_2 progeny to generate lines homozygous for either

mh[mel] or *mh[sim]* allele. To confirm that the introduced alleles encoded the expected

sequence and in the expected location, we amplified the entire region from homozygous flies

using primers that anneal outside of the homology arms (5'-AATGGATTTCGGCAAATGAG-3',

5'-GTCGTTGTAGGAGCCCATGT-3') and then sequenced across the entire region. We also

designed primers that amplified the native *mh* locus (5'-GGCCCTGCTCATATCGTATC-3',

5'-AAGAACCTTACTGCGTGCAAC-3') to confirm that our final genotypes were true

replacements. Primers can be found in Table S1.

Constructing UAS-mh and UAS-Top2 lines

We used the *ΦC31* integrase-mediated transgenesis system to introduce into the same landing

site *mh* from *D. melanogaster* or *D. simulans* downstream of an "upstream activating sequence"

or "UAS"[59]. Using the HDR plasmids as a template (see above), we PCR-amplified the 3xFLAG-

tagged *mh* coding sequence (either *D. melanogaster* or *D. simulans)* using Phusion High-

Fidelity DNA Polymerase (NEB, Ipswich, MA). We cloned the resulting PCR products into

NotI/XbaI sites of the pUASp-attB vector (*Drosophila* Genomics Resource Center, Bloomington,

IN). We confirmed the absence of PCR-introduced mutations in the cloned UAS-*mh[mel]* and

UAS-*mh[sim]* alleles by direct Sanger sequencing of the constructs (Table S1). We introduced

the constructs into *D. melanogaster yw; PBac[y⁺-attP-9A]VK00018* flies, which have an attP

transgene landing site at cytological position 75A10 on chromosome *3L* (BestGene Inc, Chino

Hills, CA). We next made each transgene homozygous. To overexpress the transgenic alleles,

we crossed these stocks to *Gal4::VP16-nos* (BDSC #64277), which drives germline expression

of transgenes downstream of UAS.

Similarly, we used the *ΦC31* integrase-mediated transgenesis system to introduce *Top2* from

D. melanogaster downstream of an UAS promoter.[59] We synthesized a codon-optimized (for

D. melanogaster) *Top2* coding sequence from *D. melanogaster* (Twist, South San Francisco,

CA). We N-terminally tagged each sequence with 3xHA along with a linker sequence

(GGTGGTTCATCA). We introduced the constructs into *D. melanogaster yw; PBac[y⁺-attP-*

9A]VK00018 flies (see above, BestGene Inc, Chino Hills, CA). We next constructed either

mh[mel]; UAS-*Top2[mel]* or *mh[sim]*; UAS-*Top2[mel]* stocks using balancer chromosomes. To

overexpress the transgenic *Top2* allele, we crossed either *mh[mel]*; UAS-*Top2[mel]* or *mh[sim]*;

UAS-*Top2[mel]* males to either *mh[mel]*; *Gal4::VP16-nos* or *mh[sim]*; *Gal4::VP16-nos* females,

respectively.

Zhr rescue stocks

To generate stocks that encode both the *X*-linked *mh*-transgene and the *X*-linked 359 satellite

deletion (*Zhr¹,* BDSC #25140), we first generated trans-heterozygote females. We crossed

these trans-heterozygote females to FM7 males and used PCR to assay individual recombinant

male progeny for the presence of both the *mh* transgene and *Zhr.* We detected the *mh*

transgenes with forward primer 5'-AAGTGTCGCGCTATTTCACC-3' and reverse primer

5'-TCACCGTCATGGTCTTTGTAGTCCAT-3'. To detect the *Zhr* mutation (*i.e.*, 359 satellite

deletion), we used forward primer 5'-TATTCTTACATCTATGTGACC-3' and reverse primer

5'-GTTTTGAGCAGCTAATTACC-3'.[6] Performing a 10-cycle PCR at an annealing temperature

of 52C yields a band only in the presence of the 11Mb 359 satellite array (Figure 3A). We

backcrossed males positive for both the *mh* transgene and *Zhr* mutation to FM7 females to

generate a permanent stock.

Additional stocks

Heterozygote *mh[mel]*/*mh[sim]* females were generated by crossing *mh[mel]* females to *mh[sim]*

males.

We used a +/FM7; +/CyO stock to generate flies encoding both the *mh* transgene at the native

locus (chromosome *X*) and the *Chk*[-/-] (*mnk*) mutation (chromosome *2*). The *mnk*[l6] stock[34] was a

gift from N. Phadnis.

To generate heterozygous *Top2* hypomorph females, we also used a +/FM7; +/CyO stock to

generate flies encoding both the *mh* transgene at the native locus (chromosome *X*) and a

heterozygous *Top2*[176]/CyO mutation (chromosome *2*). The *Top2*[176] stock[60] was a gift from P.

Geyer.

Immunoblotting

To assay 3x-FLAG MH protein abundance in the ovary, we dissected 20 ovary pairs in 1X PBS

and ground the material in RIPA buffer (Cell Signaling Technology, Danvers, MA), Protease

Inhibitor Cocktail (Roche, Basel, Switzerland), and 2X PMSF (Cell Signaling Technology,

Danvers, MA). To promote solubility, we incubated the lysate in benzonase (Sigma Aldrich, St.

Louis, MO) for 1hr at 4C. We used 20µg of lysate and probed with 1:10,000 anti-FLAG (M2,

Sigma Aldrich, St. Louis, MO) or 1:1000 anti-αTubulin (Developmental Studies Hybridoma Bank,

Iowa City, IA) and 1:1000 anti-mouse HRP secondaries (Kindle Biosciences, Greenwich, CT).

We exposed blots with Kwikquant Western Blot detection kit and imaged with a Kwikquant

imager (Kindle Biosciences, Greenwich, CT).

Fertility assays

Female fertility

To assay female fertility, we first aged virgin females 3-5 days. For each replicate vial, we crossed four virgin females to four w^{1118} males. We conducted all crosses on molasses food at 24C. We flipped the parents onto new food every three days over the course of nine days and counted all progeny that emerged.

Ovary size and mature egg counts

To determine the number of mature eggs and ovary size from focal genotypes, we first dissected ovary pairs in 1X PBS and imaged at 8X magnification with a Leica DFC7000 T camera. We quantified the area of each ovary using the polygon tool in FIJI[61] to define the borders of the tissue. For each ovary pair, we used the Freehand tool in FIJI to calculate the area (mm^2) within these boundaries for each individual ovary. After imaging, we counted the number of eggs that contain elongated dorsal appendages (stages 13 and 14).

Immunofluorescence

We conducted immunofluorescence on ovaries following the protocol described in.[62] We stained ovaries with anti-FLAG (1:3000, M2, Sigma Aldrich, St. Louis, MO) and anti-γH2Av (1:1000, a gift from R. S. Hawley). We mounted ovaries with ProLong Gold Antifade Reagent with DAPI (Thermo Fisher Scientific, Waltham, MA). We imaged slides at 63X magnification on a Leica TCS SP8 Four Channel Spectral Confocal System. For each experiment, we used the same imaging parameters across genotypes.

We conducted immunofluorescence on embryos collected in a 0-70 minute window from *mh[mel]*, *mh[sim]*, or *mh^1* females crossed to males homozygous for *P{gcid.EGFP.cid}III.2* ([55], a gift from K. McKim). We followed the protocol described in[63] to fix and stain the embryos with anti-GFP (1:1000, Aves Labs, Tigard, OR). We mounted and imaged the embryos as described above.

Analysis of cytological data

Cell death quantification

To quantify the incidence of cell death, we mounted fixed whole ovaries with ProLong Gold

Antifade Reagent with DAPI (Thermo Fisher Scientific, Waltham, MA) and imaged at 63X

magnification on a Leica TCS SP8 Four Channel Spectral Confocal System using the tile

scanning and merging feature. We identified the number of ovarioles that contained egg

chambers with >1 condensed, signal-saturated nurse cell nuclei. We then divided this number

by the total number of ovarioles present in each ovary to determine the fraction of cell death

incidence in *mh[mel]* and *mh[sim]* ovaries.

Immunofluorescence quantification

To quantify the average fluorescence of γH2Av in *mh[mel]* and *mh[sim]* ovaries we outlined a

representative stage four egg chamber with the Freehand tool in FIJI[61]. We calculated the

fluorescent signal intensity using the polygon tool in FIJI to define the borders of the tissue. We

used the Measure tool in FIJI to calculate the mean pixels within these boundaries. We

normalized the fluorescent signal intensity of *mh[mel]*, *mh[sim], and mh[1]* to the mean intensity

signal of the *mh[mel]*. Similarly, the fluorescent signal intensity of *mh[mel],Zhr* and *mh[sim],Zhr*

Was normalized to the mean intensity signal of *mh[mel],Zhr*. Finally, the fluorescent signal

intensity of *mh[mel]*; UAS-*Top2[mel]* and *mh[sim]*; UAS-*Top2[mel]* was normalized to the mean

intensity signal of *mh[mel]*; UAS-*Top2[mel]*.

FIGURE LEGENDS

Figure 1. MH evolves adaptively to preserve female fertility. (A) Counts of synonymous and

nonsynonymous polymorphic and fixed sites within and between *D. melanogaster* and

D. simulans; χ^2 test, p = 0.04. (B) Swap strategy: the *D. melanogaster* ("*D. mel*", blue) or

D. simulans ("*D. sim*," yellow) *mh* coding sequence, codon-optimized for *D. melanogaster* and

3xFLAG-tagged, replaced the native *mh* gene on the *X* chromosome. (C) Total offspring from

mh[mel] or *mh[sim]* females crossed to wildtype (*w^{1118}*) males. (D) Frequency distribution of

embryos at increasing mitotic cycle numbers collected for 70 minutes from *mh[mel]*, *mh[sim]*,

and *mh^1* females. Dashed bars from *mh^1* females correspond to embryos undergoing mitotic

catastrophe likely triggered at the first mitosis. Solid gray bars representing *mh^1*-derived

embryos are presumed to be maternal haploid. (E) Representative images and ovary size

estimates from *mh[mel]* and *mh[sim]* females. (F) Number of mature eggs per ovary pair from

mh[mel] and *mh[sim]* females. (*t*-test: "***" = p < 0.001, scale bar = 100µm)

Figure 2. MH[sim] poisons oogenesis through a DNA damage pathway. (A) Diagram of a

Drosophila ovary (above) and a single ovariole (below) with the germline stem cells in the

germarium at the anterior (left) position and the mature eggs at the posterior position (right). The

dashed box shows the developmental stages shown in images 2B-2D (Created using

BioRender.com). (B) *mh[mel]* and *mh[sim]* ovaries stained with anti-FLAG to visualize MH

localization (left). Merged images of single nuclei (*) show no MH foci on the DNA (dashed inset

boxes). Merged images of enlarged egg chambers imaged under higher laser power reveal that

MH[sim] but not MH[mel] localizes to the peripheral follicle cells (right). (C) Incidence of cell

death captured by the fraction of ovarioles with condensed nuclei (arrowheads) in *mh[mel]* and

mh[sim] ovaries. (D) γH2Av signal in *mh[mel]* and *mh[sim]* ovaries and the quantification of

normalized fluorescent signal intensity. Note that the expected γH2Av-positive cells in the

germarium in *mh[mel]* are absent under the imaging parameters used but are indeed present,

see Figure S2D. (E) Ovary size estimates from *mh[mel];Chk2$^{-/-}$* and *mh[sim];Chk2$^{-/-}$* females. (F)

Number of mature eggs per ovary pair from *mh[mel];Chk2$^{-/-}$* and *mh[sim];Chk2$^{-/-}$* females. In

panels E and F, dotted lines correspond to *mh[mel]* and *mh[sim]* averages reported in Figure 1E

and F, respectively. (*t*-test: "***" = p < 0.001, "n.s." p > 0.05, scale bar = 25µm)

Figure 3. The 359 satellite deletion rescues *mh[sim]* genome integrity and fertility. (A) The

Zhr X chromosome lacks the 11Mb pericentromeric 359 satellite array (left). A 10-cycle PCR

distinguishes between wildtype 359 copy number and the 359-deletion (*Zhr*) and validates the

recombined *mh[mel],Zhr* and recombined *mh[sim],Zhr X* chromosomes (right). (B) γH2Av signal

in *mh[mel],Zhr* and *mh[sim],Zhr* ovaries and the quantification of normalized fluorescent signal

intensity. (C) Ovary size of *mh[mel],Zhr* and *mh[sim],Zhr* females. (D) Number of mature eggs

per ovary pair from *mh[mel],Zhr* and *mh[sim],Zhr* females. (E) Progeny counts from *mh[mel],Zhr*

and *mh[sim],Zhr* females crossed to wildtype (*w[1118]*) males. In panel B, dotted lines correspond

to *mh[mel]* and *mh[sim]* averages reported in Figure 2D. In panels C, D, and E, dotted lines

correspond to *mh[mel]* and *mh[sim]* averages reported in Figure 1E, F, and C, respectively.

(*t*-test: "n.s." $p > 0.05$, scale bar = 25μm)

Figure 4. MH[sim] interferes with Top2 processing of 359 entanglements. (A) Progeny

counts from *mh[mel];Top2[-]/+* and *mh[sim];Top2[-]/+* females crossed to wildtype (*w[1118]*) males. (B)

Progeny counts from nos-Gal4-VP16 (female germline GAL4) driven *mh[mel];*UAS-*Top2* or

*mh[sim];*UAS-*Top2* females crossed to wildtype (*w[1118]*) males. (C) γH2Av signal from ovaries of

nos-Gal4-VP16 driven *mh[mel];*UAS-*Top2* or *mh[sim];*UAS-*Top2* females and quantification of

normalized fluorescent signal intensity. (D) Model of MH[sim] interference with Top2 processing

of 359 entanglements. These entanglements threaten genome integrity and ultimately, fertility.

MH[mel], in contrast, has no measurable function in the ovaries, suggesting that it avoids

interfering with 359 processing by Top2. (E) Model of MH evolution tracking 359 satellite

proliferation. In panel A, dotted lines correspond to *mh[mel]* and *mh[sim]* averages reported in

Figure 1C. In panel C, dotted lines correspond to *mh[mel]* and *mh[sim]* averages reported in

Figure 2D. (*t*-test: "***" = $p < 0.001$, "n.s." $p > 0.05$, scale bar = 25μm)

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. MH[sim] is expressed at comparable levels to MH[mel] and does not

phenocopy *mh* null. (A) Western Blot of *mh[mel]* and *mh[sim]* ovaries probed with anti-FLAG

and anti-αTubulin. (B) Proportion of female progeny from *mh[mel]* and *mh[sim]* females. (C)

Representative images of diploid embryos from crosses between *EGFP-cid* fathers and *mh[mel]*

or *mh[sim]* mothers stained with anti-GFP. Diploid embryos are GFP-positive and haploid

embryos are GFP-negative. (D) Ovary size estimates from mh^1 homozygous and $mh^1/+$

heterozygous females. (F) Number of mature eggs per ovary pair from mh^1 homozygous and

$mh^1/+$ heterozygous females. In panel B, dotted lines correspond to *mh[mel]* and *mh[sim]*

averages reported in Figure 1C. (*t*-test: "n.s." $p > 0.05$, scale bar = 25μm)

Figure S2. MH[sim] toxicity is dose-dependent, phenotypically distinct from *mh* null, and

acts through a DNA damage pathway to block oogenesis. (A) Anti-FLAG staining of *mh[mel]*

and *mh[sim]* upon overexpression under the UAS-GAL4 system. The driver nos-GAL4-VP16

expresses the *mh* transgene in the female germline only. Corresponding chromosome *X*

(wildtype) and chromosome *3* are shown above images. (B) Progeny counts of the nos-Gal4-

VP16 driven UASp-*mh[mel]* or UASp-*mh[sim]* females crossed to wildtype (w^{1118}) males. Note

that wildtype *mh* is present in these genotypes. (C) Total offspring of *mh[mel], mh[sim],* and

mh[mel]/mh[sim] heterozygous females crossed to wildtype (w^{1118}) males. (D) *Drosophila*

ovariole (above) showing the germarium where meiotic recombination occurs. Germarium of an

mh[mel] female (below) showing programmed double stranded breaks (arrowheads) occurring

in "region 2A" of the germarium detected under higher laser power relative to images displayed

in Figure 3B. These double stranded breaks are repaired via meiotic recombination pathways.

(E) γH2Av signal in mh^1 ovaries and quantification of normalized fluorescent signal intensity. (F)

Progeny counts from *mh[mel];Chk2$^{-/-}$* and *mh[sim];Chk2$^{-/-}$* females crossed to wildtype (w^{1118})

males. Note that *mh[mel];Chk2$^{-/-}$* reduced progeny counts are due to the *Chk2$^{-/-}$* mutation[64]. In

panel E, dotted lines correspond to *mh[mel]* and *mh[sim]* averages reported in Figure 2D. (*t*-test:

"***" = $p < 0.001$, "n.s." $p > 0.05$, scale bar = 25μm)

Brand and Levine Figure S1

499   SUPPLEMENTARY TABLE LEGEND

501   **Table S1. Primers used in this study.**

Table S1. Primers used in this study.

Forward	Reverse	Purpose
GATCCAGGAAATACTCGGCC	CCAAAGACTGGTCACACTGC	Primers used to amplify the *mh* ortholog in *D. simulans*
ATCCTGGCCAACAGTGACTC		Sequencing primer for *mh* ortholog in *D. simulans*
	TTTGCCCAAAATATCGAACG	Sequencing primer for *mh* ortholog in *D. simulans*
GCAGAGCAGACGAGTCAGTG		Sequencing primer for *mh* ortholog in *D. simulans*
	TTGACCTTGTGTGAGGACGA	Sequencing primer for *mh* ortholog in *D. simulans*
ACGACGCCTCTTTTTGGTAA		Sequencing primer for *mh* ortholog in *D. simulans*
	ACATACGCAACTGGTTCGAC	Sequencing primer for *mh* ortholog in *D. simulans*
CACCTTGTTGATCGTCTCCA		Sequencing primer for *mh* ortholog in *D. simulans*
	GATCCCACTCCGGACATCTA	Sequencing primer for *mh* ortholog in *D. simulans*
TTCTCCATTTTTGTCTTGTGGA		Sequencing primer for *mh* ortholog in *D. simulans*
AAGTGTCGCGCTATTTCACC	TCACCGTCATGGTCTTTGTAGTCCAT	Reverse primer anneals in FLAG-tag. Amplifies *mh[mel]* and *mh[sim]* CRISPR mutations
AATGGATTTCGGCAAATGAG	GTCGTTGTAGGAGCCCATGT	Primers anneal outside homology arms. Amplicon spans entire *mh[mel]* and *mh[sim]* CRISPR mutation
ATGCGCAGGCTAAATTCGAT		Sequencing primer for *mh[mel]* and *mh[sim]* region
GATTACCTGTTCGCCCTGAA		Sequencing primer for *mh[mel]* and *mh[sim]* region
ACCAACATCACCGTGTACCA		Sequencing primer for *mh[mel]* and *mh[sim]* region
CATTGAGCACATCGACCTGT		Sequencing primer for *mh[mel]* region
CATCGAGAACATCGATATCT		Sequencing primer for *mh[sim]* region
CTCCCTGACCAGCAAGGAT		Sequencing primer for *mh[mel]* region
CAGCTTCACCTCCAAGGAT		Sequencing primer for *mh[sim]* region
TCACGAATTCCGCCCTCT		Sequencing primer for *mh[mel]* and *mh[sim]* region
GGCCCTGCTCATATCGTATC	AAGAACCTTACTGCGTGCAAC	Primers anneal in *mh* introns. Amplifies endogenous *mh* gene in *D. melanogaster*.
TATTCTTACATCTATGTGACC	GTTTTGAGCAGCTAATTACC	Primers amplify a repeat unit of the 359 satellite array (Rosic *et al.* 2014)

LITERATURE CITED

1. Cechova, M., Harris, R.S., Tomaszkiewicz, M., Arbeithuber, B., Chiaromonte, F., and
Makova, K.D. (2019). High satellite repeat turnover in great apes studied with short- and
long-read technologies. Mol Biol Evol *36*, 2415-2431. 10.1093/molbev/msz156.
2. Jagannathan, M., Cummings, R., and Yamashita, Y.M. (2018). A conserved function for
pericentromeric satellite DNA. Elife *7*, e34122. 10.7554/eLife.34122.
3. Jagannathan, M., Warsinger-Pepe, N., Watase, G.J., and Yamashita, Y.M. (2017).
Comparative Analysis of Satellite DNA in the Drosophila melanogaster Species
Complex. G3 (Bethesda) *7*, 693-704. 10.1534/g3.116.035352.
4. Joshi, S.S., and Meller, V.H. (2017). Satellite Repeats Identify X Chromatin for Dosage
Compensation in Drosophila melanogaster Males. Curr Biol *27*, 1393-1402.
10.1016/j.cub.2017.03.078.
5. Kipling, D., Ackford, H.E., Taylor, B.A., and Cooke, H.J. (1991). Mouse minor satellite
DNA genetically maps to the centromere and is physically linked to the proximal
telomere. Genomics *11*, 235-241. 10.1016/0888-7543(91)90128-2.
6. Rosic, S., Kohler, F., and Erhardt, S. (2014). Repetitive centromeric satellite RNA is
essential for kinetochore formation and cell division. J Cell Biol *207*, 335-349.
10.1083/jcb.201404097.
7. Round, E.K., Flowers, S.K., and Richards, E.J. (1997). Arabidopsis thaliana centromere
regions: genetic map positions and repetitive DNA structure. Genome Res *7*, 1045-1053.
10.1101/gr.7.11.1045.
8. Rudd, M.K., and Willard, H.F. (2004). Analysis of the centromeric regions of the human
genome assembly. Trends Genet *20*, 529-533. 10.1016/j.tig.2004.08.008.
9. Sproul, J.S., Khost, D.E., Eickbush, D.G., Negm, S., Wei, X., Wong, I., and Larracuente,
A.M. (2020). Dynamic Evolution of Euchromatic Satellites on the X Chromosome in
Drosophila melanogaster and the simulans Clade. Mol Biol Evol *37*, 2241-2256.
10.1093/molbev/msaa078.
10. Henikoff, S., Ahmad, K., and Malik, H.S. (2001). The centromere paradox: stable
inheritance with rapidly evolving DNA. Science *293*, 1098-1102.
10.1126/science.1062939.
11. Ferree, P.M., and Prasad, S. (2012). How can satellite DNA divergence cause
reproductive isolation? Let us count the chromosomal ways. Genet Res Int *2012*,
430136. 10.1155/2012/430136.
12. Jagannathan, M., and Yamashita, Y.M. (2021). Defective Satellite DNA Clustering into
Chromocenters Underlies Hybrid Incompatibility in Drosophila. Mol Biol Evol *38*, 4977-
4986. 10.1093/molbev/msab221.
13. Bayes, J.J., and Malik, H.S. (2009). Altered heterochromatin binding by a hybrid sterility
protein in Drosophila sibling species. Science *326*, 1538-1541.
10.1126/science.1181756.
14. Kumon, T., Ma, J., Akins, R.B., Stefanik, D., Nordgren, C.E., Kim, J., Levine, M.T., and
Lampson, M.A. (2021). Parallel pathways for recruiting effector proteins determine
centromere drive and suppression. Cell *184*, 4904-4918 e4911.
10.1016/j.cell.2021.07.037.

15.   Maheshwari, S., Tan, E.H., West, A., Franklin, F.C.H., Comai, L., and Chan, S.W.L.
545   (2015). Naturally occurring differences in CENH3 affect chromosome segregation in
546   zygotic mitosis of hybrids. PLoS Genet *11*, e1004970. 10.1371/journal.pgen.1004970.
547   16.   Ferree, P.M., and Barbash, D.A. (2009). Species-specific heterochromatin prevents
548   mitotic chromosome segregation to cause hybrid lethality in Drosophila. PLoS Biol *7*,
549   e1000234. 10.1371/journal.pbio.1000234.
550   17.   Gibeaux, R., Acker, R., Kitaoka, M., Georgiou, G., van Kruijsbergen, I., Ford, B.,
551   Marcotte, E.M., Nomura, D.K., Kwon, T., Veenstra, G.J.C., and Heald, R. (2018).
552   Paternal chromosome loss and metabolic crisis contribute to hybrid inviability in
553   Xenopus. Nature *553*, 337-341. 10.1038/nature25188.
554   18.   Lohe, A.R., Hilliker, A.J., and Roberts, P.A. (1993). Mapping simple repeated DNA
555   sequences in heterochromatin of Drosophila melanogaster. Genetics *134*, 1149-1174.
556   101093/genetics.134.4.1149.
557   19.   Brutlag, D.L. (1980). Molecular arrangement and evolution of heterochromatic DNA.
558   Annu Rev Genet *14*, 121-144. 10.1146/annurev.ge.14.120180.001005.
559   20.   de Lima, L.G., Hanlon, S.L., and Gerton, J.L. (2020). Origins and Evolutionary Patterns
560   of the 1.688 Satellite DNA Family in Drosophila Phylogeny. G3 (Bethesda) *10*, 4129-
561   4146. 10.1534/g3.120.401727.
562   21.   Abad, J.P., Agudo, M., Molina, I., Losada, A., Ripoll, P., and Villasante, A. (2000).
563   Pericentromeric regions containing 1.688 satellite DNA sequences show anti-
564   kinetochore antibody staining in prometaphase chromosomes of Drosophila
565   melanogaster. Mol Gen Genet *264*, 371-377. 10.1007/s004380000331.
566   22.   Losada, A., and Villasante, A. (1996). Autosomal location of a new subtype of 1.688
567   satellite DNA of Drosophila melanogaster. Chromosome Res *4*, 372-383.
568   10.1007/BF02257273.
569   23.   Delabaere, L., Orsi, G.A., Sapey-Triomphe, L., Horard, B., Couble, P., and Loppin, B.
570   (2014). The Spartan ortholog maternal haploid is required for paternal chromosome
571   integrity in the Drosophila zygote. Curr Biol *24*, 2281-2287. 10.1016/j.cub.2014.08.010.
572   24.   Tang, X., Cao, J., Zhang, L., Huang, Y., Zhang, Q., and Rong, Y.S. (2017). Maternal
573   Haploid, a Metalloprotease Enriched at the Largest Satellite Repeat and Essential for
574   Genome Integrity in Drosophila Embryos. Genetics *206*, 1829-1839.
575   10.1534/genetics.117.200949.
576   25.   Loppin, B., Berger, F., and Couble, P. (2001). Paternal chromosome incorporation into
577   the zygote nucleus is controlled by maternal haploid in Drosophila. Dev Biol *231*, 383-
578   396. 10.1006/dbio.2000.0152.
579   26.   McDonald, J.H., and Kreitman, M. (1991). Adaptive protein evolution at the Adh locus in
580   Drosophila. Nature *351*, 652-654. 101038/351652a0.
581   27.   Garrigan, D., Kingan, S.B., Geneva, A.J., Andolfatto, P., Clark, A.G., Thornton, K.R., and
582   Presgraves, D.C. (2012). Genome sequencing reveals complex speciation in the
583   Drosophila simulans clade. Genome Res *22*, 1499-1511. 10.1101/gr.130922.111.
584   28.   Kerr, J.F., Wyllie, A.H., and Currie, A.R. (1972). Apoptosis: a basic biological
585   phenomenon with wide-ranging implications in tissue kinetics. Br J Cancer *26*, 239-257.
586   10.1038/bjc.1972.33.

587   29.   McCall, K. (2004). Eggs over easy: cell death in the Drosophila ovary. Dev Biol *274*, 3-
588   14. 10.1016/j.ydbio.2004.07.017.

589   30.   Roos, W.P., Thomas, A.D., and Kaina, B. (2016). DNA damage and the balance
590   between survival and death in cancer biology. Nat Rev Cancer *16*, 20-33.
591   10.1038/nrc.2015.2.

592   31.   Jang, J.K., Sherizen, D.E., Bhagat, R., Manheim, E.A., and McKim, K.S. (2003).
593   Relationship of DNA double-strand breaks to synapsis in Drosophila. J Cell Sci *116*,
594   3069-3077. 10.1242/jcs.00614.

595   32.   Madigan, J.P., Chotkowski, H.L., and Glaser, R.L. (2002). DNA double-strand break-
596   induced phosphorylation of Drosophila histone variant H2Av helps prevent radiation-
597   induced apoptosis. Nucleic Acids Res *30*, 3698-3705. 10.1093/nar/gkf496.

598   33.   Bakhrat, A., Pritchett, T., Peretz, G., McCall, K., and Abdu, U. (2010). Drosophila Chk2
599   and p53 proteins induce stage-specific cell death independently during oogenesis.
600   Apoptosis *15*, 1425-1434. 10.1007/s10495-010-0539-z.

601   34.   Brodsky, M.H., Weinert, B.T., Tsang, G., Rong, Y.S., McGinnis, N.M., Golic, K.G., Rio,
602   D.C., and Rubin, G.M. (2004). Drosophila melanogaster MNK/Chk2 and p53 regulate
603   multiple DNA repair and apoptotic pathways following DNA damage. Mol Cell Biol *24*,
604   1219-1231. 10.1128/MCB.24.3.1219-1231.2004.

605   35.   Sawamura, K., Yamamoto, M.T., and Watanabe, T.K. (1993). Hybrid lethal systems in
606   the Drosophila melanogaster species complex. II. The Zygotic hybrid rescue (Zhr) gene
607   of D. melanogaster. Genetics *133*, 307-313. 10.1093/genetics.133.2.307.

608   36.   Dokshin, G.A., Davis, G.M., Sawle, A.D., Eldridge, M.D., Nicholls, P.K., Gourley, T.E.,
609   Romer, K.A., Molesworth, L.W., Tatnell, H.R., Ozturk, A.R., et al. (2020). GCNA
610   Interacts with Spartan and Topoisomerase II to Regulate Genome Stability. Dev Cell *52*,
611   53-68 e56. 10.1016/j.devcel.2019.11.006.

612   37.   Lopez-Mosqueda, J., Maddi, K., Prgomet, S., Kalayil, S., Marinovic-Terzic, I., Terzic, J.,
613   and Dikic, I. (2016). SPRTN is a mammalian DNA-binding metalloprotease that resolves
614   DNA-protein crosslinks. Elife *5*, e21491. 10.7554/eLife.21491.

615   38.   Vaz, B., Popovic, M., Newman, J.A., Fielden, J., Aitkenhead, H., Halder, S., Singh, A.N.,
616   Vendrell, I., Fischer, R., Torrecilla, I., et al. (2016). Metalloprotease SPRTN/DVC1
617   Orchestrates Replication-Coupled DNA-Protein Crosslink Repair. Mol Cell *64*, 704-719.
618   10.1016/j.molcel.2016.09.032.

619   39.   Gale, K.C., and Osheroff, N. (1992). Intrinsic intermolecular DNA ligation activity of
620   eukaryotic topoisomerase II. Potential roles in recombination. J Biol Chem *267*, 12090-
621   12097.10.1016/S0021-9258(19)49810-4.

622   40.   Morimoto, S., Tsuda, M., Bunch, H., Sasanuma, H., Austin, C., and Takeda, S. (2019).
623   Type II DNA Topoisomerases Cause Spontaneous Double-Strand Breaks in Genomic
624   DNA. Genes *10*, 868. 10.3390/genes10110868.

625   41.   Pommier, Y., Sun, Y., Huang, S.N., and Nitiss, J.L. (2016). Roles of eukaryotic
626   topoisomerases in transcription, replication and genomic stability. Nat Rev Mol Cell Biol
627   *17*, 703-721. 10.1038/nrm.2016.111.

628   42.   Kas, E., and Laemmli, U.K. (1992). In vivo topoisomerase II cleavage of the Drosophila
629   histone and satellite III repeats: DNA sequence and structural characteristics. EMBO J
630   *11*, 705-716. 10.1002/j.1460-2075.1992.tb05103.x.

43. Hughes, S.E., and Hawley, R.S. (2014). Topoisomerase II is required for the proper
separation of heterochromatic regions during Drosophila melanogaster female meiosis.
PLoS Genet *10*, e1004650. 10.1371/journal.pgen.1004650.
44. Ferree, P.M., Gomez, K., Rominger, P., Howard, D., Kornfeld, H., and Barbash, D.A.
(2014). Heterochromatin position effects on circularized sex chromosomes cause filicidal
embryonic lethality in Drosophila melanogaster. Genetics *196*, 1001-1005.
10.1534/genetics.113.161075.
45. Chmátal, L., Gabriel, S.I., Mitsainas, G.P., Martínez-Vargas, J., Ventura, J., Searle, J.B.,
Schultz, R.M., and Lampson, M.A. (2014). Centromere strength provides the cell
biological basis for meiotic drive and karyotype evolution in mice. Curr Biol *24*, 2295-
2300. 10.1016/j.cub.2014.08.017.
46. Iwata-Otsubo, A., Dawicki-McKenna, J.M., Akera, T., Falk, S.J., Chmátal, L., Yang, K.,
Sullivan, B.A., Schultz, R.M., Lampson, M.A., and Black, B.E. (2017). Expanded Satellite
Repeats Amplify a Discrete CENP-A Nucleosome Assembly Site on Chromosomes that
Drive in Female Meiosis. Curr Biol *27*, 2365-2373.e2368. 10.1016/j.cub.2017.06.069.
47. Malik, H.S., and Henikoff, S. (2001). Adaptive evolution of Cid, a centromere-specific
histone in Drosophila. Genetics *157*, 1293-1298. 10.1093/genetics/157.3.1293.
48. Malik, H.S., and Henikoff, S. (2002). Conflict begets complexity: the evolution of
centromeres. Curr Opin Genet Dev *12*, 711-718. 10.1016/s0959-437x(02)00351-9.
49. Vermaak, D., Bayes, J.J., and Malik, H.S. (2009). A surrogate approach to study the
evolution of noncoding DNA elements that organize eukaryotic genomes. J Hered *100*,
624-636. 10.1093/jhered/esp063.
50. Carracedo, M.C., Asenjo, A., and Casares, P. (2000). Location of Shfr, a new gene that
rescues hybrid female viability in crosses between Drosophila simulans females and D.
melanogaster males. Heredity *84*, 630-638. 10.1046/j.1365-2540.2000.00658.x.
51. Gerard, P.R., and Presgraves, D.C. (2012). Abundant genetic variability in Drosophila
simulans for hybrid female lethality in interspecific crosses to Drosophila melanogaster.
Genet Res *94*, 1-7. 10.1017/S0016672312000031.
52. Orr, H.A. (1996). The unexpected recovery of hybrids in a Drosophila species cross: a
genetic analysis. Genet Res *67*, 11-18. 10.1017/s0016672300033437.
53. Sawamura, K., Taira, T., and Watanabe, T.K. (1993). Hybrid lethal systems in the
Drosophila melanogaster species complex. I. The maternal hybrid rescue (mhr) gene of
Drosophila simulans. Genetics *133*, 299-305. 10.1093/genetics.133.2.307.
54. Sawamura, K., and Yamamoto, M.T. (1993). Cytogenetical localization of Zygotic hybrid
rescue (Zhr), a Drosophila melanogaster gene that rescues interspecific hybrids from
embryonic lethality. Mol Gen Genet *239*, 441-449. 10.1007/BF00276943.
55. Schuh, M., Lehner, C.F., and Heidmann, S. (2007). Incorporation of Drosophila
CID/CENP-A and CENP-C into centromeres during early embryonic anaphase. Curr Biol
17, 237-243. 10.1016/j.cub.2006.11.051.
56. Pool, J.E., Corbett-Detig, R.B., Sugino, R.P., Stevens, K.A., Cardeno, C.M., Crepeau,
M.W., Duchen, P., Emerson, J.J., Saelao, P., Begun, D.J., and Langley, C.H. (2012).
Population Genomics of sub-saharan Drosophila melanogaster: African diversity and
non-African admixture. PLoS Genet *8*, e1003080. 10.1371/journal.pgen.1003080.

57. Rogers, R.L., Cridland, J.M., Shao, L., Hu, T.T., Andolfatto, P., and Thornton, K.R.
(2014). Landscape of standing variation for tandem duplications in Drosophila yakuba
and Drosophila simulans. Mol Biol Evol *31*, 1750-1766. 10.1093/molbev/msu124.
58. Chakraborty, M., Chang, C.H., Khost, D.E., Vedanayagam, J., Adrion, J.R., Liao, Y.,
Montooth, K.L., Meiklejohn, C.D., Larracuente, A.M., and Emerson, J.J. (2021).
Evolution of genome structure in the Drosophila simulans species complex. Genome
Res *31*, 380-396. 10.1101/gr.263442.120.
59. Venken, K.J., He, Y., Hoskins, R.A., and Bellen, H.J. (2006). P[acman]: a BAC
transgenic platform for targeted insertion of large DNA fragments in D. melanogaster.
Science *314*, 1747-1751. 10.1126/science.1134426.
60. Hohl, A.M., Thompson, M., Soshnev, A.A., Wu, J., Morris, J., Hsieh, T.S., Wu, C.T., and
Geyer, P.K. (2012). Restoration of topoisomerase 2 function by complementation of
defective monomers in Drosophila. Genetics *192*, 843-856.
10.1534/genetics.112.144006.
61. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods *9*, 676-682. 10.1038/nmeth.2019.
62. McKim, K.S., Joyce, E.F., and Jang, J.K. (2009). Cytological analysis of meiosis in fixed
Drosophila ovaries. Methods Mol Biol *558*, 197-216. 10.1007/978-1-60761-103-5_12.
63. Levine, M.T., Vander Wende, H.M., and Malik, H.S. (2015). Mitotic fidelity requires
transgenerational action of a testis-restricted HP1. Elife *4*, e07378. 10.7554/eLife.07378.
64. Xu, J., and Du, W. (2003). Drosophila chk2 plays an important role in a mitotic
checkpoint in syncytial embryos. FEBS Lett *545*, 209-212. 10.1016/s0014-
5793(03)00536-2.