Enhanced negative response to stress following morphine administration

increases wanting of social reward

Claudia Massaccesi [1*]

Matthaeus Willeit [2]

Boris B. Quednow [3,4]

Urs M. Nater [1]

Claus Lamm [5]

Daniel Mueller [6]

Giorgia Silani [1*]

[1] Department of Clinical and Health Psychology, Faculty of Psychology, University of Vienna,

Vienna, Austria

[2] Department of Psychiatry and Psychotherapy, Medical University of Vienna, Vienna, Austria

[3] Experimental and Clinical Pharmacopsychology, Department of Psychiatry, Psychotherapy and

Psychosomatics, Psychiatric Hospital of the University of Zurich, Zurich, Switzerland

[4] Neuroscience Center Zurich, University of Zurich and Swiss Federal Institute of Technology

Zurich, Zurich, Switzerland.

[5] Department of Cognition, Emotion, and Methods in Psychology, University of Vienna, Vienna,

Austria

[6] Institute for Clinical Chemistry, University Hospital Zurich, Zurich, Switzerland

*Correspondence

Claudia Massaccesi: claudia.massaccesi@univie.ac.at

Giorgia Silani: giorgia.silani@univie.ac.at

Abstract

Animal research suggests a central role of the μ-opioid receptor (MOR) system in mediating contact seeking and the stress-buffering function of social touch. However, the human neurochemistry of social motivation in aversive situations is still poorly understood.

In a randomized, double-blind, between-subject design, healthy female volunteers (N = 80) received either 10 mg of the μ-opioid agonist morphine sulfate or a placebo. Following psychosocial stress induction, participants engaged in a social reward task, in which the motivation (subjective ratings of wanting and physical effort) to obtain skin-to-skin social touch and the hedonic reactions (subjective ratings of liking and facial electromyography) elicited by it were assessed.

Morphine administration prevented the increase of salivary cortisol, usually observed in response to acute stress exposure. The dampened physiological reaction to the psychosocial stress was associated with increased negative mood and subsequent higher subjective wanting of the most pleasurable touch. Furthermore, participants administered with morphine displayed greater activity of the corrugator muscle during reward anticipation, possibly tracking enhanced attention toward the social stimuli.

Overall, the results provide novel evidence on the effect of exogenous opioids administration on the reactions to psychosocial stress and point to a state-dependent regulation of social motivation.

Introduction

Social behaviors such as bonding and affiliation are crucial for the survival and wellbeing of many species. By providing fundamental benefits (e.g., promoting safety and enhancing stress resilience) and by generating comfort and pleasure, social stimuli (e.g., social contact) gain rewarding value, inducing approach motivation. Inability to form and maintain social bonds contributes to a range of psychiatric and physical pathologies (e.g., Cacioppo & Cacioppo, 2014; Uchino, 2006), therefore highlighting the importance to better understand the neurobiological basis of social motivation.

The Brain Opioid Theory of Social Attachment (BOTSA), originally inspired by the analogies between opioid addiction and social dependence, pinpoint the μ-opioid receptor (MOR) system as a key mediator of bonding and affiliation (Panksepp, 1998; Panksepp et al., 1980). Specifically, BOTSA predicts that social isolation results in reduced levels of basal opioid levels, which in turn motivate social contact seeking, and that rewarding social interactions elicit release of endogenous opioids, associated with euphoria and contentment. The two-fold role of the endogenous opioid system in reward and pain (Leknes & Tracey, 2008) therefore seems to apply also to experiences of social nature. In contexts of comfort, the activity of the MOR system during social contact mediates the rewarding properties, and associated pleasure, of social stimuli, positively reinforcing exploration and affiliation. In contexts of distress, the release of endogenous opioids via social contact reduces pain and negative affect, negatively reinforcing social behaviors (Loseth et al., 2014).

In line with this model, animal studies have shown that μ-opioids are released following physical social contact, such as grooming and playing (Keverne et al., 1989; Panksepp & Bishop, 1981; Vanderschuren et al., 1995). Pharmacological studies on separation-distress in animals further

confirmed BOTSA predictions, showing that administration of MOR agonists reduces, and of

MOR antagonists increases distress responses and motivation for social contact, such as grooming

(Keverne et al., 1989; Martel et al., 1995; Panksepp et al., 1980; Fabre-Nys et al., 1982; Meller et

al., 1980; for a review see Machin & Dunbar, 2011; see Loseth et al., 2014 for a model accounting

for divergent findings in different species).

In the last decade, preliminary confirmatory evidence on the μ-opioid mediation of the rewarding

properties of social stimuli in humans have been provided. These studies mainly focused on two

fundamental components: wanting, namely the motivational drive to obtain the social stimulus,

and liking, namely the pleasure elicited once it is received. Results showed that pharmacological

challenge of the MOR system affects wanting and/or liking of different kind of social stimuli, such

as erotic pictures (Buchel et al., 2018), attractive faces (Chelnokova et al., 2014), and social touch

(Korb, Götzendorfer, et al., 2020; but see also Løseth et al., 2019). MOR activation has been also

associated with social laughter (Manninen et al., 2017), and MOR blockade was shown to reduce

feelings of social connection (Inagaki et al., 2015, 2016, 2020), as well as of interpersonal

closeness and social reward expectation (Tchalova & MacDonald, 2020). Using PET, Hsu et al.

(2013) observed MOR activation in response to both social rejection and social acceptance. In the

first case, MOR activity was correlated with reduced negative affect, while in the latter it was

associated with increased desire for social interaction. These findings are consistent with a

"protective" and an "affiliative" role of the MOR system during social distress and social comfort,

respectively. Regarding the role of the MOR system in stress regulation, a recent study showed

that MOR agonist administration reduces individuals' physiological stress response (as measured

by cortisol), as well as the perceived difficulty of the stress task (Bershad et al., 2018). However,

direct evidence of an involvement of the MOR system in downregulating stress and how this

affects social contact seeking, as shown in non-human animal species, is still lacking.

Here, we aimed at filling this knowledge gap by investigating the effect of MOR agonist

administration (morphine) on social motivation and social pleasure following stress exposure.

Using a double-blind, placebo-controlled, randomized, between-subjects design, female

participants (N = 80) were orally administered with either 10 mg morphine sulfate (a highly

selective μ-opioid agonist) or placebo. Following a psychosocial stress induction procedure, the

motivation to obtain social touch (wanting) and the pleasure elicited by receiving it (liking) were

assessed (see Figure 1 for a detailed description). In order to enhance the translational value of the

study, the following steps were taken: i) as in previous animal research, which entailed social

separation, a stressor of social nature (via the Trier Social Stress Test, Kirschbaum et al., 1993)

was employed; ii) in addition to self-reports of wanting and liking, we assessed real physical effort

(via a hand-dynamometer) and hedonic facial reactions (via facial electromyography of corrugator

and zygomaticus muscles), approximating the motivational and hedonic (i.e., wanting and liking)

measures used in animal studies; iii) to parallel grooming in animals, skin-to-skin touch

administered at different speeds (6, 21, 27 cm/s, corresponding to high, low, and very low reward

levels) was employed as a social reward.

Based on the existing human and animal literature, we hypothesized that participants administered

with morphine, compared to placebo, would show reduced physiological and subjective responses

to stress, and that this would result in decreased social motivation. As previous studies had shown

an effect of stress on wanting rather than on liking of touch (Massaccesi et al., 2021) or food (e.g.

Pool et al., 2015) reward, we did not expect changes in the hedonic reactions to social contact.

Given that MOR manipulations have been shown to induce the strongest effects on the best reward

option available (Chelnokova et al., 2014; Eikemo et al., 2016), and considering the stress

buffering function of C tactile (CT)-optimal touch (i.e., touch with stroking speed of 1 to 10 cm/s;

Morrison, 2016), we expected the predicted effects to be stronger for the most valuable social

reward (touch at 6 cm/s).

Figure 1. **Overview of the experimental session, set-up, and trial structure of the Social**
Reward task. (A) Overview of the experimental session. T1 – T7 represent the time points at
which subjective and/or physiological measures were obtained. Blue drops indicate saliva sample
collection. (B) Set-up of the Social Reward task (based on Korb, Götzendorfer, et al., 2020;
Massaccesi et al., 2021). Participants were seated in front of the monitor, holding the hand-
dynamometer in the right hand. The left arm was rested on a cushion, next to a keyboard used to
express judgements during the task. The toucher was seated on the other side of a curtain used to
limit the participants' field of view to the monitor. Touch was administered on the participants'
left forearm using the index and middle finger, at 3 stroking speeds corresponding to 3 levels of
reward (high = 6 cm/s, low = 21 cm/s, very low = 27 cm/s). (C) Trial structure of the Social Reward
task: i) announcement of the best attainable social reward (high or low); ii) rating of subjective
wanting; iii) effort task requiring to squeeze a hand-dynamometer to obtain the announced reward.
Participants' force exerted was indicated in real-time by a vertical bar filling in red. The top of the
displayed vertical bar corresponded to their previously measured maximum voluntary contraction
(MVC); iv) announcement of the reward attained (high, low, or – if insufficient effort had been
exerted – very low); v) skin-to-skin touch administration; vi) relax phase; vii) rating of subjective
liking. Facial electromyography (fEMG) was recorded during the whole task and analyzed in
reward anticipation (Anticipation Pre-Effort and Post-Effort) and consumption (Delivery and
Relax). TMT, Trail Making Test; DSST, Digit Symbol Substitution Test; TSST, Trier Social Stress
Test.

Results

Drug blinding

After completing the session, 50% of the participants correctly guessed they received morphine, indicating successful blinding. Overall, 55% of the total sample believed to have been administered with placebo, 29% with morphine, and 16% with naltrexone[1].

Effects of morphine on cognitive functions and drug side-effects

There were no significant differences in the Digit Symbol Substitution Test (DSST) and Trail Making Test (TMT) scores across groups, indicating that drug administration did not have negative effects on attention, psychomotor and processing speed, and visuo-perceptual functions (See Table S1 in *Supplementary Material*). Morphine administration significantly increased self-reported weakness (MORPH vs. PLB at T2: $t(69.9) = 2.64$, $p = 0.01$, and at T7: $t(55.96) = 2.19$, $p = 0.03$) and dry mouth (MORPH vs. PLB at T7: $t(57.70) = 2.56$, $p = 0.01$). For all side-effect measures, group average scores were generally low, and no side-effect was on average rated as moderate or strong (see Figure S2 in *Supplementary Material*).

Effects of morphine on stress response

Subjective measures

Morphine administration resulted in significantly higher scores of the Profile of Mood States (POMS) subscale "Displeasure" after stress induction (Drug*Time: $F(4,312) = 2.97$, FDR $p = 0.03$; MORPH vs. PLB at T5: FDR $p < 0.01$) (Figure 2A). A similar pattern, though not reaching statistical significance, was observed for negative mood (Drug*Time: $F(6,468) = 2.16$, FDR $p = 0.068$) (Figure 2A). No significant group differences were observed for positive mood or for the

---

[1] To reduce drug-related expectancy, participants were told they might receive an opioid agonist (morphine), antagonist (naltrexone) or placebo (but they could receive only morphine or placebo).

POMS subscales "Depression", "Vigor", and "Fatigue" (all FDR $p > .15$). Furthermore, no drug

effects were observed in anticipatory stress (PASA primary and secondary appraisal at T4; both t

< 0.60, $p > 0.55$), nor in the participants' satisfaction towards their performance expressed after

TSST completion (T5; $t(75.2) = -0.63$, $p = 0.53$).

Physiological measures

As physiological stress biomarkers, salivary cortisol (hypothalamic–pituitary–adrenal [HPA] axis

activity), salivary alpha-amylase and heart rate (autonomic nervous system [ANS] activity) had

been assessed at different time-points during the session (see Figure 1).

Morphine administration suppressed cortisol response to the TSST (Drug*Time: $F(5,375) = 7.68$,

FDR $p < 0.001$, Figure 2B). Specifically, the morphine group showed reduced salivary cortisol

compared to the placebo group at T2 (FDR $p = 0.079$), T3 (FDR $p < 0.01$), T5 (FDR $p < 0.001$),

T6 (FDR $p < 0.001$) and T7 (FDR $p < 0.001$) (Figure 2A). No drug effects were observed in

salivary alpha-amylase (all $F < 0.38$, all FDR $p > 0.62$) or in heart rate (all $F < 2.83$, all FDR $p >$

0.09) (Figure 2B).

A. Subjective response to stress

B. Physiological response to stress

Figure 2. **Effects of morphine administration on subjective and physiological stress responses.** (A) Morphine administration increased the subjective negative response to stress, as shown by elevated negative mood (measured via visual analog scales) and higher scores at the "Displeasure" subscale (POMS) immediately after the Trier Social Stress Test (TSST). No significant drug effects were observed in positive mood or in any other POMS subscales (depression, vigor, fatigue). (B) Morphine administration suppressed the hypothalamic–pituitary–adrenal (HPA) axis activity, as shown by blunted salivary cortisol before and after stress induction. No significant drug effects were observed in the autonomic nervous system (ANS) response to stress, assessed via salivary alpha-amylase (sAA) and heart rate. Grey bars represent the TSST time window (anticipation, speech, and arithmetic task); error bars represent standard error of the mean; asterisks indicate significant differences between drug groups (* $p < .05$, *** $p < .001$).

Correlations between physiological and subjective measures of stress

Given the observed opposite effect of the drug on cortisol and mood, we conducted a correlation analysis to investigate the association between physiological and subjective measures of stress. To this aim, we first computed the area under the curve in respect to increase ((AUCi, from T3 to T7; Pruessner et al., 2003) of the cortisol levels and of the negative mood ratings (VAS and POMS). Results showed a negative correlation between salivary cortisol and negative mood (VAS; r_s = -.36, *p bonferroni* = .048) as well as between cortisol and the Displeasure subscale of POMS (r_s = -.37, *p bonferroni* = .044), suggesting an inverse relationship between HPA axis and mood responses to stress.

Effects of morphine on social reward processing

Using a recently developed Social Reward task (Korb, Götzendorfer, et al., 2020; Massaccesi et al., 2021), we assessed subjective wanting and liking of social reward in different levels. Participants were announced with either high (6 cm/s, CT-optimal touch[2]) or low (21 cm/s, non-CT-optimal touch) social reward and rated their wanting of the stimulus. Based on an effort task, they could then receive the announced or the very low (27 cm/s, non-CT-optimal touch) social reward and had to rate their liking of the received stimulus.

We first assessed whether the number of high, low and very low rewards obtained during the task was similar for the two groups. A linear model on number of trials was fitted, including Drug (MORPH, PLB) and Reward Level (high, low, and very low). We observed only a main effect of Reward Level ($F(2,369.5) = 11.28$, $p < 0.001$), indicating that participants obtained more often high rewards as compared to low and very low rewards. We also tested for group differences in

---

[2] Touch administered at a velocity falling into the range of 1 to 10 cm/s activates CT afferents, a special class of nerve fibers found in human hairy skin involved in signaling the affective aspects of social touch (Morrison et al., 2010), and was shown to elicit greater pleasure, compared to touch administered at faster velocities.

the average maximum voluntary contraction (MVC), assessed before and after the Social Reward

task and used to calibrate the effort task. No significant differences emerged (all $t < -0.84$, all

$p > 0.40$), indicating no drug effects on participants' grip force at rest. Last, we tested for group

differences in the baseline activity of the corrugator and zygomaticus muscles. No significant

differences emerged (all $t < 1.27$, all $p > 0.21$).

Effects of morphine on subjective ratings of wanting and liking

To investigate the effects of morphine administration on subjective wanting and liking of social

touch under stress, we fitted two linear mixed-effects models (LMM) on the ratings of wanting

and liking, including Drug (MORPH, PLB) and Reward Level (high, low for wanting ratings, and

high, low, very low for liking ratings) as fixed effects, and by-subject random intercepts and slopes

for Reward Level.

Ratings of Wanting. Participants administered morphine expressed significantly greater wanting

of the high social reward (CT-optimal touch) compared to the placebo group (Drug*Reward Level:

$F(1,78) = 10.56$, FDR $p = 0.003$; high social reward MORPH vs. PLB: FDR $p = 0.035$) (Figure

3A). No drug effects emerged for the low social reward (FDR $p = 0.72$).

Ratings of Liking. Participants in both groups expressed greater liking for high compared to low

social rewards, which in turn were more liked than the very low social rewards (Reward Level:

$F(2,72.9) = 20.93$, FDR $p < 0.001$; high vs. low vs. very low: all FDR $p < 0.001$). Further,

participants administered morphine generally liked the interpersonal touch (across all reward

levels) more compared to the placebo group, but the effect did not reach significance (Drug:

$F(1,78) = 4.23$, FDR $p = 0.065$) (Figure 3B).

Effects of morphine on force exerted and hedonic facial reactions

We further assessed wanting of social rewards in terms of force exerted to obtain the tactile stimuli,

as well as hedonic facial reactions during reward anticipation (Anticipation Pre- and Post-Effort

phases). Hedonic facial reactions during and after consumption (Delivery and Relax phases) of the

social reward were employed as a measure of liking. Hedonic facial reactions were assessed via

facial electromyography (fEMG) of the corrugator (frowning) and zygomaticus (smiling) muscles.

Force exerted. Participants overall exerted greater force to obtain the high compared to the low

social reward (Reward Level: $F(1,78) = 14.23$, FDR $p < 0.001$). We also observed greater

difference in terms of force exerted for the high compared to the low reward in the morphine than

in the placebo group, but the effect did not reach significance (Drug*Reward Level interaction:

$F(1,78) = 4.13$, FDR $p = 0.068$) (Figure 3C).

Corrugator (frowning). During the announcement of the highest attainable social reward

(Anticipation Pre-Effort), we observed greater corrugator activity following morphine

administration, compared to placebo (Drug: $F(1,791.3) = 6.15$, FDR $p = 0.036$) (Figure 3D).

Opposite to previous evidence (e.g., Korb, Götzendorfer, et al., 2020; Korb, Massaccesi, et al.,

2020; Ree et al., 2020), in this phase greater frowning was associated with greater subjective

wanting in both groups (Wanting: $F(1,1203) = 7.94$, FDR $p = 0.036$) (Figure 3D). No significant

effects were observed in other phases of the task.

Zygomaticus (smiling). During the announcement of the obtained social reward (Anticipation Post-

Effort), activity of the zygomaticus was positively associated with subjective wanting of the social

reward (Wanting: $F(1,73.2) = 10.31$, FDR $p < 0.01$) (Figure 3E). No significant effects of drug

were observed in the activity of this muscle.

Figure 3. **Effects of morphine administration on social reward processing after stress exposure.** Morphine administration resulted in (A) significantly greater wanting for the high social reward (6 cm/s, CT-optimal touch), (B) greater liking of the social stimuli regardless of reward level (marginally significant drug main effect), and (C) greater differentiation of the high and low reward levels in terms of effort exerted (marginally significant drug by reward level interaction effect), compared to placebo. Regarding facial hedonic reactions, (D) during the announcement of the highest attainable reward (Anticipation Pre-Effort), the morphine group showed greater corrugator (CS) activity, which was generally positively associated with subjective wanting. (E) During the announcement of the gained reward (Anticipation Post-Effort), greater zygomaticus (ZM) activity was associated with greater subjective wanting, but no significant drug effects were observed for this muscle. Error bars represent standard error of the mean; black dots represent group means; colored dots represent individual means; asterisks indicate significant differences between conditions (* $p < .05$). VAS, Visual Analogue Scale; MORPH, morphine group; PLB, placebo group.

Discussion

Social motivation is a powerful force guiding behavior, as social rewards (e.g., social contact, bonding, affiliation) are fundamental to the individual's physical and psychological well-being. Despite the important role of social contact in stress resilience, the neurochemical mechanisms underlying social contact seeking following stress exposure in humans are still poorly understood. In this study, we pharmacologically challenged the μ-opioid receptor (MOR) system to investigate its role in the regulation of the motivational and hedonic components of social reward processing following stress induction. To parallel previous animal research, participants were exposed to a stressor of social nature and interpersonal touch was used as a social reward. Further, force exerted to obtain the reward as well as hedonic facial reactions during its anticipation and consumption were assessed, together with subjective ratings of wanting and liking. Following the enhancement of the MOR system activity via administration of its agonist morphine, we observed suppression of the HPA axis activity (as indicated by a reduced cortisol response) and increased negative affect in response to psychosocial stress. Notably, this increased negative response to stress following morphine administration was associated with enhanced motivation for the most pleasurable social reward.

μ-opioid agonist morphine dampens cortisol and increases negative affect in response to stress

In this study, single administration of the μ-opioid agonist morphine prior to TSST exposure resulted in blunted salivary cortisol response, indicating suppression of the HPA axis reactivity to stress. This is in line with previous animal and human evidence indicating an inhibitory role of μ-opioids on cortisol release following stress exposure (Pechnick, 1993). Recently, two studies investigated the effects of partial (buprenorphine) and full (hydromorphone) MOR agonists on psychosocial stress, induced via TSST (Bershad et al., 2016, 2018). Akin to the present findings,

reduced cortisol responses to stress were observed. Buprenorphine and hydromorphone also

reduced the perceived threat and the appraisal of how challenging the participants found the stress

task, respectively. While the authors interpret the findings on cortisol and stress appraisal as

indicators of a reduced stress response, no mood-buffering effects of the drug were observed. This

is in contrast with the current study, where we find that the dampened cortisol response is

accompanied by an enhanced aversive reaction to stress, as shown by higher negative affect

following morphine compared to placebo administration. Human cortisol permeates the blood-

brain barrier to feedback the central nervous system, reducing HPA axis activity and promoting

emotion regulation (Het et al., 2012; McEwen, 1998). The observed inverse relationship between

the HPA axis activity and negative affect possibly indicates a disruption of this feedback loop by

morphine administration and is in line with a mood-buffering effect of cortisol (Het et al., 2012;

Het & Wolf, 2007). Accordingly, it was recently shown that pharmacological HPA axis

suppression, by means of dexamethasone administration, blunts the cortisol response to stress and

increases negative mood, especially in women (Ali et al., 2017, 2020).

Our results are opposite to our *a priori* hypothesis based on previous evidence from animal studies

on separation distress indicating a reduction of stress indices, such as distress vocalizations,

following MOR agonist administration (Panksepp et al., 1980). The paradoxical morphine effect

of enhanced aversive stress reaction observed in the current study vs. the previous animal literature

may be explained by experimental differences, such as route and timing of administration of the

opioid compounds. For instance, in animal research morphine is typically delivered intravenously

after stress induction, possibly allowing the system to prepare to face the stressor via an initial

mounting of the physiological stress response. In this study, morphine was administered orally to

minimize the invasiveness of the administration procedure. Unlike intravenous administration,

per-oral morphine has a slow pharmacokinetic profile and requires an average time of 60 min to

reach the peak blood concentration. As the subjective response to acute stress, especially its effect

on mood, typically lasts only for short periods of time after laboratory stress induction, orally

administering the drug and waiting for it to reach peak concentration after the TSST would not

have been feasible. For these reasons here the compound was administered prior to stress exposure.

However, the resulting suppression of the HPA axis activity before the beginning of the stress

induction may have led to the observed difference in the mood response. Accordingly, the raising

of cortisol levels, via cortisol administration, prior to stress induction has been shown to have a

protective role in women, lowering the negative subjective reaction to stress (Het & Wolf, 2007).

Morphine-induced increased negative reaction to stress is associated with enhanced social
motivation

BOTSA posits that when facing stressors, individuals seek physical social contact with others in

order to down-regulate the negative state and re-store comfort, via endogenous release of beta-

endorphins, unless opioids are externally provided. Contrary to our initial hypotheses, morphine

increased, rather than reduced, participants' negative reactions to the TSST, and such increased

stress responses were accompanied by enhanced social motivation. Specifically, we observed

greater subjective wanting of the most pleasurable social reward (CT-optimal touch) following

morphine administration, compared to placebo. This enhanced subjective wanting was also

accompanied by greater differentiation of the high and low social rewards (CT-optimal and non-

CT-optimal touch) in terms of effort exerted to obtain them in the morphine group compared to

placebo, but the effect did not reach statistical significance.

This pattern of results is consistent with previous findings from our group indicating that, as

already shown for other kinds of reward like food and money (e.g. Kumar et al., 2014; Lewis et

al., 2014; Pool et al., 2015), stress increases wanting of rewards of social nature, such as

interpersonal touch (Massaccesi et al., 2021). Differently from the present findings, in Massaccesi

et al. (2021) the effect of enhanced wanting for the social reward was independent from reward

magnitude. In other words, participants under stress expressed higher wanting ratings for all types

of social touch compared to non-stressed individuals, regardless of reward magnitude. Here we

observed an effect of morphine and stress on the most pleasurable touch only (high reward level),

characterized by slow stroking velocity (CT-optimal touch).

This potentiation of wanting for the most pleasurable touch may be due to a specific influence of

the enhanced MOR system activity. Indeed, previous research has shown that pharmacological

challenges of the MOR system have strongest effects on rewards of greatest magnitude (e.g.,

Chelnokova et al., 2014; Eikemo et al., 2016). Therefore, it can be hypothesized that while stress

alone may result in increased motivation for all available rewarding stimuli, an additional

potentiation of motivation for the most valuable stimulus is seen for the best reward due to MOR

stimulation. This would align well with the observation that participants administered with

morphine also exerted more effort in order to obtain the high reward, as compared to the low

reward, and that this difference in force exerted was greater than in the placebo group (but with

FDR $p = 0.068$ did not reach the statistical significance). Alternatively, a reason for the partially

different result pattern might be that, differently from the current study, in Massaccesi et al. (2021)

participants were allowed to individually rank the three types of touch as high, low and very low.

The high reward was therefore not necessarily the CT-optimal touch. This could have masked a

possible impact of the type of touch. While Massaccesi et al. 2021 included different state

manipulations, this study focused on the effects of stress. Since slow CT-optimal touch is more

likely to convey affiliative intentions such as social support compared to faster touch (Kirsch et

al., 2017), here the slow CT-optimal touch was kept as fixed high reward to avoid possible

confounding effects related to the speed of stroking.

During reward anticipation (announcement of the best attainable reward), the increased subjective

wanting of social touch in the morphine group was accompanied by greater activity of the

corrugator muscle. Previous research indicates that individuals relax the corrugator during

anticipation and consumption of rewarding stimuli, showing an inverse relationship between

corrugator activity and reward wanting/liking (Korb, Götzendorfer, et al., 2020; Korb, Massaccesi,

et al., 2020; Massaccesi et al., 2021; Mayo et al., 2018; Ree et al., 2020; Sato et al., 2020). Here,

in contrast, we observed greater frowning for most wanted stimuli. Activity of the corrugator is

commonly associated with emotional processing, but it has been shown to be related also to

processes without emotional content, such as mental and physical effort, alertness, perceptual

attention, task engagement and difficulty (Cacioppo et al., 1985; Cohen et al., 1992; de Morree &

Marcora, 2010; Van Boxtel & Jessurun, 1993; Waterink & van Boxtel, 1994). As individuals are

more attentive to highly rewarding stimuli (e.g. Hickey et al., 2010; Hickey & van Zoest, 2012;

Theeuwes & Belopolsky, 2012) and, since morphine - at a similar dose - has already been shown

to enhance attentional processes such as set-shifting (Quednow et al., 2008), it is possible that the

observed pattern in the corrugator activity reflects enhanced attention/preparation towards the most

wanted stimuli. Together with the higher subjective of wanting, the greater saliency of the

anticipated social rewards, reflected in greater frowning, suggests an enhancement of social

motivation following stress exposure in the morphine group, compared to placebo.

Based on previous evidence, we predicted that the effects of morphine administration on social

touch processing under stress would be specific for the motivational component, without affecting

the hedonic aspect. Accordingly, we did not observe a significant effect of drug on the ratings of

liking or in the facial hedonic reactions during the reward consumption phase. However, we

nevertheless observed that the ratings of liking were generally higher in the morphine group

compared to the placebo group (FDR $p = 0.065$), and we therefore refrain to exclude an effect of

the manipulation on social pleasure.

Study limitations

Some limitations of the study should be considered. First, while the use of a within-subject design

is usually preferable in pharmacology, in this study we employed a between-subjects design. The

choice was mainly motivated by the fact that repeated exposure to the TSST, especially with a

short time interval, can lead to habituation of the stress response, resulting in low test re-test

reliability (e.g., Gerra et al., 2001; Petrowski et al., 2012) and thus outweighing by far the possibly

higher statistical power a within-subject design might have had in principle. Second, despite our

efforts in enhancing the social properties of the administered touch (e.g., skin-to-skin

administration), the absence of a social relationship between the toucher and the participant, as

well as other methodological aspects (e.g., use of a curtain separating participant and toucher),

might have affected our results. Nevertheless, in humans an involvement of the MOR system in

bond formation, rather than just maintenance, has also been observed (Tchalova & MacDonald,

2020). Third, the current study tested a sample of healthy female participants, therefore a

generalization to male individuals is not possible, especially given the known sex differences in

stress reactivity and MOR system functioning. Finally, it is unlikely that a complex behavior such

as social motivation can be fully explained by the activity of a single neurochemical system only.

For instance, the neuropeptides oxytocin and vasopressin are well-known for their crucial role in

bonding and reproductive functions (Donaldson & Young, 2008). Stress exposure induces a potent

activation of the dopaminergic system (Bloomfield et al., 2019; Holly & Miczek, 2016), and it has

been shown that MOR stimulation disinhibits dopaminergic neurons in the ventral tegmental area

and increases dopamine release in the nucleus accumbens (Pan, 1998). To investigate how these

systems interact in driving social contact seeking during aversive states seems therefore necessary

to understand the neurobiology of social motivation.

Conclusions

To conclude, our findings show that enhancing μ-opioid signaling before psychosocial stress

exposure increases, rather than reduces, the aversive reaction to stress, leading to an increased

subjective motivation for the attainment of highly rewarding social contact. Specifically, we

showed that morphine administration blunted cortisol reactivity to stress and increased the

negativity of the adverse experience, in line with a mood-buffering effect of cortisol. Further, the

results indicate that this enhanced morphine-induced stress response was followed by increased

motivation to attain social rewards, both in terms of subjective wanting as well as of facial

reactions during reward anticipation.

Overall, the results provide novel evidence on the modulation of the physiological and subjective

responses to social stress by the MOR system and indicate a state-dependent regulation of social

motivation. To better understand the role of μ-opioids in the potentiation of social motivation

during aversive states, further investigation of the effects of MOR agonists and antagonists

administered after stress induction, as well as of the interaction with other neurochemical systems,

is needed.

Materials & Methods

Study design

The between-subject, double-blind, placebo-controlled study consisted of one experimental session in which participants received either 10 mg morphine sulfate or a placebo (60 mg mannitol).

Participants

Based on previous work that had investigated the effects of stress on social reward processing (Massaccesi et al., 2021) and the effects of similar compounds on stress responses (Bershad et al., 2018), we aimed at collecting data from 40 participants per group. The study sample included 82 participants, of which 42 received morphine (MORPH) and 40 received a placebo (PLB). Two participants (MORPH) did not complete the session and were therefore not included in data analysis, yielding to a final sample size of 80 participants (40/group)[3]. Only female participants were included due to i) gender differences in opioid pharmacokinetics (Fillingim & Gear, 2004; Zubieta et al., 1999) and stress response (Kelly et al., 2008), ii) expected higher preference of same-sex touch in females than males (Stier & Hall, 1984; Suvilehto et al., 2015). Participants were tested during the luteal phase of their menstrual cycle, as assessed by self-report. All participants reported to be right-handed, to smoke less than ten cigarettes per week, to have no history of current or former drug abuse, to have a BMI between 17 and 35, and to be free of psychiatric or neurological disorders. Other exclusion criteria were: single or repeated use of any strong opioids in the last two years, use of hormonal contraceptives, regular intake of medications,

---

[3] Due to technical problems, EMG data from 6 (4 MORPH) participants and HR data from 1 participant (MORPH) were not recorded. Saliva samples from one participant (MORPH) are also missing.

current pregnancy or breastfeeding, suffering from impaired respiratory functions, respiratory

weakness or lung disease, injury/disease of the arms (making it impossible to squeeze with the

right hand and to be caressed on the left forearm). Participants were instructed to refrain from

eating, brushing their teeth and consuming caffeinated beverages, juices, and chewing gum in the

two hours preceding the test, as well as from smoking, doing physical activity and intaking alcohol

and medications in the 24 hours preceding the test. The two experimental groups did not differ

significantly in terms of age, BMI, autistic traits (short version of the German Autism Spectrum

Quotient, AQ-k; Freitag et al., 2007), general (State-Trate Anxiety Inventory, STAI; Spielberger

et al., 1970) and social (Liebowitz Social Anxiety Scale, LSAS; Heimberg et al., 1999) anxiety,

and social touch appreciation (Social Touch Questionnaire, STQ; Wilhelm et al., 2001) (Table 1).

The study was approved by the Ethics Committee of the Medical University of Vienna (EK

1393/2017) and was performed in line with the Declaration of Helsinki (World Medical

Association, 2013). All participants signed a consent form before taking part in the study.

Table 1. Demographic and self-reported substance use characteristics of the participants.

	MORPHINE	PLACEBO	*p* value
N	40	40	---
Age (years)	23.08 (3.08)	23.88 (3.86)	0.31
BMI (kg/m^2)	21.37 (3.38)	22.26 (2.66)	0.19
Autism (AQ-k)	6.45 (3.81)	5.65 (3.58)	0.34
Anxiety (STAI)	38.13 (9.43)	35.73 (8.27)	0.23
Social Anxiety (LSAS)	36.73 (17.87)	33.18 (19.02)	0.39
Social Touch Preferences (STQ)	25.33 (9.53)	27.10 (7.67)	0.36
Drug use (% lifetime – last year)			
Cannabis	47.5 – 25	60 – 32.5	---
Tranquilizers	7.5 – 5	5 – 2.5	---
Stimulants	25 – 10	17.5 – 7.5	---
Opiates	7.5 – 0	2.5 – 0	---
Hallucinogens	12.5 – 2.5	12.5 – 2.5	---

Other	5 – 2.5	2.5 – 0	---

Procedure

The study was conducted at the Department of Psychiatry and Psychotherapy of the Medical University of Vienna. After completing an online survey to assess their eligibility, potential participants were first invited to a health screening (~45'), including blood examination, electrocardiogram, blood pressure measurement, and psychiatric interview (Mini-International Neuropsychiatric Interview; Sheehan et al., 1998). Eligible participants were then invited to the experimental session (~210'), which always started between 11:30 and 12:30 in order to control for cortisol diurnal fluctuations (Labuschagne et al., 2019). At the beginning of the session, urine drug and pregnancy tests were administered. After baseline mood and physiological measures were collected, participants received a standardized snack, and took the assigned capsule. Throughout the session, mood and physiological measures were obtained at regular intervals. The experimental tasks were completed between 60 and 160 min after drug administration, and included economic decision making, facial mimicry, emotion recognition, stress induction and social reward processing (see Fig.1A for an overview of the session timeline). Here we will focus on the last two, while the others will be reported elsewhere. Approximately 180 min. after pill administration, a blood sample was taken to confirm drug uptake (see Table S2 of the *Supplementary Material*). After completing the experimental session, participants were debriefed and received a financial compensation of 65€. Half of the sample was tested before and half during the COVID-19 pandemic (see *Supplementary Material* for details regarding the employed safety measures and additional statistical analyses to assess possible effects of the pandemic and safety measures on the results).

Drug administration

Ten mg of morphine sulfate (Morapid®) were administered per oral to stimulate the activity of the MOR system with minimal subjective effects. Morphine is a selective MOR agonist and, for oral administration, has an average bioavailability of 30–40 %, a maximal effect (t-max) at 1–2 h after administration, and a half-life of 2–4 h (Lugo & Kern, 2002). Placebo consisted of capsules containing 650 mg of mannitol (sugar), visually identical to the ones containing morphine.

Stress induction

In order to induce a stress reaction in the participants, the Trier Social Stress Test (TSST; Kirschbaum et al., 1993) was employed. In the TSST, participants were given 3 minutes to prepare a 5-minute speech for a mock-job interview, followed by a 5-minute arithmetic task, in which they were asked to count backwards from 2043 in steps of 17 as fast and as accurate as possible. The speech and arithmetic tasks were completed in front of an evaluating panel (one male and one female confederates). Participants were told that these tasks would be video recorded via a camera located next to the examiners (no video was actually taped).

Social Reward task

Stimuli

Gentle caresses in three different speeds (CT-optimal: 6 cm/s, non-CT-optimal: 21 cm/s and 27 cm/s) were used as social rewards of different level of pleasantness (high, low, very low). Pleasantness of touch is high between 1 and 10 cm/s, and decreases between 10 and 30 cm/s (Löken et al., 2009) and CT-optimal touch has been shown to have a stress buffering effects, reducing arousal (Pawling et al., 2017) and distress following social exclusion (von Mohr et al., 2018), and increasing heart rate variability (Triscoli et al., 2017). The suitability of these stroking speeds has been confirmed in three previous independent studies from our group (Korb,

Götzendorfer, et al., 2020; Korb, Massaccesi, et al., 2020; Massaccesi et al., 2021). Caressed were

delivered over a previously-marked nine–cm area on participants' left forearm (measurement

started from the wrist towards the elbow) by a female experimenter, moving her index and middle

fingers back and forth in the marked area of the forearm (Figure 1B). Touch delivery was guided

by auditory rhythms, which matched the frequency of the stimulation, over headphones. The

experimenter administering the touch was seated on the other side of a curtain, used to limit the

participants' field of view to the monitor (Figure 1B). All experimenters were presented as trained

masseurs, wore standardized clothes (white t-shirt and beige trousers) to minimize differences in

their appearance, and underwent extensive training on the tactile stimulation delivery.

Task

The Social Reward task (Korb, Götzendorfer, et al., 2020; Massaccesi et al., 2021) consisted of

two blocks of 16 trials, separated by a 5-minute break. To avoid habituation to the touch, the site

of application (left or right area of the forearm) was alternated within the two blocks, in a counter-

balanced order. Before starting the task, participants experienced each type of touch once. Each

trial was structured as follow (see Figure 1C): i) announcement of the best attainable reward (high

or low, 16 trials each, 3s); ii) rating of subjective wanting via a VAS ranging from -10 (not at all)

to +10 (very much) (no time limit); iii) effort task (4s), requiring to squeeze a hand-dynamometer

(HD-BTA, Vernier Software & Technology, USA) with the right hand, in order to obtain the

announced reward – the applied force, displayed via an online visual-feedback, was expressed as

percentage of the participants' MVC, measured immediately before the task, and translated into

the probability of obtaining the announced reward (0–100%); iv) announcement of the reward

obtained (high, low, or – if insufficient force had been exerted – very low; 2s); v) reward delivery

(6s); vi) relax phase (5s); vii) rating of subjective liking via a VAS ranging from -10 (not at all) to

+10 (very much) (no time limit). At the end of the task, participants' MVC was measured again.

The task was implemented in MATLAB 2014b (MathWorks, Inc) and presented on an LCD

monitor with a resolution of 1280 × 1024 pixels.

Facial EMG

Facial EMG was recorded throughout the Social Reward task, using a g.USBamp amplifier (g.tec

Medical Engineering GmbH) and the software Matlab (MathWorks, Inc). Participants' face areas

were prepared using alcohol, water and an abrasive paste. Reusable Ag/AgCl electrodes were then

attached bipolarly according to guidelines (Fridlund & Cacioppo, 1986) on the left corrugator

supercilii (corrugator), and zygomaticus major (zygomaticus) muscles. A ground electrode was

attached to the participants' forehead and a reference electrode on the left mastoid. The EMG data

were sampled at 1200 Hz with impedances below 20 kΩ.

Each trial was divided in 4 epochs: Anticipation Pre-Effort (announcement of best attainable

reward, 3s), Anticipation Post-Effort (announcement of attained reward, 2s), Delivery (touch

administration, 6s), and Relax (relax phase after reward delivery, 5s). EMG was first averaged

over 1s time-windows and then over the epoch total duration. For each trial, values in the four

epochs were expressed as percentage of the average amplitude during the fixation cross at the

beginning of that trial (baseline, 2s). Outliers in baseline (defined as values 3 SDs over the subjects'

average baseline) were substituted with the average of the baseline preceding and following that

trial. The extracted epochs were visually inspected in order to identify signal artifacts which were

then removed (33 epochs for corrugator and 51 epochs for zygomaticus across 5 participants). As

data were extremely skewed, for each subject, epochs over the subject's mean ± 3 SD were

removed (average removed epochs per subject: corrugator: $M = 2.4$, $SD = 1.2$; zygomaticus: $M =$

3.1, $SD = 1.1$), and the remaining data were transformed using natural logarithmic transformation.

Subjective measures of stress and mood

During the session, participants completed self-report questionnaires assessing their mood and

subjective state at regular intervals. Happiness, calmness, relaxation, feeling good, as well as

stress, tension, anxiety and feeling bad were assessed via visual analogue scales ranging from +1

("not at all") to +101 ("very much") at 7 time points (T1-T7; see Figure 1A). Positive and negative

mood items were averaged to constitute the "Positive mood" and "Negative mood" scales used for

statistical analyses. The German short version Profile of Mood States (POMS; Albani et al., 2005),

consisting of 4 subscales for current mood (depression, vigor, fatigue, and displeasure), was

administered at 5 time points (T1, T2, T3, T5, T6; see Figure 1A). During the preparation phase

of the TSST (T4), anticipatory cognitive appraisal (PASA; Gaab et al., 2005) was also assessed.

Lastly, after completion of the TSST (T5), participants' satisfaction towards their performance at

the speech and math tasks was assessed on a VAS scale ranging from 1 ("not at all") to +101 ("very

much").

Physiological measures of stress

Heart rate was recorded using a chest strap (Polar H10; Polar Electro Oy, Kempele, Finland). Data

were collected over a 10-minute period at baseline, during TSST, and during the Social Reward

task. Values in each time window were then averaged for statistical analyses.

Saliva samples were collected via passive drool method using Salicaps (IBL, Hamburg, Germany)

at 6 time points (T1, T2, T3, T5, T6, T7; see Figure 1A). Free cortisol concentration in saliva was

determined by using commercial luminescence immunosorbent assay (LUM; IBL, Hamburg,

Germany). Salivary alpha-amylase activity was measured using a kinetic colorimetric test and

reagents obtained from DiaSys Diagnostic Systems (Holzheim, Germany). For heart rate, salivary

cortisol and alpha-amylase analyses, outliers were defined as subjects with a baseline value (T1)

3 SDs over the mean baseline of the sample. This procedure led to the exclusion of two participants

for cortisol (1 MORPH, 1 PLB) and of one participant for alpha-amylase (MORPH).

Drug effects on cognitive functions and side-effects

To assess potential drug effects on cognitive functions, participants completed the Trail Making

Test (TMT; Reitan, 1958) and the Digit Symbol Substitution Test (DSST; Wechsler, 1939) 55 min

after drug administration. Regarding subjective drug effects, participants filled out a self-report

questionnaire assessing nausea, dry mouth and other 24 possible side-effects on a 4-point Likert

scale (with the anchors 1 = "not at all" and 4 = "very much") at baseline (T1), as well as 50 min

(T2) and 160 min (T7) after drug administration (see Figure 1A).

Statistical analyses

Statistical analyses were conducted in R (R Core Team, 2019). Drug effects on subjective and

physiological state measures were analyzed using LMMs with Drug (MORPH, PLB) and Time as

fixed effects and by-subject random intercepts. Drug effects on ratings of wanting and liking, and

force exerted were analyzed with LMMs including Drug (MORPH, PLB) and Reward Level (high,

low, and very vow in the case of liking) as fixed effects, and by-subject random intercepts and

slopes for Reward Level. For EMG data, LMMs for each muscle and task phase were fitted,

including Drug (MORPH, PLB) and trial-by-trial Wanting (for anticipation) or Liking (for

consumption) as fixed effects, and by-subject random intercepts and slopes for Wanting/Liking.

In case of model unconvergence or singularity, random effects with the lowest cumulative variance

were removed and transformed into the corresponding complex random intercept (CRI; Scandola

& Tidoni, 2021). Group differences in age, BMI, and personality traits, as well as drug effects on

executive functions, PASA, satisfaction for the TSST performance and side-effects were assessed

using independent two-sided t-tests.

LMMs were computed using the lmer() function of the *lme4* package (Bates et al., 2015). Type-

III F-tests were computed with the Satterthwaite degrees of freedom approximation, using the

anova() function of the *lmerTest* package (Kuznetsova et al., 2017). Significant interactions were

further analyzed with multiple comparisons using the function emmeans() from the homonymous

package (Lenth, 2021). Results from all LMMs and multiple comparisons were controlled for the

false discovery rate (FDR) associated with multiple testing using the Benjamini–Hochberg method

(Benjamini & Hochberg, 1995). Data and analysis scripts are available online

(https://osf.io/gbd24/).

599 **Funding**

600 The study was supported by the Vienna Science and Technology Fund (WWTF) with a grant

601 (CS15-003) awarded to Giorgia Silani and Matthäus Willeit, as well as by the Uni:docs scholarship

602 and the JUWI Covid-19 fellowship of the University of Vienna, and the OeAD Marietta Blau grant

603 awarded to Claudia Massaccesi. Funders had no role in study design, data collection and analyses,

604 decision to publish or preparation of the manuscript.

605 **Acknowledgments**

606 We thank Gheorghe L. Preda and Carina Bum for their contribution to carry out the medical

607 procedures during data collection, Sebastian Korb for valuable intellectual input, Nadine Skoluda

608 for her support with the saliva analysis, and all the students who contributed to data collection.

Authors contribution

CM: Conceptualization of the study, Data curation, Software, Formal analysis, Supervision, Investigation, Visualization, Methodology, Project administration, Funding acquisition, Writing - original draft, review and editing; MW: Contribution to the conceptualization of the design, Project administration, Medical testing, Funding acquisition, Writing - review and editing; BBQ: Contribution to the conceptualization of the design, Writing - review and editing; UMN: Contribution to the conceptualization of the design, Curation of saliva analysis, Writing - review and editing; CL: Contribution to the conceptualization of the design, Writing - review and editing; DM: Curation of blood analyses; GS: Conceptualization of the study, Supervision, Methodology, Funding acquisition, Project administration, Writing - review and editing.

All authors approved the final version of the manuscript.

Competing interests

No competing interests declared.

References

Albani, C., Blaser, G., Geyer, M., Schmutzer, G., Brähler, E., Bailer, H., & Grulke, N. (2005). The German short version of "Profile of Mood States" (POMS): Psychometric evaluation in a representative sample. *PPmP - Psychotherapie · Psychosomatik · Medizinische Psychologie*, *55*(07), 324–330. https://doi.org/10.1055/s-2004-834727

Ali, N., Cooperman, C., Nitschke, J. P., Baldwin, M. W., & Pruessner, J. C. (2020). The effects of suppressing the biological stress systems on social threat-assessment following acute stress. *Psychopharmacology*, *237*(10), 3047–3056. https://doi.org/10.1007/s00213-020-05591-z

Ali, N., Nitschke, J. P., Cooperman, C., & Pruessner, J. C. (2017). Suppressing the endocrine and

autonomic stress systems does not impact the emotional stress experience after

psychosocial stress. *Psychoneuroendocrinology*, *78*, 125–130.

https://doi.org/10.1016/j.psyneuen.2017.01.015

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

https://doi.org/10.18637/jss.v067.i01

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*

(Methodological), *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bershad, A. K., Miller, M. A., Norman, G. J., & de Wit, H. (2018). Effects of opioid- and non-

opioid analgesics on responses to psychosocial stress in humans. *Hormones and*

Behavior, *102*, 41–47. https://doi.org/10.1016/j.yhbeh.2018.04.009

Bershad, A. K., Seiden, J. A., & de Wit, H. (2016). Effects of buprenorphine on responses to

social stimuli in healthy adults. *Psychoneuroendocrinology*, *63*, 43–49.

https://doi.org/10.1016/j.psyneuen.2015.09.011

Bloomfield, M. A., McCutcheon, R. A., Kempton, M., Freeman, T. P., & Howes, O. (2019). The

effects of psychosocial stress on dopaminergic function and the acute stress response.

ELife, *8*, e46797. https://doi.org/10.7554/eLife.46797

Buchel, C., Miedl, S., & Sprenger, C. (2018). Hedonic processing in humans is mediated by an

opioidergic mechanism in a mesocorticolimbic system. *ELife*, *7*, e39648.

https://doi.org/10.7554/eLife.39648

Cacioppo, J. T., & Cacioppo, S. (2014). Social Relationships and Health: The Toxic Effects of

Perceived Social Isolation. *Social and Personality Psychology Compass*, *8*(2), 58–72.

https://doi.org/10.1111/spc3.12087

Cacioppo, J. T., Petty, R. E., & Morris, K. J. (1985). Semantic, Evaluative, and Self-Referent

Processing: Memory, Cognitive Effort, and Somatovisceral Activity. *Psychophysiology*,

22(4), 371–384. https://doi.org/10.1111/j.1469-8986.1985.tb01618.x

Chelnokova, O., Laeng, B., Eikemo, M., Riegels, J., Løseth, G., Maurud, H., Willoch, F., &

Leknes, S. (2014). Rewards of beauty: The opioid system mediates social motivation in

humans. *Molecular Psychiatry*, *19*(7), 746–747. https://doi.org/10.1038/mp.2014.1

Cohen, B. H., Davidson, R. J., Senulis, J. A., Saron, C. D., & Weisman, D. R. (1992). Muscle

tension patterns during auditory attention. *Biological Psychology*, *33*(2–3), 133–156.

https://doi.org/10.1016/0301-0511(92)90028-s

de Morree, H. M., & Marcora, S. M. (2010). The face of effort: Frowning muscle activity reflects

effort during a physical task. *Biological Psychology*, *85*(3), 377–382.

https://doi.org/10.1016/j.biopsycho.2010.08.009

Donaldson, Z. R., & Young, L. J. (2008). Oxytocin, Vasopressin, and the Neurogenetics of

Sociality. *Science*, *322*(5903), 900–904. https://doi.org/10.1126/science.1158668

Eikemo, M., Løseth, G. E., Johnstone, T., Gjerstad, J., Willoch, F., & Leknes, S. (2016). Sweet

taste pleasantness is modulated by morphine and naltrexone. *Psychopharmacology*,

233(21–22), 3711–3723. https://doi.org/10.1007/s00213-016-4403-x

Fabre-Nys, C., Meller, R. E., & Keverne, E. B. (1982). Opiate antagonists stimulate affiliative

behaviour in monkeys. *Pharmacology Biochemistry and Behavior*, *16*(4), 653–659.

https://doi.org/10.1016/0091-3057(82)90432-4

Fillingim, R. B., & Gear, R. W. (2004). Sex differences in opioid analgesia: Clinical and

experimental findings. *European Journal of Pain*, *8*(5), 413–425.

https://doi.org/10.1016/j.ejpain.2004.01.007

Freitag, C. M., Retz-Junginger, P., Retz, W., Seitz, C., Palmason, H., Meyer, J., Rösler, M., &

von Gontard, A. (2007). Evaluation der deutschen Version des Autismus-Spektrum-

Quotienten (AQ)—Die Kurzversion AQ-k. *Zeitschrift Für Klinische Psychologie Und*

Psychotherapie, *36*(4), 280–289. https://doi.org/10.1026/1616-3443.36.4.280

Fridlund, A. J., & Cacioppo, J. T. (1986). Guidelines for Human Electromyographic Research.

Psychophysiology, *23*(5), 567–589. https://doi.org/10.1111/j.1469-8986.1986.tb00676.x

Gaab, J., Rohleder, N., Nater, U. M., & Ehlert, U. (2005). Psychological determinants of the

cortisol stress response: The role of anticipatory cognitive appraisal.

Psychoneuroendocrinology, *30*(6), 599–610.

https://doi.org/10.1016/j.psyneuen.2005.02.001

Gerra, G., Zaimovic, A., Mascetti, G. G., Gardini, S., Zambelli, U., Timpano, M., Raggi, M. A.,

& Brambilla, F. (2001). Neuroendocrine responses to experimentally-induced

psychological stress in healthy humans. *Psychoneuroendocrinology*, *26*(1), 91–107.

https://doi.org/10.1016/S0306-4530(00)00046-9

Heimberg, R. G., Horner, K. J., Juster, H. R., Safren, S. A., Brown, E. J., Schneier, F. R., &

Liebowitz, M. R. (1999). Psychometric properties of the Liebowitz Social Anxiety Scale.

Psychological Medicine, *29*(1), 199–212. https://doi.org/10.1017/S0033291798007879

Het, S., Schoofs, D., Rohleder, N., & Wolf, O. T. (2012). Stress-Induced Cortisol Level

Elevations Are Associated With Reduced Negative Affect After Stress: Indications for a

Mood-Buffering Cortisol Effect. *Psychosomatic Medicine*, *74*(1), 23–32.

https://doi.org/10.1097/PSY.0b013e31823a4a25

Het, S., & Wolf, O. T. (2007). Mood changes in response to psychosocial stress in healthy young

women: Effects of pretreatment with cortisol. *Behavioral Neuroscience*, *121*(1), 11–20.

https://doi.org/10.1037/0735-7044.121.1.11

Hickey, C., Chelazzi, L., & Theeuwes, J. (2010). Reward Changes Salience in Human Vision via

the Anterior Cingulate. *Journal of Neuroscience*, *30*(33), 11096–11103.

Hickey, C., & van Zoest, W. (2012). Reward creates oculomotor salience. *Current Biology*,

22(7), R219–R220. https://doi.org/10.1016/j.cub.2012.02.007

Holly, E. N., & Miczek, K. A. (2016). Ventral tegmental area dopamine revisited: Effects of

acute and repeated stress. *Psychopharmacology*, *233*(2), 163–186.

https://doi.org/10.1007/s00213-015-4151-3

Hsu, D. T., Sanford, B. J., Meyers, K. K., Love, T. M., Hazlett, K. E., Wang, H., Ni, L., Walker,

S. J., Mickey, B. J., Korycinski, S. T., Koeppe, R. A., Crocker, J. K., Langenecker, S. A.,

& Zubieta, J.-K. (2013). Response of the μ-opioid system to social rejection and

acceptance. *Molecular Psychiatry*, *18*(11), 1211–1217.

https://doi.org/10.1038/mp.2013.96

Inagaki, T. K., Hazlett, L. I., & Andreescu, C. (2020). Opioids and social bonding: Effect of

naltrexone on feelings of social connection and ventral striatum activity to close others.

Journal of Experimental Psychology: General, *149*(4), 732–745.

https://doi.org/10.1037/xge0000674

Inagaki, T. K., Irwin, M. R., & Eisenberger, N. I. (2015). Blocking opioids attenuates physical

warmth-induced feelings of social connection. *Emotion (Washington, D.C.)*, *15*(4), 494–

500. https://doi.org/10.1037/emo0000088

Inagaki, T. K., Ray, L. A., Irwin, M. R., Way, B. M., & Eisenberger, N. I. (2016). Opioids and

social bonding: Naltrexone reduces feelings of social connection. *Social Cognitive and*

Affective Neuroscience, *11*(5), 728–735. https://doi.org/10.1093/scan/nsw006

Kelly, M. M., Tyrka, A. R., Anderson, G. M., Price, L. H., & Carpenter, L. L. (2008). Sex

differences in emotional and physiological responses to the Trier Social Stress Test.

Journal of Behavior Therapy and Experimental Psychiatry, *39*(1), 87–98.

https://doi.org/10.1016/j.jbtep.2007.02.003

Keverne, E. B., Martensz, N. D., & Tuite, B. (1989). Beta-endorphin concentrations in

cerebrospinal fluid of monkeys are influenced by grooming relationships.

Psychoneuroendocrinology, *14*(1), 155–161. https://doi.org/10.1016/0306-

4530(89)90065-6

Kirsch, L. P., Krahé, C., Blom, N., Crucianelli, L., Moro, V., Jenkinson, P. M., & Fotopoulou, A.

(2017). Reading the Mind in the Touch: Neurophysiological Specificity in the

Communication of Emotions by Touch. *Neuropsychologia*.

https://doi.org/10.1016/j.neuropsychologia.2017.05.024

Kirschbaum, C., Pirke, K. M., & Hellhammer, D. H. (1993). The 'Trier Social Stress Test'—A

tool for investigating psychobiological stress responses in a laboratory setting.

Neuropsychobiology, *28*(1–2), 76–81. https://doi.org/10.1159/000119004

Korb, S., Götzendorfer, S. J., Massaccesi, C., Sezen, P., Graf, I., Willeit, M., Eisenegger, C., &

Silani, G. (2020). Dopaminergic and opioidergic regulation during anticipation and

consumption of social and nonsocial rewards. *ELife*, *9*, e55797.

https://doi.org/10.7554/eLife.55797

Korb, S., Massaccesi, C., Gartus, A., Lundström, J. N., Rumiati, R., Eisenegger, C., & Silani, G.

(2020). Facial responses of adult humans during the anticipation and consumption of

touch and food rewards. *Cognition*, *194*, 104044.

https://doi.org/10.1016/j.cognition.2019.104044

Kumar, P., Berghorst, L. H., Nickerson, L. D., Dutra, S. J., Goer, F. K., Greve, D. N., &

Pizzagalli, D. A. (2014). Differential effects of acute stress on anticipatory and

consummatory phases of reward processing. *Neuroscience*, *266*(Supplement C), 1–12.

https://doi.org/10.1016/j.neuroscience.2014.01.058

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in

Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(1), 1–26.

https://doi.org/10.18637/jss.v082.i13

Labuschagne, I., Grace, C., Rendell, P., Terrett, G., & Heinrichs, M. (2019). An introductory

guide to conducting the Trier Social Stress Test. *Neuroscience and Biobehavioral*

Reviews, *107*, 686–695. https://doi.org/10.1016/j.neubiorev.2019.09.032

Leknes, S., & Tracey, I. (2008). A common neurobiology for pain and pleasure. *Nature Reviews.*

Neuroscience, *9*(4), 314–320. https://doi.org/10.1038/nrn2333

Lenth, R. V. (2021). *emmeans: Estimated marginal means, aka least-squares means. R package*

version 1.5. 4.

Lewis, A. H., Porcelli, A. J., & Delgado, M. R. (2014). The effects of acute stress exposure on

striatal activity during Pavlovian conditioning with monetary gains and losses. *Frontiers*

in Behavioral Neuroscience, *8*. https://doi.org/10.3389/fnbeh.2014.00179

Löken, L. S., Wessberg, J., Morrison, I., McGlone, F., & Olausson, H. (2009). Coding of

766   pleasant touch by unmyelinated afferents in humans. *Nature Neuroscience*, *12*(5), 547–

767   548. https://doi.org/10.1038/nn.2312

768   Løseth, G. E., Eikemo, M., & Leknes, S. (2019). Effects of opioid receptor stimulation and

769   blockade on touch pleasantness: A double-blind randomised trial. *Social Cognitive and*

770   *Affective Neuroscience*, *14*(4), 411–422. https://doi.org/10.1093/scan/nsz022

771   Loseth, G. E., Ellingsen, D.-M., & Leknes, S. (2014). State-dependent μ-opioid modulation of

772   social motivation. *Frontiers in Behavioral Neuroscience*, *8*, 430.

773   https://doi.org/10.3389/fnbeh.2014.00430

774   Lugo, R. A., & Kern, S. E. (2002). Clinical Pharmacokinetics of Morphine. *Journal of Pain &*

775   *Palliative Care Pharmacotherapy*, *16*(4), 5–18. https://doi.org/10.1080/J354v16n04_02

776   Machin, A. J., & Dunbar, R. I. M. (2011). The brain opioid theory of social attachment: A review

777   of the evidence. *Behaviour*, *148*(9–10), 985–1025.

778   https://doi.org/10.1163/000579511X596624

779   Manninen, S., Tuominen, L., Dunbar, R. I., Karjalainen, T., Hirvonen, J., Arponen, E., Hari, R.,

780   Jääskeläinen, I. P., Sams, M., & Nummenmaa, L. (2017). Social Laughter Triggers

781   Endogenous Opioid Release in Humans. *Journal of Neuroscience*, *37*(25), 6125–6131.

782   https://doi.org/10.1523/JNEUROSCI.0688-16.2017

783   Martel, F. L., Nevison, C. M., Simpson, M. J. A., & Keverne, E. B. (1995). Effects of opioid

784   receptor blockade on the social behavior of rhesus monkeys living in large family groups.

785   *Developmental Psychobiology*, *28*(2), 71–84. https://doi.org/10.1002/dev.420280202

Massaccesi, C., Korb, S., Skoluda, N., Nater, U. M., & Silani, G. (2021). Effects of Appetitive

and Aversive Motivational States on Wanting and Liking of Interpersonal Touch.

Neuroscience, *464*, 12–25. https://doi.org/10.1016/j.neuroscience.2020.09.025

Mayo, L. M., Lindé, J., Olausson, H., Heilig, M., & Morrison, I. (2018). Putting a good face on

touch: Facial expression reflects the affective valence of caress-like touch across

modalities. *Biological Psychology*, *137*, 83–90.

https://doi.org/10.1016/j.biopsycho.2018.07.001

McEwen, B. S. (1998). Stress, adaptation, and disease. Allostasis and allostatic load. *Annals of*

the New York Academy of Sciences, *840*, 33–44. https://doi.org/10.1111/j.1749-

6632.1998.tb09546.x

Meller, R. E., Keverne, E. B., & Herbert, J. (1980). Behavioural and endocrine effects of

naltrexone in male talapoin monkeys. *Pharmacology Biochemistry and Behavior*, *13*(5),

663–672. https://doi.org/10.1016/0091-3057(80)90010-6

Morrison, I. (2016). Keep Calm and Cuddle on: Social Touch as a Stress Buffer. *Adaptive*

Human Behavior and Physiology, *2*(4), 344–362. https://doi.org/10.1007/s40750-016-

0052-x

Morrison, I., Löken, L. S., & Olausson, H. (2010). The skin as a social organ. *Experimental*

Brain Research, *204*(3), 305–314. https://doi.org/10.1007/s00221-009-2007-y

Pan, Z. Z. (1998). μ-Opposing actions of the κ-opioid receptor. *Trends in Pharmacological*

Sciences, *19*(3), 94–98. https://doi.org/10.1016/S0165-6147(98)01169-9

Panksepp, J. (1998). *Affective Neuroscience: The Foundations of Human and Animal Emotions*.

Oxford University Press.

Panksepp, J., & Bishop, P. (1981). An autoradiographic map of (3H)diprenorphine binding in rat

brain: Effects of social interaction. *Brain Research Bulletin*, *7*(4), 405–410.

Panksepp, J., Herman, B. H., Vilberg, T., Bishop, P., & DeEskinazi, F. G. (1980). Endogenous

opioids and social behavior. *Neuroscience & Biobehavioral Reviews*, *4*(4), 473–487.

https://doi.org/10.1016/0149-7634(80)90036-6

Pawling, R., Cannon, P. R., McGlone, F. P., & Walker, S. C. (2017). C-tactile afferent

stimulating touch carries a positive affective value. *PLOS ONE*, *12*(3), e0173457.

https://doi.org/10.1371/journal.pone.0173457

Pechnick, R. N. (1993). Effects of Opioids on the Hypothalamo-Pituitary-Adrenal Axis. *Annual*

Review of Pharmacology and Toxicology, *33*(1), 353–382.

https://doi.org/10.1146/annurev.pa.33.040193.002033

Petrowski, K., Wintermann, G.-B., & Siepmann, M. (2012). Cortisol response to repeated

psychosocial stress. *Applied Psychophysiology and Biofeedback*, *37*(2), 103–107.

https://doi.org/10.1007/s10484-012-9183-4

Pool, E., Brosch, T., Delplanque, S., & Sander, D. (2015). Stress increases cue-triggered

"wanting" for sweet reward in humans. *Journal of Experimental Psychology. Animal*

Learning and Cognition, *41*(2), 128–136. https://doi.org/10.1037/xan0000052

Pruessner, J. C., Kirschbaum, C., Meinlschmid, G., & Hellhammer, D. H. (2003). Two formulas

for computation of the area under the curve represent measures of total hormone

concentration versus time-dependent change. *Psychoneuroendocrinology*, *28*(7), 916–

931. https://doi.org/10.1016/S0306-4530(02)00108-7

Quednow, B. B., Csomor, P. A., Chmiel, J., Beck, T., & Vollenweider, F. X. (2008).

Sensorimotor gating and attentional set-shifting are improved by the μ-opioid receptor

831 agonist morphine in healthy human volunteers. *International Journal of*

832 *Neuropsychopharmacology*, *11*(5), 655–669.

833 https://doi.org/10.1017/S1461145707008322

834 R Core Team. (2019). *R: A language and environment for statistical computing.* R Foundation

835 for Statistical Computing.

836 Ree, A., Bendas, J., Pabel, L., Croy, I., & Sailer, U. (2020). Right between the eyes: Corrugator

837 muscle activity tracks the changing pleasantness of repeated slow stroking touch.

838 *Physiology & Behavior*, *222*, 112903. https://doi.org/10.1016/j.physbeh.2020.112903

839 Reitan, R. M. (1958). Validity of the Trail Making Test as an Indicator of Organic Brain

840 Damage. *Perceptual and Motor Skills*, *8*(3), 271–276.

841 https://doi.org/10.2466/pms.1958.8.3.271

842 Sato, W., Minemoto, K., Ikegami, A., Nakauma, M., Funami, T., & Fushiki, T. (2020). Facial

843 EMG Correlates of Subjective Hedonic Responses During Food Consumption. *Nutrients*,

844 *12*(4). https://doi.org/10.3390/nu12041174

845 Scandola, M., & Tidoni, E. (2021). *The development of a standard procedure for the optimal*

846 *reliability-feasibility trade-off in Multilevel Linear Models analyses in Psychology and*

847 *Neuroscience.* PsyArXiv. https://doi.org/10.31234/osf.io/kfhgv

848 Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T.,

849 Baker, R., & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview

850 (M.I.N.I.): The Development and Validation of a Structured Diagnostic Psychiatric

851 Interview for DSM-IV and ICD-10. *The Journal of Clinical Psychiatry*, *59*(suppl 20), 0–

852 0.

Spielberger, C. D., Gorsuch, R. L., & Lushene, R. E. (1970). *Manual for the State-Trait Anxiety*

Inventory. http://ubir.buffalo.edu/xmlui/handle/10477/2895

Stier, D. S., & Hall, J. A. (1984). Gender differences in touch: An empirical and theoretical

review. *Journal of Personality and Social Psychology*, *47*(2), 440–459.

https://doi.org/10.1037/0022-3514.47.2.440

Suvilehto, J. T., Glerean, E., Dunbar, R. I. M., Hari, R., & Nummenmaa, L. (2015). Topography

of social touching depends on emotional bonds between humans. *Proceedings of the*

National Academy of Sciences, *112*(45), 13811–13816.

https://doi.org/10.1073/pnas.1519231112

Tchalova, K., & MacDonald, G. (2020). Opioid receptor blockade inhibits self-disclosure during

a closeness-building social interaction. *Psychoneuroendocrinology*, *113*, 104559.

https://doi.org/10.1016/j.psyneuen.2019.104559

Theeuwes, J., & Belopolsky, A. V. (2012). Reward grabs the eye: Oculomotor capture by

rewarding stimuli. *Vision Research*, *74*, 80–85.

https://doi.org/10.1016/j.visres.2012.07.024

Triscoli, C., Croy, I., Steudte-Schmiedgen, S., Olausson, H., & Sailer, U. (2017). Heart rate

variability is enhanced by long-lasting pleasant touch at CT-optimized velocity.

Biological Psychology, *128*, 71–81. https://doi.org/10.1016/j.biopsycho.2017.07.007

Uchino, B. N. (2006). Social Support and Health: A Review of Physiological Processes

Potentially Underlying Links to Disease Outcomes. *Journal of Behavioral Medicine*,

29(4), 377–387. https://doi.org/10.1007/s10865-006-9056-5

Van Boxtel, A., & Jessurun, M. (1993). Amplitude and bilateral coherency of facial and jaw-

elevator EMG activity as an index of effort during a two-choice serial reaction task.

Psychophysiology, *30*(6), 589–604. https://doi.org/10.1111/j.1469-8986.1993.tb02085.x

Vanderschuren, L. J. M. J., Stein, E. A., Wiegant, V. M., & Van Ree, J. M. (1995). Social play

alters regional brain opioid receptor binding in juvenile rats. *Brain Research*, *680*(1),

148–156. https://doi.org/10.1016/0006-8993(95)00256-P

von Mohr, M., Krahé, C., Beck, B., & Fotopoulou, A. (2018). The social buffering of pain by

affective touch: A laser-evoked potential study in romantic couples. *Social Cognitive and*

Affective Neuroscience, *13*(11), 1121–1130. https://doi.org/10.1093/scan/nsy085

Waterink, W., & van Boxtel, A. (1994). Facial and jaw-elevator EMG activity in relation to

changes in performance level during a sustained information processing task. *Biological*

Psychology, *37*(3), 183–198. https://doi.org/10.1016/0301-0511(94)90001-9

Wechsler, D. (1939). *The measurement of adult intelligence* (pp. ix, 226). Williams & Wilkins

Co. https://doi.org/10.1037/10020-000

Wilhelm, F. H., Kochar, A. S., Roth, W. T., & Gross, J. J. (2001). Social anxiety and response to

touch: Incongruence between self-evaluative and physiological reactions. *Biological*

Psychology, *58*(3), 181–202. https://doi.org/10.1016/S0301-0511(01)00113-2

Zubieta, J.-K., Dannals, R. F., & Frost, J. J. (1999). Gender and Age Influences on Human Brain

Mu-Opioid Receptor Binding Measured by PET. *American Journal of Psychiatry*,

156(6), 842–848. https://doi.org/10.1176/ajp.156.6.842

Supplementary Material

Enhanced negative response to stress following morphine administration

increases wanting of social reward

C. Massaccesi, M. Willeit, B. B. Quednow, U. M. Nater, C. Lamm, D. Mueller, G. Silani

Index

1. Effects of COVID-19 pandemic

Half of the sample was collected after the COVID-19 pandemic outbreak. The implemented safety

measures, as well as the statistical analyses conducted to assess the possible effects of the pandemic

on the study dependent variables are described below.

a.   Safety measures

After the COVID-19 pandemic outbreak the following safety measures were implemented:

• All experimenters wore face masks. The evaluating panel of the TSST wore a special face

mask with a clear plastic insert on the mouth region in order to allow the participant to see the

"absence of facial feedback" (crucial for stress induction) during the stress paradigm.

• Participants were provided with clear mouth visors resting on the chin in order to minimize

the contact with the face and avoid disturbances during facial electromyography (EMG).

In order to assess participants reaction towards these safety measures, they were asked to answer

the following questions at the end of the experimental session:

1) "How well/comfortable did you feel during the study?" rated on a VAS ranging from 1 (not

all) to 101 (very much).

2) "How high do you rate the risk of infection during the study?" rated on a VAS ranging from

1 (very high) to 101 (very low).

3) "Were you afraid of being infected with COVID-19 during the study?", Yes/No

4) "How comfortable did you feel with the research team wearing a mask during the study?"

rated on a VAS ranging from 1 (not all) to 101 (very much).

As shown in Figure S1, participants did not report to feel threatened by COVID-19 infection during

the study, and to feel comfortable during the session.

Figure S1. Participants' attitudes toward COVID-19 pandemic and safety measures during the
study. (A) Ratings of wellness and comfort. (B) Perceived risk of COVID-19 infection during the
study. (C) Ratings of comfort related to use of face masks during the study. (D) Perceived fear of
having been infected with COVID-19 during the study.

b.   Additional analyses to assess the effects of the COVID-19 pandemic and employed
safety measures on responses to social rewards

In order to rule out possible effects of the pandemic and related implemented safety measures on
subjective stress response (positive and negative mood, POMS subscales), as well as on the ratings
of wanting and liking, on the force exerted to obtain the social rewards, and on facial EMG data,
by adding the covariate "COVID-19" (2 levels: pre-covid19, covid19) to the analyses. No changes
in the pattern of results were observed.

2. Drug effects on cognitive functions

Table S1. Mean (SD) scores at the Trial Making Test (TMT) part A and part B, and at the Digit
Symbol Substitution Test (DSST) across drug groups.

	MORPHINE	PLACEBO	*p* value
TMT A	26.33 (8.02)	26.40 (8.60)	0.97
TMT B	57.25 (33.05)	58.93 (21.94)	0.79
DSST	54.20 (8.99)	54.10 (9.32)	0.96

3. Drug side-effects

Figure S2. Drug side-effects assessed at baseline (T1), 60 min (T2) and 160 min (T7) after drug
administration using a 4-point Likert scale (with the anchors 1 = "not at all" and 4 = "very much").

4. Serum levels of morphine and its metabolites

A blood sample was drawn at the end of the session (~180 min after drug administration). Analyses
were performed at the Institute of Clinical Chemistry, University Hospital Zurich, using liquid
chromatography coupled to mass spectrometry (LC-MS) to identify serum levels of morphine and
its two major metabolites: morphine-3-glucuronide (M3G) and morphine-6-glucuronide (M6G).
Blood samples from 3 participants could not be obtained and 10 samples were lost because of

storage problems. Results from the available samples confirmed drug uptake, as shown in *Table*

S2.

Table S2. Serum levels (nmol/l) of morphine and its major metabolites at the end of the
experimental session (~180 min after drug administration). M3G & M6G = morphine-3/6-
glucronide.

	M	SD
Morphine	12.63	6.82
M3G	394.79	179.46
M6G	84.63	44.32