Drosophila nicotinic acetylcholine receptors and their interactions with insecticidal peptide toxins

Dagmara Korona[1]*, Benedict Dirnberger[1,2,4]*, Carlo N G Giachello[4]*, Rayner M L Queiroz[2], David-Paul Minde[2], Michael J Deery[2], Glynnis Johnson[1], Karin H Müller[3], Lucy C Firth[4], Fergus G Earley[4], Steven Russell[1] ** and Kathryn S Lilley[5] **

[1]Department of Genetics, University of Cambridge, Downing Street, Cambridge, CB2 3EH, United Kingdom

[2]Cambridge Centre for Proteomics, Department of Biochemistry, Gleeson Building, University of Cambridge, Tennis Court Road, Cambridge, CB2 1GA, United Kingdom

[3]Cambridge Advanced Imaging Centre, Department of Physiology, Development and Neuroscience/Anatomy Building, University of Cambridge, Downing Street, CB2 3DY, United Kingdom

[4]Syngenta, Jealott's Hill International Research Centre, Bracknell, RG42 6EY, United Kingdom

[5]Cambridge Centre for Proteomics, Department of Biochemistry, Gleeson Building, University of Cambridge, Tennis Court Road, Cambridge, CB2 1GA, United Kingdom. k.s.lilley@bioc.cam.ac.uk

*These authors contributed equally: Dagmara Korona, Benedict Dirnberger, Carlo N G Giachello

**Correspondence should be addressed to Kathryn S Lilley k.s.lilley@bioc.cam.ac.uk and Steven Russell sr120@cam.ac.uk

Abstract

Drosophila nicotinic acetylcholine receptors (nAChRs) are ligand-gated ion channels that represent a target for insecticides. Peptide neurotoxins are known to block nAChRs by binding to their target subunits, however, a better understanding of receptor subunit composition is needed for effective design of insecticides. To facilitate the analysis of nAChRs we used a CRISPR/Cas9 strategy to generate null alleles for all ten *nAChR* subunit genes in a common genetic background. We studied interactions of nAChR subunits with peptide neurotoxins by larval injections and styrene maleic acid lipid particles (SMALPs) pull-down assays. For the null alleles we determined the effects of α-Bungarotoxin (α-Btx) and ω-Hexatoxin-Hv1a (Hv1a) administration, identifying potential receptor subunits implicated in the binding of these toxins. We employed pull-down assays to confirm α-Btx interactions with the Dα5, Dα6, Dα7 subunits. Finally, we report the localization of fluorescent tagged endogenous Dα6 during nervous system development. Taken together this study elucidates native *Drosophila* nAChR subunit interactions with insecticidal peptide toxins and provides a resource for the *in vivo* analysis of insect nAChRs.

Keywords *Drosophila* nAChRs / insecticidal toxins / nAChR null alleles / SMALPs pull-downs

Subject Categories Neurotransmitter receptor/ Neurotoxin interactions

Introduction

Global climate change and other factors are placing increasing demands on available agricultural land to deliver efficient, reliable and sustainable food production. Insecticides are important tools in securing yields of all major crops but need to be continually replaced to overcome resistance in target species and reduce environmental impacts. In addition, new insecticides must have low toxicity to non-target species, particularly the major pollinators essential for agriculture. A large class of insecticide targets are neurotransmitter receptors such as the nicotinic acetylcholine receptors (nAChRs) located in synaptic plasma membranes (Ihara *et al*, 2020). These pentameric cys-loop ligand-gated ion channels consist of either only α-subunits or α- and β-subunits, with ligand binding sites located between two α-subunits or between α- and β-subunits. Most insect genomes, including that of the highly tractable *Drosophila melanogaster* model, harbour ten highly conserved subunit genes that assemble in various combinations to form the active receptors.

An essential pre-requisite for effective design of new insecticides targeting these receptors is understanding the subunit composition of nAChRs and their distinctive binding properties. For many reasons, including low expression in endogenous tissues or difficulties in expressing insect receptors in heterologous systems, the characterisation of functional insect receptors has been challenging (Perry *et al*, 2021; Zuo *et al*, 2021; Salgado, 2021). Even in the tractable *D. melanogaster* insect model, there has been no systematic isolation of mutations in *nAChR* subunit genes, until recently, when Perry and colleagues described the generation of a new set of null mutations in nine out of the ten *D. melanogaster* subunit genes (Perry *et al*, 2021). These mutations, however, were generated in different genetic backgrounds necessitating additional work to assay background sensitive phenotypes such as neural or behavioural defects.

Several classes of insecticide, the most effective being those in the neonicotinoid and spinosad class, have been shown to bind insect nAChRs highly selectively to block their functions (Chambers *et al*, 2019; Houchat *et al*, 2019). Recently, the binding affinity and the positive allosteric effects of ω-Hexatoxin-Hv1a (Hv1a) peptide on nAChRs has been demonstrated (Chambers *et al*, 2019) and this spider venom peptide is well known for its insecticidal effects. In addition, other peptide toxins, such the snake venom constituent, α-Bungarotoxin (α-Btx), have been widely used to probe nAChR functions, however whether α-Btx harbours a selective insecticidal property is currently unknown. Alpha-Btx is a 74 amino acid peptide that binds irreversibly to nAChR α-subunits in different species, including *D. melanogaster,* although the

exact subunit composition of target receptors is not fully understood (Schmidt-Nielsen *et al*, 1977; Dellisanti *et al*, 2007; Dacosta *et al*, 2015). Landsdell and co-workers have shown binding of α-Btx to *D. melanogaster* Dα5, Dα6, and Dα7 subunits in a heterologous S2 cell expression system (Lansdell & Millar, 2004; Lansdell *et al*, 2012) and the amino acid sequence of these subunits show strong similarity across their ligand-binding domains (LBD).

The lipid bilayer surrounding nAChRs is known to be essential for structural integrity, stability and ligand binding (Dacosta *et al*, 2013). However, this lipid requirement can make analysis of membrane protein complexes challenging. The development of methods for extracting membrane proteins from lipid bilayers using detergents and introducing them into artificial lipid nanodiscs has facilitated a much better characterisation of receptor-ligand interactions (Denisov & Sligar, 2016). The use of detergents generally used to solubilize membrane proteins, however, leads to destabilisation, aggregation and misfolding and are therefore not compatible with this type of analysis (Loo *et al*, 1996). Styrene maleic acid lipid particles (SMALPs) allow detergent-free extraction of membrane proteins in their local lipid environment and provide a promising technique for investigating receptor-ligand interactions under native conditions (Lee *et al*, 2016). This is particularly important since loss of lipids surrounding membrane proteins can lead to changes in measured binding affinities (Martens *et al*, 2018; Gault *et al*, 2020). The combination of detergent free SMALPs extraction coupled with mass spectrometry analysis provides a potential route for characterising native membrane receptor complexes (Sobotzki *et al*, 2018; Kalxdorf *et al*, 2021).

Here we report the results from a combined genetic and biochemical analysis of *D. melanogaster* nAChRs *in vivo*. Using CRISPR/Cas9 genome engineering we generated new null mutations for all ten receptor subunit genes in a uniform genetic background as well as introducing a fluorescent protein tag into the *nAChRα6* locus. We show that the null mutants in all seven α-subunit genes and two of the three β-subunit genes are viable and fertile, although we find mild morphological defects and some neurological impairment. Mutation of the remaining subunit gene, *nAChRβ1,* is recessive lethal. All nine of the viable null mutants were used to demonstrate a novel selective insecticidal effect of α-Btx on the *nAChRα5, nAChRα6* and *nAChRα7* subunits. We also applied the insecticidal Hv1a peptide to the viable null mutants, showing resistance with two subunit gene mutants: *nAChRα4* and *nAChRβ2*. In our biochemical studies we analysed receptor-ligand interactions in native conditions using SMALPs to verify the *in vivo* receptor subunit composition of the α-Btx binding target in adult neural tissue from wild-type and receptor subunit mutants. Our analysis revealed binding of α-

Btx to receptors containing Dα5, Dα6 and Dα7 subunits with the analysis of mutants in these subunits genes indicating heterogeneity in α-Btx binding nAChRs. Furthermore, we have identified specific glycosylation sites in Dα5 and Dα7 subunits which are known from other studies to play a critical role in α-Btx binding affinity (Dellisanti *et al*, 2007; Rahman *et al*, 2020). Localization studies with the Dα6 subunit tagged at the endogenous locus with a fluorescent reporter shows expression at different developmental stages in specific neuronal cells, including the Kenyon cells of the mushroom bodies, a known site of α-Btx-binding.

Results

New *D. melanogaster* nicotinic acetylcholine receptor subunit gene mutations

To investigate the role of individual nAChR subunits we used CRISPR/Cas9 to generate deletion mutations in each of the seven α-subunit and three β-subunit genes. All of the mutations were generated in virtually identical genetic backgrounds using nanos-Cas9 sources on the second or third chromosome of otherwise genetically homogeneous fly lines. In brief, for each gene we targeted exons shared between all predicted isoforms, close to the N-terminus of the protein. In order to disrupt each coding sequence and facilitate screening we introduced a visible fluorescent marker, DsRED under control of the eye-specific 3xP3 promoter at the targeted locus. Positive lines were confirmed by PCR and sequencing, and subsequently the DsRED marker was excised from the genome by Cre-Lox recombination.

For nine out of ten subunit genes we established homozygous viable and fertile stocks, the exception was the *nAChRβ1* gene which proved to be recessive lethal. Although all the other lines are viable, we noticed that most of the mutants, but particularly *nAChRα1, nAChRα2, nAChRα5* and *nAChRβ3*, exhibited a curled abdomen phenotype that is most prominent in males (approximately 25, 20, 15 and 15 % respectively, Fig 1A). It is possible that this phenotype is a result of defects in neural control of abdominal muscles and it is interesting to note that a previous analysis of an *nAChRα1* allele reports reduced male courtship and mating (Somers *et al*, 2017).

Since nAChRs are mostly found in the nervous system, we carried out basic climbing assays on the null alleles to assess potential locomotor defects (Fig 1B, Appendix Table S1). We saw little or no impact on the locomotor activity of ten day old flies with *nAChRα4, nAChRα5, nAChRα7, nAChRβ2* or *nAChRβ3* homozygous mutants, however, deletions of *nAChRα1*,

nAChRα2 and *nAChRα6* showed 50-60 % reductions in climbing ability compared to wild-type. In addition, the *nAChRα3* null mutant and heterozygotes for *nAChRβ1* exhibited a severe reduction in locomotor activity to less than 40 % of wild-type (22% and 34% respectively).

Taken together, we report the generation and validation of null mutations in all ten *D. melanogaster nAChR* subunit genes, with mild morphological defects associated with most of the new alleles and impaired locomotion observed with some mutants.

Distinct nAChR subunits mediate interactions with ω-Hexatoxin-Hv1a and α-Bungarotoxin

In order to investigate the selective contribution of each *nAChR* subunits to toxin binding *in vivo*, we injected 3[rd] instar larvae from the homozygous *nAChR* null mutants with either ω-Hexatoxin-Hv1a (Hv1a) or α-Bungarotoxin (α-Btx) dissolved in PBS. As a control, injections of PBS alone (vehicle) were performed in parallel, and all larvae survived the injection procedure and showed no detectable defects. Larval injection of 2.5 nmol/g Hv1a induced locomotor paralysis and full lethality in the control groups (*w[1118]*, *THattP40* and *THattP2*, Fig 2A, Appendix Table S2). Survival was quantified as the percentage of pupae formed after injection. Hv1a did not result in full lethality with *nAChRα4* and *nAChRβ2* homozygous mutants, since both showed an increase in survival to 42±22% (One-way ANOVA followed by Bonferroni's test, P=0.0035, Fig 2A). Mortality in all the other null mutants was comparable to controls (P>0.9).

We also observed significant toxicity following injection of 1.25 nmol/g α-Btx, with larvae exhibiting a progressive reduction in locomotion until stationary, resulting in developmental arrest and death. We found that α-Btx induced lethality is drastically reduced in the *nAChRα5*, *nAChRα6* and *nAChRα7* subunit mutants, with the survival rate significantly increased from 0% (controls) to 61±10% (P=0.001), 53±24% (P=0.0051) and 72±25% (P=0.0001) respectively (One-way ANOVA followed by Bonferroni's test, Fig 2B).

Together, these results indicate that Hv1a and α-Btx do not share the same binding target and differentially interact with the nAChR subunits *in vivo*. Since α-Btx showed a novel insecticidal effect on nAChRs we further examined its interactions biochemically.

Forming SMA-lipid particles (SMALPs) of ring-like nAChR complex structures

In order to take advantage of our new receptor subunit mutants for the biochemical analysis of native nAChR functions, we examined the composition of the receptors responsible for binding α-Btx. To address the functionality of *D. melanogaster* nAChRs isolated from endogenous membranes, we utilised detergent-free SMALPs extraction to characterise the interaction between receptor native lipid discs and the α-Btx toxin (Fig 3A).

In brief, we prepared membrane extracts from adult *D. melanogaster* heads (Depner *et al*, 2014) and generated lipid particle discs by solubilising the membrane extracts with the SMA copolymer. We used affinity beads coupled to α-Btx (Wang *et al*, 2003; Mulcahy *et al*, 2018) to enrich for nAChRs in the SMALP preparations that bound to the toxin, and performed mass spectrometric analysis of tryptic peptides generated from the enriched preparations. In parallel we processed membrane extracts without SMALP and with SMALP extracts enriched with beads alone.

We first determined whether membrane protein discs are formed from enriched membranes using the SMA copolymer. We prepared membrane enriched fractions from adult heads, solubilized these with SMA and separated the insoluble particles from the lipid discs by ultracentrifugation. We negatively stained the SMALP preparations and imaged them with transmission electron microscopy (TEM), observing irregular discs of varying shapes and sizes, with clusters containing different numbers of discs (Fig 3B).

Membrane receptors often have a unique shape in TEM images and the five subunits of a nAChRs is expected to form a ring-like structure, suggesting that the receptors are extracted as a complex. However, we did not observe pentameric ring-like structures perhaps suggesting that nAChRs are of low abundance and that analysis may benefit from enrichment. We coupled α-Btx to affinity beads to enrich nAChR complexes that bind the toxin in SMALP preparations (Fig 3C). In contrast to the unenriched samples, TEM images of the enriched preparations showed increased numbers of ring-like structures of 15 nm in diameter (Fig 3D and E). Thus our TEM analysis shows an increased number of ring-like membrane complexes in the SMALP preparations which are likely to be nAChRs.

Efficient SMALPs extraction allow to study nAChR subunits solubility

To assess to what extent the SMA copolymer solubilized nAChRs, we performed a bottom-up proteomics analysis to identify receptor subunits. Membrane preparations were solubilized in buffer with or without SMA, and affinity beads with or without α-Btx were used to assess ligand-binding to nAChR subunits. Comparing the number of proteins identified in samples solubilized either with or without 5% SMA, we observed a significantly increased identification rate of proteins dissolved in SMA by equal numbers of MS/MS spectral counts (two-tailed t-test, $P<0.01$, Fig 4A and non-significant, Fig 4B). This indicates that mass spectrometer performance was comparable during the measurements.

Sequences of membrane spanning segments of nAChR subunits, which are in close contact to the hydrophobic lipid environment, are largely composed of nonpolar side chains. Determining the average of hydrophobicity of identified protein sequences revealed significantly increased numbers of proteins with a positive hydrophobicity score in samples solubilized in SMA (two-tailed t-test, $P<0.0001$, Fig 4C), indicative of enrichment of membrane proteins. An analysis of Gene Ontology (GO) slim terms supports the conclusion that the SMALP preparations are enriched of membrane embedded and associated proteins (Fig 4D), and that these are not limited to plasma membrane proteins. In the SMA-enriched samples we found enrichment for proteins annotated with metabolic and catalytic activity terms and also enhanced response to biological stimuli (Appendix Fig S1A and B), highlighting the recovery of membrane-associated proteins.

Next, we focused on identified membrane proteins predicted to contain transmembrane helical (TMH) domains and found an increased number of proteins containing TMHs in SMA solubilized samples (Fig 4E). While the majority of these proteins contained a single TMH domain, we identified Piezo, a mechanosensory ion channel protein containing 37 predicted transmembrane helices. Both α- and β-nAChR subunits contain four TMH domains and could be solubilized in SMA. The number of β-barrel membrane spanning proteins identified was also significantly increased by SMA extraction (two-tailed t-test, $P<0.0001$, Appendix Fig S1C).

In addition, palmitoylated lipid anchor modifications to nAChR subunits has been shown to be important for receptor assembly into membranes and the formation of functional complexes (Alexander *et al*, 2010). Comparing samples solubilized with and without SMA showed a significantly increased identification of proteins which are predicted to be palmitoylated and

myristoylated (two-tailed t-test, $P<0.0001$, Appendix Fig S1D and E). In contrast, membrane

proteins that are predicted to contain a glycosylphosphatidylinositol (GPI)-anchor are equally

solubilized in both conditions (two-tailed t-test, non-significant, Appendix Fig 1F).

Focusing on the membrane receptors solubilized by SMA, we analysed the amino acid

sequence properties of identified proteins and calculated an overall solubility score (Sormanni

et al, 2015; Sormanni *et al*, 2017). Comparing the solubility to the hydrophobicity showed a

calculated R^2 of 0.56 (Fig 4F). Sequences with a score greater than 1 are highly soluble

receptors and those less than minus -1 are difficult to solubilize. As a result, samples solubilized

in SMA contain more receptors, which are difficult to solubilize. These receptors are more

hydrophobic and contain larger numbers of TMH domains (Fig 4G). Calculating an average

solubility score of -2.76 for nAChR sequences indicates that difficult to solubilize subunits are

successfully recovered with SMA (Fig 4H).

Taken together, these analyses confirm that SMA solubilizes nAChR complexes in a state

suitable for subunit identification by mass spectrometry and suggests that α-Btx interactions

can be studied with SMALP preparations.

259 Three nAChR α-subunits are targets of α-Btx

To explore native nAChR subunit interactions with α-Btx we searched for peptides from

subunit ligand-binding and cytoplasmic domains, identifying the Dα5, Dα6 and Dα7 subunits

in the α-Btx affinity bead pull-downs (Fig 5A and B, Appendix Table S3). Several other

nAChR subunit peptides could be identified in the negative controls performed without

coupling α-Btx to affinity beads (Appendix Table S4). The sequences of the ligand-binding

domains of the Dα5, Dα6 and Dα7 subunits are very similar (avg. 95.49 %) and we identified

peptides common to all three subunits (Appendix Fig S2A) as well as unique peptides within

their cytoplasmic domains (Appendix Fig S2B). However, we found no evidence of peptides

mapping to TMH domains. The ligand-binding domain of α-subunits show structural similarity

across different species (Appendix Fig S3A) and by mapping the identified peptides to known

structures we concluded they are most likely outside of the α-Btx binding sites (Appendix Fig

S3B).

To further characterize the role of the three α-subunits identified in α-Btx binding we generated

SMALP preparations and performed α-Btx affinity bead enrichments with adult head

preparations from homozygous null mutations for each of the *nAChRα5*, *nAChRα6* and

nAChRα7 subunit genes. With all three deletion mutants we observed, as expected, no

detectable peptides from the missing subunit but could still identify peptides from the other

two subunits (Fig 5C).

We compared the repertoire of proteins identified with α-Btx enrichment in wild-type with

those found in each of the three mutant lines to identify any changes in the representation of

biological pathways annotated in KEGG (Kanehisa *et al*, 2020, Fig 5D). While the enrichments

in wild-type and the mutants were broadly similar, we noticed a loss of proteins associated with

cofactor/vitamin metabolism, particularly retinol and ascorbate, in all three of the mutants as

well as proteins associated with vesicular transport. It is possible that these pathway changes

represent alterations in neurotransmitter production or trafficking. Interestingly, we also

noticed specific enrichment of cytochrome P450 related pathways in the *nAChRα6* mutants,

suggesting perturbation of neurotransmitter pathways.

In summary, our analysis indicates that a functional α-Btx binding nAChR involves the Dα5,

Dα6 and Dα7 subunits. This is entirely in line with our genetic findings described above, where

loss of each of these subunit genes conferred substantial resistance to α-Btx induced lethality.

Glycosylation sites of nAChR subunits by α-Btx binding

We next examined glycosylation sites on nAChR subunits since these are known to have an

important role in α-Btx binding affinity in other systems. For example, deglycosylation reduces

α-Btx binding in human nAChRs by more than two orders of magnitude (Dellisanti *et al*, 2007)

and α-Btx binding to loop C in *Torpedo californica* α-subunits is enhanced by N-glycosylation

of sites in these regions (Rahman *et al*, 2020). To identify specific glycosylation sites in *D.*

melanogaster nAChRs we first purified SMALP solubilized receptors with α-Btx affinity

beads, digested them into peptides and enriched for glycopeptides using HILIC resin

(Hägglund *et al*, 2004, Fig 6A).

Site-specific identification of glycans on peptides by mass spectrometry is challenging (Fang

et al, 2020) and often requires an additional deglycosylation step for glycopeptide

measurement. Deglycosylation of enriched peptides was carried out using two separate

enzymes: Endoglycosidase H (Endo H), which cleaves asparagine-linked oligosaccharides to

generate a truncated sugar molecule with one N-acetylhexosamine (HexNAc) residue, and the

endoglycosidase PNGase F, which releases the entire glycan from asparagine residues and

deaminates the sugar free asparagine to aspartic acid. While very few glycopeptides were

observed in the flow through (an average 20 glycopeptides Fig 6B), we identified a total of 397 glycopeptides after enrichment and deglycosylation with Endo H or PNGase F (Fig 6C).

Shared glycopeptides from Dα5 and Dα7 nAChR subunits were identified after enrichment and deglycosylation with Endo H or PNGase F (Fig 6D). Deglycosylation with Endo H identified modified asparagine (N2) residues on the peptide (NNGSCLYVPPGIFK), which is predicted to be part of the Dα5 and Dα7 ligand-binding domains involved in α-Btx binding. This asparagine residue was modified with an N-acetylhexosamine (HexNAc) truncated sugar chain. Releasing N-glycans after deglycosylation by PNGase F enabled us to identify a deaminated asparagine residue in the same peptide. The monoisotopic mass of this peptide changed due to the different modifications on the asparagine residue (Fig 6E).

The genome of *Caenorhabditis elegans* encodes for at least 29 nAChR subunits (Jones *et al*, 2007). The alpha-type unc-63 subunit contains an N-linked HexNAc modified asparagine residue on position 136 (Kaji *et al*, 2007). Performing a multiple sequence alignment showed that this asparagine residue is conserved between insects and nematodes (Appendix Fig S4A). Comparing identified glycosylation sites of Dα5 and Dα7 subunits to known N-linked glycosylation sites of α-subunits from *T. californica, Danio rerio, Mus musculus* or *Homo sapiens* indicates that this site is not conserved between vertebrates and invertebrates (Appendix Fig S4B).

We also identified glycosylation sites in the Dα3 (ATKATLNYTGR) and Dβ3 (VVLPENGTAR) subunits after Endo H treatment but not with PNGase F treatment, suggesting they harbour a single N-linked HexNAc modified asparagine residue (Appendix Fig S4C).

Taken together these findings suggest that the Dα5 and Dα7 subunits are modified at asparagine residues in the α-Btx ligand-binding domain with an N-linked sugar chain.

Localization of Dα6 nAChRs subunit in the brain

In order to examine the endogenous localization of an α-Btx binding receptor subunit we used CRISPR/Cas9 genome engineering to introduce in frame C-terminal fluorescence and epitope tags into the endogenous *nAChRα6* locus. The resulting line is homozygous viable and fertile, and shows no apparent phenotypes. We live imaged the unfixed brains of larvae and adults homozygous for the tagged line using confocal microscopy. In 2nd instar larvae we observed

low level well-distributed fluorescence signal throughout the ventral nerve cord (VNC), including on commissural axons, and in the developing brain (Fig 7A).

By early L3, we found more defined localization in the VNC and developing mushroom bodies (Fig 7B and D), particularly noticeable in the Kenyon cells, a known site of α-Btx binding (Su & O'Dowd, 2003). Localization in larval mushroom bodies continued to evolve, with defined expression in the Kenyon cells, calyx, peduncle, dorsal and medial lobes as well as the medulla and lamina of the emerging optic lobes (Fig 7C and E). We also observed localisation to a number of cell bodies overlying the optic lobes (Fig 7F).

Finally, in the adult brain, expression was largely restricted to the mushroom bodies particularly the Kenyon cells and connections across the midline between the β and γ lobes and the optic lobes (Fig 7G). The temporal localization of Dα6 subunit in the CNS is summarized in schematic form (Fig 7H).

Discussion

Elucidation of complex insect nAChRs heterogeneity will lead to a better understanding of selective insecticidal effects. We present a new set of null mutations in all *D. melanogaster nAChR* subunit genes and investigated insecticidal peptide toxin effects on wild-type and receptor subunit mutant larvae. Utilising biochemical approaches with SMALP pull-downs we characterised toxin binding and subunit composition of native nAChR complexes.

Our genome engineering approach generated viable and fertile mutations in nine out of the ten subunit genes encoded in the *D. melanogaster* genome and is largely concordant with the recently described work by Perry and colleagues (Perry *et al*, 2021). In both studies, null mutations in the *nAChRβ1* gene were inviable as stocks. We add to the previous work by generating viable mutations in *nAChRα5*. We observed some minor morphological defects in some of the null mutants especially in *nAChRα1, nAChRα2, nAChRα5* and *nAChRβ3* as well as locomotor defects with some alleles, particularly severely in *nAChRα3* homozygotes and *nAChRβ1* heterozygotes. The locomotor defects we observed are in agreement with previously reported neuronal phenotypes with *nAChR* subunit genes, including sleep disruption, defective jump response, memory impairment or locomotor defects (Fayyazuddin *et al*, 2006; Rohde *et al*, 2016; Somers *et al*, 2017; Tackenberg *et al*, 2020).

We used the nAChR null mutants to study insecticidal effects of the Hv1a peptide on viability

after injection into larvae and investigated whether α-Btx has any insecticidal properties. As

described by Chambers and colleagues, we confirm that Hv1a effects nAChRs (Chambers *et*

al, 2019) and our analysis shows that the Dα4 and Dβ2 subunits are involved in the insecticidal

response. We show for the first time that α-Btx has selective insecticidal effects against the

Dα5, Dα6 and Dα7 subunits, which we further characterized at the biochemical level.

The pharmacology of Hv1a and α-Btx binding has been shown to be distinctive (Chambers *et*

al, 2019), correlating with our demonstration that these two peptide toxins mediate their effects

through different receptor alpha subunits. Furthermore, resistance to neonicotinoid

insecticides, which interact most strongly with Hv1a binding, has been associated with Dβ2

(Perry *et al*, 2008; Perry *et al,* 2021), consistent with the involvement of this subunit in the

response to Hv1a. However, no resistance to neonicotinoids was seen in *D. melanogaster*

carrying a *nAChRα4* gene deletion (Perry *et al,* 2021), which could be explained if

neonicotinoids act at multiple receptor classes. Multiple binding sites for the neonicotinoid

imidacloprid can be resolved in equilibrium binding assays in many insect species (Xu *et al*,

2010) and by binding kinetics in flies (Liu & Casida, 1993).

Resistance to spinosad is strongly associated with Dα6 (Perry *et al,* 2021), and spinosad binding

is much more sensitive to the action of α-Btx than to the action of neonicotinoids (Chambers

et al, 2019), again consistent with the involvement of this subunit with sensitivity to injected

α-Btx and with the proposition that α-Btx and Hv1a act at distinct receptor classes.

nAChR subunits are known to be difficult to purify due to solubilisation issues (Cheng *et al*,

2015; Maldonado-Hernández *et al*, 2020) and the requirement for a lipid environment for

ligand binding (Dacosta *et al*, 2013) makes it challenging to study these receptors in native

conditions. We used the SMALPs extraction method for preparing membrane discs and

enriched nAChRs via α-Btx affinity purification. Electron microscopy analysis indicated that

receptor-like particles were recovered and these were substantially enriched by α-Btx pull-

down. Mass spectrometry analysis showed an enrichment for the Dα5, Dα6 and Dα7 subunits

in these preparations, which is concordant with our *in vivo* injection results and previous studies

that characterised aspects of α-Btx binding (Lansdell & Millar, 2004; Wu *et al*, 2005; Lansdell

et al, 2012).

To our knowledge this is the first report of the identification of a native endogenous α-Btx

binding nAChRs. We note however, that we cannot determine from our analysis whether all

three identified subunits are part of the same complex or if there are different receptors containing a subset of these subunits. Using chimeric receptors in a cell line system, Landsdell and colleagues reported that a combination of all three of these subunits show high affinity acetylcholine binding but α-Btx binding varied depending on receptor combinations, with Dα5 and Dα6 binding most strongly (Lansdell *et al*, 2012). In a prior study they implicated Dα6 and Dα7 (Lansdell & Millar, 2004). However, these assays were performed with 5HT3A-nAChR subunit fusions, here we provide strong evidence that these three subunits bind to α-Btx *in vitro* and *in vivo*.

In addition, glycopeptide enrichment showed site specific glycosylation modifications on the Dα5 and Dα7 nAChR subunit ligand binding domains. The unique lipid environment and glycosylation sites of nAChR α-subunits from the electric ray, *T. californica*, were found to be important for α-Btx binding activities (Quesada *et al*, 2016; Rahman *et al*, 2020), and structural studies support this conclusion (Dellisanti *et al*, 2007). Our work supports the view that there is a role for Dα5 and Dα7 glycosylation modifications in the recognition of α-Btx in *D. melanogaster*.

Our localization studies with fluorescence tagged endogenous Dα6 subunit showed relatively restricted expression in the brain and ventral nerve cord, with prominent expression in the Kenyon cells of the mushroom body, all known regions. The expression of Dα6 in Kenyon cells across development is in line with a proposed role for this subunit in memory plasticity, along with other α-subunits including Dα5, in mushroom body output neurons (Barnstedt *et al*, 2016). Thus, it is possible that retention of α-Btx binding in the absence of Dα6 may simply reflect its restricted localisation. In contrast, it is clear that Dα6 plays a major and specific role in binding to the insecticide spinosad in *D. melanogaster* since mutations in this subunit are highly resistant to the toxin (Perry *et al*, 2015).

Localization studies of Dα6 nAChRs subunit fusion protein by confocal microscopy are largely consistent with recent reports of *nAChRα6* expression derived from expression reporters (Kondo *et al*, 2020), though these studies appear to indicate wider adult brain expression than we observed, perhaps reflecting a degree of translational control or limitations in the sensitivity of our live imaging.

In conclusion, we identified ligand-binding subunit sites for a *D. melanogaster* nAChR antagonist with newly insecticidal effects. Our findings contribute to a better understanding of the role of nAChR subunits which interacts with insecticidal peptide toxins.

Materials and methods

Drosophila methods

Embryos were injected using standard procedures into the *THattP40* (*y^1 sc v^1 sev^{21}; P{$y^{+t7.7}$ $v^{+t1.8}$ nos-Cas9.R}attP40*) or *THattP2* (*y^1 sc v^1 sev^{21}; P{$y^{+t7.7}$ $v^{+t1.8}$ nos-Cas9.R}attP2*) lines expressing *nos*-Cas9 (Bloomington *Drosophila* Stock Centre). Donor DNA (500 ng/μL) in sterile H_2O was injected together with of gRNA plasmids (100 ng/μL) as described previously (Korona *et al*, 2020). Individually selected surviving adults were crossed to w^{1118} and the progeny screened for DsRED fluorescence localized mostly to the eyes of transgenic flies: positive flies were balanced and homozygous stocks established where possible. The correct localization of the insert was confirmed via PCR and sequencing. Transgenic flies were assessed for the phenotype using bright field microscope. For tagging of *nAChRα6*, the stocks were additionally subjected to Cre-recombination for marker removal and several independent lines were verified by PCR. Some of these lines were screened for YFP fluorescence using confocal microscopy. From the YFP positive balanced stocks, the viable and fertile homozygote was established. Injections were performed by the Department of Genetics Fly Facility (https://www.flyfacility.gen.cam.ac.uk). All fly stocks were maintained at 25°C on standard cornmeal medium. Larvae of 2^{nd} and 3^{rd} stage were collected, and their brains were dissected according to standard protocols. Brains were mounted in glycerol and live imaged.

Cloning of gRNAs and generation of donor vectors

Construction of *nAChR* subunits null alleles

In order to generate individual *nAChR* subunits gene deletions the open reading frame (ORF) was disrupted by introducing a visible marker harbouring DsRED marker under eye specific driver 3Px3 using CRISPR/Cas9 technology as previously described (Korona *et al*, 2020). The targeted exons are shared between different isoforms and adjacent to the N-terminus to ensure the protein translated was interrupted. The insertion sites were designed *in silico* and optimal gRNAs were chosen (Appendix Table S5) that were tested against the injection strain and cloned into pCDF3. Briefly, target specific sequences were synthesized and either 5′-

phosphorylated annealed and ligated into the *Bbs*I sites of pCDF3 precut with *Bbs*I. Positive clones were confirmed by sequencing.

For generation of donor vectors, firstly, homology arms were amplified on genomic DNA (Appendix Table S6) that, secondly, were used as a template to amplify the homology arms (Appendix Table S7) of the donor vector for CRISPR/Cas9 homologous recombination (HDR). The inserts with visible marker were amplified using as a template previously generated constructs (Korona *et al*, 2020) with appropriate primers. These fragments were used for Gibson assembly using Gibson Assembly Master Mix (New England Biolabs). PCR products were produced with the Q5 High-Fidelity 2X Master Mix (New England Biolabs). All inserts were verified by sequencing.

C-terminal tagging of Dα6 nAChRs subunit fusion protein

For tagging of Dα6 nAChRs subunit the C-terminal fusion with FSVS fluorescent protein harbouring StrepII and 3xFLAG epitope tags (3xFLAG-StrepII-Venus-StrepII) was generated for CRISPR/Cas9 mediated genome engineering (Korona *et al*, 2017; Korona *et al*, 2020). Firstly, gRNAs were designed (Appendix Table S5) and tested against the genomic DNA sequence of injection strains. The oligonucleotides were phosphorylated and ligated into *Bbs*I pre-cut pCDF3. The positive variants were confirmed by sequencing.

The donor vector to generate protein fusion with fluorescent protein harbouring epitope tags was cloned in 2 steps strategy by creating initially (A) nAChRα6-FSVS donor and then adding the removable marker to generate (B) nAChRα6-FSVS-loxP-3PX3_DsRED_loxP donor vector. At first, the homology arms were enriched on genomic DNA (Appendix Table S6) and used to amplify homology arms for donor vector nAChRα6-FSVS (Appendix Table S7) that was assembled using Gibson Assembly as described above. The FSVS tag was amplified on previously generated constructs (Korona *et al*, 2017) with appropriate overlapping oligonucleotides (Appendix Table S7). The construct was confirmed by Sanger sequencing and used as a template to generate donor vector with removable marker. The PCR fragments harbouring homology arms and FSVS tag were amplified on nAChRα6-FSVS construct, whereas the 3PX3-DsRed with adjacent loxP sites was amplified using earlier generated constructs (Korona *et al*, 2017). The final donor vector was generated using Gibson Assembly® as described above and positive variants were confirmed by sequencing.

Confocal microscopy

Localization of FSVS-tagged (3xFLAG-StrepII-Venus-StrepII) Dα6 nAChRs subunit was visualised in dissected larvae brains via monitoring the YFP fluorescence (Venus). Briefly, the larval brains were dissected and mounted in glycerol for live imaging. Images were acquired using a Leica SP8 confocal microscope (Leica microsystems) with appropriate spectral windows for mVenus, images were processed with Fiji software.

Locomotor behaviour

Adult female and male flies were collected shortly after eclosion and separated into 10 cohorts consisting of 10 flies (100 total) for each genotype. Flies were maintained at 25°C and transferred to fresh food every three days. For the climbing assay, each cohort was transferred to 10ml serological pipette, and allowed to acclimatize for five min. For each trial, flies were tapped down to the bottom of the vial, and the percentage of flies able to cross a five-ml mark successfully within 10 seconds was recorded as the climbing index. Five trials were performed for each cohort, with a 1-min recovery period between each trial. Climbing assays were performed 10 days after eclosion.

Drosophila larval injections

Injections were performed by using the Nanoliter 2000 (World Precision Instruments, Hertfordshire, United Kingdom) mounted on a micromanipulator (Narishige, London, United Kingdom). Micropipettes were pulled from glass capillary tubes (1.14 mm OD, 0.530 mm ± 25 μm ID; #4878, WPI) using a laser-based micropipette puller (Sutter P-2000, Sutter Instrument, Novato, CA, USA). Third instar larvae (wandering stage) were transferred to an adhesive surface after being quickly washed with water to remove food residues and gently dried using paper tissue. The micropipette was positioned over the approximate centre of the body, on the dorsal side, and the tip was advanced through the cuticle into the hemocoel of the larva. Larvae were injected with 69 nL of PBS (phosphate-buffered saline) supplemented with 10% (v/v) filtered food dye (PME, moss green food colouring; 0.2 µm filter). Food dye was included to aid in monitoring the success of the injection under a dissection microscope (Leica MZ65, Milton Keynes, United Kingdom). ω-hexatoxin-Hv1a (Hv1a, Syngenta, Schaffhauserstrasse, CH-4332 Stein, Switzerland) and α-Bungarotoxin (α-Btx, ab120542, Abcam, Cambridge, United Kingdom) were added to the injection mix in order to obtain a final concentration of 2.5 nmol/g and 1.25 nmol/g, respectively (average larval weight was 2.14 mg).

After injection, larvae were then gently transferred into agar/grape juice (Ritchie Products Limited, Burton-On-Trent, United Kingdom) plates and kept at 25°C. The rate of survival (expressed as percentage) was calculated as the number of living pupae, formed 1-2 days after injection, divided by the total number of injected larvae. Experiments were repeated three times independently with a total number of 10 larvae for each experimental group. Results were analysed with One-way ANOVA followed by Bonferroni's multiple comparisons test using GraphPad Prism (version 7, GraphPad Software, San Diego, California, USA).

Coupling procedure of α-Bungarotoxin to affinity beads

Coupling of α-Bungarotoxin, α-Btx (ab120542, Abcam, Cambridge, United Kingdom) to cyanogen bromide-activated (CNBr) sepharose beads 4B (C9 142-5G, Sigma-Aldrich, Haverhill, United Kingdom) was performed as described (Wang *et al*, 2003; Mulcahy *et al*, 2018). CNBr-activated sepharose 4B beads (0.25 g) were hydrated in 1.25 ml of 1 mM HCl for 1 hr at 4°C on a rotator. Beads were centrifuged for 5 min at 1500 × g, the supernatant removed and beads washed twice with 1 ml of coupling buffer (0.25 M NaHCO$_3$, 0.5 M NaCl, pH 8.3). Beads were centrifuged for 5 min at 1500 × g and the supernatant was removed. Alpha-Btx (1 mg) was resuspended in 1 ml coupling buffer and incubated together with the affinity beads at 4°C for 16 hr on a rotator. Beads were centrifuged for 5 min at 1500 × g. Coupling efficiency was determined using a Pierce^{TM} quantitative fluorometric peptide kit and used according to the manufacturer's instructions (23290, Thermo Scientific^{TM}, Bishop's Stortford, United Kingdom). Beads were blocked with 1 ml of 0.2 M glycine in 80 % coupling buffer at 4°C for 16 hr on a rotator. Beads were then centrifuged for 5 min at 1500 × g and washed with 1 ml of 0.1 M NaHCO$_3$, 0.5 M NaCl, pH 8.0. This step was repeated with 1 ml of 0.1 M NaCH$_3$CO$_2$, 0.5 M NaCl, pH 4.0. Beads were washed again in 1 ml of 0.1 M NaHCO$_3$, 0.5 M NaCl, pH 8.0. After a final wash step with 1 ml coupling buffer the beads were incubated twice for 30 min in 1 ml Tris-buffer (50 mM Tris, 150 mM NaCl, pH 8.0). The beads were centrifuged for 5 min at 1500 × g, the supernatant was removed and 20 µl Tris-buffer, pH 8.0 was added.

Membrane protein enrichment and incorporation in SMALPs

D. melanogaster heads were obtained and separated according to (Depner *et al*, 2014). In a 50 ml falcon tube approximately 6 g flies were rapidly frozen in liquid nitrogen and vortexed twice for 3 min, with the tube cooled for 30 sec in liquid nitrogen between. Heads were separated from bodies by sieving (1201124 & 1201125, Endecotts, London, United Kingdom). 1 ml of

isotonic lysis buffer (0.25 M sucrose, 50 mM TRIS/HCl pH 7.4, 10 mM HEPES pH 7.4, 2mM EDTA, Protease inhibitor) was added to approximately 0.8 g separated heads. The solution was mixed three times by vortexing and the heads were lysed with 60 strokes in a Dounce homogenizer with a pestle. Membrane protein preparation was performed by differential centrifugation-based fractionation as described (Depner *et al*, 2014; Geladaki *et al*, 2019). Membrane protein pellets were resuspended in 20 to 100 μl 5 % SMALP solution (5 % styrene maleic acid copolymer (3:1), 5 mM Tris-Base, 0.15 mM NaCl, pH 8.0). For efficient incorporation and formation of SMALPs, membrane proteins were incubated with 5 % SMALP solution for 2 hr at room temperature on a rocking platform. To separate the insoluble proteins from the soluble SMALPs a centrifugation step at $100000 \times g$ for 60 min, 4°C was performed. Supernatant containing the SMALPs was combined and used for the nAChRs pull-downs.

Enrichment of nAChRs by α-Btx pull-down

SMALPs (20-35 mg/ml) were incubated with 200 μl α-Btx conjugated affinity beads for 16 hr, 4°C on a rotator. The beads were then centrifuged for 5 min at $1500 \times g$ and washed two or three times, each for 10 min with 1 ml ice-cold TBS (50 mM Tris, 150 mM NaCl, pH 8.0) on a rotator at 4°C. Beads were centrifuged for 5 min at $1500 \times g$ and nAChRs selectively eluted twice with 100 μl 1 M carbachol (CAS 51-83-2, Insight Biotechnology Ltd, Wembley, United Kingdom). These steps were performed for 25 min at room temperature on a rotator. Beads were centrifuged for 5 min at $1500 \times g$ and eluates were combined and ice-cold 100 % acetone in the volume of four times of the sample was added to the samples, mixed by vortexing and proteins were precipitated for 16 hr at -20°C. Samples were centrifuged at $13000 \times g$ for 15 min. Supernatant was removed and dried proteins were dissolved in Laemmli buffer (1M Tris pH 6.8, 10 % SDS, 5 % glycerol, 2 % bromophenol blue). Proteins were heated at 60°C and loaded on Mini-Protean TGX precast gels (456–1084, 4-15 %, Bio-Rad Laboratories, Inc., Watford, United Kingdom).

Electron microscopy preparation

For negative staining analysis, membrane proteins were extracted with 5 % SMA and nAChRs were enriched using α-Btx affinity pull-downs. Proteins were diluted 1:10 with deionised water to approximately 0.9 mg/ml and an aliquot of the samples were absorbed onto a glow-discharged copper/carbon-film grid (EM Resolutions) for approximately 2 min at room temperature. Grids were rinsed twice in deionised water and negative staining was performed using a 2 % aqueous uranyl acetate solution. Samples were viewed in a Tecnai G2 transmission

582 electron microscope (TEM, FEI/ThermoFisher) run at 200 keV accelerating voltage using a 20

583 µm objective aperture to increase contrast; images were captured using an AMT CCD camera.

Sample preparation for liquid chromatography–mass spectrometry (LC-MS)

The protein lanes were excised from the gels and proteolytic digestion with trypsin/lys-C mix

(V5073, Promega, Southampton, United Kingdom) was performed as described (Shevchenko

et al, 2007). The gel pieces were covered with 50 mM NH_4HCO_3 / 50 % ACN and shaken for

10 min. This step was repeated with 100 % acetonitrile and finally dried in a speed vac. Samples

were reduced with 10 mM DTT in 50 mM NH_4HCO_3 at 56°C for 1 hr and alkylated with 50

mM iodoacetamide in 50mM NH_4HCO_3 at room temperature without light for 45 min. The gels

were covered with 50 mM NH_4HCO_3 and 100 % ACN and shaken for 10 min. These steps were

repeated and samples were dried in a speed vac. Trypsin/lys-C buffer was added to the sample

according to manufacturer's instructions and incubated for 45 min on ice. Next 30 µl 25 mM

NH_4HCO_3 was added and samples were incubated at 37°C for 16 hr. The gel pieces were

covered with 20 mM NH_4HCO_3 and shaken for 10 min. Supernatant with peptides was

collected. Next, the gels were covered with 50 % ACN / 5 % FA and shaken for 20 min. These

steps were repeated and peptides were dried in a speed vac. Samples for glycopeptide

enrichment were digested in-solution according to (Queiroz *et al*, 2019). Samples were reduced

and alkylated in 10 mM DTT and 50 mM iodoacetamide. Proteins were digested in final

concentration of 2.5 µg trypsin/lys-C buffer for 16 hr at 37°C.

Peptide clean-up

Peptides were desalted using C-18 stage tips according to (Rappsilber *et al*, 2007). C-18

material (three C-18 plugs were pasted in a 200 µl pipette tip, Pierce^{TM} C18 Spin Tips, 84850

Thermo Scientific^{TM}, Bishop's Stortford, United Kingdom) was equilibrated with methanol/0.1

% FA , 70 % ACN/0.1 % FA and with 0.1 % FA. Peptides were loaded on C-18 material,

washed with 0.1 % FA and eluted with 70 % ACN/0.1 % FA. Samples were dried and finally,

peptides were resuspended in 20 µl 0.1 % FA. For glycopeptide enrichment peptides were first

desalted using poros oligo r3 resin (1-339-09, Thermo Scientific^{TM}, Bishop's Stortford, United

Kingdom) as described (Gobom *et al*, 1999; Queiroz *et al*, 2019). Pierce™ centrifuge columns

(SH253723, Thermo Scientific^{TM}, Bishop's Stortford, United Kingdom) were filed with 250 µl

of poros oligo r3 resin. Columns were washed three times with 0.1 % TFA. Peptides were

loaded onto the columns and washed three times with 0.1 % TFA and subsequently eluted with 70 % ACN.

Glycopeptide enrichment

Enrichment of glycopeptides of nAChRs was performed as described (Hägglund *et al*, 2004). Micro columns were prepared with 200 μl peptide tips filled with a C8 plug and iHILIC – fusion 5μm, 100 Å silica based material (HCS 160119, Hilicon, Umeå, Sweden). Peptides were solubilized stepwise in 19 μl dH$_2$O and then in 80 μl ACN plus 1 μl TFA acid. The micro columns were cleaned with 50 μl 0.1 % TFA and three times equilibrated with 100 μl 80 % ACN, 1 % TFA. Peptides were loaded onto the micro column and washed twice with 100 μl 80 % ACN, 1 % TFA. Glycopeptides were eluted from the column using twice 40 μl 0.1 % TFA and finally with 20 μl 80 % ACN, 1 % TFA. Samples were dried in a speed vac before peptides were deglycosylated with Endo H or PNGase F according to manufacturer's instructions (P07025 & P0710S, New England Biolabs Inc., Hitchin, United Kingdom).

LC-MS/MS

Peptide samples were dissolved in 20 μl of 0.1 % (v/v) FA. Approximately 1 μg peptide solution was used for each LC-MS/MS analysis. All LC-MS/MS experiments were performed using a Dionex Ultimate 3000 RSLC nanoUPLC (Thermo Fisher Scientific Inc, Waltham, MA, USA) system and a Q Exactive™ Orbitrap mass spectrometer (Thermo Fisher Scientific Inc, Waltham, MA, USA). Separation of peptides was performed by reverse-phase chromatography at a flow rate of 300 nL/min and a Thermo Scientific reverse-phase nano Easy-spray column (Thermo Scientific PepMap C18, 2μm particle size, 100A pore size, 75 μm i.d. x 50 cm length). Peptides were loaded onto a pre-column (Thermo Scientific PepMap 100 C18, 5μm particle size, 100A pore size, 300 μm i.d. x 5mm length) from the Ultimate 3000 autosampler with 0.1 % FA for 3 min at a flow rate of 15 μL/min. After this period, the column valve was switched to allow elution of peptides from the pre-column onto the analytical column. Solvent A was water + 0.1 % FA and solvent B was 80 % ACN, 20 % water + 0.1 % FA. The linear gradient employed was 2-40 % B in 90 min (the total run time including column washing and re-equilibration was 120 min). In between runs columns were washed at least four times to avoid any carryovers. The LC eluant was sprayed into the mass spectrometer by means of an Easy-spray source (Thermo Fisher Scientific Inc.). An electrospray voltage of 2.1 kV was applied in order to ionize the eluant. All *m/z* values of eluting ions were measured in an Orbitrap mass

analyzer, set at a resolution of 35000 and scanned between *m/z* 380-1500 Data dependent scans

(Top 20) were employed to automatically isolate and generate fragment ions by higher energy

collisional dissociation (HCD, Normalised collision energy (NCE): 25 %) in the HCD collision

cell and measurement of the resulting fragment ions were performed in the Orbitrap analyser,

set at a resolution of 17500. Singly charged ions and ions with unassigned charge states were

excluded from being selected for MS/MS and a dynamic exclusion of 20 seconds was

employed.

Peptide/protein database searching

Protein identification was carried out using sequest HT or mascot search engine software

operating in Proteome Discoverer 2.3 (Eng *et al*, 1994; Koenig *et al*, 2008). Raw flies were

searched against the uniprot *Drosophila_melanogaster_*20180813 database (23297 sequences;

16110808 residues) and a common contaminant sequences database. The search parameters

using mascot algorithm were: (i) trypsin was set as the enzyme of choice, (ii) precursor ion

mass tolerance 20 ppm, (iii) fragment ion mass tolerance 0.1 Da, (iv) maximum of two missed

cleavage sites were set, (v) a minimum peptide length of six amino acids were set, (vi) fixed

cysteine static modification by carbamidomethylation, (vii) variable modification by

methionine oxidation & deamidation on asparagine and glutamine and N-acetylhexosamine

(HexNAc(1)dHex(1) + HexNAc on asparagine) as variable glycopeptide modifications, (viii)

A site probability threshold of 75 % was set, (ix) Percolator was used to assess the false

discovery rate and peptide filters were set to high confidence (FDR<1).

Data handling and statistical analysis

Protein data evaluation was performed using R 3.5.3 (Ihaka & Gentleman, 1996). Plotting of

graphs were performed in RStudio 1.3.959 (Rstudio, 2020) using ggplot2 (Ginestet, 2011) and

other R packages. In order to characterise membrane proteins the following tools were used:

(i) TMHMM - 2.0 (Krogh *et al*, 2001), (ii) PRED-TMBB2 (Tsirigos *et al*, 2016) (iii) SwissPalm

(Blanc *et al*, 2015), (iv) PredGPI (Pierleoni *et al*, 2008), (v) Gravy calculator (www.gravy-

calculator.de), (vi) Myristoylator (Bologna *et al*, 2004) (vii) Solubility scores (Sormanni *et al*,

2015; Sormanni *et al*, 2017). Analysis of gene ontology (GO) slim terms (Carbon *et al*, 2019)

were performed within proteome discoverer 2.3 (Thermo Fisher Scientific). KEGG (Kanehisa

et al, 2020) pathway enrichment analysis was performed using DAVID (Huang *et al*, 2009).

For each experimental investigation n ≥ 3 were considered and data are represented as means

± SEM. Experiments were performed in a blinded manner whenever possible. Data are

presented as mean ± SD. Statistical tests for SMALPs were performed using two-tailed t-test
with an unequal variance and P values of ≤ 0.05 were considered to be significant. In DAVID,
Fisher's exact P values are computed to measure the gene-enrichment terms. Fisher's exact P
value of 0 represents perfect enrichment of a term. Usually P value of ≤ 0.05 are to be
considered as strongly enriched. In this study the default threshold set in DAVID of 0.1 was
used. Linear regression analysis was performed in order to study the efficiency of SMALPs
extraction of membrane receptors.

Structural assessment and illustration of nAChR subunits

For structural alignment of nAChRs matchmaker command operating in UCSF Chimera X 0.91
(Goddard *et al*, 2018) was used. This command is superimposing protein structures by first
creating pairwise sequence alignments, then fitting the aligned residue pairs and displays in an
overlaid structure as a result. The following parameters were set to create the aligned structure:
(i) alignment algorithm; Needleman-Wunsch (ii) similarity matrix; BLOSUM-62. Structural
animation was performed in Blender 2.8 (www.blender.org), an open-source 3D graphics
software. For annotation of protein sequences InterProScan was used (Mitchell *et al*, 2019).
Illustrator for biological sequences (IBS) web server was used to present biological sequences
(Liu *et al*, 2015). Multiple sequence alignments were performed (Madeira *et al*, 2019) or using
BoxShade multiple sequence alignments (Swiss institute of bioinformatics).

Data availability
694

695   The mass spectrometry data from this publication have been deposited to PRIDE
696   (http://www.ebi.ac.uk/pride/archive/) with the data set identifier PXD028484. Biochemical
697   source data is provided (Table EV1).

698   **Expanded View** for this article is available online.

699   **Acknowledgements**
700

We thank Professor Tim Dafforn for kindly providing us styrene maleic acid (SMA)
copolymer, Joao A. Paulo, Ph.D. for helpful exchange and Mrs Renata Feret for technical
discussions. We are very grateful to Syngenta and the Milner Therapeutics Institute for
excellent infrastructural support. Electron microscopy was performed using the facilities at

CAIC (Cambridge Advanced Imaging Centre, University of Cambridge). Funding was
provided by BBSRC (BB/P021107/1) and Syngenta.

Author Contributions

Conflict of interest

713 The authors declare that they have no conflict of interest.

Reference

715 Alexander JK, Govind AP, Drisdel RC, Blanton MP, Vallejo Y, Lam TT, Green WN (2010)
716   Palmitoylation of nicotinic acetylcholine receptors. *J Mol Neurosci* 40: 12-40

717 Barnstedt O, Owald D, Felsenberg J, Brain R, Moszynski JP, Talbot CB, Perrat PN, Waddell
718   S (2016) Memory-Relevant Mushroom Body Output Synapses Are Cholinergic. *Neuron*
719   89: 1237–1247

720 Blanc M, David F, Abrami L, Migliozzi D, Armand F, Bürgi J, van der Goot FG (2015)
721   SwissPalm: Protein Palmitoylation database. *F1000Research* 4: 261

722 Bologna G, Yvon C, Duvaud S, Veuthey AL (2004) N-terminal myristoylation predictions by
723   ensembles of neural networks. *Proteomics* 4: 1626-1632

724 Carbon S, Douglass E, Dunn N, Good B, Harris NL, Lewis SE, Mungall CJ, Basu S,
725   Chisholm RL, Dodson RJ, *et al* (2019) The Gene Ontology Resource: 20 years and still
726   GOing strong. *Nucleic Acids Res* 47: D330-D338

727 Chambers C, Cutler P, Huang YH, Goodchild JA, Blythe J, Wang CK, Bigot A, Kaas Q,
728   Craik DJ, Sabbadin D, *et al* (2019) Insecticidal spider toxins are high affinity positive
729   allosteric modulators of the nicotinic acetylcholine receptor. *FEBS Lett* 593: 1336-1350

730 Cheng H, Fan C, Zhang SW, Wu ZS, Cui ZC, Melcher K, Zhang CH, Jiang Y, Cong Y, Xu
731   HE (2015) Crystallization scale purification of α7 nicotinic acetylcholine receptor from
732   mammalian cells using a BacMam expression system. *Acta Pharmacol Sin* 36: 1013-
733   1023

734 Dacosta CJB, Dey L, Therien JPD, Baenziger JE (2013) A distinct mechanism for activating
735   uncoupled nicotinic acetylcholine receptors. *Nat Chem Biol* 9: 701-707

736 Dacosta CJB, Free CR, Sine SM (2015) Stoichiometry for α-bungarotoxin block of α7
737   acetylcholine receptors. *Nat Commun* 6: 8057

738 Dellisanti CD, Yao Y, Stroud JC, Wang ZZ, Chen L (2007) Crystal structure of the
739   extracellular domain of nAChR α1 bound to α-bungarotoxin at 1.94 Å resolution. *Nat*
740   *Neurosci* 10, 953-962

Denisov IG & Sligar SG (2016) Nanodiscs for structural and functional studies of membrane
proteins. *Nat Struct Mol Biol* 23: 481-486

Depner H, Lützkendorf J, Babkir HA, Sigrist SJ, Holt MG (2014) Differential centrifugation-
based biochemical fractionation of the *Drosophila* adult CNS. *Nat Protoc* 9: 2796-2808

Eng JK, McCormack AL, Yates JR (1994) An approach to correlate tandem mass spectral
data of peptides with amino acid sequences in a protein database. *J Am Soc Mass*
Spectrom 5: 976-989

Fang P, Ji Y, Silbern I, Doebele C, Ninov M, Lenz C, Oellerich T, Pan KT, Urlaub H (2020)
A streamlined pipeline for multiplexed quantitative site-specific N-glycoproteomics. *Nat*
Commun 11: 5268

Fayyazuddin A, Zaheer MA, Hiesinger PR, Bellen HJ (2006) The Nicotinic Acetylcholine
Receptor Dα7 Is Required for an Escape Behavior in*Drosophila*. *PLoS Biol* 4: e63

Gault J, Liko I, Landreh M, Shutin D, Bolla JR, Jefferies D, Agasid M, Yen HY, Ladds
MJGW, Lane DP, *et al* (2020) Combining native and 'omics' mass spectrometry to
identify endogenous ligands bound to membrane proteins. *Nat Methods* 17: 505-508

Geladaki A, Britovšek NK, Breckels LM, Smith TS, Vennard OL, Mulvey CM, Crook OM,
Gatto L, Lilley KS (2019) Combining LOPIT with differential ultracentrifugation for
high-resolution spatial proteomics. *Nat Commun* 10: 331

Ginestet C (2011) ggplot2: elegant graphics for data analysis. In *ggplot2: elegant graphics*
for data analysis, Wickham H (ed), 2nd edn, pp 1-245. New York, NY: Springer-Verlag

Gobom J, Nordhoff E, Mirgorodskaya E, Ekman R, Roepstorff P (1999) Sample purification
and preparation technique based on nano-scale reversed-phase columns for the sensitive
analysis of complex peptide mixtures by matrix-assisted laser desorption/ionization
mass spectrometry. *J Mass Spectrom* 34: 105-116

Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE (2018)
UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci*
27: 14-25

Hägglund P, Bunkenborg J, Elortza F, Jensen ON, Roepstorff P (2004) A new strategy for
identification of N-glycosylated proteins and unambiguous assignment of their
glycosylation sites using HILIC enrichment and partial deglycosylation. *J Proteome Res*
3: 556-566

Houchat JN, Dissanamossi BM, Landagaray E, Mathé-Allainmat M, Cartereau A, Graton J,
Lebreton J, Le Questel JY, Thany SH (2019) Mode of action of sulfoxaflor on α-
bungarotoxin-insensitive nAChR1 and nAChR2 subtypes: Inhibitory effect of
imidacloprid. *Neurotoxicology* 74: 132-138

Huang DW, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44-57

Ihaka R & Gentleman R (1996) R: A Language for Data Analysis and Graphics. *J Comput*
Graph Stat 5: 299-314

Ihara M, Furutani S, Shigetou S, Shimada S, Niki K, Komori Y, Kamiya M, Koizumi W,
Magara L, Hikida M, *et al* (2020) Cofactor-enabled functional expression of fruit fly,
honeybee, and bumblebee nicotinic receptors reveals picomolar neonicotinoid actions.

Proc Natl Acad Sci USA 117: 16283-16291

Jones AK, Davis P, Hodgkin J, Sattelle DB (2007) The nicotinic acetylcholine receptor gene family of the nematode Caenorhabditis elegans: An update on nomenclature. *Invert Neurosci* 7:129-31

Kaji H, Kamiie JI, Kawakami H, Kido K, Yamauchi Y, Shinkawa T, Taoka M, Takahashi N, Isobe T (2007) Proteomics reveals n-linked glycoprotein diversity in *Caenorhabditis elegans* and suggests an atypical translocation mechanism for integral membrane proteins. *Mol Cell Proteomics* 6: 2100-2109

Kalxdorf M, Günthner I, Becher I, Kurzawa N, Knecht S, Savitski MM, Eberl HC, Bantscheff M (2021) Cell surface thermal proteome profiling tracks perturbations and drug targets on the plasma membrane. *Nat Methods* 18: 84-91

Kanehisa M, Furumichi M, Sato Y, Ishiguro-Watanabe M, Tanabe M (2020) KEGG: integrating viruses and cellular organisms. *Nucleic Acids Res* 49: D545-D551

Koenig T, Menze BH, Kirchner M, Monigatti F, Parker KC, Patterson T, Steen JJ, Hamprecht FA, Steen H (2008) Robust prediction of the MASCOT score for an improved quality assessment in mass spectrometric proteomics. *J Proteome Res* 7: 3708-3717

Kondo S, Takahashi T, Yamagata N, Imanishi Y, Katow H, Hiramatsu S, Lynn K, Abe A, Kumaraswamy A, Tanimoto H (2020) Neurochemical Organization of the *Drosophila* Brain Visualized by Endogenously Tagged Neurotransmitter Receptors. *Cell Rep* 30: 284-297.e5

Korona D, Koestler SA, Russell S (2017) Engineering the *Drosophila* Genome for Developmental Biology. *J Dev Biol* 5: 16

Korona D, Nightingale D, Fabre B, Nelson M, Fischer B, Johnson G, Lees J, Hubbard S, Lilley K, Russell S (2020) Characterisation of protein isoforms encoded by the *Drosophila* Glycogen Synthase kinase 3 gene *shaggy*. *PLoS One* 15: e0236679

Krogh A, Larsson B, Von Heijne G, Sonnhammer ELL (2001) Predicting transmembrane protein topology with a hidden Markov model: Application to complete genomes. *J Mol Biol* 305: 567-580

Lansdell SJ, Collins T, Goodchild J, Millar NS (2012) The *Drosophila* nicotinic acetylcholine receptor subunits Dα5 and Dα7 form functional homomeric and heteromeric ion channels. *BMC Neurosci* 13: 73

Lansdell SJ & Millar NS (2004) Molecular characterization of Dα6 and Dα7 nicotinic acetylcholine receptor subunits from *Drosophila*: Formation of a high-affinity α-bungarotoxin binding site revealed by expression of subunit chimeras. *J Neurochem* 90: 479-489

Lee SC, Knowles TJ, Postis VLG, Jamshad M, Parslow RA, Lin YP, Goldman A, Sridhar P, Overduin M, Muench SP, *et al* (2016) A method for detergent-free isolation of membrane proteins in their local lipid environment. *Nat Protoc* 11: 1149–1162

Liu MY & Casida JE (1993) High affinity binding of [3h]imidacloprid in the insect acetylcholine receptor. *Pestic Biochem Physiol* 46: 40-46

Liu W, Xie Y, Ma J, Luo X, Nie P, Zuo Z, Lahrmann U, Zhao Q, Zheng Y, Zhao Y, *et al*

(2015) IBS: An illustrator for the presentation and visualization of biological sequences.
Bioinformatics 31: 3359-61

Loo RR, Dales N, Andrews PC (1996) The effect of detergents on proteins analyzed by
electrospray ionization. *Methods Mol Biol* 61:141-160

Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN,
Potter SC, Finn RD, *et al* (2019) The EMBL-EBI search and sequence analysis tools
APIs in 2019. *Nucleic Acids Res* 47: W636-W641

Maldonado-Hernández R, Quesada O, Colón-Sáez JO, Lasalde-Dominicci JA (2020)
Sequential purification and characterization of *Torpedo californica* nAChR-DC
supplemented with CHS for high-resolution crystallization studies. *Anal Biochem* 610:
113887

Martens C, Shekhar M, Borysik AJ, Lau AM, Reading E, Tajkhorshid E, Booth PJ, Politis A
(2018) Direct protein-lipid interactions shape the conformational landscape of secondary
transporters. *Nat Commun* 9: 4151

Mitchell AL, Attwood TK, Babbitt PC, Blum M, Bork P, Bridge A, Brown SD, Chang HY,
El-Gebali S, Fraser MI, *et al* (2019) InterPro in 2019: Improving coverage, classification
and access to protein sequence annotations. *Nucleic Acids Res* 47: D351-D360

Mulcahy MJ, Paulo JA, Hawrot E (2018) Proteomic Investigation of Murine Neuronal α7-
Nicotinic Acetylcholine Receptor Interacting Proteins. *J Proteome Res* 17: 3959-3975

Perry T, Chen W, Ghazali R, Yang YT, Christesen D, Martelli F, Lumb C, Luong HNB,
Mitchell J, Holien JK, *et al* (2021) Role of nicotinic acetylcholine receptor subunits in
the mode of action of neonicotinoid, sulfoximine and spinosyn insecticides in
Drosophila melanogaster. *Insect Biochem Mol Biol* 131: 103547

Perry T, Heckel DG, McKenzie JA, Batterham P (2008) Mutations in Dα1 or Dβ2 nicotinic
acetylcholine receptor subunits can confer resistance to neonicotinoids in *Drosophila*
melanogaster. *Insect Biochem Mol Biol* 38: 520-528

Perry T, Somers J, Yang YT, Batterham P (2015) Expression of insect α6-like nicotinic
acetylcholine receptors in *Drosophila melanogaster* highlights a high level of
conservation of the receptor: spinosyn interaction. *Insect Biochem Mol Biol* 64: 106-115

Pierleoni A, Martelli PL, Casadio R (2008) PredGPI: A GPI-anchor predictor. *BMC*
Bioinformatics 9: 392

Queiroz RML, Smith T, Villanueva E, Marti-Solano M, Monti M, Pizzinga M, Mirea D-M,
Ramakrishna M, Harvey RF, Dezi V, *et al* (2019) Comprehensive identification of
RNA-protein interactions in any organism using orthogonal organic phase separation
(OOPS). *Nat Biotechnol* 37: 169-178

Quesada O, González-Freire C, Ferrer MC, Colón-Sáez JO, Fernández-García E, Mercado J,
Dávila A, Morales R, Lasalde-Dominicci JA (2016) Uncovering the lipidic basis for the
preparation of functional nicotinic acetylcholine receptor detergent complexes for
structural studies. *Sci Rep* 6: 32766

Rahman MM, Teng J, Worrell BT, Noviello CM, Lee M, Karlin A, Stowell MHB, Hibbs RE
(2020) Structure of the Native Muscle-type Nicotinic Receptor and Inhibition by Snake
Venom Toxins. *Neuron* 106: 952-962.e5

Rappsilber J, Mann M, Ishihama Y (2007) Protocol for micro-purification, enrichment, pre-
fractionation and storage of peptides for proteomics using StageTips. *Nat Protoc* 2:
1896–1906

Rohde PD, Madsen LS, Neumann Arvidson SM, Loeschcke V, Demontis D, Kristensen TN
(2016) Testing candidate genes for attention-deficit/hyperactivity disorder in fruit flies
using a high throughput assay for complex behavior. *Fly (Austin)* 10: 25-34

Rstudio Team (2020) RStudio: Integrated Development for R. *Rstudio, PBC, Boston, MA*
URL http//www.rstudio.com/

Salgado VL (2021) Selective actions of insecticides on desensitizing and non-desensitizing
nicotinic acetylcholine receptors in cockroach (*Periplaneta americana*) neurons. *Pest*
Manag Sci 77: 3663-3672

Schmidt-Nielsen BK, Gepner JI, Teng NNH, Hall LM (1977) Characterization of an α-
bungarotoxin binding component from *Drosophila melanogaster*. *J Neurochem* 29:
1013-1029

Shevchenko A, Tomas H, Havliš J, Olsen JV, Mann M (2007) In-gel digestion for mass
spectrometric characterization of proteins and proteomes. *Nat Protoc* 1: 2856–2860

Sobotzki N, Schafroth MA, Rudnicka A, Koetemann A, Marty F, Goetze S, Yamauchi Y,
Carreira EM, Wollscheid B (2018) HATRIC-based identification of receptors for orphan
ligands. *Nat Commun* 9: 1519

Somers J, Luong HNB, Mitchell J, Batterham P, Perry T (2017) Pleiotropic effects of loss of
the Dα1 subunit in *Drosophila melanogaster*: Implications for insecticide resistance.
Genetics 205: 263-271

Sormanni P, Amery L, Ekizoglou S, Vendruscolo M, Popovic B (2017) Rapid and accurate in
silico solubility screening of a monoclonal antibody library. *Sci Rep* 7: 8200

Sormanni P, Aprile FA, Vendruscolo M (2015) The CamSol method of rational design of
protein mutants with enhanced solubility. *J Mol Biol* 427: 478-490

Su H & O'Dowd DK (2003) Fast synaptic currents in *Drosophila* mushroom body kenyon
cells are mediated by α-bungarotoxin-sensitive nicotinic acetylcholine receptors and
picrotoxin-sensitive GABA receptors. *J Neurosci* 23: 9246–9253

Tackenberg MC, Giannoni-Guzmán MA, Sanchez-Perez E, Doll CA, Agosto-Rivera JL,
Broadie K, Moore D, McMahon DG (2020) Neonicotinoids disrupt circadian rhythms
and sleep in honey bees. *Sci Rep* 10: 17929

Tsirigos KD, Elofsson A, Bagos PG (2016) PRED-TMBB2: Improved topology prediction
and detection of beta-barrel outer membrane proteins. *Bioinformatics* 32: i665-i671

Wang H, Yu M, Ochani M, Amella CA, Tanovic M, Susarla S, Li JH, Wang H, Yang H,
Ulloa L, *et al* (2003) Nicotinic acetylcholine receptor α7 subunit is an essential regulator
of inflammation. *Nature* 421: 384-388

Wu P, Ma D, Pierzchala M, Wu J, Yang LC, Mai X, Chang X, Schmidt-Glenewinkel T
(2005) The Drosophila acetylcholine receptor subunit Dα5 is part of an α-bungarotoxin
binding acetylcholine receptor. *J Biol Chem* 280: 20987-20994

Xu X, Bao H, Shao X, Zhang Y, Yao X, Liu Z, Li Z (2010) Pharmacological characterization

of cis-nitromethylene neonicotinoids in relation to imidacloprid binding sites in the
brown planthopper, *Nilaparvata lugens*. *Insect Mol Biol* 19: 1-8

Zuo YY, Xue YX, Wang ZY, Ren X, Aioub AAA, Wu YD, Yang YH, Hu ZN (2021)
Knockin of the G275E mutation of the nicotinic acetylcholine receptor (nAChR) α6
confers high levels of resistance to spinosyns in *Spodoptera exigua*. *Insect Sci* 0: 1-9

Figure Legends

Figure 1. Morphological and locomotor phenotypes in *nAChR* subunit mutants.

A   Adult males from indicated *nAChR* subunit null mutants. Arrows indicate strong curled
abdomen phenotypes.
B   Graph of locomotor activity determined in climbing assays as a percentage of wild-
type. Error bars represent standard deviation from 5 replicates.

Figure 2. ω-Hexatoxin-Hv1a and α-Bungarotoxin target different nAChR subunits.

A   Bar graph of the survival rate, measured as the percentage of pupae formed, following
larval injection of 2.5 nmol/g Hv1a in the indicated homozygous lines. Mean ± SD of
3 independent replicates of 10 larvae per replicate. **P=0.0035 (one-way ANOVA
($F_{(11,24)}$=4.99, P=0.0005 with Bonferroni's multiple comparisons test). 3 independent
replicates in each group (10 injected larvae in total).
B   Survival rate following larval injection of 1.25 nmol/g α-Btx. Mean ± SD of 3
independent replicates of 10 larvae per replicate. **P<0.001, ***P=0.0001 (one-way
ANOVA ($F_{(11,24)}$= 7.921, P<0.0001, followed by Bonferroni's multiple comparisons
test). 3 independent replicates in each group (10 injected larvae in total). w^{1118} is the
wild-type base stock, *THattp40* and *THattP2* are the Cas9 lines used to establish the
mutants, w^{1118} + PBS represents the injection control.

Figure 3. Forming styrene maleic acid lipid particles (SMALPs).

A   Schematic representation of the SMALPs extraction and nAChRs pull-down for mass
spectrometric analysis.
B   Negative staining of extracted SMALPs by transmission electron microscopy. Scale bar
100 nm.
C   Fluorescence signal of uncoupled α-Btx in solution before and after coupling to affinity
beads (two-tailed t-test, ***P<0.001, n=4).
D, E Negative staining of extracted SMALPs after α-Btx coupled pull-downs. Ring-like
protein structures are boxed (Scale bar = 100 nm) with an example in the magnified
image (Scale bar = 20 nm). A top view of the nAChR structure from PDB entry 4HQP
is shown for reference.

Figure 4. Identification of proteins enriched by SMALP extraction.

A	Number of identified proteins in affinity pull-down samples solubilized with or without SMA, two-tailed t-test, **$P<0.01$, n=6 or 8 replicates per condition.
B	MS/MS spectrum counts from samples solubilized with or without SMA, ns = not significant after two-tailed t-test with n=6 or 8.
C	Calculated hydrophobicity score of amino acid residues found in protein sequences obtained with and without SMA solubilisation, ****$P<0.0001$, two-tailed t-test, n=3 per condition.
D	GO term (cellular compartment) enrichment of proteins identified with and without SMA solubilisation, n=4 or 11.
E	Predicted numbers of proteins containing transmembrane helices obtained with or without SMA solubilisation, n=4 or 8.
F, G	Analysis of solubility and hydrophobicity of receptors identified with and without SMA solubilisation ($r^2= -0.56$, $P<0.0001$, n=4) and of transmembrane receptor helices ($r^2=0.56$, $P<0.01$, n=4).
H	Solubility score of individual nAChR subunits.

Figure 5. Three nAChR α-subunits are binding to α-Bungarotoxin (α-Btx).

A	Graphical representation of ten nAChR subunits. The position of protein domains and signal peptides are shown.
B	Identified peptides of Dα5, Dα6 and Dα7 nAChR subunits in pull-downs using α-Btx affinity beads. Found peptides in ligand-binding and cytoplasmic domain are highlighted in red.
C	Numbers of identified unique peptides in wild-type pull-downs using affinity beads in absence and presence of α-Btx, n=3. Deleting *nAChRα5, nAChRα6, nAChRα7* and performing pull-downs identified unique peptides of nAChR subunits suggesting that functional complexes can be formed in null alleles, n=3.
D	KEGG pathway enrichment analysis of pull-downs in wild-type and *nAChRα5, nAChRα6, nAChRα7* null alleles, Fisher's exact test, n=3. Protein counts with *P* values of enriched pathways are shown. *P* values of ≤ 0.05 are to be considered as strongly enriched with default threshold of 0.1.

Figure 6. N-glycosylation sites in nAChR subunits.

A	Diagrammatic representation of nAChR subunit glycopeptide enrichment. Pull-downs with α-Btx affinity beads enrich for nAChRs and after proteolytic digestion glycopeptides were enriched. Glycopeptides were deglycosylated with Endo H or PNGase F and analyzed by mass spectrometry.
B	Low numbers of glycopeptides (average 20) are detected in flow through fractions.
C	Numbers of identified glycopeptides according to site probabilities are shown (n=3).
D	Shared glycopeptide identified in the ligand-binding domain of Dα5 and Dα7, an N-linked glycosylated asparagine (N) residue is highlighted.

986	E	Deglycosylated peptide with either Endo H or PNGase F and contains either an N-acetylhexosamine or is deamidated on asparagine (N2). The two different modifications on the same peptide lead to a different monoisotopic mass (MH+ [Da]). Peptide contains an additional carbamidomethyl on cysteine (C5).
987		
988		
989		
990		

Figure 7. *In vivo* imaging of endogenously tagged Dα6 nAChR subunit.

992	A-G	Live imaging of fly brains carrying a C-terminal EGFP fusion into the endogenous *nAChRα6* locus.
994	A-C	Dα6 subunit in 2nd, early and late 3rd instar larvae brain, respectively. Visible localization in ventral nerve cord (VNC), mushroom bodies (MB), and optic lobes (OL). Scale bar = 100 µm.
997	D	Dα6 subunit in mushroom bodies of 3rd instar larvae with detectable fluorescence signal in Kenyon cells (KC), calyx (CX), peduncle (PED), dorsal lobes (DL) and medial lobes (ML). Scale bar = 100 µm.
1000	E	Dα6 subunit was observed in developing optic lobes, lamina (LAM) and medulla (MED) of later 3rd instar larvae. Scale bar = 100 µm.
1002	F	Dα6 subunit on the external structures of developing lobes in later 3rd instar larvae. Scale bar = 100 µm.
1004	G	Dα6 subunit in adult fly brain, strong signal detected in mushroom bodies (MB) and optic lobe (OL). Scale bar = 100 µm.
1006	H	Schematic summary of Dα6 subunit expression during different developmental stages, 2nd and 3rd instar larvae and adult fly, (L2, L3 and Adult, respectively) in which the green lines indicate the localization of the Dα6 subunit.

1025   **Appendix Tables and Figures legends**

1026

1027   **Table of Contents**

1028

1040

1041

1042

1043   **Appendix Table S1 Climbing ability.**

10 days old flies									
number	receptor	series 1		series 2		series 3		average	standard deviation
	subunit	percentage	actual change	percentage	actual change	percentage	actual change	percent	STEV
1	nAChR α1	30	37.5	40	50	50	71.4	53	17.2
2	nAChR α2	40	50	40	50	50	71.4	57.1	12.4
3	nAChR α3	0	0	20	25	30	42.9	22.6	21.5
4	nAChR α4	70	87.5	80	100	60	85.7	91.1	7.8
5	nAChR α5	60	75	70	87.5	50	71.4	78	8.4
6	nAChR α6	40	50	50	62.5	50	71.4	61.3	10.8
7	nAChR α7	70	87.5	80	100	70	100	95.8	7.2
8	nAChR β1	30	37.5	30	37.5	20	28.6	34.5	5.2
9	nAChR β2	80	100	80	100	60	85.7	95.2	8.2
10	nAChR β3	80	100	80	100	70	100	100	0
11	WT	80	100	80	100	70	100	100	0

1044

1045

1046

1047

1048

Appendix Table S2 *Drosophila* **larval injection of ω-Hexatoxin-Hv1a & α-Bungarotoxin.**

	Strain	Injected cpd	Survival (% of pupae formed after injection)				
			Rep1	Rep2	Rep3	Average	Standard Deviation
CONTROL	w[1118]	2.5 nmol/g Hv1a	0	0	0	0	0
CONTROL - Cas9 lines	THattP40	2.5 nmol/g Hv1a	0	0	0	0	0
CONTROL - Cas9 lines	THattP2	2.5 nmol/g Hv1a	0	0	0	0	0
nAChR CRISPR mutant	nAChR α1	2.5 nmol/g Hv1a	25	0	0	8.33	14.43
nAChR CRISPR mutant	nAChR α2	2.5 nmol/g Hv1a	25	0	0	8.33	14.43
nAChR CRISPR mutant	nAChR α3	2.5 nmol/g Hv1a	0	0	0	0	0
nAChR CRISPR mutant	nAChR α4	2.5 nmol/g Hv1a	25	33.33	66.67	41.67	22.05
nAChR CRISPR mutant	nAChR α5	2.5 nmol/g Hv1a	0	0	0	0	0
nAChR CRISPR mutant	nAChR α6	2.5 nmol/g Hv1a	0	0	0	0	0
nAChR CRISPR mutant	nAChR α7	2.5 nmol/g Hv1a	0	0	0	0	0
nAChR CRISPR mutant	nAChR β2	2.5 nmol/g Hv1a	25	33.33	66.67	41.67	22.05
nAChR CRISPR mutant	nAChR β3	2.5 nmol/g Hv1a	0	33.33	0	11.11	19.25
Injection CONTROL	w[1118]	PBS	100	100	100	100	0
CONTROL	w[1118]	1.25 nmol/g α-Btx	0	0	0	0	0
CONTROL - Cas9 lines	THattP40	1.25 nmol/g α-Btx	0	0	0	0	0
CONTROL - Cas9 lines	THattP2	1.25 nmol/g α-Btx	0	0	0	0	0
nAChR CRISPR mutant	nAChR α1	1.25 nmol/g α-Btx	0	0	33.33	11.11	19.25
nAChR CRISPR mutant	nAChR α2	1.25 nmol/g α-Btx	25	0	33.33	19.44	17.35
nAChR CRISPR mutant	nAChR α3	1.25 nmol/g α-Btx	0	0	33.33	11.11	19.25
nAChR CRISPR mutant	nAChR α4	1.25 nmol/g α-Btx	0	33.33	0	11.11	19.25
nAChR CRISPR mutant	nAChR α5	1.25 nmol/g α-Btx	50	66.67	66.67	61.11	9.62
nAChR CRISPR mutant	nAChR α6	1.25 nmol/g α-Btx	25	66.67	66.67	52.78	24.06
nAChR CRISPR mutant	nAChR α7	1.25 nmol/g α-Btx	50	100	66.67	72.22	25.46
nAChR CRISPR mutant	nAChR β2	1.25 nmol/g α-Btx	0	33.33	0	11.11	19.25
nAChR CRISPR mutant	nAChR β3	1.25 nmol/g α-Btx	0	0	0	0	0
Injection CONTROL	w[1118]	PBS	100	100	100	100	0

 Appendix Table S3 Identified nAChR peptides in pull-downs with α-Bungarotoxin.

Peptides from Dα3, Dα5, Dα6, Dα7 and Dβ3 nAChR subunits are listed and found [N] times within
individual replicates. Protein domains are marked with: Ed extracellular-, Id Intracellular-, LBD ligand-
binding-, and Non-domain localization. The mass-to-charge ratio (m/z) of the precursor ions, the
protonated monoisotopic masses, the theoretical MH$^+$ masses in Dalton [Da] and peptide modifications
are listed. Peptide modifications are listed with: (C) Carbamidomethylation; (N,Q) Deamidation; (H)
N-acetylhexosamine (HexNAc); (M) Oxidation.

Appendix Table S4 Identified nAChR peptides in pull-downs without α-Bungarotoxin.

Identified peptides of nAChR subunits which are found in control pull-down samples without α-
Bungarotoxin (α-Btx).

1061

Appendix Table S3 Identified nAChR peptides in pull-downs with α-Bungarotoxin.

Subunit	Accession	Sequence	Found [N]	Domain	m/z [Da]	MH+ [Da]	Theo. MH+ [Da]	Modification
nAChRα3	Q9W3G6	ATLnYTGR	2	LBD	549.77423	1098.54119	1098.54258	H4 or position 2 , 6
nAChRα5	Q7KT97	TVYGQGDDGSIGPIGSTR	7	Id	890.42413	1779.84099	1779.85078	
nAChRα5	Q7KT97	TVYGqGDDGSIGPIGSTR	2	Id	890.93036	1780.85344	1780.83480	Q5
nAChRα5	Q7KT97	FITDQLR	7	Id	446.74768	892.48808	892.48869	
nAChRα5	Q7KT97	KHQILSDVELKER	1	Id	399.47717	1594.8869	1594.8911	
nAChRα5	Q7KT97	SSTEYELGLILK	1	Id	676.86945	1352.7316	1352.7308	
nAChRα5, α7	Q7KTF97, Q9VWI9	LEWNDMNLR	1	LBD	595.78156	1190.55583	1190.5623	
nAChRα5, α7	Q7KTF97, Q9VWI9	NnGScLYVPPGIFK	2	LBD	884.93384	1768.86040	1768.85745	H2, C5
nAChRα5, α7	Q7KTF97, Q9VWI9	NnGScLYVPPGIFK	2	LBD	783.88092	1566.75456	1566.76209	N2, C5
nAChRα5, α7	Q7KTF97, Q9VWI9	nnGScLYVPPGIFK	2	LBD	784.39069	1567.77410	1567.74611	N1, N2, C5
nAChRα6	Q7KTF9	ELQFITAR	4	Id	489.27380	977.54033	977.54146	
nAChRα6	Q7KTF9	ELqFITAR	1	Id	489.7753	978.54332	978.52547	Q3
nAChRα6	Q7KTF9	TADIHEMPPWIK	1	Non	719.36407	1437.72087	1437.71950	
nAChRα6	Q7KTF9	TILLSNR	4	Id	408.75095	816.49462	816.49378	
nAChRα6	Q7KTF9	ADDEAELIGDWK	4	Id	681.3179	1361.61.61785	1361.62195	
nAChRα6	Q7KTF9	KADDEAELIGDWK	1	Id	745.36469	1489.72209	1489.71691	
nAChRα6	Q7KTF9	KTILLSNR	2	Id	472.79971	944.59215	944.58874	
nAChRα5, α6, α7	Q7KT97, Q7KTF9, Q9VWI9	IDITWFPFDDQR	3	LBD	776.87292	1552.7386	1552.7431	
nAChRα5, α6, α7	Q7KT97, Q7KTF9, Q9VWI9	SLLANVLDIDDDFR	3	Id	803.40924	1605.8112	1605.8119	
nAChRα6, α7	Q7KTF9, Q9VWI9	IDITWFPFDDqR	1	LBD	777.37311	1553.73894	1553.7209	Q11
nAChRα7	Q9VWI9	SLLANVLDIDDDFRcNHR	1	Id	544.01544	2173.0399	2173.0455	C15
nAChRα7	Q9VWI9	KQQIQNVELKER	6	Id	504.95410	1512.84774	1512.84927	
nAChRα7	Q9VWI9	KQQIQNVELK	1	Id	614.35730	1227.70732	1227.70556	
nAChRα7	Q9VWI9	KQqIQNVELKER	1	Id	502.29062	1513.8573	1513.8333	N3
nAChRα7	Q9VWI9	QGDDGSVGPVGPAGPVVDGR	12	Id	918.44794	1835.88860	18335.88823	
nAChRα7	Q9VWI9	qGDDGSVGPVGPAGPVVDGR	1	Id	918.94897	1936.89067	1836.87225	Q1
nAChRα7	Q9VWI9	QQIQNVELK	1	Id	550.30487	1099.60246	1099.61060	
nAChRα7	Q9VWI9	EDETSDITR	1	Non	533.23535	1065.46343	1065.46947	
nAChRα7	Q9VWI9	mQRPGQVGYEcPPPPSSSSSSASGEK	1	Id	908.41028	2723.2163	2723.2036	M1; C11
nAChRα7	Q9VWI9	cASATLPHQPTYYR	2	Id	555.59827	1664.7803	1664.785	C1
nAChRα7	Q9VWI9	WITEQLKKEDETSDITR	2	Id	698.01874	2092.0417	2092.0557	
nAChRα7	Q9VWI9	WITEQLK	1	Non	459.25516	917.50304	917.50909	
nAChRα7	Q9VWI9	cNHRCASATLPHQPTYYR	1	Id	832.89795	1664.7886	1664.785	C1
nAChRβ3	Q9VPQ8	VVLPEnGTAR	1	LBD	629.83380	1258.66033	1258.66376	H6 or position 8

Appendix Table S4 Identified nAChR peptides in pull-downs without α-Bungarotoxin.

Subunit	Accession	Sequence	Found [N]	Domains	m/z [Da]	MH+ [Da]	Theo. MH+ [Da]	Modification
nAChRα1	A0A0B4KGU3	LFIQILPK	1	Ed	486.31339	971.61949	917.62882	
nAChRα1, α2	A0A0B4KGU3, P17644	LYDDLLSNYNR	1	Ed	693.3407	1385.674	1385.66957	
nAChRα2	P17644	AIDVQLSDVAK	2	Non	579.823	1158.639	1158.636	
nAChRα2	P17644	VVWTPPAIFK	1	Ed	579.3391	1157.671	1157.671	
nAChRα2, β2	P17644, P25162	LSQLIEVNLK	1	Ed	386.5676	1157.692	1157.677	
nAChRα4	A8JNX5	LVSSGYNNSLPK	10	Non	639.8441	1278.661	1278.66884	
nAChRα4	A8JNX5	LVSSGYnNSLPK	1	Non	640.3381	1279.669	1279.653	N7
nAChRα4	A8JNX5	LSQLIDVNLK	2	LBD	571.8425	1142.678	1142.678	
nAChRα4	A8JNX5	SPILNNPAFSHSK	1	Non	471.24591	1411.72318	1411.73284	
nAChRα4	A8JNX5	RPTYNFETSK	3	Non	621.80939	1242.61150	1242.61133	
nAChRα4	A8JNX5	RPTYnFETSK	3	Non	622.3102	1243.613	1243.595	N5
nAChRα4	A8JNX5	LYDDLLSNYNK	2	LBD	679.3725	1357.64910	1357.66342	
nAChRα4	A8JNX5	RPTYNFETSKLLLK	1	Non	621.8079	1242.609	1242.611	
nAChRβ1	P04755	NKNFVDLSDYWK	2	Ed	643.8068	1286.600	1286.605	
nAChRβ1	P04755	NFVDLSDYWK	4	Ed	643.8044	1286.602	1286.605	
nAChRβ1	P04755	nFVDLSDYWK	4	Ed	644.3093	1287.611	1287.589	N1
nAChRβ1	P04755	ILPPTSLVLPLIAK	1	Non	737.9825	1474.96074	1474.96071	
nAChRβ1	P04755	ATEAVEFIAEHLR	5	Id	495.92743	1485.76773	1485.76962	
nAChRβ1	P04755	VWKPDIVLFNNADGNYEVR	1	Ed	750.3779	2249.119	2249.135	
nAChRβ2	P25162	LYDDLLSNYNR	3	Ed	693.34070	1385.67412	1385.66957	
nAChRβ2	P25162	LSqLIEVNLKNqVMTTNLWVK	1	Ed	825.1133	2473.325	2473.3370	Q3; Q12
nAChRβ2	P25162	LSqLIEVNLK	2	Ed	386.5676	1157.688	1157.678	Q3

Appendix Table S5 List of gRNAs and oligonucleotides used for cloning.

nAChR subunit	gRNA target name	gRNA sequence (NGG)	oligonucleotides name (Forward/Reverse)	oligonucleotides sequence
Knockouts				
nAChRα1	Da1_111(+)	5'CGGAGATGTAGTAGTCCTGCAGG3'	41_Da1_111_F	5'GTCGCGGAGATGTAGTAGTCCTGC3'
			42_Da1_111_R	5'AAACGCAGGACTACTACATCTCCG3'
	Da1_126(-)	5'CCTGCAGGTCGATGCCCACCTCG3'	43_Da1_126_F	5'GTCGCGAGGTGGGCATCGACCTGC3'
			44_Da1_126_R	5'AAACGCAGGTCGATGCCCACCTCG3'
nAChRα2	Da2_99 (+)	5'GCTCCTCTGCGAAACCGTTCAGG3'	45_Da2_99_F	5'GTCGCTCCTCTGCGAAACCGTTC3'
			46_Da2_99_R	5'AAACGAACGGTTTCGCAGAGGAG3'
nAChRα3	Da3_18(+)	5'GTCCGGACGCCAGATGTGATCGG3'	49_Da3_18_F	5'GTCGTCCGGACGCCAGATGTGAT3'
			50_Da3_18_R	5'AAACATCACATCTGGCGTCCGGA3'
nAChRα4	Da4_19(+)	5'TTGTTGCGACGAACCATACTTGG3'	53_Da4_19_F	5'GTCGTTGTTGCGACGAACCATACT3'
			54_Da4_19_R	5'AAACAGTATGGTTCGTCGCAACAA3'
nAChRα5	Da5_232(-)	5'CCGGGGATCTTCAAGTCGACGTG3'	57_Da5_232_F	5'GTCGCACGTCGACTTGAAGATCCC3'
			58_Da5_232_R	5'AAACGGGATCTTCAAGTCGACGTG3'
	Da5_251(+)	5'CGTGCAAGATCGACATCACGTGG3'	59_Da5_251_F	5'GTCGCGTGCAAGATCGACATCACG3'
			60_Da5_251_R	5'AAACCGTGATGTCGATCTTGCACG3'
nAChRα6	Da6_70(+)	5'CGTATTCTTCTTTCCCGGCATGG3'	61_Da6_70_F	5'GTCGCGTATTCTTCTTTCCCGGCA3'
			62_Da6_70_R	5'AAACTGCCGGGAAAGAAGAATACG3'
nAChRα7	Da7_1226(+)	5'CATTGACCACCGGACGCTCCAGG3'	63_Da7_1226_F	5'GTCGCATTGACCACCGGACGCTCC3'
			64_Da7_1226_R	5'AAACGGAGCGTCCGGTGGTCAATG3'
nAChRβ1	Db1_2(+)	5'TGGAGTCTTCCTGCAAATCCTGG3'	67_Db1_2_F	5'GTCGTGGAGTCTTCCTGCAAATCC3'
			68_Db1_2_R	5'AAACGGATTTGCAGGAAGACTCCA3'
nAChRβ2	Db2_955(+)	5'TCAGACCTAACCAAACCGTCAGG3'	71_Db2_955_F	5'GTCGTCAGACCTAACCAAACCGTC3'
			72_Db2_955_R	5'AAACGACGGTTTGGTTAGGTCTGA3'
nAChRβ3	Db3_466(+)	5'CTTTGAAGTCCAGCGAGGTCTGG3'	75_Db3_466_F	5'GTCGCTTTGAAGTCCAGCGAGGTC3'
			76_Db3_466_R	5'AAACGACCTCGCTGGACTTCAAAG3'
C-terminal tagging				
nAChRα6	Da6_181(+)	5'TTGCACGATTATGTGCGGAGCGG3'	131_Da6_181_F	5'GTCGTTGCACGATTATGTGCGGAG3'
			132_Da6_181_R	5'AAACCTCCGCACATAATCGTGCAA3'
	Da6_176(+)	5'CCTTATTGCACGATTATGTGCGG3'	133_Da6_176_F	5'GTCGCCTTATTGCACGATTATGTG3'
			134_Da6_176_R	5'AAACCACATAATCGTGCAATAAGG3'

Appendix Table S6 List of oligonucleotides used for amplification from genomic DNA.

nAChR subunit	homology arm	oligonucleotides name (Forward/Reverse)	oligonucleotides sequence
		knockouts	
nAChRα1	Da1_LHA	Da1_LHA_F1	5'TGGGGCGACAAAATAGCATG3'
		Da1_LHA_R1	5'GGGGAAATGGGCCAACAAAT3'
nAChRα1	Da1_RHA	Da1_RHA_F1	5'GCAGATACTTTCCCAGCAGC3'
		Da1_RHA_R1	5'CCGCGTCCTTGACTACTTTG3'
nAChRα2	Da2_LHA	Da2_LHA_F1	5'ACGAAATGCAAAACCGAGCT3'
		Da2_LHA_R2	5'CCCAATTTGACCAACACCGT3'
nAChRα2	Da2_RHA	Da2_RHA_F1	5'GCGGGCAGAAAGGTAAACAA3'
		Da2_RHA_R1	5'TCACCTGATCACCGTCGTAG3'
nAChRα3	Da3_LHA	Da3_LHA_F1	5'CTCCAGCCGTTCCCAAATCT3'
		Da3_LHA_R1	5'CAATCTGTGGGTGGAGCAGT3'
nAChRα3	Da3_RHA	Da3_RHA_F1	5'CTGCTCGTCGAAGGGAAAGT3'
		Da3_RHA_R1	5'GATCCGAGCCAGACTAAGCC3'
nAChRα4	Da4_LHA	Da4_LHA_F1	5'GATGAACAACAGGGCAGCAA3'
		Da4_LHA_R1	5'CAAAACAACAACCGTCACGC3'
nAChRα4	Da4_RHA	Da4_RHA_F1	5'TTAGAGCGTAACAGTGGGCG3'
		Da4_RHA_R1	5'ACGCCTACAAACCGGACAAA3'
nAChRα5	Da5_LHA	Da5_LHA_F1	5'ACCGCATTCCTGTCGCATAT3'
		Da5_LHA_R1	5'CAGGACGACGTTGGCTTACT3'
nAChRα5	Da5_RHA	Da5_RHA_F1	5'GGATCTTCAAGTCGACGTGC3'
		Da5_RHA_R1	5'GAGGGTGTGGCTGGATTTTC3'
nAChRα6	Da6_LHA	Da6_LHA_F1	5'GTGTACGGGTGTGAGACAGA3'
		Da6_LHA_R1	5'TCACACATTGCTTGCCGAAA3'
nAChRα6	Da6_RHA	Da6_RHA_F1	5'GTCAGTTTCTCGCCCGAATC3'
		Da6_RHA_R1	5'CCGAGAGTTGACTGTAGCCA3'
nAChRα7	Da7_LHA	Da7_LHA_F1	5'TGTAAACCCTAGCAGTGCCA3'
		Da7_LHA_R1	5'TATGATACCGGGTGAGTGCC3'
nAChRα7	Da7_RHA	Da7_RHA_F1	5'CATCCGGTTTCCATAGGCGA3'
		Da7_RHA_R1	5'ACGGAAATCACAATGCCCCT3'
nAChRβ1	Db1_LHA	Db1_LHA_F1	5'TCATCAACAGCAGGCAGAGA3'
		Db1_LHA_R1	5'TGGCAATGAGAGCTTGGAGA3'
nAChRβ1	Db1_RHA	Db1_RHA_F1	5'CTGCAAATCCTGGCTGTTGT3'
		Db1_RHA_R1	5'GTGTGTGTGTGTGTGGTCTC3'
nAChRβ2	Db2_LHA	D2b_LHA_F1	5'TCAACTCAGGACAGCACACA3'
		D2b_LHA_R1	5'ACCACCACTTTCCTAGCTCC3'
nAChRβ2	Db2_RHA	D2b_RHA_F1	5'CCCATCGCAACTTGTAGTCG3'
		D2b_RHA_R1	5'CATTCGTCCAGGTAAGTGCG3'
nAChRβ3	Db3_LHA	Db3_LHA_F1	5'AACGGTTCCGATGACTTCCT3'
		Db3_LHA_R1	5'TGAGCATGTTGAGTTCGCAG3'
nAChRβ3	Db3_RHA	Db3_RHA_F1	5'TCCTTCGTCCTCTCCTTCGT3'
		Db3_RHA_R1	5'TTCTGCGGGAAACTACGACC3'

Appendix Table S6 Continued, List of oligonucleotides used for amplification from genomic DNA.

C-terminal tagging			
nAChR subunit	homology arm	oligonucleotides name (Forward/Reverse)	oligonucleotides sequence
nAChRα6	Da6_RHA	74_Da6_RHA_F_Gen	5'GGGTTTCTGTTCTTGCGCTG3'
		75_Da6_RHA_R_Gen	5'GCCCTGCTGATTTGTTTGCT3'
nAChRα6	Da6_LHA	76_Da6_LHA_F_Gen	5'CCGATGCTTCCGACGTATCC3'
		77_Da6_LHA_R_Gen	5'GCCATACTAGCGCATGACTCT3'

Appendix Table S7 C-terminal tagging of nAChRa6 with FSVS.

C-terminal tagging with FSVS			
nAChR subunit	fragment	site	oligonucleotides sequence
nAChRα6		105__Donor_LHA_Da6	5'CGGGCTAATTATGGGGTGTCGCCCTGTGCATGCAGAGAATGAAACC3'
	31	106_LHA_Da6_linker_R	5'CCTTGCACGATTATGTGCGGAGCTGAGAGCAGCACCGTAACCG3'
		107_linkerTag_F	5'GCTCTCAGCTCCGCACATAATCGTGCAAGGATCCGGCGGAGGGGGC3'
	32	108_RHDa6_Tag_R	5'CTAATTCGAGCGTCCTTACTTTTCGAACTGGGGATGGC3'
		109_TG-RHA_Da6_F	5'CCCCAGTTCGAAAAGTAAGGACGCTCGAATTAGGCC3'
	33	110_Donor_RHDa6	5'AAATTTTGTGTCGCCCTTGAACTCGATTTGCGCTGCTTAGCTTCATCTG3'
C-terminal tagging with FSVS-loxP-3Px3DsRED-loxP using as a template donors with FSVS tags above (fragments 31-33)			
nAChR subunit	fragment	site	oligonucleotides sequence
nAChRα6		105__Donor_LHA_Da6	5'CGGGCTAATTATGGGGTGTCGCCCTGTGCATGCAGAGAATGAAACC3'
	34	114_RHDa6_Tag_R	5'CTTTTCGAACTGGGGATGGCTCCAAGCTCC3'
		155_Marker_F1	5'CTTGGAGCCATCCCCAGTTCGAAAAGTAGTAAGGTACCGCGGGTATAAC3'
	36	157_Marker_R1	5'GGCTCTTCTATATAACTTCGTATAGCATAC3'
		165_Da6_F_marker	5'ATGCTATACGAAGTTATATAGAAGAGCCTAAGGACGCTCGAATTAGGCC3'
	36	110_Donor_RHDa6	5'AAATTTTGTGTCGCCCTTGAACTCGATTTGCGCTGCTTAGCTTCATCTG 3'

Appendix Figure S1 GO terms and predicted membrane proteins.

A, B   GO slim term for biological process and for molecular function analysed within samples solubilized without or with SMA, n=4 or 11 per conditions.

C   Predicted β-barrel membrane proteins, two-tailed t-test ****$P<0.0001$, n=6 or 10.

D   Palmitoylated proteins, two-tailed t-test ****$P<0.0001$, each n=8.

E   Myristoylated proteins, two-tailed t-test ****$P<0.0001$, n=6 or 10.

F   GPI-anchored proteins, two-tailed t-test, non-significant ns, n=6 or 10.

Appendix Figure S2 Identified peptides in ligand-binding and cytoplasmic domain.

A   Shared peptides found in the ligand-binding domains are shown in red.

B   Identified unique (/) and shared (,) peptides in cytoplasmic domains.

Appendix Figure S3 Superimposed nAChR α-subunits structure together with identified peptides.

A   Superimposed nAChR α-subunit structures from *H. sapiens* (blue, PDB 6USF) and *Torpedo californica* (red, 6UWZ). Extracellular ligand-binding domain (LBD) illustrates a structure similarity.

B   Same superimposed structures bound to α-Bungarotoxin (α-Btx, surface structure). Peptides found in LBD are highlighted in green. The homology regions of Dα6 nAChRs LBD are shown in violet.

Appendix Figure S4 Glycosylation sites of nAChR subunits.

A   Multiple sequence alignment of insect α7 nAChR subunits compared to sequences of nematodes. The glycosylated ligand-binding domain (LBD) sequence of Dα5 and Dα7 nAChR subunits are shown. Glycosylated asparagine residues highlighted in red are conserved within insects and nematodes (positions: Dα5 422 and Dα7 170 amino acids).

B   Same Dα5 and Dα7 nAChR subunit sequences compared to *T. californica*, *D. rerio*, *M. musculus* and *H. sapiens*.

C   Graphical representation of Dα3 and Dβ3 nAChR subunits. N-acetylhexosamine (H) modification on asparagine residues are highlighted and are of low site probability $\leq 80 \%$.

Figures

Table of Contents

Figure 1. Morphological and locomotor phenotypes in *nAChR* subunit mutants.

Figure 2. ω-Hexatoxin-Hv1a and α-Bungarotoxin target different nAChR subunits.

Figure 3. Forming styrene maleic acid lipid particles (SMALPs).

Figure 4. Identification of proteins enriched by SMALP extraction.

Figure 5. Three nAChR α-subunits are binding to α-bungarotoxin (α-Btx).

Figure 6. N-glycosylation sites in nAChR subunits.

Figure 7. *In vivo* **imaging of endogenously tagged Dα6 nAChR subunit.**

Appendix Figure S1 GO terms and predicted membrane proteins.

Appendix Figure S2 Identified peptides in ligand-binding and cytoplasmic domain.

Appendix Figure S3 Superimposed nAChR α-subunits structure together with identified peptides.

Appendix Figure S4 Glycosylation sites of nAChR subunits.