Maternal control over caste allocation in the ant *Cardiocondyla obscurior*

Eva Schultner[1*]

Tobias Wallner[1]

Benjamin Dofka[1]

Jeanne Bruelhart[1]

Jürgen Heinze[1]

Dalial Freitak[2]

Tamara Pokorny[1]

Jan Oettler[1*]

[1]Zoology/Evolutionary Biology, University of Regensburg, Universitätsstrasse 31,

93053 Regensburg, Germany

[2] Institute for Biology, University of Graz, Universitätsplatz 2, 8010 Graz, Austria

*corresponding authors

ORCIDs:

Eva Schultner: 0000-0002-5069-9732

Tobias Wallner: 0000-0001-9135-6456

Tamara Pokorny: 0000-0002-8183-9638

Dalial Freitak: 0000-0001-8574-0531

Jan Oettler: 0000-0002-8539-6029

Keywords:

Social insects, caste determination, development, eggs, larvae, discrimination

behavior

Author contributions:

ES and JO supervised the study and wrote the manuscript. ES analyzed and visualized

the data. TW discovered the crystalline deposits and together with BD contributed to

the rearing and behavioral experiments. JB contributed morphological data, DF

contributed protein data, TP contributed cuticular hydrocarbon data. All authors read

and approved the final version of the manuscript.

Abstract

Ant queens and workers commonly engage in conflict over reproductive allocation

because they have different fitness optima. How such conflicts are resolved depends

on the power that each party of interest holds. Here, we show that queens have

control over caste allocation in the ant *Cardiocondyla obscurior*. With the discovery

of crystalline deposits that allow the identification of queens and workers across the

entire course of development, we demonstrate that caste fate is irreversibly

determined in the egg stage. Comparison of queen and worker-destined eggs and

larvae revealed size and weight differences in late development, but no discernible

differences in traits that may be used in social interactions, including hair morphology

and cuticular odors. In line with this lack of caste-specific traits, adult workers treated

developing queens and workers indiscriminately, even in the presence of a fertile

queen. Together with previous studies showing queen control over sex allocation,

these results provide evidence that conflict over reproductive allocation is absent in

this ant.

Introduction

Ant queens and workers are fixed in irreversible, evolutionarily co-dependent roles,

analogous to the germ line and soma of multicellular organisms [1–3]. However,

compared to multicellular organisms, ant colonies can show relatively high levels of

within-unit conflict. This is because they are made up of non-clonal individuals, with

factors such as polygyny (multiple reproducing queens per colony) and polyandry

(multiple mating by queens) causing variation in relatedness among nestmates [4,5].

As a result, colony members can differ in their fitness optima regarding reproductive

allocation, creating the potential for conflict [6–8].

Whether potential conflict leads to actual conflict depends on the power each party

holds, in particular over reproduction and access to information [9–11]. Adult

reproductive power can be classified along a range of worker reproductive constraints

[12–15]: in species where workers retain fully or partly functional reproductive organs

(most ant genera), adult females can engage in conflict over reproductive dominance

or male parentage by participating in production of female and male offspring [16–

18]. In the few ant genera in which workers have lost reproductive organs entirely,

they cannot participate in reproduction directly. Nonetheless, because workers are

responsible for brood care, even in these species they can influence reproductive

allocation by manipulating the sex and caste ratios of brood, e.g by preferentially

rearing females into queens or by culling male or queen-destined brood [19–22].

Across the range of reproductive constraints, the fitness interests of developing

individuals also play an important role, in particular in the context of queen-worker

caste determination [21,23,24]. In all conflict scenarios, access to information about

brood maternity, sex and caste can be decisive, placing developing individuals at the

center of within-colony conflict [25,26].

Cardiocondyla ants represent the extreme end of the range of reproductive

constraints because workers lack reproductive organs, precluding any direct conflict

over reproduction. In *Cardiocondyla obscurior*, workers further refrain from

manipulating brood sex ratios, giving queens control over colony-level sex allocation

[27,28]. Recently, we found that in single queen colonies of this ant, production of

new queens increases as the resident queen ages, irrespective of absolute queen

fitness, queen lifespan or colony size [29]. This suggests that queens also control caste

allocation, and points towards relaxed selection on potentially costly traits related to

caste allocation conflict.

Here, we explore this idea by providing a description of the phenotypes of

developing *C. obscurior* ant queens and workers, made possible by the discovery of

crystalline deposits that distinguish castes from the egg stage onwards. We focus on

traits that are predicted to differ between castes and could transmit potential

information about caste in a social context, and test whether workers discriminate

between developing queens and workers. Our results support the hypothesis that

queens control primary caste ratios in eggs and workers refrain from manipulating

secondary caste ratios of brood, giving queens complete control over caste allocation

in this ant.

Results and Discussion

Identifying female caste in developing ants

"Auch ein blindes Huhn findet mal ein Korn"

Twenty years after the first *C. obscurior* colony was brought to the laboratory in

Regensburg, we discovered that queen and worker-destined eggs and larvae can be

distinguished by the caste-specific distribution of white spots in the posterior end

(Figures 1,2). To our knowledge, this is the first description of a discrete trait that can

be used to identify female caste over the entire course of ant development. Semi-thin

sections of third instar larvae viewed through a polarization filter revealed these spots

have a crystalline reflection. Queens exhibit a pair of clumped crystalline deposits on

the dorsal side, which become larger and more distinct as development progresses

(Figures 1,2, Figure S1). In late-instar queen larvae, crystalline deposits form large

aggregations around the developing ovaries, which already contain immature

oocytes (Figure 3). Workers do not have crystalline deposits in the egg stage (Figure

2), and small, randomly distributed deposits in larval stages (Figures 1, 3). In both

castes, crystalline deposits may consist of urate crystals, which are usually stored in

specialized cells, the urocytes, in the insect fat body [30], and which have been linked

to nutrient storage in ant larvae [31] and oocyte development in termite reproductives

[32].

Crystalline deposit patterns can be used to identify individuals by caste with high

accuracy, and workers successfully reared both castes at comparable rates in colony

fragments without queens (Table 1). This demonstrates that caste fate is fixed in the

embryo, likely under physiological control of the queen, and provides a first indication

that workers do not discriminate between developing queens and workers.

Similar crystalline deposit patterns that reliably predict caste were found in third instar

larvae of *C. nuda*, *C. venustula* as well as *C. wroughtonii,* the sister species to *C.*

obscurior (Figure S2, Table S1). No caste-specific patterns were found in *C.*

"argyrotricha" (provisional name of a recognized morphospecies to be described by

B. Seifert), *C. elegans*, *C. emeryi*, *C. minutior* and *C. thoracica* (Figure S3, Table S1),

or in any of the other 42 screened ant species (Figure S4, Table S2). This patchy

pattern does not follow the phylogenetic relationships within the *Cardiocondyla*

genus [33]. Caste-specific patterns are also not linked to the presence of the main

bacterial symbiont *Cand.* Westeberhardia cardiocondylae, an endosymbiont with a

highly reduced genome that resides in gut-associated bacteriomes, as well as in

ovaries in queens [34]. *Cardiocondyla minutior* carries the symbiont [35] but does not

display caste-specific patterns, while a *C. obscurior* lineage that is naturally free of

Cand. Westeberhardia [34,36] and has been maintained in the lab since 2011 shows

the patterns (Schultner, pers. obs.). This does not rule out a connection between

symbiont but indicates that a potential link must entail more than mere symbiont

presence. Future studies will uncover the metabolic processes from which the

deposits stem and reveal their function.

Phenotypes of developing queens and workers

General morphologies of ant larvae have been extensively described, e.g. [37–40], and non-invasive identification of larval sex has been demonstrated previously [41]. Traits that distinguish late-stage worker from sexual larvae (males, queens) have been discovered in six ant species from three subfamilies; these are largely continuous measures, including body size (*Aphaenogaster senilis* [42]; *Harpegnathos saltator* [43]; *Myrmica rubra* [44]), cuticle "shininess" (*Linepithema humile*, [45]), hair density (*Monomorium pharaonis* [46]), and hair morphology *(Acromyrmex echinatior* [47]). Apart from the crystalline deposits, *C. obscurior* queen and worker larvae appeared similar in their morphology, and neither the density nor the morphology of hairs differed between castes in any of the three instars (Figure S5). There was also no difference in size between the castes during early development (Figure 4, eggs: Linear mixed model, length: $\chi^2=0.123$, df=1, p=0.726; width: $\chi^2=0.007$, df=1, p=0.934; L1 larvae: ANOVA, head width: $F_{67,1}=3.009$, p=0.088; body length: $F_{52,1}=0.366$, p=0.548). However, as expected in an ant with morphologically distinct queen and worker castes [48], queen larvae grew larger than worker larvae as development progressed (Figure 4, L2 larvae: ANOVA, head width: $F_{60,1}=3.124$, p=0.088; body length: $F_{60,1}=14.542$, p<0.001; L3 larvae: ANOVA, head width: $F_{60,1}=17.198$, p<0.001; body length: $F_{60,1}=11.754$, p<0.01). Third instar queen larvae were also heavier than worker larvae (Figure 5A, L2 larvae: ANOVA, $F_{10,1}=3.547$, p=0.096; L3 larvae: ANOVA, $F_{10,1}=98.208$, p<0.001) but there was no difference in protein content between the two larval castes (Figure 5B, L2 larvae: ANOVA, $F_{41,1}=0.133$, p=0.717; L3 larvae: ANOVA, $F_{49,1}= 0.876$, p=0.354).

Chemical analyses of cuticular odors of queen and worker third instar larvae and adults of *C. obscurior* detected 32 compounds (Table 2). The profiles of adult queens and workers were slightly more complex than those of larvae (adults: >30 compounds including trace amounts below 0.5%; larvae: 24 compounds). Larval profiles were dominated by n-alkanes, with pentacosane and heptacosane constituting over 65% of the total profile. These two compounds were also present in high relative amounts

in adults, together with unsaturated and methyl-branched hydrocarbons. Principal

component analysis clearly separated samples by development stage on PC1, apart

from two adult samples, which clustered with larva samples for unknown reasons

(Figure 6, Table S2). Differences between development stages were confirmed by

analysis of similarities (adult vs. larva profiles: ANOSIM, R=0.628, p<0.001). PC2

separated adult castes, with a single worker sample clustering with queens (Figure 6,

Table S2, ANOSIM, R=0.548, p<0.001). In contrast, the profiles of queen larvae and

worker larvae did not differ (Figure 6, Table S2, ANOSIM, R=-0.059, p=0.651). A

similar lack of caste-specific larval odors has been demonstrated in late-stage larvae

of the ant *Aphaenogaster senilis* [42], while in *Harpegnathos saltator*, odor profiles of

queen-destined larvae were distinct from those of worker-destined larvae, allowing

workers to identify larvae by caste [43]. Ant larvae are often considered chemically

insignificant because they are easily accepted across colonies, populations and even

species (reviewed in [26]. At the same time, brood odors can play an important role

in queen fertility signaling within colonies, for example when eggs and larvae carry

odors that match those of queens [49,50]. Because of the relative difficulty of

distinguishing ant brood by developmental stage, sex and caste, detailed data remain

limited, and more studies are needed to understand the function of brood odors.

No sign of caste conflict involving brood

In line with the absence of caste-specific morphological and chemical cues, adult

workers did not adjust their behavior to the caste of encountered larvae in any of the

five tested scenarios. Workers were similarly attracted to queen and worker larvae

(Figure S6, number of contacts: ANOVA, $F_{44,1}$=3.193, p=0.077; duration of contacts:

Mann-Whitney-U-test, W=1184, p=0.072) and did not preferentially retrieve larvae of

either caste (queen larvae retrieved: 59% (40/67), worker larvae retrieved: 40%

(27/67), Chi-squared test, χ^2=2.522, df=1, p=0.113). Observations of feeding

behavior revealed no preference for either caste, with 65% (13/20) of worker larvae

and 59% (10/17) of queen larvae containing fluorescent dye in their gut two hours

after starved colonies composed of larvae, workers and queens were fed with

fluorescent honey (Fisher's exact test, odds ratio=0.774, p=0.745). There was no

difference in the overall survival of queen and worker larvae in these setups (survival

queen larvae: 68% (17/25); survival worker larvae: 80% (20/25); Fisher's exact test,

odds ratio=0.538, p=0.52). Workers also reared second instar larvae of both castes

to pupation with similar success when they were simultaneously presented with queen

and worker larvae in the presence of a mated queen (larvae reared to pupation:

queens: 38% (19/50); workers: 29% (14/48); Fisher's exact test, odds ratio=1.482,

p=0.398). Finally, when workers reared groups of first and second instar larvae of

unknown caste alone or in the presence of a fertile queen, they reared similar

proportions of new queens (L1 larvae - queen present: 42.5% queens (17/40); queen

absent: 32.6% queens (16/49); Fisher's exact test, odds ratio=1.517, p=0.382; L2

larvae - queen present: 13.7% queens (10/73); queen absent: 12.7% queens (7/55);

Fisher's exact test, odds ratio=1.089, p=1). Together, these results indicate that *C.*

obscurior workers are indifferent to the caste of developing individuals.

Conclusion

Previous studies in *C. obscurior* found that queens control egg sex ratios, and that

workers do not manipulate secondary sex ratios of brood [27,28]. In the present study

we show that female caste in this species is determined in the egg stage, indicating

that caste ratios are also under maternal control. Developing queens and workers

differed in size and weight in late larval stages, but not in cuticular odors, hair

morphology or physiology. Workers indiscriminately reared eggs and larvae of both

castes in different social contexts, including when a fertile queen was present. *C.*

obscurior queens thus appear to have control over sex and caste allocation, two

colony-level traits that often elicit conflict in social Hymenoptera.

Queen control of sex allocation has been demonstrated previously in ants [19,51–53],

but complete control over caste allocation has, to the best of our knowledge, not

been described. In other ant genera with fully sterile workers such as fire ants

(*Solenopsis*), Pharaoh ants (*Monomorium*) and big-headed ants (*Pheidole*), caste may

similarly be determined in the egg, allowing queens control over primary caste ratios

[13,54]; however, workers continue to play an important role in regulating secondary

caste ratios by culling sexual larvae in the presence of fertile queens or their

pheromonal equivalent [46,55–61].

Why then is *Cardiocondyla* different? Colonies are highly inbred due to the evolution

of wingless males with lifelong spermatogenesis, which remain in the nest over the

course of their relatively long lives to mate with multiple related queens, including

sisters and mothers [62–64]. Inbreeding results in particularly high levels of

relatedness, and consequently decreased conflict potential within colonies. *C.*

obscurior moreover has a relatively low degree of dimorphism between sexuals and

workers [48,65] and queens only have six ovarioles, producing about 1-2 eggs per

day and an average of 350 eggs over the course of their lives [29]. Accordingly,

colonies are small and contain on average one to several queens and ~30 workers

[65]). In comparison, *Monomorium*, *Solenopsis* and *Pheidole* queens are much larger

than their workers and highly fertile, with ovariole numbers ranging from 24 to over

100 [15,66], and they can produce hundreds of eggs per day, and hundreds of

thousands of eggs over the course of their lives. High queen fertility may allow these

species to adopt a strategy of brood culling to dynamically regulate sex and caste

allocation, possibly under constraints imposed by high dimorphism between sexuals

and workers such as the inability to rear sexuals when colonies are small, lack

development stages involved in nutrition processing, or resources are limited

[56,67,68]. In *C. obscurior*, low queen fertility means individuals eggs have high

relative value for all colony members, likely preventing the evolution of a similar

strategy. However, low dimorphism between sexuals and workers allows both castes

to be produced at comparable costs, so that even tiny colony fragments consisting

of only a few workers and brood can initiate a new colony. This provides an important

advantage in the exposed and often disturbed arboreal habitats where the ant thrives,

and may help explain its success as a global tramp species [69]. How the unique

combination of high relatedness, worker sterility, limited queen fertility and ecological

constraints resulted in the evolution of maternal control of egg caste, and the

physiological mechanisms underlying this process, remain to be uncovered.

Methods

Ants

Cardiocondyla obscurior and several other *Cardiocondyla* species are cosmotropical

tramp ants [70]. This myrmicine genus has evolved wingless fighter males with lifelong

spermatogenesis, which remain in the nest and mate with newly-hatched nestmate

queens [62]. *Cardiocondyla obscurior* has the smallest known ant genome (~193MB,

[65,71]). Adult queens and workers differ in size and morphology and workers lack

ovaries [48]. Larvae develop via three instars, which can be distinguished by body

shape and the degree of melanization of the mandibles [72]. All ants used for this

study were from the Old World lineage [71,73], maintained in the lab since 2010.

Stock and experimental colonies were kept in plaster-bottom nests with nest inserts

made of acrylic glass covered by dark foil in a climate chamber under a 12h/12h and

22°C/26°C night/day cycle at 70% humidity. All colonies were provided with water

and fed three times a week with honey, pieces of cockroaches and fruit flies.

Crystalline deposit patterns in queen and worker larvae of *Cardiocondyla*

Localization of crystalline deposits was characterized in eggs, first instar (L1), second

instar (L2) and third instar (L3) larvae as queen-like or worker-like by visually inspecting

brood from stock colonies under a stereomicroscope. For better detection of the

patterns, eggs and L1 larvae were submerged in a dissection dish containing 0.3%

PBT, after which they were mounted on a microscope slide and sealed with nail polish.

From each developmental stage, representative individuals with queen-like and

worker-like patterns were selected and photographed using a stereomicroscope

connected to a camera (Keyence VHX 500FD, Neu-Isenburg, Germany). Crystalline deposit patterns were characterized in the same manner in L3 larvae from eight additional *Cardiocondyla* species (*C. "argyrotricha"*, *C. elegans*, *C. emeryi*, *C. minutior*, *C. nuda*, *C. thoracica*, *C. venustula*, *C. wroughtonii*) spanning the phylogeny [33]. We further examined larvae of six species from four subfamilies available as live colonies and accessed photos of 36 species from seven subfamilies from Alex Wild's library (https://www.alexanderwild.com/Ants/Natural-History/Metamorphosis-Ant-Brood/), for a total of 51 examined species (Table S2).

In *C. obscurior*, the development of all stages was tracked to confirm that crystalline deposit patterns are associated with caste (Table 1). Eggs and L1 larvae were transferred to filter paper to remove excess buffer before being moved in groups of 10 to experimental colonies containing 10 workers each (eggs: queen-like=100, worker-like=100, L1 larvae: queen-like=100, worker-like =100). L2 and L3 larvae were transferred in groups of 10-15 to experimental colonies containing 10-12 workers (L2 larvae: queen-like=170, worker-like=175; L3 larvae: queen-like=50, worker-like=40).

We also tracked the development of L3 larvae from some of the other *Cardiocondyla* species (*C. minutior*, *C. nuda*, *C. thoracica*, *C. venustula*, *C. wroughtonii*) (Table S1). Experimental colonies were monitored three times per week and, when necessary, workers were added from the corresponding stock colonies to standardize worker number. Developing individuals were monitored until pupation, at which time they were counted and classified according to female caste, until no more brood remained. From these data, rates of survival until pupation and caste ratios were calculated. For *C. obscurior*, caste-specific survival and accuracy of caste prediction were tested with Fisher's exact tests (function fisher.test in R 4.0.3 [74]).

Larval histology

C. obscurior L3 larvae were collected from stock colonies and sorted according to crystalline deposit patterns. Sorted larvae were transferred into a fixation solution

consisting of 25% glutaraldehyde (GAH) in cacodylate buffer [(50 mM cacodylic acid,

pH 7.3) containing 150 mM Sucrose] (GAH : cacodylate buffer = 1 : 12,5) and kept

overnight at 11 °C. Samples were then rinsed in cacodylate buffer on ice, and fixated

in 4% osmium tetroxide in cacodylate buffer. After fixation, larvae were washed in

cacodylate buffer, dehydrated in a graded ethanol series and embedded in Epon.

Transversal semithin sections of 1 μm were cut and stained with methylene blue and

Azur II (Richardson's stain). Semi-thin sections were scanned with a Zeiss Primo Star

microscope and imaged with a Moticam 580 digital microscope camera.

Phenotypes of *C. obscurior* queen and worker-destined larvae

Size

Length and width of eggs with queen-like and worker-like crystalline deposit patterns

were assessed (n=39 queen-like eggs from 7 stock colonies; n=34 worker-like eggs

from 7 stock colonies). Each egg was photographed through a stereomicroscope

connected to a camera at 200x magnification and its length and width measured

using the system software (Keyence VHX 500FD, Neu-Isenburg, Germany). Caste-

specific egg lengths and widths were compared using linear mixed models with

length or width as response variables and larval caste as explanatory variable. To

account for between-colony variation, the stock colonies from which eggs originated

were implemented as random variables. Model residuals were checked for normality

using Shapiro-Wilks tests (function shapiro.test) and residual tests implemented in the

DHarma package (function testResiduals [75]) in R 4.0.3 [74]. Size differences between

castes were assessed using a Wald $\mathbf{X}^2$ test (function Anova, package car [76]).

To assess the size of queens and workers over the course of larval development, head

capsule width and body length were measured (L1 larvae: n=35 queens, n=32

workers; L2 larvae: n=30 queens, n=30 workers; L3 larvae: n=30 queens, n=30

workers). Individuals were collected from stock colonies and sorted by instar and

caste, and then anaesthetized with CO_2 for four hours. Larvae were placed on their

backs and photographed using a JEOL scanning electron microscope (L1 larvae) or a

stereomicroscope connected to a camera (L2 & L3 larvae). Measurements were taken

using ImageJ [77] after setting the appropriate scales. In each instar, head capsule

width and body length of queen and worker larvae were compared using linear

regressions with width or length as response variables and larval caste as explanatory

variable. Model residuals were checked for normality using Shapiro-Wilks tests

(function shapiro.test) and residual tests (function testResiduals [75]) in R 4.0.3 [74].

Size differences between castes were assessed using a Wald X^2 test (function Anova,

package car [76]).

Weight

Five pools of five larvae each per instar (L2, L3) and caste (queen, worker) were

collected from stock colonies and weighed on a Sartorius SC2 fine scale (Sartorius,

Göttingen, Germany). Weight differences were analysed within each instar with a

linear regression using weight per individual larva in mg as response variable and

caste as explanatory variable. Model residuals were checked for normality using the

Shapiro-Wilks test (function shapiro.test) and residual tests (function testResiduals

[75]) and differences between castes were assessed using a Wald X^2 test (function

Anova, package car [76]) in R 4.0.3 [74].

Hair density and morphology

Hair density and morphology was assessed qualitatively in queen and worker larvae

of all three instars (n=3 per caste and instar). Larvae were collected from stock

colonies and sorted by instar and caste, and then placed on a self-adhesive sample

insert and moved to -20°C for seven minutes. Freshly killed larvae were scanned with

a JEOL scanning electron microscope and density and morphology of hairs was

compared visually.

Protein content

L2 and L3 queen and worker larvae were collected from stock colonies and fixed

individually in 15 µl of 1x PBS solution and frozen at -20°C until further analysis (n=30

per instar and caste). For analysis the samples were allowed to thaw on ice and were

then homogenized using pestles. Protein concentration in the samples was measured

using the Bicinchoninic Acid Protein Assay Kit (Sigma Aldrich, St. Louis, MO, USA)

according to the manufacturer's protocol with the following modifications: half of the

volume for the microplate assay was used, resulting in 100 µl of working reagent and

12.5 µl of sample consisting of 3 µl of homogenized ant sample and 9.5 µl of 1x PBS.

All samples were run in two technical replicates at an absorbance of 526 nm using a

SpectraMax iD3 microplate reader (Molecular Devices, San Diego, CA, USA). The

average of the two technical replicates was used for further calculations. Protein

concentration was calculated using a standard curve created with BSA (bovine serum

albumin). Within each instar, protein content of queen and worker larvae was

compared using a linear regression with protein content in µg/ml as response variable

and larval caste as explanatory variable. Model residuals were checked for normality

using a Shapiro-Wilks test and residual tests implemented in the DHARMa package.

Differences between castes and instars were assessed with a Wald X^2 test. A Levene's

test was used to compare the variances of protein content between castes; this was

done separately for each instar.

Cuticular odors

Cuticular hydrocarbon (CHC) profiles of pools of L3 queen-destined larvae (20

individuals each, n=6), L3 worker-destined larvae (20 individuals each, n=8), adult

queens and adult workers (10 individuals each, n=9 for both groups) were analyzed

using gas chromatography coupled with mass-spectrometry (GC-MS). For each

sample, individuals from at least five stock colonies were pooled to account for

colony-level differences. To obtain samples free of contamination from nest material,

individuals used in analyses were collected from stock colonies, separated by caste

and kept in freshly-plastered experimental nests for ~24h prior to sampling. All
samples were frozen in glass screw top vials and stored at -20°C until extraction.

Samples were extracted in 100 µl HPLC grade hexane for 1 minute. Hexane extracts
were then fractionated using silica columns (Chromabond SiOH, 1 mL/100mg).
Columns were rinsed consecutively with 1 ml of hexane, dichloromethane, methanol
and again dichloromethane, before being conditioned with 2 ml of hexane and
adding the sample. The non-polar fraction containing hydrocarbons was eluted with
1 ml of hexane, and subsequently evaporated under a stream of nitrogen. After being
re-dissolved in 100 µl of hexane, 1 µl per sample was subjected to GCMS (Shimadzu
GCMS-QP2010 Plus) on a SGE BPX-5 column (30m × 0.25µm × 0.25mm). Helium gas
(1 mL/min) served as carrier gas, and injection was splitless. The temperature program
started at 70 °C isothermal for 1 min, after which the temperature was raised to 200
°C at 30 °C/min and then from 200 to 320 °C at 5 °C/min, where it was held for 5 min.

Compound determination of peaks was based on their characteristic mass spectra
and matching of retention indices (for methyl-branched alkanes: [78]. For peaks of a
concentration too low to yield an unambiguous mass spectrum, compounds were
considered to be identical to those present in other samples if their retention indices
matched. Compounds that could not be identified were labeled with letters, while
double-bond positions of unsaturated hydrocarbons were not further elucidated and
are indicated with "x" (Table 2). Only peaks with retention times over 15 minutes
(docosane and longer chain lengths) were used for quantitative analyses. Relative
peak areas were calculated by dividing individual peak areas by the summed area of
all hydrocarbon peaks. Values were then square-root transformed, centered and
scaled prior to principal component analysis (PCA, function prcomp in R).
Differentiation between groups was tested on square-root transformed values with
analysis of similarities (ANOSIM, function anosim in R); tests were run separately for
comparisons between developmental stages (adults vs. larvae) and castes within each
stage (adult queens vs. adult workers, queen larvae vs. worker larvae).

Worker behavior toward queen and worker larvae in different in social contexts

The behavior of adults toward queen and worker larvae was investigated in five scenarios.

First, we investigated whether isolated worker ants are differentially attracted to L3 queen and worker larvae. This was done in round observation chambers (diameter 20 mm), which were visually divided into two equally sized halves. In the center of each half, a clean piece of plastic foil with either a L3 queen or worker larva from the same stock colony was placed, so that each chamber contained one larva of each caste. Larvae were glued to the plastic foil with a small drop of sucrose solution to prevent workers from moving larvae around during the observation period. To avoid potential side bias, queen and worker larvae were randomly placed in the left and right halves of the chambers. Once larvae were placed in the chamber, one worker from the same stock colony was added and a transparent microscope slide placed on top to prevent workers from escaping. Chambers were observed for 12 minutes and the number and duration of contacts, i.e., instances of bodily contact between the adult worker and either queen or worker larva, were scored using JWatcher software [79] (n=44 chambers observed in total). The number of contacts was analysed with a linear regression using larval caste as the explanatory variable. Model residuals were checked for normality using the Shapiro-Wilks test and residual tests implemented in the DHARMa package. The duration of contacts did not follow a normal distribution and was analysed with a Mann-Whitney-U-test (function wilcox.test) in R.

Second, we tested whether worker ants retrieve L3 queen and worker larvae at different rates. From each of five stock colonies, one experimental colony was set up containing 12 workers and one mated queen. Colonies were allowed to settle for 48h. One L3 worker larva and one L3 queen larva from the original stock colonies were then added to the corresponding experimental colonies on a clean piece of filter paper. Experimental colonies were observed until the first larva was retrieved into the

nest and the caste of the retrieved larva was noted. This procedure was repeated at

least 11 times in each experimental colony, until a total of 67 retrieval events were

documented. Retrieval rates of the two castes were compared using a **X**^{2} Test.

Third, we investigated whether workers display preferential treatment towards larvae

of either caste under food limitation. Five experimental colonies were set up, each

containing one mated queen, five workers, and five L2 larvae of each caste. Colonies

were then starved of sucrose for three days. On day three, colonies were presented

with 0.25 µl of a 1.5 M sucrose solution containing 0.08 g l^{-1} Rhodamine B (Czaczkes

et al., 2019). After two hours, all larvae were counted and examined for the presence

of Rhodamine B in the digestive tract under a green laser light (555 nm) using a Zeiss

Axioplan-2 fluorescence microscope. Larvae positive for Rhodamine B were

considered "fed", while the absence of fluorescence in larvae was scored as "unfed".

The number of fed and unfed larvae of each caste, as well as the overall survival of

queen and worker larvae, was compared using Fisher's exact tests (function fisher.test

in R).

Fourth, we assessed whether workers changed their rearing behavior in the presence

of a mated queen when presented with queen and worker larvae simultaneously. Five

colonies were set up, each with 10 second instar queen and 10 second instar worker

larvae, as well as 20 workers and one mated queen from the same stock colony.

Colonies were fed as described above and all pupae counted and their sex and caste

noted. Caste-specific survival of larvae was compared using Fisher's exact test

(function fisher.test in R).

Fifth, we tested whether the presence of a mated queen influenced secondary caste

ratios in brood of unknown caste. First and second instar larvae of unknown caste

were collected from stock colonies and reared in experimental colonies with and

without queens. For each instar separately, colonies were set up containing 15 larvae

and 15 workers, as well as one mated queen (L1 n=10, L2 n=8) or no queen (L1 n=13,

L2 n=9). In each experimental colony, all individuals came from the same stock
colony. Colonies were fed as described above and all pupae counted and their sex
and caste noted. The proportion of queens produced in colonies with and without a
mated queen was compared using Fisher's exact test (function fisher.test in R).

Acknowledgements

We thank B. Lautenschläger and M. Schiwek for help with histology, E. Strohm for
help identifying the crystalline nature of the deposits, and A. Koch, K. Pogorelski, A.
Reisinger and C. Winkler for help with experiments. JO dedicates this paper to Margit
Oettler, for nature and nurture. This study was funded by DFG grants (DFG OE 549/3-
1, OE 549/3-2, OE 549/3-3).

References

1.   Maynard Smith J, Szathmary E. 1995 *The major transitions in evolution.*
Oxford, UK: Oxford University Press.

2.   Boomsma JJ, Gawne R. 2018 Superorganismality and caste differentiation as
points of no return: how the major evolutionary transitions were lost in
translation. *Biol. Rev.* **93**, 28–54. (doi:10.1111/brv.12330)

3.   Bourke AFG. 2011 *Principles of social evolution.* Oxford, UK: Oxford
University Press.

4.   Bourke AFG, Franks NR. 1995 *Social Evolution in Ants.* Princeton, NJ:
Princeton University Press.

5.   Keller L. 1995 Social life: the paradox of multiple-queen colonies. *Trends Ecol.*
Evol. **10**, 355–360.

6.   Ratnieks FLW, Reeve HK. 1992 Conflict in single-queen Hymenopteran
societies: the structure of conflict and processes that reduce conflict in
advanced eusocial species. *J. Theor. Biol.* **158**, 33–65.

7.   Sundström L, Boomsma JJ. 2001 Conflicts and alliances in insect families.

Heredity (Edinb). **86**, 515–521.

8.   Heinze J. 2004 Reproductive Conflict in Insect Societies. *Adv. Study Behav.*
487
488   **34**, 1–57. (doi:10.1016/S0065-3454(04)34001-5)

9.   Beekman M, Komdeur J, Ratnieks FLW. 2003 Reproductive conflicts in social
489
490   animals: who has power? *Trends Ecol. Evol.* **18**, 277–282. (doi:10.1016/S0169-
491   5347(03)00068-5)

10.   Beekman M, Ratnieks FLW. 2003 Power over reproduction in social
492
493   Hymenoptera. *Philos. Trans. R. Soc. B Biol. Sci.* **358**, 1741–1753.
494   (doi:10.1098/rstb.2002.1262)

11.   Ratnieks FLW, Foster KR, Wenseleers T. 2006 Conflict resolution in insect
495
496   societies. *Annu. Rev. Entomol.* **51**, 581–608.
497   (doi:10.1146/annurev.ento.51.110104.151003)

12.   Khila A, Abouheif E. 2008 Reproductive constraint is a developmental
498
499   mechanism that maintains social harmony in advanced ant societies. *Proc.*
500   *Natl. Acad. Sci.* **105**, 17884–17889. (doi:10.1073/pnas.0807351105)

13.   Khila A, Abouheif E. 2010 Evaluating the role of reproductive constraints in
501
502   ant social evolution. *Philos. Trans. R. Soc. B Biol. Sci.* **365**, 617–630.
503   (doi:10.1098/rstb.2009.0257)

14.   Ronai I, Vergoz V, Oldroyd BP. 2016 The Mechanistic, Genetic, and
504
505   Evolutionary Basis of Worker Sterility in the Social Hymenoptera. In *Advances*
506   *in the Study of Behavior*, pp. 251–317. Elsevier Ltd.
507   (doi:10.1016/bs.asb.2016.03.002)

15.   Gotoh A, Billen J, Hashim R, Ito F. 2016 Degeneration patterns of the worker
508
509   spermatheca during morphogenesis in ants (Hymenoptera: Formicidae). *Evol.*
510   *Dev.* **18**, 96–104. (doi:10.1111/ede.12182)

16.   Hammond RL, Keller L. 2004 Conflict over male parentage in social insects.
511

PLoS Biol. **2**, E248. (doi:10.1371/journal.pbio.0020248)

17.   Wenseleers T, Ratnieks FLW. 2006 Comparative analysis of worker

reproduction and policing in eusocial hymenoptera supports relatedness

theory. *Am. Nat.* **168**, E163–E179. (doi:10.1086/508619)

18.   Monnin T. 1999 Dominance hierarchy and reproductive conflicts among

subordinates in a monogynous queenless ant. *Behav. Ecol.* **10**, 323–332.

(doi:10.1093/beheco/10.3.323)

19.   Rosset H, Chapuisat M. 2006 Sex allocation conflict in ants: when the queen

rules. *Curr. Biol.* **16**, 328–331. (doi:10.1016/j.cub.2005.12.036)

20.   Helanterä H, Ratnieks FLW. 2009 Sex allocation conflict in insect societies:

who wins? *Biol. Lett.* **5**, 700–704. (doi:10.1098/rsbl.2009.0501)

21.   Bourke AFG, Ratnieks FLW. 1999 Kin conflict over caste determination in

social Hymenoptera. *Behav. Ecol. Sociobiol.* **46**, 287–297.

(doi:https://doi.org/10.1007/s002650050622)

22.   Mehdiabadi NJ, Reeve HK, Mueller UG. 2003 Queens versus workers: sex-

ratio conflict in eusocial Hymenoptera. *Trends Ecol. Evol.* **18**, 88–93.

(doi:10.1016/S0169-5347(02)00056-3)

23.   Dobata S. 2012 Arms race between selfishness and policing: two-trait

quantitative genetic model for caste fate conflict in eusocial Hymenoptera.

Evolution (N. Y). **66**, 3754–3764. (doi:10.1111/j.1558-5646.2012.01745.x)

24.   Wenseleers T, Ratnieks FLW. 2004 Tragedy of the commons in *Melipona*

bees. *Proc. R. Soc. B Biol. Sci.* **271**, S310–S312. (doi:10.1098/rsbl.2003.0159)

25.   Schultner E, Oettler J, Helanterä H. 2017 The role of brood in eusocial

Hymenoptera. *Q. Rev. Biol.* **92**, 39–78. (doi:doi.org/10.1086/690840)

26.   Schultner E, Pulliainen U. 2020 Brood recognition and discrimination in ants.

Insectes Soc. **67**, 11–34. (doi:10.1007/s00040-019-00747-3)

27. Cremer S, Heinze J. 2002 Adaptive production of fighter males: queens of the
ant *Cardiocondyla* adjust the sex ratio under local mate competition. *Proc. R.*
Soc. B Biol. Sci. **269**, 417–422. (doi:10.1098/rspb.2001.1892)

28. De Menten L, Cremer S, Heinze J, Aron S. 2005 Primary sex ratio adjustment
by ant queens in response to local mate competition. *Anim. Behav.* **69**, 1031–
1035. (doi:10.1016/j.anbehav.2004.09.005)

29. Jaimes-Niño LM, Heinze J, Oettler J. 2022 Late-life fitness gains and
reproductive death in Cardiocondyla obscurior ants. *Elife* **11**, e74695.

30. Paes de Olivaira T V., Da Cruz Landim C. 2003 Morphology and Function of
Insect Fat Body Cells: a Review. *Biociencias* **11**, 195–205.

31. Zara FJ, Caetano FH. 2004 Ultramorphology and histochemistry of fat body
cells from last Instar larval of the *Pachycondyla (=Neoponera) villosa*
(Fabricius) (Formicidae: Ponerinae). *Brazilian J. Biol.* **64**, 725–735.
(doi:10.1590/s1519-69842004000400022)

32. Elliott KL, Stay B. 2007 Juvenile hormone synthesis as related to egg
development in neotenic reproductives of the termite *Reticulitermes flavipes*,
with observations on urates in the fat body. *Gen. Comp. Endocrinol.* **152**,
102–110. (doi:10.1016/j.ygcen.2007.03.003)

33. Oettler J, Suefuji M, Heinze J. 2010 The evolution of alternative reproductive
tactics in male *Cardiocondyla* ants. *Evolution (N. Y).* **64**, 3310–3317.
(doi:10.1111/j.1558-5646.2010.01090.x)

34. Klein A *et al.* 2016 A novel intracellular mutualistic bacterium in the invasive
ant *Cardiocondyla obscurior. ISME J.* **10**, 376–388.
(doi:10.1038/ismej.2015.119)

35. Jackson R *et al.* 2022 Convergent evolution of a labile nutritional symbiosis in
ants. *ISME J.* **16**, 2114–2122. (doi:10.1038/s41396-022-01256-1)

36. Ün Ç, Hacker J, Heinze J, Oettler J, Schultner E. 2021 Endosymbionts
mediate the effects of antibiotic exposure in the tramp ant *Cardiocondyla*
obscurior. Ecol. Entomol. , 13099. (doi:10.1111/een.13099)

37. Wheeler GC, Wheeler J. 1976 Ant Larvae: Review and Synthesis. *Mem.*
Entomol. Soc. Washingt. **7**, 1–108.

38. Wheeler GC, Wheeler J. 1986 Ten-year supplement to 'Ant larvae: review and
synthesis'. *Proc. Entomol. Soc. Washingt.* **88**, 684–702.

39. Wheeler W. 1918 A study of some ant larvae, with a consideration of the
origin and meaning of the social habit among insects. *Proc. Am. Philos. Soc.*
57, 293–343.

40. Solis DR, Fox EGP, Kato LM, de Jesus CM, Yabuki AT, Campos AEDC, Bueno
OC. 2010 Morphological description of the immatures of the ant,
Monomorium floricola. J. Insect Sci. **10**, 15. (doi:10.1673/031.010.1501)

41. Penick CA, Ebie J, Moore D. 2014 A non-destructive method for identifying
the sex of ant larvae. *Insectes Soc.* **61**, 51–55. (doi:10.1007/s00040-013-0323-
5)

42. Villalta I, Amor F, Cerdá X, Boulay R. 2016 Social coercion of larval
development in an ant species. *Sci. Nat.* **103**, 18. (doi:10.1007/s00114-016-
1341-8)

43. Penick CA, Liebig J. 2017 A larval 'princess pheromone' identifies future ant
queens based on their juvenile hormone content. *Anim. Behav.* **128**, 33–40.
(doi:10.1016/j.anbehav.2017.03.029)

44. Brian MV. 1973 Queen recognition by brood-rearing workers of the ant
Myrmica rubra L. *Anim. Behav.* **21**, 691–698. (doi:10.1016/S0003-
3472(73)80093-4)

45. Passera L, Aron S, Bach D. 1995 Elimination of sexual brood in the Argentine

ant *Linepithema humile*: queen effect and brood recognition. *Entomol. Exp.*
Appl. **75**, 203–212. (doi:10.1111/j.1570-7458.1995.tb01928.x)

46.   Edwards J. 1991 Caste regulation in the pharaoh's ant *Monomorium*
pharaonis: recognition and cannibalism of sexual brood by workers. *Physiol.*
Entomol. **16**, 263–271. (doi:doi.org/10.1111/j.1365-3032.1991.tb00565.x)

47.   Adams RMM, Larsen RS, Stylianidi N, Cheung D, Qiu B, Murray S, Zhang G,
Boomsma J. 2021 Hairs distinguish castes and sexes: identifying the early
ontogenetic building blocks of a fungus-farming superorganism.
Myrmecological News **31**, 201–216. (doi:10.25849/myrmecol.news)

48.   Oettler J, Platschek T, Schmidt C, Rajakumar R, Favé M-J, Khila A, Heinze J,
Abouheif E. 2019 Interruption points in the wing gene regulatory network
underlying wing polyphenism evolved independently in male and female
morphs in *Cardiocondyla* ants. *J. Exp. Zool. Part B Mol. Dev. Evol.* **332**, 7–16.
(doi:10.1002/jez.b.22834)

49.   Holman L, Jørgensen CG, Nielsen J, d'Ettorre P. 2010 Identification of an ant
queen pheromone regulating worker sterility. *Proc. R. Soc. B Biol. Sci.* **277**,
3793–3800. (doi:10.1098/rspb.2010.0984)

50.   Endler A, Liebig J, Schmitt T, Parker JE, Jones G, Schreier P, Hölldobler B.
2004 Surface hydrocarbons of queen eggs regulate worker reproduction in a
social insect. *Proc. Natl. Acad. Sci.* **101**, 2945–2950.
(doi:10.1073/pnas.0308447101)

51.   Passera L, Aron S, Vargo EL, Keller L. 2001 Queen control of sex ratio in fire
ants. *Science (80-.).* **293**, 1308–1310. (doi:10.1126/science.1062076)

52.   De Menten L, Fournier D, Brent CS, Passera L, Vargo EL, Aron S. 2005 Dual
mechanism of queen influence over sex ratio in the ant *Pheidole pallidula*.
Behav. Ecol. Sociobiol. **58**, 527–533. (doi:10.1007/s00265-005-0964-0)

53.   Aron S, Passera L, Keller L. 1994 Queen-worker conflict over sex ratio: A
comparison of primary and secondary sex ratios in the Argentine ant,
Iridomyrmex humilis. J. Evol. Biol. **7**, 403–418. (doi:10.1007/s10958-007-0251-
x)

54.   Passera L, Suzzoni JP. 1979 Le role de la reine de *Pheidole pallidula* (Nyl.)
(Hymenoptera, Formicidae) dans la sexualisation du couvain après traitement
par l'hormone juvénile. *Insectes Soc.* **26**, 343–353. (doi:10.1007/BF02223553)

55.   Klobuchar E, Deslippe R. 2002 A queen pheromone induces workers to kill
sexual larvae in colonies of the red imported fire ant (*Solenopsis invicta*).
Naturwissenschaften **89**, 302–304. (doi:10.1007/s00114-002-0331-1)

56.   Passera L. 1980 La ponte d'œufs préorientés chez la fourmi *Pheidole pallidula*
(Nyl.) (Hymenoptera-Formicidae). *Insectes Soc.* **27**, 79–95.
(doi:10.1007/BF02224522)

57.   Vargo EL, Passera L. 1991 Pheromonal and behavioral queen control over the
production of gynes in the Argentine ant *Iridomyrmex humilis* (Mayr). *Behav.*
Ecol. Sociobiol. **28**, 161–169.

58.   Edwards J. 1987 Caste regulation in the pharaoh's ant *Monomorium*
pharaonis: the influence of queens on the production of new sexual forms.
Physiol. Entomol. **12**, 31–39. (doi:doi.org/10.1111/j.1365-
3032.1987.tb00721.x)

59.   Warner MR, Lipponen J, Linksvayer TA. 2018 Pharaoh ant colonies
dynamically regulate reproductive allocation based on colony demography.
Behav. Ecol. Sociobiol. **72**, 31. (doi:10.1007/s00265-017-2430-1)

60.   Boonen S, Billen J. 2017 Caste regulation in the ant *Monomorium pharaonis*
(L.) with emphasis on the role of queens. *Insectes Soc.* **64**, 113–121.
(doi:10.1007/s00040-016-0521-z)

61.   Oliveira RC, Warson J, Sillam-Dussès D, Herrera-Malaver B, Verstrepen K,
Millar JG, Wenseleers T. 2020 Identification of a queen pheromone mediating
the rearing of adult sexuals in the pharaoh ant *Monomorium pharaonis*. *Biol.*
Lett. **16**, 20200348. (doi:10.1098/rsbl.2020.0348rsbl20200348)

62.   Heinze J, Hölldobler B. 1993 Fighting for a harem of queens: physiology of
reproduction in *Cardiocondyla* male ants. *Proc. Natl. Acad. Sci.* **90**, 8412–
8414.

63.   Heinze J. 2016 The male has done his work - The male may go. *Curr. Opin.*
Insect Sci. **16**, 22–27. (doi:10.1016/j.cois.2016.05.005)

64.   Schmidt CV, Frohschammer S, Schrempf A, Heinze J. 2014 Virgin ant queens
mate with their own sons to avoid failure at colony foundation.
Naturwissenschaften **101**, 69–72. (doi:10.1007/s00114-013-1126-2)

65.   Schrader L *et al.* 2014 Transposable element islands facilitate adaptation to
novel environments in an invasive species. *Nat. Commun.* **5**, 5495.
(doi:10.1038/ncomms6495)

66.   Tschinkel WR. 1987 Relationship between Ovariole Number and
Spermathecal Sperm Count in Ant Queens: a New Allometry. *Ann. Entomol.*
Soc. Am. **80**, 208–211. (doi:10.1093/aesa/80.2.208)

67.   Warner MR, Kovaka K, Linksvayer TA. 2016 Late-instar ant worker larvae play a
prominent role in colony-level caste regulation. *Insectes Soc.* **63**, 1–9.
(doi:10.1007/s00040-016-0501-3)

68.   Tschinkel WR. 1995 Stimulation of fire ant queen fecundity by a highly specific
brood stage. *Ann. Entomol. Soc. Am.* **88**, 876–882.

69.   Heinze J. 2017 Life-history evolution in ants: the case of *Cardiocondyla*. *Proc.*
R. Soc. B Biol. Sci. **284**, 20161406. (doi:10.1098/rspb.2016.1406)

70.   Heinze J, Cremer S, Eckl N, Schrempf A. 2006 Stealthy invaders: The biology

of *Cardiocondyla* tramp ants. *Insectes Soc.* **53**, 1–7. (doi:10.1007/s00040-005-
0847-4)

71. Errbii M, Keilwagen J, Hoff KJ, Steffen R, Altmüller J, Oettler J, Schrader L.
2021 Transposable elements and introgression introduce genetic variation in
the invasive ant *Cardiocondyla obscurior*. *Mol. Ecol.* , mec.16099.
(doi:10.1111/mec.16099)

72. Schrempf A, Heinze J. 2006 Proximate mechanisms of male morph
determination in the ant *Cardiocondyla obscurior*. *Evol. Dev.* **8**, 266–272.
(doi:10.1111/j.1525-142X.2006.00097.x)

73. Ün Ç, Schultner E, Manzano-Marín A, Flórez L V., Seifert B, Heinze J, Oettler
J. 2021 Cytoplasmic incompatibility between Old and New World populations
of a tramp ant. *Evolution (N. Y).* **75**, 1775–1791. (doi:10.1111/evo.14261)

74. R Team. 2016 R: A Language and Environment for Statistical Computing.

75. Hartig F. 2020 DHARMa: Residual Diagnostics for Hierarchical (Multi-Level /
Mixed) Regression Models.

76. Fox J, Weisberg S. 2019 *An R Companion to Applied Regression*. Third edit.
Thousand Oaks, CA: Sage. See
https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

77. Schneider C, Rasband W, Eliceiri K. 2012 NIH Image to ImageJ: 25 years of
image analysis. *Nat. Methods* **9**, 671–675.
(doi:https://doi.org/10.1038/nmeth.2089)

78. Carlson DA, Bernier UR, Sutton BD. 1998 Elution Patterns from Capillary GC
for Methyl-Branched Alkanes. *J. Chem. Ecol.* **24**, 1845–1865.
(doi:10.1023/A:1022311701355)

79. Blumstein DT, Daniel J. 2007 *Quantifying Behavior the JWatcher way.*
Sunderland, MA: Sinauer Associates.

Figure legends

Figure 1: Crystalline deposit patterns distinguish queen- and worker-destined larvae

in the ant *Cardiocondyla obscurior*. Stereomicroscope images of queen-destined and

worker-destined larval instars show a pair of large, clumped crystalline deposits (cd)

in the posterior end of queens, and small, randomly distributed crystalline deposits

in workers (lateral and dorsal views).

Figure 2: Crystalline deposit patterns in queen- and worker-destined embryos in the

ant *Cardiocondyla obscurior*. Queen-destined embryos have crystalline deposits (cd)

localized in the last segments of their germ band (gb). Worker-destined embryos do

not show any crystalline deposits. The first segments of the embryo are separated

into the head and the three gnathal segments: mandibular (md), maxillary (mx) and

labial (lb).

Figure 3: Histological sections of queen and worker-destined third instar larvae. (A)

Queen-destined larvae show paired ovaries (ov) close to the midgut (mid). (B,C)

Ovaries in queen-destined larvae are surrounded by large crystalline deposits. (D) In

workers ovaries are missing. (E, F) In worker-destined larvae crystalline deposits are

distributed randomly and aggregate close to trophocytes (t). top row: 5x

magnification, middle row: 20x magnification, bottom row: 40x magnification.

Figure 4: Size of *Cardiocondyla obscurior* queen and worker eggs and larvae (A) egg

width, (B) egg length, (C) larva head capsule width, (D) larva body length. L1: first

instar, L2: second instar, L3: third instar

Figure 5: Weight (A) and protein content (B) of queen and worker *Cardiocondyla*

obscurior larvae. L2: second instar, L3: third instar

Figure 6: Principal component analysis of cuticular hydrocarbon profiles of queen and

worker larvae and adults in the ant *Cardiocondyla obscurior*

Figures

Figure 1: Crystalline deposit patterns distinguish queen- and worker-destined larvae in the ant *Cardiocondyla obscurior*. Stereomicroscope images of queen-destined and worker-destined larval instars show large, paired crystalline deposits (cd) in the posterior end of queens, and small, randomly distributed crystalline deposits in workers (lateral and dorsal views).

Figure 2: Crystalline deposit patterns in queen- and worker-destined embryos in the ant *Cardiocondyla obscurior*. Queen-destined embryos have crystalline deposits (cd) localized in the last segments of their germ band (gb). Worker-destined embryos do not show any crystalline deposits. The first segments of the embryo are clearly separated into the head and the three gnathal segments: mandibular (md), maxillary (mx) and labial (lb).

Figure 3: Histological sections of queen and worker-destined third instar larvae. (A) Queen-destined larvae show paired ovaries (ov) close to the midgut (mid). (B,C) Ovaries in queen-destined larvae are surrounded by large crystalline deposits. (D) Transversal section of a worker-destined L3 larva; in workers ovaries are missing. (E, F) In worker-destined larvae crystalline deposits are distributed randomly and aggregate close to trophocytes (t). top row: 5x magnification, middle row: 20x, bottom row: 40x.

Figure 4: Size of *C. obscurior* queen and worker eggs and larvae (A) egg width, (B) egg length, (C) larva head capsule width, (D) larva body length. L1: first instar, L2: second instar, L3: third instar

Figure 5: Weight (A) and protein content (B) of queen and worker ant larvae L2: second instar, L3: third instar

Figure 6: Principal component analysis of cuticular hydrocarbon profiles of queen and worker larvae and adults in the ant *Cardiocondyla obscurior*

Table 1: Localization of crystalline deposits predicts caste in eggs and larvae of the ant *Cardiocondyla obscurior*

Development Stage	Crystalline deposit pattern	n	Proportion of pupae produced	Survival differences between predicted castes		Proportion of correct predictions	Accuracy of caste prediction	
				Odds ratio	p		Odds ratio	p
egg	queen-like	100	56% (56/100)	0.664	0.157	82.1% (46/56)	83.033	**<0.001**
	worker-like	100	45% (45/100)			95.1% (39/41)		
1st instar larva	queen-like	100	47% (47/100)	0.961	1	57.4% (27/47)	28.573	**<0.001**
	worker-like	100	46% (46/100)			95.6% (44/46)		
2nd instar larva	queen-like	170	58.2% (99/170)	0.648	0.052	96.9% (96/99)	2133.229	**<0.001**
	worker-like	175	47.4% (83/175)			98.8% (82/83)		
3rd instar larva	queen-like	50	60% (30/50)	0.497	0.137	100% (30/30)	infinite	**<0.001**
	worker-like	40	42.5% (17/40)			100% (17/17)		

Note that 4 out 45 of eggs predicted to be workers developed into males; these were excluded prior to analysis of caste prediction accuracy. No males developed in any of the other setups.

Table 2: Cuticular hydrocarbon profiles of queen and worker larvae and adults of the ant *Cardiocondyla obscurior*

RI	Compound	mean relative amounts ± standard deviation (%)			
		Queen larvae	Worker larvae	Queens	Workers
2189	C22	3.78 ± 2.66	3.89 ± 2.83	2.11 ± 1.29	6.25 ± 2.49
2300	A [a]	1.2 ± 0.72	0.83 ± 0.57	0.51 ± 0.12	1.49 ± 0.52
2400	C24	2.97 ± 1.71	2.77 ± 1.68	1.63 ± 0.89	3.28 ± 0.64
2499	C25	50.49 ± 7.14	50.37 ± 4.92	26.28 ± 8.93	13.48 ± 6.99
2532	13-Me C25 [b]	tr	tr	0.51 ± 0.16	1.25 ± 0.64
2572	3-Me C25	3.27 ± 1.68	2.61 ± 1.14	4.66 ± 1.22	8.04 ± 1.28
2599	C26	3.13 ± 0.96	3.18 ± 1.19	2.47 ± 0.82	3.5 ± 0.53
2632	B	tr	tr	tr	0.64 ± 0.41
2662	C	tr	tr	tr	tr
2675	x,x-C27:2	1.1 ± 0.42	0.59 ± 0.31	7.28 ± 1.27	13.24 ± 7.64
2699	C27	17.89 ± 4.13	20.8 ± 3.68	25.68 ± 5.05	9.65 ± 6.68
2731	13-Me, 11-Me C27	1.46 ± 0.76	0.94 ± 0.44	3.3 ± 1.39	12.56 ± 5.46
2749	5-Me C27	tr	tr	tr	0.58 ± 0.26
2771	3-Me C27	1.79 ± 0.37	1.61 ± 0.35	2.66 ± 0.5	2.35 ± 0.42
2799	D [a]	1.63 ± 0.74	1.5 ± 0.75	1.89 ± 0.73	2.69 ± 0.8
2831	E	tr	tr	tr	1.06 ± 0.51
2861	4-Me C28	0.61 ± 0.27	0.6 ± 0.44	tr	tr
2880	x-C29:1	tr	tr	tr	tr
2899	C29	3.7 ± 1.33	4.3 ± 1.07	8.73 ± 2.52	4.24 ± 2.58
2929	13-Me, 11-Me C29	1.02 ± 0.53	tr	1.45 ± 0.42	5.39 ± 2.21
2972	3-Me C29	tr	tr	0.8 ± 0.31	tr
2999	C30	1.68 ± 0.81	1.94 ± 0.8	1.61 ± 1.38	4.36 ± 1.77
3006	F	tr	tr	0.59 ± 0.19	tr
3028	G	tr	tr	tr	tr
3060	H	tr	tr	tr	tr
3079	x-C31:1	tr	tr	tr	tr
3098	C31	0.57 ± 0.18	0.54 ± 0.23	1.88 ± 0.76	tr
3125	13-Me C31 [b]	0.91 ± 0.42	tr	2.28 ± 0.71	1.85 ± 0.9
3172	3-Me C31	tr	tr	tr	tr
3196	I	tr	tr	0.66 ± 0.28	0.52 ± 0.25
3297	C33	0.69 ± 0.42	0.77 ± 0.61	0.93 ± 0.52	1.16 ± 0.72
3326	13-Me C33 [b]	tr	tr	tr	0.77 ± 0.42

RI - linear retention index

tr – trace amounts (<0.5% of total profile)

[a] diagnostic ions were not detectable, but RIs indicate: A=C23, D=C28

[b] peaks may include a further monomethyl alkane (11-Me) of the respective chain length at low relative concentration

Supplementary Figure S1: Crystalline deposit patterns are variable in first instar queen-destined larvae. (A,B) String of pearl-like deposits. (C,D) Blob-like deposits.

Supplementary Figure S2: Crystalline deposit patterns distinguish queen- and worker-destined larvae in several *Cardiocondyla* species. Stereomicroscope images of queen-destined (A, B, C) and worker-destined (D, E, F) third instar larvae in *C. venustula, C. wroughtonii* and *C. nuda*.

Supplementary Figure S3: Crystalline deposit patterns in *Cardiocondyla* species without caste specific patterns. Stereomicroscope images of third instar larvae in *C. "argyrotricha"*, *C. minutior*, *C. thoracica* and *C. elegans*. No pictures were taken of *C. emeryi*.

Supplementary Figure S4: Crystalline deposit patterns in seven species from four ant subfamilies available as live colonies in the lab. Stereomicroscope images of larvae in *Camponotus mus* (Formicinae), *Formica rufibarbis* (Formicinae), *Linepithema humile* (Dolichoderinae), *Platythyrea punctata* (Ponerinae) (ventral view), *Wasmannia auropunctata* (Myrmicinae), *Temnothorax sp.* (Myrmicinae). Both representatives of the Formicinae lack visible deposits.

Supplementary Figure S5: Transmission electron microscopy reveals similar external morphologies in queen and worker destined larvae of the ant *Cardiocondyla obscurior*

Supplementary Figure S6: Workers are similarly attracted to queen and worker third instar larvae. (A) number of contacts, (B) duration of contacts

Supplementary Table S1: Crystalline deposit patterns in third instar larvae predict caste in some *Cardiocondyla* species

Species	Clade	Crystalline deposit pattern	larvae with queen-like pattern			larvae with worker-like pattern		
			survival until pupation	queens	workers	survival until pupation	queens	workers
C. wroughtonii	A	queen-like + worker-like	10% (3/30)	100% (3/3)	0% (0/3)	23% (7/30)	0% (0/7)	100% (7/7)
C. nuda	B	queen-like + worker-like	75% (12/16)	100% (12/12)	0% (0/12)	50% (10/20)	10% (1/10)	90% (9/10)
C. venustula	B	queen-like + worker-like	20% (4/20)	100% (4/4)	0% (0/4)	76% (16/21)	0% (0/16)	100% (16/16)
C. thoracica	A	worker-like	-	-	-	80% (8/10)	50% (4/8)	50% (4/8)
C. minutior	B	worker-like	-	-	-	60% (12/20)	8% (1/12)	92% (11/12)

Supplementary Table S2: Overview of ant species screened for crystalline deposit patterns

Species	Subfamily	Crystalline deposit pattern	Source
Acromyrmex versicolor	Myrmicinae	worker-like	www.alexanderwild.com
Adetomyrma goblin	Amblyoponinae	worker-like	www.alexanderwild.com
Amoldius species	Dolichoderinae	worker-like	www.alexanderwild.com
Aphaenogaster picae	Myrmicinae	worker-like	www.alexanderwild.com
Aphaenogaster tennesseenis	Myrmicinae	worker-like	www.alexanderwild.com
Azteca alfari	Dolichoderinae	worker-like	www.alexanderwild.com
Calyptomyrmex piripilis	Myrmicinae	worker-like	www.alexanderwild.com
Camponotus mus	Formicinae	no visible deposits	live colonies
Camponotus sansabeanus	Formicinae	worker-like	www.alexanderwild.com
Cardiocondyla "argyrotricha"	Myrmicinae	worker-like	live colonies
Cardiocondyla elegans	Myrmicinae	worker-like	live colonies
Cardiocondyla emeryi	Myrmicinae	worker-like	live colonies
Cardiocondyla minutior	Myrmicinae	worker-like	live colonies
Cardiocondyla nuda	Myrmicinae	worker-like + queen-like	live colonies
Cardiocondyla obscurior	Myrmicinae	worker-like + queen-like	live colonies
Cardiocondyla thoracica	Myrmicinae	worker-like	live colonies
Cardiocondyla venustula	Myrmicinae	worker-like + queen-like	live colonies
Cardiocondyla wroughtonii	Myrmicinae	worker-like + queen-like	live colonies
Doleromyrma darwiniana	Dolichoderinae	worker-like	www.alexanderwild.com
Formica rufibarbis	Formicinae	no visible deposits	live colonies
Gnamptogenys mordax	Ectatomminae	worker-like	www.alexanderwild.com
Gnamptogenys striatula	Ectatomminae	worker-like	www.alexanderwild.com
Iridomyrmex victorianus	Dolichoderinae	worker-like	www.alexanderwild.com
Linepethima humile	Dolichoderinae	worker-like	live colonies
Monomorium floricola	Myrmicinae	worker-like	www.alexanderwild.com
Myrmecia piliventris	Myrmicinae	worker-like	www.alexanderwild.com
Myrmecia pilosula	Myrmicinae	worker-like	www.alexanderwild.com
Nylanderia faisonensis	Formicinae	worker-like	www.alexanderwild.com
Ochetellus species	Dolichoderinae	worker-like	www.alexanderwild.com
Odontomachus clarus	Ponerinae	worker-like	www.alexanderwild.com
Orectognathus versicolor	Myrmicinae	worker-like	www.alexanderwild.com
Paraparatrechina	Formicinae	worker-like	www.alexanderwild.com
Pheidole dentata	Myrmicinae	worker-like	www.alexanderwild.com
Pheidole desertorum	Myrmicinae	worker-like	www.alexanderwild.com
Pheidole floridana	Myrmicinae	worker-like	www.alexanderwild.com
Platythyrea punctata	Ponerinae	whole body	live colonies
Pseudomyrmex gracilis	Pseudomyrmecinae	worker-like	www.alexanderwild.com
Pseudomyrmex simplex	Pseudomyrmecinae	whole body	www.alexanderwild.com
Pseudoponera succedanae	Pseudomyrmecinae	whole body	www.alexanderwild.com

Sericomyrmex amabilis	Myrmicinae	worker-like	www.alexanderwild.com
Stigmatomma oregonensis	Amblyoponinae	worker-like	www.alexanderwild.com
Strumigenys clypeata	Myrmicinae	worker-like	www.alexanderwild.com
Strumigenys lujae	Myrmicinae	worker-like	www.alexanderwild.com
Strumigenys rostrata	Myrmicinae	worker-like	www.alexanderwild.com
Tapinoma sessile	Dolichoderinae	worker-like	www.alexanderwild.com
Temnothorax sp.	Myrmicinae	worker-like	live colonies
Temnothorax difficilis	Myrmicinae	worker-like	www.alexanderwild.com
Temnothorax rugatulus	Myrmicinae	worker-like	www.alexanderwild.com
Turneria bidentata	Dolichoderinae	worker-like	www.alexanderwild.com
Wasmannia auropunctata	Myrmicinae	worker-like	live colonies
Wasmannia rochai	Myrmicinae	worker-like	www.alexanderwild.com