Crystalline deposits reveal caste identity of late embryos and larvae of the ant

Cardiocondyla obscurior

Tobias Wallner[1]

Eva Schultner[1]

Jan Oettler[1*]

[1]Zoology/Evolutionary Biology, University of Regensburg, Universitätsstrasse 31,

93053 Regensburg, Germany

*corresponding author

ORCID:

Tobias Wallner: 0000-0001-9135-6456

Eva Schultner: 0000-0002-5069-9732

Jan Oettler: 0000-0002-8539-6029

Keywords:

Caste; social insects, ant larvae; urate; ovarian development; eco-evo-devo

Abstract

Social insects are interesting models for the study of anticipatory developmental

plasticity because of the striking differentiation into reproductive queens and

functionally sterile workers. A few ant genera, including *Cardiocondyla*, represent the

pinnacle of social evolution in the Hymenoptera, where workers have completely lost

their reproductive organs, minimizing reproductive conflicts between queens and

workers. Here we show that late embryos and larvae of queens of the ant *C. obscurior*

can be identified by the appearance of urate deposits around the forming ovaries.

The discovery of caste-specific urate patterns in *C. obscurior* and three additional

Cardiocondyla species will facilitate future studies of developmental plasticity in ants.

Introduction

Alternative developmental trajectories leading to the division of labour between reproductive queens and non-reproductive workers form the basis of superorganismality, thereby permitting one of the major transitions in evolution [1]. Research on caste development in ants has a long tradition (e.g. [2-5], and conceptual discussions are ongoing (e.g. [6,7]. However, how the processes of caste determination and differentiation are regulated at a proximate level is not well understood.

Compared to honeybees, where caste fate is under strict nutritional control, the factors underlying caste fate in ants are more diverse, ranging from genetic to socio-environmental [8]. With this comes variation in the timing of developmental divergence (e.g. [4,9], so that ants exhibit different degrees of "reproductive constraints" [9]. In some species of Ponerine ants, workers retain full reproductive potential, including the ability to mate and store sperm. In the majority of ant species workers have lost the spermatheca but retain more or less functional ovaries capable of producing haploid, male-destined eggs. Finally, workers from 11 genera completely lack ovaries. These obligately sterile workers are an example of an extended phenotype without any direct fitness, representing a highly derived form of superorganismality. The biology of some myrmicine species with fully sterile workers has been studied extensively (*Cardiocondyla obscurior*, *Monomorium pharaonis*, *Pheidole* spec., *Solenopsis invicta*), but comparably little is known about their development. Even less is known about development in the remaining seven genera with workers lacking reproductive organs (to the best of our knowledge; Myrmicinae: *Wasmannia*, *Tetramorium*, *Pheidologetum*; Dorylinae: *Eciton*; Ponerinae: *Leptogenys*, *Hypoponera*, *Anochetus*).

Across the range of reproductive constraints, a diverse set of signals spanning nature and nurture is likely to be involved in caste-specific development. Together with the facts that ant larval mobility is variable, ant brood is reared in piles, and brood is often

relocated and can serve functional roles in the colony [10], this has rendered the study

of ant development a more complex problem compared to *Drosophila* research. In

particular, it has been difficult to study caste-specific developmental trajectories

because it is not possible to distinguish worker-destined larvae from queen-destined

and male-destined larvae (e.g. [11]. Here, we close this gap for the ant *Cardiocondyla*

obscurior by showing that queen- and worker-destined embryos and larvae can be

visually distinguished by crystalline deposits surrounding the developing ovaries of

queen-destined larvae. This discovery will facilitate the study of caste determination

and differentiation at the extreme end of the superorganismality spectrum, thus

bringing us closer to a general understanding of the mechanisms facilitating the

evolution of social insects.

Methods

Ants

C. obscurior is a cosmotropical tramp ant [12], with a very streamlined genome

(~193MB, [13]), the smallest ant genome known to date. Adult queens and workers

differ in size and morphology and workers lack ovaries. Larvae develop via three larval

instars which can be distinguished by the shape of the body and the degree of

sclerotization of the mandibles [14]. The ants used in this study were all from the Old

World lineage [13], maintained in the lab since 2010. Stock colonies were kept in a

climate chamber under a 12h/12h and 22°C/26°C night/day cycle at 70% humidity.

Experimental colonies were kept in round plaster-bottom nests with nest indentations

covered by dark foil under the same conditions. All colonies were provided with water

and fed three times a week with honey and pieces of cockroaches and fruit flies.

White crystalline spots

All larvae of *C. obscurior* exhibit white spots. After producing semi-thin sections (see

below), we used a polarization filter that revealed a crystalline reflection. These

crystalline structures are common in insect larvae and have been described as urate

crystals (see discussion). Hence, because it is a parsimonious explanation we use

"urate deposits" in the following when we refer to the white spots, even though we

are aware that this may not be correct and requires future verification.

Urate deposit localization

We characterized urate deposits in eggs and first instar (L1), second instar (L2) and

third instar (L3) larvae as unpaired (= worker-destined) or paired (= queen-destined)

by visually inspecting brood from stock colonies under a stereomicroscope. For better

detection of the patterns, eggs and L1 larvae were submerged in a dissection dish

containing PBT (0.3 %), after which they were mounted on a microscope slide and

sealed with nail polish. From each development stage, we selected and

photographed one representative individual with a paired pattern and one individual

with an unpaired pattern using a stereomicroscope connected to a camera (Keyence

VHX 500FD, Neu-Isenburg, Germany).

We additionally characterized urate patterns of third instar larvae from seven

Cardiocondyla species available in the lab. We further examined brood of six species

from four subfamilies available as live colonies. Lastly, we accessed Alex Wild's photo

library for a broader overview of species

(https://www.alexanderwild.com/Ants/Natural-History/Metamorphosis-Ant-Brood/).

Caste fate and survival according to urate deposits

We tracked development of all stages to confirm that urate localization patterns are

associated with caste in *C. obscurior*. Brood was sampled from several stock colonies

and sorted by development stage and urate pattern as described above. Eggs and L1

larvae were transferred to filter paper to remove excessive buffer. Eggs were then
transferred in groups of ~15 to experimental colonies containing 10 workers (eggs:
paired=192, unpaired=165). L1 larvae were transferred to experimental colonies
containing 10 workers; two colonies were setup with 17 queen-destined larvae each,
and two colonies with 19 and 4 for worker-destined larvae, respectively (L1:
paired=34, unpaired=23). L2 and L3 larvae were transferred in groups of ten to
experimental colonies containing 10-12 workers (L2: paired=50, unpaired=40, L3:
paired=50, unpaired=40). Experimental colonies were monitored three times per
week and, when necessary, workers added from the corresponding stock colonies to
standardize worker number. All pupae were counted and classified according to
female caste until no more brood remained. From these data, we calculated average
survival until pupation and caste ratios. Survival of castes and accuracy of caste
prediction were tested with Fisher exact tests (fisher.test in R).

We tracked development of L3 larvae from six additional *Cardiocondyla* species, *C.*
wroughtonii, *C. argyrotricha*, *C. thoracica*, *C. minutior*, *C. nuda*, *C. venustula*,
spanning the phylogeny with exception of the palearctic clade [15], to validate that
urate deposit patterns accurately predict caste.

Larval histology

L3 larvae were collected from stock colonies of *C. obscurior* and sorted according to
their urate patterns. Sorted larvae were transferred into a fixation solution consisting
of 25% glutaraldehyde (GAH) in cacodylate buffer [(50 mM cacodylic acid, pH 7.3)
containing 150 mM Sucrose] (GAH : cacodylate buffer = 1 : 12,5) and kept overnight
at 11 °C. Samples were then rinsed in cacodylate buffer on ice, and fixated in 4%
osmium tetroxide in cacodylate buffer. After fixation, larvae were washed in
cacodylate buffer, dehydrated in a graded ethanol series and embedded in Epon.
Transversal semithin sections of 1 µm were cut and stained with methylene blue and

Azur II (Richardson's stain). Semi-thin sections were scanned with a Zeiss Primo Star

microscope and imaged with a Moticam 580 digital microscope camera.

Ovary development

C. obscurior queen L2 and L3 larvae were dissected in PBT 0.3% (PBS 1x + Triton

0.3%) by cutting below the midgut with a micro scissor. The posterior part of the larvae

was cleared of excess fat tissue. The larval ovaries were placed in an ice-cooled well

filled with PBT. We chose two different stages of queen pupae based on their

development. Early-stage pupae were identified as unsclerotized white pupae with

unpigmented ommatidia and ocelli. Mid-stage pupae exhibit pigmented ommatidia

and ocelli. Ovaries from queen pupae and adults were obtained by carefully pulling

on the last tergite of the abdomen with forceps, which removes the entire

reproductive apparatus. Larval, pupal and adult ovaries were fixated in 4%

paraformaldehyde diluted in PBS for 20 min at room temperature. The fixated ovaries

were then washed three times with PBT for 15 min on a tumbler. After washing, the

ovaries were processed for staining.

Vasa protein was stained with a rat anti-vasa antibody and actin filaments were

visualized using a rabbit anti-actin antibody. Both antibodies were generated from

Drosophila [9]. Cell nuclei were stained using DAPI. The primary antibodies were

diluted 1:200 in PBT 0.3% with 5% normal goat serum overnight at room temperature

on a tumbler. To remove the primary antibody, the larval and pupal ovaries were

washed three times with PBT for 15 min. The secondary antibodies goat anti-rat and

goat anti-rabbit were used to visualize the rat anti-vasa and rabbit anti-actin

antibodies, both diluted 1:200 in PBT. After 2 h of incubation at room temperature,

the ovaries were again washed three times with PBT for 15 min each. Finally, the larval,

pupal and adult ovaries were washed with PBT containing DAPI (1:10000). Ovaries

were then mounted in Vectashield on a microscope slide and sealed with nail polish.

Images for the antibody staining were obtained using a Leica SP8 confocal

microscope under 40x and 63x objective lenses. Complete confocal series of ovarioles

of early- and mid-staged queen pupae were processed using the FIJI plugin TrakEM2

[16] for 3D rendering and analysis.

Nanos and *vasa* expression was measured in queens and workers from all larval

stages. L1 and L2 larvae were collected and transferred to individual 1.5 ml tubes (L1:

n=5 queen larvae, n=7 worker larvae; L2: n=5 queen larvae, n=5 worker larvae). L3

larvae were cut beneath the midgut at the posterior end (containing the developing

ovaries in queen larvae) and transferred to individual tubes (L3: n=5 queen larvae, n=4

worker larvae). All samples were immediately frozen in liquid nitrogen and stored at -

75°C prior to RNA extraction. Total RNA was extracted using the ReliaPrep RNA Cell

Miniprep Kit (Promega) and RNA concentrations measured using the Qubit RNA HS

Assay Kit (Invitrogen). RNA concentrations were standardized to 2.5 ng and RNA

reverse-transcribed to cDNA using the iScript gDNA Clear cDNA Synthesis Kit (Bio-

Rad Laboratories). Expression of the gene *nanos* was quantified with qPCR with the

primer pair nos_for/nos_rev and normalized with two housekeeping genes

(Y45F10D_JO1_for/Y45F10D_JO1_rev [17]) and Actin_JO1_for/Actin_JO1_rev; see

Table S1 for primer sequences). All qPCR reactions were performed in triplicates and

specificity of reactions confirmed with manual melt curve inspection. Relative target

gene expression was calculated as $2^{-\Delta Cq}$ following [18], using the geometric mean of

the two housekeeping genes for normalization.

Results

Queen and worker-destined eggs and larvae of *C. obscurior* can be distinguished, by

what appear to be urate crystals. In late embryos (Figure 1) and the three larval stages

(Figure 2, Figure S1), urate deposit patterns associated with queen caste are visible.

Queen patterns appear either as a string of pearls or are snail shell-like (Figure S2).

Handling L1 larvae results in high mortality, but L2 and L3 individuals can be attributed

to the two castes with high precision (Table 1).

Transversal semithin sections of L3 larvae revealed that worker larvae lack ovaries and

have urate deposits which are distributed randomly throughout the caudal end (Figure

3 A). These cluster in small aggregates (10-20 µm diameter) in proximity to

trophocytes (Figure 3 C, E). In queen larvae, urate crystals form voluminous

aggregations surrounding the developing ovaries, which contain several dozens of

developing oocytes (Figure 3 B, D, F). In both castes urate comes in the shape of

translucent birefringent spherocrystals.

Urate deposit patterns allowed for selection of queen-destined L2 and L3 larvae,

making it efficient to reconstruct ovarian development. In L2 larvae the developing

ovaries are visible as a bilateral cluster of germ cells located at the posterior end of

the last abdominal segments (Figure 4 A-D). The ovaries are in close proximity to the

caudal end of the central nervous system (Figure 4 D, CNS). The cluster of cells in L2

larvae then differentiate into three ovarioles per ovary in L3 larvae (Figure 4 E-H), with

the tips extending in an anterior-apical orientation (Figure S8). Disc-shaped cells can

be found which stack in a medial-lateral orientation (Figure 4 H, asterisk). These cells

are likely cap cells [19], which separate the germarium from the terminal filament [20].

In pupae, the ovaries are completely differentiated, with fully extended terminal

filaments (Figure 4 I-L). The germ cells in the germarium have differentiated into

cystocytes, which will later give rise to the oocyte with the associated nurse cells

(Figure 4 L) [21]. After reaching the adult stage, the fully matured ovarioles have egg

chambers, containing oocytes with their corresponding nurse cells (Figure 4 M-P). The

oocytes are surrounded by follicle cells, and these egg chambers are in different

stages of maturation (Figure 4 P). Reconstruction of the cystocytes inside the ovarioles

of early- and mid-staged queen pupae showed that the developing cells increase

significantly in size during pupation (Linear regression, R^2=0.9036, p<0.001) and that

cell size is highly variable (Figure S3).

The expression of the germline marker *nanos* was higher in L2 and L3 queen larvae

compared to worker larvae (Figure S4, L2: Mann-Whitney-U-test: W=24, p=0.016; L3:

W=20, p=0.016, Figure S4), but not in L1 larvae (W=25, p=0.27). *nanos* expression

increased over the course of development in queens, matching the developmental

state of the ovaries. The variation in *nanos* expression is conspicuously high in L2

queen larvae. In contrast, *vasa* expression was significantly higher in L1 (W=33,

p=0.010) and L3 queen larvae (W=20, p=0.016), but not in L2 larvae (W=17, p=0.421),

confirming that sampling based on urate patterns was successful, despite failure to

show this for L1 larvae in rearing experiments.

We found queen-like urate patterns in larvae of three additional *Cardiocondyla*

species (*C. venustula*, *C. wroughtonii* and *C. nuda*) (Figure S5, Table S2). In the

remaining four species (*C. thoracica*, *C. elegans*, *C. argyrotricha* and *C. minutior*) we

did not find obvious caste-specific patterns (Figure S6, Table S2).

39 of 41 additional species representing 30 genera from seven ant subfamilies had

larvae with obvious urate deposits (Table S3; Figure S7), but no species exhibited

paired urate deposits resembling the pattern observed in *Cardiocondyla* queen

larvae.

Discussion

"Auch ein blindes Huhn findet mal ein Korn" (German saying)

Cardiocondyla is unique among the Formicidae due to the evolution of wingless

fighter males and the repeated loss of the winged male morph across the genus [15].

Because colonies of *C. obscurior* can be easily reared and manipulated, the species

has been used to address questions in diverse biological contexts [22,23], ranging

from genome evolution [13,24], to social behavior [25], aging [26,27], symbiosis

[28,29], and phenotypic plasticity [17,30,31]. In this respect, *C. obscurior* has been

considered a "gold mine" (E. Abouheif pers. com.). Now, as is often the case in

mining, sheer luck has led to the discovery that queen and worker embryos and larvae

in some *Cardiocondyla* species can be distinguished using the localization of urate

deposits. We could not find a similar pattern in other ant genera, but 39 species from

30 genera is by far not representative of the vast diversity of ants. We hope that our

discovery inspires others to take a closer look at the brood of their favorite species.

In insects, urate is typically deposited in specialized cells – so-called urocytes [32] –

which together with trophocytes [32] and oenocytes [33] represent the three major

cell types in the fat body. Urocytes store urates either in vacuoles or as circular,

birefringent spherocrystals [32], and can occupy a large fraction of the fat body, for

instance in the termite *Mastotermes darwiniensis* [32]. In the termite *Reticulitermes*

flavipes, uric acid is recycled by gut bacteria as a nitrogen source [34], and urate

deposits have been linked to oocyte development [35]. In *Pachycondyla* ants,

urocytes specialize in the storage of nutrients and excretion products [36] and a

composition assay in *Melipona quadrifasciata* bees identified Na, Ca, Mg, P as main

elements in urocytes, as well as traces of Zn, Mn, and K [33]. In *C. obscurior* we found

circular, birefringent spherocrystals but no cell nuclei, indicating that urate is

deposited directly. Worker larvae exhibited small, randomly distributed urate clusters

which were found closely associated with what appeared to be trophocytes rather

than urocytes. These smaller clusters may originate as a by-product of metabolic

processes in trophocytes. Queen larvae showed dense aggregations of urate

surrounding the developing ovaries. These large urate deposits may constitute an

energy reservoir for cellular growth, a process that is energy-demanding since cells in

and around ovaries proliferate at rapid rates.

Caste-specific urate patterns only occurred in some *Cardiocondyla* species and in

none of the species representing other ant genera. At the moment, we do not know

why this is the case. It is unlikely that the presence of the bacterial symbiont *Cand.*

Westeberhardia cardiocondylae is directly linked, although in principle it makes for a

promising candidate. This endosymbiont with a highly eroded genome resides in

bacteriomes associated with the gut and, prominently, with the ovaries in queens, indicating a nutritional link. However, the same urate patterns occur in a *C. obscurior* lineage that is naturally free of *Cand.* Westeberhardia [29], as well as in larvae which have been experimentally cleared of their bacterial symbionts with antibiotic treatment [28]. There is also no obvious phylogenetic signal. *C. obscurior* and its sister species *C. wroughtonii* both exhibit caste-specific patterns but cluster together in clade A with *C. thoracica* and *C. argyrotricha*, both of which do not show the pattern. In clade B, *C. nuda* and *C. venustula,* which exhibit the caste-specific patterns, cluster together with *C. minutior* [15]. A welcome additional coincidence is that all four species with the distinct queen pattern have two alternative male morphs [15] and can be kept in the lab with minimum effort, making the genus perfect for comparative studies of female and male diphenic development.

Urate deposit patterns allowed us to reconstruct ovarian development in *C. obscurior.* The female reproductive apparatus is conserved in insects, and consists of two ovaries, the oviduct, the uterus, accessory glands and the spermatheca. Large variation can be found in the number of ovarioles making up each ovary, ranging from 1 in a dung beetle, 15-20 in the fruit fly *Drosophila melanogaster* [19], ~200 in honey bees [37] to more than 1000 in some termite and ant species [38]. Six ovarioles in *C. obscurior* and *Cardiocondyla nuda* queens [39] mark the lower end in ants, thus allowing detailed study of ovary development in a simple model. In *C. obscurior*, the ovaries develop from a cluster showing early differentiation in L2 larvae to ovaries with differentiated ovarioles in L3 larvae. Nevertheless, the size of urate deposits in L2 larvae is similar to that in L3 larvae. Future studies investigating development prior to the L2 stage will reveal when exactly queen and worker phenotypes diverge, and uncover how urate deposits are related to ovary formation.

Germline stem cells (GSCs) are typically located in the germarium at the tip of the ovarioles, close to the terminal filament cells ([9,20]. GSCs continuously renew themselves and produce new germ cells. They then proliferate and produce

differentiated cystoblasts, which divide synchronously to form cystocytes [21]. These

cystocytes are found in the germarium of pupal ovarioles in *C. obscurior,* with some

cells larger than others. Cystocystes go on to differentiate into oocytes with their

corresponding nurse cells. We suspect that the largest cells are pre-determined

oocytes in the pupal germarium, which allow to initiate oocyte maturation soon after

hatching. *Cardiocondyla* queens mate inside their natal nest within days of reaching

adulthood and can almost immediately begin with the production of offspring [27].

The discovery of caste-specific patterns has important implications for understanding

both the biology of *C. obscurior* and the general mechanisms underlying caste

development in ants. At the life history level, it is now possible to study questions

revolving around the phenotypes of developing queens and workers. For example,

do the larval phenotypes differ in morphology, behavior or cuticular chemistry? We

can also test whether workers discriminate between queen- and worker-destined

brood, and if larvae are treated differently depending on caste. In the future, it should

even be possible to track individual larvae without the need for tagging by using their

unique patterns for individual recognition, similar to the way fins are used to study

humpback whales [40]. Creativity limits possibilities now, not vice versa.

Because caste-specific urate patterns are already visible in late embryos, we are also

finally able to study caste development beyond differentiated features such as wing

discs in late larvae [41]. The establishment of *nanos* and *vasa* as reliable expression

markers further allows for verification of individual caste identity in molecular studies.

C. obscurior queens produce increasingly queen-biased caste ratios with age [27],

facilitating efficient sampling of queen-destined embryos, and opening numerous

avenues for the study of the mechanisms involved in ant caste determination and

differentiation, and in social insect polyphenism in general. It has already been shown

that ant and honeybee development shares some features, with an overlapping set of

genes involved in growth (e.g. TOR, Insulin-like) [42]) and juvenile hormone synthesis

and degradation [43] playing a role in developmental regulation [8,30]. Upstream, the

importance of transcription factors involved with the sex differentiation cascade also finds support in ants [17,44], and honeybees [45,46]. Recently, we postulated that a major evolutionary transition, the evolution of sex, has facilitated another major transition, the evolution of eusociality, via co-option of the same developmental switch mechanism [17]. Now we can test this hypothesis, bringing us one step closer to a conceptual understanding of how social insect polyphenism evolved.

Acknowledgements

We thank Alexandra Koch for help rearing embryos, Kristina Pogorelski and Laura Spichtinger for help rearing larvae, Birgit Lautenschläger and Maria Schiwek for help with histology, Erhard Strohm for help identifying the crystalline nature of the deposits, Christopher Winkler for taking pictures, and Lina Pedraza for providing ant species. JO dedicates this paper to Margit Oettler, for nature and nurture. This study was funded by DFG grants (DFG OE549/3-1, OE 549/3-2, OE 549/3-3).

References

1. Boomsma, J. J. & Gawne, R. 2017 Superorganismality and caste differentiation as points of no return: how the major evolutionary transitions were lost in translation. *Biological Reviews* **471**, E1. (doi:10.1111/brv.12330)

2. Light, S. F. 1942 The Determination of the Castes of Social Insects. *The Quarterly Review of Biology* **17**, 312–326. (doi:10.1086/394661)

3. Gößwald, K. & Bier, K. 1954 Untersuchungen zur Kastendetermination in der Gattung Formica. *Insect. Soc.* **1**, 229–246. (doi:10.1007/BF02222948)

4. Brian, M. V. 1973 Caste control through worker attack in the ant Myrmica. *Insect. Soc.* **20**, 87–102. (doi:10.1007/BF02223340)

5. Brian, M. V. 1975 Caste determination through a queen influence on diapause in larvae of the ant *Myrmica rubra. Entomologia Experimentalis et Applicata* **18**, 429–442. (doi:10.1111/j.1570-7458.1975.tb00420.x)

6. Trible, W. & Kronauer, D. J. C. 2017 Caste development and evolution in

ants: it's all about size. *Journal of Experimental Biology* **220**, 53–62.
(doi:10.1242/jeb.145292)

7. Abouheif, E. 2021 Ant caste evo-devo: it's not all about size. *Trends in
Ecology & Evolution* (doi:10.1016/j.tree.2021.04.002)

8. Corona, M., Libbrecht, R. & Wheeler, D. E. 2016 Molecular mechanisms
of phenotypic plasticity in social insects. *Current Opinion in Insect
Science* **13**, 55–60. (doi:10.1016/j.cois.2015.12.003)

9. Khila, A. & Abouheif, E. 2010 Evaluating the role of reproductive
constraints in ant social evolution. *Philosophical Transactions of the Royal
Society B: Biological Sciences* **365**, 617–630.
(doi:10.1098/rstb.2009.0257)

10. Schultner, E., Oettler, J. & Helanterä, H. 2017 The role of brood in
eusocial hymenoptera. *The Quarterly Review of Biology*, 1–41.

11. Pontieri, L., Rajakumar, A., Rafiqi, A. M., Larsen, R. S., Abouheif, E. &
Zhang, G. 2020 From egg to adult: a developmental table of the ant
Monomorium pharaonis. *bioRxiv*, 2020.12.22.423970.
(doi:10.1101/2020.12.22.423970)

12. Heinze, J., Cremer, S., Eckl, N. & Schrempf, A. 2006 Stealthy invaders:
the biology of *Cardiocondyla* tramp ants. *Insect. Soc.* **53**, 1–7.
(doi:10.1007/s00040-005-0847-4)

13. Errbii, M., Keilwagen, J., Hoff, K. J., Steffen, R., Altmüller, J., Oettler, J. &
Schrader, L. 2021 Transposable elements and introgression introduce
genetic variation in the invasive ant Cardiocondyla obscurior. *Molecular
Ecology* (doi:10.1111/mec.16099)

14. Schrempf, A. & Heinze, J. 2006 Proximate mechanisms of male morph
determination in the ant *Cardiocondyla obscurior*. *Evol Dev* **8**, 266–272.
(doi:10.1111/j.1525-142X.2006.00097.x)

15. Oettler, J., Suefuji, M. & Heinze, J. 2010 The evolution of alternative
reproductive tactics in male Cardiocondyla ants. *Evolution* **64**, 3310–
3317. (doi:10.1111/j.1558-5646.2010.01090.x)

16. Cardona, A., Saalfeld, S., Schindelin, J., Arganda-Carreras, I., Preibisch,
S., Longair, M., Tomancak, P., Hartenstein, V. & Douglas, R. J. 2012
TrakEM2 software for neural circuit reconstruction. *PLoS ONE* **7**, e38011.
(doi:10.1371/journal.pone.0038011)

17. Klein, A., Schultner, E., Lowak, H., Schrader, L., Heinze, J., Holman, L. &

Oettler, J. 2016 Evolution of social insect polyphenism facilitated by the
sex differentiation cascade. *PLoS Genet* **12**, e1005952–e1005952.
(doi:10.1371/journal.pgen.1005952)

18.   Schmittgen, T. D. & Livak, K. J. 2008 Analyzing real-time PCR data by the
comparative CT method. *Nat Protoc* **3**, 1101–1108.
(doi:10.1038/nprot.2008.73)

19.   Sahut-Barnola, I., Godt, D., Laski, F. A. & Couderc, J.-L. 1995 Drosophila
ovary morphogenesis: analysis of terminal filament formation and
identification of a gene required for this process. *Dev. Biol.* **170**, 127–
135. (doi:10.1006/dbio.1995.1201)

20.   Xie, T. 2013 Control of germline stem cell self-renewal and differentiation
in the Drosophila ovary: concerted actions of niche signals and intrinsic
factors. *Wiley Interdiscip Rev Dev Biol* **2**, 261–273. (doi:10.1002/wdev.60)

21.   Okada, Y., Miyazaki, S., Miyakawa, H., Ishikawa, A., Tsuji, K. & Miura, T.
2010 Ovarian development and insulin-signaling pathways during
reproductive differentiation in the queenless ponerine ant Diacamma sp.
J. Insect Physiol. **56**, 288–295. (doi:10.1016/j.jinsphys.2009.10.013)

22.   Oettler, J. 2020 *Cardiocondyla*: Heart Node Ants. In *Encyclopedia of*
Social Insects (ed C. Starr), pp. 1–4.(doi:doi.org/10.1007/978-3-319-
90306-4_19-1)

23.   Heinze, J. 2017 Life-history evolution in ants: the case of *Cardiocondyla*.
Proc. Biol. Sci. **284**, 20161406. (doi:10.1098/rspb.2016.1406)

24.   Schrader, L. et al. 2014 Transposable element islands facilitate
adaptation to novel environments in an invasive species. *Nature*
Communications **5**, 5495. (doi:10.1038/ncomms6495)

25.   de Menten, L., Cremer, S., Heinze, J. & Aron, S. 2005 Primary sex ratio
adjustment by ant queens in response to local mate competition. *Animal*
Behaviour **69**, 1031–1035. (doi:10.1016/j.anbehav.2004.09.005)

26.   Oettler, J. & Schrempf, A. 2016 Fitness and aging in Cardiocondyla
obscurior ant queens. *Current Opinion in Insect Science* **16**, 58–63.
(doi:10.1016/j.cois.2016.05.010)

27.   Jaimes Nino, L. M., Heinze, J., Oettler J., 2021 Late-life fitness gains
explain the absence of a selection shadow in ants. *bioRxiv* 2021
(doi:https://doi.org/10.1101/2021.02.24.432699)

28.   Ün, Ç., Schultner, E., Manzano-Marín, A., Flórez, L. V., Seifert, B., Heinze,

J. & Oettler, J. 2021 Cytoplasmic incompatibility between Old and New
World populations of a tramp ant. *Evolution* (doi:10.1111/evo.14261)

29.   Klein, A. et al. 2015 A novel intracellular mutualistic bacterium in the
invasive ant *Cardiocondyla obscurior. ISME J* **10**, 376–388.
(doi:10.1038/ismej.2015.119)

30.   Schrader, L., Simola, D. F., Heinze, J. & Oettler, J. 2015 Sphingolipids,
transcription factors, and conserved toolkit genes: Developmental
plasticity in the ant *Cardiocondyla obscurior. Molecular Biology and*
Evolution **32**, 1474–1486. (doi:10.1093/molbev/msv039)

31.   Schrader, L., Helanterä, H. & Oettler, J. 2017 Accelerated Evolution of
Developmentally Biased Genes in the Tetraphenic Ant Cardiocondyla
obscurior. *Molecular Biology and Evolution* **34**, 535–544.
(doi:10.1093/molbev/msw240)

32.   Costa-Leonardo, A. M., Laranjo, L. T., Janei, V. & Haifig, I. 2013 The fat
body of termites: functions and stored materials. *J. Insect Physiol.* **59**,
577–587. (doi:10.1016/j.jinsphys.2013.03.009)

33.   Furtado, W. C. A., Azevedo, D. O., Martins, G. F., Zanuncio, J. C. &
Serrão, J. E. 2013 Histochemistry and ultrastructure of urocytes in the
pupae of the stingless bee Melipona quadrifasciata (Hymenoptera:
Meliponini). *Microsc Microanal* **19**, 1502–1510.
(doi:10.1017/S1431927613013445)

34.   Potrikus, C. J. & Breznak, J. A. 1981 Gut bacteria recycle uric acid
nitrogen in termites: A strategy for nutrient conservation. *Proceedings of*
the National Academy of Sciences **78**, 4601–4605.
(doi:10.1073/pnas.78.7.4601)

35.   Elliott, K. L. & Stay, B. 2007 Juvenile hormone synthesis as related to egg
development in neotenic reproductives of the termite *Reticulitermes*
flavipes, with observations on urates in the fat body. *Gen Comp*
Endocrinol **152**, 102–110. (doi:10.1016/j.ygcen.2007.03.003)

36.   Zara, F. J. & Caetano, F. H. 2004 Ultramorphology and histochemistry of
fat body cells from last instar larval of the Pachycondyla (=Neoponera)
villosa (Fabricius) (Formicidae: Ponerinae). *Braz J Biol* **64**, 725–735.
(doi:10.1590/s1519-69842004000400022)

37.   Hartfelder, K., Tiberio, G. J., Lago, D. C., Dallacqua, R. P. & Bitondi, M.
M. G. 2018 The ovary and its genes—developmental processes
underlying the establishment and function of a highly divergent

reproductive system in the female castes of the honey bee, Apis
mellifera. *Apidologie* **49**, 49–70. (doi:10.1007/s13592-017-0548-9)

38. Church, S. H., de Medeiros, B. A. S., Donoughe, S., Márquez Reyes, N. L.
& Extavour, C. G. 2021 Repeated loss of variation in insect ovary
morphology highlights the role of development in life-history evolution.
Proc. Biol. Sci. **288**, 20210150. (doi:10.1098/rspb.2021.0150)

39. Heinze, J., Kühnholz, S., Schilder, K. & Hölldobler, B. 1993 Behavior of
ergatoid males in the ant, Cardiocondyla nuda. *Insect. Soc.* **40**, 273–282.
(doi:10.1007/BF01242363)

40. Mizroch, S. A., Beard, J. A. & Lynde, M. 1990 Computer assisted photo-
identification of humpback whales. *Rep Int Whale Comm*

41. Oettler, J., Platschek, T., Schmidt, C., Rajakumar, R., Favé, M.-J., Khila,
A., Heinze, J. & Abouheif, E. 2019 Interruption points in the wing gene
regulatory network underlying wing polyphenism evolved independently
in male and female morphs in Cardiocondyla ants. *J. Exp. Zool. (Mol.*
Dev. Evol.) **332**, 7–16. (doi:10.1002/jez.b.22834)

42. Mutti, N. S., Dolezal, A. G., Wolschin, F., Mutti, J. S., Gill, K. S. &
Amdam, G. V. 2011 IRS and TOR nutrient-signaling pathways act via
juvenile hormone to influence honey bee caste fate. *Journal of*
Experimental Biology **214**, 3977–3984. (doi:10.1242/jeb.061499)

43. Kayukawa, T., Minakuchi, C. & Namiki, T. 2012 Transcriptional regulation
of juvenile hormone-mediated induction of Krüppel homolog 1, a
repressor of insect metamorphosis. (doi:10.1073/pnas.1204951109/-
/DCSupplemental)

44. Jia, L.-Y., Chen, L., Keller, L., Wang, J., Xiao, J.-H. & Huang, D.-W. 2018
Doublesex Evolution Is Correlated with Social Complexity in Ants.
Genome Biology and Evolution **10**, 3230–3242.
(doi:10.1093/gbe/evy250)

45. Johnson, B. R. & Jasper, W. C. 2016 Complex patterns of differential
expression in candidate master regulatory genes for social behavior in
honey bees. *Behav Ecol Sociobiol* **70**, 1033–1043. (doi:10.1007/s00265-
016-2071-9)

46. Pan, L.-X., Cheng, F.-P. & Wang, Z. L. 2021 Caste-biased Expression of
fem and Amdsx Genes in Apis mellifera ligustica (Hymenoptera: Apidae).
Sociobiology **68**, 5654. (doi:10.13102/sociobiology.v68i2.5654)

Table 1: Urate deposit patterns predict caste in the ant *Cardiocondyla obscurior.*
Survival differences between castes and accuracy of caste prediction were tested
using Fisher's exact tests.
506
507

Development stage	Predicted caste	n	Survival			Produced caste		
			Proportion (%)	Differences between castes		Proportion (%)	Accuracy of caste prediction	
				Odds ratio	p		Odds ratio	p
egg	queen	165	24.9 (41/165)	1.276	0.342	78 (32/41)	5.398	<0.001
	worker	192	29.7 (57/192)			42.1 (24/57)		
1st instar larva	queen	34	5.9 (2/34)	1.512	1	0 (0/2)	-	-
	worker	23	8.7 (2/23)			100 (2/2)		
2nd instar larva	queen	50	62 (31/50)	0.334	0.019	90.3 (28/31)	97.129	<0.001
	worker	40	35 (14/40)			92.9 (13/14)		
3rd instar larva	queen	50	60 (30/50)	0.439	0.062	100 (30/30)	infinite	<0.001
	worker	40	42.5 (17/40)			100 (17/17)		

Figure legends

Figure 1: Urate deposit patterns distinguish queen- and worker-destined larvae in the

ant *Cardiocondyla obscurior*. Stereomicroscope images of worker-destined larval

instars (A, C, E) and queen-destined larval instars (B, D, F). Ur = urate.

Figure 2: Urate localization in queen- and worker-destined embryos in the ant

Cardiocondyla obscurior. Worker-destined embryos show no urate (ur) (A), while

queen-destined embryos have urate localized in the last segments of their germ band

(gb) (B). The first segments of the embryo are clearly separated into the head and the

three gnathal segments: mandibular (md), maxillary (mx) and labial (lb).

Figure 3: Histological sections of *Cardiocondyla obscurior* larvae. (A) Transversal

section of a worker-destined L3 larva. In workers ovaries are missing. (B) Queen-

destined larvae show paired ovaries (ov) close to the midgut (mid). (C) In worker-

destined larvae urate (ur) accumulates in small randomly distributed deposits. (D)

Ovaries in queen-destined larvae are surrounded by large urate deposits. (E) Small

urate deposits aggregate close to trophocytes (t). (F) Urate encloses the developing

ovary in queen larvae. Tf = terminal filament.

Figure 4: Ovarian reconstruction in queens of *Cardiocondyla obscurior*. (A-D) Early

gonadal formation in second instar larva. DAPI is used to show the cell nuclei. Vasa

stains against germ cells and actin visualizes the cytoskeleton. (A) DAPI staining, (B)

Vasa staining, (C) Actin staining, and (D) Merged image with a visible central nervous

system (CNS). (E-H) Ovarioles (ov) of a third instar queen larva. (E) DAPI, (F), Vasa, (G)

Actin and (H) Merge. The asterisk (*) shows cap cells. (I-L) Ovary of a queen pupa with

its six ovarioles. (I) DAPI, (J), Vasa, (K) Actin and (L) Merge. The germ cells have

differentiated into cystocytes in this stage, inhabiting the caudal end of the ovariole.

Tf = terminal filament. (M-P) Mature ovarioles in adult queens. (M) DAPI, (N) Vasa, (O)

Actin and (P) Merge showing the oocyte (oo) with its nurse cells (nc). The oocyte is

encircled by follicle cells (fc).

Supplementary Figure legends

Supplementary Figure 1: Dorsal and ventral view of urate patterns in worker- and

queen-destined larvae in the ant *C. obscurior*. Stereomicroscope images of worker-

destined larval instars (A, C & E) and queen-destined larval instars (B, D & F). (A, B)

show larvae in ventral orientation, (C-F) show larvae in ventral orientation. Urate

deposits are visible as bilateral symmetrical clusters in queen-destined larvae.

Supplementary Figure 2: Types of urate patterns in queen-destined larvae. (A, B) First

instar larva with string of pearl-like urate deposits. (C, D) First instar larva with snail

shell-like urate deposits.

Supplementary Figure 3: Images of queen pupae. (A) Early-stage queen pupa with

unpigmented ommatidia and ocelli. (B) Mid stage queen pupa with pigmented

ommatidia and ocelli. (C) Example of a 3D rendered ovariole with TrakEM2. Data

generated form the reconstruction was used for data analysis. Colors represent

individual cells. (D) Cell volume to cell surface in *C. obscurior* queen pupae. The plot

illustrates the different sizes of cells in developing cystocytes of queen ovarioles from

two pupal stages.

Supplementary Figure 4: Caste-specific expression of the germline marker *nanos* in

the ant *C. obscurior*. Normalized gene expression of the germline marker in queens

and workers in all three larval stages. Expression of *nanos* is significantly higher in L2

and L3 queen larvae compared to worker larvae. * $P < 0.05$, Mann-Whitney-U-test.

Supplementary Figure 5: Urate patterns distinguish queen and worker-destined larvae

in various *Cardiocondyla* species. Stereomicroscope images of queen-destined larval

instars (A, B, C) and worker-destined larval instars (D, E, F). (A) Queen larva of *C.*

venustula (B) Queen larva of *C. wroughtonii*. (C) Queen larva of *C. nuda*. (D) Worker

larva of *C. venustula*. (E) Worker larva of *C. wroughtonii*. (F) Worker larva of *C. nuda*.

Ur = urate.

Supplementary Figure 6: Urate patterns in *Cardiocondyla* species without paired urate

deposits. Stereomicroscope images of larval instars. (A) *C. argyrotricha*. (B) *C.*

minutior. (C) *C. thoracica*. (D) *C. elegans*.

Supplementary Figure 7: Urate localization patterns in seven species from four ant

subfamilies. Stereomicroscope images of larval instars. (A) *Camponotus mus*

(Formicinae). (B) *Formica rufibarbis* (Formicinae). (C) *Linepithema humile*

(Dolichoderinae). *Platythyrea punctata* (Ponerinae) (dorsal view). (E) *Wasmannia*

auropunctata (Myrmicinae). (F) *Temnothorax sp.* (Myrmicinae). Both representatives of

the Formicinae are lacking visible urate deposits.

Supplementary Figure 8: Schematic illustration of postembryonic ovarian

development in the ant *C. obscurior*. The sketched images represent the 2nd instar (A),

3rd instar (B) and pupal (C) stage of queens. In the 3rd instar the lateral oviduct (red) is

already distinct. The green coloration represents the germ cells as seen in Figure 4.

Figure 2

Figure 3

Figure 4